White-crowned Sparrow environmental songs

RESEARCH ARTICLE

Machine learning reveals relationships between song, climate, and migration in coastal

Zonotrichia leucophrys

Jiaying Yang[1], Bryan C. Carstens[1], Kaiya L. Provost[1*]

1: Museum of Biological Diversity & Department of Evolution, Ecology and Organismal

Biology, The Ohio State University, 1315 Kinnear Rd., Columbus OH, USA, 43212.

* Corresponding author: provost.27@osu.edu

ACKNOWLEDGEMENTS

We thank the Carstens Lab past and present, as well as D. Nicholson and A. Delgado for helpful

feedback. We thank two anonymous reviewers, one associate editor, and T. S. Sillett for reviews.

Funding statement: KLP was funded by NSF DEB-2016189. BCC was funded by NSF DBI-

1661029, DEB-2016189, NSF DEB-1831319, and DBI-1910623. JY was funded by the

Undergraduate Diversity at Evolution program.

Author contributions: Experimental design: JY, KLP, BCC. Data collection: JY. Methods

development: KLP, BCC. Analyses: JY, KLP. Writing: JY. Figures: JY, KLP. Editing: JY, KLP,

BCC.

Data depository: Data will be available upon acceptance on Dryad. Scripts will be available on

GitHub.

ABSTRACT

Vocalization behavior in birds, especially songs, strongly affects reproduction, but it is also

highly impacted by geographic distance, climate, and time. For this reason, phenotypic

differences in vocalizations between different bird populations are often interpreted as evidence

of lineage divergence. Previous work has demonstrated that there is extensive variation in the

songs of White-crowned Sparrow (*Zonotrichia leucophrys*) throughout the species range,

including between neighboring (and genetically distinct) subspecies *Z. l. nuttalli* and *Z. l.

pugetensis*. However, it is unknown whether the divergence in their songs correlates to

environmental or geographical factors. Previous work has been hindered by time-consuming

traditional methods to study bird songs that rely on the manual annotation of song spectrograms

into individual syllables. Here we explore the performance of automated machine learning

methods of song annotation, which can process large datasets more efficiently, paying attention

to the question of subspecies differences. We utilize a recently published artificial neural

network to automatically annotate hundreds of White-crowned Sparrow vocalizations across two

subspecies. By analyzing differences in syllable usage and composition, we find that *Z. l. nuttalli*

and *Z. l. pugetensis* have significantly different songs. Our results are consistent with the

interpretation that these differences are caused by the changes in syllables in the White-crowned

Sparrow repertoire. However, the large sample size enabled by the AI approach allows us to

demonstrate that divergence in song is correlated with environmental difference and migratory

status, but not with geographical distance. Our findings support the hypothesis that the evolution

of vocalization behavior is affected by environment, in addition to population structure.

Keywords: Bird song; Machine learning; climate; cultural drift; Bioacoustics, *Zonotrichia*

leucophrys.

LAY SUMMARY

● Birdsong is an important behavior because it is important in bird communication and

reproduction.

● White-crowned Sparrows in western North America are known to use different songs

along their range, but it is unknown if those songs vary due to the environment.

● We used machine learning to analyze these songs and found that populations of White-

crowned Sparrows can be differentiated based on their songs.

● Environmental factors during the breeding season exert a greater influence on song

evolution in migratory subspecies.

INTRODUCTION

59
Vocalization is an important method of bird communication (Catchpole and Slater 2008, Brumm

and Zollinger 2013), especially in *Passeriformes*. Unlike other sensory communication channels

(e.g., visual, chemical, tactile), vocalizations can transmit long distances and have a fast rate of

evolutionary change, due in part to cultural transmission (Slater 1986, Slater 1989, Slabbekoorn

and Smith 2002a, Podos and Warren 2007, Derryberry 2009; but see Mason et al. 2017). Bird

songs are vocalizations that are generally important for reproductive effort. While it was

previously thought that songs were mostly produced by males during the breeding seasons, with

common functions of territory defense and mating (Kroodsma and Byers 1991), more recent

observations demonstrate that females also produce songs, and that songs are produced in non-

breeding contexts (DeWolfe et al 1989, Baptista et al 1993, DeWolfe and Baptista 1995, Odom

et al 2014, Webb et al 2016, Riebel et al 2019).

Song evolution is influenced by biotic and abiotic factors. Vocal behaviors vary among

species and play a role in species recognition (Beer 1971, Emlen 1972, Baker et al. 1981b,

Freeberg et al 1995). Even closely related populations can have different songs (i.e., dialects)

that they can use to tell each other apart (Thompson et al. 1993). The trend of song development

in birds is toward complexity and variety (i.e., the anti-monotony principle; Hartshorne 1956),

mainly due to sexual selection, including male-male competition and female preference (Morton

1982, Morton 1986, Catchpole and Slater 2008). In male-male competition, listeners can

determine the singer's distance by assessing the amount of degradation of signals so that singers

tend to increase the number of different song types in their repertoire (i.e., the ranging

hypothesis; Morton 1982, Morton 1986).

The level of degradation and attenuation of signals is also affected by the habitat

environment due to different transmitting efficacy (Acoustic Adaptation Hypothesis; Morton

1975, Hunter and Krebs 1979). Lower frequencies and slower notes transmit more effectively

through denser vegetation, and birds living in those habitats have songs with those characteristics

(Slabbekoorn and Smith 2002a, Naguib 2003, Kopuchian et al. 2004, Seddon 2005, Baker 2006,

van Dongen and Mulder 2006, Derryberry 2009). Likewise, birds in urbanized habitats sing at

higher frequencies to prevent overlap with low-frequency anthropogenic noise (Derryberry et al.

2016, Derryberry et al. 2020). Geographically restricted song variants emerge through time

(Podos and Warren 2007), resulting in a pattern analogous to isolation-by-distance (Nelson et al

2004). In sum, the wide variety of bird songs likely results from any combination of sexual

selection, different habitats/environments, or other biotic and abiotic factors.

White-crowned Sparrow

One well-studied songbird is the White-crowned Sparrow (*Zonotrichia leucophrys*), which has

variable song across North America (Baptista 1989). This species is known to sing year-round,

not just during the breeding season, making them good candidates for understanding

environmental impacts on their sounds (Blanchard 1941, Baptista 1974, Brenowitz et al. 1998).

Non-breeding singing in *Z. l. pugetensis* appears to be associated with flock hierarchy formation

as well as hormone stimulation (DeWolfe and Baptista 1995), in contrast with *Z. l. nuttalli* which

sings due to a protracted breeding season.

White-crowned Sparrows are known to have divergent dialects across different populations

(Petrinovich and Patterson 1981, Baker et al. 1984, Lien and Corbin 1990). We focus here on

Zonotrichia leucophrys nuttalli and *Z. l. pugetensis*. These subspecies have within-species

dialects, as well as divergent songs across populations associated with genetic differentiation

(Lipshutz et al. 2017), suggesting that they are distinct evolutionary units. In addition, previous

estimates show that dialects present in *Z. l. pugetensis* are more widely distributed than in *Z. l.*

nuttalli (DeWolfe and Baptista 1995). Songs help to identify these two subspecies, both for

researchers and for the birds themselves. The songs of these birds vary such that the subspecies

can tell each other apart, suggesting that reproductive isolation is evolving. The subspecies have

an area of hybridization in Mendocino County, CA, where intermediate individuals are found

(Lipshutz et al 2017). Generally, subspecies *Z. l. nuttalli* (Figure 1A) and *Z. l. pugetensis* (Figure

1B) differ in the length of introductory whistle notes, presence of buzz notes, and

length/presence of terminal trill notes (Dunn et al. 1995; Baptista and King 1980), though

variation exists within populations and through time. Song is directly implicated in reducing

gene flow between these populations, which have been at least partially genetically isolated for

~45,000 years (Lipshutz et al 2017). The different songs of both *Z. l. nuttalli* and *Z. l. pugetensis*

also show a geographic pattern within the subspecies, which are proposed to have been formed

by limited or biased dispersal, morphological variation, and/or habitat (Marler and Tamura

1962, Marler and Tamura 1964, Barker and Mewaldt 1978, Baker and Thompson 1985, Nelson

2000, Nelson et al. 2004, Kroodsma et al. 1985, reviewed by Podos and Warren 2007,

Derryberry 2009).

Though the two subspecies are phenotypically similar, with only slight differences in body

size (Dunn et al. 1995), they do vary in their migratory behavior and geographic distributions.

Subspecies *Z. l. nuttalli* is a non-migratory subspecies that is distributed in coastal California

(from ~40.4 to 34.3 degrees latitude; Banks 1963; Figure 2). In contrast, *Z. l. pugetensis* is a

migratory subspecies that is distributed from British Columbia to northwest California during the

breeding season, both coastal and inland, and expands to the southern part of its range through

southern California during the non-breeding season (Banks 1963, Patten et al. 2003, Baptista

1974, Heinemann 1981; Figure 2). Subspecies *Z. l. pugetensis* thus shares its wintering grounds

with three other subspecies: *Z. l. nuttalli* , which is phenotypically similar, as well as *oriantha*

and *gambelii*, which are distinguished by differences in color of the beak and the lores (Welke et

al 2021, Lisovski et al 2019). There is evidence that White-crowned Sparrows can learn songs

during migration or on the wintering grounds (in *Z. l. oriantha*, Baptista and Morton 1988).

Despite this, *Z. l. pugetensis* populations having the same breeding-season dialect tend to migrate

to the same location in the non-breeding season (Baptista 1974, Heinemann 1981, Nelson 2000,

Nelson et al 2004) despite those locations not being segregated by song dialect (DeWolfe and

Baptista 1995).

Though there is compelling evidence that the divergence of White-crowned Sparrow songs is

caused by sexual selection (Marler and Tamula 1962, Nelson and Poesel 2013), there is less

evidence on whether the evolution of vocalization behavior is correlated to the abiotic

environment, such as climate, or geographic distance as a proxy for cultural evolution. Distinct

differences in the histories of these two subspecies may be mediating climatic adaptation. The

two subspecies overlap extensively in their non-breeding grounds. Traditionally, *Z. l. pugetensis'*

breeding and wintering grounds were thought to be distinct, with the breeding not occurring

south of Oregon and wintering only occurring in California (DeWolfe and Baptista 1995).

However, *Z. l. pugetensis* experienced a range shift east through the Cascade Mountains and

North into Alaska because of habitat change and logging regimes (Gibson et al 2013, Hunn and

Beaudette 2014, Welke et al 2021). Additionally, although *Z. l. pugetensis* is assumed to migrate

south to overlap with *Z. l. nuttalli*, birds identified as *Z. l. pugetensis* have been found *north* of

their breeding grounds during the winter (Sullivan et al 2009; GBIF 2023). As such, it is likely

that these shifts in location may be related to the adaptation of *Z. l. pugetensis* to environmental

change. In this paper, we use these White-crowned Sparrow subspecies to investigate whether

the divergence in songs within and between subspecies is correlated to these abiotic factors

(climate and geographic distance). We hypothesize that the different songs are correlated to

climates and geographical differences between the two subspecies. We also hypothesize that *Z. l.*

pugetensis songs may be influenced more by the climate than that of *Z. l. nuttalli*, under the

assumption that they are migratory to avoid harsh winter conditions (Somveille et al 2015), and

that this impact should be more apparent during the breeding season.

Machine learning

Bird songs have been studied for a long time with many methods for qualitatively and

quantitatively describing their sounds having been developed. Traditional methods require

researchers to analyze and measure song parameters manually. Since the mid 20^{th} century, the

most popular methods of analysis use spectrograms, which display frequencies through time

using fast Fourier transforms (Borror and Reese 1953; Cooley and Tukey 1965; Figure 1). Many

analyses require every individual syllable or note on the spectrogram to be identified and

annotated, or segmented, for downstream processing. Segmentation and annotation are time

intensive and the amount of data that can be processed is limited (Swiston et al. 2009).

Recently, machine learning and artificial intelligence techniques have been used broadly in

biological science (e.g., Schrider and Kern 2017, Pelletier and Carstens 2018, Viener et al 2022).

Machine learning has also been applied to syllable segmentation with high accuracy via the

neural network TweetyNet (Cohen et al. 2020). Originally trained on single species, the

algorithm has since been applied to a suite of birds (Provost et al 2022) with similarly high

performance, allowing for the use of larger datasets. It is unknown whether this and other

automatic segmentation methods effectively recapitulate known differences in populations. Here,

we examine whether we can apply automatic syllable recognition to the different songs of White-

crowned Sparrow songs.

METHODS

Recording data and metadata

We downloaded our sound data from two sources: the Borror Laboratory of Bioacoustics (BLB; blb.osu.edu) directly from the database and the Xeno-Canto database (XC; xeno-canto.org) by using *query_xc* function in the *warbleR* package version 1.1.26 (Araya-Salas and Smith-Vidaurre 2017) in R version 4.0.3 (R Core Team 2020). We used all recordings of *Z. l. pugetensis* and *Z. l. nuttalli* in the BLB. On XC, we restricted recordings to be graded as Quality "A." We transformed the XC recordings from MP3 format to WAV format by using the *readMP3* function in the *tuneR* package version 1.3.3 (Ligges et al. 2018) in R. We also converted recordings to have a sample rate of 48,000 Hz if needed using the *resamp* function in the *seewave* package version 2.2.0 (Sueur et al 2008) in R. Recordings were made between 1965 and 2021 and included metadata subspecies identification, date, locality, and recordist.

Although many recordings were identified to subspecies in the BLB and XC when we acquired them, both databases contained recordings from the region of interest that were not identified to subspecies. We assigned individuals without known subspecies identity using latitude and time of year as criteria following Lipshutz et al. (2017). We assigned all breeding season "unknown" individuals north of 40.544 °N and south of 48.461 °N to be *Z. l. pugetensis*, and all breeding season "unknown" individuals south of 38.419 °N and north of 37.803 °N to be *Z. l. nuttalli*, because they do not have overlapping distribution within these regions during summer. Based on the distribution, we also reassigned some mis-identified birds in the original datasets. We did not assign "unknown" individuals between the two regions because the two subspecies intergraded in that area. Moreover, "unknown" individuals that were further than ~110 km (one decimal degree) from the coast were not classified as either subspecies as they likely belonged to subspecies *Z. l. oriantha* or *Z. l. gambelii*. For non-breeding season birds, we

did not assign "unknown" individuals due to substantial amounts of overlap between *Z. l.*

nuttalli, Z. l. pugetensis, Z. l. gambelii, and *Z. l. oriantha* which all overwinter through the

region. Any birds where subspecies identity or time of recording was not known were excluded.

Our recordings were mostly restricted to the west coast of the United States and Canada.

However, there were three localities from Utah and eastern California that are in high-elevation

habitats (above 1900 m), which are colder, more variable in temperature, less variable in

precipitation, and drier (Fick and Hijmans 2017). As these localities were likely to influence our

results, we removed them from our downstream analyses. Our final dataset included 1,111

recordings from the breeding season (547 *Z. l. pugetensis*, 564 *Z. l. nuttalli*) and 802 recordings

from the non-breeding season (391 *Z. l. pugetensis*, 411 *Z. l. nuttalli*) for a total of 1,913.

Manual Annotation

We segmented and annotated syllables present in our recordings. Segmentation is the process of

going from a raw song file to a list of syllables present in the recording, including their time and

frequency boundaries. Annotation is then the labeling of those syllables by syllable type (see

below). For segmentation, we followed the methodology of a previous study using White-

crowned Sparrows (Provost et al. 2022). 201 of these samples were segmented for that study; we

manually added syllable type to those existing segmentations.

For manual segmentation and annotation, we input the recording to Raven Pro version

1.6.1 (K. Lisa Yang Center for Conservation Bioacoustics at the Cornell Lab of Ornithology

2019) and generated a spectrogram using a Fast Fourier Transform size of 512 samples. We

segmented the spectrogram by drawing boxes that show the maximum and minimum values of

the frequency and time of each syllable. We annotated five distinct types of syllables in these

data which represent the types of syllables previously identified in *Z. leucophrys*. First, we define

whistles (W) as long, single-note introductory phrases. Second, we define buzzes (B) as notes

that rapidly go from high to low bandwidth. Third, we define trills (T) as individual notes that

are repeated at the end of the song. Fourth, we define calls (C) as short high-pitched contact

notes which are not typically used in songs. Notes that were identified as calls were excluded

from downstream analyses. Lastly, we define all other notes as "special" syllables (S) if they did

not fall into these categories. All annotations were output as .TXT files. We then split the

original recordings and their corresponding annotation files into subsets that had approximately

equal amounts of annotated syllables and silence using previously generated scripts (Provost et

al. 2022).

Using Machine Learning for Bioacoustics

To automatically segment and annotate the White-crowned Sparrow songs, we trained a

TweetyNet model on our *Zonotrichia leucophrys* data following a previously described

procedure (Provost et al. 2022). Although the previous study did have a model trained on

Zonotrichia leucophrys, it was not trained on our specific syllable types from *Z. l. nuttalli* and *Z.*

l. pugetensis. We used vak version 0.6.0 and TweetyNet version 0.8.0 for that and we used

identical parameters to (Provost et al. 2022). We annotated 573 buzzes, 67 calls, 1,518 special

syllables, 1,513 trills, and 658 whistles for 4,329 syllables. We used 80% of those as training

data, 10% as validation data, and 10% as test data. After training, we discarded anything labeled

as a call.

After training the model, we had the model predict the locations and annotations of syllables

for all data, including data that TweetyNet had never seen before. We assessed model

performance (syllable error rates, accuracy, precision, recall, and f-score) for each subspecies

and per syllable type. We calculated precision and recall for each syllable type separately and

then averaged them to give an overall precision and recall score.

Summary statistics

Once we had syllables segmented and annotated for all recordings, we extracted bioacoustic data

for each syllable. We extracted two types of summary statistics: one set was general to all birds,

and the other was specific to White-crowned Sparrows. For the general summary statistics, data

were calculated using the *spectro_analysis* function in the *warbleR* package in R. The data

extracted from syllables include bandwidth (the difference between highest frequency and lowest

frequency of the syllable), time duration (the length of the syllable), center frequency (central

frequency between the upper and lower cutoff frequencies), average slope (slope of the change in

dominant frequency), and inflection points (the local extrema). For each recording, we calculated

the centroid value across all syllables. We then calculated a distance between centroids between

all individuals within and between subspecies, which we used as a representation of song

distance.

For the White-crowned Sparrow specific summary statistics, we calculated ten metrics

that were previously identified as being important for distinguishing the two focal subspecies

(Lipshutz et al 2017). These metrics are as follows: 1) the duration, 2) bandwidth, and 3) rate of

the terminal trill; 4) the duration of the initial whistle notes; 5) the duration of the introductory

phrase, including the initial whistle note as well as any buzzes and unique notes; 6) the

maximum and 7) minimum dominant frequency present in the song; 8) the mean syllable

duration, and 9) the mean dominant frequency of the first whistle note alone or 10) all initial

whistle notes. We omitted metrics of vocal performance in this case.

Some of these metrics required knowledge of the position of the syllables in the recording

and there were some ambiguities with how these values should be calculated. We defined the

introduction as all notes before the first trill (T) note identified in the song; if not trills were

identified, the introduction is the same length as the entire song. We defined the initial whistle

notes as being all whistles present in the introduction if whistles were present. Finally, we

defined the trill as being the duration from the first to the last trill in the song. These metrics

were calculated for each song within the recording and then averaged across all songs in a single

recording.

Principal Components Analysis and ANOVAs

We performed a principal components analysis (PCA) on the vocal summary statistics present in

our syllables using the *prcomp* function in the *stats* package version 4.1.2 in R. We centered and

scaled all data. When values for the raw statistics were missing for the PCA specifically we filled

them in with the mean value for the data. The PCA results were used to compare differences

between the subspecies and seasons.

To identify the principal components (PCs) we used downstream, we used the broken-stick

model (Bro and Smilde 2014) and kept all the initial PCs that explained more variation than

expected under that model. We used an ANOVA on both the raw statistics and results of PCA to

test the differences between the two subspecies and two seasons (breeding and non-breeding)

using the *aov* function in the *stats* package in R. Given that we examined 15 raw variables (and

one PC, see results) and would have a high false discovery rate across these ANOVAs, we used a

Bonferroni correction (Bonferroni 1936) and set our significance value to 0.0031 from an alpha

value of 0.05.

Multiple Matrix Regression with Randomization

To determine whether differences in songs were being driven by differences in either the climate

or geographic distance, we used multiple matrix regression with randomization analysis

(MMRR) in R (Wang 2020). This technique allows one to interpret correlations between

environmental features and geography to assess each's relative contribution. To calculate

geographic distances, we calculated the Euclidean distance between individuals with respect to

latitude and longitude. For climatic differences, we downloaded 19 climate variables from

WorldClim (Fick and Hijmans 2017) and extracted values at each locality in our dataset. We

excluded three individuals (two *Z. l. nuttalli*, one *Z. l. pugetensis*) for whom climate information

was lacking. We then calculated Euclidean distance between individuals with respect to those

values. MMRR models were created such that we predicted song divergence from geographic

distances and climatic differences simultaneously (i.e., song divergence ~ geographic distance +

climatic distance). We also performed a second set of MMRR models where we estimate those

parameters plus temporal distance, derived from the year of the recording (i.e., song divergence

~ geographic distance + climatic distance + time distance). *Zonotrichia leucophrys* is known to

have experienced dramatic changes in song over the period of this study and so we wished to

compare our results with and without controlling for temporal changes (Derryberry 2009). Given

that we examined nine MMRR models with and without time, we used a Bonferroni correction

and set our significance value to 0.0056 from an alpha value of 0.05.

Because *Z. l. pugetensis* is a subspecies that migrates through the year, we separated our data

by seasons: breeding (May to August) and non-breeding (September to April; Chilton et al.

2020). We performed these MMRR analyses described above with only the breeding and non-

breeding datasets separately because the different ranges due to migration of *Z. l. pugetensis* may

result in variation in geographical distance and ecological distance.

Environmental Comparisons

Given that the impact of environment on the White-crowned Sparrow songs differs between

subspecies and seasons, we wished to determine whether White-crowned Sparrows in our

different datasets occupy different climate regimes. We performed a PCA on the 19 WorldClim

variables used in our MMRRs. We did this by extracting the climate data from all 307 unique

localities that White-crowned Sparrows were found at in our dataset, and then performing a PCA

on those data only. As with the PCA on song values, we used the broken stick model to select

PCs to retain. We then used ANOVAs to compare whether seasons and subspecies differed in

their environments. Given that we examined 19 raw environmental variables (and two PCs, see

results) we used a Bonferroni correction and set our significance value to 0.0024 from an alpha

value of 0.05.

RESULTS

336
First, we analyzed how the machine learning model performed in syllable annotation. These

models had high performance across both subspecies. Our model predicted 403 syllables on the

test dataset. The syllable error rate of our trained machine learning model's annotations was -

11% for *Z. l. nuttalli* and -1% for *Z. l. pugetensis* (Table 1), such that for both subspecies, the

model predicted fewer syllables than were present via manual annotation. This suggests that the

models either miss some syllables outright or lump syllables together that should be separated.

Although the model has a better segment error rate for *Z. l. pugetensis,* the model classifies *Z. l.*

nuttalli better on all other metrics. Segmentation of the model into sounds vs background noise

performed well overall (accuracy >90%, precision >96%, recall >82%, F-score>88%). Recall

was the worst performing metric which was driven by the model labeling some syllables as

silence. When looking at annotation of known syllables only (excluding background noise)

performance is notably worse (accuracy >73%, precision >84%, recall >74%, F-score>79%).

This appears to be driven largely by trills (T) which the model struggles to classify correctly

(Figure 3). Trills are often misclassified as "special" (S) notes or silence.

PCA

We found that our PCA on syllable variables explained a high proportion of the vocal data, with

the first three PCs explaining 33%, 15%, and 11% of variation, respectively (Table 2). Under the

broken stick model these numbers were expected to be 22, 15, and 12%, respectively. As such,

we retained only PC1 for downstream analyses. Syllables with high PC1 had low whistle

dominant frequencies, few inflection points, low maximum and center frequencies, and short

syllable introduction durations. They also had wide bandwidths (Table 2).

We found that our summary statistics of song did differentiate subspecies and seasons (Table

4). PC1 was significantly higher in *Z. l. nuttalli* than *Z. l. pugetensis* (p<0.0001). Most of the

underlying song statistics that PC1 was calculated from also differed significantly between

subspecies. Subspecies *Z l. pugetensi*s had songs with steeper slopes, wider trill bandwidths,

shorter whistles, longer introductory phrases and overall syllables, higher whistle and overall

song frequencies, more inflection points, and shorter and faster trills. Likewise, PC1 was higher

in the breeding season than the non-breeding season, although this relationship was only nearly

significant after correcting for multiple tests (p=0.0104). Despite PC1 not being significant, some

of the individual statistics did show differences. Breeding season birds have narrower trill

bandwidths, longer trills, lower whistle dominant frequencies but higher minimum dominant

frequencies, steeper slopes, and fewer inflection points than non-breeding season birds.

MMRR

We found that geography explained song divergence for both species together, irrespective of

season ($R^2 > 0.01$, geography $p<0.001$; Table 5; Table 6). Climate is a significant factor during

the breeding season and year-round ($p<0.001$), but only nearly significant during the non-

breeding season (p=0.013). After we account for temporal changes, these patterns are identical,

but year of recording is also significant ($R^2 > 0.05$, time $p<0.001$, climate $p<0.016$, geography

$p<0.001$).

For *Z. l. nuttalli* alone, we found a relationship between climate and song in the non-

breeding season ($R^2 = 0.004$, climate p=0.004, geography p=0.098). However, this relationship is

not present when accounting for time, suggesting that year of recording explains this pattern (R^2

$= 0.010$, time p=0.001, climate p=0.092, geography p=0.148). There was no impact of geography

or climate on song for *Z. l. nuttalli* during the breeding season. After accounting for time, there is

an association during the breeding season between song and time, but climate and geography are

both nearly significant ($R^2 = 0.05$, time p=0.001, climate p=0.002, geography p=0.007). In

contrast for the full year, while there was no impact of geography and climate on song before

accounting for time, time itself does significantly explain song.

As for *Z. l. pugetensis*, in the non-breeding season there is an association between song

and geography ($R^2 = 0.075$, climate p=0.385, geography p=0.001); this association holds up after

accounting for time, but time is not a significant predictor. Year-round there is an association

between song and climate ($R^2 = 0.001$, climate p=0.001, geography p=0.643), but this

association is only near significant after accounting for time, with time instead explaining the

variation. Finally, while there was no impact on *Z. l. pugetensis* song in the breeding season

before accounting for time, after accounting for time both time and climate are significant. In all

instances where year of recording significantly explains song divergence, it is likely that this is

reflecting patterns of cultural evolution caused by young sparrows learning their songs from

adults near them (Baptista and Morton 1988).

Environmental Comparisons

The first three environmental PCs (Env.PCs) explained 63%, 21%, and 10% of the

environmental variation, respectively (Supplementary Material Table S1). Under the broken stick

model, these should have explained 18%, 13%, and 11%, respectively. As such we retained the

first two PCs. High values of Env.PC1 indicated environments that were cold, wet, and had little

seasonal precipitation variation (Supplementary Material Table S2, Supplementary Material

Table S3). High values of Env.PC2 were associated with temperature, specifically environments

that had elevated temperature seasonality and annual range, and little isothermality.

ANOVAs revealed that environments were significantly different between seasons and

subspecies (Supplementary Material Table S4, Supplementary Material Table S5). Subspecies

significantly differ across the Env.PC1 and Env.PC2, with *pugetensis* occupying colder, wetter,

and more variable areas. Seasons also differ in Env.PC1 such that breeding season birds are on

average in areas that are more seasonal and less isothermal. These patterns likely reflect

differences in where individual birds are located during the breeding and non-breeding seasons,

especially as *pugetensis* migrates south.

DISCUSSION

415
Our analysis demonstrates that the trained machine learning model by TweetyNet has high

accuracy with low error rates and can annotate *Z. l. nuttalli* and *Z. l. pugetensis* songs. Our

results suggest that the annotation by the machine learning model is as reliable as manual

annotations and produces comparable data, although training with more diverse syllable types

may improve the classification of our models, particularly on trill syllables. Given its

computational efficiency, machine learning can be a useful tool to automatically annotate bird

songs and extract data because it can be used to process large datasets quickly and accurately.

We found significant differences in the syllables used in the songs of *Z. l. nuttalli* and the

songs of *Z. l. pugetensis*, which corroborates previous evidence that these two taxa have

divergent songs (e.g., Lipshutz et al. 2017). Our bioacoustic metrics found that on average, *Z. l.*

nuttalli syllables were lower in pitch, had fewer inflections, and had more negative slopes

compared to *Z. l. pugetensis,* as well as differing in trill duration and bandwidth. Our analyses

cannot distinguish if this is due to subspecies using different syllables or the same syllables at

different frequencies. Preliminary analyses of syllable shape suggest that the differences are

mediated using different syllables, but further investigation is needed to confirm.

We also found significant differences between the seasons in some metrics. Syllables used in

the breeding season differed in trill bandwidth, whistle dominant frequency, slope, and

inflections compared to the non-breeding season. In other species of songbird, singing in the

non-breeding season is less consistent due to differences in hormone levels (e.g., the Song

Sparrow *Melospiza melodia*, Smith et al 1997; the Rufous-collared Sparrow *Zonotrichia*

capensis, Addis et al 2010), which is consistent with our findings. At the same time, this species

modulates its song with respect to the environment, being especially impacted by noise

transmission (Derryberry 2009, Luther et al 2016, Derryberry et al 2020). Additionally, distinct

parts of *Zonotrichia leucophrys* song appear to have distinct functions (Nelson and Poesel 2007):

introductory whistles (as well as buzzes) are highly conserved and serve as species identifiers,

terminal trills are moderately conserved and indicate dialect, and other notes in the song are

highly variable and indicate individual identity. Whether this species is responding to their

environments in different seasons or simply singing less consistently due to a lack of

testosterone, there is clearly detectable variation associated between the seasons and the

environment.

The differences in song between subspecies and seasons are compounded by the fact that the

two subspecies have different responses to climate, geographic distance, and time. Almost every

analysis showed that year of recording partially or fully predicted variation in song (except for

non-breeding season *Z. l. pugetensis*). Thus, the impact of cultural drift on this taxon cannot be

overstated and corroborates previous analyses (Derryberry 2009). Geographic distance is

important when both subspecies are analyzed simultaneously, regardless of season. Additionally,

the migratory *Z. l. pugetensis* also shows a high impact of geographic distance on songs recorded

in the non-breeding season. In contrast, there is mixed evidence whether geographic distance

explains song variation in the sedentary *Z. l. nuttalli.* The impact of climate on song is important

when both species are analyzed together year-round and during the breeding season. In contrast,

climate does not explain variation in either subspecies after accounting for time. The evidence

suggests that while cultural drift through time is important for these two taxa in explaining song

differences, overall, the migratory subspecies *Z. l. pugetensis* shows higher influence of spatial

factors, particularly geographic distance, though only during the non-breeding months. This may

be driving the patterns where both species are combined in those seasons; alternatively, the two

subspecies may still be adapted to their relatively different environments. It is likely that these

factors (i.e., migration and seasonality) have contributed to the evolution of song differences.

We performed an analysis of the environments that each of the subspecies occupy during

different seasons and found that these climates are significantly different. Year-round, *Z. l.*

nuttalli occupies warmer and drier areas than *Z. l. pugetensis* does in the summer. In contrast,

wintering *Z. l. pugetensis* experience nearly the full yearly range of temperature and precipitation

across both subspecies, and overall experience more variable and more seasonal climates than *Z.*

l. nuttalli. A synthesis across multiple birds hypothesizes that migration occurs as a mechanism

to avoid harsh winters and exploit resource surpluses (Somerville et al 2015); given that the

range of *Z. l. pugetensis* is wide-spread and has been changing in response to habitat disruption

(Welke et al 2021, Hunn and Beaudette 2014, Gibson et al 2013), we hypothesize that *Z. l.*

pugetensis as a subspecies may be more susceptible to spatial heterogeneity in their environment,

leading to more variation in their songs. In addition, because *Z. l. pugetensis* is known to have

dialects ranging over larger areas that encapsulate more variable climates (DeWolfe and Baptista

1995), it is possible that the selection pressures to adapt song to environment is higher across the

range of this subspecies.

Caveats

Several caveats are worth documenting to place our results in the appropriate context. All

analyses suggest that geographic distance is not correlated to the divergence in song across these

two subspecies, which is not consistent with our hypothesis. Some explanations are as follows.

First, although the number of recordings of *Z. l. pugetensis* in non-breeding habitats is large, they

do not cover the entire projected breeding range. Second, our recordings are all from the BLB

and XC database from 1965 to 2021, resulting in substantial cultural drift. Over the time scale of

the study, White-crowned Sparrows are known to have demonstrably changed their songs (e.g.,

Derryberry 2009). However, even if we control for the year of recording, we still find significant

relationships with spatial factors like geography and climate. Third, our analysis only used each

syllable's characteristics and types but did not analyze the syntax or order of syllables. It is

possible that they used the same syllables but different orders in some populations. Fourth,

because White-crowned Sparrows learn their songs not only directly from their parents but also

from the neighboring community (Cunningham and Baker 1983) and that individuals in hybrid

zones can learn the songs of the "incorrect" subspecies (Chilton et al 1990), it is possible that

late-hatched *Z. l. nuttalli* may be exposed to *Z. l. pugetensis* that have already arrived in their

winter habitats. It is hard to answer these questions with only the recordings and data of their

locality and habitats. However, the strong positive relationship between ecological differences

and song in the non-breeding season corresponds with our original hypothesis. Fifth and finally,

we did not investigate the other properties of the environment, such as the habitat type, instead

focusing only on abiotic climate. Further research on whether the abiotic environment affects the

evolution of bird songs directly or indirectly via biotic environmental factors is needed.

In conclusion, using machine learning for automatic analyses, we evaluated divergence

within and between two subspecies of the White-crowned Sparrow. Our results indicated that

Zonotrichia leucophrys pugetensis and *Z. l. nuttalli* have significantly different songs from each

other, and both sing different songs in breeding and non-breeding contexts. The divergence in

songs correlate with spatial factors such as geographic distance and climate, but subspecies

identity and season mediate the response.

LITERATURE CITED

508
Araya-Salas, M., and G. Smith-Vidaurre (2017). "warbleR: an r package to streamline analysis

of animal acoustic signals." https://dx.doi.org/10.1111/2041-210X.12624.

Addis, E. A., D. S. Busch, A. D. Clark, and Wingfield, J. C. (2010). Seasonal and social

modulation of testosterone in Costa Rican rufous-collared sparrows (*Zonotrichia capensis*

costaricensis). *General and comparative endocrinology* 166:581-589.

Baker, M. C. (2006). Differentiation of mating vocalizations in birds: Acoustic features in

mainland and island populations and evidence of habitat-dependent selection on songs.

Ethology 112:757-771. https://doi.org/10.1111/j.1439-0310.2006.01212.x

Baker, M. C., D. B. Thompson, and G. L. Sherman (1981b). Neighbor-stranger song

discrimination in White-crowned Sparrows. Condor 83:265-267.

Baker, M. C., and D. B. Thompson (1985). Song dialects of White-crowned Sparrows: historical

processes inferred from patterns of geographic variation. *The Condor* 87:127-141.

Baker, M. C., M. A. Cunningham, and A. D. Thompson Jr. (1984). Cultural and genetic

differentiation of two subspecies of White-crowned Sparrow. Condor 86:359-367.

Baker, M. C., and M. A. Cunningham (1985). The Biology of Bird-Song Dialects. Behavioral

and Brain Sciences 8:85-100. https://doi.org/10.1017/S0140525X00019750

Baker, M. C., and L. R. Mewaldt (1979). The use of space by White-crowned Sparrows: juvenile

and adult ranging patterns and home range versus body size comparisons in an avian

granivore community. Behavioral Ecology and Sociobiology 6:45-52.

Banks, R. C. (1963c). Geographic variation in the White-crowned Sparrow *Zonotrichia*

leucophrys. University of California Publications in Zoology 70:1-123.

Baptista, L. F. (1974). The Effects of Songs of Wintering White-crowned Sparrows on Song

Development in Sedentary Populations of the Species. Zeitschrift Für Tierpsychologie,

34:147-171. https://doi.org/10.1111/j.1439-0310.1974.tb01794.x

Baptista, L. F. (1989). The best studied songbird: White-crowned Sparrows in the wild and in the

laboratory. Birds International 1:10-21.

Baptista, L. F., and J. R. King. (1980). Geographical variation in song and song dialects of

montane white-crowned sparrows. *The Condor 82*: 267-284.

Baptista, L. F., and M. L. Morton. (1988). Song learning in montane white-crowned sparrows:

from whom and when. *Animal Behaviour 36*:1753-1764.

Baptista, L. F., P. W. Trail, B. B. DeWolfe, and M. L. Morton. (1993). Singing and its functions

in female white-crowned sparrows. *Animal behaviour 46:*511-524.

Baker, M. C., and L. R. Mewaldt. (1978). Song dialects as barriers to dispersal in white-crowned

sparrows, *Zonotrichia leucophrys nuttalli*. *Evolution*:712-722.

Blanchard, B. D., E. R. Hall, F. Richardson, F. H. Test, J. T. Boulware, L. W. Adams, and W. H.

Behle. (1941). *The white-crowned sparrows (*Zonotrichia leucophrys*) of the Pacific

seaboard: environment and annual cycle*. University of California Press.

Bonferroni, C. (1936). Teoria statistica delle classi e calcolo delle probabilita. *Pubblicazioni del

R Istituto Superiore di Scienze Economiche e Commericiali di Firenze 8*:3-62.

Beer, C. G. (1971). Individual Recognition of Voice in the Social Behavior of Birds. Advances

in the Study of Behavior, 3:27-74.

BirdLife International and Handbook of the Birds of the World (2021) Bird species distribution

maps of the world. Available at http://datazone.birdlife.org/species/requestdis.

Borror, D. J., and C. R. Reese (1953). The analysis of bird songs by means of a vibralyzer. The

Wilson Bulletin 65:271-276.

Brenowitz, E. A., L. F. Baptista, K. Lent, and J. C. Wingfield (1998). Seasonal plasticity of the

song control system in wild Nuttall's white-crowned sparrows. Journal of neurobiology,

34:69-82.

Bro, R., and A. K. Smilde. (2014). Principal component analysis. *Analytical methods 6:*2812-

2831.

Brumm, H., and S. A. Zollinger (2013). Avian Vocal Production in Noise. In Animal

Communication and Noise. (Brumm, H., Editor). Animal Signals and Communication 2:187-

227. https://doi.org/10.1007/978-3-642-41494-7_7

Catchpole, C. K., and P. J. B. Slater (2008). Bird Song: Biological Themes and Variations.

Cambridge University Press, Cambridge, UK.

Catchpole, C. K. (1987). Bird song, sexual selection and female choice. In Trends in Ecology

and Evolution 2:94-97. https://doi.org/10.1016/0169-5347(87)90165-0

Chilton, G., M. C. Baker, C. D. Barrentine, and M. A. Cunningham (2020). White-crowned

Sparrow *(Zonotrichia leucophrys)*, version 1.0. In Birds of the World (A. F. Poole and F. B.

Gill, Editors). Cornell Lab of Ornithology, Ithaca, NY, USA.

https://doi.org/10.2173/bow.whcspa.01

Chilton, G., M. R. Lein, and L. F. Baptista. (1990). Mate choice by female white-crowned

sparrows in a mixed-dialect population. *Behavioral Ecology and Sociobiology 27*:223-227.

Cohen, Y., D. A. Nicholson, A. Sanchioni, E. K. Mallaber, V. Skidanova, and T. J. Gardner

(2022) Automated annotation of birdsong with a neural network that segments spectrograms.

eLife 11:e63853. https://doi.org/10.7554/eLife.63853

Cooley, J. W., and J. W. Tukey (1965). An algorithm for the machine calculation of complex

Fourier series. Mathematics of computation 19:297-301.

Cortopassi, A. J., and L. R. Mewaldt (1965). The circumannual distribution of White-crowned

Sparrows. Bird-banding 36:141-169.

Cunningham, M. A., and M. C. Baker (1983). Vocal learning in White-crowned Sparrows:

sensitive phase and song dialects. Behavioral Ecology and Sociobiology 13:259-269.

https://doi.org/10.1007/BF00299673

Derryberry, E. P. (2009). Ecology shapes birdsong evolution: Variation in morphology and

habitat explains variation in white-crowned sparrow song. American Naturalist 174:24-33.

https://doi.org/10.1086/599298

Derryberry, E. P., R. M. Danner, J. E. Danner, G. E. Derryberry, J. N. Phillips, S. E. Lipshutz, K.

Gentry, and D. A. Luther (2016). Patterns of song across natural and anthropogenic

soundscapes suggest that White-crowned Sparrows minimize acoustic masking and

maximize signal content. PLoS ONE 11:e0154456.

Derryberry, E. P., J. N. Phillips, G. E. Derryberry, M. J. Blum, and D. Luther (2020). Singing in

a silent spring: Birds respond to a half-century soundscape reversion during the COVID-19

shutdown. Science 370:575-579.

DeWolfe, B. B., and L. F. Baptista. (1995). Singing behavior, song types on their wintering

grounds and the question of leap-frog migration in Puget Sound White-crowned Sparrows.

The Condor 97:376-389.

DeWolfe, B. B., L. F. Baptista, and L. Petrinovich. (1989). Song development and territory

establishment in Nuttall's white-crowned sparrows. *The Condor 91*:397-407.

Dunn, J. L., K. L. Garrett and J. K. Alderfer (1995). White-crowned Sparrow subspecies:

identification and distribution. Birding 27:182-200.

Evans, C. S., L. Evans, and P. Marler (1993). On the meaning of alarm calls: functional reference

in an avian vocal system. Animal Behaviour 46:23-38.

Emlen, S. T. (1972). An experimental analysis of the parameters of bird song eliciting species

recognition. *Behaviour* 41:130-171.

Fick, S.E., and R.J. Hijmans (2017). WorldClim 2: new 1 km spatial resolution climate surfaces

for global land areas. International Journal of Climatology 37:4302-4315.

Fitch, W. T. (2006). The biology and evolution of music: A comparative perspective. Cognition

100:173-215. https://doi.org/10.1016/j.cognition.2005.11.009

Freeberg, T. M., A. P. King, and M. J. West (1995). Social Malleability in Cowbirds (*Molothrus*

ater artemisiae): Species and Mate Recognition in the First 2 Years of Life. Journal of

Comparative Psychology 109:357-367. https://doi.org/10.1037/0735-7036.109.4.357

Freeman, B. G., J. Rolland, G. A. Montgomery, and D. Schluter (2022). Faster evolution of a

premating reproductive barrier is not associated with faster speciation rates in New World

passerine birds. Proceedings of the Royal Society B: Biological Sciences 289:20211514.

https://doi.org/10.1098/rspb.2021.1514

GBIF.org (28 February 2023) GBIF Occurrence Download https://doi.org/10.15468/dl.vg5xnc

Gil, D., and M. Gahr (2002). The honesty of bird song: Multiple constraints for multiple traits. In

Trends in Ecology and Evolution 17:133-141. https://doi.org/10.1016/S0169-5347(02)02410-

2

Grinnell, J., and A. H. Miller (1944). The distribution of the birds of California. Pacific Coast

Avifauna 27:523-524. Cooper Ornithological Club, Berkeley, CA, USA.

Hartshorne, C. (1956). The monotony-threshold in singing birds. The Auk 73:176-192.

Heinemann, D. (1981). Song Dialects, Migration, and Population Structure of Puget Sound

White-Crowned Sparrows. The Auk 98:512-521.

Hunn, E. S., and D. Beaudette. (2014). Apparent sympatry of two subspecies of the White-

crowned Sparrow, *Zonotrichia leucophrys pugetensis* and *gambelii*. *Washington State West*

Birds 45:132-140.

Hunter, M. L., and J. R. Krebs (1979). Geographical variation in the song of the great tit (Parus

major) in relation to ecological factors. The Journal of Animal Ecology 48:759-785.

K. Lisa Yang Center for Conservation Bioacoustics at the Cornell Lab of Ornithology. (2019).

Raven Pro: Interactive Sound Analysis Software (Version 1.6.1) [Computer software]. The

Cornell Lab of Ornithology, Ithaca, NY, USA. Available from

https://ravensoundsoftware.com/.

Kondo, N., and S. Watanabe (2009). Contact calls: Information and social function. In Japanese

Psychological Research 51:197-208. https://doi.org/10.1111/j.1468-5884.2009.00399.x

Kopuchian, C., D. Alejandro Lijtmaer,P. Luis Tubaro, and P. Handford (2004). Temporal

stability and change in a microgeographical pattern of song variation in the rufous-collared

sparrow. Animal Behaviour 68:551-559. https://doi.org/10.1016/j.anbehav.2003.10.025

Kroodsma, D., and J. Verner (1978). Complex Singing Behaviors among *Cistothorus* Wrens.

The Auk 95:703-716.

Kroodsma, D. E., M. C. Bake, L. F. Baptista, and L. Petrinovich (1985). Vocal "Dialects" in

Nuttall's White-Crowned Sparrow. Current Ornithology 2:103-133.

https://doi.org/10.1007/978-1-4613-2385-3_3

Kroodsma, D. E., and B. E. Byers. (1991). The function (s) of bird song. *American Zoologist*

31:318-328.

Kroodsma, D. E., P. W. Houlihan, P. A. Falleon, and J. A. Wells (1997). Song development by

grey catbirds. Animal Behaviour 54:457-464. https://doi.org/10.1006/anbe.1996.0387

Lein, M. R., and K. W. Corbin (1990). Song and plumage phenotypes in a contact zone between

subspecies of the White-crowned Sparrow (*Zonotrichia leucophrys*). Canadian Journal of

Zoology 68:2625-2629.

Lewis, R. N., M. Soma, S. R. de Kort, and R. T. Gilman (2021). Like Father Like Son: Cultural

and Genetic Contributions to Song Inheritance in an Estrildid Finch. Frontiers in Psychology

12:654198. https://doi.org/10.3389/fpsyg.2021.654198

Lipshutz, S. E., I. A. Overcast, M. J. Hickerson, R. T. Brumfield, and E. P. Derryberry (2017).

Behavioural response to song and genetic divergence in two subspecies of white-crowned

sparrows (*Zonotrichia leucophrys*). Molecular Ecology 26:3011-3027.

https://doi.org/10.1111/mec.14002

Lisovski, S., Z. Németh, J. C. Wingfield, J. S. Krause, K. . Hobson, N. E. Seavy, J. Gee, and M.

Ramenofsky. (2019). Migration pattern of Gambel's white-crowned sparrow along the

Pacific flyway. *Journal of Ornithology 160*:1097-1107.

Luther, D. A., J. Phillips, and E. P. Derryberry. (2016). Not so sexy in the city: urban birds adjust

songs to noise but compromise vocal performance. *Behavioral Ecology* 27:332-340.

Ligges, U., S. Krey, O. Mersmann, and S. Schnackenberg (2018). tuneR: Analysis of Music and

Speech. https://CRAN.R-project.org/package=tuneR.

Mahler, P. (1970). A comparative approach to vocal learning: song development in White-

crowned Sparrows. Journal of Comparative and Physiological Psychology 71:1-25.

Marler, P. (2004). Bird calls: their potential for behavioral neurobiology. Annals of the New

York Academy of Sciences 1016:31-44.

Marler, P., and M. Tamura (1962). Song "dialects" in three populations of White-crowned

Sparrows. The Condor 64:368-377. https://www.jstor.org/stable/1365545

Marler, P., and M. Tamura (1964). Culturally transmitted patterns of vocal behavior in sparrows.

Science, 146:1483-1486.

Marler, P., P. Mundinger, M. S. Waser, and A. N. N. Lutjen (1972). Effects of acoustical

stimulation and deprivation on song development in red-winged blackbirds (*Agelaius*

phoeniceus). Animal Behaviour 20:586-606.

Mason, N. A., K. J. Burns, J. A. Tobias, S. Claramunt, N. Seddon, and E. P. Derryberry (2017).

Song evolution, speciation, and vocal learning in passerine birds. Evolution 71:786-796.

https://doi.org/10.1111/evo.13159

Morton, E. S. (1975). Ecological sources of selection on avian sounds. *The American Naturalist*

109:17-34.

Morton, E. S. (1982). Grading, discreteness, redundancy, and motivation-structural rules.

Acoustic communication in birds. Academic Press, New York.

Morton, E. S. (1986). Predictions from the ranging hypothesis for the evolution of long distance

signals in birds. *Behaviour* 99:65-86.

Naguib, M. (2003). Reverberation of rapid and slow trills: implications for signal adaptations to

long-range communication. The Journal of the Acoustical Society of America, 113:1749-

1756.

Nelson, D. A. (2000). Song overproduction, selective attrition and song dialects in the white-

crowned sparrow. Animal Behaviour 60:887–898. https://doi.org/10.1006/anbe.2000.1560

Nelson, D. A., and A. Poesel (2013). Song sharing correlates with lifetime social pairing success

but not territory tenure in the Puget Sound white-crowned sparrow. *Behavioral Ecology and*

Sociobiology 67:993-1000.

Nelson, D. A., and A. Poesel. (2007). Segregation of information in a complex acoustic signal:

individual and dialect identity in white-crowned sparrow song. *Animal behaviour*, 74:1073-

1084.

Nelson, D. A., K. I. Hallberg, and J. A. Soha (2004). Cultural evolution of Puget Sound white-

crowned sparrow song dialects. Ethology 110:879-908. https://doi.org/10.1111/j.1439-

0310.2004.01025.x

Nice, M. M., and B. D. Blanchard (1942). The White-Crowned Sparrows (*Zonotrichia*

leucophrys) of the Pacific Seaboard: Environment and Annual Cycle. Bird-Banding 13:96-

97. https://doi.org/10.2307/4509737

Nowicki, S., S. Peters, and J. Podos (1998). Song learning, early nutrition and sexual selection

in. American Zoologist 38:179-190. https://doi.org/10.1093/icb/38.1.179

Nowicki, S., M. Westneat, and W. Hoese (1992). Birdsong: motor function and the evolution of

communication. Seminars in Neuroscience 4:385-390.

Otter, K. A., A. Mckenna, S. E. LaZerte, and S. M. Ramsay (2020). Continent-wide Shifts in

Song Dialects of White-Throated Sparrows. Current Biology 30:3231-3235.

https://doi.org/10.1016/j.cub.2020.05.084

Odom, K. J., M. L. Hall, K. Riebel, K. E. Omland, and N. E. Langmore (2014). Female song is

widespread and ancestral in songbirds. *Nature communications* 5:3379.

Päckert, M. (2018). Song: The Learned Language of Three Major Bird Clades. In Bird

Species (D. T. Tietze, Editor). Springer, Cham. https://doi.org/10.1007/978-3-319-91689-7_5

Patten, M. A., G. McCaskie, and P. Unitt (2003). Birds of the Salton Sea: Status, biogeography,

and ecology. University of California Press, Berkeley, CA, USA.

Pelletier, T. A., and B. C. Carstens. (2018). Geographical range size and latitude predict

population genetic structure in a global survey. *Biology Letters*, 14:20170566.

Petrinovich, L., and T. L. Patterson (1981). The response of White-crowned Sparrows to songs

of different dialects and subspecies. Zeitschrift für Tierpsychologie 57:1-14.

Podos, J., and S. Nowicki (2004a). Performance limits on birdsong. Nature's Music 318-342.

Podos, J., and S. Nowicki (2004b). Beaks, adaptation, and vocal evolution in Darwin's finches.

Bioscience 54:501-510.

Podos, J., and P. S. Warren (2007). The Evolution of Geographic Variation in Birdsong. In

Advances in the Study of Behavior 37:403-458. https://doi.org/10.1016/S0065-

3454(07)37009-5

Provost, K. L., J. Yang, and B. C. Carstens (2022). The impacts of transfer learning,

phylogenetic distance, and sample size on big-data bioacoustics. PLoS ONE, 17:e0278522.

https://doi.org/10.1371/journal.pone.0278522

Provost, K. L., E. A. Myers, and B. T. Smith (2021). Community phylogeographic patterns

reveal how a barrier filters and structures taxa in North American warm deserts. Journal of

Biogeography 48:1267-1283.

Robinson, F. N. (1975). Vocal Mimicry and the Evolution of Bird Song. Emu 75: 23–

27. doi:10.1071/mu9750023

Riebel, K., K. J. Odom, N. E. Langmore, and M. L. Hall. (2019). New insights from female bird

song: towards an integrated approach to studying male and female communication roles.

Biology letters 15:20190059.

Rothstein, S. I., and R. C. Fleischer (1987). Vocal dialects and their possible relation to honest

status signalling in the Brown-headed Cowbird. The Condor 89:1-23.

R Core Team. (2020). R: a language and environment for statistical computing, version 4.0.3. R

Foundation for Statistical Computing.

Schrider, D. R., and A. D. Kern. (2017). Soft sweeps are the dominant mode of adaptation in the

human genome. *Molecular biology and evolution* 34:1863-1877.

Smith, G. T., E. A. Brenowitz, and J. C. Wingfield. (1997). Roles of photoperiod and

testosterone in seasonal plasticity of the avian song control system. *Journal of neurobiology*

32:426-442.

Somveille, M., A. S. Rodrigues, and A. Manica. (2015). Why do birds migrate? A

macroecological perspective. *Global Ecology and Biogeography 24*:664-674.

Sueur, J., T. Aubin, and C. Simonis. (2008) Seewave, a free modular tool for sound analysis and

synthesis. *Bioacoustics* 18:213-226.

Sullivan, B. L., C. L. Wood, M. J. Iliff, R. E. Bonney, D. Fink, and S. Kelling. (2009). eBird: A

citizen-based bird observation network in the biological sciences. *Biological conservation*,

142:2282-2292.

Searcy, W. A., and M. Andersson (1986). Sexual selection and the evolution of song. Annual

Review of Ecology and Systematics 17:507-533.

https://doi.org/10.1146/annurev.es.17.110186.002451

Seddon, N. (2005). Ecological adaptation and species recognition drives vocal evolution in

neotropical suboscine birds. Evolution 59:200-215. https://doi.org/10.1111/j.0014-

3820.2005.tb00906.x

Slabbekoorn, H., and T. B. Smith (2002). Bird song, ecology and speciation. Philosophical

Transactions of the Royal Society B: Biological Sciences 357:493–503.

https://doi.org/10.1098/rstb.2001.1056

Slater, P. J. B. (1986). The cultural transmission of bird song. Trends in Ecology and Evolution,

1:94-97. https://doi.org/10.1016/0169-5347(86)90032-7

Slater, P. J. B. (1989). Bird song learning: causes and consequences. Ethology Ecology and

Evolution 1:19-46.

Swiston, K. A., and D. J. Mennill (2009) Comparison of manual and automated methods for

identifying target sounds in audio recordings of Pileated, Pale-billed, and putative Ivory-

billed woodpeckers. Journal of Field Ornithology 80: 42-50.

Thompson, Jr., A. D., and M. C. Baker (1993). Song dialect recognition by male White-crowned

Sparrows: effects of manipulated song components. Condor 95:414-421.

Taylor, R. S., A. C. Bramwell, R. Clemente-Carvalho, N. A. Cairns, F. Bonier, K. Dares, and S.

C. Lougheed (2021). Cytonuclear discordance in the crowned-sparrows, *Zonotrichia*

atricapilla and *Zonotrichia leucophrys*. Molecular Phylogenetics and Evolution 162:107216.

Van Dongen, W. F., and R. A. Mulder (2006). Habitat density, song structure and dialects in the

Madagascar paradise flycatcher *Terpsiphone mutata*. Journal of Avian Biology 37:349-356.

Veiner, M., J. Morimoto, E. Leadbeater, and F. Manfredini. (2022). Machine learning models

identify gene predictors of waggle dance behaviour in honeybees. *Molecular Ecology*

Resources 22:2248-2261.

Wang I.J. (2020) Topographic path analysis for modeling dispersal and functional connectivity:

calculating topographic distances using the topoDistance R package. Methods in Ecology and

Evolution 11:265-272.

Webb, W. H., D. H. Brunton, J. D. Aguirre, D. B. Thomas, M. Valcu, and J. Dale. (2016).

Female song occurs in songbirds with more elaborate female coloration and reduced sexual

dichromatism. *Frontiers in Ecology and Evolution* 4:22.

Winker, K., and D. D. Gibson. (2018). Some broad-scale effects of recent and future climate

change among migratory birds in Beringia. *Trends and traditions: avifaunal change in*

Western North America, Stud. West. Birds, 2.

Weckstein, J. D., R. M. Zink, R. C. Blackwell-Rago, and D. A. Nelson (2001). Anomalous

variation in mitochondrial genomes of White-crowned (*Zonotrichia leucophrys*) and Golden-

crowned (*Z. atricapilla*) sparrows: Pseudogenes, hybridization, or incomplete lineage

sorting? The Auk 118:231-236.

Welke, C. A., B. Graham, R. R. Conover, J. W. Rivers, and T. M. Burg (2021). Habitat-linked

genetic structure for white-crowned sparrow (*Zonotrichia leucophrys*): Local factors shape

population genetic structure. Ecology and evolution 11:11700-11717.

Figure Captions

Figure 1. Spectrogram of (A) *Z. l. nuttalli* , BLB11552, and (B) *Z. l. pugetensis,* BLB07718,

songs with manually annotation in red boxes. The X axis shows the duration time, in seconds,

and the Y axis shows the frequency, in kilohertz. Higher frequencies represent higher pitch; and

darker color represents higher amplitude i.e., louder sound. Each red box represents one

annotated syllable, with the width representing the duration while the height indicates the

differences of maximum and minimum frequency i.e., bandwidth. Note that background sounds

are not annotated. Above certain notes are annotated the note type: W = whistle, B = buzz, S =

"special" syllable, and T = trill of simple notes.

Figure 2. The geographical range of *Zonotrichia leucophrys* recordings that were used in this study. The sedentary *Z. l. nuttalli* individuals are blue circles and the migratory *Z. l. pugetensis* individuals are red triangles. Individuals in western California and Utah were outliers (see main text). Map below shows species distribution in all year (purple) and non-breeding season (light blue) (BirdLife International and Handbook of the Birds of the World 2021; Welke et al. 2021, Weckstein et al. 2001, Grinnell and Miller 1944, Taylor et al 2021, Cortopassi and Mewaldt 1965). Note that this map includes distributions for other subspecies (*gambelii* and *oriantha*). Individuals are separated by Non-Breeding Season (September to April, left two panels, saturated colors) and Breeding Season (May to August, right two panels, dull colors).

Figure 3: Overall syllable classifications by type. Arrows begin at the true syllable and end at

the predicted syllable, with line thicknesses and color indicating the percentage of time those

assignments were made. Assignments under 10% are omitted for clarity. Syllable types include

buzzes (B), trills (T), whistles (W), "special" syllables (S), and zero syllables, i.e., background

noise (Z)

824 **Tables**

826 **Table 1**. The machine learning model showed a high accuracy in annotating the syllables of both *Z. l. pugetensis* and *Z. l. nuttalli* for

827 all data.

		Subspecies		
		Both	*Z. l. nuttalli*	*Z. l. pugetensis*
	Segment Error Rate	-6.93%	-11.43%	-1.06%
	Segment Error Rate (Abs)	6.93%	11.43%	1.06%
Sound/Background	Accuracy	91.78%	92.95%	90.67%
	Precision	97.16%	98.10%	96.17%
	Recall	84.61%	86.84%	82.35%
	F-Score	90.45%	92.12%	88.72%
Syllable Type	Accuracy	87.43%	87.97%	86.92%
	Precision	86.20%	87.44%	84.71%
	Recall	76.34%	78.93%	74.65%
	F-Score	80.97%	82.97%	79.36%
Syllable Type (Silence Excluded)	Accuracy	75.15%	76.36%	73.91%
	Precision	85.71%	87.00%	84.06%
	Recall	70.96%	74.40%	68.97%
	F-Score	77.64%	80.00%	75.78%

830 **Table 2**: Principal component rotation of individual PCs axes for song across summary statistics of song properties. "BW" =

831 bandwidth. "Dur" = Duration. "Dom" = Dominant. "Freq"=Frequency.

	Principal component														
	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15
BW	0.285	0.010	0.215	-0.261	0.126	-0.066	0.119	-0.694	0.298	-0.429	0.062	-0.058	0.073	0.019	-0.001
Time	-0.227	-0.452	0.237	-0.055	0.051	0.392	0.044	-0.048	0.054	0.074	0.101	0.108	-0.178	0.677	0.064
Center Freq	-0.281	0.270	-0.152	0.323	-0.050	0.112	0.183	-0.105	-0.012	-0.407	-0.097	0.680	-0.153	-0.002	-0.016
Inflections	-0.372	0.012	0.201	-0.179	-0.002	0.117	0.058	0.117	0.195	0.039	0.190	0.185	0.794	-0.129	-0.044
Slope	-0.124	0.059	0.267	-0.594	-0.036	-0.181	-0.331	0.044	-0.547	-0.169	0.019	0.250	-0.142	-0.028	-0.008
Dur Intro	-0.244	-0.213	0.153	0.103	-0.081	-0.471	0.692	-0.080	-0.321	0.029	0.156	-0.130	-0.030	-0.031	0.017
Syllable Dur	-0.295	-0.384	0.203	0.014	-0.016	0.315	-0.062	-0.123	0.121	0.017	-0.039	-0.055	-0.325	-0.693	0.001
Whistle Dom Freq (First)	-0.399	0.205	-0.047	0.012	-0.022	-0.008	-0.110	0.041	0.088	-0.273	0.130	-0.414	-0.131	0.137	-0.689
Whistle Dom Freq (All)	-0.399	0.195	-0.048	0.033	0.023	-0.087	-0.161	0.066	0.111	-0.306	0.071	-0.365	-0.049	0.072	0.714
Max Dom Freq	-0.295	0.075	-0.091	0.256	0.006	-0.151	-0.367	-0.644	-0.190	0.455	-0.007	0.025	0.116	0.063	-0.005
Min Dom Freq	-0.119	0.300	-0.033	-0.179	0.837	0.154	0.249	0.007	-0.067	0.227	-0.059	-0.020	-0.116	-0.049	0.004
Trill BW	-0.229	0.079	-0.199	-0.466	-0.245	-0.288	0.138	-0.009	0.516	0.321	-0.267	0.153	-0.234	0.061	0.004
Trill Dur	0.020	-0.272	-0.607	-0.161	0.130	-0.074	-0.073	-0.023	0.030	-0.048	0.683	0.149	-0.065	-0.081	0.011
Trill Rate	0.112	0.317	0.517	0.199	-0.009	-0.178	-0.101	0.119	0.274	0.225	0.542	0.171	-0.273	-0.023	0.017
Whistle Dur	-0.055	-0.412	0.096	0.217	0.437	-0.536	-0.292	0.177	0.223	-0.178	-0.239	0.160	0.055	0.034	-0.089

833 **Table 3**: Principal component importance and eigenvalues of individual PCs axes for song across summary statistics of song

834 properties. Broken stick refers to expected proportion of variance explained under the broken stick model.

PC	Standard deviation	Proportion of Variance	Cumulative Proportion	Broken Stick
1	2.226	0.330	0.330	0.221
2	1.488	0.148	0.478	0.155
3	1.275	0.108	0.586	0.121
4	1.232	0.101	0.687	0.099
5	0.919	0.056	0.744	0.082
6	0.887	0.052	0.796	0.069
7	0.801	0.043	0.839	0.058
8	0.749	0.037	0.876	0.048
9	0.663	0.029	0.906	0.040
10	0.648	0.028	0.934	0.033
11	0.592	0.023	0.957	0.026
12	0.538	0.019	0.976	0.020
13	0.466	0.015	0.991	0.014
14	0.341	0.008	0.999	0.009
15	0.145	0.001	1.000	0.004

837 **Table 4**: Differences between season (breeding vs non-breeding) and subspecies (*Z. l. pugetensis* vs *Z. l. nuttalli*) using ANOVAs

838 while controlling for both. Degrees of freedom = 1 for all comparisons. Note that unknown season individuals were removed. SSQ =

839 sum of squares. MSQ = mean sum of squares. F = F-statistic. P = p-value. For these analyses, significant values have p<0.0031. We

840 indicate a near-significant p-value with bold text and a significant value with bold text plus an asterisk.

	Mean values				Model parameters: season				Model parameters: subspecies			
	breeding	non-breeding	*nuttalli*	*pugetensis*	SSQ	MSQ	F	P	SSQ	MSQ	F	P
PC1	0.255	0.01	0.874	-0.59	29	29.3	6.57	**0.01**	1037	1036.7	232.55	**0***
BW	2360	2390	2380	2360	585200	585171	0.679	0.41	320900	320875	0.372	0.542
Center Freq	3510	3440	3330	3630	2036000	2036430	3.083	0.0793	41240000	41238170	62.425	**0***
Inflections	8.63	9.36	7.36	10.5	268	268	11.1	**0***	4672	4672	193.3	**0***
Dur Intro	1.28	1.27	1.16	1.41	0.01	0.008	0.05	0.824	22.91	22.909	135.53	**0***
Syllable Dur	0.187	0.189	0.18	0.197	0.002	0.0022	0.606	0.436	0.138	0.13778	37.884	**0***
Slope	-40.7	-30.3	-47.9	-24.5	51135	51135	23.95	**0***	253743	253743	118.85	**0***
Whistle Dom Freq (First)	1.86	2.11	1.49	2.49	31.1	31.1	24.58	**0***	444.4	444.4	351.43	**0***
Whistle Dom Freq (All)	2.06	2.28	1.73	2.61	25	25	21.31	**0***	343.6	343.6	293.16	**0***
Max Dom Freq	5.21	5.33	4.93	5.63	7	7.22	2.458	0.117	219	218.64	74.428	**0***
Min Dom Freq	0.184	0.138	0.169	0.159	0.9	0.9143	5.039	**0.024**	0	0.0409	0.225	0.6351
Time	0.21	0.21	0.208	0.212	0	0.000015	0.003	0.958	0.01	0.009564	1.801	0.18
Trill BW	0.0017	0.002	0.0015	0.0022	0.00003	0.00003	27.27	**0***	0.0001676	0.0001676	147.47	**0***

	0.244	0.228	0.264	0.206	0.11	0.1106	4.617	**0.031**	1.3	1.3015	54.319	**0***
Trill Dur	0.244	0.228	0.264	0.206	0.11	0.1106	4.617	**0.031**	1.3	1.3015	54.319	**0***
Trill Rate	10.2	10.2	10	10.5	0	0.11	0.011	0.915	76	75.99	7.642	**0.006**
Whistle Dur	0.562	0.552	0.602	0.51	0.06	0.06	1.559	0.212	3.8	3.803	99.624	**0***

843 **Table 5**: MMRR analysis shows that divergence in songs of *Z. l. pugetensis* and *Z. l. nuttalli* is strongly correlated with environmental

844 differences before accounting for time. For these analyses, significant values have $p<0.0056$. We indicate a near-significant p-value

845 with bold text and a significant value with bold text plus an asterisk.

Subspecies	Season	R²	F	Coefficients		T-values		P-Values		
				Climate	Geo.	Climate	Geo.	Climate	Geo.	Overall
Both	Summer	0.008	2384.98	0.064	0.038	45.799	27.115	**0.001***	**0.001***	**0.001***
	Winter	0.041	6869.57	0.047	0.166	18.546	66.263	**0.013**	**0.001***	**0.001***
	Year	0.015	14109.1	0.072	0.069	82.157	79.380	**0.001***	**0.001***	**0.001***
N	Summer	0.003	203.976	-0.006	0.054	-1.713	16.363	0.820	0.072	0.076
	Winter	0.004	172.497	0.082	-0.055	18.525	-12.538	**0.004***	0.098	**0.031**
	Year	0.001	222.164	0.024	0.009	12.827	4.745	0.197	0.695	0.157
P	Summer	0.001	41.754	0.024	-0.003	8.923	-0.950	0.127	0.884	0.298
	Winter	0.075	3114.39	0.014	0.264	2.832	53.695	0.385	**0.001***	**0.001***
	Year	0.003	755.771	0.055	0.008	32.426	4.712	**0.001***	0.643	**0.001***

849   Table 6: MMRR analysis shows that divergence in songs of *Z. l. pugetensis* and *Z. l. nuttalli* is strongly correlated with environmental

850   differences even after accounting for time. For these analyses, significant values have p<0.0056. We indicate a near-significant p-

851   value with bold text and a significant value with bold text plus an asterisk.

Subspecies	Season	R²	F	Coefficients			T-values			P-Values			
				Time	Climate	Geo.	Time	Climate	Geo.	Time	Climate	Geo.	Overall
Both	Breeding	0.114	26395	0.327	0.039	0.031	271.742	29.514	23.202	**0.001***	**0.001***	**0.001***	**0.001***
	Non-breeding	0.052	5817	0.110	0.042	0.129	59.679	16.783	50.211	**0.001***	**0.016**	**0.001***	**0.001***
	Year	0.073	47858	0.244	0.048	0.046	337.048	56.164	54.598	**0.001***	**0.001***	**0.001***	**0.001***
N	Breeding	0.050	2764.570	0.223	-0.070	0.077	88.688	-21.033	23.624	**0.001***	**0.009**	**0.007**	**0.001***
	Non-breeding	0.010	269.207	0.080	0.048	-0.047	21.464	10.278	-10.738	**0.001***	0.092	0.148	**0.007**
	Year	0.022	3463.662	0.149	-0.024	0.027	99.686	-12.232	14.219	**0.001***	0.217	0.231	**0.001***
P	Breeding	0.133	7589.539	0.364	0.013	-0.010	150.574	5.101	-3.790	**0.001***	0.364	0.575	**0.001***
	Non-breeding	0.076	2097.494	0.037	0.012	0.240	7.680	2.473	40.915	0.342	0.447	**0.001***	**0.001***
	Year	0.088	14167.874	0.293	0.025	-0.002	202.118	15.334	-1.236	**0.001***	**0.010**	0.899	**0.001***

6