Article title: Biodiversity shows unique responses to land-use change across regional

biomes

Authors:

Peggy Bevan[1,2], Guilherme Braga Ferreira[1], Daniel J Ingram[3], Marcus Rowcliffe[2], Lucy

Young[4], Robin Freeman[2], Kate E. Jones[1].

Affiliations:

1.   Centre for Biodiversity and Environment Research, University College London,

Gower

Street, WC1E 6BT

2.   Institute of Zoology, Zoological Society of London, Wellcome Building, Outer Circle,

London NW8 7LS

3.   Durrell Institute of Conservation and Ecology, School of Anthropology and

Conservation,

University of Kent, Canterbury, CT2 7NR, UK.

4.   WWF UK, Living Planet Centre, Brewery Road, Woking, GU21 4LL

Corresponding author: Peggy Bevan, peggy.bevan.17@ucl.ac.uk

Data availability:

All data used in this manuscript is downloaded from open source locations. The PREDICTS

database can be downloaded from the Natural History Museum Data portal

(https://data.nhm.ac.uk/dataset/the-2016-release-of-the-predicts-database).

Geographical data used to delineate regional biome boundaries in this manuscript can be

downloaded from The Nature Conservancy's Geospatial Conservation Atlas

(https://geospatial.tnc.org/datasets/b1636d640ede4d6ca8f5e369f2dc368b/about).

All R code used to run analysis presented in this manuscript can be found at

https://github.com/PeggyBevan/RegionalBiomes

<u>Abstract</u>

Biogeography has a critical influence on how ecological communities respond to threats and

how effective conservation interventions are designed. For example, the resilience of

ecological communities is linked to environmental and climatic features, and the nature of

threats impacting ecosystems also varies geographically. Understanding community-level

threat responses may be most accurate at fine spatial scales, however collecting detailed

ecological data at such a high resolution would be prohibitively resource intensive. In this

study, we aim to find the spatial scale that could best capture variation in community-level

threat responses whilst keeping data collection requirements feasible. Using a database of

biodiversity records with extensive global coverage, we modelled species richness and total

abundance (the responses) across land-use types (reflecting threats), considering three

different spatial scales: biomes, biogeographical realms, and *regional biomes* (the

interaction between realm and biome). We then modelled data from three highly sampled

biomes separately to ask how responses to threat differ between regional biomes and

taxonomic group. We found strong support for regional biomes in explaining variation in

species richness and total abundance compared to biomes or realms alone. Our biome case

studies demonstrate that there is a high variation in magnitude and direction of threat

responses across both regional biomes and taxonomic group, but all groups in tropical forest

showed a consistently negative response, whilst many taxon-regional biome groups showed

no clear response to threat in temperate forest and tropical grassland. Our results suggest

that the taxon-regional biome unit has potential as a reasonable spatial and ecological scale

for understanding how ecological communities respond to threats and designing effective

conservation interventions to bend the curve on biodiversity loss.

Introduction

<u>Introduction</u>

Despite multiple internationally agreed science-based conservation targets, biodiversity

continues to decline across the globe, with the consequent loss of essential ecosystem

services (Tittensor *et al.*, 2014; Brooks *et al.*, 2015; Ceballos, Ehrlich and Dirzo, 2017; Isbell

et al., 2017; Mace *et al.*, 2018; IPBES, 2019; Ingram *et al.*, 2021). Habitat disturbance

caused by land-use change is a major factor contributing to biodiversity loss (Newbold *et al.*,

2015). However, the impact of disturbance on biodiversity (its response) varies spatially and

is contingent on factors such as the intensity of environmental change (Felipe-Lucia *et al.*,

2020), the original habitat (Monsarrat, Jarvie and Svenning, 2019) and taxon studied. Such

threat-response relationships will further be influenced by specific life-history traits,

biogeographic context, and the ecological scale being measured (*e.g.*, a community,

population or individual) (Isaac and Cowlishaw, 2004; Murphy, 2021; Suraci *et al.*, 2021).

Given the recognised variability of biodiversity's responses to threats such as habitat

disturbance, this knowledge can help improve conservation decisions. With the adoption of

new targets such as the Kunming-Montreal Global Biodiversity Framework (CBD, 2022) and

the Global Deal for Nature (Dinerstein *et al.*, 2019), it is vital to develop biodiversity

monitoring systems that can accurately measure indicators at an appropriate spatial and

ecological scale that optimises available resources with the resolution necessary to inform

successful conservation actions.

The specific location of a species or ecological community on the planet, known as its

biogeography, plays a crucial role in determining its sensitivity to an anthropogenic threat.

This could be down to the current climatic circumstances of the location, historical natural

disturbances, or the history of human activity in the area. For example, studies on

biodiversity in tropical areas have been shown to have higher levels of fragmentation-

sensitive species, stronger negative responses to land-use change and faster declining

abundance than comparable temperate areas (Betts *et al.*, 2019; Newbold *et al.*, 2020;

WWF, 2022). This could be explained by the extinction filter hypothesis, which suggests that

ecosystems with a history of disturbance will be more resilient in the present day due to the

previous filtering of sensitive species (Balmford, 1996). Tropical biomes are considered to

have a history of low natural disturbance and a stable climate, whereas temperate biomes

have been subject to elevated levels of glaciation, widespread fires, and human-caused

forest loss in the last 10,000 years (Betts *et al.*, 2019). In addition, traits typically associated

with tropical species (Stevens, 1989; Gaston, 2000; Brown, 2014), including habitat

specialisation and small range size, have also been associated with stronger negative

responses to land-use change (Newbold *et al.*, 2013). The history of land use and

environmental disturbance is not uniform across the globe and will further contribute to

modern day responses to anthropogenic threats (Goldewijk *et al.*, 2011; Yang *et al.*, 2021).

Incorporating an understanding of biogeography and historical knowledge of an environment

could play a key role in global biodiversity monitoring.

Considering the influence of biogeography on threat-response relationships, monitoring

biodiversity at the smallest possible scale would be appropriate. However, there are

currently more than 42,000 threatened species on the IUCN red list (IUCN, 2022), all of

which cannot have unique conservation plans created for them. Therefore, indicators must

be used at a higher ecological and spatial scale that strikes a balance between presenting

global trends and understanding local scale responses to human pressure (Ingram *et al.*,

2021). One way to account for biogeographic variation in threat responses is to use a

monitoring framework based on spatial units, but choosing the correct unit is challenging.

For example, creating national statistics for biodiversity is helpful for feeding into policy, but

biodiversity is not nationally constrained (Murphy, 2021). A global framework has been

suggested for monitoring biodiversity at an ecoregional level (Dinerstein *et al.*, 2017, 2019;

Smith *et al.*, 2018). This framework is powerful when measuring variables that can be

obtained using remote sensing or global species lists (*e.g.*, Dinerstein *et al.*, 2017; Smith *et*

al., 2020), but many ecoregions are lacking the required field data or access to study

population responses to human pressure at this resolution (Stephenson *et al.*, 2015). It is

essential to investigate whether broader spatial scales, already used in many global

monitoring studies (Newbold *et al.*, 2016; Blowes *et al.*, 2019; WWF, 2022), can effectively

encompass the diversity in threat-response relationships.

The broadest spatial unit of terrestrial habitats is the climatically defined biome, of which

there are 14 (Olson *et al.*, 2001). Responses of animal and plant populations to threats do

differ between biomes (Greenville *et al.*, 2018; Green *et al.*, 2020) and species richness is

particularly sensitive to land use change in tropical forest, tropical grassland and

Mediterranean biomes compared to temperate and desert biomes (Newbold *et al.*, 2020). In

addition to biomes, there are 8 biogeographic realms, which broadly follow the continents

(Olson *et al.*, 2001; UNEP-WCMC 2004). Data from the Living Planet Database (Collen *et*

al., 2009; WWF, 2022) has shown that decreases in vertebrate populations are more

pronounced in southern hemisphere realms, particularly the Neotropics (Green *et al.*, 2020;

WWF, 2022). Although there is evidence for variation in threat-response relationships

between biomes and between realms, the intersection between these two spatial units is

rarely investigated, which may mean a large amount of variation is unaccounted for when

monitoring biodiversity in biomes or realms alone.

A potential, intermediate spatial framework for monitoring biodiversity are regional biomes

(the interaction of realms and biomes) (n = 64) (Ingram *et al.*, 2021). Regional biomes each

cover 11 ecoregions on average, ranging from 1-81 ecoregions (Olson and Dinerstein,

2002). Separating biomes by biogeographic realms can account for differences in

evolutionary history, vegetative structure, threats, and socioeconomic status that occur

between realms (Moncrieff, Bond and Higgins, 2016; Allan *et al.*, 2019). Furthermore,

threats are not spread evenly across the world and can be region-specific (Lewis, Edwards

and Galbraith, 2015; Bowler *et al.*, 2020); for example, tropical forest in southeast Asia is a

threat 'hotspot', whereas large parts of the Amazon rainforest act as threat 'refugia' (Allan *et*

al., 2019). Some areas of forest biomes, particularly in the northern hemisphere, are being

afforested, whilst others experience mass deforestation (Song *et al.*, 2018). In Asian tropical

forests, species richness was found to be significantly more sensitive to disturbance

compared to similar regions in South America and Africa (Gibson *et al.*, 2011; Phillips,

Newbold and Purvis, 2017). Conversely, vertebrate abundance is declining more in the

Neotropics and Afrotropics than in the Indo-Pacific region (Green *et al.*, 2020; WWF, 2022).

These conflicting trends highlight the necessity of understanding and clearly reporting how

biogeographical variation contributes to threat responses. Regional biomes may present a

middle-ground between the finer scale monitoring of ecoregions and the coarser biome

scale, facilitating robust reporting of biodiversity responses.

Here, we use the PREDICTS database to analyse data from over 400 field studies to

explore the efficacy of regional biomes as a spatial monitoring unit for terrestrial species.

Our expectation is that there will be negative responses to human-dominated land-use types

compared to primary vegetation, but the magnitude of response will differ between regional

biomes. We therefore hypothesise that regional biomes present a more parsimonious

explanation of responses of biodiversity to land-use change than realms or biomes alone.

Secondly, as case studies we model the three most data-rich biomes (Tropical Forest,

Temperate Forest, Tropical Grasslands) in the PREDICTS database separately to explore

how responses to land-use change and use-intensity vary between regional biomes. Here,

we also consider the impact of taxonomic group (Vertebrate, Invertebrate, Plant) on the

threat-response relationship within regional biomes. We expect the highest variation but also

uncertainty in predicted responses between tropical regional biomes due to ecoregions

being more distinct in these biomes (Smith *et al.*, 2020), and smaller and less variable

responses between temperate regional biomes due to the history of natural and

anthropogenic disturbance leading to biotic homogenization (Newbold *et al.*, 2018).

Methods

The PREDICTS database

We used species abundance and occupancy data from the PREDICTS database (Hudson et

al 2017; 22678 sites, 480 sources, 666 studies, 47,000 species). We chose to use the

PREDICTS database due to high global coverage compared to other databases (Hudson *et*

al., 2017), making it suitable for a study comparing many regional biomes. The PREDICTS

project (Predicting Responses of Ecological Diversity In Changing Terrestrial Systems;

www.predicts.co.uk) is a database that collates biodiversity studies with comparable

measures of terrestrial biodiversity from sites of different land uses and land-use intensity

(Hudson *et al.*, 2017). All sampled sites are classified by stage of disturbance or recovery

which can be used as a substitute for actual temporal change at a single site (Walker *et al.*,

2010; Srivathsa *et al.*, 2018). This space-for-time substitution methods allows for specific

measurements of biodiversity over distinct land-use types, acting as a proxy for

anthropogenic pressure. PREDICTS does not include marine or freshwater data but does

have extensive global coverage and has made attempts to reduce taxonomic and

geographic bias (Hudson *et al.*, 2017).

We summarised occurrence and abundance records by site and taxon using the

predictsFunctions R package (Newbold, 2018). 'Taxa' were categorised as Vertebrate,

Invertebrates, Plants or Fungi. The PREDICTS database covers 47 out of 64 regional

biomes (70%), with some regional biomes more comprehensively sampled than others

(Table S1). For analysis, we excluded data from three biomes with fewer than 50 sites in

total from the dataset (Flooded Grasslands and Savannahs, Tundra, Deserts & Xeric

Shrublands and Mangroves). Additionally, regional biomes were considered data deficient

and excluded from analysis if they had no sites in primary vegetation land-use types (our

reference level), or no sites in any kind of managed land use (cropland, pasture, and

plantation forest). This ensured that regional biomes included in analysis had data points

(sites) across a gradient of increasingly anthropogenic land-use types. After removing data

deficient regional biomes, a total of 30 regional biomes were available for analysis (Fig. 1;

20011 sites, 545 studies).

Using the predominant land use and use-intensity classifications from the PREDICTS

database (land use: primary vegetation, mature secondary vegetation, intermediate

secondary vegetation, young secondary vegetation, plantation forest, pasture, cropland and

urban; use intensity: minimal, light, intense) (Hudson *et al.*, 2017), we created two land-use

variables – *LandUse* and *UseIntensity*. We combined the three secondary vegetation

classes (mature, intermediate, and young) into one and removed data from urban land-use

types due to the low quantity of samples compared to other land-types (917 sites from 53

studies). Because some combinations of regional biome and *LandUse* class have low

sample sizes, we tested 5 variations of this variable where categories were aggregated to

find the most parsimonious grouping that still explained variation in the model well. For

example, to make the *LandUse2* variable we grouped primary vegetation and secondary

vegetation as 'natural vegetation'. Full descriptions of each land-use type and the 5

variations of *LandUse* can be found in Table S2. *UseIntensity* is a combination of

predominant land-use type and use-intensity. To create this variable, we grouped 'minimal'

and 'light' use-intensities into 'minimal' and combined with groups from *LandUse3* (defined

as primary vegetation, secondary vegetation, and agriculture) to give categories primary

vegetation, minimal secondary vegetation, intense secondary vegetation, minimal

agriculture, intense agriculture. Primary vegetation sites were not split by use intensity as

there is no difference in biodiversity metrics between primary vegetation sites with differing

use-intensities (Newbold *et al.*, 2015).

<u>Statistical Analysis</u>

We used generalised mixed-effects models (Bolker *et al.*, 2009) to test for differences in

species richness and total abundance in response to land use, biome, realm and regional

biome (an interaction of biome and realm). Species Richness was modelled using a Poisson

distribution and total abundance was log transformed (natural logarithm) and modelled with

a gaussian distribution. In all models, we included study and taxa as random intercepts to

account for site-level differences, data collection methods and scale differences between

response metrics of Vertebrates, Invertebrates, Plants and Fungi. We ran a series of models

with species richness and total abundance against each *LandUse* variable (i.e. *LandUse1,*

LandUse2, or LandUse3 – see the previous section) and selected the best fitting variable

using Akaike Information Criterion (AIC) (Burnham, Anderson and Huyvaert, 2011). We

found that *LandUse1* (Primary vegetation, Secondary vegetation, Agriculture, Plantation

Forest, Cropland) had the most support and was used for remaining analyses (Table S3).

<u>Global analysis of regional biomes</u>

To see if including regional biome is a more parsimonious explanation for changes in

biodiversity than realm or biome, we ran a model selection process using ΔAIC values to

assess the support for each variable. The fixed effect structures tested were *LandUse,*

LandUse:Realm, LandUse:Biome, and *LandUse:RegionalBiome* and species richness and

total abundance were used as response variables. We tested model robustness in two ways.

First, we increased the sample size threshold that would allow a regional biome to be

included in the model (minimum 1, 5, 25 or 50 sites in each regional biome-land use

combination) and checked if the model selection process gave the same results in each

case. Second, we ran a series of 100 hold-out models using the 25-site threshold dataset

where each iteration randomly removed 10% of studies to see if this would change the

results of model selection.

<u>Biome-specific case studies</u>

As case studies, we investigated the differences in biodiversity response to land-use change

between regional biomes by selecting the three biomes which had three or more regional

biomes with at least 100 sites in the PREDICTS database (Fig. S1). The biomes selected

were tropical and subtropical moist broadleaf forest (Tropical Forest) (3 realms; Indo-Malay:

1781 sites; Neotropics: 2274 sites; Afrotropics: 2081 sites), temperate broadleaf and mixed

forests (Temperate Forest) (4 realms; Palearctic: 3986 sites; Neotropics: 577 sites; Nearctic:

856 sites; Australasia: 336 sites) and tropical and subtropical savannahs, grasslands and

shrublands (Tropical Grassland) (3 realms; Afrotropic: 1663 sites; Neotropics: 190 sites;

Australasia: 502 sites). For each biome and biodiversity metric (species richness and total

abundance), we ran four models that used the *LandUse* or *UseIntensity* variables and tested

if including *Realm*, which in this case represents regional biomes, increases model support.

We then predicted the average change in species richness and total abundance, relative to

primary vegetation, in each land-use type in each realm using the StatisticalModels R

package (Newbold, 2015).

In addition to the biome case studies, we did a further analysis to investigate how taxonomic

group further influences the responses of species richness to land-use change across

regional biomes. For each biome case study, we ran one model using Species Richness as

the response variable and *LandUse*, *Realm* and *Taxa* (Vertebrate, Invertebrate and Plant)

as fixed interaction terms. As above, we predicted the average change in species richness

relative to primary vegetation.

Results

Global analysis of regional biomes

Including regional biome as a variable improved model fit and better explained the effect of

land-use change on species richness and total abundance (Fig. 2). Models with realm or

biome alone had lower support and this was consistent across 100 iterations of hold-out

models (Fig. 2). Increasing the sample size threshold for regional biomes did not change the

overall outcome (Table S4). Furthermore, marginal R^2 values showed that regional biome

explained more variation in both species richness and total abundance than biome or realm

interacting with land use (Table S4).

Biome-specific case studies

We found that species richness and total abundance in each regional biome of tropical forest

showed a distinct response to land-use change. Species richness reduced in human-

dominated land-use types compared to primary vegetation in all realms but to different

degrees at each land-use type (Fig. 3a). The largest loss of species richness was seen in

Afrotropical cropland (-53% reduction compared to primary vegetation), followed by Indo-

Malayan cropland (-50%) and plantation forest (-43%). Conversely, the Neotropics did not

show a significant species richness reduction in cropland and a barely significant reduction

in plantation forest (-20%), with species richness for this regional biome being worst affected

in pasture land-use types (-30%). When predicting responses of total abundance to land-use

change in the tropical forest biome there was much higher uncertainty. The negative

responses to land-use type seen in the species richness models did not always carry over to

the abundance models, for example there was no sign of a negative response in Neotropical

pasture. However, total abundance was significantly lower in Indo-Malay plantation forest

compared to other regional biomes (Fig. S2a). For the models including *UseIntensity*, a

negative response to increased disturbance can be observed in Indo-Malayan and

Afrotropical regional biomes, with declines >50% compared to primary vegetation in both

species richness and total abundance (Fig. 3b, Fig. S2b). This extreme response is not seen

in the Neotropical regional biome.

In temperate forest regional biomes, species richness and total abundance both showed low

to no response in human-dominated land-use types compared to primary vegetation (Fig.

4a, Fig. S2c). The exception is the Palearctic regional biome, where total abundance

reduced by 55% in cropland land-use types. There were also reductions in species richness

in plantation forest and cropland in this regional biome, but the upper confidence limits just

surpassing 0 (Fig. 4a). This was the only response from any temperate forest regional biome

that showed a significant change from primary vegetation (95% confidence intervals do not

include 0). Furthermore, predictions for change in species richness and total abundance

have high levels of uncertainty for all realms except palearctic.

Similar to temperate forest, biodiversity metrics in most tropical grassland regional biomes

showed little or no response to any land-use type when compared to primary vegetation

(Fig. 5). Predictions of species richness and total abundance change showed a high degree

of uncertainty across all regional biomes, except for the Afrotropics, where there was a

strong response to intense agriculture (-67%), although this was the only regional biome with

enough datapoints to predict in this category so cannot be compared (Fig. 5b, Fig. S2f).

Species richness in Australasia showed negative responses to pasture and minimally-used

agricultural land, but with the upper confidence interval just passing zero in both cases (Fig.

5a, b).

Including taxon as an interaction term reveals distinct responses to land-use change across

different taxonomic groups within regional biomes (Fig. 6). In tropical forest, there is still a

general trend of strong negative responses to human-dominated land-use types, but for

some regional biomes there is high variation in responses. For example, afrotropical

vertebrates have no observable response to cropland and plantation forest, but plants in the

same regional biome display strong reductions in species richness as these land-use types.

Conversely, in Neotropical tropical forest, Vertebrates and Invertebrates show a negative

response to pasture land, but plants do not. In temperate forest, the addition of taxon groups

highlights the level of incomplete sampling in the dataset but reveals strong responses of

vertebrates and plants to land-use change in the Palearctic regional biome that are missed

in the unified model (as seen in Fig. 4). In tropical grassland, we observe the opposite effect,

where there are no strong responses predicted to any land-use type and increased

uncertainty in predictions, due to much lower sampling effort.

Discussion

Global analysis of regional biomes

Our results show that including regional biome when modelling the global effects of land use

on biodiversity can increase explanatory power. There is a need for a biodiversity monitoring

spatial framework that finds the balance between having feasible data collection

requirements and adequate global coverage, at a realistic scale that captures true variation

in ecological resilience to disturbance (Ingram *et al.*, 2021). It has been previously shown

that biodiversity shows different responses to disturbance across biomes (Greenville *et al.*,

2018; Blowes *et al.*, 2019; Newbold *et al.*, 2020) and across realms (Gibson *et al.*, 2011;

Phillips, Newbold and Purvis, 2017; Lambertini, 2020), but for the first time, we have shown

the important interaction between biome and realm. The combination of these two spatial

delineations gives us more information about ecological processes. This result is important

for biodiversity monitoring and conservation interventions as not only are the magnitude of

biodiversity responses to anthropogenic pressure different between regions but that

threatening processes (i.e., land-use types) are affecting regional biomes differently.

Biome-specific case studies

Whilst our global models suggest that regional biome can modulate the effect of land-use

change on species richness and abundance on a global scale, a more focused analysis

found that the influence of regional biome is stronger in some biomes than others. Generally,

species richness and total abundance in tropical biomes showed negative responses to

disturbed land-use types, the degree of which changed with realm and taxonomic group. In

temperate forest, however, there was no significant response to land-use change overall,

excluding the palearctic realm, particularly vertebrates. The observed difference between

tropical and temperate regional biomes is expected; responses to land-use change are

stronger in tropical biomes compared to temperate biomes, especially tropical forest

(Newbold *et al.*, 2020). However, ecoregions in tropical biomes are considered more

ecologically distinct than neighbouring ecoregions compared to those in temperate biomes

(Smith *et al.*, 2018). As each regional biome is a group of many ecoregions, it would follow

that predicted responses to disturbance in tropical regional biomes would contain more

uncertainty than temperate, but the opposite is true for our case studies. The high

uncertainty in predicted responses produced by the temperate forest and tropical grassland

models can be partly explained by low sample sizes and the high variation in responses of

taxonomic groups. Predictions in palearctic temperate forest have very low uncertainty

compared to all other regional biomes and is also the most extensively sampled regional

biome in the PREDICTS database (Table S1).

For tropical forest regional biomes, our model predictions suggest Indo-Malayan biodiversity

responds most strongly to disturbance, followed by the Afrotropics. There was no significant

change in species richness in the Neotropics tropical forest in any land-use type except for

pasture. Adding taxon to the model showed that these responses are nuanced and driven by

different taxonomic groups, but the patterns remain largely the same. Biodiversity in Asian

tropical forest is repeatedly found to be extremely sensitive to land-use change and

disturbance (Sodhi *et al.*, 2009; Gibson *et al.*, 2011; Phillips, Newbold and Purvis, 2017).

The islands of south east Asia have a rich geographical history which has led to this region

having the highest number of endemic species and biodiversity hotspots (Myers *et al.*, 2000;

Sodhi *et al.*, 2004) compared to other tropical regions. This area also has the highest

proportion of species under threat in the world (Allan *et al.*, 2019). This high baseline for

species richness combined with high levels of specialisation may partly explain why the

response to land-use change is so pronounced in this regional biome. Response metrics in

Neotropical tropical forest have the smallest response to disturbed land-use types compared

to similar regional biomes. However, temporal trends from the Living Planet Index suggest

that populations of vertebrates in south America have decreased by 94% since 1970

(Lambertini, 2020). In addition, other studies in this region have shown a decline in

biodiversity due to land-use change (Bogoni, Peres and Ferraz, 2020; Gaona *et al.*, 2021;

Quintero *et al.*, 2023). The difference is a demonstration of the importance of testing multiple

data types and of assessing the regional biome framework using time-series biodiversity

data and across multiple threat gradients in the future. Further research is needed to

understand how biological and taxonomic differences between these tropical forest realms

might be impacting sensitivity to land-use change.

The intensity and history of habitat disturbance in each regional biome will also impact the

observed response of biodiversity and may even cause biases in our data set. There is a

strong focus on palm oil as a driver of biodiversity loss in Asian tropics, which may lead to a

publication bias in this area. Temperate forest biomes have a long history of human

habitation and extinction events, meaning the baseline in primary vegetation is much lower

in these regional biomes compared to less historically disturbed biomes (Monsarrat and

Svenning, 2022). The lack of variation in responses observed in temperate forest regional

biomes may well be because of this lower baseline in primary vegetation. Indeed, trends of

vertebrate abundance have been increasing in the northern hemisphere since 1970 (Leung,

Greenberg and Green, 2017) as well as forest cover (Song *et al.*, 2018). The drivers that

cause differences in sensitivity to habitat disturbance are related to both biogeographic and

human history, splitting the world into regional biomes can further account for these

differences.

Although regional biomes may be a valuable spatial monitoring unit, our models were limited

by sample size for some regional biomes. The PREDICTS database is the most extensive

collection of terrestrial biodiversity records available, but there are still gaps. Under-sampling

was pronounced in non-forested temperate biomes like Montane Grasslands and

Shrublands, and realms were unevenly sampled, for example studies from Palearctic

temperate forest dominate the database but there are no studies at all on Palearctic

temperate grasslands (Table S1). Furthermore, taxonomic under sampling was highlighted

in our model that included taxonomic group. For example, Fig. 3 suggests that afrotropical

tropical forests have a greater response to Cropland compared to the neotropical regional

biome. However, this response is driven by Plants in the Afrotropics and Invertebrates and

Vertebrates in the Neotropics, making the impact of cropland hard to compare. Prioritising

data collection in under-represented regional biomes and taxonomic groups highlighted here

would enhance progress towards an effective monitoring framework (Ingram *et al.*, 2021).

Furthermore, this study focuses only on terrestrial regional biomes, as these are more

extensively sampled and we were specifically testing the effect of land-use change on

species richness and abundance. We acknowledge that marine biodiversity is showing

stronger responses to disturbance than terrestrial (Blowes *et al.*, 2019), and suggest a

similar monitoring framework could be adapted for marine ecosystems.

Conclusion

Our results show that the taxa-regional biome unit has the potential to be a powerful

framework for monitoring biodiversity and implementing conservation action. Resources for

data collection and biodiversity monitoring on the ground are sparse, and the regional biome

framework implies that monitoring studies could be more evenly spread across regional

biomes and taxonomic groups, optimising resource allocation for data collection. Here we

have highlighted that there is within-biome variation in how biodiversity is responding to

disturbance. We show that the difference between realms may be more important in tropical

biomes than temperate, and that spatial resolution should be carefully considered in any

attempts at monitoring global biodiversity trends. Although more data are needed, regional

biomes have the potential to be a meaningful scale at which to prioritise monitoring. These

results will be beneficial for informing upcoming policy discussions such as the CBD's

COP16 on the best way to monitor progress towards biodiversity targets and effect change

on the ground.

References

Allan, J.R. *et al.* (2019) 'Hotspots of human impact on threatened terrestrial vertebrates',
PLOS Biology. Edited by P. Jordano, 17(3), p. e3000158. Available at:
https://doi.org/10.1371/journal.pbio.3000158.

Balmford, A. (1996) 'Extinction filters and current resilience: the significance of past
selection pressures for conservation biology', *Trends in Ecology & Evolution*, 11(5), pp.
193–196. Available at: https://doi.org/10.1016/0169-5347(96)10026-4.

Betts, M.G. *et al.* (2019) 'Extinction filters mediate the global effects of habitat
fragmentation on animals', *Science*, 366(6470), pp. 1236–1239. Available at:
https://doi.org/10.1126/science.aax9387.

Blowes, S.A. *et al.* (2019) 'The geography of biodiversity change in marine and terrestrial
assemblages', *Science*, 366(6463), pp. 339–345. Available at:
https://doi.org/10.1126/science.aaw1620.

Bogoni, J.A., Peres, C.A. and Ferraz, K.M.P.M.B. (2020) 'Extent, intensity and drivers of
mammal defaunation: a continental-scale analysis across the Neotropics', *Scientific Reports*,
10(1), p. 14750. Available at: https://doi.org/10.1038/s41598-020-72010-w.

Bolker, B.M. *et al.* (2009) 'Generalized linear mixed models: a practical guide for ecology
and evolution', *Trends in Ecology & Evolution*, 24(3), pp. 127–135. Available at:
https://doi.org/10.1016/j.tree.2008.10.008.

Bowler, D.E. *et al.* (2020) 'Mapping human pressures on biodiversity across the planet
uncovers anthropogenic threat complexes', *People and Nature*, 2(2), pp. 380–394. Available
at: https://doi.org/10.1002/pan3.10071.

Brooks, T.M. *et al.* (2015) 'Harnessing biodiversity and conservation knowledge products to
track the Aichi Targets and Sustainable Development Goals', *Biodiversity*, 16(2–3), pp. 157–
174. Available at: https://doi.org/10.1080/14888386.2015.1075903.

Brown, J.H. (2014) 'Why are there so many species in the tropics?', *Journal of
Biogeography*, 41(1), pp. 8–22. Available at: https://doi.org/10.1111/jbi.12228.

Burnham, K.P., Anderson, D.R. and Huyvaert, K.P. (2011) 'AIC model selection and
multimodel inference in behavioral ecology: some background, observations, and
comparisons', *Behavioral Ecology and Sociobiology*, 65(1), pp. 23–35. Available at:
https://doi.org/10.1007/s00265-010-1029-6.

CBD, (Convention on Biological Diversity) (2022) *15/4. Kunming-Montreal Global
Biodiversity Framework*. CBD/COP/DEC/15/4. Available at:
https://www.cbd.int/doc/decisions/cop-15/cop-15-dec-04-en.pdf.

Ceballos, G., Ehrlich, P.R. and Dirzo, R. (2017) 'Biological annihilation via the ongoing
sixth mass extinction signaled by vertebrate population losses and declines', *Proceedings of
the National Academy of Sciences*, 114(30), pp. E6089–E6096. Available at:
https://doi.org/10.1073/pnas.1704949114.

Collen, B. *et al.* (2009) 'Monitoring Change in Vertebrate Abundance: the Living Planet
Index', *Conservation Biology*, 23(2), pp. 317–327. Available at:
https://doi.org/10.1111/j.1523-1739.2008.01117.x.

Dinerstein, E. *et al.* (2017) 'An Ecoregion-Based Approach to Protecting Half the Terrestrial
Realm', *BioScience*, 67(6), pp. 534–545. Available at: https://doi.org/10.1093/biosci/bix014.

Dinerstein, E. *et al.* (2019) 'A Global Deal For Nature: Guiding principles, milestones, and
targets', *Science Advances*, 5(4), p. eaaw2869. Available at:
https://doi.org/10.1126/sciadv.aaw2869.

Felipe-Lucia, M.R. *et al.* (2020) 'Land-use intensity alters networks between biodiversity,
ecosystem functions, and services', *Proceedings of the National Academy of Sciences*, p. 11.
Available at: https://doi.org/www.pnas.org/cgi/doi/10.1073/pnas.2016210117.

Gaona, F.P. *et al.* (2021) 'Drastic loss ofinsects (Lepidoptera: Geometridae) in urban
landscapes in a tropicalbiodiversity hotspot', *Journal of Insect Conservation*, 25(3), pp. 395–
405. Available at: https://doi.org/10.1007/s10841-021-00308-9.

Gaston, K.J. (2000) 'Global patterns in biodiversity', *Nature*, 405(6783), pp. 220–227.
Available at: https://doi.org/10.1038/35012228.

Gibson, L. *et al.* (2011) 'Primary forests are irreplaceable for sustaining tropical
biodiversity', *Nature*, 478(7369), pp. 378–381. Available at:
https://doi.org/10.1038/nature10425.

Goldewijk, K.K. *et al.* (2011) 'The HYDE 3.1 spatially explicit database of human-induced
global land-use change over the past 12,000 years', *Global Ecology and Biogeography*,
20(1), pp. 73–86. Available at: https://doi.org/10.1111/j.1466-8238.2010.00587.x.

Green, E.J. *et al.* (2020) 'Below the canopy: global trends in forest vertebrate populations and
their drivers', *Proceedings of the Royal Society B: Biological Sciences*, 287(20200533).
Available at: http://dx.doi.org/10.1098/rspb.2020.0533.

Greenville, A.C. *et al.* (2018) 'Biodiversity responds to increasing climatic extremes in a
biome-specific manner', *Science of The Total Environment*, 634, pp. 382–393. Available at:
https://doi.org/10.1016/j.scitotenv.2018.03.285.

Hudson, L.N. *et al.* (2017) 'The database of the PREDICTS (Projecting Responses of
Ecological Diversity In Changing Terrestrial Systems) project', *Ecology and Evolution*, 7(1),
pp. 145–188. Available at: https://doi.org/10.1002/ece3.2579.

Ingram, D.J. *et al.* (2021) 'Targeting Conservation Actions at Species Threat Response
Thresholds', *Trends in Ecology & Evolution*, 36(3), pp. 216–226. Available at:
https://doi.org/10.1016/j.tree.2020.11.004.

IPBES (2019) *Global assessment report on biodiversity and ecosystem services of the
Intergovernmental Science-Policy Platform on Biodiversity and Ecosystem Services. E. S.
Brondizio, J. Settele, S. Díaz, and H. T. Ngo (editors)*. Bonn, Germany: IPBES secreteriat.

Isaac, N.J.B. and Cowlishaw, G. (2004) 'How species respond to multiple extinction threats',
Proceedings of the Royal Society of London. Series B: Biological Sciences, 271(1544), pp.
1135–1141. Available at: https://doi.org/10.1098/rspb.2004.2724.

Isbell, F. *et al.* (2017) 'Linking the influence and dependence of people on biodiversity across
scales', *Nature*, 546(7656), pp. 65–72. Available at: https://doi.org/10.1038/nature22899.

IUCN (2022) *The IUCN Red List of Threatened Species. Version 2022-2*. Available at:
https://www.iucnredlist.org (Accessed: 17 August 2023).

Lambertini, M. (2020) *Living Planet Report 2020: Bending the Curve of Biodiversity Loss*.
Available at: http://www.deslibris.ca/ID/10104983 (Accessed: 1 April 2021).

Leung, B., Greenberg, D.A. and Green, D.M. (2017) 'Trends in mean growth and stability in
temperate vertebrate populations', *Diversity and Distributions*, 23(12), pp. 1372–1380.
Available at: https://doi.org/10.1111/ddi.12636.

Lewis, S.L., Edwards, D.P. and Galbraith, D. (2015) 'Increasing human dominance of
tropical forests', *Science*, 349(6250), pp. 827–832. Available at:
https://doi.org/10.1126/science.aaa9932.

Mace, G.M. *et al.* (2018) 'Aiming higher to bend the curve of biodiversity loss', *Nature*
Sustainability, 1(9), pp. 448–451. Available at: https://doi.org/10.1038/s41893-018-0130-0.

Moncrieff, G.R., Bond, W.J. and Higgins, S.I. (2016) 'Revising the biome concept for
understanding and predicting global change impacts', *Journal of Biogeography*, 43(5), pp.
863–873. Available at: https://doi.org/10.1111/jbi.12701.

Monsarrat, S., Jarvie, S. and Svenning, J.-C. (2019) 'Anthropocene refugia: integrating
history and predictive modelling to assess the space available for biodiversity in a human-
dominated world', *Philosophical Transactions of the Royal Society B: Biological Sciences*,
374(20190219). Available at: https://doi.org/10.1098/rstb.2019.0219.

Monsarrat, S. and Svenning, J.-C. (2022) 'Using recent baselines as benchmarks for
megafauna restoration places an unfair burden on the Global South', *Ecography*, 2022(4).
Available at: https://doi.org/10.1111/ecog.05795.

Murphy, S.J. (2021) 'Sampling units derived from geopolitical boundaries bias biodiversity
analyses', *Global Ecology and Biogeography*, 30(9), pp. 1876–1888. Available at:
https://doi.org/10.1111/geb.13352.

Myers, N. *et al.* (2000) 'Biodiversity hotspots for conservation priorities', *Nature*, 403(6772),
pp. 853–858. Available at: https://doi.org/10.1038/35002501.

Newbold, T. *et al.* (2013) 'Ecological traits affect the response of tropical forest bird species
to land-use intensity', *Proceedings of the Royal Society B: Biological Sciences*, 280(1750), p.
20122131. Available at: https://doi.org/10.1098/rspb.2012.2131.

Newbold, T. *et al.* (2015) 'Global effects of land use on local terrestrial biodiversity', *Nature*,
520(7545), pp. 45–50. Available at: https://doi.org/10.1038/nature14324.

Newbold, T. (2015) 'StatisticalModels'. Available at:
https://github.com/timnewbold/StatisticalModels.

Newbold, T. *et al.* (2016) 'Has land use pushed terrestrial biodiversity beyond the planetary
boundary? A global assessment', *Science*, 353(6296), pp. 288–291. Available at:
https://doi.org/10.1126/science.aaf2201.

Newbold, T. (2018) 'predictsFunctions: Functions for reading and processing the PREDICTS
data'.

Newbold, T. *et al.* (2018) 'Widespread winners and narrow-ranged losers: Land use
homogenizes biodiversity in local assemblages worldwide', *PLOS Biology*. Edited by H.
Morlon, 16(12), p. e2006841. Available at: https://doi.org/10.1371/journal.pbio.2006841.

Newbold, T. *et al.* (2020) 'Tropical and Mediterranean biodiversity is disproportionately
sensitive to land-use and climate change', *Nature Ecology & Evolution* [Preprint]. Available
at: https://doi.org/10.1038/s41559-020-01303-0.

Olson, D.M. *et al.* (2001) 'Terrestrial Ecoregions of the World: A New Map of Life on
Earth', *BioScience*, 51(11), p. 933. Available at: https://doi.org/10.1641/0006-
3568(2001)051[0933:TEOTWA]2.0.CO;2.

Olson, D.M. and Dinerstein, E. (2002) 'The Global 200: Priority Ecoregions for Global
Conservation', *Annals of the Missouri Botanical Garden*, 89(2), pp. 199–224. Available at:
https://doi.org/10.2307/3298564.

Phillips, H.R.P., Newbold, T. and Purvis, A. (2017) 'Land-use effects on local biodiversity in
tropical forests vary between continents', *Biodiversity and Conservation*, 26(9), pp. 2251–
2270. Available at: https://doi.org/10.1007/s10531-017-1356-2.

Quintero, S. *et al.* (2023) 'Effects of human-induced habitat changes on site-use patterns in
large Amazonian Forest mammals', *Biological Conservation*, 279, p. 109904. Available at:
https://doi.org/10.1016/j.biocon.2023.109904.

Smith, J.R. *et al.* (2018) 'A global test of ecoregions', *Nature Ecology & Evolution*, 2(12),
pp. 1889–1896. Available at: https://doi.org/10.1038/s41559-018-0709-x.

Smith, J.R. *et al.* (2020) 'The biogeography of ecoregions: Descriptive power across regions
and taxa', *Journal of Biogeography*, 47(7), pp. 1413–1426. Available at:
https://doi.org/10.1111/jbi.13871.

Sodhi, N.S. *et al.* (2004) 'Southeast Asian biodiversity: an impending disaster', *Trends in*
Ecology & Evolution, 19(12). Available at: https://doi.org/doi:10.1016/j.tree.2004.09.006.

Sodhi, N.S. *et al.* (2009) 'A Meta-Analysis of the Impact of Anthropogenic Forest
Disturbance on Southeast Asia's Biotas', *Biotropica*, 41(1), pp. 103–109. Available at:
https://doi.org/10.1111/j.1744-7429.2008.00460.x.

Song, X.-P. *et al.* (2018) 'Global land change from 1982 to 2016', *Nature*, 560(7720), pp.
639–643. Available at: https://doi.org/10.1038/s41586-018-0411-9.

Srivathsa, A. *et al.* (2018) 'Substituting space for time: Empirical evaluation of spatial
replication as a surrogate for temporal replication in occupancy modelling', *Journal of*
Applied Ecology, 55(2), pp. 754–765. Available at: https://doi.org/10.1111/1365-2664.13005.

Stephenson, P.J. *et al.* (2015) 'Overcoming the challenges to conservation monitoring:
integrating data from *in-situ* reporting and global data sets to measure impact and
performance', *Biodiversity*, 16(2–3), pp. 68–85. Available at:
https://doi.org/10.1080/14888386.2015.1070373.

Stevens, G.C. (1989) 'The Latitudinal Gradient in Geographical Range: How so Many
Species Coexist in the Tropics', *The American Naturalist*, 133(2), pp. 240–256. Available at:
https://doi.org/10.1086/284913.

Suraci, J.P. *et al.* (2021) 'Disturbance type and species life history predict mammal responses
to humans', *Global Change Biology*, 00, pp. 1–14. Available at:
https://doi.org/10.1111/gcb.15650.

Tittensor, D.P. *et al.* (2014) 'A mid-term analysis of progress toward international
biodiversity targets', *Science*, 346(6206), pp. 241–244. Available at:
https://doi.org/10.1126/science.1257484.

Walker, L.R. *et al.* (2010) 'The use of chronosequences in studies of ecological succession
and soil development', *Journal of Ecology*, 98(4), pp. 725–736. Available at:
https://doi.org/10.1111/j.1365-2745.2010.01664.x.

WWF (2022) *Living Planet Report 2022 - Building a nature positive society*. Gland,
Switzerland: WWF. Available at:
https://wwflpr.awsassets.panda.org/downloads/lpr_2022_full_report_1.pdf (Accessed: 17
October 2022).

Yang, X. *et al.* (2021) 'Threatened bird species are concentrated in regions with less
historical human impacts', *Biological Conservation*, 255, p. 108978. Available at:
https://doi.org/10.1016/j.biocon.2021.108978.

Data Accessibility Statement

All data used in this manuscript is downloaded from open-source locations. The PREDICTS
database can be downloaded from the Natural History Museum Data portal
(https://data.nhm.ac.uk/dataset/the-2016-release-of-the-predicts-database).
616
617 Geographical data used to delineate regional biome boundaries in this manuscript can be
618 downloaded from The Nature Conservancy's Geospatial Conservation Atlas
619 (https://geospatial.tnc.org/datasets/b1636d640ede4d6ca8f5e369f2dc368b/about).
620
621 All R code used to run analysis presented in this manuscript is accessible in a GitHub
622 repository (not included here to maintain anonymity)
623
624 Figures
625

Legend:
- Tropical and Subtropical Moist Broadleaf Forests
- Tropical and Subtropical Grasslands, Savannas and Shrublands
- Tropical and Subtropical Dry Broadleaf Forests
- Tropical and Subtropical Coniferous Forests
- Temperate Grasslands, Savannas and Shrublands
- Temperate Broadleaf and Mixed Forests
- Temperate Conifer Forests
- Montane Grasslands and Shrublands
- Mediterranean Forests, Woodlands and Scrub
- Mangroves
- Flooded Grasslands and Savannas
- Deserts and Xeric Shrublands
- Boreal Forests/Taiga
- Tundra
- Inland Water
- Rock and Ice

626 **Figure 1. Map of regional biomes.** Solid coloured areas represent biomes (see legend).

627 Black lines separate out biogeographic realms. Black dots represent sites in the PREDICTS

628 database used in our global model (Hudson et al 2014). Hashed coloured areas represent

629 regional biomes that were not included in analysis due to data deficiency (see methods).

630 Biome and biogeographic realm spatial data from (Olson *et al.*, 2001).

Figure 2. Change in Akaike's Information Criterion (ΔAIC) for Generalised Linear Mixed Models (GLMMs) predicting change in a global dataset of (a) species richness and (b) total abundance consistently suggests models including regional biome have highest support when modelling response to land-use change. Four GLMMs with differing fixed-effect structures containing Land Use (LU) interacting with Regional Biome (RB), Biome or Realm were compared using AIC values. A model with low AIC is considered to have higher support. ΔAIC is calculated as the difference in AIC from the lowest scoring model. The box and whisker plots represent a summary of ΔAIC values of 100 hold-out models, where each iteration removed 10% of studies at random as a test of model robustness. Importantly, for a given dataset in this hold-out analysis, LU:RB model had the lowest AIC in 100% of iterations. The PREDICTS dataset was subset to only include data from regional biomes with 25 data points per combination of regional biome and land-use type. Land Use is a discrete variable with Primary Vegetation, Secondary Vegetation, Plantation Forest, Pasture and Cropland as its categories (see Table S2 for descriptions).

Figure 3. Responses of biodiversity to land-use change are unique across tropical

forest regional biomes. The results of GLMMs predicting the response of species richness

across land-use types and land-use intensity, when compared to primary vegetation (PV)

across tropical forest regional biomes. Responses of species richness of plants and animals

were predicted across land-use types (a) including secondary vegetation (SV), plantation

forest (PF), pasture (Pa) and cropland (Cr). Responses were also predicted for two levels of

land-use intensity: minimal (M) or intense (I) in secondary vegetation (SV) and agriculture

(Agr) land use types (b). Each point represents mean prediction, with 75% confidence

intervals (thick whiskers) and 95% confidence intervals (thin whiskers). Responses are

considered significantly different if the 75% confidence intervals do not overlap.

Figure 4. Responses of biodiversity to land-use change are hard to predict

unique across temperate forest regional biomes. The results of GLMMs

predicting the response of species richness across land-use types and land-use

intensity, when compared to primary vegetation (PV) across temperate forest

regional biomes. Responses of species richness of plants and animals were

predicted across land-use types (a) including secondary vegetation (SV), plantation

forest (PF), pasture (Pa) and cropland (Cr). Responses were also predicted for two

levels of land-use intensity: minimal (M) or intense (I) in secondary vegetation (SV)

and agriculture (Agr) land use types (b). Each point represents mean prediction, with

75% confidence intervals (thick whiskers) and 95% confidence intervals (thin

whiskers). To keep scales consistent, upper 95% CIs greater than 100 have been

included as text instead. Responses are considered significantly different if the 75%

confidence intervals do not overlap.

Figure 5. Responses of biodiversity to land-use change are hard to predict unique

across tropical grassland regional biomes. The results of GLMMs predicting the

response of species richness across land-use types and land-use intensity, when compared

to primary vegetation (PV) across tropical grassland regional biomes. Responses of species

richness of plants and animals were predicted across land-use types (a) including secondary

vegetation (SV), plantation forest (PF), pasture (Pa) and cropland (Cr). Responses were

also predicted for two levels of land-use intensity: minimal (M) or intense (I) in secondary

vegetation (SV) and agriculture (Agr) land use types (b). Each point represents mean

prediction, with 75% confidence intervals (thick whiskers) and 95% confidence intervals (thin

whiskers). To keep scales consistent, upper 95% CIs greater than 100 have been included

as text instead. Responses are considered significantly different if the 75% confidence

intervals do not overlap.

Figure 6. Responses of species richness in individual taxa groups to land-use change

across regional biomes. The results of 6 GLMMs predicting the response of species

richness across land-use types and realm across three biomes (Tropical Forest, Temperate

Forest and Tropical grassland) and three taxon groups (Vertebrate, Invertebrate and Plants)

when compared to primary vegetation (PV) across tropical grassland regional biomes.

Responses of species richness were predicted across disturbed land-use types including

secondary vegetation (SV), plantation forest (PF), pasture (Pa) and cropland (Cr). Each

point represents mean prediction, with 75% confidence intervals (thick whiskers) and 95%

confidence intervals (thin whiskers). To keep scales consistent, upper 95% CIs greater than

695   100 have been included as text instead. Responses are considered significantly different if

696   the 75% confidence intervals do not overlap.

697