Article title: Biodiversity shows unique responses to land-use change across regional

biomes

Authors:

Peggy Bevan[1,2], Guilherme Braga Ferreira[1], Daniel J Ingram[3], Marcus Rowcliffe[2], Lucy

Young[4], Robin Freeman[2], Kate E. Jones[1].

Affiliations:

1. Centre for Biodiversity and Environment Research, University College London, Gower

Street, WC1E 6BT

2. Institute of Zoology, Zoological Society of London, Wellcome Building, Outer Circle,

London NW8 7LS

3. Durrell Institute of Conservation and Ecology, School of Anthropology and Conservation,

University of Kent, Canterbury, CT2 7NR, UK.

4. WWF UK, Living Planet Centre, Brewery Road, Woking, GU21 4LL

Corresponding author: Peggy Bevan, peggy.bevan.17@ucl.ac.uk

Data availability:
All data used in this manuscript is downloaded from open source locations. The PREDICTS
database can be downloaded from the Natural History Museum Data portal
(https://data.nhm.ac.uk/dataset/the-2016-release-of-the-predicts-database).
20
21 Geographical data used to delineate regional biome boundaries in this manuscript can be
22 downloaded from The Nature Conservancy's Geospatial Conservation Atlas
23 (https://geospatial.tnc.org/datasets/b1636d640ede4d6ca8f5e369f2dc368b/about).
24
25 All R code used to run analysis presented in this manuscript can be found at
26 https://github.com/PeggyBevan/RegionalBiomes
27

Abstract

Aim: Biogeography has a critical influence on how ecological communities respond to

threats and how effective conservation interventions are designed. For example, the

resilience of ecological communities is linked to environmental and climatic features, and the

nature of threats impacting ecosystems also varies geographically. Understanding

community-level threat responses may be most accurate at fine spatial scales, however

collecting detailed ecological data at such a high resolution would be prohibitively resource

intensive. In this study, we aim to find the spatial scale that could best capture variation in

community-level threat responses whilst keeping data collection requirements feasible.

Location: Global Terrestrial

Time period: 1980 – Present

Major Taxa studied: Terrestrial Plants & Animals

Methods: Using a database of biodiversity records with extensive global coverage, we

modelled species richness and total abundance (the responses) across land-use types

(reflecting threats), considering three different spatial scales: biomes, biogeographical

realms, and *regional biomes* (the interaction between realm and biome). We then modelled

data from three highly sampled biomes separately to ask how responses to threat differ

between regional biomes.

Results: We found strong support for regional biomes in explaining variation in species

richness and total abundance compared to biomes or realms alone. Responses to disturbed

land-use types compared to primary vegetation were more pronounced and more distinct

between regional biomes in the tropics compared to the temperate zone.

Main conclusions: Our results suggest that regional biomes have potential as a reasonable

spatial scale for understanding how ecological communities respond to threats and

designing effective conservation interventions, but the lack of variation in response to land-

use change between temperate regional biomes could mean that this framework would be

more beneficial for refining the monitoring of tropical habitats.

Introduction

Despite multiple internationally agreed science-based conservation targets, biodiversity

continues to decline across the globe, with the consequent loss of essential ecosystem

services (Tittensor *et al.*, 2014; Brooks *et al.*, 2015; Ceballos, Ehrlich and Dirzo, 2017; Isbell

et al., 2017; Mace *et al.*, 2018; IPBES, 2019; Ingram *et al.*, 2021). As new targets are

developed, such as the up-coming post-2020 Convention on Biological Diversity goals and

the proposed Global Deal for Nature (Dinerstein *et al.*, 2019; CBD, 2020), it is vital to

develop biodiversity monitoring systems that are able to quantify the relationship between

anthropogenic threat and biodiversity response metrics (*e.g.,* abundance, species richness).

Such threat-response relationships will be influenced by specific life-history traits,

biogeography and the ecological scale being measured (*e.g.*, a community, population or

individual) (Isaac and Cowlishaw, 2004; Murphy, 2021; Suraci *et al.*, 2021). Habitat

disturbance caused by land-use change is a major factor contributing to biodiversity loss

(Newbold *et al.*, 2015), but the impact of disturbance on biodiversity varies with the intensity

of change (Felipe-Lucia *et al.*, 2020) and the original habitat (Monsarrat, Jarvie and

Svenning, 2019). This variation of impact of habitat loss also means that protection

interventions should be proportionate to the threat. To create realistic predictions of

biodiversity trends, the underlying biogeographic processes contributing to variation in

threat-response relationships should be considered when modelling global biodiversity

trends.

The biogeography of a species or ecological community can impact its sensitivity to an

anthropogenic threat. Tropical biomes, particularly tropical forest, have higher species

richness, smaller species' ranges, and higher number of specialist species compared to

temperate biomes (Stevens, 1989; Gaston, 2000; Brown, 2014). Habitat specialisation and

small range size have been associated with stronger negative response to land-use change

(Newbold *et al.*, 2013), suggesting communities in tropical biomes have lower resilience to

environmental disturbance compared to communities in similar temperate biomes.

Furthermore, the history of land use and environmental disturbance is not uniform across the

globe and will further contribute to modern day responses to anthropogenic threats. Historic

anthropogenic disturbance has been more intense in temperate biomes and these areas

have already lost a large number of species (Goldewijk *et al.*, 2011). Areas with a history of

disturbance have been found to contain fewer threatened species than less disturbed

counterparts, reflecting the significance of historic disturbance on threat-sensitive species

(Yang *et al.*, 2021). Incorporating an understanding of biogeography and historical

knowledge of an environment could play an important role in global biodiversity monitoring.

Given the implications of biogeography for threat-response relationships, monitoring

biodiversity at the smallest possible scale would be appropriate. However, there are

currently 37,000 threatened species on the red list (IUCN, 2021), all of which cannot have

unique conservation plans created for them. There is a necessity to find an appropriate scale

to monitor biodiversity that finds a balance between presenting global trends and efficiently

implementing local-scale interventions (Ingram *et al.*, 2021). One way to account for

biogeographic variation in threat responses is to split the world into spatial units, however

the method for this is contended. For example, creating national statistics for biodiversity is

helpful for feeding into policy, but creates accuracy issues as biodiversity is not nationally

constrained (Murphy, 2021). Secondly, a global framework has been suggested for

monitoring biodiversity at an ecoregional level (Dinerstein *et al.*, 2017, 2019; Smith *et al.*,

2018). Ecoregions (n = 814) are described as unique spatial areas that are determined by

ecological features (Olson and Dinerstein, 2002) but are more ecologically distinct in some

regions than others (McDonald *et al.*, 2005; Smith *et al.*, 2020). The ecoregion framework is

powerful when measuring variables that can be obtained using remote sensing or global

species lists (*e.g.*, Dinerstein *et al.*, 2017; Smith *et al.*, 2020), but many ecoregions are

lacking the required field data to practically study population responses to human pressure

at this resolution (Stephenson *et al.*, 2015). There is a need to test whether coarser spatial

scales can capture variation in threat-response relationships so that data collection

resources can be prioritised, whilst still providing meaningful evidence on biodiversity trends.

The broadest spatial unit of terrestrial habitats is the climatically-defined biome, of which

there are 14 (Olson *et al.*, 2001). Responses of animal and plant populations to threats do

differ between biomes (Greenville *et al.*, 2018; Green *et al.*, 2020) and species richness is

particularly sensitive to land use change in tropical forest, tropical grassland and

Mediterranean biomes compared to temperate and desert biomes (Newbold *et al.*, 2020). In

addition to biomes, there are 8 biogeographic realms (Olson *et al.*, 2001; UNEP-WCMC

2004). Data from the Living Planet Database (Collen *et al.*, 2009; WWF, 2022) has shown

that decreases in vertebrate populations are more pronounced in southern hemisphere

realms, particularly the Neotropics (Green *et al.*, 2020; WWF, 2022). Although there is

evidence for variation in threat-response relationships between biomes and between realms,

the interaction between these two intersecting spatial units is rarely considered, which may

mean a large amount of variation is unaccounted for when monitoring biodiversity in biomes

or realms alone.

A potential, intermediate spatial framework for monitoring biodiversity are regional biomes

(the interaction of realms and biomes) (n = 64) (Ingram *et al.*, 2021). Regional biomes

contain 11 ecoregions on average, ranging from 1-81 ecoregions (Olson and Dinerstein,

2002). Separating biomes by biogeographic realms can account for differences in vegetative

structure, threats and socioeconomic status that occur between realms (Moncrieff, Bond and

Higgins, 2016; Allan *et al.*, 2019). Furthermore, threats are not spread evenly across the

world and can be region-specific (Bowler *et al.*, 2020); for example tropical forest in

southeast Asia is a threat 'hotspot', whereas large parts of the Amazon rainforest act as

threat 'refugia' (Allan *et al.*, 2019). Some areas of forest biomes are being afforested, mainly

in the northern hemisphere, whilst others experience mass deforestation (Song *et al.*, 2018).

Regional variation in sensitivity to anthropogenic pressure within biomes has been mainly

explored in tropical biomes. In Asian tropical forests, species richness was found to be

significantly more sensitive to disturbance than other regions (South America and Africa)

(Gibson *et al.*, 2011; Phillips, Newbold and Purvis, 2017). On the other hand, vertebrate

abundance is declining more in the Neotropics and Afrotropics than in the Indo-Pacific

region (Green *et al.*, 2020). Regional biomes may present a middle-ground between the finer

scale monitoring of ecoregions and the coarser biome scale, and where adequate sampling

of biodiversity responses is more achievable.

Here, we use the PREDICTS database (Hudson *et al.*, 2017) to analyse data from over 400

field studies to explore the efficacy of regional biomes as a spatial monitoring unit for

terrestrial species. We ask if regional biomes have more explanatory power than realms or

biomes alone when modelling the effect of land-use change on species richness and total

abundance. Due to the known explanatory power of biomes and realms (Phillips, Newbold

and Purvis, 2017; Newbold *et al.*, 2020), we expect the interaction between realm and biome

to have increased power in explaining variation in biodiversity metrics between different

land-use types. Secondly, as case studies we model the three most data-rich biomes

(Tropical Forest, Temperate Forest, Tropical Grasslands) in the PREDICTS database

separately to explore how responses to land-use change and use-intensity vary between

regional biomes. Our expectation is that there will be negative responses to human-

dominated land-use types compared to primary vegetation, but the magnitude of response

will differ between regional biomes. We expect the highest variation but also uncertainty in

predicted responses between tropical regional biomes due to ecoregions being more distinct

in these biomes (Smith *et al.*, 2020) and less variation in response between temperate

regional biomes due to biotic homogenization (Newbold *et al.*, 2018).

164 Methods

165 The PREDICTS database

We used species abundance and occupancy data from the PREDICTS database (Hudson et

al 2017; 22678 sites, 480 sources, 666 studies, 47,000 species). We chose to use the

PREDICTS database due to high global coverage compared to other databases (Hudson *et*

al., 2017), making it suitable for a study comparing many regional biomes. The PREDICTS

project (Predicting Responses of Ecological Diversity In Changing Terrestrial Systems;

www.predicts.co.uk) is a database that collates biodiversity studies with comparable

measures of terrestrial biodiversity from sites of different land uses and land-use intensity

(Hudson *et al.*, 2017). All sampled sites are classified by stage of disturbance or recovery

which can be used as a substitute for actual temporal change at a single site (Walker *et al.*,

2010; Srivathsa *et al.*, 2018). This space-for-time substitution methods allows for specific

measurements of biodiversity over distinct land-use types, acting as a proxy for

anthropogenic pressure. PREDICTS does not include marine or freshwater data but does

have extensive global coverage and has made attempts to reduce taxonomic and

geographic bias (Hudson *et al.*, 2017).

We summarised occurrence and abundance records by site using the *predictsFunctions* R

package (Newbold, 2018). The PREDICTS database covers 47 out of 64 regional biomes

(70%), with some regional biomes more comprehensively sampled than others (Table S1).

For analysis, we excluded data from three biomes with fewer than 50 sites in total from the

dataset (Flooded Grasslands and Savannahs, Tundra, Deserts & Xeric Shrublands and

Mangroves). Additionally, regional biomes were considered data deficient and excluded from

analysis if they had no sites in primary vegetation land-use types (our reference level), or no

sites in any kind of managed land use (cropland, pasture and plantation forest). This

ensured that regional biomes included in analysis had data points (sites) across a gradient

of increasingly anthropogenic land-use types. After removing data deficient regional biomes,

a total of 30 regional biomes were available for analysis (Fig. 1; 20555 sites, 561 studies).

Using the predominant land use and use-intensity classifications from the PREDICTS

database (land use: primary vegetation, mature secondary vegetation, intermediate

secondary vegetation, young secondary vegetation, plantation forest, pasture, cropland and

urban; use intensity: minimal, light, intense) (Hudson *et al.*, 2017), we created two land-use

variables – *LandUse* and *UseIntensity*. We combined the three secondary vegetation

classes (mature, intermediate and young) into one and removed all data from urban land-

use types. Because some combinations of regional biome and *LandUse* class have low

sample sizes, we tested 5 variations of this variable where categories were aggregated to

find the most parsimonious grouping that still explained variation in the model well. For

example, to make the *LandUse2* variable we grouped primary vegetation and secondary

vegetation as 'natural vegetation'. Full descriptions of each land-use type and the 5

variations of *LandUse* can be found in Table S2. *UseIntensity* is a combination of

predominant land-use type and use-intensity. To create this variable, we grouped 'minimal'

and 'light' use-intensities into 'minimal' and combined with groups from *LandUse3* (defined

as primary vegetation, secondary vegetation and agriculture) to give categories primary

vegetation, minimal secondary vegetation, intense secondary vegetation, minimal

agriculture, intense agriculture. Primary vegetation sites were not split by use intensity as

there is no difference in biodiversity metrics between primary vegetation sites with differing

use-intensities (Newbold *et al.*, 2015).

Statistical Analysis

We used generalised mixed-effects models (Bolker *et al.*, 2009) to test for differences in

species richness and total abundance in response to land use, biome, realm and regional

biome (an interaction of biome and realm). Both response variables (species richness and

total abundance) were log transformed (natural logarithm) and modelled with a gaussian

distribution. We included study as a random intercept to account for site-level differences

and data collection methods. We ran a series of models with species richness and total

abundance against each *LandUse* variable (i.e. *LandUse1, LandUse2, or LandUse3* – see

the previous section) and selected the best fitting variable using Akaike Information Criterion

(AIC) (Burnham, Anderson and Huyvaert, 2011). We found that *LandUse1* (Primary

vegetation, Secondary vegetation, Agriculture, Plantation Forest, Cropland) had the most

support and was used for remaining analyses (Table S3).

Global analysis of regional biomes

To see if including regional biome helps to explain more variation in biodiversity than realm

or biome, we ran a model selection process using ΔAIC values to assess the support for

each variable. The fixed effect structures tested were *LandUse, LandUse:Realm,*

LandUse:Biome, and *LandUse:RegionalBiome* and species richness and total abundance

were used as response variables. We tested model robustness in two ways. First, we

increased the sample size threshold that would allow a regional biome to be included in the

model (minimum 1, 5, 25 or 50 sites in each regional biome-land use combination) and

checked if the model selection process gave the same results in each case. Second, we ran

a series of 100 hold-out models where each iteration randomly removed 10% of studies to

see if this would change the results of model selection.

<u>Biome-specific case studies</u>

As case studies, we investigated the differences in biodiversity response to land-use change

between regional biomes by selecting the three biomes which had three or more regional

biomes with at least 100 sites in the PREDICTS database (Fig. 2). The biomes selected

were tropical and subtropical moist broadleaf forest (Tropical Forest) (3 realms; Indo-Malay:

1781 sites; Neotropics: 2274 sites; Afrotropics: 2081 sites), temperate broadleaf and mixed

forests (Temperate Forest) (4 realms; Palearctic: 3986 sites; Neotropics: 577 sites; Nearctic:

856 sites; Australasia: 336 sites) and tropical and subtropical savannahs, grasslands and

shrublands (Tropical Grassland) (3 realms; Afrotropic: 1663 sites; Neotropics: 190 sites;

Australasia: 502 sites). For each biome and biodiversity metric (species richness and total

abundance), we ran four models that used the *LandUse* or <u>*UseIntensity*</u> variables and tested

if including *Realm*, which in this case represents regional biomes. We then predicted the

average change in species richness and total abundance in each land-use type in each

realm using the *StatisticalModels* R package (Newbold, 2015).

<u>Results</u>

<u>Global analysis of regional biomes</u>

Including regional biome as a variable improved model fit and better explained the effect of

land-use change on species richness and total abundance (Fig. 3). Models with realm or

biome alone had lower support and this was consistent across 100 iterations of hold-out

models (Fig. 3). Increasing the sample size threshold for regional biomes did not change the

overall outcome (Table S4). Furthermore, marginal R^2 values showed that regional biome

explained more variation in both species richness and total abundance than biome or realm

interacting with land use (Table S4).

<u>Biome-specific case studies</u>

We found that species richness and total abundance in each regional biome of tropical forest

showed a distinct response to land-use change. Species richness reduced in human-

dominated land-use types compared to primary vegetation in all realms but to different

degrees at each land-use type (Fig 4a). The largest loss of species richness was seen in

Afrotropical cropland (-53% reduction compared to primary vegetation), followed by Indo-

Malayan cropland (-50%) and plantation forest (-43%). Conversely, the Neotropics did not

show a significant species richness reduction in cropland and a barely significant reduction

in plantation forest (-20%), with species richness for this regional biome being worst affected

in pasture land-use types (-30%). When predicting responses of total abundance to land-use

change in the tropical forest biome there was much higher uncertainty. The negative

responses to land-use type seen in the species richness models did not always carry over to

the abundance models, for example there was no sign of a negative response in Neotropical

pasture. However, total abundance was significantly lower in Indo-Malay plantation forest

compared to other regional biomes, mirroring the pattern in species richness (Fig 4c). For

the models including *UseIntensity*, a negative response to increased disturbance can be

observed in Indo-Malayan and Afrotropical regional biomes, with declines >50% compared

to primary vegetation in both species richness and total abundance (Fig 4b, d). This extreme

response is not seen in the Neotropical regional biome.

In temperate forest regional biomes, species richness and total abundance both showed low

to no response in human-dominated land-use types compared to primary vegetation (Fig 5a,

c). The exception is the Palearctic regional biome, where total abundance reduced by 55%

in cropland land-use types. There were also reductions in species richness in plantation

forest and cropland in this regional biome, but the upper confidence limits just surpassing 0

(Fig 5a). This was the only response from any temperate forest regional biome that showed

a significant change from primary vegetation (95% confidence intervals do not include 0).

Furthermore, predictions for change in species richness and total abundance have high

levels of uncertainty for all realms except palearctic.

Similar to temperate forest, biodiversity metrics in most tropical grassland regional biomes

showed little or no response to any land-use type when compared to primary vegetation

(Fig. 6). Predictions of total abundance change showed a high degree of uncertainty across

all regional biomes, except for the Afrotropics where there was a strong response to intense

agriculture (-67%), although this was the only regional biome with enough datapoints to

predict in this category so cannot be compared. Species richness in Australasia showed

negative responses to pasture and minimally used agricultural land, but with the upper

confidence interval just passing zero in both cases (Fig. 6a, b).

<u>Discussion</u>

<u>Global analysis of regional biomes</u>

Our results show that including regional biome when modelling the effects of land use on

biodiversity can increase explanatory power. There is a need for a biodiversity monitoring

spatial framework that finds the balance between having feasible data collection

requirements and adequate global coverage, at a realistic scale that captures true variation

in ecological resilience to disturbance (Ingram *et al.*, 2021). It has been previously shown

that biodiversity shows different responses to disturbance across biomes (Greenville *et al.*,

2018; Blowes *et al.*, 2019; Newbold *et al.*, 2020) and across realms (Gibson *et al.*, 2011;

Phillips, Newbold and Purvis, 2017; WWF, 2020), but for the first time, we have shown the

important interaction between biome and realm. The combination of these two spatial

delineations gives us more information about ecological processes. This result is important

for biodiversity monitoring and conservation interventions as not only are the magnitude of

biodiversity responses to anthropogenic pressure different between regions but that

threatening processes (i.e., land-use types) are affecting regional biomes differently.

<u>Biome-specific case studies</u>

Whilst our models suggest that regional biome can modulate the effect of land-use change

on species richness and abundance on a global scale, a more focused analysis found that

the influence of regional biome is stronger in some biomes than others. Generally, species

richness and total abundance in tropical biomes showed negative responses to disturbed

land-use types, the degree of which changed with realm. In temperate forest, however, there

was no significant response to land-use change overall, excluding the palearctic realm. The

observed difference between tropical and temperate regional biomes is expected as

responses to land-use change are stronger in tropical biomes compared to temperate

biomes, especially tropical forest (Newbold *et al.*, 2020). However, ecoregions in tropical

biomes are considered more ecologically distinct than neighbouring ecoregions compared to

those in temperate biomes (Smith *et al.*, 2018). As each regional biome is a group of many

ecoregions, it would follow that predicted responses to disturbance in tropical regional

biomes would contain more uncertainty than temperate, but the opposite is true for our case

studies. The high uncertainty in predicted responses produced by the temperate forest

models may be due to low sample sizes or variation in sampled taxa, considering that

predictions for palearctic temperate forest have very low uncertainty compared to all other

regional biomes, and is also the most extensively sampled regional biome in the PREDICTS

database (Table S1).

For tropical forest regional biomes, our model predictions suggest Indo-Malayan species

richness and total abundance respond most strongly to disturbance, followed by the

Afrotropics. There was no significant change in species richness in the Neotropics tropical

forest in any land-use type except for pasture. Biodiversity in Asian tropical forest is

repeatedly found to be extremely sensitive to land-use change and disturbance (Sodhi *et al.*,

2009; Gibson *et al.*, 2011; Phillips, Newbold and Purvis, 2017). The islands of south east

Asia have a rich biogeographical history which has led to this region having the highest

number of endemic species and biodiversity hotspots (Myers *et al.*, 2000; Sodhi *et al.*, 2004)

compared to other tropical regions. This area also has the highest proportion of species

under threat in the world (Allan *et al.*, 2019). This high baseline for species richness

combined with high levels of specialisation may partly explain why the response to land-use

change is so pronounced in this regional biome. Response metrics in Neotropical tropical

forest have the smallest response to disturbed land-use types compared to similar regional

biomes. However, temporal trends from the Living Planet Index suggest that populations of

vertebrates in south America have decreased by 94% since 1970 (WWF, 2020). Our results

are not directly comparable to this, but the difference is a demonstration of the importance of

testing multiple data types and of assessing the regional biome framework using time-series

biodiversity data in the future. Further research is needed to understand how biological and

taxonomic differences between these tropical forest realms might be impacting sensitivity to

land-use change.

The intensity and history of habitat disturbance in each regional biome will also impact the

observed response of biodiversity and may even cause biases in our data set. There is a

strong focus on palm oil as a driver of biodiversity loss in Asian tropics, which may lead to a

publication bias in this area. Temperate forest biomes have a long history of human

habitation and extinction events, meaning the baseline in primary vegetation is much lower

in these regional biomes compared to less historically disturbed biomes (Monsarrat and

Svenning, 2022). The lack of variation in responses observed in temperate forest regional

biomes may well be because of this lower baseline in primary vegetation. Indeed, trends of

vertebrate abundance have been increasing in the northern hemisphere since 1970 (Leung,

Greenberg and Green, 2017) as well as forest cover (Song *et al.*, 2018). The drivers that

cause differences in sensitivity to habitat disturbance are related to both biogeographic and

human history; splitting the world into regional biomes can further account for these

differences.

Although regional biomes may be a valuable spatial monitoring unit, our models were limited

by sample size for some regional biomes. The PREDICTS database is the most extensive

collection of terrestrial biodiversity records available, but there are still gaps. Under-sampling

was pronounced in non-forested temperate biomes like Montane Grasslands and

Shrublands, and realms were unevenly sampled, for example studies from Palearctic

temperate forest dominate the database but there are no studies at all on Palearctic

temperate grasslands (Table S1). Prioritising data collecting in under-represented regional

biomes highlighted here would enhance progress towards an effective monitoring framework

(Ingram *et al.*, 2021). Furthermore, taxon is an important factor to consider and would add

even more information. We could not include taxon as a predictor due to the already high

number of parameters combined with low sample sizes. This study focuses only on

terrestrial regional biomes, as these are more extensively sampled, and we were specifically

testing the effect of land-use change on species richness and abundance. We acknowledge

that marine biodiversity is showing stronger responses to disturbance than terrestrial

(Blowes *et al.*, 2019), and suggest a similar monitoring framework could be adapted for

marine ecosystems.

<u>Conclusion</u>

Our results show that regional biomes have the potential to be a powerful unit for monitoring

biodiversity and implementing conservation action. Resources for data collection and

biodiversity monitoring on the ground is sparse, and the regional biome framework implies

that monitoring studies could be more evenly spread across regional biomes, optimising

resource allocation for data collection. Here we have highlighted that there is within-biome

variation in how biodiversity is responding to disturbance. We show that the difference

between realms may be more important in tropical biomes than temperate, and that spatial

resolution should be carefully considered in any attempts at monitoring global biodiversity

trends. Although more data are needed, regional biomes have the potential to be a

meaningful scale at which to prioritise monitoring. These results will be beneficial for

informing upcoming policy discussions such as the CBD's COP16 on the best way to

monitor progress towards biodiversity targets and effect change on the ground.

Acknowledgements:

This research was supported by the Natural Environment Research Council (NERC), United

Kingdom (NE/S007229/1) and WWF-UK as part of the Biome Health Project. DJI

acknowledges support from UK Research and Innovation (Future Leaders Fellowship

awarded to DJI., Grant ref: MR/W006316/1). We thank Dr Tim Newbold for discussion and

comments on an earlier draft of this manuscript. We thank all authors that have contributed

data and time to the PREDICTS project so that research like this can continue.

Declaration of Competing Interest

The authors declare that they have no known competing financial interests or personal

relationships that could have appeared to influence the work reported in this paper.

<u>References</u>

Allan, J.R. *et al.* (2019) 'Hotspots of human impact on threatened terrestrial vertebrates',
PLOS Biology. Edited by P. Jordano, 17(3), p. e3000158. Available at:
https://doi.org/10.1371/journal.pbio.3000158.

Blowes, S.A. *et al.* (2019) 'The geography of biodiversity change in marine and terrestrial
assemblages', *Science*, 366(6463), pp. 339–345. Available at:
https://doi.org/10.1126/science.aaw1620.

Bolker, B.M. *et al.* (2009) 'Generalized linear mixed models: a practical guide for ecology
and evolution', *Trends in Ecology & Evolution*, 24(3), pp. 127–135. Available at:
https://doi.org/10.1016/j.tree.2008.10.008.

Bowler, D.E. *et al.* (2020) 'Mapping human pressures on biodiversity across the planet
uncovers anthropogenic threat complexes', *People and Nature*, 2(2), pp. 380–394. Available
at: https://doi.org/10.1002/pan3.10071.

Brooks, T.M. *et al.* (2015) 'Harnessing biodiversity and conservation knowledge products to
track the Aichi Targets and Sustainable Development Goals', *Biodiversity*, 16(2–3), pp. 157–
174. Available at: https://doi.org/10.1080/14888386.2015.1075903.

Brown, J.H. (2014) 'Why are there so many species in the tropics?', *Journal of
Biogeography*, 41(1), pp. 8–22. Available at: https://doi.org/10.1111/jbi.12228.

Burnham, K.P., Anderson, D.R. and Huyvaert, K.P. (2011) 'AIC model selection and
multimodel inference in behavioral ecology: some background, observations, and
comparisons', *Behavioral Ecology and Sociobiology*, 65(1), pp. 23–35. Available at:
https://doi.org/10.1007/s00265-010-1029-6.

CBD, (Convention on Biological Diversity) (2020) *Update of the zero draft of the post-2020
global biodiversity framework*. CBD/POST2020/PREP/2/1., p. 9.

Ceballos, G., Ehrlich, P.R. and Dirzo, R. (2017) 'Biological annihilation via the ongoing
sixth mass extinction signaled by vertebrate population losses and declines', *Proceedings of
the National Academy of Sciences*, 114(30), pp. E6089–E6096. Available at:
https://doi.org/10.1073/pnas.1704949114.

Collen, B. *et al.* (2009) 'Monitoring Change in Vertebrate Abundance: the Living Planet
Index', *Conservation Biology*, 23(2), pp. 317–327. Available at:
https://doi.org/10.1111/j.1523-1739.2008.01117.x.

Dinerstein, E. *et al.* (2017) 'An Ecoregion-Based Approach to Protecting Half the Terrestrial
Realm', *BioScience*, 67(6), pp. 534–545. Available at: https://doi.org/10.1093/biosci/bix014.

Dinerstein, E. *et al.* (2019) 'A Global Deal For Nature: Guiding principles, milestones, and
targets', *Science Advances*, 5(4), p. eaaw2869. Available at:
https://doi.org/10.1126/sciadv.aaw2869.

Felipe-Lucia, M.R. *et al.* (2020) 'Land-use intensity alters networks between biodiversity,
ecosystem functions, and services', *Proceedings of the National Academy of Sciences*, p. 11.
Available at: https://doi.org/www.pnas.org/cgi/doi/10.1073/pnas.2016210117.

Gaston, K.J. (2000) 'Global patterns in biodiversity', *Nature*, 405(6783), pp. 220–227.
Available at: https://doi.org/10.1038/35012228.

Gibson, L. *et al.* (2011) 'Primary forests are irreplaceable for sustaining tropical
biodiversity', *Nature*, 478(7369), pp. 378–381. Available at:
https://doi.org/10.1038/nature10425.

Goldewijk, K.K. *et al.* (2011) 'The HYDE 3.1 spatially explicit database of human-induced
global land-use change over the past 12,000 years', *Global Ecology and Biogeography*,
20(1), pp. 73–86. Available at: https://doi.org/10.1111/j.1466-8238.2010.00587.x.

Green, E.J. *et al.* (2020) 'Below the canopy: global trends in forest vertebrate populations and
their drivers', *Proceedings of the Royal Society B: Biological Sciences*, 287(20200533).
Available at: http://dx.doi.org/10.1098/rspb.2020.0533.

Greenville, A.C. *et al.* (2018) 'Biodiversity responds to increasing climatic extremes in a
biome-specific manner', *Science of The Total Environment*, 634, pp. 382–393. Available at:
https://doi.org/10.1016/j.scitotenv.2018.03.285.

Hudson, L.N. *et al.* (2017) 'The database of the PREDICTS (Projecting Responses of
Ecological Diversity In Changing Terrestrial Systems) project', *Ecology and Evolution*, 7(1),
pp. 145–188. Available at: https://doi.org/10.1002/ece3.2579.

Ingram, D.J. *et al.* (2021) 'Targeting Conservation Actions at Species Threat Response
Thresholds', *Trends in Ecology & Evolution*, 36(3), pp. 216–226. Available at:
https://doi.org/10.1016/j.tree.2020.11.004.

IPBES (2019) *Global assessment report on biodiversity and ecosystem services of the
Intergovernmental Science-Policy Platform on Biodiversity and Ecosystem Services. E. S.
Brondizio, J. Settele, S. Díaz, and H. T. Ngo (editors)*. Bonn, Germany: IPBES secretariat.

Isaac, N.J.B. and Cowlishaw, G. (2004) 'How species respond to multiple extinction threats',
Proceedings of the Royal Society of London. Series B: Biological Sciences, 271(1544), pp.
1135–1141. Available at: https://doi.org/10.1098/rspb.2004.2724.

Isbell, F. *et al.* (2017) 'Linking the influence and dependence of people on biodiversity across
scales', *Nature*, 546(7656), pp. 65–72. Available at: https://doi.org/10.1038/nature22899.

IUCN (2021) *The IUCN Red List of Threatened Species. Version 2021-1*. Available at:
https://www.iucnredlist.org (Accessed: 17 May 2021).

Leung, B., Greenberg, D.A. and Green, D.M. (2017) 'Trends in mean growth and stability in
temperate vertebrate populations', *Diversity and Distributions*, 23(12), pp. 1372–1380.
Available at: https://doi.org/10.1111/ddi.12636.

Mace, G.M. *et al.* (2018) 'Aiming higher to bend the curve of biodiversity loss', *Nature*
Sustainability, 1(9), pp. 448–451. Available at: https://doi.org/10.1038/s41893-018-0130-0.

McDonald, R. *et al.* (2005) 'Species compositional similarity and ecoregions: Do ecoregion
boundaries represent zones of high species turnover?', *Biological Conservation*, 126(1), pp.
24–40. Available at: https://doi.org/10.1016/j.biocon.2005.05.008.

Moncrieff, G.R., Bond, W.J. and Higgins, S.I. (2016) 'Revising the biome concept for
understanding and predicting global change impacts', *Journal of Biogeography*, 43(5), pp.
863–873. Available at: https://doi.org/10.1111/jbi.12701.

Monsarrat, S., Jarvie, S. and Svenning, J.-C. (2019) 'Anthropocene refugia: integrating
history and predictive modelling to assess the space available for biodiversity in a human-
dominated world', *Philosophical Transactions of the Royal Society B: Biological Sciences*,
374(20190219). Available at: https://doi.org/10.1098/rstb.2019.0219.

Monsarrat, S. and Svenning, J.-C. (2022) 'Using recent baselines as benchmarks for
megafauna restoration places an unfair burden on the Global South', *Ecography*, 2022(4).
Available at: https://doi.org/10.1111/ecog.05795.

Murphy, S.J. (2021) 'Sampling units derived from geopolitical boundaries bias biodiversity
analyses', *Global Ecology and Biogeography*, 30(9), pp. 1876–1888. Available at:
https://doi.org/10.1111/geb.13352.

Myers, N. *et al.* (2000) 'Biodiversity hotspots for conservation priorities', *Nature*, 403(6772),
pp. 853–858. Available at: https://doi.org/10.1038/35002501.

Newbold, T. *et al.* (2013) 'Ecological traits affect the response of tropical forest bird species
to land-use intensity', *Proceedings of the Royal Society B: Biological Sciences*, 280(1750), p.
20122131. Available at: https://doi.org/10.1098/rspb.2012.2131.

Newbold, T. *et al.* (2015) 'Global effects of land use on local terrestrial biodiversity', *Nature*,
520(7545), pp. 45–50. Available at: https://doi.org/10.1038/nature14324.

Newbold, T. (2015) 'StatisticalModels'. Available at:
https://github.com/timnewbold/StatisticalModels.

Newbold, T. (2018) 'predictsFunctions: Functions for reading and processing the PREDICTS
data'.

Newbold, T. *et al.* (2018) 'Widespread winners and narrow-ranged losers: Land use
homogenizes biodiversity in local assemblages worldwide', *PLOS Biology*. Edited by H.
Morlon, 16(12), p. e2006841. Available at: https://doi.org/10.1371/journal.pbio.2006841.

Newbold, T. *et al.* (2020) 'Tropical and Mediterranean biodiversity is disproportionately
sensitive to land-use and climate change', *Nature Ecology & Evolution* [Preprint]. Available
at: https://doi.org/10.1038/s41559-020-01303-0.

Olson, D.M. *et al.* (2001) 'Terrestrial Ecoregions of the World: A New Map of Life on
Earth', *BioScience*, 51(11), p. 933. Available at: https://doi.org/10.1641/0006-
3568(2001)051[0933:TEOTWA]2.0.CO;2.

Olson, D.M. and Dinerstein, E. (2002) 'The Global 200: Priority Ecoregions for Global
Conservation', *Annals of the Missouri Botanical Garden*, 89(2), pp. 199–224. Available at:
https://doi.org/10.2307/3298564.

Phillips, H.R.P., Newbold, T. and Purvis, A. (2017) 'Land-use effects on local biodiversity in
tropical forests vary between continents', *Biodiversity and Conservation*, 26(9), pp. 2251–
2270. Available at: https://doi.org/10.1007/s10531-017-1356-2.

Smith, J.R. *et al.* (2018) 'A global test of ecoregions', *Nature Ecology & Evolution*, 2(12),
pp. 1889–1896. Available at: https://doi.org/10.1038/s41559-018-0709-x.

Smith, J.R. *et al.* (2020) 'The biogeography of ecoregions: Descriptive power across regions
and taxa', *Journal of Biogeography*, 47(7), pp. 1413–1426. Available at:
https://doi.org/10.1111/jbi.13871.

Sodhi, N.S. *et al.* (2004) 'Southeast Asian biodiversity: an impending disaster', *Trends in
Ecology & Evolution*, 19(12). Available at: https://doi.org/doi:10.1016/j.tree.2004.09.006.

Sodhi, N.S. *et al.* (2009) 'A Meta-Analysis of the Impact of Anthropogenic Forest
Disturbance on Southeast Asia's Biotas', *Biotropica*, 41(1), pp. 103–109. Available at:
https://doi.org/10.1111/j.1744-7429.2008.00460.x.

Song, X.-P. *et al.* (2018) 'Global land change from 1982 to 2016', *Nature*, 560(7720), pp.
639–643. Available at: https://doi.org/10.1038/s41586-018-0411-9.

Srivathsa, A. *et al.* (2018) 'Substituting space for time: Empirical evaluation of spatial
replication as a surrogate for temporal replication in occupancy modelling', *Journal of
Applied Ecology*, 55(2), pp. 754–765. Available at: https://doi.org/10.1111/1365-2664.13005.

Stephenson, P.J. *et al.* (2015) 'Overcoming the challenges to conservation monitoring:
integrating data from *in-situ* reporting and global data sets to measure impact and
performance', *Biodiversity*, 16(2–3), pp. 68–85. Available at:
https://doi.org/10.1080/14888386.2015.1070373.

Stevens, G.C. (1989) 'The Latitudinal Gradient in Geographical Range: How so Many
Species Coexist in the Tropics', *The American Naturalist*, 133(2), pp. 240–256. Available at:
https://doi.org/10.1086/284913.

Suraci, J.P. *et al.* (2021) 'Disturbance type and species life history predict mammal responses
to humans', *Global Change Biology*, 00, pp. 1–14. Available at:
https://doi.org/10.1111/gcb.15650.

Tittensor, D.P. *et al.* (2014) 'A mid-term analysis of progress toward international
biodiversity targets', *Science*, 346(6206), pp. 241–244. Available at:
https://doi.org/10.1126/science.1257484.

Walker, L.R. *et al.* (2010) 'The use of chronosequences in studies of ecological succession
and soil development', *Journal of Ecology*, 98(4), pp. 725–736. Available at:
https://doi.org/10.1111/j.1365-2745.2010.01664.x.

WWF (2020) *Living Planet Report 2020: bending the curve of biodiversity loss*. Available at:
https://edepot.wur.nl/531235 (Accessed: 20 October 2020).

WWF (2022) *Living Planet Report 2022 - Building a nature positive society*. Gland,
Switzerland: WWF. Available at:
https://wwflpr.awsassets.panda.org/downloads/lpr_2022_full_report_1.pdf (Accessed: 17
October 2022).

Yang, X. *et al.* (2021) 'Threatened bird species are concentrated in regions with less
historical human impacts', *Biological Conservation*, 255, p. 108978. Available at:
https://doi.org/10.1016/j.biocon.2021.108978.

<u>Data Accessibility Statement</u>

All data used in this manuscript is downloaded from open-source locations. The PREDICTS
database can be downloaded from the Natural History Museum Data portal
(https://data.nhm.ac.uk/dataset/the-2016-release-of-the-predicts-database).
578
579   Geographical data used to delineate regional biome boundaries in this manuscript can be
580   downloaded from The Nature Conservancy's Geospatial Conservation Atlas
581   (https://geospatial.tnc.org/datasets/b1636d640ede4d6ca8f5e369f2dc368b/about).
582
583   All R code used to run analysis presented in this manuscript is accessible in a github
584   repository (not included here to maintain anonymity)
585
586   <u>Figures</u>
587

Fig 1. A map of regional biomes. Solid coloured areas represent biomes (see legend). Black lines separate out biogeographic realms. Black dots represent sites in the PREDICTS database used in our global model (Hudson et al 2014). Hashed coloured areas represent regional biomes that were not included in analysis due to data deficiency (see methods). Biome and biogeographic realm spatial data from (Olson *et al.*, 2001).

Fig. 2. Three case-studies of regional biomes: distribution of data. The PREDICTS

database was subset into studies from three biomes that included at least 25 data points in

each land use category from a minimum of three biogeographic realms. The figures show

ranges of regional biomes, coloured by realm, for (a) Tropical Forest, (b) Temperate Forest

and (c) Tropical grassland. Black dots represent a data point (record of species richness or

total abundance) from the PREDICTS dataset. Regional biome distribution data from (Olson

et al., 2001).

Fig 3. Change in Akaike's Information Criterion (ΔAIC) for Generalised Linear Mixed Models (GLMMs) predicting change in a global dataset of (a) species richness and (b) total abundance consistently suggests models including regional biome have highest support when modelling response to land-use change. Four GLMMs with differing fixed-effect structures containing Land Use (LU) interacting with Regional Biome (RB), Biome or Realm were compared using AIC values. A model with low AIC is considered to have higher support. ΔAIC is calculated as the difference in AIC from the lowest scoring model. The box and whisker plots represent a summary of ΔAIC values of 100 hold-out models, where each iteration removed 10% of studies at random as a test of model robustness. Importantly, for a given dataset in this hold-out analysis, LU:RB model had the lowest AIC in 100% of iterations. The PREDICTS dataset was subset to only include data from regional biomes with 25 data points per combination of regional biome and land-use type. Land Use is a discrete variable with Primary Vegetation, Secondary Vegetation, Plantation Forest, Pasture and Cropland as its categories (see Table S2 for descriptions).

Fig 4. Responses of biodiversity to land-use change are unique across tropical forest regional biomes. The results of GLMMs predicting the response of two biodiversity metrics across land-use types and land-use intensity, when compared to primary vegetation (PV) across tropical forest regional biomes. Responses of species richness (top row) and total abundance (bottom row) of plants and animals were predicted across land-use types (a, c) including secondary vegetation (SV), plantation forest (PF), pasture (Pa) and cropland (Cr). Responses were also predicted for two levels of land-use intensity: minimal (M) or intense (I) in secondary vegetation (SV) and agriculture (Agr) land use types. Each point represents mean prediction, with 75% confidence intervals (thick whiskers) and 95% confidence intervals (thin whiskers). Responses are considered significantly different from primary vegetation if the 95% confidence intervals do not pass 0 (dotted line).

Fig 5. Responses of biodiversity to land-use change are hard to predict across temperate forest regional biomes. The results of GLMMs predicting the response of two biodiversity metrics across land-use types and land-use intensity, when compared to primary vegetation (PV) across temperate forest regional biomes. Responses of species richness (top row) and total abundance (bottom row) of plants and animals were predicted across land-use types (a, c) including secondary vegetation (SV), plantation forest (PF), pasture (Pa) and cropland (Cr). Responses were also predicted for two levels of land-use intensity: minimal (M) or intense (I) in secondary vegetation (SV) and agriculture (Agr) land-use types. Each point represents mean prediction, with 75% confidence intervals (thick whiskers) and 95% confidence intervals (thin whiskers). To keep scales consistent, upper 95% CIs greater than 100 have been included as text instead. Responses are considered significantly different from primary vegetation if 95% confidence intervals do not pass 0 (dotted line).

Fig 6. Responses of biodiversity to land-use change are hard to predict across tropical grassland regional biomes. The results of GLMMs predicting the response of two biodiversity metrics across land-use types and land-use intensity, when compared to primary vegetation (PV) across tropical grassland regional biomes. Responses of species richness (top row) and total abundance (bottom row) of plants and animals were predicted across land-use types (a, c) including secondary vegetation (SV), plantation forest (PF), pasture (Pa) and cropland (Cr). Responses were also predicted for two levels of land-use intensity: minimal (M) or intense (I) in secondary vegetation (SV) and agriculture (Agr) land use types. Each point represents mean prediction, with 75% confidence intervals (thick whiskers) and 95% confidence intervals (thin whiskers). To keep scales consistent, upper 95% CIs greater than 100 have been included as text instead. Responses are considered significantly different from primary vegetation if 95% confidence intervals do not pass 0 (dotted line).