Transition from free-living cyanobacterium to endosymbiont enables adaptations favoring nitrogen fixation in a photosynthetic eukaryote.

Solène L.Y. Moulin [1], Sarah Frail [2], Thomas Braukmann[1,2], Jon Doenier[2], Ellen Yeh* [1,3]

[1] Department of Pathology, Stanford School of Medicine, Stanford, California, USA

[2] Department of Biochemistry, Stanford School of Medicine, Stanford, California, USA

[3] Department of Microbiology & Immunology, Stanford School of Medicine, Stanford, California, USA

* Corresponding author: 299 Campus Dr, Fairchild 317, Stanford, CA 94305; ellenyeh@stanford.edu

Author Contributions: S.L.Y.M., and E.Y. designed research; S.L.Y.M., S.F., and T.B. performed research; S.L.Y.M., S.F., T.B. and J.D analyzed data; and S.L.Y.M., and E.Y. wrote the paper.

Competing Interest Statement: The authors declare no competing interest.

Keywords: Endosymbiosis, Nitrogen fixation, cyanobacteria, diatom.

Abstract

Nitrogen is a limiting nutrient for photosynthetic productivity. Yet biological nitrogen fixation which transforms abundant atmospheric nitrogen to bioavailable ammonia is largely restricted to bacteria. *Epithemia* spp. diatoms contain obligate, nitrogen-fixing endosymbionts, or "diazoplasts", derived from cyanobacteria. They represent rare photosynthetic eukaryotes that have successfully coupled oxygenic photosynthesis with oxygen-sensitive nitrogenase activity. Here, we report a newly-isolated species, *E. clementina*, as a model to investigate endosymbiotic acquisition of this critical biological function. To detect the metabolic changes associated with endosymbiotic specialization, we compared nitrogen fixation, nitrogen utilization, and their regulatory pathways in the *Epithemia* diazoplast with its close free-living cyanobacteria relative, *Crocosphaera subtropica*. Unlike *C. subtropica*, we show that nitrogenase activity in the diazoplast is concurrent with, and even dependent on, host photosynthesis. In contrast to its expanded nitrogen-fixing activity, the diazoplast has diminished ability to utilize alternative nitrogen sources. Finally, while key regulators of nitrogen metabolism are retained in the diazoplast genome, we observed a conserved negative feedback regulation of nitrogen fixation to ammonium, but paradoxical nitrogen assimilation responses in the diazoplast compared with *C. subtropica*. Our results suggest key adaptations that facilitated the transition from free-living to endosymbiotic specialist diazotroph, including import of host carbohydrates, streamlined carbon catabolism favoring high respiratory rates, and altered nitrogen regulation to favor nitrogen transfer to the host. Elucidating how photosynthesis and nitrogen fixation are coupled in a eukaryotic cell has important implications for bioengineering nitrogen-fixing crop plants for sustainable agriculture.

Main Text

Introduction

Nitrogen composes 78% of the atmosphere; however, it is present in the form of dinitrogen (N_2), an inert gas, which cannot be directly incorporated into organic molecules such as DNA or protein. Nitrogen fixation, a critical biological reaction, is the reduction of N_2 into bioavailable ammonia catalyzed by the enzyme nitrogenase. Biological nitrogen fixation is largely restricted to certain genera of bacteria. Nitrogen availability determines photosynthetic productivity in both natural environments and agriculture, such that the activity of these nitrogen-fixing organisms, or

diazotrophs, are a critical source of bioavailable nitrogen for plants and other photosynthetic organism. To bypass the dependence of food production on biological nitrogen fixation, modern agriculture uses energy from fossil fuels to chemically fix nitrogen and produce nitrogen-rich fertilizer, which is estimated to feed 45% of the world population (1). However, synthetic fertilizers have significant negative environmental consequences including greenhouse gas emissions, soil degradation, and coastal dead zones (eutrophication) with much of the applied nitrogen polluting waterways (2–4).

While a nitrogen-fixing plant might seem an obvious solution, examples of nitrogen-fixing photosynthetic eukaryotes are rare. Attempts to introduce nitrogen fixation into plant cells by expressing nitrogenase genes have met significant challenges (5, 6). On one hand, coupling nitrogen fixation and photosynthesis in a single cell is energetically advantageous. Nitrogenase requires 16 ATP and 8 NADPH molecules to fix a single N_2 molecule, energy that can be supplied by the photosynthetic electron transfer chain. On the other hand, nitrogenase is also exquisitely sensitive to irreversible inactivation by oxygen which is produced during photosynthesis, making photosynthesis and nitrogen fixation biochemically incompatible. Instead, throughout evolution, these two biologically critical reactions have been coupled via symbiosis between nitrogen-fixing bacteria and diverse photosynthetic organisms (7).

Amongst these diverse symbiotic interactions, a few rare microalgae have acquired the cellular ability to fix nitrogen via endosymbioses of diazotrophic cyanobacteria. *Crocosphaera* spp. cyanobacteria (previously named *Cyanothece*) appear to be a privileged endosymbiotic partner (8). Several microalgae, the diatoms *Epithemia* spp. and *Climacodium frauenfeldianum* (9) as well as the coccolithophore *Braarudosphaera bigelowii* (10), have permanent endosymbionts closely related to the *Crocosphaera* genus, specifically *Crocosphaera subtropica* (*Cyanothece* sp. ATCC51142). As an endosymbiont within these photosynthetic eukaryotes, these cyanobacteria lose their photosynthetic function and become specialized for nitrogen fixation. The remarkable ability of *Crocosphaera* cyanobacteria to transition from free-living diazotrophic phototroph to endosymbiotic diazotrophic heterotroph in multiple eukaryotic lineages suggests pre-existing traits that predispose them to endosymbiotic acquisition. For example, *C. subtropica* is able to grow mixotrophically using carbon substrates which may favor its success as an endosymbiotic partner (11, 12).

Crocosphaera's partnership with *Epithemia* diatoms has been particularly successful. Diverse *Epithemia* species containing endosymbionts are widespread in freshwater habitats globally and have recently been found in marine environments as well (13). The endosymbionts, first described as "spheroid bodies" within *Epithemia gibba* (formerly *Rhopalodia gibba*), were shown to encode an intact *nif* gene cluster and demonstrate nitrogen fixation activity enabling the diatom to grow in nitrogen-depleted environments (14–17). Hereafter, we will refer to these nitrogen-fixing endosymbionts as diazoplast, combining "diazo-" meaning 'relating to the group N_2 and "-plast" meaning 'living substance'. Sequencing of diazoplast genomes from two *Epithemia* species revealed significant genome reduction, including loss of most of the genes involved in photosynthesis and an incomplete TCA cycle implying that diazoplasts are dependent on the diatom for metabolic needs(17–19). Consistent with their being obligate endosymbionts, diazoplasts have never been observed outside their diatom, and isolated diazoplasts cannot be grown in laboratory cultures outside the host cells. Diazoplasts are not only vertically transmitted during cell division but show uniparental inheritance during sexual reproduction, demonstrating robust mechanisms for inheritance of the diazoplast (20). Although *Epithemia* spp. are morphologically diverse, their diazoplasts are believed to derive from a single endosymbiotic event with a cyanobacteria from *Crocosphaera* genus (Nakayama *et al*, 2011)). Dated at 12 Mya (21), this is the most recent obligate primary endosymbiosis documented to date. Finally, several *Epithemia* strains isolated from environmental samples have been successfully grown in laboratory cultures (22). Overall, the diazoplasts of *Epithemia* diatoms are an ideal case study for i) the acquisition of nitrogen fixation function by a photosynthetic eukaryote and ii) investigating evolution associated with organellogenesis (18).

Though *Epithemia*'s diazoplasts were described nearly half a century ago, little is known about this functionally unique and ecologically widespread endosymbiotic interaction. So far, the

most detailed molecular studies have focused on the extensive genome reduction in the diazoplast suggestive of unprecedented metabolic dependence on the host cell (17, 18). While the pattern of reduction and retention in the diazoplast genome suggests new cell biology, such as the energetic coupling of diatom photosynthesis to diazoplast nitrogen fixation, functional studies to support these hypotheses have been limited. Herein, we systematically compare nitrogen fixation and nitrogen assimilation, as well as their regulatory pathways in the *Epithemia* diazoplast with a close free-living relative. The diazoplast was investigated in a newly isolated *Epithemia* species, *E. clementina*, while *C. subtropica* served as a proxy for the ancestral free-living cyanobacteria. Our studies reveal key adaptations accompanying the transition from free-living to endosymbiont that allowed diazoplasts to become specialized for nitrogen fixation in a photosynthetic eukaryote.

Results

Epithemia clementina is a new model system to study evolution of diazoplasts

Epithemia spp. diatoms are widely distributed in freshwater environments globally, and laboratory cultures have previously been established from isolates collected in the US, Germany, and Japan. Of these cultures, only one *Epithemia* strain has been deposited in a culture collection (*Rhopalodia gibba*, FD213, UTEX); however, we were unable to establish cultures from this isolate. Therefore, to obtain a strain of *Epithemia* for laboratory studies, we collected water samples from a freshwater stream near Butano State Park, CA. The samples were serially diluted over a period of 8 weeks in nitrogen- and carbon-free media to isolate diazotrophic phototrophs. Most isolates were dominated by green-colored filamentous cyanobacteria, likely *Nostoc* spp. that are ubiquitous phototrophic diazotrophs. However, a few isolates were visibly brown and consisted of diatoms resembling described *Epithemia* spp. that contained endosymbionts. Consistent with their unique ability to fix nitrogen, these diatoms were the only eukaryotic algae that were isolated from passage in nitrogen-free media.

Under light microscopy, the isolated alga was approximately 20 µm long and resembled a wedge of an orange and was thus named *Epithemia clementina* (Fig. 1*A*). Like other members of its genus, it is a pennate diatom that comprises two valves. It has an asymmetrical growth of its griddle bands and shows the presence of marginal raphes on its concave side. Morphologically, the shape and pattern associated with the valve does not match other described *Epithemia* species (Fig. S1*A*). Similar to other *Epithemia* diatoms, *E. clementina* contains endosymbionts appearing as spheroid bodies within the diatom both under light microscopy and by fluorescence microscopy after staining for nucleic acids (Fig. 1 *A* and *B*). Nucleic acid staining also revealed the presence of bacteria attached to the frustule of *E. clementina.* Consistent with previous characterization of diazoplasts, the endosymbionts were colorless in light microscopy and did not have any signal in fluorescence microscopy, indicating a lack of photosynthetic pigments (Fig. 1*C* and *D,* Fig. S1*B*).

The morphologic classification was confirmed by molecular phylogeny. Phylogenetic analysis based on a selection of conserved genes from diatoms confirmed that this undescribed species is a member of the *Epithemia* genus, most closely related to *Epithemia iriomotensisa* (Fig. 1*E*) (23). To identify the origin of the endosymbiont, we also performed a phylogenetic analysis of conserved cyanobacterial genes which showed that the *E. clementina* endosymbiont is of the same origin as "spheroid bodies" previously found in *Epithemia* spp. (Fig. 1*F*) (24). The diazoplast genome was assembled from long-read sequencing (Nanopore) polished by short-read sequencing (Illumina) of a metagenomic sample containing *E. clementina* and associated bacteria. The genome confirmed that the endosymbiont was indeed a diazoplast, containing an intact *nif* cluster as observed in other *Epithemiaceae* (17, 18). Because we had observed bacteria closely associated with *E. clementina* in the monoalgal cultures that could not be removed by dilution, detergent, or antibiotic treatment, we took advantage of our metagenomic sequencing data to characterize the bacterial community of *E. clementina*. 62 metagenome-assembled genomes (MAGs) from taxonomically diverse bacteria were obtained in addition to the diazoplast

genome, (Fig. S2, Table S1). None of these were assigned as phototrophs, though the presence of *nif* genes was detected in 2 MAGs related to *Methyloversatilis* sp. RAC08 (~90% identity) and *Hyphomicrobium* sp. DMF-1 (~75-80% identity). Based on sequencing depth, the diazoplast accounted for 19.2% of bacterial genomes detected, compared to <0.3% for the two nif-containing MAGs, suggesting that it is the primary source of nitrogenase activity in our cultures.

Diazoplast has lost temporal regulation of nitrogenase

Having established laboratory cultures of *E. clementina* to study its diazoplast, we also obtained cultures of *C. subtropica,* which is closely related to the diazoplast, to serve as a proxy for the ancestral free-living state of the diazoplast. We first compared the diurnal regulation of nitrogen fixation activity in free-living *C. subtropica* and *E. clementina*'s diazoplast. Nitrogenase is exquisitely sensitive to deactivation by oxygen. As a result, cyanobacteria like *C. subtropica* that can perform both photosynthesis and nitrogen fixation have evolved mechanisms to separate nitrogenase activity and oxygenic photosynthesis either spatially or temporally. Studies of *C. subtropica* demonstrated tight transcriptional and post-translational regulation of nitrogenase, such that nitrogen fixation only occurs at night when photosynthesis is inactive (25–28). Using an acetylene reduction assay (ARA), we confirmed that the nitrogenase activity of *C. subtropica* was restricted to nighttime with a strong diel pattern that peaked at 4 hours into the night (Fig. 2*A*) (27, 29, 30).

Because the diazoplast is no longer photosynthetic, we wondered if its nitrogenase activity was still temporally regulated. Studies of other *Epithemia* spp. have demonstrated daytime nitrogenase activity concurrent with photosynthesis, indicating remodeling of the temporal regulation (13–15). Consistent with these previous reports and in contrast to free-living *C. subtropica*, *E. clementina* showed nitrogenase activity throughout the day-night cycle with no clear diel pattern (Fig. 2*A*). In *C. subtropica,* immunoblotting for the NifH subunit of nitrogenase showed a diel pattern matching the nitrogenase activity (29, 30), while in *E. clementina*, NifH protein abundance was stable throughout the day-night cycle (Fig. 2*B* and *C*). We also assessed the abundance of HupL, the large subunit of the uptake hydrogenase which recycles hydrogen produced by nitrogenase and is often co-expressed with *nif* genes (31). HupL protein was also present exclusively at night timepoints in *C. subtropica*, while HupL protein abundance was stable in the diazoplast (Fig. 2*B*). Finally, we performed quantitative RT-PCR to detect transcript levels of the NifK subunit of nitrogenase and HupS, the small subunit of the uptake hydrogenase. Both *nifK* and *hupS* showed tightly-regulated nighttime expression in *C. subtropica* that was absent in the diazoplast (Fig. 2*D*). Taken together, the activity, protein abundance, and transcript levels of nitrogenase demonstrate that the global diurnal regulation of nitrogen fixation in free-living *C. subtropica* is lost in the diazoplast of *E. clementina*.

Diazoplast nitrogen fixation is coupled to host photosynthesis and carbohydrate storage

Biological nitrogen fixation is an energy-intensive reaction with nitrogenase consuming 16 ATP and 8 NADPH molecules for each molecule of N_2 converted to 2 ammonia molecules. Therefore, the metabolism of diazotrophs must be optimized to supply large quantities of both ATP and reducing equivalents. In free-living *Crocosphaera*, the energy for nitrogen fixation is provided by photosynthesis, which converts sunlight into chemical energy in the form of glucose. However, because photosynthesis is segregated temporally from nitrogen fixation in *Crocosphaera*, glucose produced during the day is stored in glycogen granules and catabolized at night when nitrogenase is active (27, 30). A global transcriptomic analysis of the diurnal cycle in *C. subtropica* confirmed these cellular findings, showing photosynthesis and glycogen biosynthesis genes upregulated during the day and glycogen degradation and nitrogen fixation genes (*nif*) upregulated at night (32). Finally, pathways that consume glucose, i.e. glycolysis, the oxidative pentose phosphate (OPP) pathway, and the tricarboxylic acid (TCA) cycle, were also upregulated at night. Overall, these observations suggested a model whereby glucose produced by photosynthesis is stored as glycogen, which is catabolized at night to provide ATP and reducing equivalents for nitrogenase activity.

Because the diazoplast no longer performs photosynthesis, its nitrogenase is necessarily fueled by the products of host photosynthesis, i.e. photosynthate. To investigate the metabolic coupling between host photosynthesis and diazoplast nitrogenase activity, ARAs were performed under light, dark, and light + an inhibitor of photosystem II, 3-(3,4-dichlorophenyl)-1,1-dimethylurea (DCMU). During the day, inhibition of the host photosynthetic activity by dark condition or DCMU treatment reduced nitrogenase activity by 60% (Fig. 3*A*). At night, host photosynthetic activity by addition of light provided little to no benefit for nitrogenase activity (Fig. 3*A*). These results indicate that photosynthates are directly shuttled from the host chloroplast to the diazoplast during the day, while nighttime nitrogenase activity still relies on carbohydrate storage. The new ability of daytime nitrogen fixation, concurrent with photosynthesis, permits a direct use of host photosynthate without reliance on carbohydrate storage.

We next asked what the source of energy was for nighttime nitrogen fixation in the diazoplast. Despite gene reduction in glycogen metabolism pathways, the diazoplast genome encodes at least one copy of each gene required for glycogen biosynthesis and degradation, suggesting these pathways are functional and important for diazoplast metabolism (Fig. 3*B,* Table S2). Even non-diazotrophic cyanobacteria commonly use glycogen as storage for excess photosynthate produced during the day that is respired at night (33). Therefore, we hypothesized that the diazoplast still accumulates glycogen as a storage form of host photosynthate that fuels nighttime nitrogen fixation activity. In this case, we expected that glycogen biosynthesis and degradation pathways would be regulated in a diurnal pattern. We measured the daytime and nighttime expression level of genes related to glycogen biosynthesis (glycogen synthase, glgA) and glycogen degradation (glycogen phosphorylase, glgP) in the *E. clementina* diazoplast and *C. subtropica* by RT-qPCR. Expression levels of *glgA* did not show significant variation between day and night timepoints in either *E. clementina* or *C. subtropica* (Fig. 3*C*). Expression levels of *glgP* was upregulated by over 10-fold at night compared to the day in *C. subtropica*, consistent with the catabolism of stored glycogen to provide energy for nighttime nitrogen fixation in this organism. In contrast, no diel variation of *glgP* was seen in the diazoplast. In fact, *glgP* expression, normalized to *nifK* expression, was consistently low in the diazoplast and comparable to daytime *glgP* expression levels in *C. subtropica* when the glycogen degradation pathway is effectively inactive (Fig. 3*C*). Overall, the low expression level and loss of diurnal regulation of the glycogen degradation enzyme, *glgP*, indicates that glycogen is not the source of energy for nitrogenase in the diazoplast, even during nighttime. Consistent with our results, glycogen granules were not detected in electron microscopy of diazoplasts from *Epithemia* spp. (14, 34). Instead, glycogen pathways may be retained in the diazoplast for other purposes or under different conditions. Since the diazoplast genome does not encode other pathways for carbohydrate storage, most likely, nighttime nitrogen fixation in the diazoplast requires mobilization of host storage carbohydrates. Taken together, carbohydrate utilization in the diazoplast demonstrates tight coupling between host and diazoplast metabolism beyond that expected to support daytime nitrogen fixation.

Carbon catabolism is dependent on the OPP pathway in the diazoplast
After examining the potential sources of carbohydrate for daytime and nighttime nitrogen fixation in the diazoplast, we turned our attention to carbon catabolism in the diazoplast to convert host-derived carbohydrates into ATP and reducing equivalents. *C. subtropica* has been shown to rely on glycolysis, the OPP pathway, and the TCA cycle for carbon catabolism. However, the OPP pathway is the only complete pathway in diazoplasts. Indeed, many of the TCA cycle genes and the gene encoding the "gatekeeper" enzyme of glycolysis, phosphofructokinase (PFK), are missing from the diazoplast genomes (17) (Fig. 3*B,* Table S2). Instead, in heterocysts, specialized nitrogen-fixing cells found in filamentous cyanobacteria, nitrogen fixation activity was shown to be highly dependent on glucose-6-phosphate dehydrogenase (G6PD), a key enzyme of the OPP pathway (35, 36). The dependence on OPP was proposed to maximize production of reducing equivalents for oxidative phosphorylation that both generates ATP and consumes oxygen. To investigate the role of the OPP pathway in diazoplast metabolism, we measured the expression of *zwf* which encodes G6PD in the diazoplast compared with *C. subtropica.* The

expression level of *zwf* in *E. clementina* was 2.5-3x higher than that of *C. subtropica* under nitrogen-fixing conditions, despite comparable *nifK* expression (Fig. 3*D* and 2*D*). As expected, *zwf* expression did not exhibit diel variation in the diazoplast. Consistent with the loss of complete glycolysis and TCA pathways, the moderate increase of relative *zwf* expression suggests that OPP is the main pathway for carbon catabolism in the diazoplast (Fig. 3*B*). Because G6PD utilizes glucose-6-phosphate as its substrate, the dependence on the OPP pathway in the diazoplast indicates that at least one host-derived carbohydrate is a hexose. Similar to heterocysts, carbon catabolism in the diazoplast has been streamlined to support nitrogen fixation, demonstrating another adaptation associated with transition to an endosymbiotic diazotroph.

The diazoplast responds to exogenous ammonia yet paradoxically represses ammonium uptake
Because nitrogen fixation is energy intensive, diazotrophs show feedback regulation repressing nitrogenase activity when exogenous nitrogen, either in the form of ammonia or nitrate, is available. The utilization of these nitrogen sources is dependent on several gene products (Fig. 4*A*; Table S2). Ammonia can diffuse across membranes in its neutral form or be imported in its protonated form via an ammonium transporter (Amt). Intracellular ammonia is then assimilated into amino acids by glutamine synthetase (GS) and glutamine oxoglutarate aminotransferase (GOGAT). Utilization of nitrate is dependent on uptake via an ABC transporter followed by intracellular conversion to ammonia by nitrate and nitrite reductases. Consistent with an ancestral ability to utilize ammonia and nitrate, these pathways are conserved in both *C. subtropica* and a related cyanobacterium, *Rippkaea orientalis* (*Cyanothece* sp*. PCC8801*). In contrast, only the genes for ammonia uptake and assimilation are retained in the 3 diazoplast genomes from *E. clementina*, *E. turgida, and R. gibberula* but those encoding the nitrate ABC transporter, nitrate reductase, and nitrite reductase are missing (Table S2).

Although *C. subtropica* is known to grow on nitrate or ammonium, its response to nitrogen repletion has not been documented (12, 27). We measured nitrogenase activity in *C. subtropica* in the presence of exogenous ammonium and nitrate at low (µM) and high (mM) concentrations. Neither 100 µM ammonium nor 120 µM nitrate suppressed nitrogenase activity. However, higher mM concentrations fully repressed nitrogenase activity (Fig. 4*B*). This repression was reflected in loss of *nifK* expression by RT-qPCR (Fig. 4*C*) and of NifK and HupL proteins by immunoblot (Fig. 4*D*). In *E. clementina*, repression of nitrogenase activity was observed at 100µM ammonium but unaffected by 120 µM nitrate (Fig. 4*B*). Ammonium repletion caused a 70% decrease in *nif* gene expression with a 2x decrease in NifK protein abundance as well as a decrease in HupL abundance, which accumulated as a higher molecular weight apoprotein (Fig. 4 *C* and *D*). These effects on nitrogen fixation were not observed with nitrate repletion, confirming the loss of nitrate utilization in the diazoplast. Consistent with a restricted ability to utilize nitrogen sources, diazoplast genomes are also missing functional genes for glutaminase and glutamate dehydrogenase required to catabolize amino acids for ammonia (Fig. 4*A*, Table S2). Curiously, the diazoplast was more sensitive to ammonium repletion, repressing nitrogenase activity at a lower ammonium concentration than *C. subtropica*.

Along with repression of nitrogen fixation activity, nitrogen repletion typically upregulates uptake and assimilation pathways to maximize the ability to utilize exogenous nitrogen sources. Indeed, we observed that ammonium repletion increased *amt* expression by over 2x in *C. subtropica* (Fig. 4*C*). In the absence of exogenous nitrogen sources, the baseline expression of *amt* (normalized to *nifK* expression) was 10x lower in the diazoplast than in *C. subtropica* (Fig. 4*C*). In contrast to upregulation observed in *C. subtropica*, ammonium repletion further decreased *amt* expression in the diazoplast. This paradoxical repression of *amt* may have evolved to diminish ammonium uptake into the diazoplast, favoring host nitrogen transfer. Altogether, the increased sensitivity to nitrogen fixation repression combined with a diminished ability to utilize nitrogen sources in the diazoplast may optimize the host cell's nitrogen resources.

Conserved regulatory genes show altered responses to exogenous ammonium

So far, we had observed significant integration of diazoplast and host nitrogen metabolism. Two models for regulation could account for our observations: one, the diazoplast retains cyanobacterial gene regulatory networks and senses its metabolic status (i.e. ammonia, carbon) for its response. Two, regulatory genes are transferred to the host diatom so it can directly control nitrogen metabolism in the diazoplast. Several transcription factors involved in nitrogen metabolism have been identified in cyanobacteria (Fig. 5*A*): NtcA is a global regulator of nitrogen metabolism conserved in both diazotrophic and nondiazotrophic cyanobacteria (37–40). It senses 2-oxoglutarate levels and phosphorylated P-II protein dependent on ammonia levels, effectively integrating information about the C:N balance of the cell (41–46). An important target of NtcA in diazotrophic cyanobacteria is CnfR (originally PatB), the master regulator of *nif* genes (47–49). This NtcA-CnfR regulatory network was identified in *C. subtropica* and in the diazoplast genome (Table S2). However, while the NtcA recognition sequence (TGTA-N_8-TACA) can be identified upstream of *C. subtropica CnfR* (-449 bp to -434 bp) (38, 49), it is absent from the promoter region of *CnfR* in the diazoplast. To determine whether the PII-NtcA-CnfR cascade still regulates the diazoplast *nif* cluster, we measured the expression of these key regulatory factors in nitrogen-depleted and -repleted conditions. The expression of *pII* and *ntcA* did not vary with ammonia repletion in either *C. subtropica* or *E. clementina*. While *ntcA* had a similar expression level between the two, *pII* showed a 6-10-fold lower expression level in the diazoplast. *cnfR* expression showed a clear 60% decrease upon ammonium repletion in the diazoplast, with a similar 80% decrease upon nitrate or ammonium repletion in *C. subtropica* (Fig. 5 *B* and *C*). The conserved responses in nitrogen fixation activity and its gene regulation between *C. subtropica* and the diazoplast suggest that the NtcA-CnfR regulation of the *nif* cluster has been retained in the diazoplast to activate nitrogen fixation activity in N-depleted conditions.

NtcA also regulates the nitrogen-regulated response regulator, NrrA. This transcription factor of the OmpR family coordinates nitrogen and carbon reserves (50). A *nrrA* homolog was identified in *C. subtropica* and the diazoplast genome (Table S2). Of the transcription factors that we tested, NrrA gave the most unexpected response: upon ammonia repletion, *nrrA* was upregulated in *C. subtropica* but downregulated in *E. clementina* (Fig. 5 *B* and *C*). Unfortunately, regulation of nitrogen metabolism has not been investigated in *C. subtropica,* and NrrA has diverse gene targets in cyanobacteria, so it is unclear what downstream effects result from the altered *nrrA* response. An *in silico* study, conducted in 15 different cyanobacteria, characterized target sequences of NrrA and experimentally validated them in *Synechocystis* sp. PCC 6803 (51). The proposed NrrA regulon in *C. subtropica* and *R. orientalis* included genes involved in glycogen catabolism, central carbon metabolism, arginine biosynthesis, and protein degradation. We manually screened the promoter region of the proposed targets and confirmed the presence of a potential NrrA binding sequence in the promoter region of *glgP, argG*, and *nrrA* itself in the diazoplast genomes of all *Epithemia* spp. (Fig. S3). The conservation of the binding sequences suggests that the NrrA regulon is conserved between *C. subtropica* and the diazoplast, but its response to exogenous nitrogen is altered upstream of *nrrA*. Once again, the paradoxical response of *nrrA* in the diazoplast compared to *C. subtropica* indicates key adaptations in nitrogen assimilation and storage that accompanied the endosymbiotic transition.

Discussion

Metabolic exchange is often the underlying currency of symbiotic interactions and a strong driver of evolution. Our results demonstrate that the acquisition of nitrogen fixation by *Epithemia* diatoms has led to global metabolic rearrangements in the diazoplast and a level of metabolic integration not previously seen in other diazotrophic symbioses. Whereas *Crocosphaera* cyanobacteria, quintessential generalists, are free-living phototrophs that perform nitrogen fixation exclusively at night, the diazoplast has evolved into a nitrogen-fixing specialist and obligate heterotroph endosymbiont. In reviewing our key findings below, we discuss potential mechanisms for the metabolic changes and how these changes may be tailored to support nitrogenase activity.

Mechanisms enabling daytime nitrogen fixation. The first notable change we observed was daytime nitrogen fixation activity in the diazoplast. We observed continuous nitrogen fixation activity in *E. clementina*, while recently discovered marine *Epithemia* species, *E. pelagica* and *E. catenate*, showed nitrogen fixation activity over the daytime with a hiatus of activity at night, a reversed day-night pattern compared to *C. subtropica* (13). This lax diurnal regulation is in stark contrast to *C. subtropica*, in which 30% of genes, mostly associated with photosynthesis and nitrogen fixation, show circadian expression (compared to just 2-7% of genes in the non-diazotrophic cyanobacterium, *Synechocystis* 6803) (32, 52–54). Cyanobacterial circadian clock relies on the 3 *kia* genes and *C. subtropica* has 2 copies of each gene while the diazoplast retained 1 copy of each (Table S2) (55). It is not clear how the loss of redundancy would affect circadian rhythm. Alternatively, the loss of redox signals resulting from photosynthetic activity that serve as environmental cues to synchronize light and dark periods may weaken the circadian cycle. Finally, we hypothesize that the incomplete TCA cycle in the diazoplast may lead to accumulation of 2-oxoglutarate, which is both a substrate for GS-GOGAT-mediated ammonia assimilation and a signaling metabolite that binds to NtcA and PII (Fig. 4*A*). Its accumulation would signal a high C:N ratio in the diazoplast, a metabolic state that activates nitrogen fixation. We propose that accumulation of 2-oxoglutarate could be a mechanism whereby nitrogen fixation is temporally deregulated. Regardless of the mechanism, the consistent shift toward daytime nitrogen fixation in all diazoplasts indicates a benefit of direct usage of photosynthate to fuel nitrogen fixation. The cyanobacterium *Trichodesmium* is a rare example known to perform nitrogen fixation concurrent with oxygenic photosynthesis, potentially benefitting from the absence of glycogen buildup during the day to maintain buoyancy in the water column (56, 57).

What host-provided carbohydrate(s) is utilized by the diazoplast? Second, we observed that daytime nitrogen fixation was dependent on photosynthetic activity while nighttime nitrogen fixation was no longer associated with glycogen storage and degradation, suggesting the diazoplast relies entirely on carbohydrates imported from the host. Though photosynthetic, some cyanobacteria have been observed to grow mixotrophically on exogenous carbons, usually glucose, fructose, and/or sucrose. Two transporters, GlcP and FrtABC, have been associated with glucose and fructose uptake, respectively, in cyanobacteria, but orthologs are not detected in *C. subtropica* or diazoplast genomes. Similarly, utilization of the disaccharide sucrose requires an invertase to hydrolyze it into glucose and fructose, but no invertase has been detected in *C. subtropica* or diazoplast genomes. Instead, *C. subtropica* shows robust growth on glycerol as a carbon source and utilizes glucose after a prolonged period of adaptation (11, 12). The transporters that allow *C. subtropica* to readily uptake glycerol and adapt to glucose uptake are unknown, though potential genes annotated as sugar ABC transporters are identified in both genomes. More detailed investigations of mixotrophic growth in *C. subtropica* will be needed to determine whether pre-existing transporter gene(s) were utilized to access host carbohydrates. An alternative model is that the diazoplast acquired the ability to uptake host carbohydrate(s), since we detected transporter genes present in all diazoplast genomes that were absent from free-living *Crocosphaera* genomes. Further investigation is needed, but these could be exciting candidates. We also considered the possibility that, instead of sugar compounds, ATP and NADPH could be directly imported into the diazoplast. However, none of the transporters known in mitochondria or chloroplasts were identified in diazoplast genomes, and, as will be discussed in the next section, the upregulation in OPP suggests the import of carbon compounds from the host that are metabolized through this pathway.

The OPP pathway provides reducing power and protection from oxygen. The third change we noted was an incomplete set of glycolysis and TCA cycle genes and likely dependence on an upregulated OPP pathway for carbon catabolism. While multiple enzymes in the TCA cycle were missing such that it was unlikely to be reconstituted, only PFK, the enzyme responsible for the first committed step of glycolysis, is absent from the diazoplast genome compared to *C. subtropica*. Loss of PFK is relatively common in cyanobacteria; however, we also considered the

possibility that the PFK gene may have transferred to the diatom nucleus and may be imported. As we have yet to report the *E. clementina* genome, we cannot rule this out. Instead, we compared the energetic output of OPP versus glycolysis feeding into the incomplete TCA cycle with 2-oxoglutarate as the final product. With initial flux through the OPP pathway, the overall reaction converting 2 molecules of glucose 6-phosphate into 2-oxoglutarate produces 16 NADH, 3 ATP and 7 CO_2. Glycolysis starting from the same 2 molecules of glucose 6-phosphate generates two 2-oxoglutarate, 8 NADH, 4 ATP, and 2 CO_2. Compared to glycolysis, the OPP pathway favors production of reducing equivalents which can then enter the respiration chain at the expense of fixed carbon. Electron micrographs of the *E. gibba* diazoplast show the presence of thylakoid membranes (14, 34). Unlike plant chloroplasts, both photosynthetic and respiratory complexes are found on thylakoid membranes in cyanobacteria. Therefore, in the absence of photosynthesis, diazoplast thylakoids likely were retained for respiratory activity. We propose that, in the diazoplast, metabolism of sugars via the OPP pathway and the latter "pay-off" phase of glycolysis generates reducing equivalents needed for high respiratory activity. Respiration not only provides the relatively large equivalents of ATP need for nitrogenase but also consumes oxygen as the terminal electron acceptor maintaining an anoxic microenvironment for nitrogenase activity. Consistent with this model, heterocysts are dependent on the OPP pathway for their nitrogenase activity, and genes related to respiration are upregulated at the transition from day to night in *C. subtropica* (32, 35, 36). There may be other mechanisms for scavenging oxygen in the diazoplast that we are continuing to explore.

A model for the nitrogen transfer to the host. Finally, we demonstrated that while negative feedback regulation of nitrogen fixation to ammonium was intact, and even sensitized, the same genetic regulation and environmental stimulus (i.e., ammonium repletion) resulted in paradoxical nitrogen assimilation responses and general suppression of the ammonia transporter, Amt. These results were puzzling at first but appear consistent with changes observed in diazotroph *Rhizobium* bacteria that colonize the root nodule of legumes. The *Rhizobium amt* gene is downregulated during its differentiation into nitrogen-fixing bacteroids in root nodules(58). Moreover, ectopic expression of *amt* disrupts bacteroid differentiation and the development of symbiotic nodules (59). Based on these findings, a model of nitrogen transfer in the legume nodule was proposed: ammonia produced by nitrogen-fixing bacteroids diffuses into the symbiosome space, an endosymbiotic compartment containing the bacteroids. Acidification of this space mediated by V-type H^+-ATPases traps ammonia in its protonated ammonium form at high concentrations (60). The downregulation of the bacteroid *amt* has been proposed to prevent ammonium transport back into the endosymbiotic bacteria; instead, ammonium is transported across the symbiosome membrane into the plant cell. In analogy to the root cell symbiosome, ammonium may also accumulate in the endosymbiotic compartment of the diazoplast. This ammonium accumulation around the diazoplast, is consistent with the increased sensitivity of the diazoplast to exogenous ammonium and the general suppression of *amt* expression compared to *C. subtropica*. As a result, the diazoplast is maintained in a state of nitrogen deficiency. Combined with the increased accumulation of 2-oxoglutarate (discussed above), the high C:N balance may positively regulate the expression of *nif* genes. We hypothesize that this mechanism has evolved to ensure nitrogen fixation activity to supply the host diatom's needs.

Implications for engineering nitrogen-fixing crop plants. Crop plants, particularly cereals, with the ability to fix nitrogen are the holy grail for sustainable agriculture. However, so far, attempts to transfer the genes for biological nitrogen fixation from bacteria into plants have met many challenges, including the large number of genes required, the complexity of nitrogenase assembly, the high cellular energy requirements, and exquisite sensitivity to oxygen (5, 6, 61). Ironically, the current strategy relies on targeting nitrogenase to the mitochondria where high respiratory rates in the compartment provides energy and protects it from inactivation (62–65). The diazotroph serves as a ready-made nitrogen-fixing factory, expressing all the genes required, properly regulated, and housed inside a protected membrane compartment. Given the advantages of nitrogen fixation capability, the absence of such an organelle is a mystery but may

be an accident of evolutionary timing or a human oversight (66, 67). No matter the state of diazotroph endosymbiosis in nature, new genetic tools open the opportunity to revisit this strategy in the laboratory. We propose that *Epithemia* diazoplasts could serve as a blueprint for engineering nitrogen-fixing organelles in plants. The metabolic adaptations we have observed in the diazoplast, for example, offers several options for genetic engineering of a suitable bacterial partner. We are as yet unaware of the extent of the host contribution to the endosymbiotic interaction, whether solely in the form of metabolites or also via imported gene products. However, analysis of *Epithemia* genomes is underway in our laboratory and will provide insight into gene transfers and imported gene products, if any. All together, our investigations of this native endosymbiotic interaction will be a powerful tool for bioengineering.

Materials and Methods

Strain isolation and cultivation
Environmental samples originate from a stream near Butano State Park, CA, USA and were taken during fall 2020. *Epithemia clementina* was isolated by successive dilution in a nitrogen-depleted media, under 10 µmole photon.$m^{-2}.s^{-1}$ of white light at 20°C as previously described (22) and have been since then been maintained in the same condition. The Csi-N media is detailed in Table S3. After subculturing for 8 weeks, we obtained many monoalgal isolates, some containing filamentous cyanobacteria and some containing *Epithemia* diatoms. From that point in time, cultures were started from single cell isolate of *E. clementina*. Cultures were monoalgal but not axenic, and cells formed a biofilm on the surface of the flasks. The Csi-N media was complemented with reduced nitrogen by addition of nitrate $Ca(NO_3)_2$, 4 H_2O at 60 µM final or ammonium $(NH_4)_2HPO_4$ at 100 µM final as described in (14). Conditions were changed 3 days prior to the measurements.

Crocosphaera suptropica (*Cyanothece* ATCC51142) was maintained at 30°C under day-night cycle (12 h / 12 h – 30 µmole photon.$m^{-2}.s^{-1}$ of / dark) in artificial sea water ASP2 (27) with constant bubbling of humidified ambient air. ASP2 media that contains no source of combined nitrogen, was complemented either as previously described for Csi media or with $NaNO_3$ at 17 mM final or ammonium $(NH_4)_2HPO_4$ at 2 mM final (12). Conditions were changed 3 days prior to the measurements.

For acetylene reduction assays (ARA), *E. clementina* cultures were scraped off the flasks and centrifuged (1,500xg, 4 min). They were washed once in fresh Csi-N media and resuspended in the appropriate media. From that culture, 2.3 mL was placed in an autoclaved 10 mL glass vial and sealed with a breathable sealing film. Note that the volume of culture was set to 2.3 mL to leave a 10 mL headspace in the vial. Vials were placed in their respective growth conditions for 3 days before the assay. For *C. subtropica*, fresh cultures were grown for 3 days in the desired conditions as described previously and sampled for ARA at the time of the experiment.

Observation of diazoplasts by DNA stain was performed after fixation for 10 min with 4% paraformaldehyde followed by incubation for 30 min with 1µg/mL DAPI. Fluorescence microscopy was performed with CY5 and TRITC filters for chlorophyll and phycoerythrin respectively.

Sequencing of the diazopalst
Epithemia clementina gDNA was extracted using QUIAGEN KIT TYPE. DNA fragments were prepped for Illumina sequencing with NEB ILLUMINA SEQUENCING KIT. Sequencing was performed on an Illumina NextSeq 2000 P3 paired end 150bp. Sequencing reads were trimmed, quality filtered with fastp (68) (--qualified_quality_phred 20 --unqualified_percent_limit 20), assembled with SPAades on default settings (69) and quality was assessed with FastQC (70).

For Nanopore sequencing, gDNA was isolated with circulomics nanobind plant nuclei DNA kit. Short DNA was removed with circulomics short read eliminator kit. Remaining DNA was then library prepped for nanopore sequencing using the Genomic DNA by Ligation (SQK-LSK112 kit) and loaded onto an R 9.4 flow cell. Nanopore reads were basecalled with guppy and short (< 3000 bp) and low quality (Phred quality score < 10) reads were removed with nanofilt (71).

Nanopore reads were assembled with Shasta (72), and contigs with one or more high quality (Expect value <= 10^{-10}) BLAST hits to previously published diazoplast genomes (NCBI accessions NZ_AP012549.1, NZ_AP018341.1) (17, 18) were isolated to create a fragmented and incomplete draft assembly. Nanopore and illumina reads were mapped to the draft assembly with minimap2 (73) and BWA-MEM (74) and mapped reads were re-assembled with unicycler (75) to produce a high quality circular diazoplast genome. Genome quality was assessed with Merqury and quast (76, 77). Gene annotation of the diazoplast was performed with NCBI Prokaryotic Genome Annotation Pipeline.

Phylogeny

Using the corresponding *Phaeodactylum tricornutum* or *Cyanothece* sp. ATCC 51142 gene sequences as queries, sequences of *Epithemia clementina* host genes for *psbC, rbcL* and 18S-rRNA and endosymbiont genes for *nifH* and 16S-rRNA were extracted from this assembly by command-line BLAST (78, 79). Sequence lengths extracted are as follows: 1391bp of *rbcL*, 1076bp of *psbC*, 1622bp of 18S-rRNA, 761bp of *nifH*, and 1410bp of 16S-rRNA. A nucleotide BLAST against the NCBI nr/nt database was performed to ensure correct identity of these sequences. For each gene, sequences were aligned using MAFFT v7.490 (mafft-linsi --adjustdirectionaccurately --maxiterate 1000) (80). Gaps and highly variable regions in the alignment were removed with trimAl v1.2rev59 (81) using the gappyout flag and inspected by eye to ensure proper alignment and trimming. Concatenated alignments and data partitions were generated using SequenceMatrix v1.9 (82). Phylogenetic trees were inferred using IQ-TREE 2 (83) with ModelFinder (84) automatic model selection and node support tested with 2000 iterations of rapid phylogenetic bootstraps.

Metagenomic analysis and assembly:

To assess metagenomic sequence diversity, MAGs were classified using kraken2 (85) and metagenomic diversity was visualized with Krona (86).

Metagenomic assembly was performed with flye (87) and contigs were polished 3x with racon (88). Eukaryotic contigs were removed with EukRep (89), and metagenomic binning was done with MetaBat2 (90). To investigate the possibility of nitrogen fixation by a free-living microbe, assembled contigs were searched against the UniProt reference proteomes database using Diamond (91) and results annotated as NifK (interpro IPR005976), NifD (interpro IPR005972), or NifH (interpro IPR005977) were isolated to identify putative nitrogen fixing microbes.

Nitrogenase activity assay

Vials with 2.3mL of culture were sealed and injected with 1 mL of acetylene generated using calcium carbide and water. After one hour of incubation under culture conditions or desired conditions, the reaction was blocked by injection of 150 μL of 16% paraformaldehyde and kept at 4°C until further analysis. Ethylene present in the 10 mL-head space was quantified by injection of 1mL of the headspace on a gas chromatography coupled with a flame ionization detector, Shimadzu GC-8A1F. A Porapak N 80/100 mesh 6' x 1/8" x 0.085" SSP/W column was used. The injector was set at 120°C and column at 80°C. Nitrogen was used as a carrier gas at 225kPa. Results were collected using a Shimadzu Chromatopac which performed automatic pic detection and quantification. A 1% ethylene standard gas was used to quantify the ethylene per signal detected. After gas analysis, cell were collected by centrifugation and chlorophyll was extracted with 100% ethanol for quantification (92).

Protein extraction and immunodetection

Cells were scraped and harvested at different time points of the day-night cycle by centrifugation at 3,000xg for 2 min. Pellets were deep-frozen in liquid nitrogen and stored at -80°C until protein extraction. Pellets were resuspended in 200μL of lysis buffer (10mM HEPES, 10mM EDTA, 0.5% tritonX-100, 2mM DTT). Cells were homogenized with a mix of 1mm and 0.5mM glass beads at 3000 strokes per minute for 2 minutes in a bead beater. Lysate was centrifuged at 15,000g for 5

min, 4°C and only supernatant was kept. Six volumes of cold acetone (-20°C) was added. After 1h at -20°C, proteins were pelleted at 15,000 g for 10 minutes, 4°C. Supernatant was kept for chlorophyll concentration measurement (92). Pellet was washed with 80% cold acetone, gently dried and resuspended at final concentration of 0.1 µg/mL of chlorophyll which correspond to 1 µg/mL of total protein content in 1X LDS loading buffer with 100 mM DTT. Sample were denaturated at 70°C for 30 min, and vortexed periodically. Before loading, samples were pelleted at 15,000xg for 5 min. Precision Plus Protein™ All Blue Standards was used as a molecular ladder to assess protein molecular weight.

Proteins were separated by electrophoresis on NuPage Gels, 4-12% Acrylamide using MES buffer. They were then transferred using Bio-rad Trans-blot Turbo transfer onto a nitrocellulose membrane. Membrane was blocked in LiCOR blocking buffer (0.1% Casein, 0.2x PBS, 0.01% sodium azide) for 1 hour at room temperature. The FeMo nitrogenase was revealed using polyclonal goat-raised antibody (1:500 dilution) kindly provided by Denis Dean from Virgnia Tech, US. The large subunit of the hydrogenase (HupL) was revealed with antibody (1:2500 dilution) raised in rabbit against amino acids 260 to 270 of HupL from *Anabaena* sp. PCC 7120 kindly provided by Paula Tamagnini from the Faculty of Sciences of University of Porto, Portugal. PsbA was used as an internal loading control, a 1:10,000 dilution was use of the rabbit antibody against the C-terminal of PsbA (AgriSera AB, Vanas, Sweden). Antibodies were diluted in 50% TBS-T / 50% LiCOR blocking buffer. After 2 hours with primary antibodies at room temperature, the membrane was washed 3 times with TBS-T and then incubated with LiCOR secondary antibodies (IRDye 800CW) for 1 hour (α-rabbit for PsbA and HupL and α-goat for NifDK). The membrane was rinsed twice with TBS-T and once with PBS and the signal was revealed using an infra-red LiCOR imager. Intensity of the signal was quantified using Image Studio Lite software v5.2. As FeMo nitrogenase and HupL have close molecular weight, the nitrogenase was first immunoblotted and the membrane was stripped with 0.2M NaOH for 5 minutes, washed with water twice and immunoblot was repeated for HupL.

Gene expression

After 3 days of subculturing in the desired condition, cells were harvested at 1,500x*g* for 2 minutes and resuspended in Trizol Reagent (Invitrogen™, Carlsbad CA, USA) and then lysed by a combination of flash-freezing and sonication. mRNAs were extracted using a Quiagen RNeasy® Plus Universal kit. mRNA concentration was assessed by nanodrop. Following extraction, RT-qPCR was achieved with NEB Luna® Universal One-Step RT-qPCR using 125ng of mRNA in a 10µL reaction as described in instructions. Amplification was performed and monitored in a 96-well plate using a StepOne™ Real-Time PCR system according to the instruction of the Luna kit and was followed by a melting curve. Primers used are detailed in Table S5. Specification of the amplification was validated first by PCR on gDNA and confirmed after the RT-qPCR of mRNA from analysis of the melting curves. DNA gyrase subunit B and 30S SSU ribosomal protein S1p were used as housekeeping genes. Expression was normalized to weighted average of GyrB and 30S SSU expression.

Acknowledgments

We thank Dr. Jonathan Zehr, Dr. Devaki Bhaya, and Dr. Arthur Grossman for their feedback on the project and reviews on the manuscript. We are grateful for Dr. Sharon Long, Dr. Denis Dean, and Dr. Paula Tamagnini to have shared material and equipment to make this work possible. This work was supported by the following funding sources: Burroughs Wellcome Fund (E.Y.), Chan Zuckerberg Biohub (E.Y.).

References

1. V. Smil, Enriching the Earth. MIT Press, Cambridge, MA. *Enriching the Earth. MIT Press, Cambridge, MA.* (2001).

2. A. G. Good, P. H. Beatty, Fertilizing Nature: A Tragedy of Excess in the Commons. *PLOS Biology* **9**, e1001124 (2011).

3. J. N. Galloway, *et al.*, Transformation of the Nitrogen Cycle: Recent Trends, Questions, and Potential Solutions. *Science* **320**, 889–892 (2008).

4. R. J. Diaz, R. Rosenberg, Spreading Dead Zones and Consequences for Marine Ecosystems. *Science* **321**, 926–929 (2008).

5. L. Curatti, L. M. Rubio, Challenges to develop nitrogen-fixing cereals by direct nif-gene transfer. *Plant Science* **225**, 130–137 (2014).

6. G. Soto, A. R. Fox, N. D. Ayub, Exploring the Intrinsic Limits of Nitrogenase Transfer from Bacteria to Eukaryotes. *J Mol Evol* **77**, 3–7 (2013).

7. C. Kneip, P. Lockhart, C. Voß, U.-G. Maier, Nitrogen fixation in eukaryotes – New models for symbiosis. *BMC Evolutionary Biology* **7**, 55 (2007).

8. J. Mareš, *et al.*, Taxonomic resolution of the genus Cyanothece (Chroococcales, Cyanobacteria), with a treatment on Gloeothece and three new genera, Crocosphaera, Rippkaea, and Zehria. *Journal of Phycology* **55**, 578–610 (2019).

9. E. J. Carpenter, S. Janson, Intracellular Cyanobacterial Symbionts in the Marine Diatom Climacodium Frauenfeldianum (bacillariophyceae). *Journal of Phycology* **36**, 540–544 (2000).

10. K. Hagino, R. Onuma, M. Kawachi, T. Horiguchi, Discovery of an Endosymbiotic Nitrogen-Fixing Cyanobacterium UCYN-A in Braarudosphaera bigelowii (Prymnesiophyceae). *PLOS ONE* **8**, e81749 (2013).

11. M. A. Schneegurt, D. M. Sherman, L. A. Sherman, Growth, Physiology, and Ultrastructure of a Diazotrophic Cyanobacterium, Cyanothece Sp. Strain Atcc 51142, in Mixotrophic and Chemoheterotrophic Cultures1. *Journal of Phycology* **33**, 632–642 (1997).

12. X. Feng, *et al.*, Mixotrophic and photoheterotrophic metabolism in Cyanothece sp. ATCC 51142 under continuous light. *Microbiology* **156**, 2566–2574 (2010).

13. C. R. Schvarcz, *et al.*, Overlooked and widespread pennate diatom-diazotroph symbioses in the sea. *Nat Commun* **13**, 799 (2022).

14. L. Floener, H. Bothe, Nitrogen fixation in Rhopalodia gibba, a diatom containing blue-greenish inclusions symbiotically. *Endocytobiosis & Cell Biology* **1**, 541–552 (1980).

15. J. Prechtl, C. Kneip, P. Lockhart, K. Wenderoth, U.-G. Maier, Intracellular Spheroid Bodies of Rhopalodia gibba Have Nitrogen-Fixing Apparatus of Cyanobacterial Origin. *Molecular Biology and Evolution* **21**, 1477–1481 (2004).

16. L. A. Sherman, H. Min, J. Toepel, H. B. Pakrasi, Better Living Through Cyanothece – Unicellular Diazotrophic Cyanobacteria with Highly Versatile Metabolic Systems in *Recent Advances in Phototrophic Prokaryotes*, P. C. Hallenbeck, Ed. (Springer, 2010), pp. 275–290.

17. T. Nakayama, *et al.*, Complete genome of a nonphotosynthetic cyanobacterium in a diatom reveals recent adaptations to an intracellular lifestyle. *PNAS* **111**, 11407–11412 (2014).

18. C. Kneip, C. Voβ, P. J. Lockhart, U. G. Maier, The cyanobacterial endosymbiont of the unicellular algae Rhopalodia gibba shows reductive genome evolution. *BMC Evolutionary Biology* **8**, 30 (2008).

19. T. Nakayama, Y. Inagaki, Genomic divergence within non-photosynthetic cyanobacterial endosymbionts in rhopalodiacean diatoms. *Sci Rep* **7**, 13075 (2017).

20. S. Kamakura, D. G. Mann, N. Nakamura, S. Sato, Inheritance of spheroid body and plastid in the raphid diatom Epithemia (Bacillariophyta) during sexual reproduction. *Phycologia* **0**, 1–9 (2021).

21. M. Hajós, Stratigraphy of Hungary's Miocene diatomaceous earth deposits. *Geologica Hungarica, Series Palaeontologica* **49**, 1–339 (1986).

22. S. Adler, L. M. Derch, U. G. Maier, Cultivation of the diatom Rhopalodia gibba. *Endocytobiosis & Cell Research* (2010).

23. E. C. Ruck, T. Nakov, A. J. Alverson, E. C. Theriot, Phylogeny, ecology, morphological evolution, and reclassification of the diatom orders Surirellales and Rhopalodiales. *Molecular Phylogenetics and Evolution* **103**, 155–171 (2016).

24. T. Nakayama, *et al.*, Spheroid bodies in rhopalodiacean diatoms were derived from a single endosymbiotic cyanobacterium. *J Plant Res* **124**, 93–97 (2011).

25. A. Bandyopadhyay, J. Stöckel, H. Min, L. A. Sherman, H. B. Pakrasi, High rates of photobiological H 2 production by a cyanobacterium under aerobic conditions. *Nature Communications* **1**, 139 (2010).

26. A. Bandyopadhyay, T. Elvitigala, M. Liberton, H. B. Pakrasi, Variations in the Rhythms of Respiration and Nitrogen Fixation in Members of the Unicellular Diazotrophic Cyanobacterial Genus Cyanothece. *Plant Physiology* **161**, 1334–1346 (2013).

27. K. J. Reddy, J. B. Haskell, D. M. Sherman, L. A. Sherman, Unicellular, aerobic nitrogen-fixing cyanobacteria of the genus Cyanothece. *J Bacteriol* **175**, 1284–1292 (1993).

28. K. M. Wegener, A. Nagarajan, H. B. Pakrasi, An Atypical psbA Gene Encodes a Sentinel D1 Protein to Form a Physiologically Relevant Inactive Photosystem II Complex in Cyanobacteria. *J Biol Chem* **290**, 3764–3774 (2015).

29. M. S. Colón-López, D. M. Sherman, L. A. Sherman, Transcriptional and translational regulation of nitrogenase in light-dark- and continuous-light-grown cultures of the unicellular cyanobacterium Cyanothece sp. strain ATCC 51142. *Journal of Bacteriology* **179**, 4319–4327 (1997).

30. M. A. Schneegurt, D. M. Sherman, S. Nayar, L. A. Sherman, Oscillating behavior of carbohydrate granule formation and dinitrogen fixation in the cyanobacterium Cyanothece sp. strain ATCC 51142. *J Bacteriol* **176**, 1586–1597 (1994).

31. X. Zhang, D. M. Sherman, L. A. Sherman, The Uptake Hydrogenase in the Unicellular Diazotrophic Cyanobacterium Cyanothece sp. Strain PCC 7822 Protects Nitrogenase from Oxygen Toxicity. *J Bacteriol* **196**, 840–849 (2014).

32. J. Stöckel, *et al.*, Global transcriptomic analysis of Cyanothece 51142 reveals robust diurnal oscillation of central metabolic processes. *PNAS* **105**, 6156–6161 (2008).

33. M. Cano, *et al.*, Glycogen Synthesis and Metabolite Overflow Contribute to Energy Balancing in Cyanobacteria. *Cell Reports* **23**, 667–672 (2018).

34. R. W. Drum, S. Pankratz, Fine structure of an unusual cytoplasmic inclusion in the diatom genus, Rhopalodia. *Protoplasma* **60**, 141–149 (1965).

35. F. Winkenbach, C. P. Wolk, Activities of Enzymes of the Oxidative and the Reductive Pentose Phosphate Pathways in Heterocysts of a Blue-Green Alga 1. *Plant Physiology* **52**, 480–483 (1973).

36. M. L. Summers, J. G. Wallis, E. L. Campbell, J. C. Meeks, Genetic evidence of a major role for glucose-6-phosphate dehydrogenase in nitrogen fixation and dark growth of the cyanobacterium Nostoc sp. strain ATCC 29133. *J Bacteriol* **177**, 6184–6194 (1995).

37. J. E. Frías, A. Mérida, A. Herrero, J. Martín-Nieto, E. Flores, General distribution of the nitrogen control gene ntcA in cyanobacteria. *J Bacteriol* **175**, 5710–5713 (1993).

38. I. Luque, E. Flores, A. Herrero, Molecular mechanism for the operation of nitrogen control in cyanobacteria. *The EMBO Journal* **13**, 2862–2869 (1994).

39. J. E. Frías, E. Flores, A. Herrero, Requirement of the regulatory protein NtcA for the expression of nitrogen assimilation and heterocyst development genes in the cyanobacterium Anabaena sp. PCC7120. *Molecular Microbiology* **14**, 823–832 (1994).

40. M. A. Vega-Palas, E. Flores, A. Herrero, NtcA, a global nitrogen regulator from the cyanobacterium Synechococcus that belongs to the Crp family of bacterial regulators. *Molecular Microbiology* **6**, 1853–1859 (1992).

41. M. F. Vázquez-Bermúdez, A. Herrero, E. Flores, 2-Oxoglutarate increases the binding affinity of the NtcA (nitrogen control) transcription factor for the Synechococcus glnA promoter. *FEBS Letters* **512**, 71–74 (2002).

42. I. Luque, M. F. Vázquez-Bermúdez, J. Paz-Yepes, E. Flores, A. Herrero, In vivo activity of the nitrogen control transcription factor NtcA is subjected to metabolic regulation in Synechococcus sp. strain PCC 7942. *FEMS Microbiology Letters* **236**, 47–52 (2004).

43. M. I. Muro-Pastor, J. C. Reyes, F. J. Florencio, Cyanobacteria Perceive Nitrogen Status by Sensing Intracellular 2-Oxoglutarate Levels *. *Journal of Biological Chemistry* **276**, 38320–38328 (2001).

44. H.-M. Lee, M. F. Vázquez-Bermúdez, N. T. de Marsac, The Global Nitrogen Regulator NtcA Regulates Transcription of the Signal Transducer PII (GlnB) and Influences Its Phosphorylation Level in Response to Nitrogen and Carbon Supplies in the Cyanobacterium Synechococcus sp. Strain PCC 7942. *Journal of Bacteriology* **181**, 2697–2702 (1999).

45. M. Fadi Aldehni, J. Sauer, C. Spielhaupter, R. Schmid, K. Forchhammer, Signal Transduction Protein PII Is Required for NtcA-Regulated Gene Expression during Nitrogen Deprivation in the Cyanobacterium Synechococcus elongatus Strain PCC 7942. *Journal of Bacteriology* **185**, 2582–2591 (2003).

46. J. Paz-Yepes, E. Flores, A. Herrero, Transcriptional effects of the signal transduction protein PII (glnB gene product) on NtcA-dependent genes in Synechococcus sp. PCC 7942. *FEBS Letters* **543**, 42–46 (2003).

47. J. Liang, L. Scappino, R. Haselkorn, The patB gene product, required for growth of the cyanobacterium Anabaena sp. strain PCC 7120 under nitrogen-limiting conditions, contains ferredoxin and helix-turn-helix domains. *J Bacteriol* **175**, 1697–1704 (1993).

48. K. M. Jones, W. J. Buikema, R. Haselkorn, Heterocyst-Specific Expression of patB, a Gene Required for Nitrogen Fixation in Anabaena sp. Strain PCC 7120. *Journal of Bacteriology* **185**, 2306–2314 (2003).

49. R. Tsujimoto, N. Kamiya, Y. Fujita, Transcriptional regulators ChlR and CnfR are essential for diazotrophic growth in nonheterocystous cyanobacteria. *PNAS* **111**, 6762–6767 (2014).

50. S. Ehira, M. Ohmori, NrrA, a Nitrogen-regulated Response Regulator Protein, Controls Glycogen Catabolism in the Nitrogen-fixing Cyanobacterium Anabaena sp. Strain PCC 7120. *J Biol Chem* **286**, 38109–38114 (2011).

51. D. Liu, C. Yang, The Nitrogen-regulated Response Regulator NrrA Controls Cyanophycin Synthesis and Glycogen Catabolism in the Cyanobacterium Synechocystis sp. PCC 6803. *Journal of Biological Chemistry* **289**, 2055–2071 (2014).

52. K. Kucho, *et al.*, Global Analysis of Circadian Expression in the Cyanobacterium Synechocystis sp. Strain PCC 6803. *Journal of Bacteriology* **187**, 2190–2199 (2005).

53. W. Mohr, M. P. Intermaggio, J. LaRoche, Diel rhythm of nitrogen and carbon metabolism in the unicellular, diazotrophic cyanobacterium Crocosphaera watsonii WH8501. *Environmental Microbiology* **12**, 412–421 (2010).

54. S. T. Wilson, *et al.*, Coordinated regulation of growth, activity and transcription in natural populations of the unicellular nitrogen-fixing cyanobacterium Crocosphaera. *Nat Microbiol* **2**, 1–9 (2017).

55. S. E. Cohen, S. S. Golden, Circadian Rhythms in Cyanobacteria. *Microbiology and Molecular Biology Reviews* **79**, 373–385 (2015).

56. N. A. Held, *et al.*, Dynamic diel proteome and daytime nitrogenase activity supports buoyancy in the cyanobacterium Trichodesmium. *Nat Microbiol*, 1–12 (2022).

57. W. Luo, K. Inomura, H. Zhang, Y.-W. Luo, N2 Fixation in Trichodesmium Does Not Require Spatial Segregation from Photosynthesis. *mSystems* **0**, e00538-22 (2022).

58. R. Taté, A. Riccio, M. Merrick, E. J. Patriarca, The Rhizobium etli amtB Gene Coding for an NH4+ Transporter Is Down-Regulated Early During Bacteroid Differentiation. *MPMI* **11**, 188–198 (1998).

59. R. Taté, *et al.*, Ectopic Expression of the Rhizobium etli amtB Gene Affects the Symbiosome Differentiation Process and Nodule Development. *MPMI* **12**, 515–525 (1999).

60. M. Udvardi, P. S. Poole, Transport and metabolism in legume-rhizobia symbioses. *Annual review of plant biology* **64**, 781–805 (2013).

61. C. Rogers, G. E. D. Oldroyd, Synthetic biology approaches to engineering the nitrogen symbiosis in cereals. *Journal of Experimental Botany* **65**, 1939–1946 (2014).

62. J. R. Postgate, G. W. Cooke, N. W. Pirie, G. D. H. Bell, Possibilities for the enhancement of biological nitrogen fixation. *Philosophical Transactions of the Royal Society of London. B, Biological Sciences* **281**, 249–260 (1997).

63. G. López-Torrejón, *et al.*, Expression of a functional oxygen-labile nitrogenase component in the mitochondrial matrix of aerobically grown yeast. *Nat Commun* **7**, 11426 (2016).

64. N. B. Ivleva, J. Groat, J. M. Staub, M. Stephens, Expression of Active Subunit of Nitrogenase via Integration into Plant Organelle Genome. *PLOS ONE* **11**, e0160951 (2016).

65. A. Shamseldin, Future Outlook of Transferring Biological Nitrogen Fixation (BNF) to Cereals and Challenges to Retard Achieving this Dream. *Curr Microbiol* **79**, 171 (2022).

66. J. F. Allen, J. A. Raven, Free-radical-induced mutation vs redox regulation: Costs and benefits of genes in organelles. *J Mol Evol* **42**, 482–492 (1996).

67. C. P. McKay, R. Navarro-González, "The Absence of Nitrogen-Fixing Organelles Due to Timing of the Nitrogen Crisis" in *Symbiosis: Mechanisms and Model Systems*, Cellular Origin, Life in Extreme Habitats and Astrobiology., J. Seckbach, Ed. (Springer Netherlands, 2002), pp. 221–228.

68. S. Chen, Y. Zhou, Y. Chen, J. Gu, fastp: an ultra-fast all-in-one FASTQ preprocessor. *Bioinformatics* **34**, i884–i890 (2018).

69. A. Prjibelski, D. Antipov, D. Meleshko, A. Lapidus, A. Korobeynikov, Using SPAdes De Novo Assembler. *Current Protocols in Bioinformatics* **70**, e102 (2020).

70. S. Andrews, FastQC: a quality control tool for high throughput sequence data (2010).

71. W. De Coster, S. D'Hert, D. T. Schultz, M. Cruts, C. Van Broeckhoven, NanoPack: visualizing and processing long-read sequencing data. *Bioinformatics* **34**, 2666–2669 (2018).

72. K. Shafin, *et al.*, Nanopore sequencing and the Shasta toolkit enable efficient de novo assembly of eleven human genomes. *Nat Biotechnol* **38**, 1044–1053 (2020).

73. H. Li, Minimap2: pairwise alignment for nucleotide sequences. *Bioinformatics* **34**, 3094–3100 (2018).

74. H. Li, Aligning sequence reads, clone sequences and assembly contigs with BWA-MEM (2013) https:/doi.org/10.48550/arXiv.1303.3997 (March 3, 2023).

75. R. R. Wick, L. M. Judd, C. L. Gorrie, K. E. Holt, Unicycler: Resolving bacterial genome assemblies from short and long sequencing reads. *PLOS Computational Biology* **13**, e1005595 (2017).

76. A. Rhie, B. P. Walenz, S. Koren, A. M. Phillippy, Merqury: reference-free quality, completeness, and phasing assessment for genome assemblies. *Genome Biology* **21**, 245 (2020).

77. A. Gurevich, V. Saveliev, N. Vyahhi, G. Tesler, QUAST: quality assessment tool for genome assemblies. *Bioinformatics* **29**, 1072–1075 (2013).

78. S. F. Altschul, W. Gish, W. Miller, E. W. Myers, D. J. Lipman, Basic local alignment search tool. *Journal of Molecular Biology* **215**, 403–410 (1990).

79. C. Camacho, *et al.*, BLAST+: architecture and applications. *BMC Bioinformatics* **10**, 421 (2009).

80. K. Katoh, K. Kuma, H. Toh, T. Miyata, MAFFT version 5: improvement in accuracy of multiple sequence alignment. *Nucleic Acids Research* **33**, 511–518 (2005).

81. S. Capella-Gutiérrez, J. M. Silla-Martínez, T. Gabaldón, trimAl: a tool for automated alignment trimming in large-scale phylogenetic analyses. *Bioinformatics* **25**, 1972–1973 (2009).

82. G. Vaidya, D. J. Lohman, R. Meier, SequenceMatrix: concatenation software for the fast assembly of multi-gene datasets with character set and codon information. *Cladistics* **27**, 171–180 (2011).

83. B. Q. Minh, *et al.*, IQ-TREE 2: New Models and Efficient Methods for Phylogenetic Inference in the Genomic Era. *Molecular Biology and Evolution* **37**, 1530–1534 (2020).

84. S. Kalyaanamoorthy, B. Q. Minh, T. K. F. Wong, A. von Haeseler, L. S. Jermiin, ModelFinder: fast model selection for accurate phylogenetic estimates. *Nature Methods* **14**, 587–589 (2017).

85. D. E. Wood, J. Lu, B. Langmead, Improved metagenomic analysis with Kraken 2. *Genome Biol* **20**, 257 (2019).

86. B. D. Ondov, N. H. Bergman, A. M. Phillippy, Interactive metagenomic visualization in a Web browser. *BMC Bioinformatics* **12**, 385 (2011).

87. M. Kolmogorov, *et al.*, metaFlye: scalable long-read metagenome assembly using repeat graphs. *Nat Methods* **17**, 1103–1110 (2020).

88. R. Vaser, I. Sović, N. Nagarajan, M. Šikić, Fast and accurate de novo genome assembly from long uncorrected reads. *Genome Res* **27**, 737–746 (2017).

89. P. T. West, A. J. Probst, I. V. Grigoriev, B. C. Thomas, J. F. Banfield, Genome-reconstruction for eukaryotes from complex natural microbial communities. *Genome Res* **28**, 569–580 (2018).

90. D. D. Kang, *et al.*, MetaBAT 2: an adaptive binning algorithm for robust and efficient genome reconstruction from metagenome assemblies. *PeerJ* **7**, e7359 (2019).

91. B. Buchfink, K. Reuter, H.-G. Drost, Sensitive protein alignments at tree-of-life scale using DIAMOND. *Nat Methods* **18**, 366–368 (2021).

92. R. J. Ritchie, Consistent Sets of Spectrophotometric Chlorophyll Equations for Acetone, Methanol and Ethanol Solvents. *Photosynth Res* **89**, 27–41 (2006).

Figure 1. Identification of *Epithemia clementina* (A) Light microscopy of *Epithemia clementina* apex (top) and valve/ventral region (bottom) with "spheroid bodies" (black arrow). (B) Epifluorescence microscopy of *E. clementina* comparing brightfield (left), DAPI stain (middle), and chlorophyll florescence (right). (C) Light microscopy of isolated endosymbionts released from crushed *E. clementina* (left), compared with free-living *C. subtropica* (right). (D) Epifluorescence microscopy of isolated endosymbionts (top) and *C. subptropica* (bottom) comparing brightfield (left), chlorophyll (Chl) fluorescence (middle), and phycoerythrin (PE) fluorescence (right). (E) Multigene phylogeny of *E. clementina* and related diatoms based on the ribosomal small subunit RNA (18S rRNA), psbC, and rbcL genes. (F) Multigene phylogeny of the *E. clementina* endosymbionts and related cyanobacteria, based on the ribosomal small subunit RNA (16S rRNA) and *nifH* genes. Accession numbers for all sequences are provided in Source Data file (Table S3)*.

Figure 2. Diel pattern of nitrogen fixation in *E. clementina* and *C. subtropica* ATCC 51142
(A) Nitrogenase activity assessed by acetylene reduction assay over a diel cycle in *E. clementina* (brown) and *C. subtropica* ATCC 51142 (green) normalized by chlorophyll content. (B) Corresponding immunoblot for nitrogenase (NifDK) and hydrogenase uptake (HupL). (C) Quantification of NifDK to PsbA ratio from B. (D) Relative gene expression of *nifK* and *hupS* in *E. clementina* (brown) and *C. subtropica* (green) during early day (open bars) and early night (solid bars). Values are mean of 3 biological replicates ± SD.

Figure 3. Carbohydrate metabolism associated with nitrogen fixation activity in the diazoplast (A) Nitrogenase activity assessed by acetylene reduction assay at day and nighttime points in *E. clementina,* comparing photosynthetic conditions (light) and non-photosynthetic conditions (dark or DCMU treatment). (B) Annotated carbon metabolism pathways in diazoplast, showing proteins only present in *C. subtropica* and absent in diazoplast's genome (red). See Table S1 for corresponding gene ID numbers. (C-D) Relative gene expression of representative genes in glycogen biosynthesis (*glgA*), glycogen degradation (*glgP*), and OPP (*zwf*) pathways in *E. clementina* (brown) and *C. subtropica* (green) during early day (open bars) and early night (solid bars). Values are mean of 3 biological replicates ± SD.

Figure 4. Effects of nitrogen repletion on nitrogen fixation (A) Annotated nitrogen utilization pathways showing genes absent in the diazoplast but present in *C. subtropica* (red). See Table S1 for corresponding gene ID numbers. (B) Relative nitrogenase activity assessed by acetylene reduction assay upon repletion with nitrate or ammonium. Data were normalized by the mean of N-free samples. Measurements were collected at 2 hours into the day for *E. clementina* and 4 hours into the night for *C. subtropica*. (C) Relative expression of *nifK* in corresponding samples from B. For *C. Subtropica*, only samples in high concentrations of nitrate and ammonium were measured.   (D) Immunoblots for NifDK, HupL, and PsbA proteins in corresponding samples from B. (E) Same as C for relative expression of *amt*. Values are mean of 3 biological replicates ± SD.

Figure 5. Effect of nitrogen repletion on regulatory genes (A) Regulation network for nitrogen fixation, nitrogen assimilation, and associated carbon metabolism in *C. subtropica*. See Table S1 for corresponding gene ID. (B) Relative gene expression of nitrogen regulators in *E. Clementina*. Samples were taken at 2 hours into the day. (C) Same as B for *C. subtropica*. Samples were taken at 4 hours into the night using only high concentrations of nitrate and ammonium. Values are mean of 3 biological replicates ± SD.

Supplemental materials

Fig S1: Supplemental micrographs of *E. clementina*

(A) Micrography of *Epithemia clementina* shell with diazoplasts (indicated by the black triangles)
(B) Fluorescence micrography of *E. clementina* diazoplast (top) and *C. subtropica* (bottom)
showing fluorescence of chlorophyll and phycoerythrin

Fig. S2: Diversity of the bacterial community. Krona visualization of kraken assignment of metagenome-assembled genomes. See Table S1 for detailed abundance.

Fig. S3: Regulon of NrrA in Cyanothece and diazoplast. Alignment of identified NrrA binding sequences in promoter region of *glgP*, *NrrA*, *ArgG* and an *Acetyltransferase* in two free-living Cyanothece (*C. subtropica* and *Rippkaea orientalis*) and 3 diazoplasts (*E. clementina*, *E. turgida*, *R. gibberula*). Number indicates the position from the start codon.

Table S1: MAGs analysis and taxonomic assignment.

Table S2: NCBI gene IDs. Genes present ID corresponding to protein involved in carbon metabolism, nitrogen metabolism, nitrogen regulation and circadian clock.

Table S3: Media recipe for Csi-N

Table S4: Source data for phylogeny

Table S5: List of primer used for RT-qPCR

Primer name	Primer sequence - Diazoplast	Primer sequence - *C. subtropica*
gyrB-For	GGTGGTAAATTCGGGGGAGG	TCG CTA TCA ACG GGC ATT AG
gyrB-Rev	TTCCCCTTTCGTAGCGTTGG	GCG CCC TCT CTC TAC TTT ATA C
30S-For	CACTCACGAGGATTTTGCGGC	CCT CCA GTC CAA CGA AAC AA
30S-Rev	CGTCTTCAGGATCGTCAACTCGG	GTT CCC AAG CTC GCA TAT ACT
NifK-For	CGTTGATCTGTTCCACCAGCC	CGG TGA TCC TGA TCT CGT AAT G
NifK-Rev	GTTGGCAAGCTTTAGCGGGG	CTT CAA ATT CCT TGG TGC TGT T
HupS-For	GAACCGGTTTGCTGGCCG	GGC CTG TTC TGG GAA TAC AA
HupS-Rev	GGTATTTAGCCCCTAAAGCTCCCCC	TAG GGA GGG ATG CCA TAA GA
glgA-For	CTCGTTTCGAACCTTGTGGC	CTT GGG AAG GAT TCC GCT TTA
glgA-Rev	CTCGTTTCGAACCTTGTGGC	ATT CGG CTG CTG ACT GAT AC
glgP-For	CGACCTAAGCCTCCGTTACC	GAC TCG GAA ACG GAG GAT TAG
glgP-Rev	TCGGCTGAATTCCTCATGGG	AAC TCG TAG CGG ATA CCA TAA C
zwf-For	GTTAACCCCTGGCTCTTCCC	GCC CAT CAA ACC AGT CCT AAT
zwf-Rev	GAACCTCCTAACGCTCTGGG	GTA GTT CTG ACC CTG GCA TTT
Amt-For	GCAGCCGATGAAATGGTTCC	CGG GTT CGG GAT GTT AGA A
Amt-Rev	GGATACCCCATCACAACCGG	ATA AGC AAG GGT GGC TAA GG
cnfR-For	GCGGACTTACCTGATGATGC	GGG AAG CTT GTA CCG TGT TA
cnfR-Rev	AGTTGTTCTTCCCACTCCGG	AAT GGT TCC TTG TCC TTG TTT G
NrrA-For	AAGGACTTAACTCCGGAGCC	CTT CAC CAA CAA CGG CAA TC
NrrA-Rev	TCTCCTCGGTACTGGACTCG	TGC ATT CCG AAG GGT TTA GT
NtcA-For	CCTAGCAGCTGTTTTTCGCC	GTG TTT GGC GTA CTG TCT TTG
NtcA-Rev	CGGGAAAGTTTTACGGCTCC	AGG GCT TGT TCG ACT TGT T
PII-For	CGGCGACCAATTTGTTGACC	GAA GCG ATT ATC GAC CCT TTA
PII-Rev	TCAGAAGTGCGGGGATTTGG	GTC CAA ACC CTC GAA CTT CA