Regulation of B cell function and expression of CD11c, T-bet, and FcLR5 after infection and in response to different activation pathways

Linn Kleberg[1,2,3], Alan-Dine Corey-Ghaouzi[1,2,3], Maximilian Julius Lautenbach[1,2,3], Anna Färnert[1,2,3], Christopher Sundling[1,2,3,*]

[1] Division of Infectious Diseases, Department of Medicine, Solna, Karolinska Institutet, Sweden
[2] Center for Molecular Medicine, Karolinska Institutet, Sweden
[3] Department of Infectious Diseases, Karolinska University Hospital, Sweden

[*]Corresponding author: Christopher Sundling, Karolinska University Hospital, BioClinicum J7:20, Visionsgatan 4, 171 64 Solna, Sweden, christopher.sundling@ki.se

Keywords: Atypical B cells, double-negative B cells, Malaria, FcRL5, CD11c, T-bet, B cells

Abstract

CD11c, FcRL5, or T-bet are commonly expressed by B cells expanding during inflammation, where they can make up >30% of mature B cells. However, the association between the proteins and differentiation and function in the host response remain largely unclear. We have assessed the co-expression of CD11c, T-bet and FcRL5 over one year in patients treated for malaria. We observed dynamic co-expression profiles that changed over time, suggesting ongoing differentiation. We used an in vitro B cell culture system to determine how stimulation via the B cell receptor (BCR), toll-like receptor 9 (TLR9), and different cytokines could influence CD11c, T-bet, and FcRL5 expression. We observed different expression dynamics for all markers, but a largely overlapping regulation of CD11c and FcRL5 in response to BCR and TLR9 activation, while T-bet was strongly dependent on IFN-γ signalling. Investigating plasma cell differentiation and antigen-presenting cell functions, there was no association between marker expression and antibody secretion or T cell help, although the functions were associated with TLR9-signalling and B cell-derived IL-6 production, respectively. These results point to CD11c, FcRL5, and T-bet expression representing different activation stages and highlight the value of using multiple markers when assessing B cell differentiation.

Introduction

B cells are important contributors to long-lived immunity through their differentiation to antibody-secreting plasma cells and long-lived memory B cells. This process is initiated in secondary lymphoid organs where antigen-activated mature naïve B cells either directly differentiate to plasma cells (T-cell-independent), in response to high-valent antigen or strong and pattern-recognition-receptor (PRR)-signal, or migrate to T cell-B cell interfollicular to interact with activated cognate CD4[+] T cells (T-cell dependent response) [1]. In the T-cell-dependent response, B cells present peptide antigens to the cognate T cells via the major histocompatibility complex (MHC) class 2. Both the T- and B-cells then deliver additional activation and regulatory signals to each other via co-stimulatory molecules (such as CD40-CD40L, ICOS-ICOSL, CD80/86-CD28) and cytokines [2]. At this stage, B cells can undergo cytokine-dependent class-switch recombination [3], enter the B cell follicle and establish germinal centres to undergo isotype- and affinity-dependent selection [4–6], eventually producing high-affinity memory B cells and plasma cells [7], or they can directly differentiate to relatively low-affinity extrafollicular memory B cells and plasma cells [8].

For certain infectious diseases with a strong or prolonged inflammatory component, such as malaria, the process of generating long-lived B cell-mediated immunity has been suggested to be inefficient [9]. This has in part been associated with the expansion of a B cell subset lacking the expression of CD21 and CD27 while upregulating the integrin CD11c, the transcription factor T-bet and the IgG receptor FcRL5, in addition to numerous other receptors associated with cell regulation and migration [10–12]. These cells have been suggested to be exhausted or dysfunctional, in part due to reduced responsiveness to BCR-ligation and upregulation of several inhibitory receptors [12,13]. However, a recent study by Ambegaonkar et al. suggests it could have more to do with how the antigen is presented [14], and it remains unclear what function and purpose these cells play in the immune response [15]. Some studies have suggested that the cells could be primed to migrate to sites of infection or inflammation to provide localized differentiation to plasma cells or to provide antigen-presenting functions to local CD4[+] T cells [15–17]. In support of such functions, CD21[−]CD27[−] B cells in malaria display an upregulation of the plasma cell-associated genes *PRDM1* and *CD38* and differentiate into antibody-secreting cells following interaction with T-follicular helper (T_FH) cells [18]. Additionally, mouse and human studies have indicated more efficient antigen uptake and upregulation of MHC2 and co-stimulatory molecules, potentially translating to an enhanced capacity to drive CD4[+] T cell responses [14,18,19].

It has become clear that CD11c[+] or T-bet[+] B cells are not necessarily only associated with severe inflammatory disease, as they are found at low levels during steady-state that then increase with normal ageing of the host [20–22]. These levels are, however, greatly expanded in many different infections and following vaccination [23]. Additionally, cells with a similar phenotype are also expanded in several different autoimmune diseases that have an inflammatory component [24]. Despite sharing many phenotypic and transcriptomic characteristics between different types of diseases [25], there are also differences that appear disease-specific, such as the expression of FcRL4 in HIV infection and FcRL5 in malaria [12,26]. An additional source of variation could come from how the cells are defined in different studies, where varying combinations of IgD, CD21, CD27, CD11c, CD85j, T-bet, CXCR3, and FcRL5/FcRL4 have been used. This has also led to the cells having been given many different names, such as atypical B cells, age-associated B cells, exhausted B cells, tissue-like B cells, double negative (DN) and DN2 B cells [10,12,26–32]. We have previously shown that the

80   co-expression of several of these markers is changing over time after parasitic infection [27]. It is
81   therefore possible that they could represent different stages of differentiation or alternatively that
82   the cells have received different stimuli prior to their differentiation.
83
In this study, we have investigated the regulation of CD11c, T-bet, and FCRL5, three of the most used
markers of these alternative lineage B cells. We have measured their expression patterns in malaria
patients from diagnosis and up to 12 months after treatment. We have also used an in vitro culture
system to determine how stimulation via the BCR, TLR9 or the response to different cytokines
influences their expression. Additionally, we have assessed their functional properties, including
differentiation to antibody-secreting cells and B cell-dependent $CD4^+$ T cell help.
90

Results

Co-expression of CD11c, T-bet, and FcRL5 on B cells after diagnosis with Plasmodium parasite
infection

To evaluate the dynamics and co-expression of CD11c, T-bet, and FcRL5, we analysed peripheral
blood mononuclear cells (PBMCs) using flow cytometry in repeated samples up to 12 months after
diagnosis with *Plasmodium* parasite infections (**Figure 1A**). Samples from ten patients and six healthy
controls were included. Six patients were symptomatic with *P. falciparum* parasites (parasitaemia
ranging from 0.1-7%), two were symptomatic with *P. ovale* or *P. vivax*, and two were asymptomatic
(one *P. malariae* and one *P. ovale*). Six individuals originated from malaria-endemic areas. From six
of the patients, PBMC samples were available from all three time points tested. Cell marker
expression over time and patient characteristics are available in **supplementary figure 1**.

Switched atypical memory B cells can be classified by their lack of IgD and CD27 expression and then
also be referred to as double-negative (DN) B cells [31]. We observed that the percentage of DN cells
was higher (approximately 30-50%) among B cells expressing CD11c, T-bet, or FcRL5 (called DN2 cells
[31]) compared with those that did not express any of the three markers, where only 10-15% of the
cells were IgD and CD27 negative (**Figure 1B and supplementary figure 2**). The frequency of switch
memory (IgD⁻CD27⁺, approximately 20% of the cells) and unswitched memory (IgD⁺CD27⁺,
approximately 15% of the cells) remained similar between CD11c, T-bet, or FcRL5 positive and
negative cells, and it was mainly the naïve cells (IgD⁺CD27⁻) that had a reduced frequency compared
with the DN phenotype (**Supplementary figure 2**).

The frequency of CD11c, T-bet, and FcRL5 expressing B cells among $CD38^{lo/int}CD24^{lo/int}$ mature B cells
was significantly increased in individuals at diagnosis compared with healthy controls (**Figure 1C**).
For CD11c and FcRL5 the expression remained significantly higher also at 10 days after diagnosis and
treatment initiation but then declined over the following 12 months. At diagnosis, an average of
30.4% of the mature B cells expressed CD11c, T-bet, and/or FcRL5 (**Figure 1D**). The majority of these
cells (13.9%) expressed all three markers, followed by CD11c single-positive cells (6.3%) and FcRL5
single-positive cells (4.2%). CD11c and FcRL5 double-positive cells only made up 2.6% of the mature
B cells at diagnosis but then increased in frequency to 8.2% of cells at the 10-day time point.
Similarly, FcRL5 single-positive cells also increase to 9.1% of the mature B cells at the 10-day time-
point after which both FcRL5 single and FcRL5 and CD11c double-positive cells declined to baseline
levels at 12 months. The triple-positive cells were reduced to 5.1% at 10 days and then further to
4.5% at 12 months (**Figure 1D**). Together these data show that the commonly used atypical B cell
markers CD11c, T-bet, and FcRL5 are not mutually exclusive, but rather have a dynamic co-
expression pattern over time following acute infection.

128
Figure 1. Expression of CD11c, T-bet, and FcRL5 on B cells following diagnosis of *Plasmodium*
parasite infection. *(A) Schematic presentation of the donors and time points. PBMCs from healthy*
controls (HC, n=6), and Plasmodium parasite-infected longitudinal samples at diagnosis (n=7), 10
days after treatment initiation (n=10), and after 12 months (n=6) were assessed by flow cytometry.
(B) Mature B cells (Live, CD19+CD20+CD38int/lowCD24int/low) were analyzed for expression of CD11c, T-
bet, and FcRL5 and further gated for CD27 and IgD. (C) CD11c+, T-bet+, and FcRL5+ B cell frequencies
were determined at the diagnosis, 10-day, and 12-month time points (red) and in healthy controls
(blue). The mean and 95% confidence interval are indicated. (D) Average frequency of co-expression
of CD11c, T-bet, and FcRL5 among mature B cells at the different time points after infection. The
Kruskal-Wallis test followed by Dunn's posttest was used to determine p-values between healthy
*controls and parasite-infected donors at different time-points after infection. *p<0.05, **p<0.01,*
****p<0.001. ns indicates not significant, p>0.05.*

141
Differential kinetics of CD11c, T-bet, and FcRL5 after stimulation
To better understand the regulation of CD11c, T-bet, and FcRL5 in B cells, we adapted an in vitro
culture system described by Ambegaonkar et al. [33] to generate B cells with an atypical phenotype.
They had previously shown that both naïve and memory B cells could upregulate T-bet following in
vitro stimulation via the BCR, TLR9, and by adding IFN-γ [33]. We observed a similar strong
upregulation of T-bet in both naïve and memory B cells sorted from healthy blood donors, while the
memory B cells were somewhat more efficient at upregulating CD11c and FcRL5 after 2 days of
stimulation (**supplementary figure 3**). As both naïve and memory B cells could efficiently respond to
the stimuli, and in vivo generated CD11c-, T-bet-, and FcRL5-expressing cells are likely generated
from both naïve and memory B cells [23], we used total B cells isolated from healthy blood donors to
further characterise the response to stimulation (**Figure 2A**). Cell marker expression was evaluated
by flow cytometry after stimulation with anti-Ig (BCR ligation), CpG-C (TLR9-ligation), and the pro-
inflammatory cytokine IFN-γ in comparison to unstimulated cells (**Figure 2B**). The stimulated cells
adopted a CD19Hi phenotype, with a significant percentage of cells having upregulated CD19 on day
2 and day 6 after stimulation compared to unstimulated controls. CD11c and FcRL5 expression was
also significantly higher among stimulated cells at both time points. The T-bet expression on the
other hand was highly upregulated by day 2 but then greatly reduced by day 6, although it remained
significantly higher than unstimulated cells.
160

161 Given the possible difference in kinetics between the markers over time, we evaluated protein levels
162 at 0, 2, 4, and 6 days after stimulation (**Figure 2C**). The percentage of CD11c⁺ B cells increased over
163 time and plateaued between 4 and 6 days of stimulation. FcRL5 and CD19 on the other hand peaked
164 at day 4 after which they started declining. T-bet was rapidly upregulated and peaked at 2 days of
165 stimulation, after which the protein levels declined.
166
167 To assess if the changes in protein levels were associated with cell division, B cells were labelled with
168 cell trace violet (CTV). The B cells stimulated with anti-Ig, CpG-C, and IFN-γ displayed measurable
169 proliferation in the form of CTV dilution at day 4 after stimulation while unstimulated cells remained
170 undivided (**Figure 2D**), clearly indicating that cell division was not a requirement for upregulation of
171 CD19, CD11c, T-bet, or FcRL5.
172

173
Figure 2. In vitro generation of B cells expressing high levels of CD19, CD11c, T-bet, and FcRL5.
*(**A**) Schematic representation of experimental setup with B cells sorted from buffy coats using*
*RosetteSep, followed by stimulation with anti-Ig, CpG-C, and IFN-γ for 2, 4, and 6 days. (**B**) The cells*
were evaluated by flow cytometry for expression of CD19, CD11c, T-bet, and FcRL5 at 2 days (n=19-
20 donors) and 6 days cells (n=10-11 donors) after stimulation (+, orange solid line and circles) or left

*unstimulated (–, dashed blue line and boxes) for unlabelled B cells and (**C**) at 0, 2, 4, and 6 days for*
*cell-trace violet labelled B cells (n=6-9 donors). (**D**) B cell division was determined by dilution of cell-*
trace violet at 0, 2, 4, and 6 days (n=6-9 donors). Data in B indicates individual donors and is pooled
from multiple experiments. Data in C and D indicates the mean with a 95% confidence interval and is
pooled from two separate experiments. Paired t-tests were used to determine p-values between
*unstimulated and anti-Ig/CpG-C/IFN-γ stimulated cells at 2 and 6 days. **p<0.01, ***p<0.001,*
*****p<0.0001.*
186

Role of BCR- and TLR9-ligation on B cell levels of CD19, CD11c, T-bet, and FcRL5

To evaluate the specific influence of BCR- and TLR9-ligation on cell marker expression, stimulation
conditions without anti-Ig or CpG-C, in combination with IFN-γ, were compared to the condition with
all three signals in sorted B cells from healthy blood donors (**Figure 3 and Supplementary table 1**).
CpG-C was shown to be critical in inducing CD19^{hi} cells. On day 2, the removal of either anti-Ig or
CpG-C led to a significant decrease in CD11c levels whereas, on day 6, only the condition without
anti-Ig had significantly fewer CD11c^{+} cells. For T-bet levels, the addition of anti-Ig or CpG-C led to a
significant increase of positive cells on day 2. On day 6, however, T-bet was down-regulated to
baseline levels in all conditions. Both anti-Ig and CpG-C contributed to the up-regulation of FcRL5 on
day 2, but no significant difference between stimulated and unstimulated cells was observed on day
6. In summary, TLR9-ligation was required for the upregulation of all four markers assessed here,
while BCR-ligation led to up-regulation of CD11c, T-bet, and FcRL5, but not CD19.
199
200

Figure 3. Role of BCR- and TLR9-ligation on CD19, CD11c, T-bet, and FcRL5 expression.

*Sorted total B cells from buffy coats were stimulated with combinations of anti-Ig, CpG-C, and IFN-γ for (**A**) 2 days (n=19-20) and (**B**) 6 days (n=10-11) after which expression of CD19, CD11c, T-bet, and FcRL5 was measured by flow cytometry. Representative histograms indicate marker expression. The line indicates the cut-off for positive expression. Numbers indicate the stimulations (see Supplementary table 1). P-values were calculated using matched pair one-way ANOVA with Geisser-Greenhouse correction followed by Sidak's post hoc test. Unstimulated cells (blue boxes) were included for visual reference. Stimulated groups (open circles, stim 3 and 4) were compared with anti-Ig, CpG-C, IFN-γ (orange circles) stimulated cells. *p<0.05, ***p<0.001, ****p<0.0001. The absence of * indicates p>0.05.*

Differential impact of cytokines on B cell CD19, CD11c, T-bet, and FcRL5 levels

To further dissect the impact of the cytokines on the levels of CD19, CD11c, T-bet, and FcRL5, B cells sorted from healthy blood donors were cultured with anti-Ig, CpG-C, and either IFN-γ, IL-21, or IL-10 (**Figure 4 and Supplementary table 1**). None of the tested cytokines led to the upregulation of CD19 or FcRL5. The addition of IFN-γ resulted in a significant increase in CD11c levels, but only on day 2, while stimulation with IL-21 led to a significant increase on both days 2 and 6. IL-10 on the other hand had no significant effect on CD11c, although the mean CD11c$^+$ frequency was somewhat higher than anti-Ig and CpG-C alone on day 2. In contrast, IL-10 stimulation led to a reduction of T-bet$^+$ cells compared with anti-Ig and CpG-C stimulated cells alone. Only IFN-γ out of the 3 cytokines resulted in a significant increase in T-bet$^+$ B cells on day 2, and a slightly higher percentage of cells remaining on day 6. This effect was not adversely affected by IL-10 (**Supplementary Figure 4**). In summary, T-bet expression was highly dependent on IFN-γ, while CD11c was upregulated in response to both IFN-γ and IL-21. CD19 and FcRL5 on the other hand were not influenced by the tested cytokines.

Figure 4. Role of IFN-γ, IL-21, and IL-10 on CD19, CD11c, T-bet, and FcRL5 expression.

Sorted total B cells from buffy coats were stimulated with combinations of anti-Ig, CpG-C, IFN-γ, IL-
*21, and IL-10 for (**A**) 2 days (n=10-20) and (**B**) 6 days (n=10-11) after which expression of CD19,*
CD11c, T-bet, and FcRL5 was measured by flow cytometry. Representative histograms indicate
marker expression. The line indicates the cut-off for positive expression. Numbers indicate the
stimulations. P-values were calculated using matched pair one-way ANOVA with Geisser-Greenhouse
correction followed by Sidak´s post hoc test or in the case of missing data, by mixed-effects analysis.
Unstimulated cells (blue boxes) were included for visual reference. All stimulated groups had received
anti-Ig and CpG-C. Stimulations 2 (orange circles), 6, and 7 (open circles) were compared with anti-Ig
*and CpG-C stimulated cells alone (grey circles). *p<0.05, **p<0.01, ****p<0.0001. The absence of **
indicates p>0.05.

Dynamic CD11c, T-bet, and FcRL5 co-expression profiles in response to stimulation and time

Different combinations of markers are used to describe the B cells that expand during inflammation.
Here we analysed how the co-expression of CD11c, T-bet, and FcRL5 change depending on
stimulatory conditions, at two different time points (**Figure 5**). The largest proportion of triple-
positive cells was generated by stimulating the B cells with anti-Ig, CpG-C, and IFN-γ for 2 days
(approximately 26%). This condition also had the lowest percentage of cells not expressing any of
CD11c, T-bet, or FcRL5 at both 2 and 6 days of stimulation. Compared to 2 days of stimulation, there
were few cells expressing all three markers at 6 days. This was mainly explained by the rapid
reduction of T-bet⁺ cells after day 2. Instead, the day 6 time-point was characterised by similar

frequencies of CD11c single- or CD11c and FcRL5 double-positive B cells. Most of the unstimulated cells remained triple-negative at both time points (88.6% and 76.6% respectively).

Figure 5. Stimulation-dependent co-expression of CD11c T-bet, and FcRL5.
*The average frequency of CD11c, T-bet, and FcRL5 co-expression on B cells sorted from buffy coats following stimulation for (**A**) 2 days (n=18) and (**B**) 6 days (n=9) with different combinations of anti-Ig, CpG-C, and IFN-γ.*

Plasma cell differentiation and antibody production are dependent on CpG-C and B cell proliferation

The functional properties of double negative or atypical (CD21⁻CD27⁻IgG⁺ and IgD⁻CD27⁻T-bet⁺) B cells remain unclear. It has been suggested that the cells contribute to the circulating antibody pool [34] and that they could represent an intermediate plasma cell differentiation stage [35]. We, therefore, assessed plasma cell differentiation and antibody production in the context of anti-Ig, CpG-C, and IFN-γ stimulation and CD11c, T-bet, and FcRL5 expression. We cultured CTV-labelled and unlabelled B cells sorted from healthy blood donors over 6 days with different combinations of anti-Ig, CpG-C, IFN-γ, and IL-21. We then assessed CTV dilution (indicating proliferation), plasma cell differentiation at 0, 2, 4, and 6 days of culture, and antibody production at 6 days of culture (**Figure 6A**). After subtracting media controls, we compared pooled log2 transformed antibody concentrations (**Figure 6B**). anti-Ig, CpG-C, and IFN-γ stimulated cells (stim. 2) produced significantly more antibodies compared to unstimulated cells (stim. 1). Exchanging IFN-γ with IL-21 (stim. 7) led to further increased antibody levels, although IFN-γ did not have a detrimental effect on the antibody levels (stim. 7 vs stim. 9). Antibody production was however completely abolished in the absence of CpG-C, independent on other stimuli (stim. 4 and 11), indicating that CpG-C was required to plasma cell differentiation.

We next assessed plasma cell differentiation by flow cytometry. Plasma cells were identified by high levels of CD27, which was only observed in CTVˡᵒʷ proliferating cells (**Figure 6C**). Quantifying CTVˡᵒʷCD27ʰⁱ plasma cells after stimulation showed that plasma cells became detectable and reached peak levels at day 4 or stimulation. Plasma cells were, however, only observed after stimulation with anti-Ig, CpG-C, and IFN-γ (stim. 2) and not anti-Ig and IFN-γ alone (stim. 4), or in unstimulated cells. This corroborates the antibody data and shows that CpG-C was critical for B cell proliferation and plasma cell differentiation (**Figure 6D**).

Since it has been indicated that T-bet contributes to plasma cell differentiation [35], we assessed if
CD11c, T-bet, or FcRL5 were specifically enriched in plasma cells compared with total live B cells at
day 4 of culture, when all markers were expressed and at peak of plasma cell frequencies (**Figure**
6E). Cells expressing either CD11c, T-bet, or FcRL5 were similarly enriched among the two
compartments, indicating that none of the proteins was specifically required for plasma cell
differentiation (**Figure 6E**).
292

293
294
Figure 6. CpG-C is required for plasma cell differentiation and antibody production.
*(**A**) Schematic illustration of B cell cultures with cell-trace violet labelled and unlabelled B cells sorted*
*from buffy coats, stimulated for 0, 2, 4, and 6 days before (**B**) measuring IgG1, 2, 3, 4, IgA, and IgM*
antibody levels in culture supernatants associated with different combinations of stimulation with
*anti-Ig, CpG-C, IFN-γ, and IL-21 as indicated (n=10). (**C**) Representative flow cytometry plots from day*
*6 and (**D**) summary graph indicating the frequency of cell-trace violet low and CD27 high plasma*
blasts at 0, 2, 4, and 6 days with no stimulation (blue boxes), anti-Ig+CpG-C+IFN-γ(orange circles), or
*anti-Ig+IFN-γ stimulation (grey triangles) (n=9). (**E**) Percent CD11c, T-bet, or FcRL5 expressing B cells*
within total live B cells (empty circle) or CTV^{low}CD27^{hi} cells (green circle) at day 4 after culture with
anti-Ig, CpG-C, and IFN-γ. P-values were calculated using repeated-measures one-way ANOVA with
Geisser-Greenhouse correction followed by Holm-Sidak's post hoc test. ns=not significant p>0.05,
*p<0.05, ***p<0.001.
307
B cell stimulation leads to the upregulation of co-stimulatory molecules and cytokine secretion
resulting in increased T cell help
In addition to plasma cell differentiation, it has been proposed that CD11c[+] B cells can provide
enhanced antigen-presenting functions to CD4[+] T cells in mice and humans [19,21]. To assess to
what extent B cells stimulated with different combinations of anti-Ig, CpG-C, and IFN-γ could drive B
cell-dependent CD4[+] T cell help, we first stimulated B cells from healthy blood donors for 2 days. We
then phenotyped the cells, measured cytokines in the supernatants and washed, counted, and co-
cultured the B cells at a 1:5 ratio with autologous purified CD4[+] T cells in the presence of cytostim,
crosslinking MHC2 and the TCR (**Figure 7A**). To measure proliferation, the CD4[+] T cells were stained
with cell trace violet (CTV) (**Figure 7B**). As a negative control, CD4[+] T cells were cultured with

cytostim in the absence of B cells, and as a positive control CD4$^+$ T cells were cultured with beads
coated with anti-CD3 and anti-CD28. Unstimulated B cells co-cultured with CD4$^+$ T cells had a similar
frequency of CTV low cells as the negative control, while > 90% of the cells had divided in response
to the positive control (**Figure 7B**). B cells pre-stimulated with anti-Ig, CpG-C and IFN-γ (stim. 2)
induced significantly more T cell proliferation (CTV^{low} cells) compared with co-cultures with
unstimulated B cells. There were also significantly more dividing T cells in the co-cultures with B cells
pre-stimulated with anti-Ig and IFN-γ (stim. 4) when compared with unstimulated cells (**Figure 7C**).
325
In vivo, B cells and CD4$^+$ T cells interact via MHC2 and the TCR, with additional survival and polarising
signals provided via cytokine secretion and co-stimulatory molecules, such as CD80 and CD86 on the
B cells [36,37]. To better understand what factors were responsible for the increased B cell-
dependent CD4$^+$ T cell proliferation, we assessed the levels of CD80 and HLA-DR after stimulation
(**Figure 7D**). CD80 was strongly upregulated by anti-Ig, CpG-C, and IFN-γ (stim. 2). The upregulation
was independent of IFN-γ as stimulation with anti-Ig and CpG (stim. 5) reached a similar frequency of
CD80$^+$ cells, while anti-Ig and CpG-C seemed to have an additive effect (**Figure 7D**). All B cells express
HLA-DR, so therefore we assessed the surface levels (median fluorescent intensity, MFI) following
stimulation after subtracting with the MFI of unstimulated cells (**Figure 7E**). All stimulations that we
tested had an average HLA-DR log2(MFI^{stim}–MFI^{unstim})>0, indicating upregulation of HLA-DR in
response to stimulation. Cells stimulated with anti-Ig and CpG (stim. 5) had a similar HLA-DR MFI
level as triple-stimulated cells, while the condition gave significantly more upregulation compared
with CpG and IFN-γ (stim. 3), anti-Ig and IFN-γ (stim. 4), or CpG-C alone (stim. 6) (**Figure 7E**).
339
We next assessed a panel of 13 cytokines in the culture supernatants from day 2 of the B cell culture
with the rationale that this could indicate which cytokines were produced by the B cells at the time
of co-culture with the T cells. Cytokines that were detected above media background levels
(log2[pg/ml^{stim}–pg/ml^{media}]>0) were pooled together to give a bulk measure of B cell cytokine
production (**Figure 7F**). IFN-γ was omitted from the cytokine pool data, since it was spiked in for
some stimulations, and is shown as relative concentration (MFI) in **Supplementary Figure 5** together
with the other cytokines measured. All stimulations had significantly more cytokines in the culture
supernatant compared with unstimulated cells (p<0.0001). anti-Ig, CpG-C, and IFN-γ (stim. 2)
stimulated cells had significantly more cytokines than all other conditions (stim. 2 vs stim. 3 p<0.05,
stim. 2 vs stim. 4-6 p<0.001). CpG-C + IFN-γ (stim. 3) and anti-Ig + CpG-C (stim. 5) had similar levels of
cytokines (p>0.05) and anti-Ig + CpG-C (stim. 5) significantly more than CpG-C alone (stim. 6 p<0.05),
indicating that anti-Ig, CpG-C, and IFN-γ all contributed to increasing overall cytokine production
(**Figure 7F**). Cells stimulated with anti-Ig and IFN-γ (stim. 4) had the least total cytokine levels of the
stimulated cells. This was mainly due to no detectable production of IL-10, which made up 33-39 %
of the measured cytokines in the other stimulations, where CpG-C was present. IL-6 was the other
main cytokine produced by the B cells, making up 39-64 % of the measured cytokines for the
stimulated cells (**Figure 7F**).
357
To test if IL-6 was potentially important for the improved CD4$^+$ T cell help, IL-6 blocking antibodies or
isotype control antibodies were added to co-cultures of pre-stimulated B cells and CTV-labelled CD4$^+$
T cells. Both the isotype antibody-treated and untreated co-cultures had a significantly higher
frequency of dividing CTV^{low} T cells when cultured with anti-Ig, CpG-C, and IFN-γ pre-stimulated B

cells compared with unstimulated B cells (**Figure 7G**). Treatment with IL-6 blocking antibodies,
however, abolished this effect by significantly reducing the frequency of CTV^{low} T cells to baseline
levels, comparable to T cells co-cultured with unstimulated B cells (**Figure 7G**). This indicates that IL-
6 was a key factor in the B cell-driven CD4$^+$ T cell proliferation.
366

Figure 7. B cell-dependent CD4$^+$ T cell proliferation is mediated via IL-6.
(A) Schematic illustration of B cell-T cell co-culture experiments of cells sorted from buffy coats. (B)
representative histograms and (C) data for cell-trace violet dilution among sorted CD4$^+$ T cells co-
cultured for 5 days with autologous pre-stimulated B cells in the presence of cytostim (n=10). B cells
were pre-stimulated for 2 days as indicated, followed by washing, counting, and re-seeding at a 1:5
B:T ratio. (D) Frequency of CD80-positive B cells after two days of stimulation (n=19). (E) B cell
surface levels of HLA-DR, calculated as log2[MFI^{stim}]–log2[MFI^{unstim}], after 2 days of culture (n=19). (F)
Cytokine levels for sCD40L, APRIL, IL-12p70, IL-2, IL-6, IL-10, IL-4, IL-13, and TNF-α in culture
supernatants calculated as log2[pg/mL in stimulated]–log2[pg/mL in culture media] following 2 days
of culture (n=10). (G) The same co-culture setup as in (C) with no stimulation (blue boxes) or anti-
Ig+CpG-C+IFN-γ (orange circles) pre-stimulated B cells in the presence of autologous CD4$^+$ T cells and
No antibody, isotype control antibody, or IL-6-blocking antibody (n=5). P-values were calculated
using repeated-measures one-way ANOVA with Geisser-Greenhouse correction followed by Sidak's
post hoc test. For (C), all comparisons were against unstimulated cells. ns=not significant p>0.05,
*p<0.05, **p<0.01, ***p<0.001, ****p<0.0001.
384

Discussion

In this study, we have described how the cell surface proteins CD11c and FcRL5 and the intracellular transcription factor T-bet are dynamically regulated after infection and following in vitro stimulation via different activation pathways. These proteins are commonly used in different combinations together with CD21, CD27, and IgD to identify a subset of B cells expanding during inflammatory conditions [23]. In a previous study of CD11c$^+$ B cells in malaria, we observed that these cells displayed different co-expression with FcRL5 and CXCR3 over time [27]. This led us to hypothesize that the expression of these markers could be regulated via different activation pathways, leading to different patterns of expression over time.

We first investigated the co-expression during and up to 12 months after diagnosis with *Plasmodium* parasite infection. The frequency of cells expressing either CD11c, T-bet, or FcRL5 was the same at approximately 30% of the mature B cells at diagnosis and at 10 days after treatment initiation. This was then reduced to 18% at 12 months. However, despite a similar frequency of B cells with these markers at diagnosis and the 10-day time-point, there was a substantial shift in the co-expression pattern with mostly triple-positive cells at diagnosis followed by CD11c and FcRL5-double positive cells at the 10-day time-point. This observation could have several different explanations, such as a more transient expression of T-bet compared with CD11c and FcRL5, that cells expressing T-bet leave the circulation, that T-bet-expressing cells preferentially differentiate to plasma cells, or that they rapidly die.

We have previously shown that many of the dividing B cells during and after acute malaria express CD11c and are enriched for parasite-specific reactivities, indicating a BCR-dependent upregulation of CD11c [27]. Similar observations of recently activated B cells being enriched for CD11c$^+$ vaccine or infection-specific B cells have also been made with antigen-specific tetramer staining [28,38,39]. It is further also consistent with stimulation of the BCR in vitro leading to CD11c upregulation [21]. FcRL5 has been shown to be transiently upregulated via BCR-ligation where it upon co-stimulation via the BCR and TLR9 led to a strong proliferation of naïve B cells [40]. FcRL5 is an IgG-receptor, enabling B cells to respond to immune complexes [41]. In conjunction with CD21 (complement receptor 2), which binds C3-fragments this effect is further enhanced [42], suggesting that early upregulation of FcRL5 could be important to prime B cells for more efficient uptake and subsequent responses to opsonized antigens. CD11c is an integrin and potentially has complementary roles to FcRL5 in enhancing phagocytosis and cell migration [43] that could improve the early B cell response to infection.

During malaria, the high abundance of parasites and subsequent pathogen-associated molecular patterns (PAMPs), such as parasite DNA, will stimulate innate immune cells, including monocytes, DCs, and B cells via the expression of endosomal TLR9 and other pattern-recognition receptors [44–47]. Signaling via TLR9 is important for parasite control in both mice [48] and humans, where different polymorphisms in human TLR9 greatly influence symptomatic disease [49]. Binding to the BCR, followed by subsequent antigen uptake into endosomes and signaling via TLR9 constitutes early cellular events that the B cells can do independently of other cellular interactions. In this study, we found that these events were the primary influence on B cell expression of CD19, FcRL5, CD11c, HLA-DR, CD80, and B cell-derived cytokine production, indicating a role for all these effects in the early B cell response to infection. BCR-ligation also leads to the upregulation of CCR7 and EBI2, which will

guide the activated B cells to the B cell-T cell interfollicular region in secondary lymphoid organs
[50,51], where they can interact with cognate CD4+ T cells [51,52].

Prior to their migration to the B cell-T cell interfollicular region, CD4+ T cells are primed by activated
DCs that have taken up and recognized PAMPs. In response to intracellular pathogens, such as
Plasmodium parasites, DCs produce IL-12, which in turn will polarize the CD4+ T cells to become IFN-
γ secreting T_H1 cells [53]. IL-12 together with IFN-γ can then further generate $T_{FH}1$ cells producing
both IFN-γ and IL-21 [54]. During malaria, circulating $T_{FH}1$ cells, identified by CXCR3-expression are
expanded in peripheral blood and are shown to provide suboptimal B cell help compared with
conventional CXCR3-negative cT_{FH} cells [55]. Especially in children where $T_{FH}1$ cells made up a larger
fraction of circulating T_{FH} cells [56]. IFN-γ has previously been shown to be important for T-bet
expression in human B cells [11] and $T_{FH}1$ cells likely constitute an important driver for the expansion
of T-bet+ B cells in malaria and other inflammatory diseases [57]. Mouse models have in addition to
IFN-γ also indicated a role for IL-21 in the generation of CD11c+T-bet+ B cells [58–60]. We similarly
show that IFN-γ is critical for optimal T-bet expression but that it is not induced by IL-21 in the
absence of IFN-γ. CD11c on the other hand was more strongly upregulated in response to IL-21 than
IFN-γ, indicating separate roles of the cytokines and consistent with different regulations of CD11c,
FcRL5, and T-bet. In support of this, other studies in mice have identified a population of CD11c+ B
cells in the absence of T-bet [60,61], and recently, Gao at al. and Gu et al. identified CD11c to likely
be regulated via the transcription factor Zeb2 in humans and mice, respectively [62,63].

IL-21 is a key cytokine in germinal centre B cell survival and plasma cell differentiation [64]. It is
mainly produced by CD4+ T_{FH} cells and its induction is dependent on interaction with cognate B cells
[65]. It was shown that CD11c+ B cells can become plasma cells in the presence of IL-21 and T cell
help [18,21,57,66]. Other studies have indicated a contribution of CD21−CD27− or CD11c+T-bet+ B
cells to the circulating antibody pool [34,67,68], indicating either active plasma cell differentiation or
that the cells are coming from the same lineage as antibody-secreting cells. Further supporting an
active plasma cell differentiation, it has been suggested that T-bet+ B cells in systemic lupus
erythematosus (SLE) could correspond to intermediate-stage plasma cells and that regulation of the
IFN-γR by T-bet is important for subsequent full plasma cell differentiation [35]. We observed an
enhanced effect of IL-21 on antibody production that was unaffected by the presence or absence of
IFN-γ in the stimulation and subsequent varying levels of T-bet (lower in the absence of IFN-γ).
Additionally, when measuring the expression of CD11c, T-bet, and FcRL5 in plasma cells or other
dividing cells responding to the stimulations, there was no enrichment of the markers among plasma
cells, suggesting that none of the proteins was actively required for plasma cell differentiation in
vitro. However, *in vivo* it is possible that co-expression of these markers is a common occurrence
during inflammatory conditions as antigen-specific B cells would upregulate CD11c and FcRL5, after
which they would migrate to the interfollicular region where they would encounter T cells producing
IFN-γ and IL-21, driving upregulation of T-bet and potentially leading to extrafollicular plasma cell
differentiation.

In response to the proinflammatory environment generated by the malaria parasites, there is also an
expanded anti-inflammatory response characterized by regulatory cells [69,70], modulation of
innate immune responses [71–73] and secretion of soluble anti-inflammatory proteins, such as IL-10

[71,74]. In mice, an increased level of IL-10 is shown to inhibit T-cell derived IFN-γ, leading to
reduced T-bet[+] B cells [75,76], indicating a potentially dynamic regulation of T-bet[+] B cells *in vivo*. We
also observed a reduced T-bet induction in B cells stimulated with IL-10, but not when co-stimulated
with IFN-γ. It has been suggested in mouse and human studies of malaria that both B cells and T cells
are major sources of IL-10 and are needed for disease control [69,77,78]. We observed a strong
induction of IL-10 in all stimulations containing CpG-C, consistent with TLR-ligands being major
inducers of IL-10 production in B cells [79,80]. Polymorphisms in both TLR9 and IL-10 are associated
with clinical presentation in malaria, highlighting the importance of this pathway in malaria disease
control [49,81]. In addition to IL-10, we also observed a large production of IL-6 from the stimulated
B cells. The IL-6 levels were also higher with CpG-C stimulation, but they were not completely CpG-
dependent, as observed for IL-10. IL-6 is a pleiotropic cytokine that affects both hematopoietic and
non-haematopoietic cells. It is an important survival and differentiation factor for B cells [82] and
CD4+ T cells, where it enhances proliferation, and influences migration and polarization [83]. By
blocking IL-6, we completely abrogated the B cell-mediated CD4[+] T cell proliferation, indicating that
IL-6 was a key mediator of how the B cells provided T cell help in our in vitro system.
489
It has been shown that CD11c[+] B cells can provide enhanced APC functions in mice [19]. We also
found that in vitro-stimulated B cells expressing CD11c, T-bet, and FcRL5 could support improved
CD4[+] T cell survival and proliferation via the production of IL-6. However, there was no obvious
association with T-bet, as stimulations lacking IFN-γ, and therefore were relatively low for T-bet,
provided T cell help equally well as stimulation with IFN-γ and high T-bet levels. Instead, the
improved T cell proliferation was mainly associated with B cells pre-stimulated via the BCR. This was
not clearly associated with specific up-regulation of CD11c, FcRL5 or specific levels of IL-6 either. It is
therefore likely that additional factors, driven by BCR-signalling and not identified here, can provide
further T cell help in conjunction with IL-6.
499
B cells expressing CD11c, T-bet, and or FcRL5 remain an enigmatic part of the B cell response. In this
study, we have evaluated co-expression after infection and triggered different activation pathways
in vitro to learn about how the markers are dynamically regulated over time. We found that the
stimuli leading to the upregulation of CD11c and FcRL5 were relatively overlapping, with BCR and
TLR9-ligation being the most important, while T-bet was dependent on IFN-γ. We also investigated
the stimulated B cells functionally and observed a clear dependency of plasma cell differentiation on
proliferation, while T cell help was dependent on the production of IL-6. However, the effector
functions were not clearly associated with given patterns of CD11c, T-bet, or FcRL5 expression,
potentially indicating that the upregulation of these markers is part of a normal response and not a
specific B cell lineage. However, to better understand how these cells behave in humans *in vivo*,
further studies are needed where B cells with different receptor expressions, potentially indicating
different stages of differentiation or time after antigen encounter, are characterised for specificity,
phenotype, and function.
513

Materials and Methods

Clinical samples

Adult travellers returning to Sweden and presenting with malaria at the emergency department at
Karolinska University Hospital and migrants identified through malaria screening were enrolled in a
prospective study. Malaria parasite species and parasitaemia were determined by microscopy
and/or qPCR by the Karolinska University Hospital lab. Blood samples were collected at the time of
diagnosis, after 10 days (10D), and again after 12 months (12M). In this study, samples from 10
patients, at three different time points, and 6 healthy controls were included for analysis. Venous
blood was collected in EDTA vacutainer tubes (BD) and peripheral blood mononuclear cells (PBMCs)
were isolated by density gradient centrifugation using Ficoll–Paque Plus (Cytiva). Briefly, the blood
was transferred to 50 ml tubes and diluted 1:1 with phosphate-buffered saline (PBS) supplemented
with 2% heat-inactivated fetal bovine serum (FBS), both from Thermo Fisher Scientific. 15 ml of
Ficoll-Paque was added to 50 ml Leucosep tubes and centrifuged at 1000 g for 1 minute at room
temperature (RT). The diluted blood was poured into the Leucosep tubes on top of the Leucosep
filter and centrifuged at 800 g for 15 minutes at RT without breaking at the end. The mononuclear
cell layer at the plasma-Ficoll interface was removed using a Pasteur pipette. The cells were diluted
to 50 ml with PBS + 2% FBS and centrifuged at 300 g for 8 min at RT. Supernatants were discarded
and cell pellets were resuspended in 30 ml PBS + 2% FBS and centrifuged again at 250 g for 10
minutes at RT. After the final wash, the cells were resuspended in PBS + 2% FBS and counted using a
Countess II (Invitrogen) following a 1:1 dilution with trypan blue. The cells were centrifuged again
and resuspended at 5×10^6 cells/ml in freezing medium (FBS + 10% DMSO) and placed inside a
CoolCell at −80°C overnight, after which the cells were transferred to liquid nitrogen for long-term
storage.

Samples – in vitro cultures

For the in vitro parts of the study, cells were sorted from buffy coats from healthy donors. Buffy
coats were obtained from the Karolinska University Hospital blood bank. Complete media that was
used in all cultures consisted of RPMI-medium supplemented with 10% heat-inactivated FBS, 2mM
L-glutamine, 100 U/ml of penicillin, 100 µg/ml of streptomycin, 10 mM HEPES and 0.05 mM 2-
Mercaptoethanol (all from ThermoFisher Scientific).

Total B cell and CD4[+] T cell isolation

B cells and CD4[+] T cells were isolated from buffy coats by negative selection with the RosetteSep
Human B Cell Enrichment Cocktail or RosetteSep Human CD4[+] T Cell Enrichment Cocktail
respectively, using 50 ml SepMate tubes (all from Stemcell technologies). Briefly, 25 µl of the Cell
Enrichment Cocktail was added for each ml of blood and incubated for 20 minutes at RT. Two
volumes of PBS was then added to the sample. 15 ml Ficoll–Paque Plus (GE Healthcare) was added
to SepMate tubes and centrifuged at 1000 g for 1 minute (RT). Samples were poured into the
SepMate tubes and centrifuged at 1200 g for 10 min with the break on. The cell fraction was
transferred to new tubes and washed twice with PBS + 2% FBS (300 g, 10 min, brake at 5, RT). For
two donors the B cells were isolated using MACS cell sorting with anti-human CD19 microbeads and
MS columns from Miltenyi Biotec, according to the manufacturer's instructions. Isolated cells were
frozen and stored as previously described. B cell and CD4[+] T cell purity was assessed by flow
cytometry and were on average±SD 90.8±5.0% and 90.5±4.8%, respectively.

Naïve and memory B cell isolation

Memory B cells and naïve B cells were isolated from RosetteSep pre-sorted total B cells via positive
selection using the EasySep™ Human Memory B Cell Isolation Kit (STEMCELL™ Technologies), and
via negative selection using the EasySep™ Human Naive B Cell Isolation Kit (STEMCELL™
Technologies), following the manufacturer's protocol. Briefly, total B cells were prepared as
previously described and resuspended at a concentration of 1×10^8 cells/mL. CD27 Positive Selection
Cocktail was then added at 50 µL/mL and incubated for 5 minutes, after which 100 µL/mL of
Releasable RapidSpheres were added. Total B cells were then incubated in an EasySep magnet for 5
min. The negative fraction was poured into a new tube and kept for naïve B cell isolation. The tube
containing isolated memory B cells was incubated with 100 µL/mL of Release buffer and 50 µL/mL of
Dextran RapidSphere for 5 minutes in an Easysep magnet. The isolated memory B cells released
from the selection beads were poured into a new tube and directly used for stimulation. The cells in
the negative fraction (from the memory B cell isolation) were resuspended at 5×10^7 cells/mL and 50
µL/mL of Cocktail Enhancer and Isolation Cocktail were added. After 5 minutes of incubation, 50
µL/mL of RapidSpheres was added and tubes were placed in an Easysep magnet for 5 minutes.
Negatively selected Naïve B cells were poured into new tubes and were directly used for stimulation.
Memory and Naïve B cell purity was assessed by flow cytometry and were on average±SD 86.4±5.2
% and 92.7±6.9 %, respectively. The buffer used during both isolations was PBS + 2% FBS.

B cell stimulation

Previously sorted total B cells were thawed, moved to a 15 ml Falcon tube and 13 ml of complete
media was added dropwise before mixing and resting the cells on ice for 20 min. The sorted B cells
were then centrifuged at 300 g for 5 min, washed once in complete media (300 g 5 min),
resuspended in 1 ml complete media and counted. Stimulations of purified B cells were done in a 96-
well round-bottom tissue-culture plate (Techno Plastic Products). The stimuli included α-Ig
(IgA+IgG+IgM [H+L] AffiniPureF(ab')2 Fragment goat-anti-human [Jackson ImmunoResearch
Laboratories]) at 1.25 µg/ml, CpG-C (Invivogen) at 2.5 µg/ml, IFN-γ (Peprotech) at 25 ng/ml, IL-21
(Peprotech) at 10 ng/ml and IL-10 (Peprotech) at 10 ng/ml. The stimuli were added in different
combinations (**Supplementary Table 1**) to wells seeded with 1×10^5 cells in a total volume of 200 µl
complete media. Unstimulated cells were cultured in complete media. Plates were incubated at 37°C
and 5% CO_2.

Flow cytometry

For flow cytometry, the cells were washed twice in PBS and then incubated with Live/Dead stain
(ThermoFisher Scientific), diluted 1:1200 in PBS, for 20 min on ice. Cells were then washed twice in
PBS + 2% FBS before incubation for 20 min on ice with an antibody mix targeting cell surface
markers. For intracellular markers, the cells were washed once after surface staining, then incubated
for 30 min at RT with the FoxP3 Fix/Perm buffer (eBioscience), washed once in FoxP3 Wash/perm
buffer, and incubated for 30 min at RT with an antibody mix targeting intracellular markers. The cells
were then washed again in Wash/perm buffer followed by one wash in PBS + 2% FBS. Samples were
then resuspended in 300 µl PBS + 2% FBS together with 5 µl CountBright™ Absolute Counting Beads
(ThermoFisher Scientific) and run on a 5-laser BD Fortessa (BD Biosciences). See **Supplementary**
Tables 2-7 for the different antibody panels.

Cell Trace Violet

For tracking cell division, the cells were stained with Cell Trace Violet (CTV) (Thermo Fisher

Scientific). The cells were suspended at a density of 2×10^6 cells per ml in PBS. CTV was added to the

cell suspension at 1 mM and incubated for 7 min at 37°C in a ventilated incubator. Staining was

stopped with FBS (3 times the staining volume) and PBS was added up to 15 ml. The cells were then

centrifuged (300 g 5 min) and washed once in PBS (300 g 5 min) and once in complete media. The

cells were then rested for 1 h in complete media at 37°C 5% CO_2 before further simulation or

analysis.

Antibody measurement

Antibody levels in the supernatants from B cells that had been stimulated for 6 days were measured

using the LEGENDplex™ Human Immunoglobulin Isotyping Panel (6-plex) (Biolegend), a bead-base

immunoassay, according to the manufacturer's protocol with some modifications. The assay was

performed in a U-bottom 96 well plate, the plate was wrapped in aluminium foil during all

incubations. Samples were diluted 1:100 in PBS + 2% FBS. 25 μl diluted sample was mixed with 25 μl

assay buffer and 12.5 μl pre-mixed beads. A standard 1:4 serial dilutions were prepared, and 25 μl

standard was mixed with 25 μl assay buffer and 12.5 μl pre-mixed beads. The plates were incubated

on a plate shaker at 700 rpm for 2 h, RT. Plates were then centrifuged at 650 g, 1 min and washed

with 200 μl wash buffer. A detection antibody was added at 1:1 in PBS, 25 μl/well. The plates were

then incubated on a plate shaker at 700 rpm for 1 h, RT. Streptavidin-PE was then added at 1:1 in

PBS, 25 μl/well (no wash in between), and again incubated on a plate shaker at 700 rpm for 30 min,

RT. The plates were then washed twice, as before, and resuspended in 200 μl PBS. The fluorescent

signal was measured on a BD 4-laser Fortessa using the 488 nm laser (forward scatter vs side scatter)

to separate beads based on size and granularity and the 640 nm laser (780/60 filter) to separate

beads based on cytokine specificity. The 561 nm laser (582/16) filter was then used to detect the

signal, translating to the amount of cytokine. The data was analyzed using the LEGENDplex™ data

analysis software.

Cytokine measurement

635 Cytokine levels in the supernatants from B cells that had been stimulated for 48 hr were measured

636 using the LEGENDplex™ Human B cell Panel (13-plex) (Biolegend), using the same procedure as

637 described for the antibody measurements, but with the addition of 12.5 μl Matrix B7 to each

638 sample. Supernatants were not diluted prior to being added to the plate and mixed with pre-mixed

639 beads, assay buffer and Matrix B7.

Co-cultures – in vitro stimulated B cells and CD4[+] T cells

In vitro stimulated B cells were co-cultured with autologous CD4[+] T cells and a superantigen to

explore the capacity of in vitro stimulated B cells to induce CD4[+] T cell proliferation. B cells were

stimulated in vitro (as previously described) for 48 h. The cells were then washed twice in fresh

complete media and counted. 20'000 B cells from each condition were cultured for five days with

100'000 CTV-labelled CD4[+] T cells along with 1 μl CytoStim™ (200X dilution, Miltenyi Biotec).

TransAct (300X dilution, Miltenyi Biotec) was used as a positive control for CD4[+] T cell proliferation.

To test the hypothesis that IL-6 signalling influences CD4[+] T cell proliferation, anti-hIL-6-IgG

(Invivogen) or mouse IgG1 kappa Isotype Control (eBioscience) were added at 1 µg/ml to the co-
culture system at the start, and another 1 µg/ml of anti-hIL-6-IgG (Invivogen) or mouse IgG1 kappa
Isotype Control were added to the wells after 2 days of co-culture.
652
Data analysis
Statistical analyses and graphs were done using GraphPad Prism (version 9). Data is generally
presented as individual data points or mean with error bars indicating the 95% confidence interval.
Groups were compared using repeated-measures one-way ANOVA with Geisser-Green house
correction of unequal variance or a mixed-effects model if including missing data, followed by
Sidak's test for multiple comparisons, or a Kruskal-Wallis or Friedman test followed by Dunn's post
hoc test for non-parametric data. A p-value < 0.05 was considered significant. Analysis of flow
cytometry data was done with FlowJo (version 10.8.1, BD). Alluvial plots were generated using the
ggalluvial package [84] in R (version 4.1.1, www.r-project.org).
662
663   **Data availability statement**
664   Data is available upon reasonable request to the corresponding author (christopher.sundling@ki.se).
665
666   **Conflict of interest disclosure**
667   The authors declare no competing interests.
668
669   **Ethics approval statement**
670   This study was performed in accordance with the declaration of Helsinki and following ethical
671   approval (2006/893-31/4 with addendum 2019-03436) by the Swedish Ethical Review Authority.
672
673   **Patient consent statement**
674   All study participants were included following informed consent.
675

Author contributions
LK planned and performed experiments, performed analysis, and wrote the first draft. ADCG
planned and performed experiments and wrote parts of the first draft. MJL performed analysis and
wrote parts of the first draft. AF was responsible for the malaria cohort and supervised students. CS
planned the study, performed analysis, supervised students, and wrote parts of the first draft. All
authors contributed to editing the first draft.

Acknowledgements
We would like to acknowledge the contribution of all the study participants. We would also like to
thank all involved clinicians and nurses, especially Irene Nordling and Debbie Ribjer for sampling and
Fariba Foroogh for malaria sample processing and organization. We would also like to thank Nora
Shahidi for performing experiments and Isabelle Eliasson for collecting patient information. Many
figures were created with Biorender.com. ADCG and MJL were supported by PhD grants from
Karolinska Institutet to CS (2020-00878) and to AF (2019-00992), respectively. The research was
supported by grants from the Swedish Research Council (2019-01940), the Magnus Bergvall
Foundation (2017-02043 and 2018-02656), and the Åke Wiberg Foundation (M18-0076) to CS and by
grants from the Swedish Research Council (2015-02977, 2018-02688, and 2018-04468), the

Stockholm Region (20150135 and 20180409), and the Marianne and Marcus Wallenberg Foundation to AF.

References

1. Akkaya M, Kwak K, Pierce SK. B cell memory: building two walls of protection against pathogens. Nat Rev Immunol. 2019;20: 229–238. doi:10.1038/s41577-019-0244-2

2. de Silva NS, Klein U. Dynamics of B cells in germinal centres. Nat Rev Immunol. 2015;15: 137–148. doi:10.1038/nri3804

3. Roco JA, Mesin L, Binder SC, Nefzger C, Gonzalez-Figueroa P, Canete PF, et al. Class-Switch Recombination Occurs Infrequently in Germinal Centers. Immunity. 2019;51: 337-350.e7. doi:10.1016/j.immuni.2019.07.001

4. Shinnakasu R, Inoue T, Kometani K, Moriyama S, Adachi Y, Nakayama M, et al. Regulated selection of germinal-center cells into the memory B cell compartment. Nat Immunol. 2016;17: 861–869. doi:10.1038/ni.3460

5. Sundling C, Lau AWY, Bourne K, Young C, Laurianto C, Hermes JR, et al. Positive selection of IgG+ over IgM+ B cells in the germinal center reaction. Immunity. 2021;54: 988-1001.e5. doi:10.1016/j.immuni.2021.03.013

6. Kräutler NJ, Suan D, Butt D, Bourne K, Hermes JR, Chan TD, et al. Differentiation of germinal center B cells into plasma cells is initiated by high-affinity antigen and completed by Tfh cells. J Exp Med. 2017;214: 1259–1267. doi:10.1084/jem.20161533

7. Lau AW, Brink R. Selection in the germinal center. Current Opinion in Immunology. Elsevier Ltd; 2020. pp. 29–34. doi:10.1016/j.coi.2019.11.001

8. Elsner RA, Shlomchik MJ. Germinal Center and Extrafollicular B Cell Responses in Vaccination, Immunity, and Autoimmunity. Immunity. 2020;53: 1136–1150. doi:10.1016/J.IMMUNI.2020.11.006

9. Portugal S, Obeng-Adjei N, Moir S, Crompton PD, Pierce SK. Atypical memory B cells in human chronic infectious diseases: An interim report. Cell Immunol. 2017;321: 18–25. doi:10.1016/j.cellimm.2017.07.003

10. Weiss GE, Crompton PD, Li S, Walsh LA, Moir S, Traore B, et al. Atypical Memory B Cells Are Greatly Expanded in Individuals Living in a Malaria-Endemic Area. The Journal of Immunology. 2009;183: 2176–2182. doi:10.4049/jimmunol.0901297

11. Obeng-Adjei N, Portugal S, Holla P, Li S, Sohn H, Ambegaonkar A, et al. Malaria-induced interferon-gamma drives the expansion of Tbethi atypical memory B cells. PLoS Pathog. 2017;13: e1006576. doi:10.1371/journal.ppat.1006576

12. Sullivan RT, Kim CC, Fontana MF, Feeney ME, Jagannathan P, Boyle MJ, et al. FCRL5 Delineates Functionally Impaired Memory B Cells Associated with Plasmodium falciparum Exposure. PLoS Pathog. 2015;11: e1004894. doi:10.1371/journal.ppat.1004894

13. Portugal S, Tipton CM, Sohn H, Kone Y, Wang J, Li S, et al. Malaria-associated atypical memory B cells exhibit markedly reduced B cell receptor signaling and effector function. Elife. 2015;4. doi:10.7554/eLife.07218

14. Ambegaonkar AA, Kwak K, Sohn H, Manzella-Lapeira J, Brzostowski J, Pierce SK. Expression of inhibitory receptors by B cells in chronic human infectious diseases restricts responses to membrane-associated antigens. Sci Adv. 2020;6. doi:10.1126/SCIADV.ABA6493

15. Holla P, Ambegaonkar A, Sohn H, Pierce SK. Exhaustion may not be in the human B cell vocabulary, at least not in malaria. Immunol Rev. 2019 [cited 9 Oct 2019]. doi:10.1111/imr.12809

16. MacLean AJ, Richmond N, Koneva L, Attar M, Medina CAP, Thornton EE, et al. Secondary influenza challenge triggers resident memory B cell migration and rapid relocation to boost

antibody secretion at infected sites. Immunity. 2022;55: P718-733.E8.
doi:10.1016/J.IMMUNI.2022.03.003

17. Johnson JL, Rosenthal RL, Knox JJ, Myles A, Naradikian MS, Madej J, et al. The Transcription
Factor T-bet Resolves Memory B Cell Subsets with Distinct Tissue Distributions and Antibody
Specificities in Mice and Humans. Immunity. 2020;52: 842-855.e6.
doi:10.1016/j.immuni.2020.03.020

18. Hopp CS, Skinner J, Anzick SL, Tipton CM, Peterson ME, Li S, et al. Atypical B cells up-regulate
costimulatory molecules during malaria and secrete antibodies with T follicular helper cell
support. Sci Immunol. 2022;7: eabn1250. doi:10.1126/sciimmunol.abn1250

19. Rubtsov A v, Rubtsova K, Kappler JW, Jacobelli J, Friedman RS, Marrack P. CD11c-Expressing B
Cells Are Located at the T Cell/B Cell Border in Spleen and Are Potent APCs. Journal of
Immunology. 2015;195: 71–79. doi:10.4049/jimmunol.1500055

20. Cancro MP. Age-Associated B Cells. Annu Rev Immunol. 2020;38: 315–340.
doi:10.1146/annurev-immunol-092419-031130

21. Golinski M-L, Demeules M, Derambure C, Riou G, Maho-Vaillant M, Boyer O, et al. CD11c+ B
Cells Are Mainly Memory Cells, Precursors of Antibody Secreting Cells in Healthy Donors.
Front Immunol. 2020;11: 32. doi:10.3389/fimmu.2020.00032

22. Colonna-Romano G, Bulati M, Aquino A, Pellicanò M, Vitello S, Lio D, et al. A double-negative
(IgD-CD27-) B cell population is increased in the peripheral blood of elderly people. Mech
Ageing Dev. 2009;130: 681–690. doi:10.1016/J.MAD.2009.08.003

23. Courey-Ghaouzi AD, Kleberg L, Sundling C. Alternative B Cell Differentiation During Infection
and Inflammation. Front Immunol. 2022;13: 3192. doi:10.3389/FIMMU.2022.908034

24. Mouat IC, Goldberg E, Horwitz MS. Age-associated B cells in autoimmune diseases. Cellular
and Molecular Life Sciences. 2022;79: 1–11. doi:10.1007/S00018-022-04433-9

25. Holla P, Dizon B, Ambegaonkar AA, Rogel N, Goldschmidt E, Boddapati AK, et al. Shared
transcriptional profiles of atypical B cells suggest common drivers of expansion and function
in malaria, HIV, and autoimmunity. Sci Adv. 2021;7: eabg8384. doi:10.1126/sciadv.abg8384

26. Moir S, Ho J, Malaspina A, Wang W, DiPoto AC, O'Shea MA, et al. Evidence for HIV-associated
B cell exhaustion in a dysfunctional memory B cell compartment in HIV-infected viremic
individuals. J Exp Med. 2008;205: 1797–1805. doi:10.1084/jem.20072683

27. Sundling C, Rönnberg C, Yman V, Asghar M, Jahnmatz P, Lakshmikanth T, et al. B cell profiling
in malaria reveals expansion and remodeling of CD11c+ B cell subsets. JCI Insight. 2019;4:
e126492. doi:10.1172/jci.insight.126492

28. Sutton HJ, Aye R, Idris AH, Vistein R, Nduati E, Kai O, et al. Atypical B cells are part of an
alternative lineage of B cells that participates in responses to vaccination and infection in
humans. Cell Rep. 2021;34: 108684. doi:10.1016/j.celrep.2020.108684

29. Knox JJ, Buggert M, Kardava L, Seaton KE, Eller MA, Canaday DH, et al. T-bet+ B cells are
induced by human viral infections and dominate the HIV gp140 response. JCI Insight. 2017;2:
e92943. doi:10.1172/jci.insight.92943

30. Gjertsson I, McGrath S, Grimstad K, Jonsson CA, Camponeschi A, Thorarinsdottir K, et al. A
close-up on the expanding landscape of CD21–/low B cells in humans. Clin Exp Immunol.
2022; 1–13. doi:10.1093/CEI/UXAC103

31. Jenks SA, Cashman KS, Zumaquero E, Marigorta UM, Patel A v., Wang X, et al. Distinct
Effector B Cells Induced by Unregulated Toll-like Receptor 7 Contribute to Pathogenic
Responses in Systemic Lupus Erythematosus. Immunity. 2018;49: 725-739.e6.
doi:10.1016/j.immuni.2018.08.015

32. Ehrhardt GRA, Hsu JT, Gartland L, Leu C-M, Zhang S, Davis RS, et al. Expression of the
immunoregulatory molecule FcRH4 defines a distinctive tissue-based population of memory
B cells. J Exp Med. 2005;202: 783–791. doi:10.1084/jem.20050879

33.   Ambegaonkar AA, Nagata S, Pierce SK, Sohn H. The Differentiation in vitro of Human Tonsil B
Cells With the Phenotypic and Functional Characteristics of T-bet+ Atypical Memory B Cells in
Malaria. Front Immunol. 2019;10: 852. doi:10.3389/fimmu.2019.00852

34.   Muellenbeck MF, Ueberheide B, Amulic B, Epp A, Fenyo D, Busse CE, et al. Atypical and
classical memory B cells produce *Plasmodium falciparum* neutralizing antibodies. J Exp Med.
2013;210: 389–399. doi:10.1084/jem.20121970

35.   Stone SL, Peel JN, Scharer CD, Risley CA, Chisolm DA, Schultz MD, et al. T-bet Transcription
Factor Promotes Antibody-Secreting Cell Differentiation by Limiting the Inflammatory Effects
of IFN-γ on B Cells. Immunity. 2019;50: 1172-1187.e7. doi:10.1016/j.immuni.2019.04.004

36.   Dustin ML, Shaw AS. Costimulation: Building an immunological synapse. Science (1979).
1999;283: 649–650. doi:10.1126/SCIENCE.283.5402.649

37.   Damle NK, Klussman K, Linsley PS, Aruffo A. Differential costimulatory effects of adhesion
molecules B7, ICAM-1, LFA-3, and VCAM-1 on resting and antigen-primed CD4+ T
lymphocytes. The Journal of Immunology. 1992;148: 1985–1992.
doi:10.4049/JIMMUNOL.148.7.1985

38.   Aye R, Sutton HJ, Nduati EW, Kai O, Mwacharo J, Musyoki J, et al. Malaria exposure drives
both cognate and bystander human B cells to adopt an atypical phenotype. Eur J Immunol.
2020;50: 1187–1194. doi:10.1002/eji.201948473

39.   Steuten J, Bos A V., Kuijper LH, Claireaux M, Olijhoek W, Elias G, et al. Distinct dynamics of
antigen-specific induction and differentiation of different CD11c+ Tbet+ B cell subsets.
Journal of Allergy and Clinical Immunology. 2023 [cited 9 Mar 2023].
doi:10.1016/J.JACI.2023.02.020

40.   Dement-Brown J, Newton CS, Ise T, Damdinsuren B, Nagata S, Tolnay M. Fc receptor-like 5
promotes B cell proliferation and drives the development of cells displaying switched
isotypes. J Leukoc Biol. 2012;91: 59–67. doi:10.1189/jlb.0211096

41.   Wilson TJ, Fuchs A, Colonna M. Cutting Edge: Human FcRL4 and FcRL5 Are Receptors for IgA
and IgG. The Journal of Immunology. 2012;188: 4741–4745. doi:10.4049/jimmunol.1102651

42.   Franco A, Kraus Z, Li H, Seibert N, Dement-Brown J, Tolnay M. CD21 and FCRL5 form a
receptor complex with robust B-cell activating capacity. Int Immunol. 2018;30: 569–578.
doi:10.1093/intimm/dxy052

43.   Schittenhelm L, Hilkens CM, Morrison VL. β2 integrins as regulators of dendritic cell,
monocyte, and macrophage function. Front Immunol. 2017;8: 1866.
doi:10.3389/FIMMU.2017.01866/BIBTEX

44.   Coban C, Ishii KJ, Kawai T, Hemmi H, Sato S, Uematsu S, et al. Toll-like receptor 9 mediates
innate immune activation by the malaria pigment hemozoin. Journal of Experimental
Medicine. 2005;201: 19–25. doi:10.1084/JEM.20041836

45.   Parroche P, Lauw FN, Goutagny N, Latz E, Monks BG, Visintin A, et al. Malaria hemozoin is
immunologically inert but radically enhances innate responses by presenting malaria DNA to
Toll-like receptor 9. Proc Natl Acad Sci U S A. 2007;104: 1919–1924.
doi:10.1073/PNAS.0608745104

46.   Bernasconi NL, Onai N, Lanzavecchia A. A role for Toll-like receptors in acquired immunity:
up-regulation of TLR9 by BCR triggering in naive B cells and constitutive expression in
memory B cells. Blood. 2003;101: 4500–4504. doi:10.1182/BLOOD-2002-11-3569

47.   Pohl K, Cockburn IA. Innate immunity to malaria: The good, the bad and the unknown. Front
Immunol. 2022;13: 4417. doi:10.3389/FIMMU.2022.914598/BIBTEX

48.   Gowda NM, Wu X, Gowda DC. TLR9 and MyD88 Are Crucial for the Development of
Protective Immunity to Malaria. The Journal of Immunology. 2012;188: 5073–5085.
doi:10.4049/JIMMUNOL.1102143

49.   Omar AH, Yasunami M, Yamazaki A, Shibata H, Ofori MF, Akanmori BD, et al. Toll-like
receptor 9 (TLR9) polymorphism associated with symptomatic malaria: A cohort study. Malar
J. 2012;11: 1–10. doi:10.1186/1475-2875-11-168/FIGURES/4

50. Kelsoe G, Zheng B. Sites of B-cell activation in vivo. Curr Opin Immunol. 1993;5: 418–422.
doi:10.1016/0952-7915(93)90062-W

51. Cyster JG, Allen CDC. B Cell Responses: Cell Interaction Dynamics and Decisions. Cell. Cell
Press; 2019. pp. 524–540. doi:10.1016/j.cell.2019.03.016

52. Okada T, Miller MJ, Parker I, Krummel MF, Neighbors M, Hartley SB, et al. Antigen-engaged B
cells undergo chemotaxis toward the T zone and form motile conjugates with helper T cells.
PLoS Biol. 2005;3: 1047–1061. doi:10.1371/journal.pbio.0030150

53. Heufler C, Koch F, Stanzl U, Topar G, Wysocka M, Trinchieri G, et al. Interleukin-12 is
produced by dendritic cells and mediates T helper 1 development as well as interferon-γ
production by T helper 1 cells. Eur J Immunol. 1996;26: 659–668.
doi:10.1002/EJI.1830260323

54. Yu S, Jia L, Zhang Y, Zhong J, Yang B, Wu C. IL-12 induced the generation of IL-21- and IFN-γ-
co-expressing poly-functional CD4+ T cells from human naive CD4+ T cells.
https://doi.org/101080/1538410120151093703. 2015;14: 3362–3372.
doi:10.1080/15384101.2015.1093703

55. Obeng-Adjei N, Portugal S, Tran TM, Yazew TB, Skinner J, Li S, et al. Circulating Th1-Cell-type
Tfh Cells that Exhibit Impaired B Cell Help Are Preferentially Activated during Acute Malaria in
Children. Cell Rep. 2015;13: 425–439. doi:10.1016/j.celrep.2015.09.004

56. Oyong DA, Loughland JR, Soon MSF, Chan JA, Andrew D, Wines BD, et al. Adults with
Plasmodium falciparum malaria have higher magnitude and quality of circulating T-follicular
helper cells compared to children. EBioMedicine. 2022;75.
doi:10.1016/J.EBIOM.2021.103784/ATTACHMENT/7123BDD3-C980-4D0F-9BB1-
4BE90A2ACDC9/MMC1.DOCX

57. Keller B, Strohmeier V, Harder I, Unger S, Payne KJ, Andrieux G, et al. The expansion of human
T-bethighCD21low B cells is T cell dependent. Sci Immunol. 2021;6: eabh0891.
doi:10.1126/sciimmunol.abh0891

58. Rubtsova K, Rubtsov A v., van Dyk LF, Kappler JW, Marrack P. T-box transcription factor T-bet,
a key player in a unique type of B-cell activation essential for effective viral clearance. Proc
Natl Acad Sci U S A. 2013;110: E3216–E3224.
doi:10.1073/PNAS.1312348110/SUPPL_FILE/PNAS.201312348SI.PDF

59. Naradikian MS, Myles A, Beiting DP, Roberts KJ, Dawson L, Herati RS, et al. Cutting Edge: IL-4,
IL-21, and IFN-γ Interact To Govern T-bet and CD11c Expression in TLR-Activated B Cells. The
Journal of Immunology. 2016;197: 1023–1028. doi:10.4049/jimmunol.1600522

60. Levack RC, Newell KL, Popescu M, Cabrera-Martinez B, Winslow GM. CD11c + T-bet + B Cells
Require IL-21 and IFN-γ from Type 1 T Follicular Helper Cells and Intrinsic Bcl-6 Expression but
Develop Normally in the Absence of T-bet. The Journal of Immunology. 2020;205: 1050–
1058. doi:10.4049/jimmunol.2000206

61. Du SW, Arkatkar T, Jacobs HM, Rawlings DJ, Jackson SW. Generation of functional murine
CD11c+ age-associated B cells in the absence of B cell T-bet expression. Eur J Immunol.
2019;49: 170–178. doi:10.1002/eji.201847641

62. Gao X, Shen Q, Roco JA, Frith K, Munier CML, Nekrasov M, et al. ZEB2 regulates the
development of CD11c+ atypical B cells. bioRxiv. 2022; 2022.09.01.506173.
doi:10.1101/2022.09.01.506173

63. Gu S, Han X, Yao C, Ding H, Hou G, Qu B, et al. The transcription factor Zeb2 drives
differentiation of age-associated B cells. bioRxiv. 2021; 2021.07.24.453633.
doi:10.1101/2021.07.24.453633

64. Tangye SG, Ma CS. Regulation of the germinal center and humoral immunity by interleukin-
21. Journal of Experimental Medicine. 2020;217. doi:10.1084/JEM.20191638/132621

65. Kerfoot SM, Yaari G, Patel JR, Johnson KL, Gonzalez DG, Kleinstein SH, et al. Germinal Center
B Cell and T Follicular Helper Cell Development Initiates in the Interfollicular Zone. Immunity.
2011;34: 947–960. doi:10.1016/j.immuni.2011.03.024

66.　Wang S, Wang J, Kumar V, Karnell JL, Naiman B, Gross PS, et al. IL-21 drives expansion and
plasma cell differentiation of autoreactive CD11chiT-bet+B cells in SLE. Nat Commun. 2018;9:
1758. doi:10.1038/s41467-018-03750-7
67.　Rivera-Correa J, Guthmiller JJ, Vijay R, Fernandez-Arias C, Pardo-Ruge MA, Gonzalez S, et al.
Plasmodium DNA-mediated TLR9 activation of T-bet+B cells contributes to autoimmune
anaemia during malaria. Nat Commun. 2017;8: 1282. doi:10.1038/s41467-017-01476-6
68.　Rivera-Correa J, Mackroth MS, Jacobs T, zur Wiesch JS, Rolling T, Rodriguez A. Atypical
memory b-cells are associated with plasmodium falciparum anemia through anti-
phosphatidylserine antibodies. Elife. 2019;8: e48309. doi:10.7554/ELIFE.48309
69.　Dooley NL, Chabikwa TG, Pava Z, Loughland JR, Hamelink J, Berry K, et al. Malaria drives
unique regulatory responses across multiple immune cell subsets. bioRxiv. 2022;
2022.11.16.516822. doi:10.1101/2022.11.16.516822
70.　Zander RA, Guthmiller JJ, Graham AC, Pope RL, Burke BE, Carr DJJ, et al. Type I Interferons
Induce T Regulatory 1 Responses and Restrict Humoral Immunity during Experimental
Malaria. Stevenson MM, editor. PLoS Pathog. 2016;12: e1005945.
doi:10.1371/journal.ppat.1005945
71.　Lautenbach MJ, Yman V, Silva CS, Kadri N, Broumou I, Chan S, et al. Systems analysis shows a
role of cytophilic antibodies in shaping innate tolerance to malaria. Cell Rep. 2022;39:
110709. doi:10.1016/J.CELREP.2022.110709
72.　Guha R, Mathioudaki A, Doumbo S, Doumtabe D, Skinner J, Arora G, et al. Plasmodium
falciparum malaria drives epigenetic reprogramming of human monocytes toward a
regulatory phenotype. Kazura JW, editor. PLoS Pathog. 2021;17: e1009430.
doi:10.1371/journal.ppat.1009430
73.　Nahrendorf W, Ivens A, Spence PJ. Inducible mechanisms of disease tolerance provide an
alternative strategy of acquired immunity to malaria. Elife. 2021;10. doi:10.7554/ELIFE.63838
74.　Peyron F, Burdin N, Ringwald P, Vuillez JP, Rousset F, Banchereau J. High levels of circulating
IL-10 in human malaria. Clin Exp Immunol. 1994;95: 300–3. doi:10.1111/j.1365-
2249.1994.tb06527.x
75.　Kumar R, Ng S, Engwerda C. The Role of IL-10 in Malaria: A Double Edged Sword. Front
Immunol. 2019;10: 229. doi:10.3389/fimmu.2019.00229
76.　Guthmiller JJ, Graham AC, Zander RA, Pope RL, Butler NS. Cutting Edge: IL-10 Is Essential for
the Generation of Germinal Center B Cell Responses and Anti-Plasmodium Humoral
Immunity. The Journal of Immunology. 2017;198: 617–622. doi:10.4049/JIMMUNOL.1601762
77.　Han X, Yang J, Zhang Y, Zhang Y, Cao H, Cao Y, et al. Potential role for regulatory B cells as a
major source of interleukin-10 in spleen from Plasmodium chabaudi-infected mice. Infect
Immun. 2018;86. doi:10.1128/IAI.00016-18
78.　Freitas do Rosário AP, Lamb T, Spence P, Stephens R, Lang A, Roers A, et al. IL-27 Promotes IL-
10 Production by Effector Th1 CD4+ T Cells: A Critical Mechanism for Protection from Severe
Immunopathology during Malaria Infection. The Journal of Immunology. 2012;188: 1178–
1190. doi:10.4049/JIMMUNOL.1102755
79.　Barr TA, Brown S, Ryan G, Zhao J, Gray D. TLR-mediated stimulation of APC: Distinct cytokine
responses of B cells and dendritic cells. Eur J Immunol. 2007;37: 3040–3053.
doi:10.1002/EJI.200636483
80.　Mauri C, Bosma A. Immune Regulatory Function of B Cells. https://doi.org/101146/annurev-
immunol-020711-074934. 2012;30: 221–241. doi:10.1146/ANNUREV-IMMUNOL-020711-
074934
81.　Zhang G, Manaca MN, McNamara-Smith M, Mayor A, Nhabomba A, Berthoud TK, et al.
Interleukin-10 (IL-10) polymorphisms are associated with IL-10: Production and clinical
malaria in young children. Infect Immun. 2012;80: 2316–2322. doi:10.1128/IAI.00261-12

82.   Hirano T, Yasukawa K, Harada H, Taga T, Watanabe Y, Matsuda T, et al. Complementary DNA
944   for a novel human interleukin (BSF-2) that induces B lymphocytes to produce
945   immunoglobulin. Nature 1986 324:6092. 1986;324: 73–76. doi:10.1038/324073a0
946   83.   Dienz O, Rincon M. The effects of IL-6 on CD4 T cell responses. Clinical Immunology.
947   2009;130: 27–33. doi:10.1016/J.CLIM.2008.08.018
948   84.   Brunson JC. ggalluvial: Layered Grammar for Alluvial Plots. J Open Source Softw. 2020;5:
949   2017. doi:10.21105/JOSS.02017

951