Further evidences of an emerging stingless bee-yeast symbiosis

Gabriela Toninato de Paula[1], Weilan Gomes da Paixão Melo[1,2], Ivan de Castro[3], Cristiano

Menezes[4], Camila Raquel Paludo[5], Carlos Augusto Rosa[6], Mônica Tallarico Pupo[1*]

[1]School of Pharmaceutical Sciences of Ribeirão Preto, University of São Paulo, Ribeirão

Preto, SP, Brazil.

[2]State University of the Tocantina Region of Maranhão, Estreito, MA, Brazil.

[3]School of Medical Sciences of Ribeirão Preto, University of São Paulo, Ribeirão Preto,

SP, Brazil.

[4]Brazilian Agricultural Research Corporation, Embrapa Meio Ambiente, Jaguariúna, SP,

Brazil.

[5]Federal University of Mato Grosso, Barra do Garças, MT, Brazil.

[6]Departamento de Microbiologia, ICB, Universidade Federal de Minas Gerais, Belo

Horizonte, MG, Brazil.

* Correspondence:

Mônica Tallarico Pupo

mtpupo@fcfrp.usp.br

Abstract

Symbiotic interactions between microorganisms and social insects have been described as crucial for the maintenance of these multitrophic systems, as observed for the stingless bee *Scaptotrigona depilis* and the yeast *Zygosaccharomyces* sp. The larvae of *S. depilis* ingest fungal filaments of *Zygosaccharomyces* sp. to obtain ergosterol, which is the precursor for the biosynthesis of ecdysteroids that modulate insect metamorphosis. In this work we verified that nutritional fungal symbioses also occur in other species of stingless bees. We analyzed brood cell samples from 19 species of stingless bees collected in Brazil. The osmophilic yeast *Zygosaccharomyces* spp. was isolated from eight bee species, namely *Scaptotrigona bipuctata*, *S. postica*, *S. tubiba*, *Tetragona clavipes*, *Melipona quadrifasciata*, *M. fasciculata*, *M. bicolor* and *Partamona helleri*. These yeasts form pseudohyphae and also accumulate ergosterol in lipid droplets, similar to the pattern observed for *S. depilis*. The phylogenetic analyses including various *Zygosaccharomyces* revealed that strains isolated from the brood cells formed a branch separated from the previously described *Zygosaccharomyces* species, suggesting that they are new species of this genus and reinforcing the symbiotic interaction with the host insects.

Importance

Benefits exchanged in insect–fungus mutualisms include nutrition, protection, and dispersal. Fungal nutritional roles are well described for some eusocial insects, such as fungus growing ants and termites, but similar interaction in stingless bees was so far observed just in *Scaptotrigona depilis*. Here we expand the knowledge of yeast-bee

symbiosis by analyzing the presence, cell morphologies, lipid accumulation and

phylogenetic relationships of fungi isolated from brood cells and other locations of bee

colonies. *Zygosaccharomyces* isolates were recovered from 42% of the bee species

assessed, and probably represent new species showing pseudohyphae formation and lipid

accumulation similar to *S. depilis* associated *Zygosaccharomyces* strains. The phylogenetic

analyses suggested an evolutionary adaptation of *Zygosaccharomyces* spp. to the brood cell

environment to provide nutritional benefits for the developing insect. Stingless bees play

important ecosystem services, and our results raise the concern that fungicidal agents used

in agriculture could disrupt this symbiosis, impacting bee health.

Keywords: stingless bees, fungi, yeast, *Zygosaccharomyces*, symbiosis, brood cell

Eusocial insects establish several symbiotic interactions with microorganisms, ranging from

obligate mutualisms to specialized parasitism (1, 2). The nutritional benefits provided by

fungal mutualists is well described for fungus-growing ants, native to the Neotropics, and

termites from Africa and Asia (3). Ants of the subtribe Attina cultivate fungi of the families

Agaricaceae and Pterulaceae for food (4). Similarly, termites of the subfamily

Macrotermitinae cultivate fungi of the genus *Termitomyces*, which are not only the main

food source but also provide digestive services (5). Other microbial symbionts also play

different roles in such multitrophic interactions, like production of chemical defenses

against entomopathogens (1, 2).

Stingless bees (SBs) (Apidae: Apinae: Meliponini) are a monophyletic group of eusocial

insects that belong to a larger group of corbiculate bees (6). Meliponini have pantropical

distribution and interact with various microorganisms such as bacteria, yeasts, filamentous

fungi, and viruses. Indeed, microbial fermentation contributes to important

physicochemical characteristics and to the preservation of pollen and honey (7).

Interestingly, the benefits provided by a yeast of the genus *Zygosaccharomyces* for the host

Scaptotrigona depilis has been reported as the first example of nutritional symbiosis in SBs.

This osmophilic yeast grows inside the brood cells of *S. depilis* and is eaten by the larvae,

an essential process for larval metamorphosis. The fungus accumulates ergosterol, which is

used by the developing insect as a precursor for ecdysteroid biosynthesis leading to the

proper pupation (8, 9). However, this remarkable interaction had been observed just for one

bee species so far, lacking generality among Meliponini (10).

This unprecedented finding of a yeast important for the larval development raised the

hypothesis that similar symbiosis might be more spread among SBs. Here, we describe the

occurrence of *Zygosaccharomyces* isolates in the brood cells of different SBs species, as

well as the phylogenetic relationships and morphological characteristics of these yeasts in

comparison with other species isolated from other locations in the colonies of SBs.

Different nest sites of 19 species of SBs, distributed among 12 genera, were assessed for

isolation of fungi, resulting in 32 fungal isolates. Twenty-seven were obtained from brood

cells, while four were isolated from honey and one from cerumen. All isolates had the 26S,

18S rRNA gene and ITS regions sequenced, resulting in 23 isolates belonging to the genus

Zygosaccharomyces, seven were identified as *Monascus* spp., one as *Xerochrysium* sp. and

one as *Leiothecium* sp. (Supplementary Material - Table S1). Fungal filaments were

observed and isolated from the brood cells of eight species of SBs distributed in four genera and eight species (*Scaptotrigona bipuctata*, *S. postica*, *S. tubiba*, *Tetragona clavipes*, *Melipona quadrifasciata*, *M. fasciculata*, *M. bicolor*, and *Partamona helleri*). All collected fungal filaments were identified as belonging to the genus *Zygosaccharomyces*, representing 42% incidence rate of *Zygosaccharomyces* sp. in brood cells of sampled SBs species. *Zygosaccharomyces* were also isolated from honey of four species and from cerumen of one species (Table S1), while fungal filaments were not observed in the brood cells of 10 SBs species. Samples with low amounts of brood cells and the different larval stages could be limiting factors for the observation of *Zygosaccharomyces*, in addition to the difficulty of culturing this microorganism in less complex culture media. Fungal filaments were observed in SBs with disk-shaped brood cells structure (Figure S1A and S1C) and a larger diameter, but were not observed in other species with smaller diameters (Figure S1B, S1D and S1E) and cluster-shaped brood cells (Figure S1D and S1E). The size and shape of the brood cells may influence the visibility of the symbiont fungus.

Cell morphologies of *Zygosaccharomyces* spp. varied according to the site of isolation. Isolates from brood cells form pseudohyphae in glucose-rich culture medium (30 G), while the others have spherical and ovoid cells under the same growth conditions (Figure S2). Pseudohyphae formation can be triggered by low nitrogen levels and is a form of foraging (11). The location of these *Zygosaccharomyces* strains may be a determining factor in the expression of pseudohyphae. The brood cell is composed of cerumen and larval food, and their composition may vary in nature. Cerumen consists only of waxes and resins (12), while larval food consists of water (40-60%), sugars (5-12%), and free amino acids (0.2-1.3%) (13). These water-soluble components make up to 70% of the larval food

composition. In addition, pollen and lipids are also present (13). This specific environment

with high sugar concentration and low nitrogen content may require some adaptation of the

microorganisms, such as the formation of pseudohyphae. It is also hypothesized that

pseudohyphae are more easily ingested by the larvae since they are present on the surface

of the larval food and have long filaments. This particular morphology of these yeasts

(Figure 1) could favor the availability and uptake of nutrients for the larval development.

Zygosaccharomyces strains from brood cells of different bees showed accumulation of

intracellular lipids, as visualized by fluorescence microscopy (Figure 1D). GC-MS analyses

of cell extracts of *Zygosaccharomyces* spp. confirmed the presence of ergosterol in all the

samples (Figure S3), as previously described for *Zygosaccharomyces* sp. isolated from *S.*

depilis (9), reinforcing the nutritional function of the yeasts for SBs larvae.

Zygosaccharomyces strains from the brood cells had the lowest query coverage and identity

percentages with species from the Genbank database and may represent new species.

Phylogenetic tree from Bayesian analyses using the sequences of the D1/D2 domain of the

26S gene was performed on *Zygosaccharomyces* spp. (Figure 2). A combined phylogeny

was also constructed using a dataset of the 26S and 18S genes (Figure S4). Phylogenies

show a branch consisting of *Zygosaccharomyces* spp. isolated exclusively from the brood

cells of different species of SBs. The other lineages of *Zygosaccharomyces* isolated from

different nest sites are phylogenetically distant. The branch consists only of

Zygosaccharomyces spp. strains isolated from the brood cells contain *Zygosaccharomyces*

sp. associated with *S. depilis* (9), and is divided into several groups. One group, which is

quite clear and distinct from the others, consists of strains of *Zygosaccharomyces* spp.,

isolated from different species of the genus *Scaptotrigona*, suggesting that these strains

belong to the same yeast species previously described for *S. depilis*. A second group

consists of *Zygosaccharomyces* strains derived from the brood cells of several bee species

of the genus *Melipona*, as well as the species *T. clavipes* and *P. helleri*. In contrast to

Scaptotrigona isolates, *Zygosaccharomyces* spp. isolated from *Melipona* species probably

belong to different species. In turn, *Zygosaccharomyces* isolates from different colonies of

the species *T. clavipes* appear to belong to the same species, since they are closely related.

Altogether, our findings suggest a widespread symbiotic relationship between species of

Zygosaccharomyces and SBs, and expand the current knowledge about the classical and

non-classical mutualisms between fungi and insects (14). Meliponini are important

pollinators of native plants and agricultural crops, playing important ecosystem services.

Declines in these pollinators are caused by several factors, including deforestation and

pesticides used in agriculture. Pesticides can directly affect bees (15), and our results raise

the hypothesis that fungicides used in agriculture could disrupt the bee-yeast symbiosis,

indirectly affecting SBs health. Further studies are needed to evaluate the toxicity of these

compounds in this symbiotic system to help define public policies for rational use of

pesticides in agricultural crops.

Data availability. The complete data set is presented in the text and the supplemental

material.

Acknowledgments

This research was supported by São Paulo Research Foundation (FAPESP) grants

#2013/50954-0 (MTP), #2013/07600-3 (MTP; CEPID-CIBFar), #2018/03650-0 (GTP),

#2015/01001-6 (WGPM), by the Conselho Nacional de Pesquisa e Desenvolvimento

Tecnológico (CNPq) grants #303792/2018-2 and #307893/2022-7 (MTP); #408733/2021-7

and #406564/2022-1 (CAR), and by Fundação de Amparo à Pesquisa do Estado de Minas

Gerais (FAPEMIG) grant APQ-01525-14 (CAR). This study was financed in part by the

Coordenação de Aperfeiçoamento de Pessoal de Nível Superior - Brasil (CAPES) - Finance

Code 001. Claudia C. de Macedo and Izabel Cristina C. Turatti are acknowledged by their

technical support. We also acknowledge the facilities of the Laboratório Multiusuário de

Microscopia Confocal – LMMC, FAPESP #2004/08868-0.

164　Author Contributions

Conceptualization – MTP, GTP, WGPM, CM, CRP, CAR; Data curation – MTP; Formal

analysis – MTP, GTP, WGPM; Funding acquisition – MTP, CAR; Investigation – GTP,

WGPM; Methodology – GTP, WGPM, IC, CM; Project administration – MTP; Resources

– MTP; Supervision - MTP; Validation – GTP, WGPM; Visualization – MTP, WGPM;

Writing - original draft – MTP, GTP, WGPM; Writing - review & editing – MTP, GTP,

WGPM, CM, CRP, CAR.

We declare no conflict of interest.

References

1.   van Arnam EB, Currie CR, Clardy J. 2018. Defense contracts: molecular protection in insect-microbe symbioses. Chem Soc Rev 47:1638–1651.

2.   Menegatti C, Fukuda TTH, Pupo MT. 2021. Chemical ecology in insect-microbe interactions in the Neotropics. Planta Med 87:38–48.

3.   Mueller UG, Schultz TR, Currie CR, Adams RMM, Malloch D. 2001. The origin of the attine ant-fungus mutualism. Q Rev Biol 76:169–198.

4.   Bizarria R, Pagnocca FC, Rodrigues A. 2022. Yeasts in the attine ant–fungus mutualism: Diversity, functional roles, and putative biotechnological applications. Yeast 39:25–39.

5.   de Fine Licht HH, Andersen A, Aanen DK. 2005. *Termitomyces* sp. associated with the termite *Macrotermes natalensis* has a heterothallic mating system and multinucleate cells. Mycol Res 109:314–318.

6.   Romiguier J, Cameron SA, Woodard SH, Fischman BJ, Keller L, Praz CJ. 2016. Phylogenomics controlling for base compositional bias reveals a single origin of eusociality in corbiculate bees. Mol Biol Evol 33:670–678.

7.   de Paula GT, Menezes C, Pupo MT, Rosa CA. 2021. Stingless bees and microbial interactions. Curr Opin Insect Sci 44:41–47.

8.   Menezes C, Vollet-Neto A, Marsaioli AJ, Zampieri D, Fontoura IC, Luchessi AD, Imperatriz-Fonseca VL. 2015. A Brazilian social bee must cultivate fungus to survive. Curr Biol 25:2851–2855.

9.   Paludo CR, Menezes C, Silva-Junior EA, Vollet-Neto A, Andrade-Dominguez A,

Pishchany G, Khadempour L, do Nascimento FS, Currie CR, Kolter R, Clardy J, Pupo MT.

2018. Stingless bee larvae require fungal steroid to pupate. Sci Rep 8(1):1122.

10.   Roubik DW. 2023. Stingless bee (Apidae: Apinae: Meliponini) ecology. Annu Rev

Entomol 68:231–256.

11.   Cullen PJ, Sprague GF. 2012. The regulation of filamentous growth in yeast.

Genetics 190(1):23–49.

12.   Roubik DW. 2006. Stingless bee nesting biology. Apidologie 37:124–143.

13.   Hartfelder K, Engels W. 1989. The composition of larval food in stingless bees:

evaluating nutritional balance by chemosystematic methods. Insect Soc 36:1–14.

14.   Biedermann PHW, Vega FE. 2020. Ecology and evolution of insect–fungus

mutualisms. Annu Rev Entomol 65:431–455.

15.   Cham KO, Nocelli RCF, Borges LO, Viana-Silva FEC, Tonelli CAM, Malaspina O,

Menezes C, Rosa-Fontana AS, Blochtein B, Freitas BM, Pires CSS, Oliveira FF, Contrera

FAL, Torezani KRS, Ribeiro MDF, Siqueira MAL, Rocha MCLSA. 2019. Pesticide

exposure assessment paradigm for stingless bees. Environ Entomol 48:36–48.

Figure 1. Images of *Zygosaccharomyces* spp. isolates from stingless bees *S. bipunctata*, *S. tubiba*, *M. bicolor*, *T. clavipes* and *P. helleri*. (A) Fungal filaments present in the brood cells. (B) Microbial growth in Petri dish (14 days) (C) Optical microscopy images (100x). (D) Fluorescence microscopy images. Stingless bees' photography: credit Cristiano Menezes.

Figure 2. Phylogenetic tree from Bayesian analysis based on sequences of the D1/D2 domain of the 26S rRNA gene of *Zygosaccharomyces* spp. strains isolated from various stingless bee species (highlighted in bold) and from previously described *Zygosaccharomyces* species retrieved from GenBank. Strains isolated from brood cells are highlighted in color, grouped by stingless bee genera. Numbers on branches indicate PP/ML bootstrap values of support for each clade. A total of 540 aligned positions were analyzed. The scale bar represents 0.01 substitutions per nucleotide position. BC: Brood Cell.

(A)   **(B)**   **(C)**   **(D)**

S. bipunctata

S. tubiba

M. bicolor

T. clavipes

P. helleri