DEK oncoprotein participates in heterochromatin replication via SUMO-dependent nuclear bodies

Agnieszka Pierzynska-Mach[1, #], Christina Czada[2, #], Christopher Vogel[2], Eva Gwosch[2], Xenia Osswald[2], Denis Bartoschek[2], Alberto Diaspro[1,3], Ferdinand Kappes[4, #, *] and Elisa Ferrando-May[2,5, #, *]

[1] Nanoscopy & NIC@IIT, Istituto Italiano di Tecnologia, Genoa, Italy
[2] Department of Biology, Bioimaging Center, University of Konstanz, Konstanz, Germany
[3] DIFILAB, Department of Physics, University of Genoa, Genoa, Italy
[4] Duke Kunshan University, Division of Natural and Applied Sciences, Kunshan, P.R. China
[5] German Cancer Research Center, Heidelberg, Germany

* Correspondence should be sent to: elisa.may@uni-konstanz.de

ferdinand.kappes@dukekunshan.edu.cn

[#] Equal contribution

Running title: SUMO-dependent formation of DEK bodies

Summary statement: The human oncoprotein DEK forms nuclear bodies that are regulated by the SUMO pathway and localize to active sites of constitutive heterochromatin replication.

Key words: oncogene, breast cancer, replication stress, histone modification, superresolution microscopy, siRNA screen.

Abstract

The correct inheritance of chromatin structure is key for maintaining genome function and preventing cellular transformation. DEK, a conserved chromatin protein, has recognized tumor-promoting properties, its overexpression being associated with poor prognosis in various cancer types. At the cellular level, DEK displays pleiotropic functions, influencing differentiation, apoptosis, and stemness, but a characteristic oncogenic mechanism remains elusive. Here, we report the identification of DEK bodies, focal assemblies of DEK occurring at specific, yet unidentified sites of heterochromatin replication. In these bodies, DEK localizes in direct proximity to active replisomes suggesting a function in the early maturation of heterochromatin. A high-throughput siRNA screen identifies SUMO as a major regulator of DEK body formation, linking DEK to the SUMO network that controls chromatin states and cell fate. This work combines and refines our previous data on DEK as a factor essential for heterochromatin integrity and facilitating replication under stress and delineates an avenue of further study for unraveling DEK´s contribution to cancer development.

Introduction

Replication of genomes is a fundamental process requiring meticulously orchestrated, yet adaptable mechanisms in space and time. The replication machinery encounters a range of different chromatin structures, some of which are prone to delaying, impairing, or even stalling replication fork progression - a state referred to as DNA replication stress (Maya-Mendoza et al., 2018) and often encountered in cancer cells (Boyer et al., 2016; Saxena and Zou, 2022). Overcoming such obstacles requires the coordinated action of a number of protective chromatin factors, often governed by targeted post-translational modifications, with poly(ADP)ribosylation and SUMOylation playing prominent roles (Abbas, 2021; Stewart-Morgan et al., 2020). Active DNA replication forks, as visualized by incorporation of DNA precursors (BrdU, EdU or other thymidine analogues) or *via* marking DNA replication factors (e.g. Proliferating Cell Nuclear Antigen, PCNA) (Chagin et al., 2016; Jackson and Pombo, 1998) appear as a characteristic pattern of nuclear foci that changes over time and typically reflects the chromatin structure being replicated (Aladjem, 2007; Casas-Delucchi et al., 2012; Heinz et al., 2018; O'Keefe et al., 1992; Sehwaiger et al., 2009). In general, euchromatin undergoes DNA replication during early S-phase, facultative heterochromatin during middle S-phase, and constitutive heterochromatin during late S-phase (Chadwick and Willard, 2004; Fragkos et al., 2015; Ryba et al., 2010; Su et al., 2020). However, some "hard-to-replicate" DNA structures, e.g., inactivated X chromosome, telomeres, centromeres, or certain DNA secondary structures, such as G-quadruplexes, four-way junction DNA and others, pose a particular challenge to the DNA replication machinery, and, in turn, require fine-tuned action of additional accessory factors (Mirkin and Mirkin, 2007).

We and others have identified the DEK oncogene, a unique and multifunctional non-histone chromosomal protein, as a supportive factor of DNA replication, particularly in scenarios where the replication machinery is under stress (Deutzmann et al., 2015; Ganz et al., 2019). Although Isolation of Proteins on Nascent DNA (iPOND) studies support no direct association of DEK with the replisome, DEK is enriched in maturing chromatin (Alabert et al., 2014; Aranda et al., 2014; Garcia et al., 2017; Lossaint et al., 2013; Ribeyre et al., 2016; Sirbu et al., 2013), in line with its post-translational modification-dependent

function as histone chaperone and its role in the maintenance of chromatin structure (Alexiadis et al., 2000; Böhm et al., 2005; Cavellán et al., 2006; Gamble and Fisher, 2007; Guo et al., 2021a; Hollenbach et al., 2002; Kappes et al., 2001; Kappes et al., 2004a; Kappes et al., 2004b; Kappes et al., 2011a; Privette Vinnedge et al., 2013; Saha et al., 2013; Sawatsubashi et al., 2010; Tabbert et al., 2006; Waidmann et al., 2014; Waldmann et al., 2002; Waldmann et al., 2003). Under moderate to severe replication stress, DEK is highly enriched at maturing chromatin, suggesting a protective role at the replication fork. Our previous research demonstrated that down-regulation of DEK in cells under moderate short-term replication stress *via* aphidicolin and camptothecin treatment results in the aggravation of replication stress phenotypes. Replication fork progression, proliferative capability and clonogenic survival were impaired and cells accumulated replication-born DNA damage and transmitted it to daughter cells (Deutzmann et al., 2015). Such fork-protective properties of DEK are modulated by PARP1/2 activity depending on the type and extent of replication stress (Ganz et al., 2019). At the same time, DEK appears to safeguard DNA replication also under steady-state, unchallenged conditions, as DNA fiber assays showed a mild decrease in fork speed when DEK expression is downregulated in untreated cells (Deutzmann et al., 2015). Collectively, these findings suggest that DEK is not a direct component of the replisome but binds in the vicinity of the fork to facilitate fork progression by supporting chromatin maturation. Evidence for a role of DEK in DNA replication via its properties as a chromatin architectural protein comes also from *in vitro* studies: experiments with SV40 mini-chromosomes showed that DEK introduces positive supercoils to chromatin and reduces the replication efficiency of chromatin in a reconstituted system (Alexiadis et al., 2000; Waldmann et al., 2002). Furthermore, DEK preferentially associates with four-way junctions over duplex DNA and with supercoiled over linear DNA, both structures that can induce replication stress *per se* or can be consequences of replication stress (Böhm et al., 2005; Waldmann et al., 2003).

Under steady-state conditions DEK associates with chromatin throughout the cell cycle (Kappes et al., 2001; Matrka et al., 2015), typically observed as pan-nuclear punctate pattern in immunofluorescence images and further supported by the requirement for

medium to high salt concentrations for its extraction from nuclei preparations (Kappes et al., 2001; Kappes et al., 2004a; Kappes et al., 2008; Özçelik et al., 2022; Saha et al., 2013), all in line with its pleiotropic involvement in multiple nuclear processes, ranging from transcriptional regulation, DNA damage response and repair, apoptosis, cell stemness and chromatin maintenance (Privette Vinnedge et al., 2013; Waldmann et al., 2004). More detailed analyses revealed that DEK can associate both with euchromatin and heterochromatin (Capitano et al., 2022; Gamble and Fisher, 2007; Guo et al., 2021a; Hollenbach et al., 2002; Hu et al., 2005; Ivanauskiene et al., 2014; Kappes et al., 2001; Kappes et al., 2004a; Kappes et al., 2004b; Kappes et al., 2008; Kappes et al., 2011b; Karam et al., 2014; Privette Vinnedge et al., 2013; Saha et al., 2013; Sandén and Gullberg, 2015; Sandén et al., 2014; Sawatsubashi et al., 2010; Waidmann et al., 2014; Zhou et al., 2022). However, qualitative and quantitative differences in the specific chromatin distribution of DEK amongst cell types and conditions are evident. Whereas DEK was identified as a suppressor of variegation (Su(var)), thus as a factor important for maintenance of heterochromatin (Capitano et al., 2019; Kappes et al., 2011a; Saha et al., 2013), other studies show preferential association with euchromatin (Hu et al., 2007; Ivanauskiene et al., 2014; Sandén et al., 2014; Yang et al., 2022), pinpointing to cell-type-specific post-translational modifications on DEK as critical for its specific chromatin association. Indeed, DEK is targeted by a plethora of post-translational modifications, with biological effects for acetylation, phosphorylation and poly(ADP)ribosylation having been reported (Cleary et al., 2005; Fahrer et al., 2010; Kappes et al., 2004a; Kappes et al., 2004b; Kappes et al., 2008; Mor-Vaknin et al., 2011; Tabbert et al., 2006). Moreover, DEK contains an RNA binding motif and is found associated with cellular RNA (Baltz et al., 2012; Beckmann et al., 2015; Castello et al., 2012; Guo et al., 2021a; Guo et al., 2021b). These data delineate DEK as a multifaceted protein subject to complex regulation and impacting multiple nuclear processes.

To deepen our insight in DEK´s functional mechanisms, in particular related to DNA replication, we investigated the dynamic localization of DEK in the nucleus throughout the cell cycle by live and high-resolution microscopy including optical nanoscopy (Diaspro and Bianchini, 2020). We discovered that, in late S-phase, DEK consistently forms

discrete foci, now termed DEK bodies, which locate in close proximity of active heterochromatic replication sites. We then conducted an imaging-based siRNA screen to identify cellular regulators involved in the formation of DEK bodies. Besides methyltransferases, acetylases and proteins involved in checkpoint regulation, we surprisingly identified SUMO and ubiquitin enzymes, with SUMO1/3 as top candidates. Driven by this finding, we show that DEK is targeted by SUMOylation *in vitro* and in cells, and that SUMO modification of DEK is required, at least in part, for the establishment of DEK bodies. Our data refine our previous knowledge about DEK´s involvement in DNA replication and put forward DEK as a co-factor in the duplication of specific heterochromatic regions including pericentric chromatin. These findings provide further insights into the post-translational-dependent mechanisms by which DEK contributes to heterochromatin formation and maintenance with consequences for a variety of cellular processes.

Results

A sub-population of DEK molecules assembles into complex nuclear bodies at sites of late DNA replication.

To obtain precise insights into the distribution of DEK on chromatin in the context of DNA replication, we first investigated its sub-nuclear localization throughout the cell cycle along with EdU or PCNA as replication fork markers. Confocal imaging of endogenous DEK in EdU-labeled MCF10A cells (**Fig. 1A**) revealed an archetypical, mostly uniform and punctuate nuclear pattern of DEK in G1/G2 and early to mid S-phase, in full agreement with previous observations as outlined above. In late S-phase, however, as far undescribed, distinct focal assemblies of DEK became visible which located in spatial proximity or within DNA replication foci as assessed either by EdU pulse labeling or by PCNA immunofluorescence (**Fig.1B, C**). We made the same observation in wild-type U2-OS cells **(Fig. 1D upper row)** and primary BJ-5ta cells **(Fig. S1A, B, C)**, where S-phase was assessed by PCNA staining. Interestingly, in contrast to non-malignant MCF10A cells, no or very few foci occurred under identical conditions in MCF7 (non-invasive and poorly aggressive with low metastatic potential) or MDA-MB-231 (invasive metastatic) breast cancer cells (**Fig. S1D, E**), suggesting cell-type dependent differences and a potential correlation of DEK foci formation with malignancy. Focal accumulations of DEK in late S-phase were observed also with a previously established U2-OS cell line carrying a TALEN-mediated knock-in of eGFP-DEK, U2-OS KI eGFP-DEK cells (Ganz et al., 2019) (**Fig. 1D lower panel, Fig. S2A, B)** and when DEK was expressed transiently in U2-OS cells (**Fig. S2C).** Such accumulations were not detected in eGFP-only expressing cells **(Fig. S2D)**. Thus, formation of late replication-associated foci is a specific feature of DEK, possibly regulated by cellular state, for which the U2-OS KI eGFP-DEK cell line is a valid reporter. We term these foci henceforth "DEK bodies".

To get deeper structural insight into DEK bodies and their spatial relation to active replication sites we turned to super-resolution imaging. We visualized firstly the granular fine structure of endogenous DEK bodies in MCF10A cells by Stimulated Emission Depletion (STED) and single-molecule localization (SML) microscopy **(Fig. 2A, B)**. Secondly, we investigated the organization of nascent DNA and replication forks within

DEK bodies of wild-type U2-OS cells by labeling with EdU and PCNA-specific antibodies followed by 3D Structured Illumination Microscopy (3D-SIM). The data show that DEK, PCNA, and EdU signals do not colocalize but are juxtaposed within DEK bodies (**Fig. 2C, D**). Intensity line plots (**Fig. 2E**) highlight the close spatial proximity, which was also observed in primary BJ-5ta foreskin fibroblasts (**Fig. S1**). This result is in line with previous data showing that DEK is not part of the replisome but binds to nascent chromatin in the vicinity of the replication fork (Ganz et al., 2019).

DEK bodies assemble at replicating heterochromatic regions

The occurrence of DEK in a subpopulation of PCNA- and EdU-positive nuclear bodies in late S-phase let us hypothesize that DEK accumulates at sites of replicating heterochromatin. To test this hypothesis, we conducted immunofluorescence analyses of three well-established chromatin markers in S-phase cells - H3K9ac for euchromatin (Wang et al., 2008), H3K27me3 for facultative heterochromatin (Trojer and Reinberg, 2007), and H3K9me3 for constitutive heterochromatin (Nakayama et al., 2001) (**Fig. 3A, B and Fig. S3 A-C**). We observed that within DEK bodies, DEK-specific signals did not overlap with H3K9ac labeled chromatin, were spatially juxtaposed to H3K27me3 signals, and colocalized with H3K9me3 (**Fig. 3A**), strongly suggesting that DEK bodies represent regions of replicating constitutive heterochromatin. This conclusion was corroborated by Manders' coefficient analysis yielding the highest overlap for DEK and H3K9me3. The values for the coefficient M_1 indicating the fraction of DEK bodies containing the different modified histones were as follows: $M_{1(DEK\&H3K9me3)} = 0.87 \pm 0.09$, $M_{1(DEK\&H3K9ac)} = 0.22 \pm 0.12$; and $M_{1(DEK\&H3K27me3)} = 0.37 \pm 0.06$; with n=8 from two independent experiments (**Fig. 3B**). Further evidence was obtained by Proximity Ligation Assay (PLA) confirming direct interaction of DEK with H3K9me3 but not with H3K27me3 (**Fig. 3C, D**), while fluorescence recovery after photobleaching (FRAP) showed a reduced mobility of DEK in the bodies with respect to the bulk chromatin, consistent with a high level of chromatin compaction in these structures (**Fig. 3E-G**). In line with these data, DEK bodies were associated with heterochromatin, in particular with H3K9me3, also in U2-OS wild type and U2-OS KI eGFP-DEK cells **(Fig. S4)**.

Finally, we investigated the localization of DEK bodies with respect to known heterochromatic compartments such as centromeres, the Barr body, and the nuclear periphery. Maximum intensity projections of confocal Z-stacks revealed that the epigenetic mark for centromeres, the histone H3 variant CENP-A, reproducibly decorates DEK bodies from both sides (**Fig. 4A**). We did not observe spatial proximity of DEK and CENP-A outside of late S (**Fig. 4B**). A staining pattern of similar symmetry was observed for *Xist* RNA, a selective marker for the Barr body (Cerase et al., 2015) (**Fig. 4C**). No such pattern was observed in male BPH-1 cells (data not shown). Similarly, we found DEK bodies in close proximity of the nuclear periphery and Lamin-A positive nuclear invaginations only in S-phase **(Fig. 4D, E)**. Furthermore, DEK bodies corresponded to regions of dense chromatin, as shown by DNA staining with ToPro3 **(Fig. 4F)**. Altogether, our collective microscopy data strongly support a role for DEK bodies in the replication of heterochromatic chromatin regions in late S-phase.

DEK body dynamics is affected by DNA replication inhibitors

Having assessed that DEK bodies are implicated in the replication of heterochromatin in fixed cells, we were interested in exploring and manipulating their dynamics in living cells. To this end, we transiently expressed mRFP-PCNA in U2-OS KI eGFP-DEK cells *via* a PCNA chromobody plasmid and performed time-lapse imaging experiments at a spinning disk confocal microscope (**Fig. 5A**). DEK body formation was evaluated manually and occurred reproducibly in every cell in late S-phase. 2.7 ± 1 DEK bodies formed per cell and persisted for 48 ± 30 min until G2 entry (**Fig. 5B, C**). All DEK bodies colocalized with PCNA-positive replication foci, but only 45% of the latter contained DEK. A similar behavior was observed also in MCF10A cells transiently expressing fluorescent fusions of DEK and PCNA, with a higher number of DEK bodies (6.7 ± 2.2) that persisted for a longer time (189 ± 38 min). 82 ± 18% of DEK bodies colocalized with PCNA-marked replication foci in these cells (object-based analysis of 19 DEK bodies) **(Fig. S5)**.

To determine whether impairment of DNA replication would affect DEK body formation, we employed hydroxyurea (10 µM), aphidicolin (200 nM) and camptothecin (100 nM), three well-established replication fork inhibitors acting by different mechanisms which at

these low doses slow down but do not arrest replication **(Fig. 6A)** (Deutzmann et al., 2015; Vesela et al., 2017). The treatments, in particular aphidicolin and camptothecin, did not affect viability, but triggered accumulation of cells in G2, followed by normal mitoses (**Fig. 6B upper panels**). Of the three substances, aphidicolin showed the most prominent effect reducing DEK body number by 47%, and prolonging DEK body lifetime by 28%. Camptothecin had a similar but more moderate effect, and hydroxyurea did not alter DEK body dynamics (**Fig.6B lower panels**). The differential response to the replication stress inducers strongly suggests that DEK bodies are implicated in the replication of DNA structures particularly susceptible to aphidicolin, such as common fragile sites that in turn can coincide with highly repetitive heterochromatic regions like centromeres or ribosomal DNA (Lukusa and Fryns, 2008), in line with our data on the colocalization of DEK bodies with heterochromatin histone marks.

Down-regulation of the SUMO pathway increases DEK body number in an siRNA screen for DEK body regulators

As our time-lapse microscopy experiments could measure the effect of mild replication stress on DEK body formation, we chose to perform an imaging-based unbiased approach to identify DEK body regulatory factors. To this end, we set up a high-throughput siRNA screen targeting genes from a recently published DEK interactome (Smith et al., 2018), DEK-related factors derived from the literature, and genes selected from a commercial library targeting the DNA damage response that responded to the gene ontology terms "DNA replication" and "S-phase". The final library consisted of 332 genes (678 siRNAs total, at least two siRNAs per gene, **Supplementary Table S1**).

Experimental conditions and a suitable image analysis pipeline using CellProfiler software were established in two pilot screens including aphidicolin and camptothecin as positive controls (**Supplementary Material and Methods**, **Supplementary Table S2** and **Fig. S6**). The primary screen targeting the chosen gene panel was performed in duplicate and a Z-score of the log-transformed mean number of DEK bodies per cell was calculated for each siRNA (**Fig. 7A**). This resulted in 51 candidate siRNAs enhancing (up-regulators)

and 29 candidates reducing DEK body numbers (down-regulators) corresponding to 44 and 28 different genes, respectively (**Fig. 7B, C**). The number of DEK bodies per cell was the most sensitive and robust readout for DEK body formation also in the automated approach, reproducing the results from the manual experiments (**Fig. S6D**). To further confirm the identified hits, we conducted a low-throughput secondary screen with 28 siRNAs (**Fig. 7D**). In general, the values for the Z-score obtained in this secondary screen were smaller and more variable, most likely due to the different, less automated imaging system and the manual dispensing of transfection reagents. Still, the effect of up- and down-regulation of DEK body number was confirmed (**Fig. 7D**). Importantly the secondary screen validated four top DEK body up-regulators from the primary screen belonging to the SUMO pathway, among them the E1 activating enzyme SAE1 and the E2 conjugating enzyme UBE2I. Our screening approach thus determined that the SUMO pathway plays a major role in the regulation of DEK body formation.

DEK is a SUMO target and SUMOylation is required for the formation of DEK bodies

As both screening approaches showed that down-regulation of the SUMOylation pathway resulted in the up-regulation of the number of DEK bodies in U2-OS KI eGFP-DEK cells, we hypothesized that DEK may be a target of SUMOylation and that, in turn, its SUMOylation state may modulate the ability to form DEK bodies in late S-phase. Indeed, the DEK protein sequence contains a SUMOylation consensus motif ψ-K-X-E/D (Zhao et al., 2014) at lysine 261 (AKRE, https://sumo.biocuckoo.cn/online.php) and DEK has been shown to be SUMOylated under stress conditions in a mass spectrometry study of H_2O_2-treated SH-SY5Y neuroblastoma cells (Grant, 2010). To test if DEK can be SUMOylated under steady-state conditions, we focused on SUMO 1 and 3, as both were strong hits in the screens. We first conducted co-immunoprecipitations using DEK-antibodies, showing direct interaction of DEK with SUMO1 and SUMO2/3 (**Fig. 8A**). To narrow down a specific SUMO paralog targeting DEK, we used HeLa cells stably expressing 6xHis-SUMO-1, 6xHis-SUMO-2, or 6xHis-SUMO-3 allowing for subsequent nickel affinity purification of SUMOylated factors. This approach revealed modification of DEK by all SUMOs, yet with

the strongest activity for SUMO-3 (which also generates multimers) (**Fig. 8B, C**). As we could confirm the existence of SUMOylated DEK in cells under physiological conditions, we were next interested to map the specific SUMOylation sites in the DEK primary sequence. As indicated above, DEK harbors one SUMOylation consensus motif at position lysine 261. However, mutation of this site did not yield substantial reduction in SUMOylation in *in vitro* SUMOylation assays using recombinant GST-tagged DEK as acceptor (**Fig. S8**). Therefore, we screened the DEK sequence for additional potential SUMOylation sites, which revealed amino acids 61, 62 and 144, 145 as suitable SUMO attachment sites. Interestingly, mutation of these sites individually, as was the case with 261, did not result in a substantial reduction of SUMOylation activity on DEK (**Fig. S8**). However, DEK molecules carrying mutations at all five sites showed an 80% reduction in SUMOylation (**Fig. 8D, E**), highlighting that most of the potential SUMOylation sites were captured by our mutational approach. Given that we had created a DEK mutant that is substantially less susceptible to SUMOylation, we then investigated the ability of such a mutant to form DEK bodies in cells. Therefore, we established a stable cell line expressing this mutant as eGFP-fusion and monitored the occurrence of DEK bodies as described above. No DEK body formation was observed in cells expressing SUMO-less DEK (GST-DEK SUMO mut, **Fig. 8F)**. This result points at a complex regulation of DEK bodies by SUMO, where SUMOylation of DEK seems to be required for DEK body formation while DEK body numbers increase when the overall SUMOylation capacity of the cell is reduced. In summary, these data uncover a yet unrecognized link between SUMO and DEK and establish SUMOylation as a crucial regulator of DEK body formation.

Discussion

DEK bodies reflect the formation of transient DEK multimers in the nucleus during late DNA replication

We report on the discovery of DEK bodies, DNA replication-dependent focal assemblies of the oncoprotein DEK in the cell nucleus. DEK bodies were observed in every cell during mid-late S-phase in primary, immortalized and in transformed, yet non-malignant cell lines. This finding was somewhat surprising as the large majority of immunofluorescence studies have described DEK as a factor dispersed in an uniform punctuate pattern in the nucleus (Hu et al., 2007; Kappes et al., 2001; Matrka et al., 2015). Intriguingly, the existence of non-covalent DEK multimers was proposed in the past based on biochemical analyses of DEK-chromatin complexes (Kappes et al., 2004b), but had eluded demonstration in live cells so far. We interpret DEK bodies as the cellular correlate of these non-covalent multimers, whose detection by microscopy was most likely facilitated by fusion of DEK to GFP. This reporter has a propensity to form aggregates at high concentrations (Stepanenko et al., 2021) contributing to slow down the dynamics of DEK bodies and make them amenable to microscopic imaging. The focused inspection of high resolution immunofluorescence images revealed that DEK bodies are distinct sites of active replication containing endogenous DEK in close spatial proximity to nascent DNA and the replication processivity factor PCNA consistent with previous iPOND analyses (Alabert et al., 2014; Aranda et al., 2014; Garcia et al., 2017; Lossaint et al., 2013; Ribeyre et al., 2016; Sirbu et al., 2013). The number and persistence of DEK bodies varied substantially between MCF10-A and U2-OS cells, the former showing a more pronounced expression of these bodies. Unfortunately, a direct quantitative comparison of DEK bodies between these two cell lines is not appropriate as DEK was expressed endogenously in U2-OS and transiently in MFC-10A cells, and imaging was performed with different microscopes in different laboratories. But in both cases, DEK bodies were sparingly observed also in mid S-phase, probably reflecting cell-to-cell variability in replication timing (Zhang et al., 2017).

Previous data suggesting that DEK may occur in defined nuclear substructures included coimmunoprecipitation and spurious colocalization with PML (Ivanauskiene et al., 2014),

and the redistribution to interchromatin granule clusters in the presence of deacetylase inhibitors (Clarey 2005, McGarvey 2000), but no link to the cell cycle or DNA replication was made in these studies.

In general, the formation of nuclear bodies has emerged as an organizing principle for the orchestration of genome-associated functions in time and space. Different nuclear bodies have been described in which proteins, often together with RNAs, are sequestered and released according to demand, including the nucleolus, nuclear speckles and paraspeckles, PML, Cajal bodies, 53BP1 and other bodies (Dundr, 2012). Post-translational modifications, as well as weak intermolecular interactions mediated by intrinsically disordered protein domains, seem to play a fundamental role in the genesis and dynamic composition of these membrane-less nuclear condensates. It is thus not surprising that DEK, a DNA and RNA-binding protein carrying several post-translational modifications and an intrinsically disordered region (aa 170-270) is prone to form nuclear bodies.

DEK bodies form at heterochromatic regions

Our data provide strong evidence that DEK bodies are sites of heterochromatin replication. These include their narrow window of appearance during late S-phase, the association with heterochromatin histone marker H3K9me3, the high level of chromatin compaction and the reduced mobility of DEK in bodies, and the local proximity to the Barr body and centromeric chromatin CENP-A. The lack of colocalization between CENP-A and DEK reported previously (Ivanauskiene et al., 2014) was most likely due to the fact that endogenous DEK bodies are transient structures visible only in late S-phase, and may thus escaped detection.

The reproducibility with which DEK bodies form in consecutive cell cycles support the notion that they reflect a conserved and necessary passage in genome duplication. The striking appearance of the centromeric and *Xist*-specific signals as dots decorating DEK bodies (Fig. 4A, C) are suggestive of newly replicated chromosomes tethered to these

bodies at regions rich of secondary structures that are preferred DEK substrates *in vitro*, are difficult to replicate and prone to breakage. In line with this observation, downregulation of DEK resulted in the formation of anaphase bridges, the appearance of DNA breaks on mitotic chromosomes and of 53BP1 bodies in daughter cells (Deutzmann et al., 2015). While, in the latter study we described how DEK affects DNA replication globally, the results obtained here suggest a more specialized function of the protein, at specific genomic regions including the centromere and possibly other constitutive heterochromatic regions. These findings may explain the relatively mild effects of DEK downregulation on replication fork progression as assessed by bulk fiber assays. The assessment of DEK´s influence on the processivity of the replication fork therefore awaits DNA combing experiments coupled with the detection of heterochromatic fibers.

DEK bodies are novel players in the SUMO-dependent regulation of heterochromatin replication

DEK body up-regulating siRNAs show a remarkable enrichment in proteins involved in post-translational modification pathways. Histone acetyltransferases (HATs) ACTL6A, KAT2B, TAF5L and TAF6L, histone deacetylase HDAC2, histone methyltransferases SUV39H1/2 and SUMOylation-related proteins UBE2I, SUMO1 and SAE1 are among the top hits. Acetylation of DEK by KAT2B/PCAF has been shown previously (Cleary et al., 2005). In this study, overexpression of KAT2B led to an accumulation of DEK in interchromatin granule clusters (IGCs), which are nuclear domains enriched in pre-mRNA splicing factors. Pharmacological inhibition of KAT2B in turn, blocked the relocation of DEK into IGCs, opposite to the effect of downregulation of KAT2B on DEK bodies. This finding further underscores that DEK bodies are distinct from ICGs, which are more abundant (20-30 clusters per cell) and are stable during the whole interphase (Mintz et al., 1999).

Here, we focused on SUMOylation as a post-translational modification that has recently emerged as a major regulator of heterochromatin formation and replication. We

substantiate the result from the siRNA screen by showing that DEK is modified by SUMO in cells, and that SUMO-less DEK does not form bodies. Recent data also show that SUV39H1 promotes SUMOylation and *de novo* targeting of HP1α to pericentric heterochromatin, independently of its methyltransferase activity. The latter is required in a second step to maintain and propagate heterochromatin (Maison et al., 2016). This knowledge aligns well with our screening data, where downregulation of both SUV39H1 and SUMO increase the number of DEK bodies suggesting the DEK plays a role in "seeding" constitutive heterochromatin after DNA replication (Maison et al., 2016). Our data also support and refine the conclusions from our previous study showing that DEK is essential for heterochromatin integrity, and has a Su(var)like activity by enhancing HP1α binding to H3K9me3 (Kappes et al., 2011a).

Heterochromatin preferentially harbors repetitive sequences causing non-B DNA secondary structures. Their replication requires specialized polymerases that can sustain fork progression in the presence of difficult-to-replicate regions, but on the other hand lack proofreading activity and are therefore error-prone and mutagenic (reviewed in (Tsao and Eckert, 2018)). The switch from replicative polymerase to these specialized, or translesion polymerases, at the replisome is regulated by ubiquitylation and SUMOylation (Despras et al., 2016; Mailand et al., 2013). In particular, SUMOylation is required on the one hand, for driving specialized polymerases to nascent DNA while, on the other hand, enabling discharge from the replisome after bypass of the problematic site to limit replication errors and prevent mutagenesis, as shown for Pol η (Despras et al., 2016; Guérillon et al., 2020). We hypothesize first, that DEK in bodies may participate in the maturation of nascent heterochromatin after replication involving specialized polymerases and that this step requires SUMOylation of DEK, explaining why a non-SUMOylatable mutant of DEK does not form bodies. Secondly, we interpret the increase in the number of DEK bodies observed after downregulation of the SUMO pathway as consequence of impaired turnover of these specialized replisomes and the replacement by proof-reading replicative polymerases. Interestingly, also Polη was reported to be SUMOylated at multiple sites which can substitute for each other, so that simultaneous

mutation of all SUMOylation sites was reported to be necessary to achieve dysregulation of the Polη dynamics (Guérillon et al., 2020).

At present, we have no straightforward interpretation for the reduction in DEK bodies that seems to result from increased DNA replication stress, as highlighted by the lack of DEK bodies in highly cancerous MDA-MB-231 cells and in the presence of aphidicolin, and - to a lesser extent - camptothecin. We speculate that under these conditions replisomes at difficult-to-replicate-regions would be more likely to collapse leading to DNA breakage before chromatin maturation, thus impairing DEK body assembly. Accordingly, we found increased γH2AX foci after DEK downregulation in a previous study (Kappes et al., 2008). The propensity to DNA strand break formation is a well-established feature of difficult-to-replicate-regions such a common fragile sites (CFS) and their stability has been shown to depend on specialized polymerases (Bergoglio et al., 2013; Bhat et al., 2013).

In conclusion, the link established here between DEK and the establishment of constitutive heterochromatin during replication sheds new light on the mechanisms of action of this so far enigmatic oncoprotein. Some of its pleiotropic effects may be associated with the seeding of (constitutive) heterochromatin, a process that broadly impacts the cell´s epigenetic landscape and transcriptional program and is essential for maintaining genomic stability (Penagos-Puig and Furlan-Magaril, 2020).

Materials and Methods

Cell culture

Mammary epithelial, non-transformed cell line MCF10A (ATCC CRL-10317) was grown in DMEM:F-12 (Dulbecco's Modified Eagle Medium:Nutrient Mixture F-12) (1 : 1) medium (Gibco, 11330057) supplemented with 5% horse serum (HS), 2 mM L-glutamine and 1% penicillin/streptomycin (Sigma-Aldrich, G6784), 10 µg/ml insulin (Sigma-Aldrich, I9278), 0,5 µg/ml hydrocortisone (Sigma-Aldrich, H0888), and 20 ng/ml of human Epidermal Growth Factor (hEGF, Sigma-Aldrich, E9644) added freshly before the use of full medium. MCF7 cells and MDA-MB-231 cells (ATTC HTB-26) were grown in DMEM medium (Gibco, 11330057) supplemented with 10% fetal bovine serum (FBS, Euroclone, ECS0180L), 1% penicillin/streptomycin (Sigma-Aldrich, G6784) and 2 mM L-glutamine. Cells were grown on 10 cm^2 dishes (about 10-12 cell passages) at 37°C in 5% CO_2. For any experiments on fixed cells, cells were plated on glass coverslips coated with 0.5% (w/v) pork gelatine (Sigma-Aldrich, G2500) dissolved in phosphate buffer saline (PBS) and autoclaved. U2-OS osteosarcoma (kind gift from G. Marra, University of Zurich, Switzerland) and U2-OS KI eGFP-DEK cells (described in (Ganz et al., 2019)) were cultured in McCoy's 5a modified medium (Gibco, Life Technologies) supplemented with 10 % FBS (Capricorn Scientific), 100 U/ml penicillin and 100 µg/ml streptomycin (both Gibco, Life Technologies). For selection after transfection, Geneticin (Thermo Fisher Scientific) was added at concentrations of 200 µg/ml for U2OS KI eGFP-DEK WT, eGFP-DEK SUMOmut Δ61,62, eGFP-DEK SUMOmut Δ261 and eGFP-DEK SUMOmut Δ61,62,144,145,261 cells. During all experiments using siRNA-mediated protein down-regulation penicillin and streptomycin were omitted from the medium. HeLa S3 cervical adenocarcinoma cells and HeLa 6xHis-SUMO cells (kind gift from Ronald T. Hay, University of Dundee, UK) were cultured in DMEM medium (Gibco, Life Technologies) supplemented with 10% FBS (Capricorn Scientific), 100 U/ml penicillin, 100 µg/ml streptomycin (both Gibco, Life Technologies) and 6 mM L-glutamine (Gibco, Life Technologies). hTERT immortalized BJ-5ta foreskin fibroblasts were cultured in a 4:1 mixture of DMEM medium and Medium 199 (Gibco, Life Technologies) supplemented with 10% FBS, 4 mM L-glutamine and 10 µg/ml hygromycin B (Calbiochem).

Transfection and cell sorting

For detection of endogenous PCNA, U2-OS KI eGFP-DEK or MCF10A cells were transfected with a PCNA-specific chromobody expression vector (pCCC-TagRFP, Chromotek) using Effectene (Qiagen U2-OS KI eGFP-DEK) or Lipofectamine 3000 (Invitrogen) according to the manufacturer´s instructions. For U2-OS KI eGFP-DEK, the medium was exchanged 4 h after transfection and the cells were incubated for further 20 h before imaging. For MCF10A cells, the medium was exchanged about 12 h post-transfection, and experiments were carried out between 24 to 48 h after transfection. For the generation of U2-OS cell lines stably expressing eGFP-DEK mutants, pEGFP-N1-hDEK plasmids encoding the mutated DEK versions were transfected into U2-OS DEK knockout cells using Lipofectamine 3000 reagent (Invitrogen). 3-9 days after transfection (depending on cell growth and viability) selection was carried out with geneticin at a final concentration of 200 µg/ml. To create cell lines with a uniform GFP fluorescent signal, the stable transfected cell lines were bulk sorted for a moderate fluorescence signal using a FACSAria IIIu Cell Sorter (BD Biosciences).

Immunochemical methods

All antibodies used in this study are listed in **Supplementary Table 3**.

For the detection of nascent DNA MCF10A and U2-OS cells were incubated with EdU (10 µM) for 25 min or 10 min respectively. After fixation and washings, the incorporated precursor was detected with the Click-iT EdU Imaging Kit or Click-iT EdU PLUS Imaging Kit, Molecular Probes with Alexa Fluor 647 or Alexa Fluor 594 azide dye. When necessary, nuclei were counterstained with TO-PRO™-3 Iodide (Invitrogen, 1:2000). For immunofluorescence detection of DEK or PCNA cells were fixed with 4% PFA/PBS (20 min) followed by quenching with 50 mM NH_4Cl/PBS (10 min). Cells were permeabilized in methanol (5 min, -20°C), incubated in 1% BSA/PBS for 30 min followed by incubation with the primary antibody dilution (diluent 1% BSA/PBS) in a dark, humid chamber (overnight, 4 °C). After washing with PBS for three times, cells were incubated with the appropriate secondary antibody (diluent 1% BSA/PBS) in the dark chamber (1 h). Coverslips were mounted either in ProLong™ Diamond Antifade Mountant (Invitrogen, P36961) or Aqua Polymount (Polysciences, 186065).

RNA-FISH

Xist RNA was detected with Stellaris® RNA FISH (Human XIST with Quasar® 570 Dye, Biosearch Technologies, SMF-2038-1) according to the manufacturer's instructions. After the hybridization step, Click-iT EdU detection was performed with the use of Alexa Fluor 647 azide. The samples were washed with PBS and mounted in ProLong™ Diamond Antifade Mountant.

Proximity Ligation Assay

Proximity Ligation Assay (PLA) was performed in MCF10A cells using the DuoLink® in situ Orange detection reagent (DUO92102, Sigma-Aldrich) according to the manufacturer's instructions. Cells were incubated for 60 min at 37°C in a humidified chamber in PLA blocking solution, followed by 1h at RT with antibody solution containing primary antibodies. After washing, the ligation and amplification steps were performed. Anti-mouse Alexa Fluor 488 was used as secondary antibody. Nascent DNA was detected as described above. Samples were mounted in ProLong™ Diamond Antifade Mountant.

Confocal and widefield microscopy

Confocal images depicting MCF10A cells were acquired with a Leica TCS SP5 confocal laser-scanning microscope, using an HCX PL APO 100x/1.40/0.70 oil immersion objective lens (Leica Microsystems, Mannheim, Germany). Excitation was provided with a white laser, enabling the best condition as function of the utilized dye, namely: 488 nm (for Alexa Fluor 488), 532 nm (for Atto 532), 554 nm (for Quasar 579), 594 nm (for Alexa Fluor 594), and 647 nm (for Alexa Fluor 647 and Atto 647N), with relevant emission detection bands 495 – 525 nm (for Alexa Fluor 488), 540 – 620 nm (for Atto 532), 560 – 625 nm (for Quasar 570), 655 – 725 nm (for Alexa Fluor 647 and Atto 647N). Signals in different channels were detected with photomultipliers (PMT) or Hybrid Detectors (HyD). Typical pixel size is 40 nm. For images showing U2-OS cells a Zeiss LSM880 was used equipped with a Plan-Apochromat 40x/1.4 Oil DIC M27 objective and 405 nm (UV-photodiode), 488 nm (Argon ion), 561 nm (DPSS) and 633 nm (HeNe) laser lines. 12 bit images were recorded at a resolution of 1024x1024 pixels, with 2x digital zoom and a

scanning speed of 7. Live cell imaging of MCF10A was performed with a NIKON A1R confocal microscope equipped with a stage top incubator and a hardware autofocus. Images were acquired in 1024 x 1024 pixel format, with a pixel size of about 85 nm, without averaging. Images were acquired every 5 min or 10 min over 10-12 hours. Excitation was provided with a CW Diode Laser (Coherent, Cube 488-50) at 488 nm for eGFP (DEK-eGFP) and a CW Diode Pumped Solid State Laser (DPSS, Melles Griot, 85YCA) at 561 nm for RFP (pCellCycleChromobody®-TagRFP). Widefield images of MCF10A cells were acquired on an AxioObserver or a CellObserver HS microscope (Zeiss) equipped with a PlanApochromat 40X/1.40 oil objective, a HXP 120 mercury arc lamp and an Axiocam MRm CCD camera (1300 X 1030 pixels). Fluorescence signals were detected with 493/517 nm (GFP and AlexaFluor 488), 590/612 nm (AlexaFluor 568) and 640/690 nm (AlexaFluor 647) filter sets. DEK bodies in U2-OS cells stably expressing eGFP-DEK or mutants thereof were imaged on a Celldiscoverer 7 widefield imaging platform (Zeiss) equipped with a Plan-Apochromat 20x/0.7 air objective and an Axiocam 512 camera. Binning was set to 2,2 and the exposure time to 200 ms. Cells were plated in 96-well imaging plates (Ibidi), 10 positions were recorded per well and images acquired every 12 min for a total duration of 22 h.

STED microscopy

STED measurements were performed on a Leica TCS SP5 gated-STED microscope, using an HCX PL APO 100x100/1.40/0.70 oil immersion objective lens (Leica Microsystems, Mannheim, Germany). Emission depletion was performed with 592 nm CW STED laser. The 592nm STED laser power was 50% of 350 mW maximum power (about 175 mW). Excitation was provided with a white laser at 488 nm wavelength for Alexa Fluor 488 and with emission detection band 495 – 525 nm, with 1.50 ns - 9.50 ns time gating using a hybrid detector (Leica Microsystems). Typical pixel size is 40 nm, with the pixel dwell time of 2 µs and 48 lines averaging.

SMLM Imaging

For the super-resolution SMLM imaging of DEK, MCF10A cells were washed with pre-warmed PBS 3x, and fixed with formaldehyde (3,7%, methanol free). After subsequent

blocking with blocking buffer (0,1% Triton X-100 and 3% w/v of bovine serum albumin (BSA)) for 1h at room temperature, samples were incubated with the primary antibody (overnight at +4°C). F(ab')2-Goat anti-Mouse IgG (H+L) Cross-Adsorbed Secondary Antibody, Alexa Fluor 647 was used as secondary antibody (ThermoFisher Scientific, A21237, 1:250, 45min in RT). Samples were post-fixed in PFA 2% for 5 min and stored in PBS at +4°C until applying GLOX buffer containing a glucose oxidase-based oxygen scavenging system for imaging (Bates et al., 2007). A commercial N-STORM & N-SIM TIRF Eclipse Ti2 microscope equipped with an oil immersion objective (CFI SR HP Apochromat TIRF 100XC Oil, NA 1.49) was used to acquire 20000 frames at a frame rate of 30 ms using oblique incidence excitation and sCMOS camera (ORCA-Flash4.0, Hamamatsu Photonics K.K.). Alexa Fluor 647 labeled samples were switched into dark state with the 647 nm laser and reactivated with the 405 nm laser. Imaging cycles consisted of 1 activation frame followed by 3 read-out frames. Acquisition was performed using the continuous STORM filter and hardware autofocus. Image reconstruction was performed with the NIS-Elements and N-STORM Analysis imaging software.

Fluorescence Recovery After Photobleaching

FRAP measurements were performed on a Leica TCS SP5 confocal laser scanning microscope. The experiments were conducted under a controlled environment in a dedicated live-cell imaging chamber (5 % of CO_2, 37°C, 90 % humidity). Briefly, regions of interest (ROI) of 2 µm x 2 µm were selected and photobleached with 100% laser power (55 µW @488nm). Subsequently, fluorescence recovery was monitored for about 3 min, all traces were min/max normalized, averaged and the standard error was calculated. The parameters used for the FRAP measurement are listed in Table 1.

Table 1. Parameters used in FRAP experiments and data acquisition

	DEK-eGFP
FRAP image acquisition	512 x 512 pixels
	40 nm pixel size
	Excitation wavelength: 488 nm 10%
	Emission: 500 nm - 580 nm
	1400 Hz scan speed
Pre-photobleaching	2 frames (time/frame 0,37 s)
Photobleaching protocol	5 frames (time/frame 0,37 s); 2 x 2 µm square ROI; 100% 488 nm laser
FRAP imaging	90 frames (time/frame 2s); 10% 488 laser

Image analysis

Image analysis was performed using ImageJ (http://imagej.nih.gov/ij/) and Leica LAS X F (Leica Microsystems GmbH). The number of DEK bodies was obtained from Z-stack Maximum projections on which was performed Find Maxima function (with the prominence about 80-90 depending on the level of immunofluorescence staining).

For finding the number of DEK bodies colocalizing with PCNA DNA replication foci, the object-based analysis was performed by using an ImageJ plugin JaCoP (Bolte and Cordelieres, 2006). For the analysis of FRAP data, first the photobleached region within cell nucleus was cropped. After this, the mean intensity of the ROI along the time experiment was plotted, and the curves were combined and normalized to the minimum and maximum of the combined curve. Finally, each of the curves was fit. The tau parameter was obtained from the average of all fits. This procedure was performed for the case of DEK within DEK bodies and for DEK within the other chromatin regions.

Preparation of cell extracts, immunoprecipitation, and immunoblot analysis

Whole cell lysates were prepared *via* scraping cells in PBS containing protease inhibitors (Complete Protease inhibitor cocktail, Roche). Cells were pelleted and resuspended in hot SDS lysis buffer (0.5 % SDS, 50 mM Tris pH 7.5), incubated for 10 min at 95°C, centrifuged, resulting samples were frozen in liquid nitrogen and stored at -80°C.

For preparation of cell lysates for immunoprecipitation, cells were harvested in hypotonic buffer (20 mM HEPES pH 7.4, 20 mM NaCl, 5 mM $MgCl_2$, 10 mM NaF, 1 mM Na-vanadate, 1 mM PMSF, 1x Complete Protease Inhibitor Cocktail (Roche)) supplemented with 10 mM N-Ethylmaleimide (NEM). After centrifugation (5 min, 500 x g, 4 °C), the resulting cell pellet was resuspended in hypotonic buffer to 1×10^6 cells per 100 µl. For extraction of chromatin-bound DEK, 450 mM NaCl and 0.5% NP-40 was added, incubated for 15 min on ice and harvested at 10,000 x g for 10 min at 4 °C.

For preparation of cell lysates for nickel affinity chromatography (6xHis-SUMO pulldown), a previously published protocol was carried out (Tatham et al., 2009). HeLa cells stably expressing 6xHis-SUMO-1 (accession number P62165), 6xHis-SUMO-2, (accession number CAG46970) or 6xHis-SUMO-3 (accession number CAA67897) as well as HeLa

Rachel cells as negative control were grown in 10 cm cell culture dishes to 80% confluency. Cells were harvested *via* scraping in 5 ml PBS containing 10 mM NEM. Crude cell extracts for visualization of SUMO conjugates were prepared with 1 ml of cell suspension. Crude samples were centrifuged (2 min, 1000 x g, RT), the supernatant was removed, and the pellet was resuspended in 200 µl 2x Laemmli sample buffer (200 mM imidazole, 5% SDS, 150 mM Tris-HCL pH 6.7, 30% glycerol, 0.0025% bromophenol blue). Samples were boiled for 5 min at 95°C, then mechanically disrupted by passing through a 20G needle 20 times using a 1-ml syringe and supplemented with 750 mM β-mercaptoethanol. The remaining 4 ml cell suspension was centrifuged (5 min, 3000 x g, RT) and the pellet was dissolved in 5 ml of cell lysis buffer (6 M Guanidinium-HCl, 10 mM Tris, 100 mM sodium phosphate buffer pH 8.0, 0.1% Triton X-100, 10 mM NEM). Samples were supplemented with 5 mM β-mercaptoethanol and 5 mM imidazole and sonicated on ice for 40 s at medium power. Cell debris was removed by centrifugation (15 min, 3000 g, RT) and the supernatant was stored at -20 °C.

For immunoprecipitation all buffers were supplemented with 10 mM NEM. 6 µl of polyclonal rabbit α-DEK K-877 (Kappes et al., 2008) was used for 100 µl cell lysate. After 30 min incubation on ice, 100 µl of a 25 % Protein A/G Plus-Agarose (SantaCruz) was added and rotated for 1 h at 4 °C. After washing three times with extraction buffer (20 mM HEPES pH 7.4, 300 mM sucrose, 450 mM NaCl, 0.5 mM $MgCl_2$), immune complexes were eluted from the beads with 2 % SDS and 5 % β-mercaptoethanol for 1 h at 37 °C. IP samples were concentrated using methanol-chloroform protein precipitation (Wessel and Flügge, 1984), solubilized in 1x SDS sample buffer and boiled for 10 min at 99 °C. Samples were separated by SDS-PAGE, blotted to nitrocellulose and analyzed by immunoblotting.

For nickel affinity chromatography for purification of 6xHis-SUMO proteins all buffers were supplemented with 10 mM NEM. The prepared cell lysates were subjected to chromatography using HisPur Ni-NTA Resin (Thermo Fisher Scientific). The lysate was added to 100 µl packed volume of beads per sample. Bead slurry was pre-washed three times with ten bead volumes of cell lysis buffer (6 M Guanidinium-HCl, 10 mM Tris, 100 mM sodium phosphate buffer pH 8.0, 0.1% Triton X-100). After tumbling the mixture overnight at 4°C, the bead suspension was centrifuged (2 min, 750 g, RT) and the

supernatant carefully removed. Beads were washed once with 4 ml of cell lysis buffer containing 5 mM β-mercaptoethanol and once with 4 ml of 8.0 pH wash buffer (8 M Urea, 10 mM Tris, 100 mM sodium phosphate buffer pH 8.0, 0.1% Triton X-100, 5 mM β-mercaptoethanol). Subsequently, the beads were washed three times with 4 ml of 6.3 pH wash buffer (8 M Urea, 10 mM Tris, 100 mM sodium phosphate buffer pH 6.3, 0.1% Triton X-100, 5 mM β-mercaptoethanol). Finally, the beads were resuspended in 1.5 ml of 6.3 pH wash buffer and then transferred to 1.5 ml centrifuge tubes. The wash buffer was completely removed after centrifugation (1 min, 6000 x g, RT) and elution of 6xHis-SUMO proteins was performed with 100 µl of elution buffer (200 mM imidazole, 5% SDS, 150 mM Tris-HCL pH 6.7, 30% glycerol, 720 mM β-mercaptoethanol, 0.0025% bromophenol blue) per sample. The beads were boiled for 2 min at 95°C and then incubated for 20 min at RT. The eluted fraction and the crude cell extracts were separated by SDS-PAGE, blotted to nitrocellulose and analyzed by immunoblotting.

For analysis of whole cell lysates 40 µg of protein were loaded onto a 12 % SDS gel and separated by electrophoresis. Proteins were transferred onto a nitrocellulose membrane using semi-dry Western blotting.

For analysis of eluted 6xHis-SUMO proteins 100 µl of eluted fraction and input/crude samples were loaded onto a 10 % SDS gel and separated by electrophoresis. Proteins were transferred onto a nitrocellulose membrane using wet Western blotting.

DEK-IP eluates were separated on a 12.5 % SDS gel. Membranes were blocked in 5 % milk/TBS-T and incubated with primary antibodies overnight at 4 °C under mild agitation (anti-DEK K-877). Membranes were washed and incubated with HRP-conjugated secondary antibodies for 1 h at RT (Goat-anti-Rabbit-HRP, Goat-anti-Mouse-HRP, both Dako Denmark). Marker lanes were incubated with Streptavidin-biotinylated-HRP (GE Healthcare). Proteins were visualized using chemiluminescent detection with ECL Prime Western Blotting Detection Reagents (Amersham RPN2232).

Site-directed mutagenesis:

Nucleobase mutations of the DEK sequence were introduced using an improved QuickChange site-directed mutagenesis protocol (Zheng et al., 2004). The pEGFP-N1-hDEK plasmid (DEK WT sequence inserted into eGFP reporter plasmid "peGFP-N1";

Addgene 6085-1) was used as a template for mutagenesis PCR. The potential SUMOylation site 1 (bases 181 – 186) and the SUMO consensus motif (bases 778 – 789) were mutated using overlapping primers containing the desired base change. For the SUMOylation site 1 two codons were mutated (lysines 61 and 62 to alanine) in two rounds of mutagenesis. For the SUMO consensus motif one round of mutagenesis was performed (lysine 261 to alanine). For SUMO consensus motif mutation, the plasmid DNA was denatured prior to the PCR using 0.2 M NaOH to reduce the amount of colonies containing unmutated wildtype-plasmid after transformation. Primer sequences are listed in **Supplementary Table S4**.

Expression and purification of recombinant GST-tagged proteins from *E. coli*:
The mutated DEK sequences (DEK SUMOmut 61-62, 144, 145 or 261) were inserted into a GST expression plasmid (pGEX 4T-1), transformed into *E.coli* BL21(DE3) (Thermo Fisher Scientific) and purified as described before (Ganz et al., 2019). Briefly, protein expression was induced with 0.5 mM IPTG (Sigma-Aldrich) for 1.5 h. Bacteria were harvested *via* centrifugation at 4600 g for 15 min at 4°C and resuspended in resuspension buffer (20 mM Tris-HCl (PH 8.0), 1 M NaCl, 0.5 mM EDTA, 1 mM DTT) and frozen in liquid nitrogen. After thawing, bacteria were sonicated on ice and 0.5 % NP-40 (Sigma-Aldrich) was added and centrifuged at 18000g for 30 min at 4°C. The supernatant was added to 200 µl Glutathione-Sepharose 4B-beads (GE Healthcare) and incubated for 2 h at 4°C. After centrifugation, beads were washed with wash buffer with decreasing NaCl concentrations (500 mM, 300 mM, 20 mM) and GST-tagged DEK was eluted with 200 µl elution buffer (200 mM Tris-HCl (pH 8.0), 200 mM NaCl, 40 mM reduced glutathione (Sigma-Aldrich), 10 % glycerol) for 1 h at 4°C. The supernatant was frozen in liquid nitrogen and stored at -80°C.

In vitro SUMOylation:
For in vitro SUMOylation, recombinant SUMO E1, SUMO E2 (UBC9), SUMO-1 and SUMO-2 were purchased from Enzo Life Sciences (Lausen, Switzerland). Reactions were performed for 2 hours at 30°C in *in vitro* buffer (20 mM Tris-HCl (pH 7.6), 50 mM NaCl, 10 mM $MgCl_2$, 0.1 mM dithiothreitol (DTT), 1x protease inhibitor cocktail

(cOmplete™, Mini, EDTA-free Protease Inhibitor Cocktail, Roche) with or without 4 mM ATP) as described before (Aichem et al., 2019). Reaction was terminated by addition of 5x SDS sample buffer with 4 % β-mercaptoethanol and boiled for 5 min at 95°C. 20 µl reactions were mixed containing 0.71 µg recombinant protein (GST, GST-DEK, GST-DEK SUMOmut 61-62, GST-DEK SUMOmut 261 or GST-DEK Sumo mut 61,62,144, 145, 261), 450 ng SUMO E1, 1 µg SUMO E2, 3 µg SUMO-1 or SUMO2.

Acknowledgements

The authors thank Mario Faretta (European Institute of Oncology, Milan, Italy) for MCF10A cells; Elena Cerutti (Istituto Italiano di Tecnologia -IIT, Genova, Italy) for help with RNA-FISH; Annette Aichem (Biotechnology Institute Thurgau, Tägerwilen Switzerland) for help with *in vitro* SUMOylation and Christian Tischer (EMBL Heidelberg, Germany) for help with image data analysis; the core facilities FlowKon and Bioimaging Center at the University of Konstanz (Germany) for assistance in microscopy and flow cytometry experiments and the Advanced Light Microscopy Facility (ALMF) of EMBL for hosting and supporting the primary siRNA screen; Michele Oneto (IIT), Marco Scotto (IIT) Luca Lanzanò (University of Catania, Italy) and Martin Stöckl (University of Konstanz) for fruitful discussions.

Competing interests

No competing interests declared.

Funding

This research was supported by the European Union's Horizon 2020 research and innovation programme under the Marie Skłodowska-Curie (841661, to A.P-M.); by the Deutsche Forschungsgemeinschaft DFG Research Training Group 1331 (RTG1331, to C.V.); the DFG Collaborative Research Center 969 (CRC969, to E.F.-M.); the Excellence Initiative of the German Federal and State Governments through funds of the Konstanz Research School Chemical Biology (to C.C.); the Evangelisches Studienwerk Villingen (to C.C.); DFG grant KA 2799/1 (to F.K); the START Program of the Faculty of Medicine, RWTH Aachen University (to F.K) and by a Wang-Chai Biochemistry grant from Duke Kunshan University (to F.K). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Data availability

R code used for DEK body analysis is provided at https://github.com/CP-Vogel/R-DEK-body-script.git.

References

Abbas, T. (2021). The Role of Ubiquitination and SUMOylation in DNA Replication. *Curr. Issues Mol. Biol.* **40**, 189–220.

Aichem, A., Sailer, C., Ryu, S., Catone, N., Stankovic-Valentin, N., Schmidtke, G., Melchior, F., Stengel, F. and Groettrup, M. (2019). The ubiquitin-like modifier FAT10 interferes with SUMO activation. *Nat. Commun.* **10**,.

Alabert, C., Bukowski-Wills, J. C., Lee, S. B., Kustatscher, G., Nakamura, K., De Lima Alves, F., Menard, P., Mejlvang, J., Rappsilber, J. and Groth, A. (2014). Nascent chromatin capture proteomics determines chromatin dynamics during DNA replication and identifies unknown fork components. *Nat. Cell Biol.* **16**, 281–291.

Aladjem, M. I. (2007). Replication in context: Dynamic regulation of DNA replication patterns in metazoans. *Nat. Rev. Genet.* **8**, 588–600.

Alexiadis, V., Waldmann, T., Andersen, J., Mann, M., Knippers, R. and Gruss, C. (2000). The protein encoded by the proto-oncogene DEK changes the topology of chromatin and reduces the efficiency of DNA replication in a chromatin- specific manner. *Genes Dev.* **14**, 1308–1312.

Aranda, S., Rutishauser, D. and Ernfors, P. (2014). Identification of a large protein network involved in epigenetic transmission in replicating DNA of embryonic stem cells. *Nucleic Acids Res.* **42**, 6972–6986.

Baltz, A. G., Munschauer, M., Schwanhäusser, B., Vasile, A., Murakawa, Y., Schueler, M., Youngs, N., Penfold-Brown, D., Drew, K., Milek, M., et al. (2012). The mRNA-bound proteome and its global occupancy profile on protein-coding transcripts. *Mol. Cell* **46**, 674–690.

Bates, M., Huang, B., Dempsey, G. T. and Zhuang, X. (2007). Multicolor super-resolution imaging with photo-switchable fluorescent probes. *Science* **317**, 1749–53.

Beckmann, B. M., Horos, R., Fischer, B., Castello, A., Eichelbaum, K., Alleaume, A.-M., Schwarzl, T., Curk, T., Foehr, S., Huber, W., et al. (2015). The RNA-binding proteomes from yeast to man harbour conserved enigmRBPs. *Nat. Commun.* **6**, 10127.

Bergoglio, V., Boyer, A. S., Walsh, E., Naim, V., Legube, G., Lee, M. Y. W. T., Rey, L., Rosselli, F., Cazaux, C., Eckert, K. A., et al. (2013). DNA synthesis by pol η promotes fragile site stability by preventing under-replicated DNA in mitosis. *J. Cell Biol.* **201**, 395–408.

Bhat, A., Andersen, P. L., Qin, Z. and Xiao, W. (2013). Rev3, the catalytic subunit of Polζ, is required for maintaining fragile site stability in human cells. *Nucleic Acids Res.* **41**, 2328–2339.

Böhm, F., Kappes, F., Scholten, I., Richter, N., Matsuo, H., Knippers, R. and Waldmann, T. (2005). The SAF-box domain of chromatin protein DEK. *Nucleic Acids Res.* **33**, 1101–1110.

Boyer, A. S., Walter, D. and Sørensen, C. S. (2016). DNA replication and cancer: From dysfunctional replication origin activities to therapeutic opportunities. *Semin. Cancer Biol.* **37**–**38**, 16–25.

Capitano, M. L., Markovitz, D. M., Hal, E., Invest, J. C., Capitano, M. L., Mor-vaknin, N., Saha, A. K., Cooper, S., Legendre, M., Guo, H., et al. (2019). Secreted nuclear protein DEK regulates hematopoiesis through CXCR2 signaling Find the latest version : Secreted nuclear protein DEK regulates hematopoiesis through CXCR2 signaling. **129**, 2555–2570.

Capitano, M. L., Sammour, Y., Ropa, J., Legendre, M., Mor-Vaknin, N. and Markovitz, D. M. (2022). DEK, a nuclear protein, is chemotactic for hematopoietic stem/progenitor cells acting through CXCR2 and Gαi signaling. *J. Leukoc. Biol.* **112**, 449–456.

Casas-Delucchi, C. S., Van Bemmel, J. G., Haase, S., Herce, H. D., Nowak, D., Meilinger, D., Stear, J. H., Leonhardt, H. and Cardoso, M. C. (2012). Histone hypoacetylation is required

to maintain late replication timing of constitutive heterochromatin. *Nucleic Acids Res.* **40**, 159–169.

Castello, A., Fischer, B., Eichelbaum, K., Horos, R., Beckmann, B. M., Strein, C., Davey, N. E., Humphreys, D. T., Preiss, T., Steinmetz, L. M., et al. (2012). Insights into RNA biology from an atlas of mammalian mRNA-binding proteins. *Cell* **149**, 1393–1406.

Cavellán, E., Asp, P., Percipalle, P. and Farrants, A. K. Ö. (2006). The WSTF-SNF2h chromatin remodeling complex interacts with several nuclear proteins in transcription. *J. Biol. Chem.* **281**, 16264–16271.

Cerase, A., Pintacuda, G., Tattermusch, A. and Avner, P. (2015). Xist localization and function: New insights from multiple levels. *Genome Biol.* **16**, 1–12.

Chadwick, B. P. and Willard, H. F. (2004). Multiple spatially distinct types of facultative heterochromatin on the human inactive X chromosome. *Proc. Natl. Acad. Sci. U. S. A.* **101**, 17450–17455.

Chagin, V. O., Casas-Delucchi, C. S., Reinhart, M., Schermelleh, L., Markaki, Y., Maiser, a, Bolius, J. J., Bensimon, a, Fillies, M., Domaing, P., et al. (2016). 4D Visualization of replication foci in mammalian cells corresponding to individual replicons. *Nat. Commun.* **7**, 11231.

Cleary, J., Sitwala, K. V., Khodadoust, M. S., Kwok, R. P. S., Mor-Vaknin, N., Cebrat, M., Cole, P. A. and Markovitz, D. M. (2005). p300/CBP-associated factor drives DEK into interchromatin granule clusters. *J. Biol. Chem.* **280**, 31760–31767.

Despras, E., Sittewelle, M., Pouvelle, C., Delrieu, N., Cordonnier, A. M. and Kannouche, P. L. (2016). Rad18-dependent SUMOylation of human specialized DNA polymerase eta is required to prevent under-replicated DNA. *Nat. Commun.* **7**, 13326.

Deutzmann, A., Ganz, M., Schönenberger, F., Vervoorts, J., Kappes, F. and Ferrando-May, E. (2015). The human oncoprotein and chromatin architectural factor DEK counteracts DNA replication stress. *Oncogene* **34**, 4270–4277.

Diaspro, A. and Bianchini, P. (2020). Optical nanoscopy. *Riv. del Nuovo Cim.* **43**, 385–455.

Dundr, M. (2012). Nuclear bodies: multifunctional companions of the genome. *Curr. Opin. Cell Biol.* **24**, 415–422.

Fahrer, J., Popp, O., Malanga, M., Beneke, S., Markovitz, D. M., Ferrando-May, E., Bürkle, A. and Kappes, F. (2010). High-affinity interaction of poly(ADP-ribose) and the human DEK oncoprotein depends upon chain length. *Biochemistry* **49**, 7119–7130.

Fragkos, M., Ganier, O., Coulombe, P. and Méchali, M. (2015). DNA replication origin activation in space and time. *Nat. Rev. Mol. Cell Biol.* **16**, 360–374.

Gamble, M. J. and Fisher, R. P. (2007). SET and PARP1 remove DEK from chromatin to permit access by the transcription machinery. *Nat. Struct. Mol. Biol.* **14**, 548–555.

Ganz, M., Vogel, C., Czada, C., Jörke, V., Gwosch, E. C., Kleiner, R., Pierzynska-Mach, A., Cella Zanacchi, F., Diaspro, A., Kappes, F., et al. (2019). The oncoprotein DEK affects the outcome of PARP1/2 inhibition during mild replication stress. *PLoS One* **14**, e0213130.

Garcia, P. A. A., Hoover, M. E., Zhang, P., Nagarajan, P., Freitas, M. A. and Parthun, M. R. (2017). Identification of multiple roles for histone acetyltransferase 1 in replication-coupled chromatin assembly. *Nucleic Acids Res.* **45**, 9319–9335.

Grant, M. M. (2010). Identification of SUMOylated proteins in neuroblastoma cells after treatment with hydrogen peroxide or ascorbate. *BMB Rep.* **43**, 720–725.

Guérillon, C., Smedegaard, S., Hendriks, I. A., Nielsen, M. L. and Mailand, N. (2020). Multisite SUMOylation restrains DNA polymerase η interactions with DNA damage sites. *J. Biol. Chem.* **295**, 8350–8362.

Guo, H., Prell, M., Königs, H., Xu, N., Waldmann, T., Hermans-Sachweh, B., Ferrando-May, E., Lüscher, B. and Kappes, F. (2021a). Bacterial Growth Inhibition Screen (BGIS) identifies a loss-of-function mutant of the DEK oncogene, indicating DNA modulating activities of DEK in chromatin. *FEBS Lett.* **595**, 1438–1453.

Guo, H., Xu, N., Prell, M., Königs, H., Hermanns-Sachweh, B., Lüscher, B. and Kappes, F. (2021b). Bacterial Growth Inhibition Screen (BGIS): harnessing recombinant protein toxicity for rapid and unbiased interrogation of protein function. *FEBS Lett.* **595**, 1422–1437.

Heinz, K. S., Casas-Delucchi, C. S., Török, T., Cmarko, D., Rapp, A., Raska, I. and Cardoso, M. C. (2018). Peripheral re-localization of constitutive heterochromatin advances its replication timing and impairs maintenance of silencing marks. *Nucleic Acids Res.* **46**, 6112–6128.

Hollenbach, A. D., McPherson, C. J., Mientjes, E. J., Iyengar, R. and Grosveld, G. (2002). Daxx and histone deacetylase II associate with chromatin through an interaction with core histones and the chromatin-associated protein Dek. *J. Cell Sci.* **115**, 3319–3330.

Hu, H. G., Illges, H., Gruss, C. and Knippers, R. (2005). Distribution of the chromatin protein DEK distinguishes active and inactive CD21/CR2 gene in pre- and mature B lymphocytes. *Int. Immunol.* **17**, 789–796.

Hu, H. G., Scholten, I., Gruss, C. and Knippers, R. (2007). The distribution of the DEK protein in mammalian chromatin. *Biochem. Biophys. Res. Commun.* **358**, 1008–1014.

Ivanauskiene, K., Delbarre, E., McGhie, J. D., Kuntziger, T., Wong, L. H. and Collas, P. (2014). The PML-associated protein DEK regulates the balance of H3.3 loading on chromatin and is important for telomere integrity. *Genome Res.* **24**, 1584–1594.

Jackson, D. A. and Pombo, A. (1998). Replicon clusters are stable units of chromosome structure: Evidence that nuclear organization contributes to the efficient activation and propagation of S phase in human cells. *J. Cell Biol.* **140**, 1285–1295.

Kamentsky, L., Jones, T. R., Fraser, A., Bray, M. A., Logan, D. J., Madden, K. L., Ljosa, V., Rueden, C., Eliceiri, K. W. and Carpenter, A. E. (2011). Improved structure, function and compatibility for cellprofiler: Modular high-throughput image analysis software. *Bioinformatics* **27**, 1179–1180.

Kappes, F., Burger, K., Baack, M., Fackelmayer, F. O. and Gruss, C. (2001). Subcellular Localization of the Human Proto-oncogene Protein DEK. *J. Biol. Chem.* **276**, 26317–26323.

Kappes, F., Damoc, C., Knippers, R., Przybylski, M., Pinna, L. A. and Gruss, C. (2004a). Phosphorylation by Protein Kinase CK2 Changes the DNA Binding Properties of the Human Chromatin Protein DEK. *Mol. Cell. Biol.* **24**, 6011–6020.

Kappes, F., Scholten, I., Richter, N., Gruss, C. and Waldmann, T. (2004b). Functional Domains of the Ubiquitous Chromatin Protein DEK. *Mol. Cell. Biol.* **24**, 6000–6010.

Kappes, F., Fahrer, J., Khodadoust, M. S., Tabbert, A., Strasser, C., Mor-Vaknin, N., Moreno-Villanueva, M., Burkle, A., Markovitz, D. M. and Ferrando-May, E. (2008). DEK Is a Poly(ADP-Ribose) Acceptor in Apoptosis and Mediates Resistance to Genotoxic Stress. *Mol. Cell. Biol.* **28**, 3245–3257.

Kappes, F., Waldmann, T., Mathew, V., Yu, J., Zhang, L., Khodadoust, M. S., Chinnaiyan, A. M., Luger, K., Erhardt, S., Schneider, R., et al. (2011a). The DEK oncoprotein is a Su(var) that is essential to heterochromatin integrity. *Genes Dev.* **25**, 673–678.

Kappes, F., Khodadoust, M. S., Yu, L., Kim, D. S. L., Fullen, D. R., Markovitz, D. M. and Ma, L. (2011b). DEK expression in melanocytic lesions. *Hum. Pathol.* **42**, 932–938.

Karam, M., Thenoz, M., Capraro, V., Robin, J. P., Pinatel, C., Lançon, A., Galia, P., Sibon, D., Thomas, X., Ducastelle-Lepretre, S., et al. (2014). Chromatin redistribution of the DEK oncoprotein represses hTERT transcription in leukemias. *Neoplasia (United States)* **16**, 21–30.

Lossaint, G., Larroque, M., Ribeyre, C., Bec, N., Larroque, C., Décaillet, C., Gari, K. and Constantinou, A. (2013). FANCD2 Binds MCM Proteins and Controls Replisome Function upon Activation of S Phase Checkpoint Signaling. *Mol. Cell* **51**, 678–690.

Lukusa, T. and Fryns, J. P. (2008). Human chromosome fragility. *Biochim. Biophys. Acta - Gene Regul. Mech.* **1779**, 3–16.

Mailand, N., Gibbs-Seymour, I. and Bekker-Jensen, S. (2013). Regulation of PCNA-protein

interactions for genome stability. *Nat. Rev. Mol. Cell Biol.* **14**, 269–282.

Matrka, M. C., Hennigan, R. F., Kappes, F., Delay, M. L., Lambert, P. F., Aronow, B. J. and Wells, S. I. (2015). DEK over-expression promotes mitotic defects and micronucleus formation. *Cell Cycle* **14**, 3939–3953.

Maya-Mendoza, A., Moudry, P., Merchut-Maya, J. M., Lee, M., Strauss, R. and Bartek, J. (2018). High speed of fork progression induces DNA replication stress and genomic instability. *Nature* **559**, 279–284.

Mintz, P. J., Patterson, S. D., Neuwald, A. F., Spahr, C. S. and Spector, D. L. (1999). Purification and biochemical characterization of interchromatin granule clusters. *EMBO J.* **18**, 4308–4320.

Mirkin, E. V. and Mirkin, S. M. (2007). Replication Fork Stalling at Natural Impediments. *Microbiol. Mol. Biol. Rev.* **71**, 13–35.

Mor-Vaknin, N., Kappes, F., Dick, A. E., Legendre, M., Damoc, C., Teitz-Tennenbaum, S., Kwok, R., Ferrando-May, E., Adams, B. S. and Markovitz, D. M. (2011). DEK in the synovium of patients with juvenile idiopathic arthritis: characterization of DEK antibodies and posttranslational modification of the DEK autoantigen. *Arthritis Rheum.* **63**, 556–567.

Nakayama, J., Rice, J. C., Strahl, B. D., Allis, C. D. and Grewal, S. I. S. (2001). Role of Histone H3 Lysine 9 Methylation in Epigenetic Control of Heterochromatin Assembly. *Science (80-.).* **292**, 110–113.

Neumann, B., Held, M., Liebel, U., Erfle, H., Rogers, P., Pepperkok, R. and Ellenberg, J. (2006). High-throughput RNAi screening by time-lapse imaging of live human cells. *Nat. Methods* **3**, 385–390.

O'Keefe, R. T., Henderson, S. C. and Spector, D. L. (1992). Dynamic organization of DNA replication in mammalian cell nuclei: Spatially and temporally defined replication of chromosome-specific α- satellite DNA sequences. *J. Cell Biol.* **116**, 1095–1110.

Özçelik, E., Kalaycı, A., Çelik, B., Avcı, A., Akyol, H., Kılıç, İ. B., Güzel, T., Çetin, M., Öztürk, M. T., Çalışkaner, Z. O., et al. (2022). Doxorubicin induces prolonged DNA damage signal in cells overexpressing DEK isoform-2. *PLoS One* **17**, e0275476.

Penagos-Puig, A. and Furlan-Magaril, M. (2020). Heterochromatin as an Important Driver of Genome Organization. *Front. Cell Dev. Biol.* **8**, 1–10.

Privette Vinnedge, L. M., Kappes, F., Nassar, N. and Wells, S. I. (2013). Stacking the DEK: From chromatin topology to cancer stem cells. *Cell Cycle* **12**, 51–66.

Ribeyre, C., Zellweger, R., Chauvin, M., Bec, N., Larroque, C., Lopes, M. and Constantinou, A. (2016). Nascent DNA Proteomics Reveals a Chromatin Remodeler Required for Topoisomerase I Loading at Replication Forks. *Cell Rep.* **15**, 300–309.

Ryba, T., Hiratani, I., Lu, J., Itoh, M., Kulik, M., Zhang, J., Schulz, T. C., Robins, A. J., Dalton, S. and Gilbert, D. M. (2010). Evolutionarily conserved replication timing profiles predict long-range chromatin interactions and distinguish closely related cell types. *Genome Res.* **20**, 761–770.

Saha, A. K., Kappes, F., Mundade, A., Deutzmann, A., Rosmarin, D. M., Legendre, M., Chatain, N., Al-Obaidi, Z., Adams, B. S., Ploegh, H. L., et al. (2013). Intercellular trafficking of the nuclear oncoprotein DEK. *Proc. Natl. Acad. Sci. U. S. A.* **110**, 6847–52.

Sandén, C. and Gullberg, U. (2015). The DEK oncoprotein and its emerging roles in gene regulation. *Leukemia* **29**, 1632–1636.

Sandén, C., Järvstråt, L., Lennartsson, A., Brattås, P. L., Nilsson, B. and Gullberg, U. (2014). The DEK oncoprotein binds to highly and ubiquitously expressed genes with a dual role in their transcriptional regulation. *Mol. Cancer* **13**, 1–13.

Sawatsubashi, S., Murata, T., Lim, J., Fujiki, R., Ito, S., Suzuki, E., Tanabe, M., Zhao, Y., Kimura, S., Fujiyama, S., et al. (2010). A histone chaperone, DEK, transcriptionally coactivates a nuclear receptor. *Genes Dev.* **24**, 159–170.

Saxena, S. and Zou, L. (2022). Hallmarks of DNA replication stress. *Mol. Cell* **82**, 2298–2314.

Sehwaiger, M., Stadler, M. B., Bell, O., Kohler, H., Oakeley, E. J. and Sehübeler, D. (2009). Chromatin state marks cell-type- and gender-specific replication of the Drosophila genome. *Genes Dev.* **23**, 589–601.

Sirbu, B. M., McDonald, W. H., Dungrawala, H., Badu-Nkansah, A., Kavanaugh, G. M., Chen, Y., Tabb, D. L. and Cortez, D. (2013). Identification of proteins at active, stalled, and collapsed replication forks using isolation of proteins on nascent DNA (iPOND) coupled with mass spectrometry. *J. Biol. Chem.* **288**, 31458–31467.

Smith, E. A., Krumpelbeck, E. F., Jegga, A. G., Prell, M., Matrka, M. M., Kappes, F., Greis, K. D., Ali, A. M., Meetei, A. R. and Wells, S. I. (2018). The nuclear DEK interactome supports multi-functionality "The DEK Interactome." *Proteins Struct. Funct. Bioinforma.* **86**, 88–97.

Stepanenko, O. V, Sulatsky, M. I., Mikhailova, E. V, Kuznetsova, I. M., Turoverov, K. K., Stepanenko, O. V and Sulatskaya, A. I. (2021). New findings on GFP-like protein application as fluorescent tags: Fibrillogenesis, oligomerization, and amorphous aggregation. *Int. J. Biol. Macromol.* **192**, 1304–1310.

Stewart-Morgan, K. R., Petryk, N. and Groth, A. (2020). Chromatin replication and epigenetic cell memory. *Nat. Cell Biol.* **22**, 361–371.

Su, Q. P., Zhao, Z. W., Meng, L., Ding, M., Zhang, W., Li, Y., Liu, M., Li, R., Gao, Y. Q., Xie, X. S., et al. (2020). Superresolution imaging reveals spatiotemporal propagation of human replication foci mediated by CTCF-organized chromatin structures. *Proc. Natl. Acad. Sci. U. S. A.* **117**, 15036–15046.

Tabbert, A., Kappes, F., Knippers, R., Kellermann, J., Lottspeich, F. and Ferrando-May, E. (2006). Hypophosphorylation of the architectural chromatin protein DEK in death-receptor-induced apoptosis revealed by the isotope coded protein label proteomic platform. *Proteomics* **6**, 5758–5772.

Tatham, M. H., Rodriguez, M. S., Xirodimas, D. P. and Hay, R. T. (2009). Detection of protein SUMOylation in vivo. *Nat. Protoc.* **4**, 1363–1371.

Trojer, P. and Reinberg, D. (2007). Facultative Heterochromatin: Is There a Distinctive Molecular Signature? *Mol. Cell* **28**, 1–13.

Tsao, W.-C. and Eckert, K. A. (2018). Detours to Replication: Functions of Specialized DNA Polymerases during Oncogene-induced Replication Stress. *Int. J. Mol. Sci.* **19**,.

Vesela, E., Chroma, K., Turi, Z. and Mistrik, M. (2017). Common Chemical Inductors of Replication Stress: Focus on Cell-Based Studies. *Biomolecules* **7**, 19.

Waidmann, S., Kusenda, B., Mayerhofer, J., Mechtler, K. and Jonak, C. (2014). A DEK domain-containing protein modulates chromatin structure and function in Arabidopsis. *Plant Cell* **26**, 4328–4344.

Waldmann, T., Eckerich, C., Baack, M. and Gruss, C. (2002). The ubiquitous chromatin protein DEK alters the structure of DNA by introducing positive supercoils. *J. Biol. Chem.* **277**, 24988–24994.

Waldmann, T., Baack, M., Richter, N. and Gruss, C. (2003). Structure-specific binding of the proto-oncogene protein DEK to DNA. *Nucleic Acids Res.* **31**, 7003–7010.

Waldmann, T., Scholten, I., Kappes, F., Hu, H. G. and Knippers, R. (2004). The DEK protein-an abundant and ubiquitous constituent of mammalian chromatin. *Gene* **343**, 1–9.

Wang, Z., Zang, C., Rosenfeld, J. A., Schones, D. E., Barski, A., Cuddapah, S., Cui, K., Roh, T. Y., Peng, W., Zhang, M. Q., et al. (2008). Combinatorial patterns of histone acetylations and methylations in the human genome. *Nat. Genet.* **40**, 897–903.

Wessel, D. and Flügge, U. I. (1984). A method for the quantitative recovery of protein in dilute solution in the presence of detergents and lipids. *Anal. Biochem.* **138**, 141–143.

Yang, Y.-S., Jia, X.-Z., Lu, Q.-Y., Cai, S.-L., Huang, X.-T., Yang, S.-H., Wood, C., Wang, Y.-H., Zhou, J.-J., Chen, Y.-D., et al. (2022). Exosomal DEK removes chemoradiotherapy resistance by triggering quiescence exit of breast cancer stem cells. *Oncogene* **41**, 2624–2637.

Zhang, X. D. (2011). Illustration of SSMD, z score, SSMD*, z* score, and t statistic for hit selection in RNAi high-throughput screens. *J. Biomol. Screen.* **16**, 775–785.

Zhang, Q., Bassetti, F., Gherardi, M. and Lagomarsino, M. C. (2017). Cell-to-cell variability and robustness in S-phase duration from genome replication kinetics. *Nucleic Acids Res.* **45**, 8190–8198.

Zhao, Q., Xie, Y., Zheng, Y., Jiang, S., Liu, W., Mu, W., Liu, Z., Zhao, Y., Xue, Y. and Ren, J. (2014). GPS-SUMO: A tool for the prediction of sumoylation sites and SUMO-interaction motifs. *Nucleic Acids Res.* **42**, 325–330.

Zheng, L., Baumann, U. and Reymond, J. L. (2004). An efficient one-step site-directed and site-saturation mutagenesis protocol. *Nucleic Acids Res.* **32**,.

Zhou, J., Zhao, L., Wu, Y., Zhang, X., Cheng, S., Wei, F., Zhang, Y., Zhu, H., Zhou, Y., Feng, Z., et al. (2022). A DEK domain-containing protein GhDEK2D mediated Gossypium hirsutum enhanced resistance to Verticillium dahliae. *Plant Signal. Behav.* **17**,.

FIGURES

Figure 1: DEK forms distinct foci, DEK bodies, which colocalize with nascent DNA markers.

A. Widefield images of MCF10A cells. Nascent DNA was pulse-labeled with EdU for 25 min followed by detection via click chemistry and immunofluorescence staining using

DEK-specific antibodies. Average number of DEK foci per cell: 6.7 ± 2.2 (n = 46 cells). Magenta: EdU. Green: DEK. Scale bar: 5 µm.

B. Immunofluorescence analysis of DEK and EdU distribution pattern in late S-phase MCF10A cells as described in (A) (maximum projection). Arrows point to sites of DEK and EdU colocalization. Scale bar: 5 µm.

C. Immunofluorescence and colocalization analysis of DEK (green) and PCNA (magenta) in a late S-phase MCF10A cell nucleus. Dashed boxes indicate two selected ROIs. The intensity profiles are measured along the lines shown in the enlarged ROIs. The fluorescence intensity was normalized to 1. Scale bars: 5 µm.

D. Widefield fluorescence images of immunolabelled U2-OS wild type cells (upper row) and U2-OS KI eGFP-DEK cells (bottom row). Endogenous DEK (green) and PCNA (magenta) were visualized by specific antibodies. In each image one prominent DEK body is marked by a white arrow. Scale bar: 5 µm.

Figure 2: DEK bodies localize in the vicinity of the replication fork

A. Confocal and STED super-resolution images of an MCF10A cell nucleus in late S-phase labeled with anti-DEK antibodies. The STED image reveals a granular internal structure of DEK bodies (insets). Scale bar: 5 µm.

B. dSTORM images of an MCF10A cell nucleus in late S-phase labeled with anti-DEK antibodies. Left: widefield conventional image of DEK immunolabelling. Middle: dSTORM image reconstructed from single-molecule localization microscopy

acquisition. Scale bar: 2 µm. Right: magnified image of the ROI shown in the middle containing a DEK body. Scale bar: 0.5 µm.

C. Pseudo-widefield representation of a representative U2-OS nucleus in late S-phase imaged in 3D-SIM mode. U2-OS cells were pulse-labelled with EdU for 10 min. DEK (green) and PCNA (magenta) were visualized *via* indirect immunofluorescence microscopy and EdU (cyan) using click chemistry.

D. Single Z-slice from a middle section of the super-resolved image stack after reconstruction and registration. Scale bar: 5 µm. The magnified inset shows one DEK-positive large replication focus. Scale bar: 0.5 µm.

E. Fluorescence intensity profiles of DEK, PCNA and EdU. Interpolated intensity profiles were calculated along the line shown in (D) and normalized to the min/max values.

Figure 3: DEK bodies position within constitutive heterochromatin regions in late S-phase.

A. Confocal images of late S-phase MCF10A cell nuclei labelled with EdU and antibodies specific for DEK and for three post-translational histone modifications (H3K9ac - left, H3K27me3 - centre, H3K9me3 - right). ROIs containing DEK bodies are marked by a dashed line and displayed as magnified images below. The fluorescence intensity profiles were calculated along the lines delimited by the white arrowheads and normalized to 1. Scale bars: 5 µm.

B. Analysis of colocalization of DEK bodies and chromatin marks using Manders' coefficients. In the box plots, the red line indicates the average value, the asterisk

marks the median, the box boundaries represent the interquartile range, and the whiskers above and below the box indicate minimum and maximum values. Graphs on the bottom schematically represent localization pattern.

C. Confocal image of a late S-phase MCF10A cell nucleus after PLA with antibodies specific for DEK and H3K9me3. The PLA positive signal (magenta) indicates direct interaction. Scale bar: 5 µm.

D. Confocal image of a late S-phase MCF10A cell nucleus after PLA with antibodies specific for DEK and H3K27me3. No PLA positive signal is detected. Scale bar: 5 µm.

E. FRAP analysis of DEK-eGFP mobility transiently expressed in MCF10A cells in bulk chromatin and in DEK bodies. Bleach-ROIs (2 µm x 2 µm) are indicated by dashed squares. Fluorescence recovery was monitored for 3 min. Scale bar: 5 µm.

F. DEK displays a lower mobility in bodies than in bulk chromatin. Normalized FRAP recovery curves based on the data from (E). Data points show the average of 6 experiments for DEK bodies and 14 experiments for DEK in chromatin. Error bars show the SEM.

G. Halftimes of recovery for eGFP-DEK in chromatin ($\tau\frac{1}{2}$ = 10.6 s ± 3.1 s), and in DEK bodies ($\tau\frac{1}{2}$ = 17.1 s ± 7 s) calculated from the FRAP data in (F).

Figure 4: DEK bodies are spatially associated with replicating heterochromatin.

A, B. Maximum intensity projection images of representative MCF10A cell nuclei in late S-phase (A), and in the absence of DNA replication (B, non-S-phase, EdU-negative) labelled with anti-DEK, anti-CENP-A antibodies, and EdU-AF467. ROIs are marked by a dashed line and displayed as magnified images below. The fluorescence intensity profiles were calculated along the lines delimited by the white arrowheads and normalized between 0 and 1. Scale bars: 5 µm.

C. Maximum intensity projection image of a representative MCF10A cell nucleus in late S-phase labelled with anti-DEK antibodies, a *Xist* RNA-specific FISH probe, and EdU-AF467. The ROI marked by a dashed line is displayed as magnified image below. The fluorescence intensity profile was calculated along the lines delimited by the white arrowheads and normalized between 0 and 1. Scale bars: 5 µm.

D, E. Confocal images of representative MCF10A cell nuclei in late S-phase (D) and in the absence of replication (E, non-S-phase, EdU-negative) labelled with anti-DEK and

anti-Lamin A antibodies, and EdU-AF467. Arrows in (D) indicate the juxtaposition of DEK and Lamin A at the periphery of the cell nucleus and at nuclear lamina invaginations, which is not observed in non-S-phase cells (E). Scale bar: 5 µm

F. Confocal image of a representative MCF10A cell nucleus in late S-phase labeled with DEK-specific antibodies, with the DNA dye ToPRo3 and EdU-AF647. The ROI marked by a dashed line is displayed as magnified image below and shows a DEK body overlapping with a region of dense chromatin. The fluorescence intensity profile was calculated along the lines delimited by the white arrowheads and normalized between 0 and 1. Scale bars: 5 µm.

Figure 5: DEK bodies form during late S-phase of the cell cycle.

A. Time-lapse fluorescence microscopy of U2-OS KI eGFP-DEK cells transiently expressing mRFP-PCNA. Images were acquired at a spinning disk confocal microscope. The figure shows a representative image sequence. Magenta: PCNA chromobody, green: eGFP-DEK. Scale bar: 5 μm.

B. Mean durations of cell cycle phases and mean DEK body lifetime. A total of 56 cells were scored manually from two independent experiments. Mid S, late S, G2 and M-phases were identified according to the PCNA localization pattern. Only images with clearly detectable DEK bodies were scored as positive. DEK bodies appear reproducibly in late S-phase (continuous green line), and sporadically also in mid-S-phase (dashed green line).

C. Quantification of PCNA body and DEK body numbers in late S-phase using the same data set as in (B). Only PCNA or DEK bodies were included in the evaluation that persisted for at least two consecutive frames. The violin plot shows the density distribution of the data, the black line indicates the median, the box the interquartile range (IQR) and the whiskers the IQR ± 1.5.

Figure 6: Formation of DEK bodies is dependent on DNA replication.

A. Time-lapse fluorescence microscopy of U2-OS KI eGFP-DEK cells treated for 1 h with either 10 μM HU, or 200 nM APH, or 100 nM CPT or left untreated (Ctrl). Images were acquired for 24 h with a spinning disk confocal microscope. A representative time series is shown for each treatment condition. Frames with DEK bodies are highlighted by a red outline. Scale bar: 10 μm.

B. Quantification of G2-phase and mitosis duration, DEK body number and lifetime in time series data as in (A) (15-132 cells, single experiment). The violin plots show the density distribution of the data points, the black line indicates the median, the box the interquartile range (IQR) and the whiskers the IQR ± 1.5. (*) The control and CPT data sets contained one outlier each, which was omitted from the quantification.

Figure 7: siRNA screen identifies positive and negative regulators of DEK body formation.

A. Plot of the log2-transformed ranked mean Z-score data of the primary screen. The screen was performed on 4 field of view per well, with two wells per siRNA (for each

2 replicates), yielding 16 Z-scores for each siRNA. Dark grey dots: Top scorer (3 ≤ Z-score ≤ -2). Blue dots: Target genes included in the pilot screen. sNEG1: negative control siRNA.

B. List of the strongest DEK body upregulating and downregulating genes based on the Z-score of the corresponding siRNAs. For the up-regulators, ranks 30-34 and 36-43 are not shown for better readability. Mean number of DEK bodies (DB) and standard deviation (SD) are shown. Highlighted are genes belonging to the SUMO pathway.

C. Representative widefield images from the primary screen showing U2-OS KI eGFP-DEK transfected with a negative control siRNA sNEG1 (s813) and siRNAs targeting the two strongest DEK body upregulating (left) and downregulating (right) genes. Magnified insets with DEK body-positive nuclei are shown. Scale bars: 100 μm; inset scale bars: 10. μm

D. Comparison of z-score data from the primary and the secondary (validation) screen. The log2-transformed mean Z-score for the indicated siRNAs is shown. The primary screen was performed in duplicate, the validation screen in triplicate. Knockdown efficiency of SUMO1 and SUMO 3 siRNAs was confirmed by Western Blot (Fig. S7)

Figure 8: DEK is SUMOylated in cells and SUMOylation is involved in the formation of DEK bodies.

A. Immunoblot analysis of co-immunoprecipitations from U2-OS whole cell lysates using DEK-, and SUMO-specific antibodies.

B. Immunoblot analysis of His-SUMO pulldown experiments from HeLa cells expressing 6xHis-SUMO-1, 6xHis-SUMO-2, 6xHis-SUMO-3, or control cells after nickel affinity chromatography. Purified proteins were analyzed with DEK and SUMO-specific antibodies.

C. Densitometric quantification of mono-SUMOylated DEK signals from (B, left panel). DEK signals in pulldown samples were normalized to the corresponding input samples. Error bars indicate the SD. **p<0.01, *p<0.05.

D. *In vitro* SUMOylation assay: Immunoblot analysis of *in vitro* SUMOylation reactions containing recombinant GST-tagged DEK WT, mutant, or GST only proteins as indicated and using DEK-specific antibodies for detection.

E. Densitometric quantification of mono-SUMOylated DEK from *in vitro* samples as in (D). Error bars indicate the SD.

F. Widefield microscopy images of U2-OS KI eGFP-DEK or a cell line stably expressing a wild-type or a SUMOylation deficient DEK mutant - DEKmut (Δ61,62,144,145,261) cells. Scale bars: 20 μm.

SUPPLEMENTAL INFORMATION

DEK oncoprotein participates in heterochromatin replication via SUMO-dependent nuclear bodies

Agnieszka Pierzynska-Mach[1, #], Christina Czada[2, #], Christopher Vogel[2], Eva Gwosch[2], Xenia Osswald[2], Denis Bartoschek[2], Alberto Diaspro[1,3], Ferdinand Kappes[4, #, *] and Elisa Ferrando-May[2,5, #, *]

Supplementary Figures S1-S8

Supplementary Table S1-4

Supplementary Materials and Methods

Supplementary Figure S1: Endogenous DEK forms DEK foci in primary BJ5-ta cells, yet not in malignant MCF7 and MDA-MB-231 cells.

A. Immunofluorescence analysis of DEK (green) and PCNA (magenta) distribution in a BJ5-TA fibroblast nucleus by super-resolution microscopy (3D-SIM) *via* indirect immunofluorescence. Pseudo-widefield representation of one slice.

B. Image of a single Z-slice from a middle section of the super-resolved image stack after reconstruction and registration. Scale bar: 5 µm. The magnified inset shows one DEK-positive large replication focus. Scale bar: 0.5 µm.

C. Fluorescence intensity profiles of DEK and PCNA. Interpolated intensity profiles were calculated along the line indicated in (B) and normalized to the min/max values.

D, E Occurrence of DEK bodies is reduced in (D) MCF7 and (E) MDA-MB-231 cells. Immunofluorescence analysis of DEK and EdU distribution pattern in late S-phase cells was performed as described in Fig. 1. Scale bar: 5 µm.

Supplementary Figure S2: Colocalization of eGFP-DEK and PCNA during late S-phase in U2-OS cells carrying a TALEN-mediated eGFP-DEK genomic knock in.

A. Representative confocal images of U2-OS KI eGFP-DEK cells in different cell cycle phases labeled with PCNA-specific antibodies. Similar focal accumulations of eGFP-DEK (green) and PCNA (magenta) can be observed in late S-phase. DNA was counterstained with Hoechst33342 (cyan). Scale bar: 10 μm.

B. Colocalization analysis of PCNA and DEK bodies in U2-OS KI eGFP-DEK cells. Scale bar: 10 μm. Representative images are shown, and intensity profile was analyzed along the white line displayed in the right panel.

C. DEK body formation is independent of the DEK expression level. U2-OS wild type cells were transfected with plasmids either coding for pEGFP-N1-hDEK or pENmRFP-PCNAL2 or double transfected with both plasmids and incubated for 24 h for maximal protein expression. U2-OS KI eGFP-DEK cells were transfected with pEN-mRFP-PCNAL2 only. Cells were monitored for 24 h (37°C, 5% CO_2). Concomitant with DEK foci formation, relocation of RFP-PCNA (magenta) can be detected both in transiently transfected cells and in the stable cell line. Single transfected cells (upper two panels) serve as control cells. Scale bar: 10 µm.

D. DEK foci formation is not an eGFP artifact. U2-OS cells were double transfected with pEGFP-C1 and pENmRFP-PCNAL2 and incubated for 24 h. During RFP-PCNA body formation, an equally distributed eGFP signal can be visualized. Scale bar: 10 µm.

Supplementary Figure S3: Localization analysis of DEK and histone marks in S-phase (data set corresponding to Fig. 3A).
Confocal images of MCF10A cells labeled with antibodies specific for DEK and histone post-translational modifications H3K9ac (A), H3K27me3 (B), and H3K9me3 (C) during S-phase. Scale bars = 5 µm.

Supplementary Figure S4: DEK bodies co-localize with heterochromatin in U2-OS cells.

A, B. Representative confocal images of U2-OS KI eGFP-DEK labelled with H3K9me3 specific antibodies (magenta) (A) and of U2-OS WT cells (B) additionally labelled with antibodies specific for DEK (green). DNA was counterstained using Hochst33342 (cyan). Scale bar: 10 μm.

C, D. Confocal images of single U2-OS KI eGFP-DEK and U2-OS WT cells from (A) and (B) with the respective fluorescence intensity profiles for the analysis of H3K9me3 (magenta) and DEK (green) colocalization. The white line is given as distance in the graph.

Supplementary Figure S5: DEK bodies form during late S-phase of the cell cycle in transiently transfected MCF10A cells.

Time-lapse fluorescence imaging of MCF10A cells co-transfected with plasmids encoding eGFP-DEK and mRFP-PCNA. The images were acquired at a scanning confocal microscope. Frames were captured every 10 min for 12h. DEK bodies can be traced for about 189 ± 38 min (13 measurements from 4 independent samples).

Supplementary Figure S6: Design of a high-throughput siRNA screening approach for the identification of DEK body regulators.

A. Schematic of the experimental protocols used for the siRNA screens. The pilot and primary screens relied on automated procedures and solid-phase reverse transfection of the siRNAs. Imaging was performed with a high-content widefield microscope. For the validation screen, cells and transfection reagents were dispensed manually. Imaging was performed on a spinning disk confocal microscope with an automated stage. Two pilot screens with slightly different imaging conditions were performed to optimize imaging parameters. The primary screens were performed in duplicate, and the validation screen in triplicate.

B. Main steps of the image analysis pipeline. Example images of cells transfected with a negative control siRNA are shown. Nuclei with detected DEK bodies are outlined in red, nuclei that passed quality control in white. Nuclei that were rejected because of cells were dead or mitotic are not outlined. DEK bodies are encircled in red. Scale bar: 20 µm.

C. Scheme of the gap-filling algorithm used after image analysis. The algorithm removed frames in which detection of DEK bodies within one track failed (false negatives), and sorted out frames in which foci were detected in one frame only (false positives).

D. Quantification of DEK body parameters in image series from one of the pilot screens processed as in (A) and (C). Calculations were done using R. Violin plots display the mean number of DEK bodies per cell (left panel), and the length of the time windows in which DEK bodies appeared (DEK body lifetime, right panel) for the indicated treatments/siRNAs. In total, image data from 12 fields of view (4 positions per well, 3 wells per condition), each containing several tracks, were evaluated. Values from tracks within a field of view were averaged. The violin plot represents the density distribution of these averages, the black line indicates the median, the box the interquartile range (IQR) and the whiskers the IQR ± 1.5.

E. Representative exemplary images of U2-OS KI eGFP-DEK cells treated with sNEG1, sDEK and the DEK body up-regulating siRNA sUBE2I and the down-regulating siRNA sRPA1. Images were taken at a spinning disk confocal microscope. Scale bar: 10 µm.

Supplementary Figure S7: SUMO1/3 siRNAs are effective in down-regulating SUMO1/3 protein levels.

A. Western blot analysis of SUMO1 and SUMO2/3/4 protein levels after siRNA-mediated knockdown of gene expression. Whole cell lysates from siRNA-transfected U2OS KI eGFP-DEK cells were subjected to Western blotting using SUMO1- and SUMO2/3-specific antibodies, respectively. As negative control the sNEG1 siRNA was used, PCNA served as loading control. Quantified SUMO signals are marked with an arrow. The experiment was performed once.
B. Densitometric quantification of (A). SUMO-specific signals were normalized to PCNA and are plotted as percentage of their respective controls (PCNA).

Supplementary Figure S8: Mutation of individual DEK SUMOylation sites does not lead to reduced DEK SUMOylation in vitro.

Samples containing recombinant GST-tagged DEK WT, mutant, or GST only proteins as indicated were subjected to *in vitro* SUMOylation assays as described in Fig. 8. Proteins were detected with DEK-, GST-, and SUMO-specific antibodies. The black arrow indicates GST-DEK signals, red arrows the mono- and di-SUMOylated variants.

Supplementary Table 1. siRNA library of the primary screen

	siRNA ID	Gene ID	Full gene name		siRNA ID	Gene ID	Full gene name
1	s859	ABCF1	ATP binding cassette subfamily F member 1	342	s30287	MTPAP	mitochondrial poly(A) polymerase
2	s860			343	s30288		
3	s962	ACTL6A	actin like 6A	344	s223607	MTREX	Mtr4 exosome RNA helicase
4	s963			345	s23982		
5	s1008	ADAR	adenosine deaminase, RNA specific	346	s9129	MYC	MYC proto-oncogene, bHLH transcription factor
6	s1009			347	s9130		
7	s1445	APEX1	apurinic/apyrimidinic endodeoxyribonuclease 1	348	s9280	NASP	nuclear autoantigenic sperm protein
8	s1446			349	s9282		
9	s1447			350	s30492	NAT10	N-acetyltransferase 10
10	s47251	APLF	aprataxin and PNKP like factor	351	s30493		
11	s47253			352	s9291	NBN	nibrin
12	s15785	ARID1A	AT-rich interaction domain 1A	353	s9293		
13	s15786			354	s9312	NCL	nucleolin
14	s31344	ASF1B	anti-silencing function 1B histone chaperone	355	s9313		
15	s31345			356	s22929	NCOA6	nuclear receptor coactivator 6
16	s31346			357	s22930		
17	s1709	ATM	ATM serine/threonine kinase	358	s28072	NIP7	NIP7, nucleolar pre-rRNA processing protein
18	s57221			359	s28074		
19	s227305	ATR	ATR serine/threonine kinase	360	s26364	NOB1	NIN1 (RPN12) binding protein 1 homolog
20	s56826			361	s26365		
21	s57368	ATRX	ATRX, chromatin remodeler	362	s36145	NOL9	nucleolar protein 9
22	s59082			363	s36146		
23	s22059	BAZ1A	bromodomain adjacent to zinc finger domain 1A	364	s28205	NOP16	NOP16 nucleolar protein
24	s22060			365	s28206		
25	s17209	BAZ1B	bromodomain adjacent to zinc finger domain 1B	366	s9676	NPM1	nucleophosmin 1
26	s17210			367	s9677		
27	s1997	BLM	BLM RecQ like helicase	368	s20257	NPM3	nucleophosmin/nucleoplasmin 3
28	s1999			369	s20258		
29	s457	BRCA1	BRCA1, DNA repair associated	370	s29683	NSUN2	NOP2/Sun RNA methyltransferase family member 2
30	s458			371	s29684		
31	s2083	BRCA2	BRCA2, DNA repair associated	372	s9892	ORC1	origin recognition complex subunit 1
32	s224695			373	s9893		
33	s38384	BRIP1	BRCA1 interacting protein C-terminal helicase 1	374	s9895	ORC2	origin recognition complex subunit 2
34	s38385			375	s9896		
35	s226839	BRIX1	BRX1, biogenesis of ribosomes	376	s24167	ORC3	origin recognition complex subunit 3
36	s30678			377	s24168		
37	s2134	BYSL	bystin like	378	s9898	ORC4	origin recognition complex subunit 4
38	s2135			379	s9900		
39	s8377	CAPRIN1	cell cycle associated protein 1	380	s9901	ORC5	origin recognition complex subunit 5
40	s8378			381	s9903		
41	s21550	CBX1	chromobox 1	382	s24164	ORC6	origin recognition complex subunit 6
42	s21551			383	s24165		
43	s223227	CBX3	chromobox 3	384	s9978	PA2G4	proliferation-associated 2G4
44	s22356			385	s9980		
45	s23883	CBX5	chromobox 5	386	s25664	PABPC1	poly(A) binding protein cytoplasmic 1
46	s23884			387	s25665		
47	s225505	CCDC124	coiled-coil domain containing 124	388	s25666		
48	s41755			389	s16693	PABPC4	poly(A) binding protein cytoplasmic 4
49	s229	CCND1	cyclin D1	390	s52787		
50	s230			391	s29268	PAF1	PAF1 homolog, Paf1/RNA polymerase II complex component
51	s225066	CCNH	cyclin H	392	s29269		
52	s2537			393	s1097	PARP1	poly(ADP-ribose) polymerase 1
53	s15830	CDC45	cell division cycle 45	394	s1099		
54	s15831			395	s19502	PARP2	poly(ADP-ribose) polymerase 2
55	s463	CDK1	cyclin dependent kinase 1	396	s19504		
56	s464			397	s19505	PARP3	poly(ADP-ribose) polymerase family member 3
57	s204	CDK2	cyclin dependent kinase 2	398	s19506		
58	s205			399	s18861	PCLAF	PCNA clamp associated factor
59	s2829	CDK7	cyclin dependent kinase 7	400	s18863		
60	s2830			401	s10134	PCNA	proliferating cell nuclear antigen
61	s416	CDKN1A	cyclin dependent kinase inhibitor 1A	402	s10135		
62	s417			403	s25745	PELP1	proline, glutamate and leucine rich protein 1
63	s2838	CDKN1B	cyclin dependent kinase inhibitor 1B	404	s25746		
64	s2839			405	s38846	PHF6	PHD finger protein 6

#	ID	Gene	Description
65	s37723	CDT1	chromatin licensing and DNA replication factor 1
66	s37724		
67	s2909	CENPB	centromere protein B
68	s2911		
69	s225845	CENPX	centromere protein X
70	s47345		
71	s19499	CHAF1A	chromatin assembly factor 1 subunit A
72	s19501		
73	s15705	CHAF1B	chromatin assembly factor 1 subunit B
74	s15706		
75	s503	CHEK1	checkpoint kinase 1
76	s504		
77	s22120	CHEK2	checkpoint kinase 2
78	s22121		
79	s14676	COL14A1	collagen type XIV alpha 1 chain
80	s14677		
81	s3495	CREBBP	CREB binding protein
82	s3497		
83	s3639	CSNK2A2	casein kinase 2 alpha 2
84	s3641		
85	s3642	CSNK2B	casein kinase 2 beta
86	s3643		
87	s3644		
88	s37000	CTC1	CST telomere replication complex component 1
89	s37001		
90	s3935	DAXX	death domain associated protein
91	s3937		
92	s3979	DDB1	damage specific DNA binding protein 1
93	s3980		
94	s3982	DDB2	damage specific DNA binding protein 2
95	s3984		
96	s17564	DDX21	DExD-box helicase 21
97	s17565		
98	s32631	DDX24	DEAD-box helicase 24
99	s32632		
100	s224194	DDX47	DEAD-box helicase 47
101	s27672		
102	s35416	DDX50	DExD-box helicase 50
103	s35417		
104	s21782	DDX52	DExD-box helicase 52
105	s21784		
106	s35469	DDX54	DEAD-box helicase 54
107	s35470		
108	s33616	DDX55	DEAD-box helicase 55
109	s33617		
110	s15457	DEK	DEK proto-oncogene
111	s15458		
112	s15459		
113	s22643	DHX30	DExH-box helicase 30
114	s22645		
115	s33511	DHX37	DEAH-box helicase 37
116	s33512		
117	s21855	DIDO1	death inducer-obliterator 1
118	s225469		
119	s4173	DNA2	DNA replication helicase/nuclease 2
120	s4175		
121	s4215	DNMT1	DNA methyltransferase 1
122	s4216		
123	s226739	DTL	denticleless E3 ubiquitin protein ligase homolog
124	s28247		
125	s4390	DUT	deoxyuridine triphosphatase
126	s4391		
127	s44664	E2F7	E2F transcription factor 7
128	s44665		
129	s21591	EBNA1BP2	EBNA1 binding protein 2
130	s21592		
131	s4556	EIF2S1	eukaryotic translation

#	ID	Gene	Description
406	s38848		
407	s194691	PML	promyelocytic leukemia
408	s194692		
409	s22287	PNKP	polynucleotide kinase 3'-phosphatase
410	s22288		
411	s32352	PNO1	partner of NOB1 homolog
412	s32353		
413	s10772	POLA1	DNA polymerase alpha 1, catalytic subunit
414	s10774		
415	s24280	POLA2	DNA polymerase alpha 2, accessory subunit
416	s24282		
417	s10775	POLB	DNA polymerase beta
418	s10776		
419	s614	POLD1	DNA polymerase delta 1, catalytic subunit
420	s615		
421	s616		
422	s10778	POLD2	DNA polymerase delta 2, accessory subunit
423	s10779		
424	s21045	POLD3	DNA polymerase delta 3, accessory subunit
425	s21046		
426	s10781	POLE	DNA polymerase epsilon, catalytic subunit
427	s10783		
428	s10785	POLE2	DNA polymerase epsilon 2, accessory subunit
429	s10786		
430	s10787	POLG	DNA polymerase gamma, catalytic subunit
431	s10788		
432	s10789		
433	s22169	POLG2	DNA polymerase gamma 2, accessory subunit
434	s22171		
435	s10790	POLH	DNA polymerase eta
436	s10791		
437	s22122	POLI	DNA polymerase iota
438	s22123		
439	s28115	POLK	DNA polymerase kappa
440	s28116		
441	s26196	POLL	DNA polymerase lambda
442	s26197		
443	s51480	POLN	DNA polymerase nu
444	s51481		
445	s21059	POLQ	DNA polymerase theta
446	s21060		
447	s719	PPP1CC	protein phosphatase 1 catalytic subunit gamma
448	s721		
449	s17267	PRC1	protein regulator of cytokinesis 1
450	s17268		
451	s11050	PRIM1	DNA primase subunit 1
452	s11051		
453	s11053	PRIM2	DNA primase subunit 2
454	s11054		
455	s61393	PRKDC	protein kinase, DNA-activated, catalytic subunit
456	s775		
457	s16335	PRKRA	protein activator of interferon induced protein kinase EIF2AK2
458	s16336		
459	s22035	PSIP1	PC4 and SFRS1 interacting protein 1
460	s22036		
461	s19272	PUM3	pumilio RNA binding family member 3
462	s19274		
463	s38873	PYM1	PYM homolog 1, exon junction complex associated factor
464	s38874		
465	s38875		
466	s20341	RACK1	receptor for activated C kinase 1
467	s20342		
468	s11589	RAD1	RAD1 checkpoint DNA exonuclease
469	s11590		
470	s11723	RAD17	RAD17 checkpoint clamp loader component
471	s11724		
472	s32295	RAD18	RAD18, E3 ubiquitin protein ligase

132	s4557		initiation factor 2 subunit alpha
133	s17003	EIF2S2	eukaryotic translation
134	s17004		initiation factor 2 subunit beta
135	s16490	EIF3A	eukaryotic translation
136	s16491		initiation factor 3 subunit A
137	s7586	EIF6	eukaryotic translation
138	s7588		initiation factor 6
139	s20415	EMG1	EMG1, N1-specific
140	s20416		pseudouridine methyltransferase
141	s4695	EP300	E1A binding protein p300
142	s4696		
143	s4785	ERCC1	ERCC excision repair 1, endonuclease non-catalytic subunit
144	s4786		
145	s230766	ERCC2	ERCC excision repair 2, TFIIH core complex helicase subunit
146	s4787		
147	s4796	ERCC3	ERCC excision repair 3, TFIIH core complex helicase subunit
148	s4798		
149	s4799	ERCC4	ERCC excision repair 4, endonuclease catalytic subunit
150	s4801		
151	s4805	ERCC6	ERCC excision repair 6, chromatin remodeling factor
152	s4806		
153	s3104	ERCC8	ERCC excision repair 8, CSA ubiquitin ligase complex subunit
154	s3105		
155	s10738	EXOSC10	exosome component 10
156	s10739		
157	s23751	EXOSC2	exosome component 2
158	s23752		
159	s223551	EXOSC7	exosome component 7
160	s22837		
161	s22838		
162	s4913	EZH1	enhancer of zeste 1 polycomb repressive complex 2 subunit
163	s4914		
164	s4916	EZH2	enhancer of zeste 2 polycomb repressive complex 2 subunit
165	s4918		
166	s163	FANCA	FA complementation group A
167	s164		
168	s4985	FANCC	FA complementation group C
169	s4987		
170	s4988	FANCD2	FA complementation group D2
171	s4989		
172	s4991	FANCE	FA complementation group E
173	s4992		
174	s5015	FANCF	FA complementation group F
175	s5016		
176	s5019	FANCG	FA complementation group G
177	s5020		
178	s33619	FANCM	FA complementation group M
179	s33620		
180	s33621		
181	s5039	FAU	FAU, ubiquitin like and ribosomal protein S30 fusion
182	s5040		
183	s39592	FBH1	F-box DNA helicase 1
184	s39593		
185	s30663	FBXW7	F-box and WD repeat domain containing 7
186	s30665		
187	s5103	FEN1	flap structure-specific endonuclease 1
188	s5104		
189	s42145	FTSJ3	FtsJ RNA methyltransferase

473	s32296		
474	s11725	RAD21	RAD21 cohesin complex component
475	s11727		
476	s55074	RAD21L1	RAD21 cohesin complex component like 1
477	s55075		
478	s791	RAD50	RAD50 double strand break repair protein
479	s793		
480	s11734	RAD51	RAD51 recombinase
481	s11736		
482	s11746	RAD52	RAD52 homolog, DNA repair protein
483	s11747		
484	s11719	RAD9A	RAD9 checkpoint clamp component A
485	s11720		
486	s224514		
487	s44687	RAD9B	RAD9 checkpoint clamp component B
488	s44688		
489	s11837	RBBP4	RB binding protein 4, chromatin remodeling factor
490	s55169		
491	s19398	REC8	REC8 meiotic recombination protein
492	s19400		
493	s11902	RECQL	RecQ like helicase
494	s11904		
495	s17989	RECQL5	RecQ like helicase 5
496	s17990		
497	s28165	REV1	REV1, DNA directed polymerase
498	s28167		
499	s11939	REV3L	REV3 like, DNA directed polymerase zeta catalytic subunit
500	s11940		
501	s11941	RFC1	replication factor C subunit 1
502	s224529		
503	s11945	RFC2	replication factor C subunit 2
504	s11946		
505	s11947	RFC3	replication factor C subunit 3
506	s11949		
507	s11951	RFC4	replication factor C subunit 4
508	s11952		
509	s11953	RFC5	replication factor C subunit 5
510	s11955		
511	s39523	RIOX2	ribosomal oxygenase 2
512	s39524		
513	s20657	RNASEH2A	ribonuclease H2 subunit A
514	s20658		
515	s12074	RNF4	ring finger protein 4
516	s12075		
517	s12128	RPA1	replication protein A1
518	s12129		
519	s12131	RPA2	replication protein A2
520	s12132		
521	s26742	RPA4	replication protein A4
522	s26744		
523	s38769	RPAIN	RPA interacting protein
524	s38771		
525	s12357	RRM1	ribonucleotide reductase catalytic subunit M1
526	s12359		
527	s12360	RRM2	ribonucleotide reductase regulatory subunit M2
528	s12362		
529	s26992	RRM2B	ribonucleotide reductase regulatory TP53 inducible subunit M2B
530	s26993		

#	ID	Symbol	Description
190	s42147		homolog 3
191	s19754	G3BP1	G3BP stress granule assembly factor 1
192	s19756		
193	s19206	G3BP2	G3BP stress granule assembly factor 2
194	s19207		
195	s19030	GINS1	GINS complex subunit 1
196	s19031		
197	s28485	GINS2	GINS complex subunit 2
198	s28486		
199	s34942	GINS3	GINS complex subunit 3
200	s34943		
201	s38849	GINS4	GINS complex subunit 4
202	s38850		
203	s26649	GNL2	G protein nucleolar 2
204	s26650		
205	s25421	GNL3	G protein nucleolar 3
206	s25422		
207	s29189	GNL3L	G protein nucleolar 3 like
208	s29190		
209	s38171	GRWD1	glutamate rich WD repeat containing 1
210	s38172		
211	s24099	GTPBP4	GTP binding protein 4
212	s24100		
213	s17117	H1-10	H1.10 linker histone
214	s17118		
215	s74	HDAC1	histone deacetylase 1
216	s75		
217	s38332	HDAC10	histone deacetylase 10
218	s38334		
219	s6493	HDAC2	histone deacetylase 2
220	s6494		
221	s19459	HDAC6	histone deacetylase 6
222	s19461		
223	s23868	HEY1	hes related family bHLH transcription factor with YRPW motif 1
224	s23869		
225	s223604	HEY2	hes related family bHLH transcription factor with YRPW motif 2
226	s23929		
227	s223702	HEYL	hes related family bHLH transcription factor with YRPW motif-like
228	s25474		
229	s14520	HIRA	histone cell cycle regulator
230	s14522		
231	s194488	H1-4	H1.4 linker histone, cluster member
232	s6402		
233	s15858	H2BC14	H2B clustered histone 14
234	s194895		
235	s15857	H2AC20	H2A clustered histone 20
236	s194887		
237	s30813	HJURP	Holliday junction recognition protein
238	s30814		
239	s30815		
240	s27105	HP1BP3	heterochromatin protein 1 binding protein 3
241	s27106		
242	s6993	HSP90AA1	heat shock protein 90 alpha family class A member 1
243	s6994		
244	s6999	HSP90AB1	heat shock protein 90 alpha family class B member 1
245	s7000		
531	s25187	RSL1D1	ribosomal L1 domain containing 1
532	s25189		
533	s24739	RTTN	rotatin
534	s24741		
535	s19550	SAE1	SUMO1 activating enzyme subunit 1
536	s19552		
537	s25208	SENP3	SUMO specific peptidase 3
538	s25209		
539	s25142	SERBP1	SERPINE1 mRNA binding protein 1
540	s25144		
541	s12705	SET	SET nuclear proto-oncogene
542	s12706		
543	s19110	SETDB1	SET domain bifurcated 1
544	s19112		
545	s38215	SETDB2	SET domain bifurcated 2
546	s38217		
547	s12708	SETMAR	SET domain and mariner transposase fusion gene
548	s12709		
549	s23770	SIRT1	sirtuin 1
550	s23771		
551	s39053	SLX4	SLX4 structure-specific endonuclease subunit
552	s39054		
553	s13131	SMARCA1	SWI/SNF related, matrix associated, actin dependent regulator of chromatin, subfamily a, member 1
554	s13132		
555	s13139	SMARCA4	SWI/SNF related, matrix associated, actin dependent regulator of chromatin, subfamily a, member 4
556	s13140		
557	s16082	SMARCA5	SWI/SNF related, matrix associated, actin dependent regulator of chromatin, subfamily a, member 5
558	s16083		
559	s26996	SMARCAL1	SWI/SNF related, matrix associated, actin dependent regulator of chromatin, subfamily a like 1
560	s26997		
561	s15751	SMC1A	structural maintenance of chromosomes 1A
562	s15752		
563	s25864	SMC1B	structural maintenance of chromosomes 1B
564	s25866		
565	s17426	SMC3	structural maintenance of chromosomes 3
566	s17427		
567	s25747	SND1	staphylococcal nuclease and tudor domain containing 1
568	s25749		
569	s813	sNEG1	-
570	s210	SRP14	signal recognition particle 14
571	s224681		
572	s13435	SRP9	signal recognition particle 9
573	s227428		
574	s13451	SRPK1	SRSF protein kinase 1
575	s224686		
576	s12731	SRSF3	serine and arginine rich splicing factor 3
577	s12733		
578	s12734	SRSF4	serine and arginine rich splicing factor 4
579	s12735		
580	s13490	SSRP1	structure specific recognition protein 1
581	s13491		
582	s20074	STAG1	stromal antigen 1
583	s20076		
584	s21089	STAG2	stromal antigen 2
585	s21090		
586	s21086	STAG3	stromal antigen 3

#	Probe	Gene	Description
246	s14373	HSP90B1	heat shock protein 90 beta family member 1
247	s14375		
248	s7066	HUS1	HUS1 checkpoint clamp component
249	s7067		
250	s20916	IGF2BP1	insulin like growth factor 2 mRNA binding protein 1
251	s20918		
252	s7246	IGHMBP2	immunoglobulin mu DNA binding protein 2
253	s7247		
254	s7398	ILF2	interleukin enhancer binding factor 2
255	s7399		
256	s7401	ILF3	interleukin enhancer binding factor 3
257	s7403		
258	s29255	INO80	INO80 complex subunit
259	s29256		
260	s195940	INO80E	INO80 complex subunit E
261	s49403		
262	s16895	KAT2B	lysine acetyltransferase 2B
263	s16896		
264	s18197	KIF23	kinesin family member 23
265	s225150		
266	s22725	KIN	Kin17 DNA and RNA binding protein
267	s22726		
268	s51989	KMT5A	lysine methyltransferase 5A
269	s51990		
270	s27462	KMT5B	lysine methyltransferase 5B
271	s27463		
272	s39429	KMT5C	lysine methyltransferase 5C
273	s39430		
274	s35206	KRI1	KRI1 homolog
275	s35208		
276	s21893	KRR1	KRR1, small subunit processome component homolog
277	s21894		
278	s23665	LARP1	La ribonucleoprotein domain family member 1
279	s23667		
280	s37861	LAS1L	LAS1 like, ribosome biogenesis factor
281	s37862		
282	s35540	LBHD1	LBH domain containing 1
283	s35542		
284	s8174	LIG1	DNA ligase 1
285	s8175		
286	s8176	LIG3	DNA ligase 3
287	s8177		
288	s8179	LIG4	DNA ligase 4
289	s8180		
290	s38853	LLPH	LLP homolog, long-term synaptic facilitation factor
291	s56478		
292	s30777	LSG1	large 60S subunit nuclear export GTPase 1
293	s30779		
294	s39727	LTV1	LTV1 ribosome biogenesis factor
295	s39729		
296	s31155	LYAR	Ly1 antibody reactive
297	s31156		
298	s20467	MAD2L2	mitotic arrest deficient 2 like 2
299	s20468		
300	s11138	MAPK1	mitogen-activated protein kinase 1
301	s11139		
302	s11140	MAPK3	mitogen-activated protein kinase 3
303	s11141		
304	s17080	MBD2	methyl-CpG binding domain protein 2
305	s225056		
306	s17076	MBD4	methyl-CpG binding domain 4, DNA glycosylase
307	s17077		
308	s30853	MCM10	minichromosome maintenance 10 replication initiation factor
309	s30854		
310	s8586	MCM2	minichromosome

#	Probe	Gene	Description
587	s21088		
588	s13546	STAU1	staufen double-stranded RNA binding protein 1
589	s13548		
590	s14607	SUMO1	small ubiquitin-like modifier 1
591	s14608		
592	s13179	SUMO2	small ubiquitin-like modifier 2
593	s13180		
594	s13176	SUMO3	small ubiquitin-like modifier 3
595	s13177		
596	s13178		
597	s51833	SUMO4	small ubiquitin-like modifier 4
598	s51835		
599	s22113	SUPT16H	SPT16 homolog, facilitates chromatin remodeling subunit
600	s22115		
601	s13659	SUV39H1	suppressor of variegation 3-9 homolog 1
602	s13660		
603	s36183	SUV39H2	suppressor of variegation 3-9 homolog 2
604	s36184		
605	s36185		
606	s25811	TAF5L	TATA-box binding protein associated factor 5 like
607	s25812		
608	s20882	TAF6L	TATA-box binding protein associated factor 6 like
609	s20883		
610	s20824	TBL3	transducin beta like 3
611	s20825		
612	s13991	TERF2	telomeric repeat binding factor 2
613	s13993		
614	s29665	TEX10	testis expressed 10
615	s29666		
616	s14003	TFAP2A	transcription factor AP-2 alpha
617	s14004		
618	s19395	THOC1	THO complex 1
619	s19396		
620	s40362	TICRR	TOPBP1 interacting checkpoint and replication regulator
621	s40363		
622	s29864	TIPIN	TIMELESS interacting protein
623	s29865		
624	s230547	TMA16	translation machinery associated 16 homolog
625	s30718		
626	s9520	TONSL	tonsoku like, DNA repair protein
627	s9521		
628	s14304	TOP1	DNA topoisomerase I
629	s14305		
630	s14307	TOP2A	DNA topoisomerase II alpha
631	s14309		
632	s14310	TOP3A	DNA topoisomerase III alpha
633	s224746		
634	s17098	TOP3B	DNA topoisomerase III beta
635	s228087		
636	s606	TP53	tumor protein p53
637	s607		
638	s229446	TREX1	three prime repair exonuclease 1
639	s229448		
640	s37740	TRMT1L	tRNA methyltransferase 1 like
641	s37741		
642	s31336	TSR1	TSR1, ribosome maturation factor
643	s31337		
644	s14556	UBA52	ubiquitin A-52 residue ribosomal protein fusion product 1
645	s14557		
646	s14558		
647	s228085	UBB	ubiquitin B
648	s338		
649	s14560	UBC	ubiquitin C
650	s14561		
651	s14565	UBE2A	ubiquitin conjugating enzyme E2 A

311	s8587		maintenance complex component 2
312	s8590	MCM3	minichromosome
313	s8591		maintenance complex component 3
314	s8593	MCM4	minichromosome
315	s8594		maintenance complex component 4
316	s8595	MCM5	minichromosome
317	s8596		maintenance complex component 5
318	s8598	MCM6	minichromosome
319	s8600		maintenance complex component 6
320	s224035	MCM7	minichromosome
321	s8602		maintenance complex component 7
322	s39083	MCM8	minichromosome
323	s39084		maintenance 8 homologous recombination repair factor
324	s48537	MCM9	minichromosome
325	s48538		maintenance 9 homologous recombination repair factor
326	s8682	MEN1	menin 1
327	s8684		
328	s40987	MGME1	mitochondrial genome
329	s40988		maintenance exonuclease 1
330	s25711	MLH3	mutL homolog 3
331	s25713		
332	s48436	MMS22L	MMS22 like, DNA repair
333	s48437		protein
334	s8898	MNAT1	MNAT1, CDK activating
335	s8900		kinase assembly factor
336	s8960	MRE11	MRE11 homolog, double
337	s8961		strand break repair nuclease
338	s30362	MRM3	mitochondrial rRNA
339	s30363		methyltransferase 3
340	s27567	MRTO4	MRT4 homolog, ribosome
341	s27568		maturation factor

652	s14566		
653	s14569	UBE2B	ubiquitin conjugating enzyme E2 B
654	s14570		
655	s14590	UBE2I	ubiquitin conjugating enzyme E2 I
656	s14591		
657	s14723	USP1	ubiquitin specific peptidase 1
658	s14724		
659	s17367	USP10	ubiquitin specific peptidase 10
660	s17368		
661	s33026	WDR18	WD repeat domain 18
662	s33027		
663	s21378	WDR3	WD repeat domain 3
664	s21379		
665	s14907	WRN	Werner syndrome RecQ like
666	s14908		helicase
667	s32337	WRNIP1	Werner helicase interacting protein
668	s32338		1
669	s14925	XPA	XPA, DNA damage recognition
670	s14926		and repair factor
671	s14928	XPC	XPC complex subunit, DNA
672	s14930		damage recognition and repair factor
673	s14952	XRCC5	X-ray repair cross complementing
674	s14953		5
675	s52594	XRCC6	X-ray repair cross complementing
676	s5456		6
677	s5457		
678	s35020	YTHDC2	YTH domain containing 2
679	s35021		
680	s40699	ZNF830	zinc finger protein 830
681	s40701		
682	s224929	ZRANB3	zinc finger RANBP2-type
683	s38487		containing 3

Supplementary Table 2. Treatments/siRNAs included in the pilot screen

identifier	treatment type	target gene name	gene id	comment	control type
untreated	-	-	-	gelatin/sucrose coating only	negative
mock	transfection reagent	-	-	coating + transfection reagent	negative
sNEG1	siRNA	-	-	non-targeting siRNA	negative
sNEG9	siRNA	-	-	non-targeting siRNA	negative
sDEK	siRNA	DEK	DEK	-	transfection
APH	drug	-	-	200 nM APH	positive
CPT	drug	-	-	100 nM CPT	positive
sPOLD	siRNA	DNA polymerase δ	POLD1	catalytic subunit	positive
sPOLA	siRNA	DNA polymerase α	POLA1	catalytic subunit	positive
sKIF11	siRNA	kinesin family member 11	KIF11	microtubule motor protein	phenotypic
sAURKB	siRNA	aurora kinase B	AURKB	binds to mitotic spindle	phenotypic

Supplementary Table 3. Antibodies used in this study.

Whole cell lysates:

anti-DEK rabbit polyclonal K-877	1:20.000	F. Kappes
anti-GFP mouse monoclonal	1:1.000	Roche (1 814 460)
Goat-anti-rabbit-HRP	1:2.000	Dako Denmark (P0448)
Goat-anti-mouse-HRP	1:2.000	Dako Denmark (P0447)
Streptavidin-biotinylated-HRP	1:15.000	GE Healthcare

DEK-IP:

anti-DEK mouse monoclonal	1:500	BD (610948)
anti-SUMO-2/3/4 mouse monoclonal Clone C-3	1:500	Santa Cruz (sc-393144)
anti SUMO-1 mouse monoclonal Clone D-11	1:200	Santa Cruz (sc-5308)
Veriblot for IP Detection Reagent (HRP)	1:200	Abcam (ab131366)
Streptavidin-biotinylated-HRP	1:50.000	GE Healthcare

6His-SUMO Ni-NTA chromatography:

anti-DEK rabbit polyclonal K-877	1:20.000	F. Kappes
anti-SUMO-2/3/4 mouse monoclonal Clone C-3	1:500	Santa Cruz (sc-393144)
anti SUMO-1 mouse monoclonal Clone D-11	1:200	Santa Cruz (sc-5308)
Goat-anti-rabbit-HRP	1:2.000	Dako Denmark
Goat-anti-mouse-HRP	1:2.000	Dako Denmark
Streptavidin-biotinylated-HRP	1:5.000	GE Healthcare

In vitro SUMOylation:

anti-DEK rabbit polyclonal K-877	1:20.000	F. Kappes
Anti-GST mouse monoclonal	1:2.000	Santa Cruz (sc-138)
anti-SUMO-2/3 rabbit monoclonal Clone 18H8	1:500	Cell Signaling (#4971)
anti SUMO-1 mouse monoclonal Clone D-11	1:200	Santa Cruz (sc-5308)
Goat-anti-rabbit-HRP	1:2.000	Dako Denmark
Goat-anti-mouse-HRP	1:2.000	Dako Denmark
Streptavidin-biotinylated-HRP	1:25.000	GE Healthcare

Immunofluoresence MCF10A

anti-DEK mouse	1:50	Santa Cruz (sc-136222)
anti-H3K9ac rabbit	1:250	Invitrogen (710293)
anti-H3K9me3 rabbit	1:500	Abcam (ab8898)
anti-H3K27me3 rabbit	1:1000	Abcam (ab192985)
anti-lamin A rabbit	1:1000	Abcam (ab253000)
anti-PCNA rabbit	1:50	Sigma-Aldrich (HPA030522)
anti-CENP-A rabbit	1:400	Cell Signaling (2186)
2° anti-mouse Alexa Fluor 488	1:250	Invitrogen (A11001)
2° anti-rabbit Alexa Fluor 488	1:250	Invitrogen (A11008)
2° anti-rabbit Atto 532	1:200	Rockland (611-152-122)
2° anti-rabbit Atto 594	1:200	Sigma-Aldrich (77671)
2° FluoTag®-X2 IgG1 anti-Mouse Atto 647N	1:500	NanoTag Biotechnologies (N2002)
2° anti-mouse IgG F(ab')2-goat Alexa Flure 647	1:240	ThermoFisher (A21237)

Supplementary Table 4. Primer sequences for the site-directed mutagenesis of the DEK primary sequence.

Target site	Primer name	Primer sequence 5' – 3'
SUMOylation site 1	SUMO1 Nr.1-fwd	CGTGGAAGGCAAGAGGGAAAAGGCAAAAGTAGAGAGG
	SUMO1 Nr. 1-rev	GCATTGTCAACCTCTCTACTTTTGCCTTTTCCCTCTTGC
	SUMO1 Nr. 2-fwd	GGAAGGCAAGAGGGAAAAGGCAGCAGTAGAGAGG
	SUMO1 Nr. 2-rev	CTTGCATTGTCAACCTCTCTACTGCTGCCTTTTCCCTC
SUMO consensus motif	SUMO3-fwd	GAGGAGCCACCAAAAAAGACAGCCGCAAGAGAAAAACCTAAAC
	SUMO3-rev	GTAGCTTTCTGTTTAGGTTTTTCTCTTGCGGCTGTCTTTTTGG

Supplementary Materials and Methods

High-throughput siRNA screen coupled with time lapse fluorescence microscopy

U2-OS KI eGFP-DEK cells were transfected with siRNAs by solid-phase reverse transfection (Neumann et al., 2006): For the primary screen, the customized siRNA library comprised 678 unique siRNAs distributed on eight 96-well plates (the complete library can be found in **Supplementary Table 1**). In each plate, four wells contained the sNEG1 negative control, two wells the sPOLD1 (s616) positive control, two wells the sDEK transfection control (s15459) siRNA and three wells were left empty. The remaining wells contained the library siRNAs in alphabetical order. Before preparing the imaging plates, siRNAs of the mother plate were dissolved in 33 µl ddH$_2$O to a final concentration of 3 µM and stored at -20 °C.

For the pilot screen, a 3 µM dilution of each control siRNA was distributed on a 96-well imaging plate (see **Supplementary Table 2**) in technical triplicates. Plates were sealed directly after distribution of siRNAs and stored at -20 °C until further use.

The following protocol was used for the coating of eight identical 96-well imaging plates from one unique siRNA layout. First, the siRNA plate was thawed and 5 µl of the 3 µM siRNA dilution were transferred into a V-shaped 96-well plate. Then, 11 µl of a mixture of 554 µl of 0.4 M sucrose in Opti-MEM, 323 µl of Lipofectamine 2000 and 323 µl of ddH2O were distributed into another 96-well plate. 7 µl of this transfection reagent mixture were transferred into the previously prepared siRNA plate and mixed 8 times. After 20 min of incubation, 7 µl of sterile filtered 0.2% gelatin (0.45 µm pore size) were added to the wells and mixed 8 times. This final transfection mix was diluted 1:51 in water in several steps. From this diluted transfection mix, 50 µl were distributed into each well of the 8 96-well COC imaging plates. In 100 µl medium the final siRNA concentration per well was 7.7 nM. Plates were immediately lyophilized (37 °C, 1 h) and stored in plastic boxes containing drying pearls.

Cell seeding and time-lapse microscopy

For the first four plates of the primary screen (layout no. 1-4, replicate 1), 4000 U2-OS GFP-DEK cells per well were seeded in 100 µl McCoy's medium (without antibiotics) into

the coated 96- well imaging plate using an automated cell seeding device and incubated for 48 h at 37 °C, 5% CO_2. For all other plates, 3500 cells were seeded because cell confluency was slightly too high in some control wells. On the day of imaging, medium was exchanged with 100 µl pre- warmed phenol red free CO_2-independent imaging medium (Gibco) supplemented with 10% FBS, 1.5 mM L-glutamine and 2.3 g/l D-glucose. The full incubation IXM XLS high-content widefield microscope (Molecular Devices) was equipped with a PlanApochromat 20X/0.75 air objective and a sCMOS camera (4.66 megapixels). For GFP-DEK detection a 488 nm solid-state light source, the GFP filter set and a 690 nm laser autofocusing system were used while exposure time was set to 150 ms. For the time lapse experiment, four imaging field of views (FOVs) per well were set and images acquired every 12 min for 22 h. Two replicate plates per siRNA layout were seeded and imaged, for layout no. 5 three replicates. For the pilot screen, 2500 – 4000 cells per well were seeded, and 200 nM APH or 100 nM CPT were added to the imaging medium in the designated wells 1 h prior to imaging. For the time lapse experiment, 4-9 imaging FOVs per well were set and images acquired every 5 - 10 min for 24 - 48 h.

Image analysis using CellProfiler

The open-source software CellProfiler (v2.1.0 (Kamentsky et al., 2011)) was used for automated image analysis via a remote cluster to reduce computing time. The image analysis pipeline encompassed background correction, nuclei segmentation, filtering of dead and mitotic cells, intensity-based detection of DEK bodies and tracking of nuclei *via* the minimal overlap method.

The following steps were performed sequentially for every image of a time lapse. Images of the same time lapse were correlated using an object (here nuclei) tracking module. Then, the raw image was corrected for uneven illumination due to vignetting and for background fluorescence. As the effect was now limited to the corners of the image, the top- and leftmost 30, the rightmost 60 and the bottommost 160 pixels (px) were cropped ("cropped image"). The cropped image was smoothed by applying a Gaussian filter and

a threshold was calculated automatically with the global Otsu method to identify nuclei. Subsequently, the cropped image was masked with the nuclei outlines and the median intensity of the remaining background signal was subtracted from the cropped image ("corrected image"). Next, nuclei were segmented similarly by applying a Gaussian filter and calculating the threshold with the global Otsu method. Using this mask on the corrected image, nuclei of dead or mitotic cells were filtered out when meeting following requirements: Too high standard deviation and upper quartile of intensity and too small nuclear area. The filtered nuclei were tracked from image to the next image within a time lapse using a simple overlap method with a maximum pixel distance of 15. Afterwards, DEK bodies were identified: the corrected image was smoothed again by applying a Gaussian filter and DEK bodies were enhanced by performing a top-hat morphological operation. To avoid false-positive detection of peripheral micronuclei as DEK bodies, these protrusions were removed, and nuclei outlines were smoothed by performing an opening morphological operation. DEK bodies were identified within the smoothed outlines using a manual threshold (0.004 AU) and a diameter of 3 - 10 px. Finally, for each image CSV files containing image, nuclei and DEK body measurements were saved before the next image was analyzed. Aggregation of these raw data and evaluation of DEK body measurements was performed with the statistical software package R.

Data aggregation and transformation using R

The open-source programming language R with additional packages was used to aggregate the data. The goal was to obtain a single measurement per treatments/siRNA from the complex, time-resolved raw data of the technical (FOVs) and biological (plates) replicates that can be used to score DEK body regulators. As CellProfiler tracks the nuclei over subsequent frames/images, these track identifiers were used to summarize relevant measurements such as DEK body lifetime, number of DEK bodies per track etc. We noticed, however, that the CellProfiler pipeline not always managed to identify DEK bodies in all of the nuclei within a DEK body-containing sequence. We therefore wrote a custom function which used the closing morphological operation to fill these gaps with the DEK body counts of the previous or following frame (see lines 31-46 in the R code). In cases where a second, shorter DEK bodies-containing sequence was detected within the

same track, the function also served to remove these likely false-negative DEK bodies by keeping only the longest sequence. Additionally, a similar function was implemented to calculate the mean DEK body number within this longest sequence, which was used as the main readout to score hits (see lines 50-60 in the R code). DEK body related measurements were then further aggregated over FOVs to investigate the variability within the same siRNAs/treatments. Afterwards, the aggregated data of all replicates was concatenated. For the primary screen, quality control was then performed based on the FOV summaries. FOVs with a small integrated nuclear fluorescence intensity (very small and/or dark nuclei), very low cell number, blurry images and a negative slope of their linear growth function were filtered out. Furthermore, all wells that were contaminated with bacteria or yeast after the time lapse (as seen under a brightfield benchtop microscope) were filtered out manually.

Lastly, Z-scores of the FOV-based summaries of the mean number of DEK bodies were calculated. The Z-score can be used when the data distribution is unimodal and variability of the control measurement (here sNEG1) does not change in between replicates (Zhang, 2011). Therefore, the data was log (base 2)-transformed and the mean (X_{sNEG1}) and the standard deviation (SD_{sNEG1}) of all negative control (sNEG1) FOVs from the same replicate were calculated. For all FOVs of the other siRNAs of this replicate a Z-score was computed:

$$Z\text{-}score = \frac{log_2(X_i) - log_2(\bar{X}_{sNEG1})}{log_2(SD_{sNEG1})}$$

The mean of all Z-scores of the same siRNA was finally calculated. By ordering this mean DEK body number by size it was possible to score DEK body up-regulators (highest Z-score) and down-regulators (lowest Z-score). R code provided at https://github.com/CP-Vogel/R-DEK-body-script.git.

Low-throughput siRNA validation screen coupled with time-lapse fluorescence microscopy

For siRNA delivery, a liquid-phase instead of a solid-phase reverse transfection procedure was employed. Transfection mixes were freshly prepared for each replicate before cell seeding. Validation siRNAs were picked from the primary screen mother plates into a V-shaped 96-well plate and diluted to 100 nM with MQ H_2O. Two wells were supplied with the same validation siRNA, four wells with sNEG1 negative control, two wells with sDEK transfection control (s15459) siRNA and four wells were left empty. The new mother plate was stored at -20 °C until the day of transfection. At the day of transfection, 8 µl of a 1:20 dilution of Lipofectamine RNAiMAX in Opti-MEM was added in each well of a 96-well imaging plate (Ibidi). 25 µl of diluted siRNAs were mixed with the pre-distributed transfection reagent dilution. After 15 min incubation, 5500 U2-OS GFP-DEK cells in 207 µl medium (no antibiotics) were added for a final siRNA concentration of 10.4 nM and cells were incubated for 48 h at 37°C, 5% CO2.

48 h after transfection, medium was exchanged with 200 µl of pre-warmed phenol red free imaging medium (Hyclone, no antibiotics) and equilibrated in a pre-warmed CellObserver HSI microscope (Zeiss) equipped with a CSU-X1 spinning disc unit (Yokogawa) and a full incubator for 1 h at 37 °C, 5% CO2, humid atmosphere. Images were acquired with a PlanApochromat 20X/0.8 air objective, a 488 nm OPSL laser for GFP-DEK detection and an Evolve EMCCD camera (Photometrics, 512 x 512 pixels). Exposure time and EM gain were set to 150 ms and 700, respectively. For the time-lapse experiment, 6 imaging FOVs per well were set, the focus adjusted manually and images acquired every 12 min for 24 h. Three replicate plates were acquired in total.

In contrast to the primary screen, the image analysis was run on a newer version of CellProfiler (v.2.2.0 instead of 2.1.0) and locally on a workstation instead of a remote cluster. The pipeline was adjusted from the primary screen to meet the needs of the new microscope setup. Raw data analysis was performed as for the primary screen with minor modifications. Calculation of Z-scores and siRNA summaries was performed as described above.