Dissection of the role of a SH3 domain in the evolution of binding preference of paralogous proteins

Pascale Lemieux[1-4], David Bradley[1-5], Alexandre K Dubé[1-5], Ugo Dionne[1,2,6] and Christian R Landry[1-5]

[1] Institut de Biologie Intégrative et des Systèmes (IBIS), 1030, avenue de la Médecine, Université Laval, Québec (Québec), Canada, G1V 0A6

[2] Regroupement Québécois de Recherche sur la Fonction, l'Ingénierie et les Applications des Protéines, (PROTEO), 1045 Avenue de la Médecine, Université Laval, Québec (Québec), Canada, G1V 0A6

[3] Centre de recherche en données massives (CRDM), 1065, avenue de la Médecine, Université Laval, Québec (Québec), Canada, G1V 0A6

[4] Département de biochimie, microbiologie et bio-informatique, 1045 Avenue de la Médecine, Université Laval, Québec (Québec), Canada, G1V 0A6

[5] Département de biologie, 1045 Avenue de la Médecine, Université Laval, Québec (Québec), Canada, G1V 0A6

[6] Centre de Recherche du Centre Hospitalier Universitaire (CHU) de Québec-Université Laval, Québec, QC, Canada. Current address: Lunenfeld-Tanenbaum Research Institute, Sinai Health, Toronto, ON, Canada.

Authors for correspondence: pascale.lemieux.4@ulaval.ca or Christian.landry@bio.ulaval.ca

Keywords: SRC Homology 3 (SH3) domain; gene duplication; ancestral sequence reconstruction; myosins;

1 **Abstract**
2
Protein-protein interactions (PPIs) drive many cellular processes. Some PPIs are directed by
Src homology 3 (SH3) domains that bind proline-rich motifs on other proteins. The evolution
of the binding specificity of SH3 domains is not completely understood, particularly following
gene duplication. Paralogous genes accumulate mutations that can modify protein functions
and, for SH3 domains, their binding preferences. Here, we examined how the binding of the
SH3 domains of two paralogous yeast type I myosins, Myo3 and Myo5, evolved following
duplication. We found that the paralogs have different SH3-dependent interaction profiles.
However, by swapping SH3 domains between the paralogs and by characterizing the SH3
domains freed from their protein context, we find that very few of the differences in interactions,
if any, depend on the SH3 domains themselves. We used ancestral sequence reconstruction
to resurrect the pre-duplication SH3 domains and examined, moving back in time, how the
binding preference changed. Although the closest ancestor of the two domains had a very
similar binding preference as the extant ones, older ancestral domains displayed a gradual
loss of interaction with the modern interaction partners when inserted in the extant paralogs.
Molecular docking and experimental characterization of the free ancestral domains showed
that their affinity with the proline motifs is likely not the cause for this loss of binding. Taken
together, our results suggest that intramolecular interaction between a SH3 and its host protein
evolves through time, making domains not functionally equivalent even when they have the
same binding specificity.

Introduction
2
Protein domains are important structural and functional components of most proteins and often
have modular and specific functions (Doolittle 1995). Among these domains are PDZ, SH2
and SH3 domains (Kuriyan and Cowburn 1997; Castagnoli et al. 2004) that direct protein-
protein interactions (PPIs) by binding specific short linear motifs present on other proteins.
SH3 domains (hereafter referred to as SH3s) have become powerful models to study the
evolution of binding affinity and specificity because they are common and involved in many
signaling pathways in eukaryotes (Kay et al. 2000; Dionne et al. 2022).
10
SH3s are numerous in the budding yeast (n = 27) and human proteomes (n > 300) (Xin et al.
2013; Teyra et al. 2017), largely as a result of the duplication of the genes that code for their
host proteins (Vogel and Chothia 2006). Since the binding specificity of SH3s is key for the
function of SH3-containing proteins, substantial effort has been put in place to map the
determinants of their binding specificity. SH3 binding motifs most often correspond to PXXP
(where X is any amino acid) and fall into canonical groups (Kuriyan and Cowburn 1997;
Tonikian et al. 2009; Teyra et al. 2017). These groups sometimes overlap such that a given
SH3 shares binding specificity with others. This, in principle, allows different SH3s to bind
overlapping sets of proteins. However, it has been suggested that negative selection could act
to limit spurious binding with non-cognate partners, which would prevent crosstalk between
pathways with independent functions (Zarrinpar et al. 2003). Several other layers of regulation,
including other interaction domains and protein colocalization, therefore combine with motif
recognition to direct PPIs and allow a greater specificity (Dionne et al. 2022). This was recently
illustrated by large-scale PPI binding efforts that have shown that SH3-directed PPIs *in vivo*
also depend on the protein host and the position of the domain in the protein, which was
defined as the protein context (Dionne et al. 2021). Since there are many mechanisms that
contribute to the specificity of PPIs among SH3-containing proteins, one important question is
how much do the protein-binding domains themselves contribute to the divergence of PPIs
among paralogous proteins?
30
One powerful way to answer this question is to explore the evolution of the domains that have
been preserved after their recent duplication. These domains and their host proteins had the
same amino acid sequence at the moment of duplication, which makes it easier to understand
how a relatively small number of amino acid changes contribute to a shift in specificity. Among
the 27 SH3s present in 18 proteins of the budding yeast *Saccharomyces cerevisiae* are four
pairs of paralogous domains that originated from a whole-genome duplication (WGD) event
100-150 million years ago (Wolfe and Shields 1997; Marcet-Houben and Gabaldón 2015;
Wolfe 2015). Here, we focus on one of these protein pairs, the type I myosins Myo3 and Myo5.
39
Myo3 and Myo5 localize to actin cortical patches and are involved in endocytosis and
exocytosis (Mochida et al. 2002). They both contain C-terminal SH3s and have overlapping
sets of physical interactions in the Arp2/3 complex (Evangelista et al. 2000; Tarassov et al.
2008) and in actin polymerization networks (Lechler et al. 2000). From a genetics point of
view, these two genes are at least partially redundant. The single deletion of either *MYO3* or
MYO5 has a slight effect on growth, but a double deletion leads to severe growth and actin
cytoskeleton assembly defects (Goodson and Spudich 1995). Myo3 and Myo5 SH3s are a
powerful model to study paralogous evolution because their multiple shared PPIs are critical

for some cellular processes. In addition, they also have specific PPIs, which could provide the resolution needed to dissect the divergence of specificity early after duplication.

In this study, we focus on the divergence of PPIs between Myo3 and Myo5 and examine the contribution of their SH3s to this divergence. We first examined the protein interaction network *in vivo* of the two paralogous proteins to identify their SH3-dependent PPIs. We then swapped the domains between the two paralogs to examine their contribution to PPI divergence. We also used resurrected ancestral SH3s to examine how the two domains have diverged since their duplication and pushed further in time to examine how they have evolved prior to the duplication event. Finally, to examine how the SH3s themselves may have changed, independently of the rest of the proteins, we examined the PPI profile of the SH3s *in silico* and free from their protein context.

Results & Discussion

SH3 domains play a major role in the function of Myo3 and Myo5 *in vivo.*

The two paralogous genes *MYO3* and *MYO5* originated from the yeast WGD (Wolfe and Shields 1997). The corresponding amino acid sequences are 76% identical. They contain a N-terminal disordered region, a myosin motor, an ATP and actin-binding domain, and a C-terminal SH3 (Myo3 : 1123-1181, Myo5 : 1088-1146). Eleven of the 59 residues of their SH3 are different (Figure S1), which results in very similar structures as shown by the low root-mean-square deviation of atomic positions (RMSD = 0.386 Å, Figure S1). RMDS < 3Å is only observed between closely homologous structures (Chothia and Lesk 1986) .

We first compared the PPI profiles of Myo3 and Myo5 *in vivo* in yeast. We used them as baits in a screen with a set of 296 proteins as preys. We measured pairwise PPIs using the dihydrofolate reductase protein-fragment complementation assay (DHFR PCA) (Tarassov et al. 2008) (Figure 1A). The preys were selected based on previously detected PPIs and the signaling network of either or both paralogs including 93 known interaction partners (see methods). Also included are proteins that have been shown to interact ubiquitously in DHFR PCA and which can be used to detect changes in protein abundance (Tarassov et al. 2008; Levy et al. 2014) as PCA signal correlates with the amount of reconstituted DHFR enzyme (Freschi et al. 2013). We refer to them as abundance controls. We included variants of Myo3 and Myo5, as baits, for which the SH3s were encoded in codon-optimized sequences (optSH3s) for expression in *S. cerevisiae* since we also test codon-optimized ancestral SH3s below. Also, because our goal was to examine the role of the SH3s in the divergence of PPIs, the experiment included the SH3-depleted paralog variants for which a short flexible peptide (GGSSGGGG) was used to replace the SH3s. We used these constructs as baits to determine which PPIs are SH3-dependent (Figure 1A) (Dionne et al. 2021).

We detected most of the previously reported PPIs (117/206) and the majority of those detected by more than one method (78/95) using the BioGRID as a reference (Stark et al. 2006). Excluding the abundance controls for which information is absent from the reference dataset, only one interaction was detected for the first time (Figure S2). Myo3 and Myo5 share most of their interaction partners (Figure 1B). PPI mapping thus confirms their common function in the actin cytoskeleton organization (Goodson and Spudich 1995; Lechler et al. 2000) and their functional redundancy. Also, several PPIs and the quantitative nature of the PCA signal were

1 validated by performing small-scale validation experiments in individual liquid cultures
2 (Spearman, optSH3s r = 0.92, p = 3.3×10^{-6}, Figure S2).
3
The comparison between the median interaction scores (PPI score, see method) of the WT
codon and the optSH3 reveals very strong correlations for both paralog (Spearman, r ≥ 0.99,
p ≤ 1.6×10^{-6}) and the optSH3s and WT SH3s PPI scores have very similar medians (WT Myo3
: 0.764, optMyo3 : 0.756, WT Myo5 : 0.765, optMyo5 : 0.769) suggesting that the codon
optimization has no effect on the DHFR PCA results. Thus, the data obtained with the optSH3
paralog variants were used as reference to analyze further experiments testing codon-
optimized SH3s. Henceforth, we refer to the WT domains as extant domains (extantSH3s) as
we also discuss ancestral domains. We identified SH3-dependent partners by analyzing
DHFR PCA signals of the optSH3 paralog variants compared to the SH3-depleted variants
(Figure 1C). PPIs were classified as SH3-dependent when a significant difference in PPI
scores was observed between the SH3-depleted and optSH3 paralog variants (Wilcoxon,
Benjamini-Hochberg corrected p-value < 0.05). We observed for Myo3, 25 SH3-dependent
PPIs and 21 for Myo5, which completely overlap with Myo3 SH3-dependent PPIs (Figure 1B).
Also, the paralogs interact with each other displaying a modest but significant SH3-dependent
nature. Some of these SH3 dependencies support results from previous studies. For example,
as we observe here when the entire domain is removed, a previously characterized mutation
disrupts SH3-dependent PPIs (W36S) between Myo3 and Las17 (Evangelista et al. 2000) and
Myo5 with Bbc1 and Vrp1 (Geli et al. 2000; Mochida et al. 2002)(Figure 1B).
22
We manually reviewed the function of each interaction partner using the *Saccharomyces*
Genome Database (SGD) (Cherry et al. 2012). This analysis revealed that the SH3-dependent
partners are mostly involved in the cytoskeleton assembly whereas the SH3-independent
partners are involved in broader cellular functions or are the abundance controls (Figure 1B).
More than half (14/23) of the SH3-independent PPIs involve the abundance controls, which
are not expected to be dependent on the bait protein, leaving only nine specific SH3-
independent PPIs. Also, it is worth mentioning that other SH3-independent PPIs partners
could interact ubiquitously as a paralog of Ssb1, Ssb2, is an abundance control and that Ssa1
is a cytosolic chaperone (Werner-Washburne et al. 1987; Nelson et al. 1992) (Figure 1B). The
finding that many of the Myo3 and Myo5 PPIs depend on their SH3s confirms that the SH3s
have an important role in the function of these two proteins (Geli et al. 2000).
34
To validate our observations on SH3 dependency, the putative proline-rich binding motifs were
predicted on the SH3-dependent interaction partners. Predictions were done using a
specificity matrix of the preferred binding peptides determined by peptide phage display
(Tonikian et al. 2009). A motif prediction score (Maximum Motif Similarity (MSS) score)
between 0 and 1 was obtained for each prey by identifying the protein sub-sequence with the
strongest match to the Myo3/Myo5 specificity matrix. A positive correlation was observed
between the SH3 contribution to the interaction, i.e. the difference of PCA signal between the
optSH3 and SH3-depleted protein variants (Δ(PPI$_{opt}$ - PPI$_{SH3\text{-}depleted}$) median score), and the
motif prediction score for both paralogs (Spearman, Myo3 r = 0.64 p = 3.2×10^{-4}, Myo5 r = 0.51
p = 9.0×10^{-3}, Figure 1D). This supports our prediction that SH3-dependent PPIs depend on
the match of the cognate Myo3 and Myo5 SH3 binding motifs to their interaction partners,
although other major factors such as prey abundance could contribute to variation in PCA
signal. To further validate the SH3 and motif dependency of PPIs, we used the same linker
sequences used above to delete the SH3s but this time to replace the motifs with the highest

MSS score on interaction partners, creating motifΔ preys. We successfully modified nine of
these motifs and measured their impact on PPIs by DHFR PCA. A significant decrease in
interaction strength for both paralogs was observed for six out of nine partners when the top-
scoring proline motif was replaced with the linker. SH3-dependent PPIs are therefore most
likely dependent on the top-scoring proline motif on the interaction partners (Figure 1E). Cases
where no dependency or partial loss of PPI is observed, could be due to redundancy in the
poly-proline motifs as many of the interaction partners contain multiple proline-rich motifs
(Figure S3).

Figure 1. Characterization of the interactome of Myo3 and Myo5. A. Schematic
representation of the experimental design. The baits are shown in circular shape and preys
are shown in rectangle shape. PPIs are represented by the lines and the color of the prey
nodes shows the binding preference of each paralog. The left panel represents the WT PPI
network of the paralogs and the right panel shows the PPI network of the SH3-depleted
paralog variants. The comparison between both experiments allows us to identify SH3-
dependent interactions. **B.** Myo3 and Myo5 PPI network. The bait proteins are displayed in
the center of the network. The edges show the SH3 contribution to the PPI: $\Delta(\text{PPI}_{WT}$ -
$\text{PPI}_{depleted})$ median score. The vertical dash line separates the SH3-dependent, for one or both
paralog, and SH3-independent preys as determined by Wilcoxon tests. Node colors indicate
functions (Cherry et al. 2012). White nodes are abundance controls. **C.** Scatter plots
comparing the PPI scores of the optSH3 paralog variants with the SH3-depleted paralogs.

Myo3 data are shown on the left and Myo5 data are shown on the right as it is for panel D and
E. Each data point represents the median score (8 or 10 biological replicates) of an interaction
with an individual partner. Triangle-shaped data points are PPIs with fewer replicates (n ≤ 2)
which prevented us from concluding if they were SH3-dependent. The labeled data points are
preys that were used to confirm the proline motif predictions in panel E. **D.** Scatter plot
comparing the Δ(PPI$_{opt}$ - PPI$_{depleted}$) median score with the corresponding motif prediction score
of the preys (Maximum MSS). The color scale represents the p-value of the motif predictions.
E. PPI scores (48 biological replicates) of the WT preys and motifΔ preys with the extant
paralogs. Wilcoxon tests were performed between the extant and the motifΔ preys and
significance levels are show for each significant comparison (*** : p <= 0.001, **** : p <=
0.0001).
12
13   <u>SH3-dependent interaction differences between paralogs are not explained by the SH3</u>
14   <u>contribution to the PPIs.</u>
15
We compared the Myo3 and Myo5 interaction datasets and we observed that Myo5 raw PCA
signal is stronger on average than Myo3 raw PCA signal (Figure S2, Wilcoxon $p < 2.2 \times 10^{-16}$).
Since DHFR PCA measures the amount of protein complex formed (Remy and Michnick 2007;
Freschi et al. 2013), differences in binding could in principle result from differences in bait
abundance. In order to confirm that this bias is due to protein abundance, we measured protein
abundance using Myo3 and Myo5 GFP-tagged strains (Figure 2A). The results reveal that
Myo5 is slightly more abundant than Myo3 (Figure 2A, Student's t-test, $p = 6.1 \times 10^{-4}$), as
reported in public databases (Cherry et al. 2012; Breker et al. 2013). We also included the
SH3-depleted proteins to examine if the deletion of the SH3s affected abundance. Dionne et
al. showed that SH3 depletion had minimal to no effect on abundance by Western blotting for
many SH3-containing proteins. We observe the same results here using flow cytometry
(Figure 2A, Student's t-test, Myo3 p = 0.92, Myo5 p = 0.46). Thus, to compare the interaction
strength of the two paralogs, we had to ensure that the normalization method to compute the
PPI score eliminated the effect of bait abundance on the interaction data (see method).
Indeed, the median PPI scores of the abundance controls with the paralog variants confirm
that the abundance bias was removed by the normalization (Figure S2, Wilcoxon p = 0.96).
32
We quantified the similarity in binding profiles of Myo3 and Myo5 by directly comparing the
normalized PPI scores of the two extant proteins. This comparison reveals a strong correlation
for SH3-dependent interaction scores (Spearman, r = 0.91, $p = 1.4 \times 10^{-6}$) and an even stronger
correlation for SH3-independent ones (Spearman, r = 0.99, $p = 4.4 \times 10^{-6}$) (Figure 2B).
However, there remain a few important differences among the SH3-dependent PPIs. For
example, Bzz1 is a strong binding partner for Myo5 but a weak one for Myo3, and we observe
the opposite pattern for Pkh2 (Figure 2B). Most of the differences between the two paralogs
are observed for PPIs with weaker signals. DHFR PCA signals can reach saturation for
stronger PPIs (Remy and Michnick 2007), potentially preventing us from detecting divergence
for the high-intensity binding partners. Furthermore, if the divergence in SH3s is contributing
to the divergence of binding, we would expect those to be for cases of weak PPIs as SH3s
form weak transient PPIs with specific proline motifs (Dalgarno et al. 1997; Kay et al. 2000).
Sequence divergence in the SH3s could thus affect these PPIs most strongly. These results
suggest that the difference in binding between Myo3 and Myo5 could potentially be driven by
changes in their SH3.
48

In order to determine whether the SH3s are sufficient or not to confer PPI specificity to the
paralogs, we swapped the SH3s between them. A paralogous SH3 inserted in the non-native
paralog is referred to as a swapSH3 paralog variant. If the SH3s are sufficient to confer PPI
specificity, swapping the domains will also switch the SH3-dependent PPI profile of the
paralogs. Any differences caused by the paralogous SH3s could result from sequence
changes following their last common ancestor resulting in gains or loss of interaction strength
of either paralog (Figure 2C). To differentiate gains and losses of PPIs, the ancestral domain
at the last common ancestor node (DupSH3) was resurrected and its codon-optimized
sequence was also inserted in both paralogs. The ancestral sequence reconstruction (ASR)
was performed using a maximum likelihood-generated phylogeny across fungal Myo3/Myo5
orthologs (Figure S1, Data S2). The Myo3/Myo5 tree topology is supported by strong
phylogenetic branch supports (aLRT), and all ancestral sequences tested were reconstructed
with high posterior probability (Figure S1). Myo3 and Myo5 SH3 orthologous sequences are
mostly conserved, but 20/59 positions diverged since their last common ancestor in at least
one ortholog (Figure S1). Therefore, the DupSH3 has high sequence identity with both extant
SH3s (53/59 Myo3, 54/59 Myo5, Figure S1). The DupSH3 sequence was reconstructed with
high confidence (aLTR = 96.6) and 58/59 positions are predicted with 95% or higher posterior
probability (Figure S1).
19
Swapping the SH3s and inserting DupSH3 in Myo3 and Myo5 reveal that the interaction
profiles of the two paralogs are not affected by the difference in the sequence of their
paralogous SH3s and their ancestor (Figure 2D & 2E). The paralogs with swapped SH3s have
almost identical SH3-dependent interaction profiles as their native ones (Figure 2D,
Spearman, Myo3 r = 0.99 p = 8.4×10^{-9}, Myo5 r = 1 p = 1.5×10^{-6}). The same result is obtained
with the insertion of DupSH3 in both paralogs (Figure 2E, Spearman, Myo3 r = 1 p < 2.2×10^{-16},
Myo5 r = 0.99 p = 1.7×10^{-6}). Overall, these results show that, although different in sequence,
the two paralogous domains and their last common ancestor SH3 do not contribute to
differences in binding between the two paralogous proteins Myo3 and Myo5.

Figure 2. SH3 domains are not sufficient to explain the PPI preferences of the paralogs.
A. Expression level of extant paralogs and SH3-depleted protein variants tagged with GFP. The median cell fluorescence signal (arbitrary units) is measured for 5000 individual cells from three independent cultures (individual data point). The negative control is a strain not expressing GFP (strain BY4741). Student's t-test p-values are shown above each comparison. **B.** Comparison of the optSH3 variants interaction profiles of Myo3 and Myo5. The colors represent the SH3 dependency of the PPIs. Triangle-shape marks are PPIs with fewer replicates (n ≤ 2) for one of the paralogs. **C.** Schematic representation of the expected PPI network of the swapSH3 and DupSH3 paralog variants compared to the extant PPI network under different evolutionary scenarios. The PPIs are represented by lines and the colors of the nodes show the extant paralog partners. The swapSH3 determines which paralog-specific PPI can be recovered only by the contribution of its native SH3. The PPIs observed with DupSH3 identifies which paralog-specific PPI was gained or lost following the duplication event. **D and E.** Comparison of the PPIs of Myo3 and Myo5 containing various SH3s. The host protein is shown at the top of each plot. **D** shows the comparisons between the swapSH3 (y-axis) and the optSH3 (x-axis). **E** shows the comparisons of the DupSH3 (y-axis) with optMyo3 or optMyo5 (x-axis) in both paralogs. Spearman correlation coefficients are shown on the plots.

Ancestral SH3 domains progressively lose SH3-dependent PPIs.

Our results comparing Myo3 and Myo5 PPIs showed that they have quantitatively slightly different interaction profiles, especially for SH3-dependent PPIs (Figure 2B). However, none of the differences observed appears to be driven by the divergence of the domains themselves. Surprised by this, we wondered how robust PPIs of these two paralogs were to the identity of their SH3s. We therefore resurrected more ancestral sequences at speciation nodes (AncA, AncB, AncC) that predate the last common ancestor of Myo3 and Myo5 and

examined PPIs again (Figure 3A). AncA is the SH3 that was present in the last common ancestor of all the WGD species. Some of these species lost the second copy of the Myo3/Myo5 orthologs. AncB and AncC are, respectively, the last common ancestors for the *Saccharomycetaceae* (family) and *Saccharomycetales* (order) clades. The same ancestral sequence reconstruction method as the one used for DupSH3 was applied to infer AncA, AncB and AncC domains. High aLRT and strong topology support was obtained even for the oldest node used for the ancestral reconstruction (aLTR, AncA = 85.6, AncB = 89, AncC = 99.9, Figure S1). However, AncC's posterior probability distribution is slightly lower than the more recent ancestral SH3s, which is expected due to the higher sequence diversity used for the reconstruction (Figure S1, Data S2). Sequence identity gradually decreases to 55% between the extant Myo3 SH3 and AncC, and to 50% for Myo5 SH3 compared to AncC (Figure S1). To support the ASR predictions, we used AlphaFold2 to predict the ancestral SH3 3D structures (Figure 3B, Data S3) (Jumper et al. 2021). The predicted structures show very low RMSD when compared to the extantSH3s, indicating overall structure conservation despite the low sequence identity.

By measuring PPIs of Myo3 and Myo5 containing these ancestral sequences, we observe a general and progressive loss of signal as domains get more ancient, specifically for SH3-dependent PPIs (Figure 3C). For the most ancient SH3, the PPI network shows a significant interaction loss for its SH3-dependent PPIs compared to the optSH3s (paired-Wilcoxon, Myo3 $p = 5.5 \times 10^{-6}$, Myo5 $p = 2.4 \times 10^{-7}$). However, no difference is observed for the SH3-independent median PPI scores across SH3-depleted, AncC and optSH3 variants (Kruskal-Wallis, Myo3 $p = 0.47$, Myo5 $p = 0.98$). The pattern of loss for AncC PPI indicates that the ancestral domain loses PPIs in a similar way to the SH3-depleted paralogs (Figure 3C). Since SH3-independent PPIs with the abundance controls appear to be preserved, the gradual loss of interaction is unlikely to be caused by protein instability and degradation. To validate this hypothesis, we tagged the paralog variants with GFP and measured protein abundance. Protein abundance was not significantly different compared to the extant paralogs for any of the variants (Figure S2). Therefore, the results point towards an actual loss of binding by the ancestral domains. An exception in the general tendency is observed for the prey Ste20, which gains interaction with the AncC paralog variants (see below).

1
Figure 3. Ancestral SH3 domains affect the PPIs of Myo3 and Myo5. A. Simplified
phylogenetic relationship of yeast where nodes represent an extant SH3 (blue and red) or an
ancestral SH3 that was reconstructed with an estimation of the time at which these domains
existed. **B.** Superimposed extantMyo3 SH3 structure (blue, PDB:1RUW) and the AlphaFold2
predicted structures of each reconstructed ancestral SH3 showing high structural conservation
as estimated by RMSD. **C.** Heatmap showing the PPIs of SH3 variants inserted in Myo3 and
Myo5 with each of their interacting partners (preys, y-axis). The preys are ordered by SH3-
depleted Myo5 PCA signal and by interaction type with the paralogs (right annotation, cyan :
SH3-dependent for one or both paralogs, gray : SH3-independent, black : abundance
controls). The tile color corresponds to the median PPI score and the red x marks correspond
to a significant difference ($p < 0.05$, Wilcoxon test, Benjamini-Hochberg correction) in
interaction strength compared to the control interaction (dashed red frame). Missing data are
shown in gray.
15
16 <u>SH3 variants affinity to proline motifs does not explain the divergence between the</u>
17 <u>paralogous PPIs.</u>

The general decrease of binding for ancestral sequences, particularly of AncC, could be caused by a diminished binding affinity of this ancestral domain with the extant binding motifs. Using the binding motif prediction on SH3-dependent preys validated above (Figure 1D, E), we performed *in silico* SH3-peptide docking with the ancestral and extant SH3s to test this hypothesis.

The experimental structures of the extant Myo3 (PDB:1RUW) and Myo5 (PDB:1YP5) SH3s and the predicted AncC structure (Figure 3B, Data S3) were used for computational molecular docking (Geng et al. 2017) with peptides corresponding to binding motifs from the preys. Multiple structures were examined each time and they were sorted into clusters based on RMSD. The clusters and structures were ranked by the docking scoring algorithm (Geng et al. 2017). Next, a global minimization step was applied to the best structures of each cluster and the SH3-peptide interaction energies were computed (ΔG (kcal/mol)) (Schymkowitz et al. 2005). In 43% of the docking peptide-SH3 combinations, the best-ranked cluster included the lowest ΔG structures (Myo3: 8/28, Myo5: 15/28, AncC: 13/28). Additionally, we observed various conformations in the docked-peptide structures even if these structures are grouped in the same cluster (Figure 4A). We thus computed the median ΔG of the best structures of each cluster to consider this variability (see methods). Using the median ΔG to compare the clusters, we confirmed that the ranking of the clusters is a good indicator of the binding energy. Indeed, the first rank cluster shows a lower distribution of median ΔG compared to clusters with a higher rank for each SH3 (Figure S4). Thus, we continued the analysis with the median ΔG of the top cluster of each docking as a proxy for the affinity.

We observe significant correlations between the median ΔGs and the PPI scores obtained by DHFR PCA for both paralogs (Spearman, Myo3 r = -0.49 p = 0.02, Myo5 r = -0.52 p = 0.012) (Figure 4B). This result supports the binding motif predictions illustrated in Figure 1D given that the peptides used for structural modeling derive from our binding motif predictions. To examine if the paralogous SH3s have different binding affinities to the proline motifs, we compared the median ΔGs between Myo3 and Myo5 SH3s (Figure 4C). This comparison suggests that Myo5 SH3 binds globally more strongly to the proline motifs (predicted and confirmed) than Myo3 SH3 (paired-Wilcoxon, p = 0.018). In addition, the comparisons between AncC and the extantSH3s median ΔGs do not explain the data collected *in vivo.* Indeed, there is no significant decrease in binding affinity computed for AncC compared to the extantSH3s (paired-Wilcoxon, Myo3 p = 0.83, Myo5 p = 0.057) while a general decrease in binding is detected *in vivo* for the AncC-paralogs variants. We note however that in these comparisons, AncC appears to have median ΔGs more similar to that of Myo3 than of Myo5 SH3s, although the differences are not significant. Thus, even if the docking reveals a globally stronger binding of the proline motifs to Myo5 SH3 compared to Myo3 SH3, it does not explain the binding profile of the extant SH3s *in vivo*.

The conflicting results between the PCA experiment and the molecular docking do not allow us to conclude as to whether the SH3 affinity to the proline motifs is one of the main factors shaping the PPIs of Myo3 and Myo5, nor to explain the loss of interaction shown with the replacement of native SH3 by the AncC variant in the paralogs. The remaining hypothesis is that PPIs of the ancestral SH3 with the preys are reduced because a SH3 also interacts with its host protein in a way that decreases its ability to bind. The context in which the SH3 is found may therefore be a strong determinant of its ability to bind their recognition motif, as

1　recently shown by Dionne et al. 2021 for more distantly-related SH3s. We therefore performed
2　experiments with the SH3s free of their protein context.
3

4
Figure 4. Contribution of SH3 binding to their motifs estimated by molecular docking.
A. Superimposed structures of the ten best-scored docking structures of Osh2 binding peptide
on Myo3 SH3. The peptides show a variety of binding conformations whereas the SH3
structures remain similar. **B.** Scatter plots showing the relationship between the computed
median ΔGs of the docked peptide-SH3 complex and the corresponding PPI scores of the
extant paralogs as measured by DHFR PCA. Yellow data points correspond to proteins for
which the proline motifs have been confirmed *in vivo* (Figure 1E). Spearman correlation
coefficients (r) and p-values appear at the top of each plot. **C.** Boxplot comparing the
distribution of median ΔG values for each SH3s. The lines connect median ΔGs obtained with
the docking of the same peptide. Paired-Wilcoxon tests p-values are labeled on top of the
boxplot.
16
17　<u>Free SH3s lose paralog-specific PPIs pattern highlighting the role of protein context on SH3-</u>
18　<u>dependent PPIs.</u>
19
The SH3s were fused directly to the DHFR fragments to measure PPIs, which has been done
for instance to identify binding motifs (Wu et al. 2007; Tonikian et al. 2009). All the SH3 variants
described previously were expressed by fusing them to the DHFR F[1,2] and we tested PPIs
with the preys for which we successfully identified a binding motif (n = 7). The free SH3s were
all tagged at either terminus to test the effect of the position of the DHFR fragment on the
SH3s PPIs. The SH3 coding sequences were inserted at a genomic locus regulated by the
GAL1 promoter and the expression was under the control of a synthetic circuit induced by β-
estradiol (Aranda-Díaz et al. 2017) (Figure 5A). This allowed us to test PPIs at various bait
expression levels to ensure that we measure PPIs below PCA signal saturation. Analysis of
the data shows that the DHFR F[1,2] C-tag empty construct causes less background growth
than the DHFR F[1,2] N-tag empty construct (Wilcoxon, $p < 2.2 \times 10^{-16}$), so we focused on the
DHFR F[1,2] C-tag constructs.
32
We tested multiple expression levels by changing β-estradiol concentrations and we observed
that median PPI scores reach saturation in a prey-dependent manner. To compare the
different SH3 variants, we choose a β-estradiol concentration in the range of maximum
sensitivity of the assay, below saturation (see methods, Figure S5). The score difference Δ(WT
preys - motifΔ preys) median PPI score in the complete paralogs measured above and the
free SH3s shows a significant correlation for optMyo3 (Spearman, r = 0.89, p = 0.012) and the
same tendency for optMyo5 (Spearman, r = 0.64, p = 0.14, Figure S5) suggesting that the free
SH3s bind the preys in a similar manner as the SH3s in their host protein context.

The comparison of the median PPI scores between free optMyo3 and free optMyo5 shows a strong significant correlation (Spearman, r = 0.89 p = 0.012, Figure 5B). This is in agreement with the experiments above where the two domains have the same PPI profiles when inserted in Myo3 or Myo5. We observe no bias in interaction strength toward Myo5 SH3 compared to Myo3 SH3 when they are expressed isolated from the proteins (Figure 5B). This supports our results of the swapping experiment that showed that the affinity of the Myo5 SH3 for the prey tested is not higher than the affinity of the Myo3 SH3. However, we tested only seven preys in this experiment, which reduces the resolution to observe divergence. Even if we find that the free paralogous domains generate similar PPIs, the free SH3s do not recreate the paralog SH3-dependent PPI profiles. For example, the PPIs of the paralogs with Osh2 are very strong and are highly affected by the SH3 depletion (Figure 3C), but the free SH3s are not able to generate as strong PPIs (Figure 5B). This can be explained by the competition happening between the free SH3 and the extant paralogs for the binding of their target peptides in those strains.The strains expressing the free SH3s are also expressing the two WT extant paralogs.

As observed previously, the SH3s are causing differences in the PPI scores, but, unlike in the experiments with the entire proteins, we do not detect a systematic decrease in PPI scores of the free AncC compared to the free optSH3s (Figure 5C). In some cases, free AncC creates stronger PPIs than the free optSH3s. Even if it seems that the N-tag free AncC loses PPIs, the statistical test is not significant (Figure S5, Kruskall-Wallis, p = 0.44), probably because of the limited number of preys included in the experiment and the generally low signal for this construct (Figure S5). Likewise, there is no apparent decrease in PCA signal for the C-tag free AncC (Figure S5, Kruskall-Wallis, p = 0.27). This validates the docking results, which predicted that the binding affinity of AncC would be similar to those of the extant domains.

An additional finding we wanted to confirm with these experiments was the stronger interaction between Ste20 and AncC paralog variants when the complete paralogs were considered. We identified the binding motif mediating this PPI (Figure S3) and we observed a stronger interaction with the free AncC compared to the optSH3s (Wilcoxon, p = 1.6×10^{-4}, Figure 5C). However, it is difficult to fully rationalize this result as molecular docking does not predict a higher affinity between AncC and Ste20 motif (Figure 4C). Indeed, it is possible that AncC and Ste20 do not interact as a typical SH3-peptide interaction, thus the docking structure restraints used would not be appropriate to study AncC-Ste20 interaction.

Overall, while ancestral SH3s tend to lose PPIs when introduced in the extant protein, they do maintain an ability to interact with their partners when expressed alone. It is well established that SH3 domains have an intrinsic binding specificity that is encoded in their sequence (Teyra et al. 2017; Brown et al. 2018). However, our results suggest that the ability of the domain to bind motifs on other proteins is allosterically regulated by its host protein. Inserting an ancestral sequence into an extant protein could disrupt this allosteric regulation, leading to the general loss of PPIs despite the fact that the domains themselves maintain their ability to bind. This type of regulation implies that coevolution takes place between an SH3 and the rest of its protein such that the intrinsic binding specificity of a domain is not, alone, sufficient to dictate the specificity of the protein.

Figure 5. Characterization of free SH3 binding profiles. A. Schematic representation of the
DHFR PCA experimental using free SH3s with an inducible promoter. The transcription factor
GEM binds to β-estradiol and induces the expression of the SH3 coding sequence fused with
the DHFR F[1,2] via the *GAL1* promoter. The expression of the construct increases with the
amount of β-estradiol, leading to an increase of the DHFR PCA signal. **B.** Scatter plot
comparing the PPI profile of the free paralogous SH3s and that of the complete paralogs
(entire proteins with the optSH3s). Spearman correlation coefficients (r) and p-values for each
PCA experiment are shown. **C.** Boxplot of the PPI scores for several preys (x-axis) with three
SH3 variants. Kruskall-Wallis tests compare the scores among the three variants for each
prey. P-value significance levels are labeled at the top of each group (ns : p > 0.05, * : p <=
0.05, ** : p <= 0.01, *** : p <= 0.001, **** : p <= 0.0001).

Conclusion

We compared the interaction profiles of two recently evolved paralogous proteins, Myo3 and
Myo5, that contain SH3s to examine if the sequence divergence of the domains contributed
to the divergence in binding to other proteins *in vivo*. Our results show that Myo3 and Myo5
have largely overlapping sets of interaction partners but some of their PPIs, which we showed
to be SH3-dependent, are biased for one paralog or the other. To test if the SH3 domains are
causing these differences in interactions, we swapped the two domains between Myo3 and
Myo5. This experiment showed that the two domains are functionally equivalent since they
had the same PPI profiles. The functional equivalence of the two paralogous SH3s since their
duplication was also confirmed by inserting the ancestral sequence reconstruction of the SH3
at the paralogs' last common ancestor node in the extant paralog. Again, the interaction
profiles of the two paralogs were not affected.

To examine further the evolution of the SH3s, we looked at more ancient resurrected domains.
Surprisingly, we found that in general Myo3 and Myo5 lose their SH3-dependent PPIs when
their SH3s were replaced with older ancestral sequences. However, molecular docking
showed that the affinity of the most ancient domain reconstructed, AncC, for the proline motifs
is not lower than the one of the extant SH3s. Experiments with the SH3 domains isolated from
their host proteins revealed similar results. This suggests that although the SH3s may have
maintained their binding specificity for millions of years, the way they interact with their host
proteins may have changed. Allosteric interactions between the host proteins and its SH3
could be disrupted when inserting the ancestral domains into extant proteins, leading to
weakened interactions. Our results thus strengthen the postulate that protein domains are not

isolated features of proteins (Melero et al. 2014; Vishwanath et al. 2018) and that evolution is shaped by coevolution between the domains and the rest of the protein for proper functioning *in vivo*.

Author contributions

PL, DB, AKD and CRL were involved in the design of the study. CRL directed this work. DB performed the phylogenetic analyses and proline motif predictions. PL and AKD performed the experiments. PL performed the molecular docking results and wrote the first draft of the manuscript. DB wrote sections of the manuscript. UD designed part of the molecular biology material. PL, DB, AKD, UD and CRL contributed to revising, reading and approving the final version of the manuscript.

Data Availability

Strains and plasmids are available upon request. File S1 contains Tables S1-S12. Tables S1-S4 contain (solid and liquid) results from all the DHFR PCA experiments described. Table S5 contains the proline motif prediction information. Table S6 contains the processed flow cytometry data. Table S7 contains a summary of the molecular docking *in silico* experiments. Tables S8 and S9 contain respectively the oligonucleotides and the SH3 DNA sequences. Tables S10 and S11 describe respectively the bait and prey strains used in this study. Table S12 describes all reagents and resources used in this paper. File S2 contains additional details on the methods. Raw data for the DHFR PCA, AlphaFold2 structure predictions, molecular docking and the detailed data from the phylogeny are available on Dryad at : https://doi.org/10.5061/dryad.sj3tx968m (Data S1-4). Code used to analyze the raw data and to generate all figures can be found at https://github.com/Landrylab/Lemieux_et_al2023.

Acknowledgments

We thank Soham Dibyachintan, Philippe Després, Romain Durand and Isabelle Gagnon-Arsenault for valuable comments on the manuscript and on the experimental design. We also thank Patrick Lagüe for his insight on molecular docking design. We are also grateful for the support of all the LandryLab members. Some of the figures were created with BioRender and all molecular graphics were performed with UCSF ChimeraX,.

Funding

This work was supported by a Canadian Institutes of Health Research (CIHR) Foundation grant 387697 and a HFSP grant (RGP0034/2018) to CRL. CRL holds the Canada Research Chair in Cellular Systems and Synthetic Biology. PL was supported by an Alexander Graham Bell master's scholarship from NSERC (BESC M), a graduate scholarship from PROTEO and a Leadership and Sustainable Development Scholarship from Université Laval. DB is funded by the European Molecular Biology Organization (EMBO) Long-Term Fellowship (ALTF 1069-2019).

Conflict of interest

The authors declare no conflict of interest.

1 **Methods**
2
3 Sequence analysis
4
5 Multiple sequence alignments (MSA) were performed with MUSCLE v3.8.1551 (Edgar 2004)
6 using the sequences retrieved from the SGD (Cherry et al. 2012).
7
8 Strain construction
9
Yeast strains used for the DHFR PCA experiments were retrieved from the Yeast Protein
Interactome Collection (Tarassov et al. 2008) or constructed. Tables S12 provide a list of the
strain backgrounds and plasmids used to construct the strains. All of the SH3 codon optimized
strains (14) were constructed in this study, while WT codon SH3 bait strains (8) were retrieved
from Dionne et al. 2021. Bait strains were constructed using strain BY4741 (Mat**a**, *his3Δ leu2Δ*
met15Δ ura3Δ) according to Tarassov et al. 2008 method (Table S10). Of the prey strains, 6
were individually reconstructed using BY4742 (Matα, *his3Δ leu2Δ lys2Δ ura3Δ*) and 290 were
directly retrieved from the Yeast Protein Interactome Collection (Table S11). Preys of interest
were selected based on a previous study (Dionne et al. 2021) and manually curated from the
SGD (Cherry et al. 2012). CRISPR-Cas9 genome editing and homologous recombination
were used to construct the paralog variants, the motifΔ preys and the free SH3 constructs.
The appropriate media were used to select for the positive clones. NAT (Cedarlane Labs),
HYG (Bioshop Canada) and G418 (Bioshop Canada) selections were used at final
concentrations of 100 µg/ml, 250 µg/ml and 200 µg/ml respectively. The detailed methods are
available in File S2 and illustrated by Figure S6.
25
26 DHFR PCA growth conditions
27
Bait and prey strains were grown on YPD medium (1% yeast extract, 2% tryptone, 2% glucose,
and 2% agar (for solid medium)) containing their specific selection antibiotic. NAT and HYG
are used for diploid selection. DHFR PCA selection was made on a synthetic medium (PCA
medium, 0.67% yeast nitrogen base without amino acids and without ammonium sulfate, 2%
glucose, 2.5% noble agar, drop-out without adenine, methionine, and lysine) containing
200 µg/ml methotrexate (MTX, Bioshop Canada) diluted in dimethyl sulfoxide (DMSO,
Bioshop Canada).
35
36 Protein-fragment Complementation Assays and analyses
37
DHFR PCA protocol is based on Tarassov et al. 2008 and it was published as a visualized
protocol (Rochette et al. 2015). The principal purpose of DHFR PCA experiments is to assay
the interaction strength between two proteins *in vivo*, by the complementation of the DHFR
F[1,2] and DHFR F[3]. The PCA signal is proportional to the amount of reconstituted DHFR in
the cell. The detailed protocol is described in File S2. Matting between the baits and preys
was performed in random arrays (384 format) on a selective medium resulting in eight or ten
replicates for each PPI. The arrays were condensed into 1536 format and then replicated on
MTX medium for the assay. Pictures of the microbial growth at the final time point were
acquired and used for the analyses.
47

For each plate picture, the software Pyphe quantified the colonies' area ('pyphe-quantify batch --grid auto_1536 --t 1 --d 3 --s 0.05' (Kamrad et al. 2020)). Raw data from the prey array, diploid selection and PCA selection are available in Data S1. Positions were ignored when there was no growth on the diploid selection or on the prey array. The area values were transformed in $\log_2$ scale. Then, the background and aberrant data were assessed and removed ($3 < \log_2(area) < 13.13$, Figure S2). A standardization using the maximum and minimum logarithm area values on each plate transformed all $\log_2(area)$ on a scale between 0 and 1 (PPI score, see equation below).

$$PPI\ score\ = \frac{log2(area)\ -\ log2(min.area)}{log2(max.area)\ -\ log2(min.\ area)}$$

If there were two or less biological replicates left after data filtering, the PPI scores were ignored. All the statistical analyses performed on PPI scores are illustrated in the figures or mentioned in the main text. The control group of the statistical tests for the comparison between SH3 variants within one paralog was the optSH3 variant. We choose optSH3 as the control group because ancestral sequences are also codon optimized. Many PPIs were validated in liquid PCA experiments. The PPI network is visualized with Cytoscape (Figure 1B) (Shannon et al. 2003).

Comparison of interactome with literature

As a quality control of the data, the BioGRID database (v4.4.209 (Stark et al. 2006)) was used as a reference for verification of previously described PPIs. The physical PPIs in both directions (A-B and B-A) were retrieved and used for comparison (Figure S2).

Proline motif prediction

Position weight matrix (PWM) specificity models for yeast SH3s were obtained from Tonikian et al. 2009. With the PWMs, a maximum matrix similarity (MSS) score was computed for each SH3-dependent paralog PPI partner (Dionne et al. 2021). The scores are a value between 0 and 1, where 1 represents a perfect match to the PWM (Kel et al. 2003). For each k-mer (k = number of PWM columns) in the preys, MSS scores were obtained and the maximum MSS k-mer across the whole protein sequence was identified as the predicted binding motif. This procedure was realized for the SH3s of Myo3 and Myo5. The p-value of the prediction was assigned by testing if the maximum MSS score for one prey is higher than the MSSs obtained for 10,000 random peptides. An empirical p-value of 0.05 indicates that the predicted binding motif had a higher MSS score than 95% of the random peptides tested against. Proline motif predictions were used to design nine motifΔ preys for which seven were confirmed experimentally (Figure 1E). The maximum MSS motif positions are shown in Figure S3 (Brennan 2018).

GFP strain construction and cytometry fluorescence assay

The green fluorescent protein (GFP) DNA sequence and the HPHNT1 resistance cassette were amplified by PCR for genomic integration (File S2). The primers (Table S1) were designed to add 40-bp homology arms at each side of the GFP-HPHNT1 amplicon. The DNA fragments were integrated by homologous recombination at the 3'-end of the paralogs replacing the DHFR F[1,2] fragment and the NATMX4 resistance cassette. Hygromycin

1 selection was applied to the cells and the genomic integration was validated by PCR and
2 Sanger sequencing.
3
Validated GFP-tagged strains were grown at 30°C in synthetic medium (SC pH 6 HYG, 0.174
% yeast nitrogen base without amino acids and without ammonium sulfate, 2% glucose, 1%
succinic acid, 0.6% NaOH, 0.1% MSG) overnight in triplicates. The cultures were diluted in
liquid PCA medium without MTX (0.67% yeast nitrogen base without amino acids and without
ammonium sulfate, 2% glucose, drop-out without adenine, and 2% DMSO (Bioshop Canada))
to an OD of 0.1, then grown again until the exponential phase was reached (OD = 0.5 - 0.7).
The cells were diluted to an OD of 0.05 in sterile water, then the fluorescent intensity of 5000
cells per replicates was measured by a Guava EasyCyte HT cytometer (blue laser, λ = 488
nm).
13
14 <u>Ancestral sequence reconstruction</u>
15
Ortholog protein sequences of Myo3 and Myo5 were retrieved for fungal species using
Ensembl Compara (Vilella et al. 2009) (Data S2), ensuring systematic annotations. A MSA of
the full-length protein sequences was then produced with MAFFT L-INS-i (v7.453 (Katoh et
al. 2005)). Redundant sequences were filtered at a sequence identity threshold of 90% and
heavily gapped positions outside of the SH3 were manually trimmed from the alignment. This
processed MSA was used to compute a phylogenetic tree using a maximum-likelihood based
approach with IQ-TREE2 (Minh et al. 2020)(Figure S1). The best-fitting amino acid substitution
model (LG+G4) was searched automatically within IQ-TREE2 and phylogenetic branch
support was assessed using the SH-aLRT test (Anisimova et al. 2011). The tree is represented
in Figure S1 (Letunic and Bork 2021). Ancestral sequences were reconstructed for the SH3
portion of the alignment with the SH3 MSA and the phylogenetic tree using FAST-ML (v3.11
(Pupko et al. 2000)). The ancestral sequence reconstruction (ASR) was performed at every
tree node using an LG substitution matrix, maximum-likelihood reconstruction of insertions
and deletions, and a joint approach for ASR ('--jointReconstruction yes'). The nodes of interest
were chosen using the taxonomic annotations provided by Ensembl Compara. The ancestral
sequences at the node of the most recent ancestor for the clades *Saccharomyces* (Dup1,
Dup2 and Dup3, 1st to 3rd most likely sequences), *Saccharomycetaceae* (AncB) and
Saccharomycetales (AncC) were selected. Some species in the WGD clade lost the additional
copy of *MYO3/MYO5*, so we also selected the WGD node as another ancestral sequence
(AncA). The nodes' age were estimated with TimeTree (Kumar et al. 2017). The boundaries
of the Myo3/Myo5 SH3s were determined from the SMART database (V8.0 (Letunic et al.
2021)). Visualization of the MSAs (Figure S1) were realized with the R package ggmsa (Zhou
et al. 2022).
39
40 <u>Molecular docking</u>
41
The experimental structures of Myo3 (PDB:1RUW) and Myo5 (PDB:1YP5) SH3s were used
for the docking. AncC AlphaFold2 (Jumper et al. 2021)(AF2) best-ranked model was also used
(Figure 3C). The structure prediction has a high coverage and pIDDT across the entire
structure (Data S3). 28 predicted proline motif structures were computed with AF2 and the
best-ranked model for each motif was used for molecular docking (Data S4) (Tsaban et al.
2022). The molecular docking of the motif predicted structures on the SH3s was done using

Haddock2.4 (Dominguez et al. 2003) with parameters optimized for protein-peptide docking (Geng et al. 2017). Ambiguous interaction restraints were defined based on previous work (Janz et al. 2007) (Figure S4) and all docking parameters are documented in the run.cns files (Data S4). 400 structures resulted from each docking. They were split in clusters according to Haddock2.4 algorithm. The structures were ranked within each cluster, and the clusters were also ranked between each other by the scoring algorithm. The 10 best structures of each cluster, based on Haddock2.4 scoring algorithm, were kept for further computations. The global energy of docking structures was minimized with FoldX ('RepairPDB' (Schymkowitz et al. 2005)) 10 times (Usmanova et al. 2018). Then, the interaction energy (ΔG) was computed with FoldX ('AnalyseComplex' (Schymkowitz et al. 2005)) for each energy minimized docking structure. The median ΔG across the 10 best structures was computed for each cluster. All raw and transformed docking data are available in Data S4. Also, all protein visualizations were realized with UCSF ChimeraX (Pettersen et al. 2021).

Figure S1. Phylogeny of Myo3/Myo5 orthologs and ancestral sequence reconstruction.
A. Phylogeny of Myo3 and Myo5 fungal orthologs. The topology obtained respects the fungal taxonomy (Schoch et al. 2020). Myo3 and Myo5 are in mirror clades indicating gene duplication (DupSH3). The branches used for the ASR are differently colored and the resurrection nodes are labeled by a star (DupSH3, AncA, AncB, AncC). aLRT values on each branch show overall strong support for the topology. **B.** Superposition of the experimental

structures of Myo3 (PDB:1RUW) and Myo5 (PDB:1YP5) extantSH3s shows their high level of
similarity (Root Mean Square Deviation of atomic position (RMSD) shown). **C.** Histograms of
the highest posterior probability amino acid reconstructed at each position for each ancestral
SH3 sequence. The bin width is 0.05. **D.** Multiple sequence alignment of the SH3s belonging
to the *Saccharomyces* clade used to reconstruct DupSH3 with conservation chart below. SH3
secondary structures are annotated above. **E.** Multiple sequence alignment of the extant and
ancestral reconstructed SH3s. Dup1, Dup2 and Dup3 are the three most likely sequences at
the last common ancestor node. Dup1 is the one used throughout the study (DupSH3).
Position 31 is the only one changing among Dup1, Dup2 and Dup3.

Figure S2. DHFR PCA quality control and validation experiments. A. Distribution of the
colony areas on a $\log_2$ scale for the first DHFR PCA experiment (complete paralog with SH3
variants). Each line represents the distribution for one plate of 1,536 colonies. The peak
centered at 0 represents the empty positions (no growth) and thus, the background. Colonies
with residual growth constitute the middle peak (no PPI) while colonies for which an actual PPI
is detected from the peak just above 10. The positions with a colony size between the gray
lines are used for normalization (see method). The red line shows the threshold used for
identification of true PPIs. **B.** Violin plot showing the distributions of the colony areas on a $\log_2$

scale for data used for normalization. Myo5 shows larger median colony areas than Myo3
(Wilcoxon, p < 2.2x10⁻¹⁶). **C.** Boxplot comparing the median PPI scores of the abundance
control with the WT baits. No significant difference is observed (Wilcoxon, p = 0.96). **D.** Scatter
plot comparing PPI scores obtained for both technical replicates. **E.** PPIs detected in this study
compared with PPIs deposited in the BioGRID database according to the experimental
methods which previously identified the PPIs (v4.4.209 (Stark et al. 2006)). **F.** Protein
abundance of the various constructs measured by flow cytometry of GFP fusion proteins. The
median of fluorescence intensity for 5000 cells in three replicates is shown. The mean of all
SH3 variants for each paralog is represented by a dashed line. Multiple Student's t-test p-
values, Benjamini-Hochberg corrected, comparing the extant paralogs (extantSH3) with the
paralogs containing SH3 variants are labeled. **G.** Scatter plot showing median DHFR PCA PPI
score obtained from growth on solid medium plates and in individual liquid cultures from
replicated crosses (see method). Growth in liquid cultures was quantified by the logistic area
under the curve (AUC) after 65 hours of growth. Different colors and shapes are used to show
the paralog variants. Specific and global Spearman correlation coefficients (r) are shown. The
preys chosen for this experiment are described in Table S4.

Figure S3. Characterization of the proline motif role in PPIs *in vivo*. A. Scatter plot
comparing the median optSH3 PPI score for each SH3-dependent interaction with their
corresponding motif prediction score (Maximum MSS). The color scale represents the p-value
of the motif predictions which acts as a confidence measure. **B.** Representation of the potential
binding motif (PXXP) positions on the protein sequence of the tested preys. The motifs with
the highest MSS score which were deleted in each prey *in vivo* are shown in yellow. **C.** PPI
scores of the preys with the binding motifs replaced with a flexible linker compared to the WT
preys with paralog variant AncC and SH3-depleted. P-values from Wilcoxon tests are shown
(* : p <= 0.05, ** : p <= 0.01, *** : p <= 0.001, **** : p <= 0.0001) for each pair. Each interaction
was tested in 48 biological replicates.

Figure S4. Characterization of the role of the proline motif *in silico*. A. Illustration of the ambiguous interaction restraints used for the docking on the Myo3 SH3 structure (PDB:1RUW). Red residues were used as active residues by Haddock2.4. **B.** Violin plot showing the distribution of median ΔGs obtained for the ten best structures of each cluster per peptide (n = 28) for each SH3 variant. The black line is the generalized linear regression of the median ΔGs over each cluster. There is a significant positive relationship between the mean ΔG and the cluster rank for all SH3s tested.

Figure S5. Description of the free SH3s DHFR PCA results. A. Scatter plot comparing the replicated PPI scores at each β-estradiol concentration tested (μM). High correlations are

observed from 20 µM β-estradiol and higher. For 40 µM and 80 µM conditions, the technical replicates 2 grew less than the replicates 1 which we observe by correlations deviated from the perfect diagonal. Thus, the technical replicates 2 for 40 µM and 80 µM β-estradiol conditions were removed from the analysis. **B.** Plots showing median PPI scores (y-axis) at different β-estradiol concentrations (x-axis, in a $\log_2$ scale) for each prey. The position of the DHFR F[1,2] fragment is shown independently. The colors represent the different baits. The vertical gray line on each plot shows the β-estradiol concentration at which the sensitivity of the assay was maximal for the prey tested. The median PPI scores obtained at these concentrations are used to compare the different preys. **C.** Scatter plot showing the effect of the motifΔ preys on the PPI scores (Δ(WT prey - motifΔ prey)) for the free SH3s and the complete paralog DHFR PCA experiments. The SH3 variant and the DHFR F[1,2] position are indicated by the row and columns. The prey standard names labeled each data point. Spearman correlation coefficients and p-values are shown for each case. **D.** Heatmap showing the median PPI scores measured between preys (y-axis) and free SH3s (x-axis) tagged with the DHFR F[1,2] in N- or C-terminus and at the β-estradiol concentrations determined in panel B. Disregarding the empty construct, the median PPI scores are not different across the free SH3s (Kruskall-Wallis, C-tag p = 0.27, N-tag p = 0.44).

Figure S6. Illustration of the strain construction strategies. A. General strategy to insert the SH3 sequences in a genomic locus using a fragment with homology arms (40 bp) for a CRISPR-Cas9 guided homologous recombination (Ryan et al. 2016). **B.** Strategy for the construction of the motifΔ preys. The NATMX4 cassette was amplified for genomic integration at the locus of interest in reverse orientation (NATMX4r). The prey sequence was used for mutagenesis to replace the PXXP motif with a stuffer and homology arms were added to target the locus of interest. This fragment was used for a CRISPR-Cas9 guided recombination using a gRNA targeting the NATMX4r cassette. **C.** Strategy for the construction of the empty LandingPad yeast strains. Fragments were amplified from genomic DNA and from pAG25-DHFR F[1,2] for fusion PCR to add homology arms (40 bp) for integration at the *GAL1* locus. Homologous recombination transformations were performed to insert the empty LandingPads in the PL0001 background (Table S12). The strategy in panel A is repeated to insert the SH3 sequences at the LandingPad locus.

<u>References</u>

Anisimova M, Gil M, Dufayard J-F, Dessimoz C, Gascuel O. 2011. Survey of branch support
methods demonstrates accuracy, power, and robustness of fast likelihood-based
approximation schemes. *Syst. Biol.* 60:685–699.

Aranda-Díaz A, Mace K, Zuleta I, Harrigan P, El-Samad H. 2017. Robust Synthetic Circuits
for Two-Dimensional Control of Gene Expression in Yeast. *ACS Synth. Biol.* 6:545–554.

Breker M, Gymrek M, Schuldiner M. 2013. A novel single-cell screening platform reveals
proteome plasticity during yeast stress responses. *J. Cell Biol.* 200:839–850.

Brennan P. 2018. drawProteins: a Bioconductor/R package for reproducible and
programmatic generation of protein schematics. *F1000Res.* 7:1105.

Brown T, Brown N, Stollar EJ. 2018. Most yeast SH3 domains bind peptide targets with high
intrinsic specificity. *PLoS One* 13:e0193128.

Castagnoli L, Costantini A, Dall'Armi C, Gonfloni S, Montecchi-Palazzi L, Panni S, Paoluzi S,
Santonico E, Cesareni G. 2004. Selectivity and promiscuity in the interaction network
mediated by protein recognition modules. *FEBS Lett.* 567:74–79.

Cherry JM, Hong EL, Amundsen C, Balakrishnan R, Binkley G, Chan ET, Christie KR,
Costanzo MC, Dwight SS, Engel SR, et al. 2012. Saccharomyces Genome Database:
the genomics resource of budding yeast. *Nucleic Acids Res.* 40:D700–D705.

Chothia C, Lesk AM. 1986. The relation between the divergence of sequence and structure
in proteins. *EMBO J.* 5:823–826.

Dalgarno DC, Botfield MC, Rickles RJ. 1997. SH3 domains and drug design: ligands,
structure, and biological function. *Biopolymers* 43:383–400.

Dionne U, Bourgault É, Dubé AK, Bradley D, Chartier FJM, Dandage R, Dibyachintan S,
Després PC, Gish GD, Pham NTH, et al. 2021. Protein context shapes the specificity of
SH3 domain-mediated interactions in vivo. *Nat. Commun.* 12:1597.

Dionne U, Percival LJ, Chartier FJM, Landry CR, Bisson N. 2022. SRC homology 3
domains: multifaceted binding modules. *Trends Biochem. Sci.* 47:772–784.

Dominguez C, Boelens R, Bonvin AMJJ. 2003. HADDOCK: a protein-protein docking
approach based on biochemical or biophysical information. *J. Am. Chem. Soc.*
125:1731–1737.

Doolittle RF. 1995. The multiplicity of domains in proteins. *Annu. Rev. Biochem.* 64:287–314.

Edgar RC. 2004. MUSCLE: multiple sequence alignment with high accuracy and high
throughput. *Nucleic Acids Res.* 32:1792–1797.

Evangelista M, Klebl BM, Tong AH, Webb BA, Leeuw T, Leberer E, Whiteway M, Thomas
DY, Boone C. 2000. A role for myosin-I in actin assembly through interactions with
Vrp1p, Bee1p, and the Arp2/3 complex. *J. Cell Biol.* 148:353–362.

Freschi L, Torres-Quiroz F, Dubé AK, Landry CR. 2013. qPCA: a scalable assay to measure
the perturbation of protein--protein interactions in living cells. *Mol. Biosyst.* 9:36–43.

Geli MI, Lombardi R, Schmelzl B, Riezman H. 2000. An intact SH3 domain is required for
myosin I-induced actin polymerization. *EMBO J.* 19:4281–4291.

Geng C, Narasimhan S, Rodrigues JPGLM, Bonvin AMJJ. 2017. Information-Driven,
Ensemble Flexible Peptide Docking Using HADDOCK. In: Schueler-Furman O, London
N, editors. Modeling Peptide-Protein Interactions: Methods and Protocols. New York,
NY: Springer New York. p. 109–138.

Goodson HV, Spudich JA. 1995. Identification and molecular characterization of a yeast
myosin I. *Cell Motil. Cytoskeleton* 30:73–84.

Janz JM, Sakmar TP, Christopher Min K. 2007. A Novel Interaction between Atrophin-
interacting Protein 4 and Î²-p21-activated Kinase-interactive Exchange Factor Is
Mediated by an SH3 Domain. *J. Biol. Chem.* 282:28893–28903.

Jumper J, Evans R, Pritzel A, Green T, Figurnov M, Ronneberger O, Tunyasuvunakool K,
Bates R, Žídek A, Potapenko A, et al. 2021. Highly accurate protein structure prediction
with AlphaFold. *Nature* 596:583–589.

Kamrad S, Rodríguez-López M, Cotobal C, Correia-Melo C, Bähler J. 2020. Pyphe, a python
toolbox for assessing microbial growth and cell viability in high-throughput colony
screens. Available from:
https://www.researchgate.net/publication/342228895_Pyphe_a_python_toolbox_for_ass
essing_microbial_growth_and_cell_viability_in_high-throughput_colony_screens

Katoh K, Kuma K-I, Toh H, Miyata T. 2005. MAFFT version 5: improvement in accuracy of
multiple sequence alignment. *Nucleic Acids Res.* 33:511–518.

Kay BK, Williamson MP, Sudol M. 2000. The importance of being proline: the interaction of
proline-rich motifs in signaling proteins with their cognate domains. *FASEB J.* 14:231–
241.

Kel AE, Gössling E, Reuter I, Cheremushkin E, Kel-Margoulis OV, Wingender E. 2003.
MATCH: A tool for searching transcription factor binding sites in DNA sequences.
Nucleic Acids Res. 31:3576–3579.

Kumar S, Stecher G, Suleski M, Hedges SB. 2017. TimeTree: A Resource for Timelines,
Timetrees, and Divergence Times. *Mol. Biol. Evol.* 34:1812–1819.

Kuriyan J, Cowburn D. 1997. Modular peptide recognition domains in eukaryotic signaling.
Annu. Rev. Biophys. Biomol. Struct. 26:259–288.

Lechler T, Shevchenko A, Li R. 2000. Direct involvement of yeast type I myosins in Cdc42-
dependent actin polymerization. *J. Cell Biol.* 148:363–373.

Letunic I, Bork P. 2021. Interactive Tree Of Life (iTOL) v5: an online tool for phylogenetic
tree display and annotation. *Nucleic Acids Res.* 49:W293–W296.

Letunic I, Khedkar S, Bork P. 2021. SMART: recent updates, new developments and status
in 2020. *Nucleic Acids Res.* 49:D458–D460.

Levy ED, Kowarzyk J, Michnick SW. 2014. High-resolution mapping of protein concentration
reveals principles of proteome architecture and adaptation. *Cell Rep.* 7:1333–1340.

Marcet-Houben M, Gabaldón T. 2015. Beyond the Whole-Genome Duplication: Phylogenetic
Evidence for an Ancient Interspecies Hybridization in the Baker's Yeast Lineage. *PLoS
Biol.* 13:e1002220.

Melero C, Ollikainen N, Harwood I, Karpiak J, Kortemme T. 2014. Quantification of the
transferability of a designed protein specificity switch reveals extensive epistasis in

molecular recognition. *Proc. Natl. Acad. Sci. U. S. A.* 111:15426–15431.

Minh BQ, Schmidt HA, Chernomor O, Schrempf D, Woodhams MD, von Haeseler A, Lanfear R. 2020. IQ-TREE 2: New Models and Efficient Methods for Phylogenetic Inference in the Genomic Era. *Mol. Biol. Evol.* 37:1530–1534.

Mochida J, Yamamoto T, Fujimura-Kamada K, Tanaka K. 2002. The novel adaptor protein, Mti1p, and Vrp1p, a homolog of Wiskott-Aldrich syndrome protein-interacting protein (WIP), may antagonistically regulate type I myosins in Saccharomyces cerevisiae. *Genetics* 160:923–934.

Nelson RJ, Ziegelhoffer T, Nicolet C, Werner-Washburne M, Craig EA. 1992. The translation machinery and 70 kd heat shock protein cooperate in protein synthesis. *Cell* 71:97–105.

Pettersen EF, Goddard TD, Huang CC, Meng EC, Couch GS, Croll TI, Morris JH, Ferrin TE. 2021. UCSF ChimeraX: Structure visualization for researchers, educators, and developers. *Protein Sci.* 30:70–82.

Pupko T, Pe'er I, Shamir R, Graur D. 2000. A fast algorithm for joint reconstruction of ancestral amino acid sequences. *Mol. Biol. Evol.* 17:890–896.

Remy I, Michnick SW. 2007. Application of protein-fragment complementation assays in cell biology. *Biotechniques* 42:137, 139, 141 passim.

Rochette S, Diss G, Filteau M, Leducq J-B, Dubé AK, Landry CR. 2015. Genome-wide protein-protein interaction screening by protein-fragment complementation assay (PCA) in living cells. *J. Vis. Exp.* [Internet]. Available from: http://dx.doi.org/10.3791/52255

Ryan OW, Poddar S, Cate JHD. 2016. CRISPR-Cas9 Genome Engineering in Saccharomyces cerevisiae Cells. *Cold Spring Harb. Protoc.* [Internet] 2016. Available from: http://dx.doi.org/10.1101/pdb.prot086827

Schoch CL, Ciufo S, Domrachev M, Hotton CL, Kannan S, Khovanskaya R, Leipe D, Mcveigh R, O'Neill K, Robbertse B, et al. 2020. NCBI Taxonomy: a comprehensive update on curation, resources and tools. *Database* [Internet] 2020. Available from: http://dx.doi.org/10.1093/database/baaa062

Schymkowitz J, Borg J, Stricher F, Nys R, Rousseau F, Serrano L. 2005. The FoldX web server: an online force field. *Nucleic Acids Res.* 33:W382–W388.

Shannon P, Markiel A, Ozier O, Baliga NS, Wang JT, Ramage D, Amin N, Schwikowski B, Ideker T. 2003. Cytoscape: a software environment for integrated models of biomolecular interaction networks. *Genome Res.* 13:2498–2504.

Stark C, Breitkreutz B-J, Reguly T, Boucher L, Breitkreutz A, Tyers M. 2006. BioGRID: a general repository for interaction datasets. *Nucleic Acids Res.* 34:D535–D539.

Tarassov K, Messier V, Landry CR, Radinovic S, Serna Molina MM, Shames I, Malitskaya Y, Vogel J, Bussey H, Michnick SW. 2008. An in vivo map of the yeast protein interactome. *Science* 320:1465–1470.

Teyra J, Huang H, Jain S, Guan X, Dong A, Liu Y, Tempel W, Min J, Tong Y, Kim PM, et al. 2017. Comprehensive Analysis of the Human SH3 Domain Family Reveals a Wide Variety of Non-canonical Specificities. *Structure* 25:1598–1610.e3.

Tonikian R, Xin X, Toret CP, Gfeller D, Landgraf C, Panni S, Paoluzi S, Castagnoli L, Currell B, Seshagiri S, et al. 2009. Bayesian modeling of the yeast SH3 domain interactome

predicts spatiotemporal dynamics of endocytosis proteins. *PLoS Biol.* 7:e1000218.

Tsaban T, Varga JK, Avraham O, Ben-Aharon Z, Khramushin A, Schueler-Furman O. 2022. Harnessing protein folding neural networks for peptide-protein docking. *Nat. Commun.* 13:176.

Usmanova DR, Bogatyreva NS, Ariño Bernad J, Eremina AA, Gorshkova AA, Kanevskiy GM, Lonishin LR, Meister AV, Yakupova AG, Kondrashov FA, et al. 2018. Self-consistency test reveals systematic bias in programs for prediction change of stability upon mutation. *Bioinformatics* 34:3653–3658.

Vilella AJ, Severin J, Ureta-Vidal A, Heng L, Durbin R, Birney E. 2009. EnsemblCompara GeneTrees: Complete, duplication-aware phylogenetic trees in vertebrates. *Genome Res.* 19:327–335.

Vishwanath S, de Brevern AG, Srinivasan N. 2018. Same but not alike: Structure, flexibility and energetics of domains in multi-domain proteins are influenced by the presence of other domains. *PLoS Comput. Biol.* 14:e1006008.

Vogel C, Chothia C. 2006. Protein family expansions and biological complexity. *PLoS Comput. Biol.* 2:e48.

Werner-Washburne M, Stone DE, Craig EA. 1987. Complex interactions among members of an essential subfamily of hsp70 genes in Saccharomyces cerevisiae. *Mol. Cell. Biol.* 7:2568–2577.

Wolfe KH. 2015. Origin of the Yeast Whole-Genome Duplication. *PLoS Biol.* 13:e1002221.

Wolfe KH, Shields DC. 1997. Molecular evidence for an ancient duplication of the entire yeast genome. *Nature* 387:708–713.

Wu C, Ma MH, Brown KR, Geisler M, Li L, Tzeng E, Jia CYH, Jurisica I, Li SS-C. 2007. Systematic identification of SH3 domain-mediated human protein-protein interactions by peptide array target screening. *Proteomics* 7:1775–1785.

Xin X, Gfeller D, Cheng J, Tonikian R, Sun L, Guo A, Lopez L, Pavlenco A, Akintobi A, Zhang Y, et al. 2013. SH3 interactome conserves general function over specific form. *Mol. Syst. Biol.* 9:652.

Zarrinpar A, Park S-H, Lim WA. 2003. Optimization of specificity in a cellular protein interaction network by negative selection. *Nature* 426:676–680.

Zhou L, Feng T, Xu S, Gao F, Lam TT, Wang Q, Wu T, Huang H, Zhan L, Li L, et al. 2022. ggmsa: a visual exploration tool for multiple sequence alignment and associated data. *Brief. Bioinform.* [Internet] 23. Available from: http://dx.doi.org/10.1093/bib/bbac222