The Complexity of Functional Connectivity Profiles of the Subgenual Anterior Cingulate Cortex and Dorsal Lateral Prefrontal Cortex in Major Depressive Disorder: a DIRECT Consortium Study

Xiao Chen[1,2,3,4,5], Bin Lu[1,2,3,4], Yu-Wei Wang[1,2,3,4], Xue-Ying Li[1,2,3,4], Zi-Han Wang[1,2,3,4], Hui-Xian Li[1,2,3,4], Yi-Fan Liao[1,2,3,4], Daniel M. Blumberger[5,6], Francisco Xavier Castellanos[7,8], Li-Ping Cao[9], Guan-Mao Chen[10], Jian-Shan Chen[9], Tao Chen[11], Tao-Lin Chen[12,13], Yu-Qi Cheng[14], Zhao-Song Chu[14], Shi-Xian Cui[1,15,16], Xi-Long Cui[17], Zhao-Yu Deng[1,2,3,4], Qi-Yong Gong[12,13], Wen-Bin Guo[17], Can-Can He[18], Zheng-Jia-Yi Hu[1,15,16], Qian Huang[19], Xin-Lei Ji[17], Feng-Nan Jia[20], Li Kuang[19], Bao-Juan Li[21], Feng Li[22], Tao Li[23,24], Tao Lian[1,2,3], Xiao-Yun Liu[25], Yan-Song Liu[20], Zhe-Ning Liu[17], Yi-Cheng Long[17], Jian-Ping Lu[26], Jiang Qiu[27], Xiao-Xiao Shan[17], Tian-Mei Si[28], Peng-Feng Sun[29], Chuan-Yue Wang[22], Hua-Ning Wang[21], Xiang Wang[17], Ying Wang[10], Xiao-Ping Wu[29], Xin-Ran Wu[27], Yan-Kun Wu[28], Chun-Ming Xie[18], Guang-Rong Xie[17], Peng Xie[30,31,32], Xiu-Feng Xu[14], Zhen-Peng Xue[26], Hong Yang[11], Jian Yang[31], Hua Yu[23,24], Yong-Qiang Yu[33,34,35], Min-Lan Yuan[36], Yong-Gui Yuan[25], Yu-Feng Zang[37,38], Ai-Xia Zhang[39], Ke-Rang Zhang[39], Wei Zhang[36], Zi-Jing Zhang[1,2,3], Jing-Ping Zhao[17], Jia-Jia Zhu[33,34,35], Xi-Nian Zuo[40,41], the DIRECT Consortium[#], Chao-Gan Yan[1,2,3,4,15,16]*

[1]CAS Key Laboratory of Behavioral Science, Institute of Psychology, Beijing 100101, China

[2]International Big-Data Center for Depression Research, Chinese Academy of Sciences, Beijing 100101, China

[3]Magnetic Resonance Imaging Research Center, Institute of Psychology, Chinese Academy of Sciences, Beijing 100101, China

[4]Department of Psychology, University of Chinese Academy of Sciences, Beijing 100049, China

[5]Temerty Centre for Therapeutic Brain Intervention, Campbell Family Research Institute, Centre for Addiction and Mental Health, Toronto M6J1H4, Ontario, Canada

[6]Department of Psychiatry, Temerty Faculty of Medicine, University of Toronto, Toronto M5S1A1,

Ontario, Canada

[7]Department of Child and Adolescent Psychiatry, NYU Grossman School of Medicine, New York, NY, USA

[8]Nathan Kline Institute for Psychiatric Research, Orangeburg, NY, USA

[9]Affiliated Brain Hospital of Guangzhou Medical University, Guangzhou 510370, China

[10]The First Affiliated Hospital of Jinan University, Guangzhou, Guangdong 250024, China

[11]Department of Radiology, The First Affiliated Hospital, College of Medicine, Zhejiang University,

Hangzhou, Zhejiang 310058, China

[12]Huaxi MR Research Center (HMRRC), Department of Radiology, West China Hospital of Sichuan

University, Chengdu, Sichuan 610044, China

[13]Research Unit of Psychoradiology, Chinese Academy of Medical Sciences, Chengdu, Sichuan 610052,

China

[14]Department of Psychiatry, First Affiliated Hospital of Kunming Medical University, Kunming, Yunnan

650032, China

[15]Sino-Danish College, University of Chinese Academy of Sciences, Beijing 101408, China

[16]Sino-Danish Center for Education and Research, Graduate University of Chinese Academy of Sciences,

Beijing 101408, China

[17]Department of Psychiatry, and National Clinical Research Center for Mental Disorders, The Second

Xiangya Hospital of Central South University, Changsha 410011, Hunan, China

[18]Department of Neurology, Affiliated ZhongDa Hospital of Southeast University, Nanjing, Jiangsu

210009, China

[19]Department of Psychiatry, The First Affiliated Hospital of Chongqing Medical University, Chongqing

400042, China

[20]Department of Clinical Psychology, Suzhou Psychiatric Hospital, The Affiliated Guangji Hospital of

Soochow University, Suzhou, Jiangsu 215003, China

[21]Xijing Hospital of Air Force Military Medical University, Xi'an, Shaanxi 710032, China

[22]Beijing Anding Hospital, Capital Medical University, Beijing 100120, China

[23]Affiliated Mental Health Center & Hangzhou Seventh People's Hospital, Zhejiang University School of

Medicine, Hangzhou, Zhejiang 310063, China

[24]Mental Health Center and Psychiatric Laboratory, West China Hospital of Sichuan University, Chengdu,

Sichuan 610044, China

[25]Department of Psychosomatics and Psychiatry, Zhongda Hospital, School of Medicine, Southeast University, Nanjing, Jiangsu 210009, China

[26]Shenzhen Kangning Hospital, Shenzhen, Guangzhou 518020, China

[27]Faculty of Psychology, Southwest University, Chongqing 400715, China

[28]National Clinical Research Center for Mental Disorders (Peking University Sixth Hospital) & Key Laboratory of Mental Health, Ministry of Health (Peking University), Beijing 100191, China

[29]Xi'an Central Hospital, Xi'an, Shaanxi 710004, China

[30]Institute of Neuroscience, Chongqing Medical University, Chongqing 400016, China

[31]Chongqing Key Laboratory of Neurobiology, Chongqing 400000, China

[32]Department of Neurology, The First Affiliated Hospital of Chongqing Medical University, Chongqing 400042, China

[33]Department of Radiology, The First Affiliated Hospital of Anhui Medical University, Hefei 230022, China

[34]Research Center of Clinical Medical Imaging, Anhui Province, Hefei 230032, China

[35]Anhui Provincial Institute of Translational Medicine, Hefei 230032, China

[36]West China Hospital of Sichuan University, Chengdu, Sichuan 610044, China

[37]Center for Cognition and Brain Disorders, The Affiliated Hospital of Hangzhou Normal University, Hangzhou, Zhejiang 310018, China

[38]Zhejiang Key Laboratory for Research in Assessment of Cognitive Impairments, Hangzhou, Zhejiang 310000, China

[39]First Hospital of Shanxi Medical University, Taiyuan, Shanxi 030001, China

[40]Developmental Population Neuroscience Research Center, IDG/McGovern Institute for Brain Research, Beijing Normal University, Beijing 100091, China

[41]National Basic Science Data Center, Beijing 100038, China

[42]Department of Psychiatry, Sir Run Run Shaw Hospital, Zhejiang University School of Medicine, Hangzhou 310020, Zhejiang, China

[43]Shanghai Mental Health Center, Shanghai Jiao Tong University School of Medicine, Shanghai 200030, China

[44]Department of Diagnostics, Affiliated Hospital, Hangzhou Normal University Medical School,

Hangzhou, Zhejiang 311121, China

[45]Department of Psychiatry, First Affiliated Hospital, China Medical University, Shenyang, Liaoning

110122, China

[46]Early Intervention Unit, Department of Psychiatry, Affiliated Nanjing Brain Hospital, Nanjing Medical

University, Nanjing 210024, China

[47]Department of Neurology, The First Affiliated Hospital of Anhui Medical University, Hefei, Anhui

230022, China

[48]Department of Psychiatry, Shanghai Jiao Tong University School of Medicine, Shanghai 200025, China

*Corresponding author:

Chao-Gan Yan, Ph. D.

CAS Key Laboratory of Behavioral Science, Institute of Psychology, Beijing, China

16 Lincui Road, Chaoyang District, Beijing 100101, China

Tel: +86-10-64101582

E-mail: yancg@psych.ac.cn

[#]DIRECT consortium: Chao-Gan Yan, Ph.D.[1,2,3,4,15,16], Xiao Chen, Ph.D.[1,2,3,4,5], Li-Ping Cao, M.D.[9], Wei

Chen, M.D.[42], Yu-Qi Cheng, M.D.[14], Yi-Ru Fang, M.D.[43], Qi-Yong Gong, M.D.[12,13], Wen-Bin Guo, M.D.[17], Li

Kuang, M.D.[19], Bao-Juan Li, M.D.[21], Tao Li, M.D.[23,24], Yan-Song Liu, M.D.[20], Zhe-Ning Liu, M.D.[17], Jian-Ping

Lu, M.D.[26], Qing-Hua Luo, M.D.[19], Hua-Qing Meng, M.D.[19], Dai-Hui Peng, M.D.[43], Jiang Qiu, Ph.D.[27], Yue-

Di Shen, M.D.[44], Tian-Mei Si, M.D.[28], Yan-Qing Tang, M.D.[45], Chuan-Yue Wang, M.D.[22], Fei Wang, M.D.[45,46],

Hua-Ning Wang, M.D.[21], Kai Wang, M.D.[47], Xiang Wang, M.D.[17], Ying Wang, M.D.[10], Xiao-Ping Wu, M.D.[29],

Chun-Ming Xie, M.D.[18], Guang-Rong Xie, M.D.[17], Peng Xie, M.D.[30,31,32], Xiu-Feng Xu, M.D.[14], Hong Yang,

M.D.[11], Jian Yang, M.D.[31], Shu-Qiao Yao, M.D.[17], Yong-Qiang Yu, M.D.[33,34,35], Yong-Gui Yuan, M.D.[25], Ke-

Rang Zhang, M.D.[39], Wei Zhang, M.D.[36], Zhi-Jun Zhang, M.D.[18], Jun-Juan Zhu, M.D.[48], Xi-Nian Zuo,

Ph.D.[40,41], Jing-Ping Zhao, M.D.[17], Yu-Feng Zang, M.D.[37,38]

Abstract

The subgenual anterior cingulate cortex (sgACC) appears to play a central role in the pathophysiology of major depressive disorder (MDD). To wit, its functional interactive profile with the left dorsal lateral prefrontal cortex (DLPFC) has been shown to be related to treatment outcomes with transcranial magnetic stimulation (TMS) treatment outcomes. Nevertheless, previous research on sgACC functional connectivity (FC) in MDD has yielded inconsistent results, partly due to small sample sizes and limited statistical power of prior work. Here, leveraging a large multi-site sample (1660 MDD patients vs. 1341 healthy controls) from Phase II of the Depression Imaging REsearch ConsorTium (DIRECT), we systematically delineated case-control difference maps of sgACC FC and examined their clinical relevance to previously identified TMS targets. We also investigated case-control FC difference maps of left DLPFC sub-fields. In MDD patients we found significantly increased FC between sgACC and thalamus and reduced FC to a broad array of brain regions, including somatosensory area, occipital lobe, medial and lateral temporal lobe, and insular cortex, when global signal regression (GSR) was not implemented. Intriguingly, we found enhanced left DLPFC-sgACC FC in MDD patients when GSR was performed. We leveraged an prior independent sample to explore the possible relationship between the case-control differences regarding sgACC's FC profiles and the treatment out comes of TMS. In sites in which open TMS treatment was administered, case-control differences in sgACC FC, with GSR, were related to clinical improvement. Next we tested whether the position of peak of the FC maps (previously identified TMS target) could be altered in MDD patients as compred with healthy controls (HC)s. We found the optimized TMS target differed in MDD patients. Several DLPFC sub-fields yielded case-control differences in whole-brain FC maps. In summary, we reliably delineated MDD-related abnormalities of sgACC FC profiles in a large sample. GSR was essential in applying case-control difference maps to identify optimized TMS targets. Our results highlight the functional heterogeneity of the left DLPFC and of precise TMS targets therein.

Keywords: major depressive disorder, transcranial magnetic stimulation, functional connectivity, global signal regression

Introduction

Major depressive disorder (MDD) is a common and debilitating psychiatric disorder projected

to be the most burdensome worldwide within this decade (1). Despite extensive research,

the pathophysiology of MDD remains elusive. Nevertheless, a key locus appears to be the

subgenual anterior cingulate cortex (sgACC), which shows reproducible metabolic

hyperactivity (2), is targeted in deep brain stimulation (3, 4), and has been implicated in

emotional responses, motivation, and rumination (5). Furthermore, identifying an optimized

neuromodulation target in the left dorsal lateral prefrontal cortex (DLPFC) on the basis of

sgACC-related functional connectivity (FC) and understanding MDD-related functional

abnormalities in DLPFC sub-fields are considered crucial for developing effective depression

treatments (6, 7). However, given inconsistent results from small sample-sized studies (8-14),

we set out to establish a large sample in which we could seek to identify reliable abnormalities

in sgACC and DLPFC FC and better understand the potential impact of such FC profiles on

neuromodulation target identification for treating depression.

Multiple studies have investigated FC abnormalities in MDD using resting-state functional

magnetic resonance imaging (rsfMRI). Abnormal FC between sgACC and amygdala, thalamus,

temporal gyrus, lingual gyrus, cerebellum, dorsal lateral prefrontal cortex (DLPFC), and default

mode network (DMN) regions such as medial and dorsal medial prefrontal cortex, precuneus,

and parahippocampus have been reported (8, 9, 12-19). However, results have been highly

inconsistent, making it challenging to integrate findings and generate clear profiles of sgACC-

related FC abnormalities. Such lack of reproducibility could be partially due to small sample

size and low statistical power of clinical imaging studies (20, 21). Another possible reason is

the implementation of global signal regression (GSR). GSR is the removal of the mean signal

averaged across the entire brain via linear regression and one of the most controvirtial

preprocessing steps in rsfMRI (22, 23). On the one hand, GSR can increase the specificity of FC

by efficiently reducing the impact of non-neuronal nuisance variables, such as head motion

and respiration (24, 25). On the other hand, GSR has been criticized for artificially shifting the

distribution of the originally derived FCs and introducing spurious anti-correlations (26).

Accordingly, not all previous studies implemented GSR, which might significantly alter case-

control comparison results (27).

Maps of sgACC-related FC abnormalities may be clinically useful for predicting repetitive

transcranial magnetic stimulation (rTMS) treatment outcomes (28-30). Specifically, the anti-

correlation between sgACC and left DLPFC has been associated with clinical improvement after

receiving rTMS treatment (31-35). This has led to the intriguing notion that the FC between

sgACC and left DLPFC could be leveraged to identify more precise rTMS targets and improve

the efficacy of rTMS delivered to the left DLPFCr (6, 36). Indeed, a group-wise sgACC target has

been identified (33), and several individualized FC-guided TMS target identification algorithms

have been proposed following this rationale (36-39). It is worth noting that the existing sgACC

group target was based on the functional connectome of a large cohort of healthy adults.

Subtle but significant case-control differences in resting-state FC profiles have been identified

in a large sample of MDD patients (40). Thus, understanding the profiles of sgACC FC map case-

control differences and their impact on potential targets of rTMS to the left DLPFC could be a

critical step toward developing optimized rTMS site identification methods.

The left DLPFC is an anatomically extensive region consisting primarily of two Brodmann areas

(BA 9 and BA 46) (41). Several rTMS targeting approaches have been implemented in the

literature, including 1) scalp-based measures ("5 cm" method and F3 Beam method); 2)

anatomical targeting (leveraging neuro-navigation to target an anatomical coordinate in BA 9

or BA 46); and 3) group-wise connectivity-based targets (the anti-correlation peak of the

group-wise mean FC map) (6, 33, 42-44). These methods have generated multiple TMS target

sites, with the result that the location of an optimal left DLPFC TMS target remains unsettled.

Evaluating case-control differences of common rTMS sites' FC maps might help delineate the

MDD-related functional abnormalities of various DLPFC sub-fields.

To address the issue of sample-size, we initiated the Depression Imaging REsearch ConsorTium

(DIRECT) (45) and and carried out an initial meta/mega-analysis (N_{MDD} = 1300), referred to as

REST-meta-MDD (40). Studies from the initially shared DIRECT data revealed multiple MDD-

related abnormalities including in network FC, whole-brain FC topological and dynamic

features, and functional lateralization (40, 46-52). The DIRECT Phase I neuroimaging data were

preprocessed using a volume-based approach (DPARSF (53, 54)), producing voxel-wise BOLD

time series at each participating hospital which were then shared. In the second phase of

DIRECT, we pooled a larger MDD sample (N_{MDD} = 1660) which was preprocessed with a state-

of-the-art pipeline, DPABISurf (55). DIRECT Phase II will share voxel/vertex level BOLD time

series, allowing more flexible and thorough investigations. The aim of the current study is

threefold, to: 1) leverage this highly powered sample to derive a reliable sgACC-related FC

abnormality profile in MDD; 2) evaluate the potential impact of such an abnormality on rTMS

target identification and explore its potential clinical usage; 3) delineate MDD-related FC

abnormalities of commonly used rTMS localization methods within the left DLPFC.

Materials and Methods

Study sample

The DIRECT Phase II initiative comprises 23 cohorts and includes rsfMRI and T1 structural scans

from 1660 MDD patients and 1341 HCs. Demographic and clinical characteristics for each

sample are presented in Figure 1. Site information, sample size, and previous publications

based on the shared data are listed in Table S1. Patients were diagnosed with MDD based on

ICD 10 or DSM-IV. All participants provided written informed consent, and local institutional

review boards approved each study from all included cohorts.

Image preprocessing

Acquisition parameters and scanners for all cohorts are provided in Table S2. All rsfMRI and

structural MRI scans were preprocessed at each site using the same DPABISurf protocol, an

rsfMRI data analysis toolbox evolved from DPABI/DPARSF (53-55) (For details, see SI).

Computation of sgACC and DLPFC FC maps

Although recent studies have attempted to identify personalized TMS targets using surface-

based algorithms (37), most previous studies have reported targets in volume-based MNI

space with sgACC ROIs defined as a sphere in volume-based space (31, 56). To better compare

our results with the existing literature, we used the volume-based preprocessed imaging data

from DPABISurf.

Because no consensus has been achieved on the anatomical definition of the sgACC, we tested

two sets of definitions: (1) a 10 mm sphere located on the average MNI coordinates based on

prior studies showing reduced glucose metabolism or blood flow after receiving an

antidepressant treatment (MNI coordinates: x = 6, y = 16, z = -10. For details, please refer to

Fox et al., 2012 (31).); and (2) a mask denoting the anatomical boundary of BA 25 (57).

Two BA areas in the DLPFC, BA9 and BA46, have been proposed to correspond to different FC

profiles. As a result, we defined a priori ROIs by placing 20 mm spheres centered along the left

hemisphere at BA9 (MNI coordinates: x = -36, y = 39, z = 43) and BA46 (MNI coordinates: x = -

44, y = 40, z = 29) (6). Additionally, we evaluated several TMS targets from the literature: 1)

the "5 cm" method (MNI coordinates: x = -41, y = 16, z = 54) (36); 2) the F3 Beam group-

average target (MNI coordinates: x = −37, y = 26, z = 49) (29); and 3) the sgACC group target

(MNI coordinates: x = −42, y = 44, z = 30) (33).

The time series of sgACC and DLPFC were determined for each individual by spatially averaging

the preprocessed rsfMRI time series across all voxels in the above-mentioned masks . We then

calculated whole-brain FC maps in volume-based MNI space. FC was calculated using Pearson's

correlation and underwent Fisher's r-to-z transformation. All FC maps were smoothed with a

6 mm full-width half maximum (FWHM) kernel size.

Harmonization of batch effects

We used ComBat (58) to control potential site and scanner biases. ComBat models site-specific scaling factors by introducing terms for biological variables and scanners in a multivariate linear mixed effects regression. By employing empirical Bayesian criteria, it improves the estimation of studies with small sample sizes. ComBat assumes that the expected FC value can be modeled as a linear combination of batch effects and biological effects. Specifically, the error term in this model is modulated by additional site-specific scaling factors. Let y denote a given FC value:

$$y = \alpha + X^T\beta + \gamma + \delta\varepsilon \tag{1}$$

where $\boldsymbol{\alpha}$ denotes the constant term; X^T denotes the design matrix for covariates of interest (diagnosis, age, sex, education, head motion); $\boldsymbol{\beta}$ is a vector of regression coefficients corresponding to X; $\boldsymbol{\gamma}$ stands for addictive site effects (location parameter) while $\boldsymbol{\delta}$ represents multiplicative site effects (scale parameter). Accordingly, harmonized FC values were defined as:

$$y^{ComBat} = \frac{y - \hat{\alpha} - X^T\hat{\beta} - \hat{\gamma}}{\hat{\delta}} + \alpha + X^T\hat{\beta} \tag{2}$$

where $\hat{\delta}$ and $\hat{\gamma}$ stand for the empirical Bayesian estimates of $\boldsymbol{\delta}$ and $\boldsymbol{\gamma}$, respectively. y^ComBat was calculated with the "yw_Harmonization" function in DPABISurf (http://rfmri.org/DPABISurf). ComBat models biological and non-biological effects and algebraically removes the estimated additive and multiplicative batch effects.

Group difference maps

We used a generalized linear model (GLM) to examine differences in the FC maps of sgACC and DLPFC seeds between MDD patients and HCs. Cohen's f^2 was also calculated to characterize effect sizes of this group difference effect. The GLM model includes age, sex, education, and head motion as covariates:

$$y = \mu + X\beta + \epsilon \tag{3}$$

where y denotes the FC value of a given voxel from a given participant; $\boldsymbol{\mu}$ stands for the

275 constant term; X represents the design matrix for the covariates of interest (diagnosis, age, sex,

276 education, and head motion); $\boldsymbol{\beta}$ is a vector of regression coefficients corresponding to X; and

277 $\boldsymbol{\epsilon}$ is a vector of residuals that follow $N(0, \sigma^2)$. Multiple comparison correction was conducted

278 using false discovery rate (FDR) correction at $q < 0.05$.

Group difference targets

We first defined a left DLPFC ROI by combining four 20 mm radius spheres centered in BA9

(MNI coordinates: x = -36, y = 39, z = 43), BA46 (MNI coordinates: x = -44, y = 40, z = 29), the

"5 cm" method site (MNI coordinates: x = -44, y = 40, z = 29), and the F3 Beam site (MNI

coordinates: x = −37, y = 26, z = 49) (37). Then we identified the continuous clusters that

survived FDR correction within this DLPFC ROI and used the peaks of these clusters as the

"group difference target."

Correlation between group differences and clinical efficiency

An openly shared dataset was leveraged to explore the clinical relevance of our present

findings (33). This dataset comprises 25 medication resistant MDD patients who underwent 4

to 7 weeks of daily repetitive TMS applied over the left DLPFC. Patients' TMS sites were

recorded using their structural MRI images and a frameless neuronavigation system.

Treatment response was assessed with the 24-item Hamilton Depression Rating Scale (HAMD).

We only used the shared MNI coordinates of TMS sites and the corresponding reductions in

HAMD scores from each patient. For more details of this dataset, please refer to Weigand et

al. (33). We first extracted the mean t-values of 4 mm radius spheres centered on the TMS

sites, then examined the Pearson correlations between these t-values and HAMD score

reductions.

sgACC group target in MDD and HC

Since the existing sgACC group target was identified using a cohort of healthy young adults (33)

and the significant FC profile differences between MDD and HC (59), we searched for the peak

sgACC anti-correlation within the abovementioned DLPFC ROIs in both MDD and HC groups.

We first averaged whole-brain FC maps across all MDD patients and HCs, then simply searched

for the peak anticorrelated voxel to define the putative target site.

Clustering of DLPFC group difference FC maps

Given the similarities among some DLPFC ROI group difference FC maps, we sought to identify

common clusters using Ward's hierarchical clustering method (60). Specifically, spatial

correlations were calculated between each pair of group difference t maps of DLPFC FC profiles,

with the maps that showed the largest similarity grouped into clusters. We evaluated the

strength of clustering with the gap statistic (61).

Network-wise group differences

To further interpret group difference maps, we extracted the mean FC values of seven

networks using Schaefer's 400 parcellation atlas (62). A GLM model identical to model (3) was

constructed to characterize case-control differences for each network. Bonferroni multiple

comparison correction was conducted ($p < 0.05/7$). All codes have been made openly available

(https://github.com/Chaogan-Yan/PaperScripts/tree/master/Chen_2023). Image results were

rendered on the brain surface using BrainNet Viewer (63). Statistics represented in graphical

format were generated in R GNU v4.2.1 using the" ggplot" package (64, 65).

Results

Group difference maps of sgACC FC

We found significant MDD-related hyperconnectivity with the sgACC in bilateral DLPFC,

temporal parietal junction and occipital lobe, as well as hypoconnectivity in bilateral temporal

lobe, left inferior frontal gyrus and left postcentral gyrus when preprocessing included GSR.

When GSR was not included in preprocessing, MDD-related sgACC FC alterations showed a

predominance of hypoconnectivity. Abnormally decreased FCs were revealed across the

central gyrus, occipital lobe, insular cortex, temporal lobe and a small portion of the frontal

lobe. Without GSR, MDD-related hyperconnectivity was limited to subcortical regions (Figure

2A). Of note, these results were obtained when the sgACC was defined as a sphere. When a

more subtle, anatomically-based definition (a BA 25 mask) was used, no significant MDD-

related FC abnormality was detected when GSR was performed. When GSR was not included

in preprocessing, hypoconnectivity predominated across the brain, except for subcortical

regions, similar to results for the spherical mask (Figure S1). Network-wise FC analyses showed

that MDD patients' FC between sgACC and the limbic network (LN) was significantly reduced

compared to HC (t = -4.357, $p_{corrected}$ < 0.001, Cohen's f^2 = 0.007). Without GSR, MDD patients

showed decreased FC between sgACC and all brain networks (all $p_{corrected}$ < 0.05) except for the

frontoparietal network (FPN) (Figure 2B).

Correlation between group differences and clinical efficiency

Using an openly shared clinical dataset, we examined the relationship between group

differences (t-values) and clinical effects (HAMD score reductions) to test the clinical relevance

of group difference maps. Group differences were positively correlated with HAMD score

reductions (r = 0.448, p = 0.025, Figure 2C).

sgACC anti-correlation peaks in MDD and HCs

The above results highlighted the group differences in sgACC FC profiles between MDD and

HCs. Since the existing sgACC group target was based on a cohort of young, healthy adults, we

sought to examine potential differences of anti-correlation peaks extracted from the mean

sgACC FC maps of MDD and HC groups (Figure 3). We found that the anti-correlation peak of

MDD patients (MNI coordinates: x = -46, y = 4, z = 40) was more posterior compared to HCs

(MNI coordinates: x = -44, y = 38, z = 30), probably due to less anticorrelated FC within the left

DLPFC in MDD patients. The anti-correlation peak extracted from HCs was closer to that

reported in the existing literature (MNI coordinates: x = −42, y = 44, z = 30).

Identifying the group difference target

We identified two contiguous clusters of voxels that showed significant group differences in the left DLPFC. The peaks of these clusters were defined as target coordinates. Group difference target 1 (MNI coordinates: x = -44, y = 38, z = 32; t = 3.277, p < 0.001, Cohen's f2 = 0.004) was ventral to group difference target 2 (MNI coordinates: x = -34, y = 36, z = 40; t = 3.670, p < 0.001, Cohen's f2 = 0.005) (Figure 4).

Group difference maps of different DLPFC ROI FC profiles

When GSR was implemented, FC maps for most DLPFC ROIs showed significant MDD-related abnormalities (FDR q < 0.05) except for Group Difference Target 2 and BA 9 (Figure 4A). Although each ROIs' MDD-related abnormality map was unique, these MDD-related abnormality maps demonstrated one of three general topographic patterns (Figure 4B). The gap statistic quantified an optimal three-cluster solution (Figure S2). Cluster 1 included Group Difference Target 1, BA 46, and the sgACC group target. MDD patients showed enhanced FCs between the medial prefrontal cortex, posterior cingulate cortex, temporal parietal junction, and lateral temporal cortex and decreased FCs between the right inferior prefrontal cortex and the DLPFC ROIs. Cluster 2 consisted of Group Difference Target 2 and BA 9. No significant MDD-related abnormalities were revealed. Nevertheless, enhanced FC between the medial orbitofrontal cortex and occipital cortex were observed when a more liberal threshold was implemented (Gaussian random field correction: voxel-wise p < 0.01, cluster-wise p < 0.05, two-tailed). Cluster 3 included the F3 Beam and the "5 cm" Method ROIs. For these, we found enhanced FCs between the dorsal anterior cingulate cortex, insular, and occipital cortex, and reduced FCs between the lateral temporal cortex and posterior cingulate cortex and the DLPFC ROIs in MDD patients compared with HCs.

Group differences of network-wise FC of different DLPFC ROIs

All three clusters showed group differences in distinct network-wise FC patterns (Figure 6). MDD patients showed enhanced FCs between the Default Mode Network (DMN) and three

Cluster 1 ROIs (Group Difference Target 1: $t = 3.483$, $p_{corrected} = 0.004$, Cohen's $f^2 = 0.004$; BA46:

$t = 3.791$, $p_{corrected} = 0.001$, Cohen's $f^2 = 0.005$; sgACC group target: $t = 3.810$, $p_{corrected} = 0.001$,

Cohen's $f^2 = 0.005$). We only found significantly reduced FC between the Ventral Attention

Network (VAN) and the sgACC group target ($t = -3.057$, $p_{corrected} = 0.016$, Cohen's $f^2 = 0.003$) but

no significant alterations in two other DLPFC ROIs (all $p_{corrected} > 0.05$). DLPFC Cluster 2 ROIs

showed significantly increased LN FC (Group Difference Target 2: $t = 2.753$, $p_{corrected} = 0.042$,

Cohen's $f^2 = 0.003$; $t = 2.897$, $p_{corrected} = 0.027$, Cohen's $f^2 = 0.003$). Only Group Difference Target

2 yielded significantly enhanced somatomotor network (SMN) FC ($t = 2.716$, $p_{corrected} = 0.047$,

Cohen's $f^2 = 0.003$). The network-wise FC profiles of Cluster 3's DLPFC ROIs were generally

similar, but significant FC differences were found in different brain networks. The F3 Beam ROI

showed significantly increased FC to the visual network (VN) ($t = 3.032$, $p_{corrected} = 0.017$,

Cohen's $f^2 = 0.003$) while the "5 cm" Method ROI showed significantly enhanced FCs to the

VAN ($t = 3.512$, $p_{corrected} = 0.003$, Cohen's $f^2 = 0.004$) and reduced FC to the DMN ($t = -2.779$,

$p_{corrected} = 0.038$, Cohen's $f^2 = 0.003$).

Discussion

In the present study, we leveraged a large multi-site sample (1660 MDD patients and 1341 HCs)

and found abnormal sgACC FC in MDD. Specifically, we found reduced FC between sgACC and

a broad array of brain regions when GSR was not conducted, while FC between sgACC and the

left DLPFC was enhanced in MDD patients when GSR was implemented. We identified two

clusters in the left DLPFC which could serve as potential rTMS treatment targets in depression.

In a dataset reporting effects of open rTMS treatment, group differences in sgACC-DLPFC FC

were related to clinical improvement. Interestingly, the locations of anti-correlation peaks

differed substantially in MDD patients compared to HCs. In examining the group difference

maps of various DLPFC ROI FC profiles we found that the DLPFC ROIs could be grouped into

three clusters, which also corresponded to network-wise FCs. These results suggest that group

differences in sgACC FC may be clinically relevant. As the field moves towards identifying

individualized TMS sites, such group differences might be useful for developing next-

generation target-localization algorithms.

MDD-related sgACC FC abnormalities

Our results add to a growing literature showing functional network abnormalities involving the sgACC in MDD (8, 10-12, 16, 18, 66-68). Nevertheless, notable discrepancies in the type of abnormality (enhanced/reduced) and specific brain regions showing altered sgACC FCs have been reported. With an unprecedented sample size, our results have provided the most robust evidence to date. However, the effect sizes of MDD-related sgACC FC abnormalities were small (Cohen's $f^2 < 0.01$). The combination of the limited power of small sample sizes in previous studies and a small effect-sized case-control difference were likely to produce false positive findings (20, 69, 70), which may not be able to be reproduced in the present study. Furthermore, we robustly demonstrated that GSR dramatically changed the group difference maps of sgACC FC profiles. GSR is effective at removing non-neuronal physiological noises such as signal contributions from respiratory (71), but can alter the BOLD signal covariances due to its less specific nature (72). One recent study examined the relationship between the sgACC-DLPFC anti-correlation and the treatment outcomes of TMS using a large dataset (THREE-D) (73) and found that such correlation was largely carried by respiration-related signal patterns (32). Accordingly, most prior studies that intended to identify the TMS targets guided by sgACC-related FCs tend to implement GSR to remove the non-neural noises as much as possible, while other studies that were intended to delineate the general case-control differences prefer not to implement GSR. Here, with an unprecendented large sample, we showed that GSR might induce notable alterations in case-control differences of FC maps (27).

MDD patients showed enhanced FC between sgACC and thalamus regardless of whether GSR was implemented. This result is consistent with previous studies in adolescents (18) and adults (56) with MDD. We found decreased FC between sgACC and the limbic network both with and without GSR. Previous studies have reported abnormal FCs between sgACC and limbic areas and some subcortical regions, such as the amygdala (8, 19) and parahippocampus regions (12). Together, these results are consistent with a model highlighting sgACC as a critical hub in an "extended medial network," which also encompasses limbic, thalamic, and striatal regions and

plays a key role in the pathophysiology of MDD (74, 75). This "extended medial network"

overlaps substantially with the DMN. Indeed, decreased FCs between sgACC and DMN regions,

such as the medial prefrontal cortex, precuneus, temporal gyrus, and parahippocampus

regions, were revealed in MDD relative to HCs when GSR was not implemented. Such

abnormalities have been previously reported (9, 11, 14, 29), which led to the hypothesis that

abnormally enhanced FC between sgACC and DMN are the network underpinnings of

rumination (76). However, inconsistent with the prediction of this model, we found reduced,

instead of enhanced FC between sgACC and DMN. Based on a large multi-site sample of MDD

patients from the first phase of DIRECT, we previously demonstrated that MDD was

characterized by reduced FC within DMN (40, 45). One notable issue is that both the first phase

and the current second phase of the DIRECT data were solely comprised of Chinese samples,

while most studies reporting enhanced sgACC-DMN FCs in MDD recruited Caucasian samples.

Different prevalence rates (77, 78), heterogeneous symptoms (79), and different risk alleles

(80) have been reported in Caucasian and Eastern Asian groups. As a result, racial differences

may have contributed to this discrepancy.

Clinical relevance of MDD-related sgACC FC abnormalities

Although case-control comparisons must be interpreted with caution when GSR has been

implemented, our results support the clinical relevance of such analyses with regard to rTMS

treatment outcomes. Substantial evidence has shown that sgACC FC profiles in MDD patients

are related to rTMS treatment response (30, 81). Our results add to this conceptualization by

showing that the degree of group differences in the sgACC FC profiles were positively related

to the clinical improvements after receiving rTMS to the left DLPFC. Normative connectome

datasets have been used to identify group-wise anti-correlation peaks as TMS targets (31, 33,

36). Here, we found that the locations of such anticorrelated peaks can differ substantially in

a large clinical cohort. Specifically, FC of the more anterior part of the left DLPFC was

abnormally elevated in MDD patients, and thus less anticorrelated to sgACC relative to HCs,

which resulted in a more posterior peak in patients than in HCs.

Interestingly, the thresholded group difference maps revealed two clusters showing enhanced

FC to sgACC in the left DLPFC. Philip and colleagues reported that the FC between the left

DLPFC and sgACC were reduced after receiving 40 daily sessions of 5 Hz TMS to the left DLPFC.

Such a reduction in FC was also related to reduced self-reported symptom scores (30).

Identifying optimum targets in the left DLPFC from multiple clusters is challenging (37).

Spherical ROIs centered at the peaks of these two clusters corresponded to different FC

profiles and group difference maps. Criteria to determine the optimal target might require

more benchmarking using clinical data. Defining a reliable TMS target solely based on one

individual's rsfMRI data (typical duration ~8 mins in clinical practice) is challenging. The

replicability of FC derived from fMRI images is poor (82), and current available individual-level

network parcellation algorithms often require a large number of fMRI images (usually more

than one hour) (83). Given the high reliability of group difference maps derived from a large

sample ($N_{patients}$ > 1500), it might be possible to leverage this map to generate better group

based targets and improve outcomes in the clinical practice.

Different functional network profiles of different DLPFC ROIs

Most clinical trials have focused on applying TMS to the left DLPFC with the belief that high-

frequency rTMS can enhance hypoactivity during depressive episodes (7, 84). The DLPFC is

anatomically extensive but it can be represented by BA 46 and BA 9 (41). It remains unclear

which DLPFC sub-field is the best target for TMS. The current FDA-approved protocol (i.e., "5

cm" method) results in different stimulation sites in DLPFC for different patients, which may

contribute to the heterogeneity in the effect sizes of antidepressant responses in prior trials

(85, 86). In the current study, we underscore the potential importance of individualized,

precise identification of stimulation sites for rTMS by showing that different DLPFC ROIs

correspond to different group difference FC profiles. In line with our findings, Siddiqi and

colleagues found that the correlation maps between stimulations' connectomes and symptom

responses could be grouped into two clusters, indicating that different response patterns are

related to different TMS targets (87). Our results regarding network-wise FC suggest that BA

46 is hyperconnected to the DMN while BA 9 is hyperconnected to the LN in MDD relative to

HCs. The DMN has been proposed to be involved in pathological ruminations (76) that can

occur in some patient with MDD, while the LN might be related to anhedonia, decreased

motivation, and changes in sleep and appetite patterns (3). We speculate that stimulating one

site might be effective for a specific sub-group of patients with a particular cluster of symptoms

(e.g., rumination vs. anhedonia) while other patients might not respond. We note that

leveraging anatomical landmarks to precisely target the boundary between BA 9 and BA 46 did

not consistently outperform the "5 cm" method or the F3 Beam method (88, 89). Our results

suggest that the FC profiles of these two regions could be a critical factor in determining the

treatment outcome of TMS.

Definition of sgACC

It may seem counterintuitive that only spherical-defined sgACC could generate significant

group difference maps when GSR was implemented. We note that the anatomical definition

of the sgACC remains controversial. The sgACC was first implicated in the physiopathology of

MDD based on its increases glucose metabolism (90) and rsfMRI FCs (9) in MDD patients who

a re not respondent to anti-depressant medications and a general reduction in glucose

metabolism (91) and blood flow (92) after anti-depressant treatments (93). Most evidence

supporting this hypothesis come from functional imaging research (e.g., PET, fMRI). As a result,

the brain region showing abnormal metabolism or blood flow may not overlap precisely with

the anatomical landmarks of BA 25. Thus, a less specific definition of sgACC (i.e., a sphere) may

better capture the brain region in which functional activity is altered, despite its lower

anatomic accuracy. Recent studies have applied a more sophisticated targeting algorithm that

searched for smaller sub-units within the broader anatomical sgACC region and led to

markedly improved treatment outcome (39). Another issue is the poor signal-to-noise ratio of

sgACC in fMRI (94). Calculating whole-brain FC maps between one given subject's sgACC and

a large independent cohort (so-called "seed map") and then using this seed map to substitute

the sgACC seed region may help address this issue and generate a more reproducible DLPFC

target (36, 37). Further research is needed to develop the optimal way to define the sgACC in

clinical practice.

Limitations

Several limitations need to be considered. First, we noted that the inconsistency between our results and the broader literature could partly be due to racial differences of the samples (i.e., Eastern Asian vs. Caucasian). Consortiums that intend to pool existing neuroimaging data worldwide, such as the Enhancing NeuroImaging Genetics through Meta-Analysis (ENIGMA) Major Depressive Disorder (MDD) consortium (46), have accumulated large-scale data primarily from Western countries and published several high-impact studies delineating MDD-related anatomical abnormalities (95, 96). Future collaborations between the DIRECT and the ENIGMA-MDD consortiums may help address potential cultural, genetic, and environmental mechanisms involved in more diverse groups of MDD patients. Second, the current study applied a volume-based preprocessing approach to facilitate comparison with previous literature. Surface-based preprocessing strategies have provided more accurate and detailed representations of cortical and subcortical structures (97). Recent research has begun to explore surface-based rTMS target identification algorithms and has shown promising clinical relevance (34, 35, 98). Future research should consider using surface-based case-control difference maps to develop new ways to identify TMS targets.

Conclusion

In summary, we leveraged a large sample of MDD patients to fully delineate group differences of sgACC FC maps between MDD patients and HCs. We also evaluated the clinical relevance of TMS treatment outcomes as a function of TMS target sites. Moreover, we considered several common left DLPFC ROIs' case-control comparison results of FC maps, which underscored the functional heterogeneity within the anatomical landmarks of the left DLPFC and highlighted the necessity of identifying individualized DLPFC TMS sites.

Conflict of interest

The authors declare no conflict of interest.

Acknowledgements

This work was funded by the Sci-Tech Innovation 2030 - Major Project of Brain Science and Brain-inspired Intelligence Technology (No. 2021ZD0200600), the National Key R&D Program of China (No. 2017YFC1309902), the National Natural Science Foundation of China (No. 82122035, No. 81671774 and No. 81630031), the 13th Five-year Informatization Plan of Chinese Academy of Sciences (No. XXH13505), the Key Research Program of the Chinese Academy of Sciences (No. ZDBS-SSW-JSC006), Beijing Nova Program of Science and Technology (No. Z191100001119104), the Scientific Foundation of Institute of Psychology, Chinese Academy of Sciences (No. E2CX4425YZ), the China Postdoctoral Science Foundation (No. 2019M660847), the China National Postdoctoral Program for Innovative Talents (No. BX20200360), the Scientific Foundation of Institute of Psychology, Chinese Academy of Sciences (No. Y9CX422005), and the Special Research Assistant Program of the Chinese Academy of Sciences (No. E2CX0624). The support provided by China Scholarship Council (CSC, No. 202104910248) during a visit of Xiao Chen to the Centre for Addiction and Mental Health is acknowledged.

References

1. G. S. Malhi, J. J. Mann, Depression. *Lancet (London, England)* **392**, 2299-2312 (2018).
2. W. C. Drevets, J. L. Price, M. L. Furey, Brain structural and functional abnormalities in mood disorders: implications for neurocircuitry models of depression. *Brain structure & function* **213**, 93-118 (2008).
3. H. S. Mayberg, Modulating dysfunctional limbic-cortical circuits in depression: towards development of brain-based algorithms for diagnosis and optimised treatment. *British medical bulletin* **65**, 193-207 (2003).
4. H. S. Mayberg *et al.*, Deep brain stimulation for treatment-resistant depression. *Neuron* **45**, 651-660 (2005).
5. S. G. Disner, C. G. Beevers, E. A. P. Haigh, A. T. Beck, Neural mechanisms of the cognitive model of depression. *Nat Rev Neurosci* **12**, 467-477 (2011).
6. R. F. H. Cash *et al.*, Using Brain Imaging to Improve Spatial Targeting of Transcranial Magnetic Stimulation for Depression. *Biol Psychiatry* 10.1016/j.biopsych.2020.05.033 (2020).
7. P. B. Fitzgerald, Targeting repetitive transcranial magnetic stimulation in depression: do we really know what we are stimulating and how best to do it? *Brain Stimul* **14**,

573   730-736 (2021).

8. C. G. Connolly *et al.*, Resting-State Functional Connectivity of Subgenual Anterior Cingulate Cortex in Depressed Adolescents. *Biological Psychiatry* **74**, 898-907 (2013).

9. M. D. Greicius *et al.*, Resting-state functional connectivity in major depression: abnormally increased contributions from subgenual cingulate cortex and thalamus. *Biol Psychiatry* **62**, 429-437 (2007).

10. H. Wu *et al.*, Changed Hub and Corresponding Functional Connectivity of Subgenual Anterior Cingulate Cortex in Major Depressive Disorder. *Front Neuroanat* **10**, 120 (2016).

11. C. G. Davey, B. J. Harrison, M. Yucel, N. B. Allen, Regionally specific alterations in functional connectivity of the anterior cingulate cortex in major depressive disorder. *Psychol Med* **42**, 2071-2081 (2012).

12. B. Cheng *et al.*, Functional connectivity patterns of the subgenual anterior cingulate cortex in first-episode refractory major depressive disorder. *Brain imaging and behavior* **15**, 2397-2405 (2021).

13. Y. I. Sheline, J. L. Price, Z. Yan, M. A. Mintun, Resting-state functional MRI in depression unmasks increased connectivity between networks via the dorsal nexus. *Proc Natl Acad Sci U S A* **107**, 11020-11025 (2010).

14. M. G. Berman *et al.*, Depression, rumination and the default network. *Social cognitive and affective neuroscience* **6**, 548-555 (2011).

15. X. Wu *et al.*, Dysfunction of the cingulo-opercular network in first-episode medication-naive patients with major depressive disorder. *J Affect Disord* **200**, 275-283 (2016).

16. K. Sudheimer *et al.*, Decreased hypothalamic functional connectivity with subgenual cortex in psychotic major depression. *Neuropsychopharmacology* **40**, 849-860 (2015).

17. C. G. Davey, M. Yücel, N. B. Allen, B. J. Harrison, Task-related deactivation and functional connectivity of the subgenual cingulate cortex in major depressive disorder. *Frontiers in psychiatry* **3**, 14 (2012).

18. M. S. Gaffrey *et al.*, Subgenual cingulate connectivity in children with a history of preschool-depression. *Neuroreport* **21**, 1182-1188 (2010).

19. T. C. Ho *et al.*, Functional connectivity of negative emotional processing in adolescent depression. *J Affect Disord* **155**, 65-74 (2014).

20. K. S. Button *et al.*, Power failure: why small sample size undermines the reliability of neuroscience. *Nat Rev Neurosci* **14**, 365-376 (2013).

21. J. Algermissen, D. M. A. Mehler, May the power be with you: are there highly powered studies in neuroscience, and how can we get more of them? *J Neurophysiol* **119**, 2114-2117 (2018).

22. K. Murphy, M. D. Fox, Towards a Consensus Regarding Global Signal Regression for Resting State Functional Connectivity MRI. *Neuroimage* 10.1016/j.neuroimage.2016.11.052 (2016).

23. L. Q. Uddin, Mixed Signals: On Separating Brain Signal from Noise. *Trends Cogn Sci* **21**, 405-406 (2017).

24. M. D. Fox, D. Zhang, A. Z. Snyder, M. E. Raichle, The global signal and observed anticorrelated resting state brain networks. *J Neurophysiol* **101**, 3270-3283 (2009).

25. R. Ciric *et al.*, Mitigating head motion artifact in functional connectivity MRI. *Nature*
protocols **13**, 2801-2826 (2018).
26. K. Murphy, R. M. Birn, D. A. Handwerker, T. B. Jones, P. A. Bandettini, The impact of
global signal regression on resting state correlations: Are anti-correlated networks
introduced? *NeuroImage* **44**, 893-905 (2009).
27. L. Parkes, B. Fulcher, M. Yücel, A. Fornito, An evaluation of the efficacy, reliability, and
sensitivity of motion correction strategies for resting-state functional MRI.
NeuroImage **171**, 415-436 (2018).
28. R. Ge, J. Downar, D. M. Blumberger, Z. J. Daskalakis, F. Vila-Rodriguez, Functional
connectivity of the anterior cingulate cortex predicts treatment outcome for rTMS in
treatment-resistant depression at 3-month follow-up. *Brain Stimul* **13**, 206-214 (2020).
29. R. F. H. Cash *et al.*, A multivariate neuroimaging biomarker of individual outcome to
transcranial magnetic stimulation in depression. *Hum Brain Mapp* **40**, 4618-4629
(2019).
30. N. S. Philip *et al.*, Network Mechanisms of Clinical Response to Transcranial Magnetic
Stimulation in Posttraumatic Stress Disorder and Major Depressive Disorder. *Biol*
Psychiatry **83**, 263-272 (2018).
31. M. D. Fox, R. L. Buckner, M. P. White, M. D. Greicius, A. Pascual-Leone, Efficacy of
transcranial magnetic stimulation targets for depression is related to intrinsic
functional connectivity with the subgenual cingulate. *Biol Psychiatry* **72**, 595-603
(2012).
32. I. G. Elbau *et al.*, Functional Connectivity Mapping for rTMS Target Selection in
Depression. *Am J Psychiatry* **180**, 230-240 (2023).
33. A. Weigand *et al.*, Prospective Validation That Subgenual Connectivity Predicts
Antidepressant Efficacy of Transcranial Magnetic Stimulation Sites. *Biol Psychiatry* **84**,
28-37 (2018).
34. R. F. H. Cash, L. Cocchi, J. Lv, P. B. Fitzgerald, A. Zalesky, Functional Magnetic Resonance
Imaging-Guided Personalization of Transcranial Magnetic Stimulation Treatment for
Depression. *JAMA psychiatry* **78**, 337-339 (2021).
35. S. H. Siddiqi, A. Weigand, A. Pascual-Leone, M. D. Fox, Identification of Personalized
Transcranial Magnetic Stimulation Targets Based on Subgenual Cingulate Connectivity:
An Independent Replication. *Biol Psychiatry* **90**, e55-e56 (2021).
36. M. D. Fox, H. Liu, A. Pascual-Leone, Identification of reproducible individualized targets
for treatment of depression with TMS based on intrinsic connectivity. *Neuroimage* **66**,
151-160 (2013).
37. R. F. H. Cash *et al.*, Personalized connectivity-guided DLPFC-TMS for depression:
Advancing computational feasibility, precision and reproducibility. *Hum Brain Mapp*
10.1002/hbm.25330 (2021).
38. E. J. Cole *et al.*, Stanford Neuromodulation Therapy (SNT): A Double-Blind Randomized
Controlled Trial. *Am J Psychiatry* 10.1176/appi.ajp.2021.20101429,
appiajp202120101429 (2021).
39. E. J. Cole *et al.*, Stanford Accelerated Intelligent Neuromodulation Therapy for
Treatment-Resistant Depression. *Am J Psychiatry* **177**, 716-726 (2020).

40.   C. G. Yan *et al.*, Reduced default mode network functional connectivity in patients with
660   recurrent major depressive disorder. *Proc Natl Acad Sci U S A*
661   10.1073/pnas.1900390116 (2019).

662   41.   G. Rajkowska, P. S. Goldman-Rakic, Cytoarchitectonic definition of prefrontal areas in
663   the normal human cortex: II. Variability in locations of areas 9 and 46 and relationship
664   to the Talairach Coordinate System. *Cerebral cortex (New York, N.Y. : 1991)* **5**, 323-337
665   (1995).

666   42.   W. Beam, J. J. Borckardt, S. T. Reeves, M. S. George, An efficient and accurate new
667   method for locating the F3 position for prefrontal TMS applications. *Brain Stimul* **2**, 50-
668   54 (2009).

669   43.   M. S. George *et al.*, Daily repetitive transcranial magnetic stimulation (rTMS) improves
670   mood in depression. *Neuroreport* **6**, 1853-1856 (1995).

671   44.   M. L. Paillère Martinot *et al.*, Influence of prefrontal target region on the efficacy of
672   repetitive transcranial magnetic stimulation in patients with medication-resistant
673   depression: a [(18)F]-fluorodeoxyglucose PET and MRI study. *The international journal
674   of neuropsychopharmacology* **13**, 45-59 (2010).

675   45.   X. Chen *et al.*, The DIRECT consortium and the REST-meta-MDD project: towards
676   neuroimaging biomarkers of major depressive disorder. *Psychoradiology* **2**, 32-42
677   (2022).

678   46.   L. Schmaal *et al.*, ENIGMA MDD: seven years of global neuroimaging studies of major
679   depression through worldwide data sharing. *Transl Psychiatry* **10**, 172 (2020).

680   47.   H. Yang *et al.*, Disrupted intrinsic functional brain topology in patients with major
681   depressive disorder. *Mol Psychiatry* 10.1038/s41380-021-01247-2 (2021).

682   48.   Y. Long *et al.*, Altered resting-state dynamic functional brain networks in major
683   depressive disorder: Findings from the REST-meta-MDD consortium. *Neuroimage Clin*
684   **26**, 102163 (2020).

685   49.   Y. D. Ding *et al.*, Disrupted hemispheric connectivity specialization in patients with
686   major depressive disorder: Evidence from the REST-meta-MDD Project. *J Affect Disord*
687   **284**, 217-228 (2021).

688   50.   K. Deng *et al.*, Impaired robust interhemispheric function integration of depressive
689   brain from REST-meta-MDD database in China. *Bipolar disorders* 10.1111/bdi.13139
690   (2021).

691   51.   S. Liang *et al.*, Biotypes of major depressive disorder: Neuroimaging evidence from
692   resting-state default mode network patterns. *Neuroimage Clin* **28**, 102514 (2020).

693   52.   P. H. Liu *et al.*, Brain structural alterations in MDD patients with gastrointestinal
694   symptoms: Evidence from the REST-meta-MDD project. *Progress in neuro-
695   psychopharmacology & biological psychiatry* **111**, 110386 (2021).

696   53.   C.-G. Yan, X.-D. Wang, X.-N. Zuo, Y.-F. Zang, DPABI: Data Processing & Analysis for
697   (Resting-State) Brain Imaging. *Neuroinformatics* **14**, 339-351 (2016).

698   54.   C.-G. Yan, Y.-F. Zang, DPARSF: A MATLAB Toolbox for "Pipeline" Data Analysis of Resting-
699   State fMRI. *Frontiers in systems neuroscience* **4**, 13 (2010).

700   55.   C.-G. Yan, X.-D. Wang, B. Lu, DPABISurf: data processing & analysis for brain imaging
701   on surface. *Science Bulletin* https://doi.org/10.1016/j.scib.2021.09.016 (2021).

56. C. Liston *et al.*, Default mode network mechanisms of transcranial magnetic
stimulation in depression. *Biol Psychiatry* **76**, 517-526 (2014).

57. J. K. Mai, M. Majtanik, *Human Brain in Standard MNI Space: A Comprehensive Pocket*
Atlas (Elsevier Science, 2017).

58. W. E. Johnson, C. Li, A. Rabinovic, Adjusting batch effects in microarray expression data
using empirical Bayes methods. *Biostatistics* **8**, 118-127 (2007).

59. C. G. Yan *et al.*, Reduced default mode network functional connectivity in patients with
recurrent major depressive disorder. *Proc Natl Acad Sci U S A* **116**, 9078-9083 (2019).

60. F. Murtagh, P. Legendre, Ward's Hierarchical Agglomerative Clustering Method: Which
Algorithms Implement Ward's Criterion? *Journal of Classification* **31**, 274-295 (2014).

61. R. Tibshirani, G. Walther, T. Hastie, Estimating the number of clusters in a data set via
the gap statistic. *Journal of the Royal Statistical Society: Series B (Statistical*
Methodology) **63**, 411-423 (2001).

62. A. Schaefer *et al.*, Local-Global Parcellation of the Human Cerebral Cortex from
Intrinsic Functional Connectivity MRI. *Cerebral cortex (New York, N.Y. : 1991)* **28**, 3095-
3114 (2018).

63. M. Xia, J. Wang, Y. He, BrainNet Viewer: a network visualization tool for human brain
connectomics. *PLoS One* **8**, e68910 (2013).

64. R. Computing, R: A language and environment for statistical computing. *Vienna: R Core*
Team (2013).

65. R. A. M. Villanueva, Z. J. Chen (2019) ggplot2: elegant graphics for data analysis.
(Taylor & Francis).

66. M. Greicius, Resting-state functional connectivity in neuropsychiatric disorders.
Current opinion in neurology **21**, 424-430 (2008).

67. K. R. Cullen *et al.*, A preliminary study of functional connectivity in comorbid
adolescent depression. *Neuroscience Letters* **460**, 227-231 (2009).

68. C. Strikwerda-Brown *et al.*, Mapping the relationship between subgenual cingulate
cortex functional connectivity and depressive symptoms across adolescence. *Social*
cognitive and affective neuroscience **10**, 961-968 (2015).

69. J. P. A. Ioannidis, Why Most Published Research Findings Are False. *PLOS Medicine* **2**,
e124 (2005).

70. X. Chen, B. Lu, C. G. Yan, Reproducibility of R-fMRI metrics on the impact of different
strategies for multiple comparison correction and sample sizes. *Hum Brain Mapp* **39**,
300-318 (2018).

71. J. D. Power, M. Plitt, T. O. Laumann, A. Martin, Sources and implications of whole-brain
fMRI signals in humans. *Neuroimage* **146**, 609-625 (2017).

72. M. F. Glasser *et al.*, Using temporal ICA to selectively remove global noise while
preserving global signal in functional MRI data. *Neuroimage* **181**, 692-717 (2018).

73. D. M. Blumberger *et al.*, Effectiveness of theta burst versus high-frequency repetitive
transcranial magnetic stimulation in patients with depression (THREE-D): a
randomised non-inferiority trial. *Lancet (London, England)* **391**, 1683-1692 (2018).

74. J. L. Price, W. C. Drevets, Neurocircuitry of mood disorders. *Neuropsychopharmacology*
35, 192-216 (2010).

75.   J. L. Price, W. C. Drevets, Neural circuits underlying the pathophysiology of mood
disorders. *Trends Cogn Sci* **16**, 61-71 (2012).

76.   J. P. Hamilton, M. Farmer, P. Fogelman, I. H. Gotlib, Depressive Rumination, the Default-
Mode Network, and the Dark Matter of Clinical Neuroscience. *Biol Psychiatry* **78**, 224-
230 (2015).

77.   G. Parker, B. Chan, D. Hadzi-Pavlovic, Lower rates of depression in westernised Chinese
in the US. *J Affect Disord* **104**, 175-178 (2007).

78.   G. Parker, G. Gladstone, K. T. Chee, Depression in the planet's largest ethnic group: the
Chinese. *Am J Psychiatry* **158**, 857-864 (2001).

79.   P. Qiu, E. D. Caine, F. Hou, C. Cerulli, M. N. Wittink, Depression as seen through the
eyes of rural Chinese women: Implications for help-seeking and the future of mental
health care in China. *J Affect Disord* **227**, 38-47 (2018).

80.   C. Kiyohara, K. Yoshimasu, Association between major depressive disorder and a
functional polymorphism of the 5-hydroxytryptamine (serotonin) transporter gene: a
meta-analysis. *Psychiatr Genet* **20**, 49-58 (2010).

81.   R. Ge *et al.*, Predictive Value of Acute Neuroplastic Response to rTMS in Treatment
Outcome in Depression: A Concurrent TMS-fMRI Trial. *Am J Psychiatry*
10.1176/appi.ajp.21050541, appiajp21050541 (2022).

82.   X.-N. Zuo, X.-X. Xing, Test-retest reliabilities of resting-state FMRI measurements in
human brain functional connectomics: A systems neuroscience perspective.
Neuroscience & Biobehavioral Reviews **45**, 100-118 (2014).

83.   D. Wang *et al.*, Parcellating cortical functional networks in individuals. *Nat Neurosci* **18**,
1853-1860 (2015).

84.   P. B. Fitzgerald *et al.*, An analysis of functional neuroimaging studies of dorsolateral
prefrontal cortical activity in depression. *Psychiatry Res* **148**, 33-45 (2006).

85.   K. A. Johnson *et al.*, Prefrontal rTMS for treating depression: location and intensity
results from the OPT-TMS multi-site clinical trial. *Brain Stimul* **6**, 108-117 (2013).

86.   T. Perera *et al.*, The Clinical TMS Society Consensus Review and Treatment
Recommendations for TMS Therapy for Major Depressive Disorder. *Brain Stimul* **9**,
336-346 (2016).

87.   S. H. Siddiqi *et al.*, Distinct Symptom-Specific Treatment Targets for Circuit-Based
Neuromodulation. *Am J Psychiatry* **177**, 435-446 (2020).

88.   P. B. Fitzgerald *et al.*, A randomized trial of rTMS targeted with MRI based neuro-
navigation in treatment-resistant depression. *Neuropsychopharmacology* **34**, 1255-
1262 (2009).

89.   T. Hebel *et al.*, A direct comparison of neuronavigated and non-neuronavigated
intermittent theta burst stimulation in the treatment of depression. *Brain Stimul* **14**,
335-343 (2021).

90.   S. H. Kennedy *et al.*, Differences in brain glucose metabolism between responders to
CBT and venlafaxine in a 16-week randomized controlled trial. *Am J Psychiatry* **164**,
778-788 (2007).

91.   H. S. Mayberg *et al.*, Reciprocal limbic-cortical function and negative mood: converging
PET findings in depression and normal sadness. *Am J Psychiatry* **156**, 675-682 (1999).

92.   M. S. Nobler *et al.*, Decreased regional brain metabolism after ect. *Am J Psychiatry* **158**,
305-308 (2001).

93.   H. S. Mayberg, Targeted electrode-based modulation of neural circuits for depression.
The Journal of clinical investigation **119**, 717-725 (2009).

94.   J. G. Ojemann *et al.*, Anatomic localization and quantitative analysis of gradient
refocused echo-planar fMRI susceptibility artifacts. *Neuroimage* **6**, 156-167 (1997).

95.   L. Schmaal *et al.*, Cortical abnormalities in adults and adolescents with major
depression based on brain scans from 20 cohorts worldwide in the ENIGMA Major
Depressive Disorder Working Group. *Mol Psychiatry* 10.1038/mp.2016.60 (2016).

96.   L. Schmaal *et al.*, Subcortical brain alterations in major depressive disorder: findings
from the ENIGMA Major Depressive Disorder working group. *Mol Psychiatry* **21**, 806-
812 (2016).

97.   O. Esteban *et al.*, fMRIPrep: a robust preprocessing pipeline for functional MRI. *Nat*
Methods **16**, 111-116 (2019).

98.   C. J. Lynch *et al.*, Automated optimization of TMS coil placement for personalized
functional network engagement. *Neuron* **110**, 3263-3277.e3264 (2022).

  Figure Legends

Figure 1. Depression Imaging REsearch ConsorTium (DIRECT) Phase II sample characteristics.

(A) Sample sizes of each site; (B) Number of male/female subjects irrespective of diagnosis; (C)

Violin plots depicting the distribution of age (in years). Solid black lines indicate the mean, 25th

and 75th percentiles; (D) Violin plots show education distribution (in years). Solid black lines

indicate the mean, 25th and 75th percentiles; (E) Violin plots depicting the distribution of scores

of the Hamilton Depression Rating Scale (HAMD). Solid black lines indicate the mean, 25th and

75th percentiles.

Figure 2. Group difference maps of subgenual anterior cingulate cortex (sgACC) functional

connectivity (FC) profiles. (A) t maps of MDD-related sgACC FC abnormalities with and without

global signal regression (GSR). (B) Group differences of FCs between sgACC and visual network

(VN), somatomotor network (SMN), dorsal attention network (DAN), ventral attention

network (VAN), limbic network (LN), frontoparietal network (FPN), and default mode network

(DMN). Asterisks indicate significance after Bonferroni correction. (C) The group difference of

sgACC FC maps was positively related to clinical improvement following transcranial magnetic

stimulation (TMS). Data were extracted from an independent openly shared dataset (33). The

sizes of the spheres indicate the reduction rates of the Hamilton Depression Rating Scale

(HAMD) and the group difference map after global signal regression (GSR) was rendered on

the surface.

Figure 3. The position of the anti-correlation peak in group mean functional connectivity (FC) differed between major depressive disorder (MDD) patients and healthy controls (HCs). The group mean FC maps of MDD patients (the MDD peak and group difference targets) or HCs (the HC peak and Weigand et al. group target) after GSR were rendered on the surface. The "group difference targets" and a previously reported subgenual anterior cingulate cortex (sgACC) target calculated based on a large cohort of healthy adults (33) were also plotted for illustration. Peaks were searched within the left dorsal lateral prefrontal cortex (DLPFC) mask.

839 Figure 4. Two clusters showed significant case-control differences in functional connectivity

840 (FC) with the subgenual anterior cingulate cortex (sgACC). The peaks of these two clusters were

841 chosen as "group difference targets."

846 Figure 5. The group difference maps of different left dorsal lateral prefrontal cortex (DLPFC)

847 regions of interest (ROI) could be grouped into 3 clusters. (A) MDD-related DLPFC FC

848 abnormalities. (B) The hierarchical clustering of left DLPFC ROIs' case-control difference maps.

849 †: Corrected with Gaussian random field correction at the threshold of voxel-wise p < 0.01 and

850 cluster-wise p < 0.05, two-tailed.

Figure 6. Group differences of FCs between the different left dorsal lateral prefrontal cortex (DLPFC) regions of interest (ROI) and visual network (VN), somatomotor network (SMN), dorsal attention network (DAN), ventral attention network (VAN), limbic network (LN), frontoparietal network (FPN), and default mode network (DMN). Asterisks indicate significance after Bonferroni correction.