Subgenual Anterior Cingulate Cortex Functional Connectivity Abnormalities in Depression: Insights from Brain Imaging Big Data and Precision-Guided Personalized Intervention via Transcranial Magnetic Stimulation

Xiao Chen[1,2,3,4,5]#, Bin Lu[1,3,4,5]#, Yu-Wei Wang[1,3,4,5], Xue-Ying Li[1,3,4,5], Zi-Han Wang[1,3,4,5], Hui-Xian Li[6], Yi-Fan Liao[1,3,4,5], Daniel M. Blumberger[2,7], Francisco Xavier Castellanos[8,9], Eduardo A. Garza-Villarreal[10], Li-Ping Cao[11], Guan-Mao Chen[12], Jian-Shan Chen[11], Tao Chen[13], Tao-Lin Chen[14,15], Yan-Rong Chen[1,3,4,5], Yu-Qi Cheng[16], Zhao-Song Chu[16], Shi-Xian Cui[1,17,18], Xi-Long Cui[19], Zhao-Yu Deng[1,3,4,5], Qing-Lin Gao[1,3,4,5], Qi-Yong Gong[14,15], Wen-Bin Guo[19], Can-Can He[20], Zheng-Jia-Yi Hu[1,17,18], Qian Huang[21], Xin-Lei Ji[19], Feng-Nan Jia[22], Li Kuang[21], Bao-Juan Li[23], Feng Li[24], Tao Li[25,26], Xue Li[1,3,4,5], Tao Lian[1,3,4,5], Xiao-Yun Liu[27], Yan-Song Liu[22], Zhe-Ning Liu[19], Yi-Cheng Long[19], Jian-Ping Lu[28], Jiang Qiu[29], Xiao-Xiao Shan[19], Tian-Mei Si[30], Peng-Feng Sun[31], Chuan-Yue Wang[24], Han-Lin Wang[1,3,4,5], Xiang Wang[19], Ying Wang[12], Chen-Nan Wu[1,3,4,5], Xiao-Ping Wu[31], Xin-Ran Wu[29], Yan-Kun Wu[30], Chun-Ming Xie[20], Guang-Rong Xie[19], Peng Xie[32,33,34], Xiu-Feng Xu[16], Zhen-Peng Xue[28], Hong Yang[13], Jian Yang[33], Hua Yu[25,26], Yong-Qiang Yu[35,36,37], Min-Lan Yuan[38], Yong-Gui Yuan[27], Yu-Feng Zang[39,40], Ai-Xia Zhang[41], Ke-Rang Zhang[41], Wei Zhang[38], Zi-Jing Zhang[1,3,4,5], Jing-Ping Zhao[19], Jia-Jia Zhu[35,36,37], Xi-Nian Zuo[42,43], the DIRECT Consortium, Hua-Ning Wang[23], Chao-Gan Yan[1,3,4,5,17,18]*

[1]CAS Key Laboratory of Behavioral Science, Institute of Psychology, Beijing 100101, China

[2]Temerty Centre for Therapeutic Brain Intervention, Campbell Family Research Institute, Centre for Addiction and Mental Health, Toronto, Ontario M6J1H4, Canada

[3]Magnetic Resonance Imaging Research Center, Institute of Psychology, Chinese Academy of Sciences, Beijing 100101, China

[4]Department of Psychology, University of Chinese Academy of Sciences, Beijing 100049, China

[5]International Big-Data Center for Depression Research, Chinese Academy of Sciences, Beijing 100101, China

[6]The Third Affiliated Hospital of Zhengzhou University, Zhengzhou, Henan 450052, China

[7]Department of Psychiatry, Temerty Faculty of Medicine, University of Toronto, Toronto, Ontario

M5S1A1, Canada

[8]Department of Child and Adolescent Psychiatry, NYU Grossman School of Medicine, New York, NY

10016, USA

[9]Nathan Kline Institute for Psychiatric Research, Orangeburg, NY 10962, USA

[10]Instituto de Neurobiologia, Universidad Nacional Autonoma de Mexico camus Juriquilla, Queretaro

7630, Mexico

[11]Affiliated Brain Hospital of Guangzhou Medical University, Guangzhou, Guangdong 510370, China

[12]The First Affiliated Hospital of Jinan University, Guangzhou, Guangdong 250024, China

[13]Department of Radiology, The First Affiliated Hospital, College of Medicine, Zhejiang University,

Hangzhou, Zhejiang 310058, China

[14]Department of Radiology and Huaxi MR Research Center (HMRRC), Functional and Molecular

Imaging Key Laboratory of Sichuan Province, West China Hospital of Sichuan University, Chengdu,

Sichuan 610041, China

[15]Research Unit of Psychoradiology, Chinese Academy of Medical Sciences, Chengdu, Sichuan 610041,

China

[16]Department of Psychiatry, First Affiliated Hospital of Kunming Medical University, Kunming, Yunnan

650032, China

[17]Sino-Danish College, University of Chinese Academy of Sciences, Beijing 101408, China

[18]Sino-Danish Center for Education and Research, Graduate University of Chinese Academy of Sciences,

Beijing 101408, China

[19]Department of Psychiatry, and National Clinical Research Center for Mental Disorders, The Second

Xiangya Hospital of Central South University, Changsha, Hunan 410011, China

[20]Department of Neurology, Affiliated ZhongDa Hospital of Southeast University, Nanjing, Jiangsu

210009, China

[21]Department of Psychiatry, The First Affiliated Hospital of Chongqing Medical University, Chongqing

400042, China

[22]Department of Clinical Psychology, Suzhou Psychiatric Hospital, The Affiliated Guangji Hospital of

Soochow University, Suzhou, Jiangsu 215003, China

[23]Xijing Hospital of Air Force Military Medical University, Xi'an, Shaanxi 710032, China

[24]Beijing Anding Hospital, Capital Medical University, Beijing 100120, China

[25]Affiliated Mental Health Center & Hangzhou Seventh People's Hospital, Zhejiang University School of

Medicine, Hangzhou, Zhejiang 310063, China

[26]Mental Health Center and Psychiatric Laboratory, West China Hospital of Sichuan University,

Chengdu, Sichuan 610044, China

[27]Department of Psychosomatics and Psychiatry, Zhongda Hospital, School of Medicine, Southeast

University, Nanjing, Jiangsu 210009, China

[28]Shenzhen Kangning Hospital, Shenzhen, Guangzhou 518020, China

[29]Faculty of Psychology, Southwest University, Chongqing 400715, China

[30]National Clinical Research Center for Mental Disorders (Peking University Sixth Hospital) & Key

Laboratory of Mental Health, Ministry of Health (Peking University), Beijing 100191, China

[31]Xi'an Central Hospital, Xi'an, Shaanxi 710004, China

[32]Institute of Neuroscience, Chongqing Medical University, Chongqing 400016, China

[33]Chongqing Key Laboratory of Neurobiology, Chongqing 400000, China

[34]Department of Neurology, The First Affiliated Hospital of Chongqing Medical University, Chongqing

400042, China

[35]Department of Radiology, The First Affiliated Hospital of Anhui Medical University, Hefei, Anhui

230022, China

[36]Research Center of Clinical Medical Imaging, Hefei, Anhui 230032, China

[37]Anhui Provincial Institute of Translational Medicine, Hefei, Anhui 230032, China

[38]West China Hospital of Sichuan University, Chengdu, Sichuan 610044, China

[39]Center for Cognition and Brain Disorders, The Affiliated Hospital of Hangzhou Normal University,

Hangzhou, Zhejiang 310018, China

[40]Zhejiang Key Laboratory for Research in Assessment of Cognitive Impairments, Hangzhou, Zhejiang

310000, China

[41]First Hospital of Shanxi Medical University, Taiyuan, Shanxi 030001, China

[42]Developmental Population Neuroscience Research Center, IDG/McGovern Institute for Brain

Research, Beijing Normal University, Beijing 100091, China

[43]National Basic Science Data Center, Beijing 100038, China

[44]Department of Psychiatry, Sir Run Run Shaw Hospital, Zhejiang University School of Medicine,

Hangzhou, Zhejiang 310020, China

[45]Shanghai Mental Health Center, Shanghai Jiao Tong University School of Medicine, Shanghai 200030,

China

[46]Department of Diagnostics, Affiliated Hospital, Hangzhou Normal University Medical School,

Hangzhou, Zhejiang 311121, China

[47]Department of Psychiatry, First Affiliated Hospital, China Medical University, Shenyang, Liaoning

110122, China

[48]Early Intervention Unit, Department of Psychiatry, Affiliated Nanjing Brain Hospital, Nanjing Medical

University, Nanjing, Jiangsu 210024, China

[49]Department of Neurology, The First Affiliated Hospital of Anhui Medical University, Hefei, Anhui

230022, China

[50]Department of Psychiatry, Shanghai Jiao Tong University School of Medicine, Shanghai 200025, China

*Corresponding author:

Chao-Gan Yan, Ph. D.

CAS Key Laboratory of Behavioral Science, Institute of Psychology, Beijing, China

16 Lincui Road, Chaoyang District, Beijing 100101, China

Tel: +86-10-64101582

E-mail: yancg@psych.ac.cn

#Xiao Chen and Bin Lu contributed equally to this work.

DIRECT consortium: Chao-Gan Yan, Ph.D.[1,3,4,5,17,18], Xiao Chen, Ph.D.[1,2,3,4,5], Li-Ping Cao, M.D.[11], Wei

Chen, M.D.[44], Yu-Qi Cheng, M.D.[16], Yi-Ru Fang, M.D.[45], Qi-Yong Gong, M.D.[14,15], Wen-Bin Guo, M.D.[19],

Li Kuang, M.D.[21], Bao-Juan Li, M.D.[23], Tao Li, M.D.[25,26], Yan-Song Liu, M.D.[22], Zhe-Ning Liu, M.D.[19],

Jian-Ping Lu, M.D.[28], Qing-Hua Luo, M.D.[21], Hua-Qing Meng, M.D.[21], Dai-Hui Peng, M.D.[45], Jiang Qiu,

Ph.D.[29], Yue-Di Shen, M.D.[46], Tian-Mei Si, M.D.[30], Yan-Qing Tang, M.D.[47], Chuan-Yue Wang, M.D.[24], Fei

Wang, M.D.[47,48], Hua-Ning Wang, M.D.[23], Kai Wang, M.D.[49], Xiang Wang, M.D.[19], Ying Wang, M.D.[12],

Xiao-Ping Wu, M.D.[31], Chun-Ming Xie, M.D.[20], Guang-Rong Xie, M.D.[19], Peng Xie, M.D.[32,33,34], Xiu-Feng

Xu, M.D.[16], Hong Yang, M.D.[13], Jian Yang, M.D.[33], Shu-Qiao Yao, M.D.[19], Yong-Qiang Yu, M.D.[35,36,37],

Yong-Gui Yuan, M.D.[27], Ke-Rang Zhang, M.D.[41], Wei Zhang, M.D.[38], Zhi-Jun Zhang, M.D.[20], Jun-Juan Zhu,

M.D.[50], Xi-Nian Zuo, Ph.D.[42,43], Jing-Ping Zhao, M.D.[19], Yu-Feng Zang, M.D.[39,40]

Abstract

Background The subgenual anterior cingulate cortex (sgACC) plays a central role in the pathophysiology of major depressive disorder (MDD), and its functional interactive profile with the left dorsal lateral prefrontal cortex (DLPFC) is associated with transcranial magnetic stimulation (TMS) treatment outcomes. Nevertheless, previous research on sgACC functional connectivity (FC) in MDD has yielded inconsistent results, partly due to small sample sizes and limited statistical power. Furthermore, calculating sgACC-FC to target TMS individually is challenging.

Methods Leveraging a large multi-site cross-sectional sample (1660 MDD patients vs. 1341 healthy controls) from Phase II of the Depression Imaging REsearch ConsorTium (DIRECT), we systematically delineated case-control difference maps of sgACC-FC. Then, we explored the potential impact of such group-level abnormality profiles on the TMS target localization and clinical efficacy. Next, we developed an MDD big data-guided individualized TMS targeting algorithm to integrate group-level statistical maps with individual-level brain activity to localize TMS targets individually.

Results We found an enhanced sgACC-DLPFC FC in MDD patients compared to healthy controls (HC). Such group differences altered the position of the sgACC anti-correlation peak in the left DLPFC. In two independent clinical samples, we showed that the magnitude of TMS targets' case-control differences in sgACC FC was related to clinical improvement. The MDD big data-guided individualized TMS targeting algorithm may generate individualized TMS targets that are clinically superior to group-level targets.

Interpretation We reliably delineated MDD-related abnormalities of sgACC-FC profiles in a large, independently ascertained sample and demonstrated the potential impact of such case-control differences on FC-guided localization of TMS targets.

Funding Ministry of Science and Technology of the People's Republic of China, National Natural Science Foundation of China, and Chinese Academy of Sciences

Keywords: major depressive disorder, transcranial magnetic stimulation, individualization, subgenual anterior cingulate cortex, functional connectivity, dual regression

Introduction

Major depressive disorder (MDD) is a common and debilitating psychiatric disorder projected to be the most burdensome condition worldwide by 2030 [1]. Despite extensive research, the pathophysiology of MDD remains elusive. Nevertheless, a key putative brain region or network hub appears to be the subgenual anterior cingulate cortex (sgACC), which shows reproducible metabolic hyperactivity [2], has been implicated in emotional responses, motivation, and rumination in MDD [3], and it has been shown to be an important target in deep brain stimulation and transcranial magnetic stimulation (TMS) [4,5]. Repetitive TMS above 5 Hz on the left dorsolateral prefrontal cortex (DLPFC) indirectly stimulates the sgACC, and the closer to the sgACC target, the better the clinical outcome [6]. Accordingly, identifying an optimized neuromodulation target in the left DLPFC based on sgACC-related functional connectivity (FC) is crucial for developing effective depression treatments [7,8]. In light of inconsistent findings derived from studies with small sample sizes [9-15], we set out to establish a large sample to identify a reliable abnormal sgACC-DLPFC FC profile in MDD and further integrate this profile with individual brain activity to generate individualized neuromodulation targets for treating depression.

Numerous investigations have delved into FC abnormalities in MDD using resting-state functional magnetic resonance imaging (R-fMRI). Abnormal FCs between sgACC and amygdala, thalamus, temporal gyrus, lingual gyrus, cerebellum, DLPFC, and default mode network (DMN) regions such as medial and dorsal medial prefrontal cortex, precuneus, and parahippocampus have been reported [9,10,13-20]. However, findings have been inconsistent, making integrating findings and generating precise profiles of sgACC-related FC abnormalities challenging. This deficiency in reproducibility could be partially due to small sample sizes, differences in preprocessing pipelines, and low statistical power of clinical imaging studies [21,22]. To address the issue of limited sample size, we initiated the Depression Imaging REsearch ConsorTium (DIRECT) [23] and conducted an initial meta/mega-analysis (N_{MDD} = 1300), referred to as REST-meta-MDD [24]. DIRECT Phase I shared ROI-level signals, thus enabling the

investigation of multiple MDD-related abnormalities in network FC, FC topological and

dynamic features, and functional lateralization [24-31]. In DIRECT Phase II data reporting, we

pooled an expanded MDD sample (N_{MDD} = 1660), which was preprocessed with a

surface-based pipeline, DPABISurf [32]. DIRECT Phase II shared voxel/vertex level BOLD time

series, allowing more flexible and thorough investigations. Leveraging the most

comprehensive MDD R-fMRI dataset to date encompassing depression patients and healthy

controls, we can determine an aberrant sgACC-FC profile associated with MDD, characterized

by superior reproducibility and low risk of false positives.

Maps of sgACC-related FC abnormalities are clinically useful for predicting repetitive TMS

(rTMS) treatment outcomes in MDD patients [33-35]. Specifically, the anti-correlation between

sgACC and left DLPFC has been associated with clinical improvement from rTMS treatment

[36-40]. This has led to the intriguing notion that the FC between sgACC and left DLPFC could be

leveraged to identify more precise rTMS targets and improve the efficacy of rTMS delivered

to the left DLPFC [7,41]. Researchers have identified a group-wise TMS target [38], which was the

most anticorrelated DLPFC site to sgACC in the mean FC map from a large cohort of healthy

adults. Subtle but significant case-control differences in resting-state FC profiles have been

identified in a large sample of MDD patients [24]. Thus, understanding the profiles of sgACC FC

case-control differences and their impact on potential targets for rTMS applied to the left

DLPFC could be a critical step toward developing optimized rTMS target site identification

methods.

Individual human brains exhibit highly heterogeneous functional organization [42], with the

DLPFC regions exhibiting the highest level of interindividual variation in cytoarchitecture,

brain function, and network connectivity profiles [8,43]. While several individualized FC-guided

TMS target identification algorithms have been proposed [36,41,44-48], the target localizations of

most existing TMS protocols have not been individualized. The major obstacles to identifying

individualized TMS locations are the low signal-to-noise ratio in the sgACC area and the poor

reproducibility of individual FC maps [49,50]. The high reliability and statistical power of the

DIRECT MDD cohort (N_{MDD} = 1660) allow the integration of group-level statistical maps and

individual functional brain images to achieve precise and reliable TMS localization. Here, we

propose an MDD big data-guided individualized TMS targeting algorithm based on dual

regression (DR), which was initially developed for mapping group-level independent

component analysis (ICA) results onto individual brains [51]. During individualized target

localization, the DR calculation is entirely confined to the DLPFC region, which has a high

signal-to-noise ratio, avoiding noisy signals from the sgACC region. Thus, this approach

enhances the efficacy and reliability of individualization approaches for identifying TMS

targets [52].

In the present study, we leverage a large-scale multi-center sample (DIRECT Phase II, 1660

MDD patients and 1341 healthy controls (HCs)) to derive a reliable sgACC-related FC

abnormality profile for MDD. Next, we showed that such case-control difference profiles may

be related to the clinical efficiency of TMS and that the positions of the sgACC

anti-correlation peaks might be different in the MDD patients as compared to the HCs. In

light of this, we developed an MDD big data-guided individualized TMS targeting algorithm

that may boost the clinical efficiency of TMS. We hypothesized that MDD patients would

show a significantly abnormal sgACC-FC profile, especially in the left DLPFC. We also

hypothesized that our newly developed DR-based approach would outperform traditional

TMS group targets. To our knowledge, this is the first study to show the possible implications

of the case-control abnormalities regarding the sgACC-FC profiles on the TMS target

localization and to integrate large-scale group-level statistical maps with individual-level

spontaneous brain activity to achieve individualized TMS targeting in MDD.

Materials and methods

Study sample

This study utilized four independent datasets. The first dataset ("DIRECT") is a large-scale,

multi-site consortium sharing standardized preprocessed R-fMRI time series. Building on the

initial success of DIRECT Phase I (the REST-meta-MDD Project) [24], consortium members and

international collaborators met on May 11th-12th, 2019, and agreed to launch DIRECT Phase II,

which comprises 23 case-control designed datasets, including R-fMRI and T1 structural scans

from 1660 MDD patients and 1341 HCs. Researchers from each site took a 2-day DPABISurf

training course on September 14th-15th, 2019, to harmonize the organization and

preprocessing of R-fMRI/T1 structural data. Demographic and clinical characteristics for each

sample are presented in Figure 1 and Table 1. Site information, sample size, and previous

publications based on the shared data are listed in Table S1. All participants were asked to

self-report their sex (biological attribute) as part of the case report form (CRF). All

participants in DIRECT Phase II were East Asian. Patients were diagnosed with MDD based on

ICD 10 or DSM-IV. Healthy controls matched with MDD patients by age, sex ratio, and

educational levels were recruited at each site. All participants provided written informed

consent, and local institutional review boards approved each study from all included cohorts.

The analysis plan of the current study has been reviewed and approved by the Institutional

Review Board of the Institute of Psychology, Chinese Academy of Sciences (No. H21102).

Data will be made available to the public as outlined in the Data Sharing Statement.

Figure 1. Sample characteristics of the DIRECT dataset. (A) Sample sizes of each site; (B) Number of

male/female subjects irrespective of diagnosis; (C) Violin plots depicting the age distribution (in years).

Solid black lines indicate the mean, 25[th], and 75[th] percentiles; (D) Violin plots show education

distribution (in years). Solid black lines indicate the mean, 25[th], and 75[th] percentiles; (E) Violin plots

depicting the distribution of scores of the Hamilton Depression Rating Scale (HAMD). Solid black lines

indicate the mean, 25[th], and 75[th] percentiles.

The second dataset ("TRD-TMS") comprises 25 medication-treatment-resistant MDD (TRD)

patients who underwent 4 to 7 weeks of daily repetitive TMS applied over the left DLPFC.

Patients' TMS sites were recorded using their structural MRI images and a frameless

neuronavigation system. Treatment response was assessed with the 24-item Hamilton

Depression Rating Scale (HAMD). The targets for rTMS stimulation were determined using

the 5.5-cm method. Only TMS outcomes and target coordinates were openly shared for this

dataset. We obtained access to these data from the supplementary materials of Weigand et

al. [38]. For more details on this dataset, please refer to Weigand et al. [38].

The third dataset ("SID-TMS") consists of 28 MDD patients with suicidal ideation who

underwent 10 daily sessions of rTMS over the left DLPFC for 5 consecutive days. Clinical

efficacy was evaluated using a 17-item HAMD. The TMS outcomes, neuroimaging data,

participants' demographic information, and target coordinates for this dataset were made

available upon request, enabling evaluation of the performance of the MDD big data-guided

individualized TMS targeting algorithm. The targets for individualized rTMS stimulation were

determined by identifying the peak subunits in the DLPFC area with the most negative

connections to the sgACC area in the original study. For more details on this dataset, see Li et

al. [53].

The fourth dataset ("CUD-TMS") comprises 27 cocaine use disorder (CUD) patients who

underwent two daily sessions of rTMS treatment over the left DLPFC in an acute phase and

two weekly sessions of rTMS treatment in a maintenance phase. The rTMS treatment was

delivered at the left DLPFC using either the 5.5 cm anatomic criterion or the Beam F3

method. Depressive symptoms were a secondary treatment outcome in the original study. A

subsample of 16 individuals, all with baseline HAMD scores above 7, was used for further

calculation of individualized TMS targets. The TMS outcomes, neuroimaging data,

participants' demographic info, and target coordinates for this dataset were openly shared

(https://openneuro.org/datasets/ds003037/versions/1.0.0). For more details on this dataset,

see Garza-Villarreal et al. [54].

Approach

The study's first objective was to delineate case-control differences in the sgACC-FC profile

and explore its implication in identifying FC-guided individualized TMS targets. Accordingly,

we conducted a generalized linear model (GLM) to compare voxel-wise sgACC-FC maps of

MDD patients and HCs in the DIRECT dataset. We then demonstrated the association

between clinical improvement and group differences in TMS targeting sgACC-FCs by

leveraging the TRD-TMS and SID-TMS datasets. Given that the peak sgACC anticorrelation of

a normative connectome within the left DLPFC was usually selected as the FC-guided sgACC

group target, we showed the impact of case-control differences on such group targets by

separately identifying the peak sgACC anticorrelation in the mean sgACC-FC maps of the

MDD group and HC group from the DIRECT dataset. Finally, we identified individualized

optimal targets using the MDD big data-guided individualized TMS targeting algorithm

guided by statistical maps (e.g., group difference map, mean sgACC-FC maps). We validated

the clinical effectiveness of the individualized approach by computing the correlation

between clinical outcomes and the distance between the actual TMS sites and the identified

individualized targets in the SID-TMS and CUD-TMS datasets. All statistical tests conducted in

the current study were two-sided.

Power calculations for primary hypotheses

The primary outcome of the current study is the case-control differences regarding the

sgACC-FC profiles. Estimates of the effect size (Cohen's d = 0.186) of MDD patients'

abnormalities in FCs are drawn directly from our prior research based on the DIRECT Phase I

dataset [24]. Power calculation was performed using R version 4.3.1 [55] with pwr [56]. A sample of

455 patients will achieve 80% power with a 5% Type I error rate.

Image preprocessing

Acquisition parameters and scanners for all cohorts are provided in Table S2. All R-fMRI and

structural MRI scans were preprocessed at each site using the same DPABISurf protocol, an

R-fMRI data analysis toolbox evolved from DPABI/DPARSF [32,57,58] (For details, see SI). Given

the controversy regarding global signal regression (GSR) and its essential role in identifying

TMS targets [41], we performed preprocessing pipelines with and without GSR.

FC maps of sgACC

Although recent studies have attempted to identify personalized TMS targets using surface-based algorithms [44], most previous studies have reported targets in volume-based MNI space with sgACC ROIs defined as a sphere in volume-based space [36,59]. As a result, we used the volume-based preprocessed imaging data from DPABISurf to better compare our results with the existing literature.

We defined the sgACC as a 10 mm diameter sphere located on the average MNI coordinates based on prior studies showing reduced glucose metabolism or blood flow after receiving an antidepressant treatment (MNI coordinates: x = 6, y = 16, z = -10. For details, please refer to Fox et al., 2012 [36]). The sgACC time series were determined for each individual by spatially averaging the preprocessed R-fMRI time series across all voxels in the abovementioned masks. We then calculated whole-brain FC maps in volume-based MNI space. FC was calculated using Pearson's correlation and underwent Fisher's r-to-z transformation. All FC maps were smoothed with a 6 mm full-width half maximum (FWHM) kernel size. We used ComBat [60] to control potential site and scanner biases (For details, see SI).

Group difference maps of sgACC-FC profiles

We used a voxel-wise GLM to examine differences in the FC maps of sgACC between MDD patients and HCs in DIRECT Phase II. Cohen's f^2 was calculated to characterize the effect sizes of this group difference effect. The GLM model includes age, sex, education, and head motion as covariates:

$$y = \mu + X\beta + \epsilon \qquad (1)$$

where y denotes the FC value of a given voxel from a given participant; μ stands for the constant term; X represents the design matrix for the covariates of interest (diagnosis, age, sex, education, and head motion); β is a vector of regression coefficients corresponding to X; and ϵ is a vector of residuals that follow $N(0, \sigma^2)$. Multiple comparison correction was conducted using false discovery rate (FDR) correction at $q < 0.05$.

To further interpret group difference maps, we extracted the mean FC values of seven

networks using Schaefer's 400 parcellation atlas [61]. A GLM model identical to model (1) was

constructed to characterize case-control differences for each network. Bonferroni multiple

comparison correction was conducted ($p < 0.05/7$). We further explored the effect of the

identified DLPFC clusters in several subgroups. Specifically, patients who were in their first

episode and had never received any antidepressant medication treatment (first episode drug

naïve, FEDN, N = 484) and patients who had undergone more than one episode (recurrent, N

= 439) were selected and compared. Three contrasts, FEDN vs. HC, recurrent vs. HC, and

FEDN vs. recurrent, were analyzed.

Relationship between group differences in TMS targets' sgACC FCs and clinical outcomes

To explore the relationship between group differences in TMS targets' sgACC FCs and clinical

outcomes, we first extracted the mean t-values from the group difference map of 8 mm

radius spheres centered at each targeting coordinate in the TRD-TMS dataset, then examined

the Pearson correlations between these t-values and HAMD score reductions. We

anticipated that greater group differences in sgACC-FC at the target location (i.e., higher t

values) would be related to better TMS therapeutic effects (higher HAMD reductions). To test

the robustness of our findings, we also used spheres with 2 mm, 4 mm, and 10 mm radiuses

to extract the t-values of group differences.

Group targets based on mean sgACC-FC maps

The prior group-level DLPFC TMS target had been derived from a cohort of healthy young

adults [38]; here, we separately averaged whole-brain sgACC-FC maps across all the DIRECT

participants in the MDD and HC groups. We then searched for the peak sgACC anticorrelated

voxel within the DLPFC area (i.e., Brodmann area (BA) 46) as the mean sgACC-FC guided TMS

targets for the MDD and HC groups.

Identification of individualized TMS targets

The reliable statistical maps from the DIRECT big sample best reflect the probability of

MDD-related abnormalities in sgACC-FC. Therefore, we can use these maps to guide the

identification of individualized abnormalities by combining this big-data-based abnormality

information with the individualized R-fMRI data from a given patient, obtaining reliable

statistical maps from the DIRECT sample. We used the dual regression approach to identify

individualized TMS targets guided by group-level statistical maps in the SID-TMS and

CUD-TMS datasets. Dual regression is a common method in independent component analysis

(ICA) for projecting group-level independent components (e.g., functional networks) onto

the individual subject level (see Figure S4 for details). In the first step of the MDD big

data-guided individualized TMS targeting algorithm, a group-level statistical spatial map (e.g.,

the sgACC-FC group difference map reflecting the probability of MDD-related abnormalities

in sgACC-FC) was used as a spatial regressor in the GLM to identify the temporal dynamic of

the group-level map (similar to spatial correlation with the abnormality spatial map). A time

series associated with the spatial map of MDD-related FC abnormalities was generated. In

the second step, the derived time series was used as a temporal regressor in the GLM to

identify an individual-level spatial map (similar to the temporal correlation with the previous

time series). This spatial map can be considered the best-individualized abnormality guided

by big-data-based abnormality. Given our prior knowledge of DLPFC TMS treatment in MDD,

we confined the big-data-based abnormality dual regression to the DLPFC area. That is, we

use the group DLPFC abnormality probability map to find the individualized DLPFC target in a

given MDD patient. The final coordinates for the individualized TMS targets are defined as

the centroids of the largest clusters within this DLPFC region on the individual-level spatial

maps. Additionally, we calculated the individualized target coordinates using the seed map

approach, following the methods described by Fox et al. [41] and Cash et al. [44]. In the seed-map

approach, a seed time series is extracted by computing a weighted average time series of all

voxels within the seed map (e.g., a group average map of sgACC-FC, but excluding the DLPFC

area). Subsequently, Pearson's correlation coefficients are computed between this extracted

time series and all other DLPFC voxels. The final TMS target is the most negatively

functionally connected cluster in the DLPFC area. Of note, in the seed map approach, the

goal of the first step is to find the most sgACC-like time series, which is not confined to the

noisy sgACC area. Following this rationale, the DLFPC time series should not be included to

avoid biasing the estimation of the sgACC-like time series. Thus, the DLPFC area was

excluded. Thus, the exclusion and inclusion of DLPFC differs between the seed map approach

and the DR approach due to the different underlying rationales. Details of the individualized

TMS target localization algorithms are provided in the supplementary materials.

Clinical efficacy of the group-level and individualized TMS targets

We leveraged the SID-TMS and CUD-TMS datasets to evaluate the clinical significance of

individualized TMS targets. We identified the proposed individualized TMS targets from the

MDD big data-guided individualized TMS targeting algorithm and calculated the targeting

offset (i.e., Euclidean distance between the individualized optimal TMS targets and the actual

stimulation coordinates) for each patient. Subsequently, we calculated the Pearson

correlations between clinical improvement (i.e., HAMD reductions) and targeting offset. We

anticipated a negative correlation between clinical outcomes and target offset (i.e., the

closer the actual stimulation target was to the individualized target from the MDD big

data-guided individualized TMS targeting algorithm, the higher the clinical improvement).

Age, sex, and head motion were included as covariates in the regression models when

calculating correlations between targeting offsets and clinical improvement.

Results

Group difference maps of sgACC-FC

In the large-scale DIRECT Phase II dataset, we found significant MDD-related

hyperconnectivity with the sgACC in bilateral DLPFC, temporal parietal junction, and occipital

lobe, as well as hypoconnectivity in the bilateral temporal lobe, left inferior frontal gyrus, and

left postcentral gyrus when preprocessing included GSR (Figure 2A). When GSR was not

included in preprocessing, MDD-related sgACC FC alterations showed predominantly

hypoconnectivity. Such abnormally decreased FCs were found across the central gyrus,

occipital lobe, insular cortex, temporal lobe, and a small portion of the frontal lobe. Without

GSR, MDD-related hyperconnectivity was limited to subcortical regions (Figure S1A). Given

that significant case-control differences in the DLPFC area were revealed only when GSR was

implemented, subsequent analyses were based on results with GSR. The uncorrected group

difference maps calculated in the volume space showed remarkable similarity with those in

the surface space (Figure S2). Network-wise FC analyses showed that MDD patients' FC

between sgACC and the limbic network (LN) was significantly reduced compared to HC

($t(2880)$ = -4.122, $p_{corrected}$ < 0.001, Cohen's d = 0.171). The FC between sgACC and the

frontoparietal network (FPN) was enhanced and approached significance ($t(2880)$ = 2.419,

$p_{corrected}$ = 0.055, Cohen's d = 0.090, Figure 2B). Without GSR, MDD patients showed

decreased FC between sgACC and all brain networks (all $p_{corrected}$ < 0.05) except for the FPN

(Figure S1B). We identified two contiguous clusters of voxels that showed significant group

differences in the left DLPFC. Group difference cluster 1 (MNI coordinates: x = -44, y = 38, z =

32; $t(2880)$ = 3.277, p < 0.001, Cohen's d = 0.141) was ventral to group difference cluster 2

(MNI coordinates: x = -34, y = 36, z = 40; $t(2880)$ = 3.670, p < 0.001, Cohen's d = 0.126)

(Figure 2C). In subgroup analyses, when GSR was performed, FEDN patients showed

enhanced FCs in both clusters (cluster 1: $t(1790)$ = 2.282, p = 0.023, Cohen's d = 0.124;

cluster 2: $t(1790)$ = 2.273, p = 0.023, Cohen's d = 0.123). There was a significant

enhancement in the DLPFC cluster 2 between the recurrent MDD patients and HCs ($t(1745)$ =

2.765, p = 0.006, Cohen's d = 0.159) while cluster 1 approached significance ($t(1745)$ = 1.864,

p = 0.063, Cohen's d = 0.107). No significant difference was revealed between the FEDN and

recurrent patients (see Figure 3).

Figure 2. Group differences of subgenual anterior cingulate cortex (sgACC) functional connectivity (FC) profiles are related to TMS treatment efficacy, demonstrating clinical significance. (A) Two-sample *t*-test maps of MDD-related sgACC FC abnormalities with global signal regression (GSR) implemented. (B) Group differences of FCs between sgACC and visual network (VN), somatomotor network (SMN), dorsal attention network (DAN), ventral attention network (VAN), limbic network (LN), frontoparietal network (FPN), and default mode network (DMN). (C) Two clusters showed significant case-control differences in sgACC-FC. Abbreviations: DLPFC, dorsal lateral prefrontal cortex; L, left hemisphere; R, right hemisphere. *: significant after Bonferroni correction; #: approaching significance.

Figure 3. Subgroup differences regarding two clusters found in DLPFC. Abbreviations: FEDN, first

episode drug naïve; HC, healthy control.

sgACC anticorrelation peaks in MDD and HCs

Our results highlighted the case-control differences in sgACC-FC profiles. Since the prior

sgACC group target (MNI coordinates: x = −42, y = 44, z = 30) had been based on a cohort of

young, healthy adults [38], we sought to examine potential differences in anticorrelation peaks

extracted from the mean sgACC FC maps of MDD and HC groups (Figure 4B-C) in the DIRECT

dataset. We found that the anticorrelation peak of MDD patients (MNI coordinates: x = -40, y

= 50, z = 22) differed from that of HCs (MNI coordinates: x = -42, y = 38, z = 32), probably due

to abnormal FCs within the left DLPFC in MDD patients. The anticorrelation peak extracted

from the DIRECT HCs was closer to the previously reported locus [38] (Figure 4).

Figure 4. The peak of the sgACC anticorrelation in MDD patients differed from that in HCs.

sgACC-FC group differences correlate with TMS treatment outcomes

Using the TRD-TMS dataset, we examined the relationship between group differences

(*t*-values) and clinical outcomes (HAMD reductions) to test the clinical relevance of the group

difference maps. Group differences were positively correlated with HAMD score reductions

($r(23) = 0.448$, $p = 0.025$, Figure 5A), suggesting that group-level difference maps may be

useful for enhancing the outcomes of TMS for treating MDD by improving target localization.

Significant associations were also observed using different radius settings (Figure S11).

The same trend was observed in the SID-TMS dataset, albeit it failed to achieve significance

($r(26) = 0.304$, $p = 0.108$, Figure 5).

Figure 5. The group difference regarding sgACC-FC to the TMS targets was correlated with clinical

efficacy. (A) The TMS targets were extracted from the TRD-TMS dataset [38]. The sizes of the spheres

indicate the magnitudes in Hamilton Depression Rating Scale (HAMD) reductions, with the group

difference map rendered on the surface. The scatter plot depicted that the magnitudes of the group

differences in the FC between TMS targets and sgACC were positively related to clinical improvements

in the TRD dataset. (B) Findings were replicated in the SID-TMS dataset [53]. Abbreviations: FC,

functional connectivity; TMS, transcranial magnetic stimulation.

Individualized targets suggest higher clinical efficacy than group targets

We utilized the MDD big data-guided individualized TMS targeting algorithm to calculate

individualized optimal targets for each SID-TMS and CUD-TMS dataset participant. Clinical

relevance of the targets was determined by the correlation between target offset and clinical

improvement. The individualized target locations derived from the sgACC-FC group

difference map are illustrated in Figure 6. The individualized targets (SID-TMS: $r(26) = -0.562$,

$p = 0.002$; CUD-TMS: $r(14) = -0.511$, $p = 0.037$) outperformed their corresponding group-level

targets (SID-TMS: $r(26) = -0.349$, $p = 0.044$; CUD-TMS: $r(14) = -0.167$, $p = 0.293$) in both

datasets (Figure 6). The DR-based individualized targets derived from the MDD or HC

group-average sgACC-FC maps also outperformed their corresponding group-level targets

and seed map-based targets (Figure S6-8). Among all the individualized targets, the DR-based

targets guided by the group difference map achieved the highest clinical efficacy.

Figure 6. Individualized targets derived from the group difference map exhibited greater clinical

efficacy than the corresponding group targets. Clinical efficacy was characterized by computing the

correlations between the offset distances of TMS targets and the clinical improvements observed in

the CUD-TMS dataset. The TMS targeting offset distance was defined as the Euclidean distance

between the actual rTMS stimulation coordinates and the individualized or group targets. Clinical

improvement was defined as the HAMD reduction during rTMS treatment, adjusted for age, sex, and

head motion. The locations of the targets are displayed on the cortex. The sizes of the spheres indicate

the magnitudes of Hamilton Depression Rating Scale (HAMD) reductions. (A-B) Clinical efficacy of the

individualized and group targets in the SID-TMS sample. (C-D) Clinical efficacy of the individualized and

group targets in the CUD-TMS sample. *$p < 0.05$. ** $p < 0.01$.

Discussion

In the present study, we leveraged a large multi-site fMRI sample (1660 MDD patients and

1341 HCs) and three independent TMS datasets to delineate abnormalities in sgACC FC in

MDD and explore their potential impact on the localization of TMS targets. Specifically, with

GSR implemented, we found enhanced FCs between sgACC and left DLPFC, bilateral

supplementary motor areas and inferior parietal lobes, thalamus, and visual areas, and

decreased FCs between sgACC and left anterior insula, left superior temporal lobe, and

bilateral temporal poles in MDD patients. Patients with MDD exhibited significantly reduced

FC between the sgACC and the DMN, while FC between the sgACC and the FPN was only

marginally increased. Crucially, we showed that the clinical outcomes of TMS treatments

were related to the magnitude of the case-control differences in the FCs between sgACC and

TMS targets. Furthermore, such group difference profiles altered the position of the sgACC

anti-correlation peak in the left DLPFC. Additionally, the MDD big data-guided individualized

TMS targeting algorithm to identify individualized TMS targets showed better clinical efficacy

than TMS targets based on group sgACC-FC profiles.

MDD-related FC abnormalities of sgACC

Our results add to a growing literature documenting functional network abnormalities

involving the sgACC in MDD [9,11-13,17,19,62-64]. Nevertheless, notable discrepancies in the type of

abnormality (enhanced/reduced) and specific brain regions showing altered sgACC FCs have

been reported. Considering the small effect sizes (Cohen's $f^2 < 0.01$) of MDD-related

sgACC-FC abnormalities, the limited sample sizes in previous studies entail a high risk of false

positive findings [21,65,66]. Another potential source of heterogeneity in previous findings may

be whether or not they applied GSR (See supplementary materials for detailed discussion).

With an unprecedented sample size, our results provide among the most robust evidence to

date. Specifically, MDD patients showed enhanced sgACC-thalamus FC and decreased

sgACC-limbic network FC regardless of whether GSR was implemented. These results are

consistent with previous studies in adolescents [19] and adults [59] with MDD. Previous studies

have reported abnormal FCs between sgACC and limbic areas and some subcortical regions,

such as the amygdala [9,20] and parahippocampus regions [13]. Together, the present results are

consistent with a model highlighting sgACC as a critical hub in an "extended medial network,"

which also encompasses limbic, thalamic, and striatal regions and plays a key role in the

pathophysiology of MDD [67,68]. This "extended medial network" overlaps substantially with

the DMN. Indeed, decreased FCs between sgACC and DMN regions, such as the medial

prefrontal cortex, precuneus, temporal gyrus, and parahippocampus regions, were revealed

in MDD relative to HCs when GSR was not implemented. Such abnormalities have been

previously reported [10,12,15,34], which led to the hypothesis that abnormally enhanced FC

between sgACC and DMN are the network underpinnings of rumination [69]. However,

contrary to the aforementioned hypothesis, we found reduced, instead of enhanced FC

between sgACC and DMN. Similarly, in the first phase of DIRECT, we demonstrated that MDD

was characterized by reduced FC within DMN [23,24]. We note that the first and second phases

of the DIRECT data are solely comprised of Chinese samples, while most studies that have

reported enhanced sgACC-DMN FCs have been in Caucasian samples. Different prevalence

rates [70,71], heterogeneous symptoms [72], and different risk alleles [73] have been reported in

Caucasian and Eastern Asian groups. Accordingly, we cannot exclude racial differences

contributing to this discrepancy. It is worth noting that we found enhanced FCs between the

visual region and sgACC in MDD patients relative to HCs when GSR was implemented, while

significantly reduced sgACC-visual region FCs were revealed when GSR was not performed.

Most of DIRECT II sites' R-fMRI data were collected with participants' eyes closed (22 out of

23 sites). Prior research had shown that participants are more likely to fall asleep when their

eyes are closed during data acquisition, and drowsiness may alter FC patterns in visual

regions [74]. Thus, it is possible that abnormalities in visual region FCs may be due to MDD

patients' lower levels of wakefulness.

Clinical relevance of abnormal sgACC-DLPFC anticorrelation in MDD

Once the group difference maps of sgACC FC profiles were delineated, we further explored

the impact of such an abnormality, especially in DLPFC, on identifying TMS targets. Most

clinical trials have focused on applying TMS to the left DLPFC based on the hypothesis that

high-frequency rTMS will enhance hypoactivity during depressive episodes [8,75]. The DLPFC is

anatomically extensive [76]. However, which DLPFC sub-field is the best target for TMS remains

unclear. The current FDA-approved protocol (i.e., the "5 cm" method) leads to large

interindividual variation in stimulation sites, which may contribute to the heterogeneity in

the effect sizes of antidepressant responses in prior trials [77,78].

Previous targeting approaches leveraging anatomical landmarks has not consistently

outperformed the "5 cm" method or the F3 Beam method [79,80]. Considering the

unsatisfactory effect of TMS target localization based on brain anatomical parcellation,

group-level normative sgACC anticorrelation peaks based on healthy population datasets

have been frequently used as TMS targets in recent years [36,38,41]. However, in the present

study, we found that the locations of such anticorrelated peaks differ substantially between

MDD and HC samples when measured in a large clinical cohort. Therefore, it might be

problematic to identify TMS targets based solely on sgACC-FC profiles in healthy or

depressed individuals. Intriguingly, we found that the magnitudes of case-control differences

in TMS targets' FCs to sgACC were positively related to the clinical improvements after

receiving rTMS to the left DLPFC. Such correlation implies that the case-control differences in

the FC between sgACC and the left DLPFC might bear important information that could be

leveraged to guide the identification of reliable, individualized TMS targets.

The MDD big data-guided individualized TMS targeting algorithm may improve the clinical

efficacy of TMS targets

In the current study, we developed an MDD big data-guided individualized TMS targeting

algorithm to individualize the TMS targets derived from group-level statistical maps. The

proposed approach takes advantage of the high signal-to-noise ratio and reliability of large

sample statistical maps while integrating individual spontaneous brain activity of individuals

with MDD. Most existing individualized TMS target localization algorithms are based on

calculating sgACC FCs using densely sampled MRI images from single subjects [81]. However,

individual MRI images tend to be noisy and unreliable [82,83], especially in the sgACC region. Air

in the sinuses often introduces susceptibility artifacts due to the different magnetic

properties of air and brain tissue. Signal loss and geometric distortion are common in areas

close to air-filled sinuses, such as the inferior frontal cortex, including the sgACC [84]. The seed

map approach has been proposed to alleviate such difficulties due to the subpar image

quality of the sgACC region [41,44]. In the seed map approach, all voxels within the seed map

(except for the DLPFC region) were used to extract the seed time series to improve its

signal-to-noise ratio. However, considering that the weight (e.g., FC value) of the sgACC area

is usually extremely high in the seed map, the derived seed time series remains somewhat

similar to the noisy sgACC time series and doesn't achieve the best TMS localization. A

cutting-edge MDD TMS therapy combining FC-guided target localization, high dose, and

intermittent theta-burst stimulation (iTBS) was reported to be highly effective in a

randomized, double-blinded, sham-controlled clinical trial [45,46]. However, the targeting

algorithm relied on hierarchical agglomerative clustering in the sgACC area which has a low

signal-to-noise ratio. Nevertheless, the final target was still determined according to

individual-level sgACC-DLPFC FCs. For the proposed MDD big data-guided individualized TMS

targeting algorithm, we view the reliable statistical maps from the DIRECT big sample as the

best reflection of the probability of MDD-related abnormalities in sgACC-FC. Therefore, we

used these maps to guide the identification of individualized abnormalities by combining this

big-data-based abnormality information with the individualized R-fMRI data from a given

patient. Given the a priori knowledge of DLPFC TMS treatment in MDD, we confined the

big-data-based abnormality dual regression only within the DLPFC area, which is less affected

by susceptibility artifacts. In this way, the superior signal quality of DLPFC and the effective

and reliable properties of the dual regression algorithm enhance the accuracy of target

localization.

Encouragingly, the DR-based individualization targets enhanced the clinical significance of

corresponding group-level targets, regardless of the template used. This result supports the

generalizability and extensibility of the algorithm, offering the potential for TMS targeting

based on other circuits and biomarkers. Considering most existing TMS research still relies on

the traditional 5 cm or Beam F3 methods for targeting, one approach based on our data to

improve targeting would be simply shifting the target to a more anterior and lateral position.

However, such a simple shift was not supported by the present study. Instead, 25% of targets

individualized using the group difference map were more medial than the original targets,

and 50% of targets individualized by the group difference map were more posterior than the

original targets. Therefore, the MDD big data-guided individualized TMS targeting algorithm

does not simply set a more anterior and lateral coordinate in the BA46 area as the

individualized optimization target for subjects. Among the group-level templates used in the

MDD big data-guided individualized TMS targeting algorithm, the sgACC-FC group difference

map performed the best, instead of the commonly used average sgACC based on healthy

individuals. This may reflect the abnormal posterior shift of the MDD sgACC anticorrelation

peak we found, and it emphasizes the immense clinical value of examining spontaneous

brain activity differences between MDD and HC in a large sample. Previous studies either

included only a small quantity of MDD functional MRI data [46,47,85] or developed targeting

algorithms based on large-scale HC samples [49]. In this study, we utilized an unprecedented

amount of functional MRI data from MDD and HC, obtained a reliable group-level difference

map of sgACC-FC, and successfully validated its potential in treating MDD with TMS.

Limitations

Several limitations need to be considered. First, we noted that the inconsistency between

our results and the broader literature could partly be due to racial differences of the samples

(i.e., Eastern Asian vs. Caucasian). Efforts that intend to pool existing neuroimaging data

worldwide, such as the Enhancing NeuroImaging Genetics through Meta-Analysis (ENIGMA)

Major Depressive Disorder (MDD) consortium [25], have accumulated large-scale data primarily

from Western countries and published several high-impact studies delineating MDD-related

anatomical abnormalities [86,87]. Planned collaborations between the DIRECT and the

ENIGMA-MDD consortiums are ongoing to help address potential cultural, genetic, and

environmental mechanisms in more diverse groups of MDD patients. Second, we performed

volume-based preprocessing to facilitate comparison with the previous literature.

Surface-based preprocessing strategies have provided more accurate and detailed

representations of cortical and subcortical structures [88]. Recent research has begun to

explore surface-based rTMS target identification algorithms and has shown promising clinical

relevance [39,40,89]. Future research should consider using surface-based case-control difference

maps to further refine ways to identify TMS targets. Given the established involvement of

the left DLPFC in existing TMS treatment protocols, the present study restricted investigation

to within this area.

We noted that other brain regions (e.g., angular gyrus and supplementary motor area)

showed significant case-control differences worth further research and may serve as

potential targets for neuromodulation [90,91 92]. The MDD big data-guided individualized TMS

targeting algorithm can be readily transferred to other neural circuits or other brain

imaging-derived feature maps (e.g., ICA, functional gradient, normative modeling). The

clinical efficacy of these alternative targets is worthy of future investigations. Identifying a

reliable personalized TMS target solely based on an individual's R-fMRI data (around 8 mins

of fMRI scan in clinical practice) is challenging. Due to the poor replicability of FC [93], existing

individual-level network parcellation algorithms need a large quantity of fMRI images

(usually more than one hour of scanning time) [94]. The present study utilized three

independent TMS samples to validate the efficacy of the individualized algorithm.

Nevertheless, the two TMS datasets used to validate the individualized TMS targets are

limited in sample size [37,95]. Publicly available TMS brain imaging datasets could be used for

independent validation of target localization algorithms to reduce the false positive rate;

however, access to such datasets remains challenging. In addition, prospective, double-blind

clinical trials are warranted to compare the treatment outcome across different rTMS

targeting algorithms (e.g., traditional anatomical landmark-based targeting, group-level

sgACC-FC targeting, and MDD big data-guided individualized TMS targeting algorithm).

Therefore, we call upon researchers involved in this field to publicly share data on TMS

targets, clinical efficacy, and brain imaging, and we will also openly share data from our

related prospective studies [96]. The large sample size of the DIRECT consortium aggregated

dataset allows for intriguing analyses, such as bio-subgroups of MDD patients. Indeed, some

previous DIRECT studies have shown that MDD patients can be subgrouped [30,97]. Future

studies may further determine whether bio-types could be achieved using sgACC [91]. Most of

the DIRECT II R-fMRI data were acquired when participants were instructed to close their

eyes, which has been shown to be associated with an increased likelihood of sleep during

scanning [74]. Instructing participants to keep their eyes open and look at fixation can help

prevent participants from falling asleep and is easy to apply. Since large multi-site R-fMRI

data aggregation endeavors such as DIRECT are prone to be biased by non-neurophysical

factors such as head motion, sleepiness, etc. [98], it is important to prospectively apply

well-designed standard operation procedures in future large-scale multi-site scientific

projects.

Conclusion

In summary, we leveraged a large sample of MDD patients to fully delineate group

differences in sgACC-FC maps between MDD patients and HCs. We next demonstrated the

impact of such case-control differences on group TMS targets based on sgACC-FC profiles by

showing that the magnitudes of case-control differences in the FC between sgACC and TMS

targets were positively associated with clinical outcomes and the peak sgACC anticorrelation

locations were different in MDD patients as compared to HCs. Moreover, we developed an

MDD big data-guided individualized TMS targeting algorithm to identify individualized TMS

targets and demonstrated that this approach may improve clinical efficacy compared to

group targets based on sgACC-FC profiles.

Contributors

C-GY, Y-FZ, X-NZ, XC, and BL conceived and designed the study. C-GY, Y-FZ, J-PZ, and X-NZ coordinated the collaboration. JQ, LK, T-MS, TL, K-RZ, Z-NL, L-PC, JY, X-PW, Y-GY, C-YW, C-MX, G-RX, Y-SL, Y-QY, XW, YW, X-FX, Y-QC, Q-YG, W-BG, J-PZ, YH, H-NW, B-JL, WZ, and J-PL supervised data collection. X-RW, QH, Y-KW, HY, A-XZ, Y-CL, J-SC, P-FS, X-YL, FL, C-CH, X-LC, F-NJ, J-JZ, X-LJ, G-MC, Z-SC, T-LC, X-XS, TC, B-JL, M-LY, Z-PX and BL organized the data. XC and C-GY analyzed and interpreted the data. BL and C-GY designed the dual regression-based TMS targeting algorithm. H-NW and B-JL collected the SID-TMS dataset. XC, BL, and C-GY drafted the manuscript. All authors revised the manuscript for important intellectual content and approved the final submitted version. XC, BL, and C-GY accessed and verified the data. XC, BL, and C-GY had full access to all the data in the study. C-GY had final responsibility for the decision to submit for publication. All authors were responsible for the final decision to submit for publication and have seen and approved the manuscript.

Conflict of interest

DMB receives research support from the Canadian Institutes of Health Research (CIHR), National Institutes of Health – US (NIH), Brain Canada Foundation, and the Temerty Family through the CAMH Foundation and the Campbell Family Research Institute. He received research support and in-kind equipment support for an investigator-initiated study from Brainsway Ltd., and he was the site principal investigator for three sponsor-initiated studies for Brainsway Ltd. He received in-kind equipment support from Magventure for investigator-initiated studies. He received medication supplies for an investigator-initiated trial from Indivior. He has participated in an advisory board for Janssen. He has participated in an advisory board for Welcony Inc. No other conflicts of interest.

Inclusion and Ethics

Local researchers were included throughout the research process. Research protocols have been approved by local ethics review committees. No potential risk was involved in the current research, and all local researchers have discussed and approved the research protocol. All participants provided written informed consent, and local institutional review boards approved each study from all included cohorts. The analysis plan of the current study has been reviewed and approved by the Institutional Review Board of the Institute of Psychology, Chinese Academy of Sciences (No. H21102).

Data Sharing

According to the success of the data sharing model of DIRECT Phase I data (REST-meta-MDD, http://rfmri.org/REST-meta-MDD), DIRECT Phase II data will also have 2 sharing stages. 1) Stage 1: coordinated sharing upon the publication of this announcing manuscript. To reduce conflict among the researchers, the consortium will review and coordinate the proposals submitted by interested researchers. The interested researchers first send a letter of intent to rfmrilab@gmail.com. Then, the consortium will send all the approved proposals to the applicant. The applicant should submit a new innovative proposal while avoiding conflict with approved proposals. The consortium would review and approve this proposal if there is no conflict. Once approved, this proposal would enter the pool of approved proposals and prevent future conflict. 2) Stage 2: unrestricted sharing after January 1, 2026. The researchers can perform any analyses of interest while not violating ethics. All codes have been made openly available (https://github.com/Chaogan-Yan/PaperScripts/tree/master/Chen_2023).

Acknowledgments

This work was funded by the Sci-Tech Innovation 2030 - Major Project of Brain Science and

Brain-inspired Intelligence Technology (No. 2021ZD0200600), the National Key R&D Program

of China (No. 2017YFC1309902), the National Natural Science Foundation of China (No.

82122035, No. 81671774 and No. 81630031), the Key Research Program of the Chinese

Academy of Sciences (No. ZDBS-SSW-JSC006), Beijing Nova Program of Science and

Technology (No. Z191100001119104 and 20230484465), Beijing Natural Science Foundation

(J230040), the Scientific Foundation of Institute of Psychology, Chinese Academy of Sciences

(No. E2CX4425YZ and No. Y9CX422005), the China Postdoctoral Science Foundation (No.

2019M660847), the China National Postdoctoral Program for Innovative Talents (No.

BX20200360), the Special Research Assistant Program of the Chinese Academy of Sciences

(No. E2CX0624) and the Key R&D Program of Sichuan Province (No. 2023YFS0076). The

support provided by the China Scholarship Council (CSC, No. 202104910248) during a visit of

Xiao Chen to the Centre for Addiction and Mental Health is acknowledged. In addition, we

would like to acknowledge the valuable insights provided by Prof. B.T. Thomas Yeo, whose

comments helped to shape our understanding of the data and refine our analysis.

References

1.   Malhi GS, Mann JJ. Depression. *Lancet (London, England)* 2018; **392**(10161): 2299-312.
2.   Drevets WC, Price JL, Furey ML. Brain structural and functional abnormalities in mood
disorders: implications for neurocircuitry models of depression. *Brain structure & function*
2008; **213**(1-2): 93-118.
3.   Disner SG, Beevers CG, Haigh EAP, Beck AT. Neural mechanisms of the cognitive model of
depression. *Nat Rev Neurosci* 2011; **12**(8): 467-77.
4.   Mayberg HS. Modulating dysfunctional limbic-cortical circuits in depression: towards
development of brain-based algorithms for diagnosis and optimised treatment. *British*
medical bulletin 2003; **65**: 193-207.
5.   Mayberg HS, Lozano AM, Voon V, et al. Deep brain stimulation for treatment-resistant
depression. *Neuron* 2005; **45**(5): 651-60.
6.   Cash RFH, Zalesky A. Personalized and Circuit-Based Transcranial Magnetic Stimulation:
Evidence, Controversies, and Opportunities. *Biol Psychiatry* 2024; **95**(6): 510-22.
7.   Cash RFH, Weigand A, Zalesky A, et al. Using Brain Imaging to Improve Spatial Targeting
of Transcranial Magnetic Stimulation for Depression. *Biol Psychiatry* 2020.
8.   Fitzgerald PB. Targeting repetitive transcranial magnetic stimulation in depression: do
we really know what we are stimulating and how best to do it? *Brain Stimul* 2021; **14**(3):

730-6.

9.   Connolly CG, Wu J, Ho TC, et al. Resting-State Functional Connectivity of Subgenual
Anterior Cingulate Cortex in Depressed Adolescents. *Biological Psychiatry* 2013; **74**(12):
898-907.

10.   Greicius MD, Flores BH, Menon V, et al. Resting-state functional connectivity in major
depression: abnormally increased contributions from subgenual cingulate cortex and
thalamus. *Biol Psychiatry* 2007; **62**(5): 429-37.

11.   Wu H, Sun H, Xu J, et al. Changed Hub and Corresponding Functional Connectivity of
Subgenual Anterior Cingulate Cortex in Major Depressive Disorder. *Front Neuroanat* 2016; **10**:
120.

12.   Davey CG, Harrison BJ, Yucel M, Allen NB. Regionally specific alterations in functional
connectivity of the anterior cingulate cortex in major depressive disorder. *Psychol Med* 2012;
42(10): 2071-81.

13.   Cheng B, Meng Y, Zuo Y, et al. Functional connectivity patterns of the subgenual anterior
cingulate cortex in first-episode refractory major depressive disorder. *Brain imaging and*
behavior 2021; **15**(5): 2397-405.

14.   Sheline YI, Price JL, Yan Z, Mintun MA. Resting-state functional MRI in depression
unmasks increased connectivity between networks via the dorsal nexus. *Proc Natl Acad Sci U*
S A 2010; **107**(24): 11020-5.

15.   Berman MG, Peltier S, Nee DE, Kross E, Deldin PJ, Jonides J. Depression, rumination and
the default network. *Social cognitive and affective neuroscience* 2011; **6**(5): 548-55.

16.   Wu X, Lin P, Yang J, Song H, Yang R, Yang J. Dysfunction of the cingulo-opercular network
in first-episode medication-naive patients with major depressive disorder. *J Affect Disord*
2016; **200**: 275-83.

17.   Sudheimer K, Keller J, Gomez R, et al. Decreased hypothalamic functional connectivity
with subgenual cortex in psychotic major depression. *Neuropsychopharmacology* 2015; **40**(4):
849-60.

18.   Davey CG, Yücel M, Allen NB, Harrison BJ. Task-related deactivation and functional
connectivity of the subgenual cingulate cortex in major depressive disorder. *Frontiers in*
psychiatry 2012; **3**: 14.

19.   Gaffrey MS, Luby JL, Repovš G, et al. Subgenual cingulate connectivity in children with a
history of preschool-depression. *Neuroreport* 2010; **21**(18): 1182-8.

20.   Ho TC, Yang G, Wu J, et al. Functional connectivity of negative emotional processing in
adolescent depression. *J Affect Disord* 2014; **155**: 65-74.

21.   Button KS, Ioannidis JPA, Mokrysz C, et al. Power failure: why small sample size
undermines the reliability of neuroscience. *Nat Rev Neurosci* 2013; **14**(5): 365-76.

22.   Algermissen J, Mehler DMA. May the power be with you: are there highly powered
studies in neuroscience, and how can we get more of them? *J Neurophysiol* 2018; **119**(6):
2114-7.

23.   Chen X, Lu B, Li H-X, et al. The DIRECT consortium and the REST-meta-MDD project:
towards neuroimaging biomarkers of major depressive disorder. *Psychoradiology* 2022; **2**(1):
32-42.

24.   Yan CG, Chen X, Li L, et al. Reduced default mode network functional connectivity in

patients with recurrent major depressive disorder. *Proc Natl Acad Sci U S A* 2019.

25.   Schmaal L, Pozzi E, T CH, et al. ENIGMA MDD: seven years of global neuroimaging

studies of major depression through worldwide data sharing. *Transl Psychiatry* 2020; **10**(1):

172.

26.   Yang H, Chen X, Chen ZB, et al. Disrupted intrinsic functional brain topology in patients

with major depressive disorder. *Mol Psychiatry* 2021.

27.   Long Y, Cao H, Yan C, et al. Altered resting-state dynamic functional brain networks in

major depressive disorder: Findings from the REST-meta-MDD consortium. *Neuroimage Clin*

2020; **26**: 102163.

28.   Ding YD, Yang R, Yan CG, et al. Disrupted hemispheric connectivity specialization in

patients with major depressive disorder: Evidence from the REST-meta-MDD Project. *J Affect*

Disord 2021; **284**: 217-28.

29.   Deng K, Yue JH, Xu J, et al. Impaired robust interhemispheric function integration of

depressive brain from REST-meta-MDD database in China. *Bipolar disorders* 2021.

30.   Liang S, Deng W, Li X, et al. Biotypes of major depressive disorder: Neuroimaging

evidence from resting-state default mode network patterns. *Neuroimage Clin* 2020; **28**:

102514.

31.   Liu PH, Li Y, Zhang AX, et al. Brain structural alterations in MDD patients with

gastrointestinal symptoms: Evidence from the REST-meta-MDD project. *Progress in*

neuro-psychopharmacology & biological psychiatry 2021; **111**: 110386.

32.   Yan C-G, Wang X-D, Lu B. DPABISurf: data processing & analysis for brain imaging on

surface. *Science Bulletin* 2021.

33.   Ge R, Downar J, Blumberger DM, Daskalakis ZJ, Vila-Rodriguez F. Functional connectivity

of the anterior cingulate cortex predicts treatment outcome for rTMS in treatment-resistant

depression at 3-month follow-up. *Brain Stimul* 2020; **13**(1): 206-14.

34.   Cash RFH, Cocchi L, Anderson R, et al. A multivariate neuroimaging biomarker of

individual outcome to transcranial magnetic stimulation in depression. *Hum Brain Mapp*

2019; **40**(16): 4618-29.

35.   Philip NS, Barredo J, van 't Wout-Frank M, Tyrka AR, Price LH, Carpenter LL. Network

Mechanisms of Clinical Response to Transcranial Magnetic Stimulation in Posttraumatic

Stress Disorder and Major Depressive Disorder. *Biol Psychiatry* 2018; **83**(3): 263-72.

36.   Fox MD, Buckner RL, White MP, Greicius MD, Pascual-Leone A. Efficacy of transcranial

magnetic stimulation targets for depression is related to intrinsic functional connectivity with

the subgenual cingulate. *Biol Psychiatry* 2012; **72**(7): 595-603.

37.   Elbau IG, Lynch CJ, Downar J, et al. Functional Connectivity Mapping for rTMS Target

Selection in Depression. *Am J Psychiatry* 2023; **180**(3): 230-40.

38.   Weigand A, Horn A, Caballero R, et al. Prospective Validation That Subgenual

Connectivity Predicts Antidepressant Efficacy of Transcranial Magnetic Stimulation Sites. *Biol*

Psychiatry 2018; **84**(1): 28-37.

39.   Cash RFH, Cocchi L, Lv J, Fitzgerald PB, Zalesky A. Functional Magnetic Resonance

Imaging-Guided Personalization of Transcranial Magnetic Stimulation Treatment for

Depression. *JAMA psychiatry* 2021; **78**(3): 337-9.

40.   Siddiqi SH, Weigand A, Pascual-Leone A, Fox MD. Identification of Personalized

Transcranial Magnetic Stimulation Targets Based on Subgenual Cingulate Connectivity: An

Independent Replication. *Biol Psychiatry* 2021; **90**(10): e55-e6.

41.   Fox MD, Liu H, Pascual-Leone A. Identification of reproducible individualized targets for

treatment of depression with TMS based on intrinsic connectivity. *Neuroimage* 2013; **66**:

151-60.

42.   Gordon EM, Nelson SM. Three types of individual variation in brain networks revealed

by single-subject functional connectivity analyses. *Current Opinion in Behavioral Sciences*

2021; **40**: 79-86.

43.   Doucet GE, Lee WH, Frangou S. Evaluation of the spatial variability in the major

resting-state networks across human brain functional atlases. *Hum Brain Mapp* 2019; **40**(15):

4577-87.

44.   Cash RFH, Cocchi L, Lv J, Wu Y, Fitzgerald PB, Zalesky A. Personalized connectivity-guided

DLPFC-TMS for depression: Advancing computational feasibility, precision and reproducibility.

Hum Brain Mapp 2021.

45.   Cole EJ, Phillips AL, Bentzley BS, et al. Stanford Neuromodulation Therapy (SNT): A

Double-Blind Randomized Controlled Trial. *Am J Psychiatry* 2021: appiajp202120101429.

46.   Cole EJ, Stimpson KH, Bentzley BS, et al. Stanford Accelerated Intelligent

Neuromodulation Therapy for Treatment-Resistant Depression. *Am J Psychiatry* 2020; **177**(8):

716-26.

47.   Williams NR, Sudheimer KD, Bentzley BS, et al. High-dose spaced theta-burst TMS as a

rapid-acting antidepressant in highly refractory depression. *Brain* 2018; **141**(3): e18.

48.   Cash RFH, Zalesky A, Thomson RH, Tian Y, Cocchi L, Fitzgerald PB. Subgenual Functional

Connectivity Predicts Antidepressant Treatment Response to Transcranial Magnetic

Stimulation: Independent Validation and Evaluation of Personalization. *Biol Psychiatry* 2019;

86(2): e5-e7.

49.   Cash RFH, Cocchi L, Lv J, Wu Y, Fitzgerald PB, Zalesky A. Personalized connectivity-guided

DLPFC-TMS for depression: Advancing computational feasibility, precision and reproducibility.

Hum Brain Mapp 2021; **42**(13): 4155-72.

50.   Ojemann JG, Akbudak E, Snyder AZ, McKinstry RC, Raichle ME, Conturo TE. Anatomic

localization and quantitative analysis of gradient refocused echo-planar fMRI susceptibility

artifacts. *Neuroimage* 1997; **6**(3): 156-67.

51.   Beckmann CF, Mackay CE, Filippini N, Smith SM. Group comparison of resting-state FMRI

data using multi-subject ICA and dual regression. *NeuroImage* 2009; **47**.

52.   Zuo XN, Kelly C, Adelstein JS, Klein DF, Castellanos FX, Milham MP. Reliable intrinsic

connectivity networks: test-retest evaluation using ICA and dual regression approach.

Neuroimage 2010; **49**(3): 2163-77.

53.   Li B, Zhao N, Tang N, et al. Targeting suicidal ideation in major depressive disorder with

MRI-navigated Stanford accelerated intelligent neuromodulation therapy. *Transl Psychiatry*

2024; **14**(1): 21.

54.   Garza-Villarreal EA, Alcala-Lozano R, Fernandez-Lozano S, et al. Clinical and Functional

Connectivity Outcomes of 5-Hz Repetitive Transcranial Magnetic Stimulation as an Add-on

Treatment in Cocaine Use Disorder: A Double-Blind Randomized Controlled Trial. *Biological*

psychiatry Cognitive neuroscience and neuroimaging 2021; **6**(7): 745-57.

55. R Core Team. R: A language and environment for statistical computing. Vienna, Austria:
Vienna: R Foundation for Statistical Computing; 2013.
56. Champely S, Ekstrom C, Dalgaard P, et al. pwr: Basic functions for power analysis. 2017.
57. Yan C-G, Wang X-D, Zuo X-N, Zang Y-F. DPABI: Data Processing & Analysis for
(Resting-State) Brain Imaging. *Neuroinformatics* 2016; **14**(3): 339-51.
58. Yan C-G, Zang Y-F. DPARSF: A MATLAB Toolbox for "Pipeline" Data Analysis of
Resting-State fMRI. *Frontiers in systems neuroscience* 2010; **4**: 13.
59. Liston C, Chen AC, Zebley BD, et al. Default mode network mechanisms of transcranial
magnetic stimulation in depression. *Biol Psychiatry* 2014; **76**(7): 517-26.
60. Johnson WE, Li C, Rabinovic A. Adjusting batch effects in microarray expression data
using empirical Bayes methods. *Biostatistics* 2007; **8**(1): 118-27.
61. Schaefer A, Kong R, Gordon EM, et al. Local-Global Parcellation of the Human Cerebral
Cortex from Intrinsic Functional Connectivity MRI. *Cerebral cortex (New York, NY : 1991)*
2018; **28**(9): 3095-114.
62. Greicius M. Resting-state functional connectivity in neuropsychiatric disorders. *Current*
opinion in neurology 2008; **21**(4): 424-30.
63. Cullen KR, Gee DG, Klimes-Dougan B, et al. A preliminary study of functional
connectivity in comorbid adolescent depression. *Neuroscience Letters* 2009; **460**(3): 227-31.
64. Strikwerda-Brown C, Davey CG, Whittle S, et al. Mapping the relationship between
subgenual cingulate cortex functional connectivity and depressive symptoms across
adolescence. *Social cognitive and affective neuroscience* 2015; **10**(7): 961-8.
65. Ioannidis JPA. Why Most Published Research Findings Are False. *PLOS Medicine* 2005;
2(8): e124.
66. Chen X, Lu B, Yan CG. Reproducibility of R-fMRI metrics on the impact of different
strategies for multiple comparison correction and sample sizes. *Hum Brain Mapp* 2018; **39**(1):
300-18.
67. Price JL, Drevets WC. Neurocircuitry of mood disorders. *Neuropsychopharmacology*
2010; **35**(1): 192-216.
68. Price JL, Drevets WC. Neural circuits underlying the pathophysiology of mood disorders.
Trends Cogn Sci 2012; **16**(1): 61-71.
69. Hamilton JP, Farmer M, Fogelman P, Gotlib IH. Depressive Rumination, the Default-Mode
Network, and the Dark Matter of Clinical Neuroscience. *Biol Psychiatry* 2015; **78**(4): 224-30.
70. Parker G, Chan B, Hadzi-Pavlovic D. Lower rates of depression in westernised Chinese in
the US. *J Affect Disord* 2007; **104**(1-3): 175-8.
71. Parker G, Gladstone G, Chee KT. Depression in the planet's largest ethnic group: the
Chinese. *Am J Psychiatry* 2001; **158**(6): 857-64.
72. Qiu P, Caine ED, Hou F, Cerulli C, Wittink MN. Depression as seen through the eyes of
rural Chinese women: Implications for help-seeking and the future of mental health care in
China. *J Affect Disord* 2018; **227**: 38-47.
73. Kiyohara C, Yoshimasu K. Association between major depressive disorder and a
functional polymorphism of the 5-hydroxytryptamine (serotonin) transporter gene: a
meta-analysis. *Psychiatr Genet* 2010; **20**(2): 49-58.
74. Tagliazucchi E, Laufs H. Decoding wakefulness levels from typical fMRI resting-state data

reveals reliable drifts between wakefulness and sleep. *Neuron* 2014; **82**(3): 695-708.

75.   Fitzgerald PB, Oxley TJ, Laird AR, Kulkarni J, Egan GF, Daskalakis ZJ. An analysis of

functional neuroimaging studies of dorsolateral prefrontal cortical activity in depression.

Psychiatry Res 2006; **148**(1): 33-45.

76.   Rajkowska G, Goldman-Rakic PS. Cytoarchitectonic definition of prefrontal areas in the

normal human cortex: II. Variability in locations of areas 9 and 46 and relationship to the

Talairach Coordinate System. *Cerebral cortex (New York, NY : 1991)* 1995; **5**(4): 323-37.

77.   Johnson KA, Baig M, Ramsey D, et al. Prefrontal rTMS for treating depression: location

and intensity results from the OPT-TMS multi-site clinical trial. *Brain Stimul* 2013; **6**(2):

108-17.

78.   Perera T, George MS, Grammer G, Janicak PG, Pascual-Leone A, Wirecki TS. The Clinical

TMS Society Consensus Review and Treatment Recommendations for TMS Therapy for Major

Depressive Disorder. *Brain Stimul* 2016; **9**(3): 336-46.

79.   Fitzgerald PB, Hoy K, McQueen S, et al. A randomized trial of rTMS targeted with MRI

based neuro-navigation in treatment-resistant depression. *Neuropsychopharmacology* 2009;

34(5): 1255-62.

80.   Hebel T, Göllnitz A, Schoisswohl S, et al. A direct comparison of neuronavigated and

non-neuronavigated intermittent theta burst stimulation in the treatment of depression.

Brain Stimul 2021; **14**(2): 335-43.

81.   Modak A, Fitzgerald PB. Personalising transcranial magnetic stimulation for depression

using neuroimaging: A systematic review. *The World Journal of Biological Psychiatry* 2021;

22(9): 647-69.

82.   Dubois J, Adolphs R. Building a Science of Individual Differences from fMRI. *Trends in*

Cognitive Sciences 2016; **20**(6): 425-43.

83.   Zuo XN, Xing XX. Test-retest reliabilities of resting-state FMRI measurements in human

brain functional connectomics: a systems neuroscience perspective. *Neurosci Biobehav Rev*

2014; **45**: 100-18.

84.   Zald DH, Rauch S. The orbitofrontal cortex: Oxford University Press, USA; 2006.

85.   Siddiqi SH, Trapp NT, Hacker CD, et al. Repetitive Transcranial Magnetic Stimulation with

Resting-State Network Targeting for Treatment-Resistant Depression in Traumatic Brain Injury:

A Randomized, Controlled, Double-Blinded Pilot Study. *J Neurotrauma* 2019; **36**(8): 1361-74.

86.   Schmaal L, Hibar DP, Samann PG, et al. Cortical abnormalities in adults and adolescents

with major depression based on brain scans from 20 cohorts worldwide in the ENIGMA

Major Depressive Disorder Working Group. *Mol Psychiatry* 2016.

87.   Schmaal L, Veltman DJ, van Erp TG, et al. Subcortical brain alterations in major

depressive disorder: findings from the ENIGMA Major Depressive Disorder working group.

Mol Psychiatry 2016; **21**(6): 806-12.

88.   Esteban O, Markiewicz CJ, Blair RW, et al. fMRIPrep: a robust preprocessing pipeline for

functional MRI. *Nat Methods* 2019; **16**(1): 111-6.

89.   Lynch CJ, Elbau IG, Ng TH, et al. Automated optimization of TMS coil placement for

personalized functional network engagement. *Neuron* 2022; **110**(20): 3263-77.e4.

90.   Downar J, Daskalakis ZJ. New targets for rTMS in depression: a review of convergent

evidence. *Brain Stimul* 2013; **6**(3): 231-40.

91.   Siddiqi SH, Khosravani S, Rolston JD, Fox MD. The future of brain circuit-targeted
therapeutics. *Neuropsychopharmacology* 2023.
92.   Dunlop K, Sheen J, Schulze L, et al. Dorsomedial prefrontal cortex repetitive transcranial
magnetic stimulation for treatment-refractory major depressive disorder: A three-arm,
blinded, randomized controlled trial. *Brain Stimul* 2020; **13**(2): 337-40.
93.   Zuo X-N, Xing X-X. Test-retest reliabilies of resting-state FMRI measurements in human
brain functional connectomics: A systems neuroscience perspective. *Neuroscience &*
Biobehavioral Reviews 2014; **45**: 100-18.
94.   Wang D, Buckner RL, Fox MD, et al. Parcellating cortical functional networks in
individuals. *Nat Neurosci* 2015; **18**(12): 1853-60.
95.   Marek S, Tervo-Clemmens B, Calabro FJ, et al. Reproducible brain-wide association
studies require thousands of individuals. *Nature* 2022; **603**(7902): 654-60.
96.   Lu B, Chen X, Xavier Castellanos F, et al. The power of many brains: Catalyzing
neuropsychiatric discovery through open neuroimaging data and large-scale collaboration.
Sci Bull (Beijing) 2024; **69**(10): 1536-55.
97.   Chen X, Lu B, Li HX, et al. The DIRECT consortium and the REST-meta-MDD project:
towards neuroimaging biomarkers of major depressive disorder. *Psychoradiology* 2022; **2**(1):
32-42.
98.   Wang YW, Chen X, Yan CG. Comprehensive evaluation of harmonization on functional
brain imaging for multisite data-fusion. *Neuroimage* 2023: 120089.
1032

1033   Table 1. Demographic and clinical data for all samples included in the DIRECT II project

Site	Age (HC)[1]	Age (MDD)[1]	% Male (HC)[2]	% Female (HC)[2]	% Male (MDD)[2]	% Female (MDD)[2]	Education (HC, years)[1]	Education (MDD, years)[1]	Age at the onset of the first episode (years)[1]	Episode number[1]	Full episode durations (months)[1]	HAMD-17 score[1]	N (HC)	N (MDD)
1	39.33 (15.80)	38.67 (13.50)	87 (35.22%)	160 (64.78%)	98 (35.13%)	181 (64.87%)	13.05 (3.89)	12.03 (3.62)	35.75 (13.47)	1.19 (0.40)	49.01 (64.25)	20.96 (5.60)	247	279
2	20.90 (2.89)	19.02 (3.79)	24 (38.71%)	38 (61.29%)	37 (27.41%)	98 (72.59%)	13.73 (1.80)	11.97 (2.55)	16.28 (4.05)	1.07 (0.35)	16.94 (19.74)	18.06 (8.99)	62	135
3	26.50 (11.22)	34.40 (10.56)	5 (25.00%)	15 (75.00%)	5 (25.00%)	15 (75.00%)	15.40 (1.47)	14.90 (2.97)	35.55 (11.96)	1.15 (0.49)	14.50 (29.14)	29.37 (6.41)	20	20
4	29.89 (9.88)	30.87 (10.39)	9 (33.33%)	18 (66.67%)	13 (41.94%)	18 (58.06%)	13.04 (3.80)	12.65 (4.22)	26.80 (10.89)	2.52 (3.66)	42.03 (58.57)	21.79 (5.74)	27	31
5	36.79 (9.07)	33.59 (9.85)	32 (56.14%)	25 (43.86%)	60 (46.88%)	68 (53.13%)	14.37 (2.64)	13.07 (3.43)	NA	1.00 (0.00)	NA	26.59 (3.76)	57	128
6	23.75 (5.15)	30.90 (9.29)	45 (44.55%)	56 (55.45%)	31 (52.54%)	28 (47.46%)	13.88 (2.31)	12.03 (3.06)	28.14 (9.46)	1.87 (1.12)	44.14 (61.28)	20.51 (6.71)	101	59
7	19.80 (3.85)	20.00 (6.51)	8 (26.67%)	22 (73.33%)	5 (23.81%)	16 (76.19%)	11.90 (2.55)	10.81 (2.86)	18.81 (7.19)	1.10 (0.30)	15.60 (16.23)	24.10 (12.68)	30	21
8	31.93 (9.66)	32.84 (8.95)	18 (42.86%)	24 (57.14%)	15 (34.88%)	28 (65.12%)	15.40 (1.86)	15.79 (1.82)	30.60 (8.78)	3.65 (2.29)	21.51 (11.65)	23.84 (3.88)	42	43
9	36.12 (13.85)	45.11 (14.32)	10 (40.00%)	15 (60.00%)	19 (40.43%)	28 (59.57%)	13.56 (4.01)	10.83 (4.52)	42.19 (14.23)	1.43 (0.77)	36.07 (53.22)	NA	25	47

10	36.49 (12.64)	34.00 (11.81)	30 (44.78%)	37 (55.22%)	36 (45.00%)	44 (55.00%)	14.58 (2.30)	13.25 (3.08)	26.15 (10.47)	2.30 (1.78)	92.00 (96.70)	14.58 (8.21)	67	80
11	36.62 (12.11)	38.29 (12.75)	30 (44.12%)	38 (55.88%)	55 (48.67%)	58 (51.33%)	12.96 (3.57)	10.96 (3.48)	32.21 (12.46)	3.36 (3.63)	75.51 (93.37)	23.54 (9.23)	68	113
12	29.59 (5.00)	30.47 (7.20)	15 (46.88%)	17 (53.13%)	21 (33.87%)	41 (66.13%)	14.59 (2.82)	13.73 (3.39)	29.63 (7.11)	1.00 (0.00)	6.10 (4.22)	21.27 (3.47)	32	62
13	35.45 (13.76)	38.00 (14.56)	9 (31.03%)	20 (68.97%)	13 (40.63%)	19 (59.38%)	11.28 (4.31)	12.03 (3.87)	33.09 (13.93)	1.45 (1.12)	NA	18.66 (9.66)	29	32
14	38.10 (11.96)	43.50 (11.28)	65 (45.14%)	79 (54.86%)	47 (34.56%)	89 (65.44%)	13.24 (4.20)	9.07 (3.60)	37.52 (11.50)	2.55 (2.15)	67.16 (77.29)	30.04 (10.52)	144	136
15	21.51 (4.58)	22.06 (4.70)	23 (65.71%)	12 (34.29%)	7 (20.59%)	27 (79.41%)	12.57 (1.42)	11.85 (2.39)	21.82 (5.19)	2.50 (2.81)	12.76 (15.11)	NA	35	34
16	25.73 (7.38)	26.07 (7.66)	44 (54.32%)	37 (45.68%)	37 (44.58%)	46 (55.42%)	15.53 (2.81)	13.81 (2.60)	23.02 (8.11)	1.73 (1.11)	30.18 (44.24)	25.42 (3.78)	81	83
17	34.73 (9.66)	35.21 (9.43)	19 (38.78%)	30 (61.22%)	15 (35.71%)	27 (64.29%)	14.31 (4.21)	12.50 (4.09)	35.00 (9.53)	1.00 (0.00)	3.79 (2.62)	23.67 (5.25)	49	42
18	32.19 (11.83)	35.25 (13.15)	13 (40.63%)	19 (59.38%)	12 (50.00%)	12 (50.00%)	12.53 (3.03)	13.42 (4.47)	NA	1.58 (0.72)	17.77 (19.57)	22.38 (4.23)	32	24
19	30.94 (7.32)	27.03 (11.04)	16 (50.00%)	16 (50.00%)	5 (15.63%)	27 (84.38%)	13.94 (2.84)	12.69 (3.18)	25.58 (10.86)	1.28 (0.46)	12.88 (14.38)	23.06 (5.66)	32	32
20	31.40 (10.99)	27.80 (8.99)	5 (25.00%)	15 (75.00%)	4 (20.00%)	16 (80.00%)	16.10 (3.21)	13.60 (4.36)	26.65 (9.21)	1.30 (0.47)	14.55 (15.14)	24.85 (6.47)	20	20
21	32.80 (11.06)	32.52 (11.44)	31 (55.36%)	25 (44.64%)	17 (28.33%)	43 (71.67%)	15.88 (4.59)	12.20 (4.12)	31.44 (11.70)	NaN (NaN)	35.79 (44.90)	22.71 (2.62)	56	60
22	34.67 (13.54)	35.74 (10.02)	8 (33.33%)	16 (66.67%)	10 (28.57%)	25 (71.43%)	14.21 (2.86)	13.20 (3.75)	31.23 (7.91)	1.06 (1.35)	5.59 (14.93)	22.29 (6.04)	24	35

23	13.57 (2.18)	13.91 (1.48)	14 (50.00%)	14 (50.00%)	8 (13.79%)	50 (86.21%)	7.57 (2.18)	7.91 (1.48)	NA	NA	NA	NA	28	58

1034   1 Mean (SD)

1035   2 N (%)

1036   NA: data were missing for this site.