Functional Connectivity Abnormalities of the Subgenual Anterior Cingulate Cortex: Implications for Transcranial Magnetic Stimulation in Depression

Xiao Chen[1,2,3,4,5]#, Bin Lu[1,3,4,5]#, Yu-Wei Wang[1,3,4,5], Xue-Ying Li[1,3,4,5], Zi-Han Wang[1,3,4,5], Hui-Xian Li[6], Yi-Fan Liao[1,3,4,5], Daniel M. Blumberger[2,7], Francisco Xavier Castellanos[8,9], Li-Ping Cao[10], Guan-Mao Chen[11], Jian-Shan Chen[10], Tao Chen[12], Tao-Lin Chen[13,14], Yan-Rong Chen[1,3,4,5], Yu-Qi Cheng[15], Zhao-Song Chu[15], Shi-Xian Cui[1,16,17], Xi-Long Cui[18], Zhao-Yu Deng[1,3,4,5], Qing-Lin Gao[1,3,4,5], Qi-Yong Gong[13,14], Wen-Bin Guo[18], Can-Can He[19], Zheng-Jia-Yi Hu[1,16,17], Qian Huang[20], Xin-Lei Ji[18], Feng-Nan Jia[21], Li Kuang[20], Bao-Juan Li[22], Feng Li[23], Tao Li[24,25], Xue Li[1,3,4,5], Tao Lian[1,3,4,5], Xiao-Yun Liu[26], Yan-Song Liu[21], Zhe-Ning Liu[18], Yi-Cheng Long[18], Jian-Ping Lu[27], Jiang Qiu[28], Xiao-Xiao Shan[18], Tian-Mei Si[29], Peng-Feng Sun[30], Chuan-Yue Wang[23], Han-Lin Wang[1,3,4,5], Hua-Ning Wang[22], Xiang Wang[18], Ying Wang[11], Chen-Nan Wu[1,3,4,5], Xiao-Ping Wu[30], Xin-Ran Wu[28], Yan-Kun Wu[29], Chun-Ming Xie[19], Guang-Rong Xie[18], Peng Xie[31,32,33], Xiu-Feng Xu[15], Zhen-Peng Xue[27], Hong Yang[12], Jian Yang[32], Hua Yu[24,25], Yong-Qiang Yu[34,35,36], Min-Lan Yuan[37], Yong-Gui Yuan[26], Yu-Feng Zang[38,39], Ai-Xia Zhang[40], Ke-Rang Zhang[40], Wei Zhang[37], Zi-Jing Zhang[1,3,4,5], Jing-Ping Zhao[18], Jia-Jia Zhu[34,35,36], Xi-Nian Zuo[41,42], the DIRECT Consortium, Chao-Gan Yan[1,3,4,5,16,17]*

[1]CAS Key Laboratory of Behavioral Science, Institute of Psychology, Beijing 100101, China

[2]Temerty Centre for Therapeutic Brain Intervention, Campbell Family Research Institute, Centre for Addiction and Mental Health, Toronto, Ontario M6J1H4, Canada

[3]Magnetic Resonance Imaging Research Center, Institute of Psychology, Chinese Academy of Sciences, Beijing 100101, China

[4]Department of Psychology, University of Chinese Academy of Sciences, Beijing 100049, China

[5]International Big-Data Center for Depression Research, Chinese Academy of Sciences, Beijing 100101, China

[6]The Third Affiliated Hospital of Zhengzhou University, Zhengzhou, Henan 450052, China

[7]Department of Psychiatry, Temerty Faculty of Medicine, University of Toronto, Toronto, Ontario M5S1A1, Canada

[8]Department of Child and Adolescent Psychiatry, NYU Grossman School of Medicine, New York, NY

10016, USA

[9]Nathan Kline Institute for Psychiatric Research, Orangeburg, NY 10962, USA

[10]Affiliated Brain Hospital of Guangzhou Medical University, Guangzhou, Guangdong 510370, China

[11]The First Affiliated Hospital of Jinan University, Guangzhou, Guangdong 250024, China

[12]Department of Radiology, The First Affiliated Hospital, College of Medicine, Zhejiang University,

Hangzhou, Zhejiang 310058, China

[13]Huaxi MR Research Center (HMRRC), Department of Radiology, West China Hospital of Sichuan

University, Chengdu, Sichuan 610044, China

[14]Research Unit of Psychoradiology, Chinese Academy of Medical Sciences, Chengdu, Sichuan 610052,

China

[15]Department of Psychiatry, First Affiliated Hospital of Kunming Medical University, Kunming, Yunnan

650032, China

[16]Sino-Danish College, University of Chinese Academy of Sciences, Beijing 101408, China

[17]Sino-Danish Center for Education and Research, Graduate University of Chinese Academy of Sciences,

Beijing 101408, China

[18]Department of Psychiatry, and National Clinical Research Center for Mental Disorders, The Second

Xiangya Hospital of Central South University, Changsha, Hunan 410011, China

[19]Department of Neurology, Affiliated ZhongDa Hospital of Southeast University, Nanjing, Jiangsu

210009, China

[20]Department of Psychiatry, The First Affiliated Hospital of Chongqing Medical University, Chongqing

400042, China

[21]Department of Clinical Psychology, Suzhou Psychiatric Hospital, The Affiliated Guangji Hospital of

Soochow University, Suzhou, Jiangsu 215003, China

[22]Xijing Hospital of Air Force Military Medical University, Xi'an, Shaanxi 710032, China

[23]Beijing Anding Hospital, Capital Medical University, Beijing 100120, China

[24]Affiliated Mental Health Center & Hangzhou Seventh People's Hospital, Zhejiang University School of

Medicine, Hangzhou, Zhejiang 310063, China

[25]Mental Health Center and Psychiatric Laboratory, West China Hospital of Sichuan University,

Chengdu, Sichuan 610044, China

[26]Department of Psychosomatics and Psychiatry, Zhongda Hospital, School of Medicine, Southeast University, Nanjing, Jiangsu 210009, China

[27]Shenzhen Kangning Hospital, Shenzhen, Guangzhou 518020, China

[28]Faculty of Psychology, Southwest University, Chongqing 400715, China

[29]National Clinical Research Center for Mental Disorders (Peking University Sixth Hospital) & Key Laboratory of Mental Health, Ministry of Health (Peking University), Beijing 100191, China

[30]Xi'an Central Hospital, Xi'an, Shaanxi 710004, China

[31]Institute of Neuroscience, Chongqing Medical University, Chongqing 400016, China

[32]Chongqing Key Laboratory of Neurobiology, Chongqing 400000, China

[33]Department of Neurology, The First Affiliated Hospital of Chongqing Medical University, Chongqing 400042, China

[34]Department of Radiology, The First Affiliated Hospital of Anhui Medical University, Hefei, Anhui 230022, China

[35]Research Center of Clinical Medical Imaging, Hefei, Anhui 230032, China

[36]Anhui Provincial Institute of Translational Medicine, Hefei, Anhui 230032, China

[37]West China Hospital of Sichuan University, Chengdu, Sichuan 610044, China

[38]Center for Cognition and Brain Disorders, The Affiliated Hospital of Hangzhou Normal University, Hangzhou, Zhejiang 310018, China

[39]Zhejiang Key Laboratory for Research in Assessment of Cognitive Impairments, Hangzhou, Zhejiang 310000, China

[40]First Hospital of Shanxi Medical University, Taiyuan, Shanxi 030001, China

[41]Developmental Population Neuroscience Research Center, IDG/McGovern Institute for Brain Research, Beijing Normal University, Beijing 100091, China

[42]National Basic Science Data Center, Beijing 100038, China

[43]Department of Psychiatry, Sir Run Run Shaw Hospital, Zhejiang University School of Medicine, Hangzhou, Zhejiang 310020, China

[44]Shanghai Mental Health Center, Shanghai Jiao Tong University School of Medicine, Shanghai 200030, China

[45]Department of Diagnostics, Affiliated Hospital, Hangzhou Normal University Medical School,

Hangzhou, Zhejiang 311121, China

[46]Department of Psychiatry, First Affiliated Hospital, China Medical University, Shenyang, Liaoning

110122, China

[47]Early Intervention Unit, Department of Psychiatry, Affiliated Nanjing Brain Hospital, Nanjing Medical

University, Nanjing, Jiangsu 210024, China

[48]Department of Neurology, The First Affiliated Hospital of Anhui Medical University, Hefei, Anhui

230022, China

[49]Department of Psychiatry, Shanghai Jiao Tong University School of Medicine, Shanghai 200025, China

*Corresponding author:

Chao-Gan Yan, Ph. D.

CAS Key Laboratory of Behavioral Science, Institute of Psychology, Beijing, China

16 Lincui Road, Chaoyang District, Beijing 100101, China

Tel: +86-10-64101582

E-mail: yancg@psych.ac.cn

#Xiao Chen and Bin Lu contributed equally to this work.

DIRECT consortium: Chao-Gan Yan, Ph.D.[1,3,4,5,16,17], Xiao Chen, Ph.D.[1,2,3,4,5], Li-Ping Cao, M.D.[10], Wei

Chen, M.D.[43], Yu-Qi Cheng, M.D.[15], Yi-Ru Fang, M.D.[44], Qi-Yong Gong, M.D.[13,14], Wen-Bin Guo, M.D.[18],

Li Kuang, M.D.[20], Bao-Juan Li, M.D.[22], Tao Li, M.D.[24,25], Yan-Song Liu, M.D.[21], Zhe-Ning Liu, M.D.[18],

Jian-Ping Lu, M.D.[27], Qing-Hua Luo, M.D.[20], Hua-Qing Meng, M.D.[20], Dai-Hui Peng, M.D.[44], Jiang Qiu,

Ph.D.[28], Yue-Di Shen, M.D.[45], Tian-Mei Si, M.D.[29], Yan-Qing Tang, M.D.[46], Chuan-Yue Wang, M.D.[23], Fei

Wang, M.D.[46,47], Hua-Ning Wang, M.D.[22], Kai Wang, M.D.[48], Xiang Wang, M.D.[18], Ying Wang, M.D.[11],

Xiao-Ping Wu, M.D.[30], Chun-Ming Xie, M.D.[19], Guang-Rong Xie, M.D.[18], Peng Xie, M.D.[31,32,33], Xiu-Feng

Xu, M.D.[15], Hong Yang, M.D.[12], Jian Yang, M.D.[32], Shu-Qiao Yao, M.D.[18], Yong-Qiang Yu, M.D.[34,35,36],

Yong-Gui Yuan, M.D.[26], Ke-Rang Zhang, M.D.[40], Wei Zhang, M.D.[37], Zhi-Jun Zhang, M.D.[19], Jun-Juan Zhu,

M.D.[49], Xi-Nian Zuo, Ph.D.[41,42], Jing-Ping Zhao, M.D.[18], Yu-Feng Zang, M.D.[38,39]

Abstract

Background: The subgenual anterior cingulate cortex (sgACC) plays a central role in the

pathophysiology of major depressive disorder (MDD), and its functional interactive profile

with the left dorsal lateral prefrontal cortex (DLPFC) is associated with transcranial magnetic

stimulation (TMS) treatment outcomes. Nevertheless, previous research on sgACC functional

connectivity (FC) in MDD has yielded inconsistent results, partly due to small sample sizes

and limited statistical power. Furthermore, calculating sgACC-FC to target TMS on an

individual level is challenging because of the low signal-to-noise ratio and the poor

replicability of individualized functional brain images.

Methods: Leveraging a large multi-site cross-sectional sample (1660 MDD patients vs. 1341

healthy controls) from Phase II of the Depression Imaging REsearch ConsorTium (DIRECT), we

systematically delineated case-control difference maps of sgACC-FC. Then, in a sample of 25

individuals with treatment-resistant depression who had received repetitive TMS (rTMS)

treatment, we examined the relationship between case-control differences in FCs between

sgACC and their specific TMS targets and treatment outcomes. Next, we tested whether the

position of the group mean FC-based target (previously determined in healthy participants)

differed in MDD patients. Finally, we developed a dual regression (DR) based approach to

integrate group-level statistical maps with individual-level spontaneous brain activity to

evaluate individualized TMS target localization in MDD. We tested this approach in a sample

of 16 individuals who had received rTMS.

Results: We found enhanced sgACC-DLPFC FC in MDD patients. The magnitude of

case-control differences in FC between sgACC and TMS targets was related to clinical

improvement. We found different peak sgACC anticorrelation locations in mean FC maps of

MDD patients and HCs. More effective TMS targets were closer to individualized DR-based

loci than to group-level targets.

Conclusion: In summary, we reliably delineated MDD-related abnormalities of sgACC-FC

profiles in a large independently ascertained sample and demonstrated the potential impact

of such case-control differences on FC-guided localization of TMS targets. The proposed

individualized approach for TMS targeting has the potential to improve TMS treatment

outcome and warrants prospective clinical trials.

Keywords: major depressive disorder, transcranial magnetic stimulation, individualization,

subgenual anterior cingulate cortex, functional connectivity, dual regression

Introduction

Major depressive disorder (MDD) is a common and debilitating psychiatric disorder projected to be the most burdensome condition worldwide by 2030 [1]. Despite extensive research, the pathophysiology of MDD remains elusive. Nevertheless, a key putative brain region appears to be the subgenual anterior cingulate cortex (sgACC), which shows reproducible metabolic hyperactivity [2], is targeted by effective deep brain stimulation [3,4], and has been implicated in emotional responses, motivation, and rumination [5]. Accordingly, identifying an optimized neuromodulation target in the left dorsal lateral prefrontal cortex (DLPFC) based on sgACC-related functional connectivity (FC) is crucial for developing effective depression treatments [6,7]. In light of inconsistent findings derived from studies with small sample sizes [8-14], we set out to establish a large sample to identify a reliable abnormal sgACC-DLPFC FC profile in MDD and further integrate this profile with individual brain activity to generate individualized neuromodulation targets for treating depression.

Numerous investigations have delved into FC abnormalities in MDD using resting-state functional magnetic resonance imaging (R-fMRI). Abnormal FCs between sgACC and amygdala, thalamus, temporal gyrus, lingual gyrus, cerebellum, DLPFC, and default mode network (DMN) regions such as medial and dorsal medial prefrontal cortex, precuneus, and parahippocampus have been reported [8,9,12-19]. However, findings have been inconsistent, making integrating findings and generating precise profiles of sgACC-related FC abnormalities challenging. This deficiency in reproducibility could be partially due to small sample sizes and low statistical power of clinical imaging studies [20,21]. To address the issue of limited sample size, we initiated the Depression Imaging REsearch ConsorTium (DIRECT) [22] and conducted an initial meta/mega-analysis (N_{MDD} = 1300), referred to as REST-meta-MDD [23]. DIRECT Phase I shared ROI-level signals, thus enabling the investigation of multiple MDD-related abnormalities in network FC, FC topological and dynamic features, and functional lateralization [23-30]. In DIRECT Phase II data reporting, we pooled an expanded MDD sample (N_{MDD} = 1660), which was preprocessed with a surface-based pipeline, DPABISurf [31]. DIRECT

Phase II shared voxel/vertex level BOLD time series, allowing more flexible and thorough

investigations. Leveraging the most comprehensive MDD R-fMRI dataset to date

encompassing depression patients and healthy controls, we can determine an aberrant

sgACC-FC profile associated with depression, characterized by superior reproducibility and

low risk of false positives.

Maps of sgACC-related FC abnormalities are clinically useful for predicting repetitive

transcranial magnetic stimulation (rTMS) treatment outcomes in MDD patients [32-34].

Specifically, the anticorrelation between sgACC and left DLPFC has been associated with

clinical improvement from rTMS treatment [35-39]. This has led to the intriguing notion that the

FC between sgACC and left DLPFC could be leveraged to identify more precise rTMS targets

and improve the efficacy of rTMS delivered to the left DLPFC [6,40]. Researchers have identified

a group-wise TMS target [37], which was the most anticorrelated DLPFC site to sgACC in the

mean FC map from a large cohort of healthy adults. Subtle but significant case-control

differences in resting-state FC profiles have been identified in a large sample of MDD

patients[23]. Thus, understanding the profiles of sgACC FC case-control differences and their

impact on potential targets for rTMS applied to the left DLPFC could be a critical step toward

developing optimized rTMS target site identification methods.

Individual human brains exhibit highly heterogeneous functional organization [41], with the

DLPFC regions exhibiting the highest level of interindividual variation in cytoarchitecture,

brain function, and network connectivity profiles [7,42]. While several individualized FC-guided

TMS target identification algorithms have been proposed [35,40,43-47], the target localizations of

most existing TMS protocols have not been individualized. The major obstacles to identifying

individualized TMS locations are the low signal-to-noise ratio in the sgACC area and the poor

reproducibility of individual FC maps [48,49]. The high reliability and statistical power of the

DIRECT MDD cohort (N_{MDD} = 1660) allow the integration of group-level statistical maps and

individual functional brain images to achieve precise and reliable TMS localization. Here, we

propose an algorithm based on dual regression (DR), which was initially developed for

mapping group-level independent component analysis (ICA) results onto individual brains [50].

Utilizing the DR-based algorithm avoids calculating FC based on the individual level time

series of a noisy area such as the sgACC. The computation of this algorithm was based on DR

of fMRI signals from the DLPFC region, which has a higher signal-to-noise ratio compared to

the sgACC.   Thus, this algorithm enhances the efficacy and reliability of individualization

approaches for identifying TMS targets [51].

In the present study, we leverage a large-scale multi-center sample (DIRECT Phase II, 1660

MDD patients and 1341 healthy controls (HCs)) to derive a reliable sgACC-related FC

abnormality profile for MDD. We aimed to assess the role of the sgACC-related FC in the

precise localization of TMS targets. To facilitate individualized localization, we aimed to

develop a DR-based targeting algorithm that significantly enhanced the correlation between

target position and the reduction of clinical symptoms. We hypothesized that MDD patients

will show a significantly abnormal sgACC-FC profile, especially in the left DLPFC. We also

hypothesized that our newly-developed DR-based algorithm will outperform traditional TMS

group targets. To our knowledge, this is the first study to utilize a dual regression algorithm

to integrate large-scale group-level statistical maps with individual-level spontaneous brain

activity to achieve individualized TMS targeting in MDD.

Materials and methods

Study sample

This study utilized three independent datasets. The first dataset ("DIRECT") is a large-scale,

multi-sited consortium sharing standardized preprocessed R-fMRI time series. Building on

the initial success of DIRECT Phase I, consortium members and international collaborators

met on May 11[th]-12[th], 2019, and agreed to launch DIRECT Phase II, which comprises 23

case-control designed datasets, including R-fMRI and T1 structural scans from 1660 MDD

patients and 1341 HCs. Researchers from each site later took a 2-day DPABISurf training

course on September 14th-15th, 2019, to harmonize the organization and preprocessing of

R-fMRI/T1 structural data. Demographic and clinical characteristics for each sample are

presented in Figure 1 and Table 1. Site information, sample size, and previous publications

based on the shared data are listed in Table S1. All participants were asked to self-report

their sex (biological attribute) as part of the case report form (CRF). All participants in DIRECT

Phase II were East Asian. Patients were diagnosed with MDD based on ICD 10 or DSM-IV.

Healthy controls who were matched with MDD patients by age, sex ratio, and educational

levels were recruited at each site. All participants provided written informed consent, and

local institutional review boards approved each study from all included cohorts. The analysis

plan of the current study has been reviewed and approved by the Institutional Review Board

of Institute of Psychology, Chinese Academy of Sciences (No. H21102). Data will be made

available to the public as outlined in the Data Sharing Statement.

The second dataset ("TRD-TMS") comprises 25 medication-treatment-resistant MDD (TRD)

patients who underwent 4 to 7 weeks of daily repetitive TMS applied over the left DLPFC.

Patients' TMS sites were recorded using their structural MRI images and a frameless

neuronavigation system. Treatment response was assessed with the 24-item Hamilton

Depression Rating Scale (HAMD). Only TMS outcomes and target coordinates were openly

shared for this dataset. We get access to these data from the supplementary materials of

Weigand et al. [37] For more details on this dataset, please refer to Weigand et al. [37]

The third dataset ("CUD-TMS") comprises 27 cocaine use disorder (CUD) patients who

underwent two daily sessions of rTMS treatment over the left DLPFC in an acute phase and

two weekly sessions of rTMS treatment in a maintenance phase. The TMS outcomes,

neuroimaging data, participants' demographic info, and target coordinates for this dataset

were openly shared. Depressive symptoms were a secondary treatment outcome in the

original study and were used to validate the localization algorithm's clinical relevance in the

current study. In addition, this dataset shared participants' baseline R-fMRI data

(https://openneuro.org/datasets/ds003037/versions/1.0.0), which allowed for the

calculation of individualized optimal TMS targets. For more details on this dataset, see

Garza-Villarreal et al. [52]

Figure 1. Sample characteristics of the DIRECT dataset. (A) Sample sizes of each site; (B) Number of

male/female subjects irrespective of diagnosis; (C) Violin plots depicting the age distribution (in years).

Solid black lines indicate the mean, 25th, and 75th percentiles; (D) Violin plots show education

distribution (in years). Solid black lines indicate the mean, 25th, and 75th percentiles; (E) Violin plots

depicting the distribution of scores of the Hamilton Depression Rating Scale (HAMD). Solid black lines

indicate the mean, 25th, and 75th percentiles.

Approach

The study's first objective was to delineate case-control differences in sgACC-FC profile and

explore its implication in identifying FC-guided individualized TMS targets. Accordingly, we

conducted a generalized linear model (GLM) to compare voxel-wise sgACC-FC maps of MDD

patients and HCs in the DIRECT dataset. We then demonstrated the association between

clinical improvement and group differences in TMS targeting sgACC-FCs by leveraging the

TRD-TMS dataset. Given that the peak sgACC anticorrelation of a normative connectome

within the left DLPFC was usually selected as the FC-guided sgACC group target, we showed

the impact of case-control differences on such group targets by separately identifying the

peak sgACC anticorrelation in the mean sgACC-FC maps of the MDD group and HC group

from the DIRECT dataset. Finally, we identified individualized optimal targets using a

DR-based algorithm guided by statistical maps (e.g., group difference map, mean sgACC-FC

maps). We validated the clinical effectiveness of the individualized approach by computing

the correlation between clinical outcome and the distance between the actual TMS sites and

the identified individualized targets in the CUD-TMS dataset. All statistical tests conducted in

the current study were two-sided.

Power calculations for primary hypothses

The primary outcome of the current study is the case-control differences regarding the

sgACC-FC profiles. Estimates of the effect size (Cohen's d = 0.186) of MDD patients'

abnormalities in FCs draw directly from our prior research based on the DIRECT Phase I

dataset [23]. Power calculation was performed using R version 4.3.1 [53] with pwr [54]. A sample of

455 patients will achieve 80% power with a 5% Type I error rate.

Image preprocessing

Acquisition parameters and scanners for all cohorts are provided in Table S2. All R-fMRI and

structural MRI scans were preprocessed at each site using the same DPABISurf protocol, an

R-fMRI data analysis toolbox evolved from DPABI/DPARSF [31,55,56] (For details, see SI). Given

the controversy regarding global signal regression (GSR) and its essential role in identifying

TMS targets [40], we performed preprocessing pipelines with and without GSR.

FC maps of sgACC

Although recent studies have attempted to identify personalized TMS targets using surface-based algorithms [43], most previous studies have reported targets in volume-based MNI space with sgACC ROIs defined as a sphere in volume-based space [35,57]. To better compare our results with the existing literature, we used the volume-based preprocessed imaging data from DPABISurf.

We defined the sgACC as a 10 mm diameter sphere located on the average MNI coordinates based on prior studies showing reduced glucose metabolism or blood flow after receiving an antidepressant treatment (MNI coordinates: x = 6, y = 16, z = -10. For details, please refer to Fox et al., 2012 [35]). The sgACC time series were determined for each individual by spatially averaging the preprocessed R-fMRI time series across all voxels in the abovementioned masks. We then calculated whole-brain FC maps in volume-based MNI space. FC was calculated using Pearson's correlation and underwent Fisher's r-to-z transformation. All FC maps were smoothed with a 6 mm full-width half maximum (FWHM) kernel size. We used ComBat [58] to control potential site and scanner biases (For details, see SI).

Group difference maps of sgACC-FC profiles

We used a voxel-wise GLM to examine differences in the FC maps of sgACC between MDD patients and HCs in DIRECT Phase II. Cohen's f^2 was calculated to characterize the effect sizes of this group difference effect. The GLM model includes age, sex, education, and head motion as covariates:

$$y = \mu + X\beta + \epsilon \tag{1}$$

where y denotes the FC value of a given voxel from a given participant; μ stands for the constant term; X represents the design matrix for the covariates of interest (diagnosis, age, sex, education, and head motion); β is a vector of regression coefficients corresponding to X; and ϵ is a vector of residuals that follow $N(0, \sigma^2)$. Multiple comparison correction was conducted using false discovery rate (FDR) correction at $q < 0.05$.

Relationship between group differences in TMS targets' sgACC FCs and clinical outcomes

To explore the relationship between group differences in TMS targets' sgACC FCs and clinical

outcomes, we first extracted the mean t-values from the group difference map of 4 mm

radius spheres centered at each targeting coordinate in the TRD-TMS dataset, then examined

the Pearson correlations between these t-values and HAMD score reductions. We

anticipated that greater group differences in sgACC-FC at the target location (i.e., higher t

values), would be related to better TMS therapeutic effect (higher HAMD reductions).

Group targets based on mean sgACC-FC maps

The prior group-level DLPFC TMS target had been derived from a cohort of healthy young

adults [37]; here, we averaged whole-brain sgACC-FC maps across all the DIRECT participants in

the MDD and HC groups separately. We then searched for the peak sgACC anticorrelated

voxel within the DLPFC area (i.e., Brodmann area (BA) 46) as the mean sgACC-FC guided TMS

targets for the MDD and HC groups.

Identification of individualized TMS targets by dual regression

Given the greater reliability of statistical maps from a large cohort of subjects (N_{MDD} = 1660 in

the DIRECT dataset), we used DR to identify individualized TMS targets guided by group-level

statistical maps (Figure 2).

We used a two-step algorithm to back-reconstruct the corresponding subject-specific spatial

patterns based on group-level components [50] (see Figure S4 for details). First, an

individual-level time course measuring how the group-level template fitted into the brain

BOLD signal spatial pattern at every time point was calculated using least-squares for each

participant. We used three group-level sgACC FC statistical maps (i.e., the group difference

map, the mean MDD sgACC-FC map, and the mean HC sgACC-FC map) as group templates in

the DR-based algorithm. As the two clusters that survived FDR correction (q < 0.05) were too

small to be used as group templates in the DR computation, we expanded them by applying

a more liberal multiple comparison correction threshold (FDR $q < 0.1$). In addition, we

reversed the mean MDD/HC sgACC-FC maps to generate the most sgACC-anticorrelated brain

area in DLPFC before conducting regressions. Second, the time course derived in the first

step was used as a temporal regressor to estimate subject-specific spatial maps. The

subject-specific maps were considered the best approximation of the group-level template

for single participants. This benefits from the group-level prior information based on a large

sample and the subject-specific spontaneous brain activity. The center of mass of the largest

cluster in the subject-specific maps was calculated to determine the final TMS targeting

coordinates. These clusters were extracted from the thresholded subject-specific maps in

which only the top-n voxels in the DLPFC mask were retained. The thresholds were related to

the type and size of the group template input into the DR algorithm and incrementally varied

between 5 and 100 voxels. The threshold for the DR approach guided by the FDR-corrected

group difference map was set to 10, and the threshold for the DR approach guided by mean

sgACC-FC maps was set to 50 voxels (Figure S5). We chose BA 46 as the DLPFC mask in the DR

approach because prior studies have reported that BA 46 is more anticorrelated to sgACC

than alternative DLPFC definitions[35] (e.g., BA 9, Beam F3, or the 5 cm method) and suggested

that a more anterior and lateral location was related to better treatment response in MDD

TMS treatment [7,37].

Figure 2. Schematic diagram of the statistical map guided individualized rTMS targeting algorithms. (A)

The sgACC-FC group difference map guided individualized target localization. (B) The mean MDD

sgACC-FC map guided individualized precise target localization. (C) The mean HC sgACC-FC map guided

individualized precise target localization.

Clinical effectiveness of the group-level and individualized TMS targets

We leveraged the CUD-TMS dataset to evaluate the clinical significance of individualized TMS

targets. In the CUD-TMS dataset, rTMS was delivered at the left DLPFC using either the 5.5

cm anatomic criterion or the Beam F3 method. We selected participants with a baseline

HAMD score greater than 7, yielding a subsample of 16 individuals, and calculated their

maximum HAMD reductions during the entire treatment course as an index of TMS

therapeutic efficacy. We identified the proposed DR-based individualized TMS targets guided

by the group difference map or the mean sgACC-FC maps and calculated the targeting offset

(i.e., Euclidean distance between the individualized optimal TMS targets and the actual

stimulation coordinates) for each patient. Subsequently, we calculated the Pearson

correlations between clinical improvement (i.e., HAMD reductions) and targeting offset. We

anticipated a negative correlation between clinical outcomes and target offset (i.e., the

closer the actual stimulation target was to the individualized DR-based target, the higher the

clinical improvement). Age, sex, education, and head motion were included as covariates in

the regression models when calculating correlations between targeting offsets and clinical

improvement.

Results

Group difference maps of sgACC-FC

In the large-scale DIRECT dataset, we found significant MDD-related hyperconnectivity with

the sgACC in bilateral DLPFC, temporal parietal junction, and occipital lobe, as well as

hypoconnectivity in the bilateral temporal lobe, left inferior frontal gyrus, and left postcentral

gyrus when preprocessing included GSR (Figure 3A). When GSR was not included in

preprocessing, MDD-related sgACC FC alterations showed predominantly hypoconnectivity.

Such abnormally decreased FCs were found across the central gyrus, occipital lobe, insular

cortex, temporal lobe, and a small portion of the frontal lobe. Without GSR, MDD-related

hyperconnectivity was limited to subcortical regions (Figure S1). Given that significant

case-control differences in the DLPFC area were revealed only when GSR was implemented,

subsequent analyses were solely based on results with GSR. We thus identified two

contiguous clusters of voxels that showed significant group differences in the left DLPFC.

Group difference cluster 1 (MNI coordinates: x = -44, y = 38, z = 32; $t(2880) = 3.277$, $p < 0.001$,

Cohen's $f^2 = 0.004$) was ventral to group difference cluster 2 (MNI coordinates: x = -34, y = 36,

z = 40; $t(2880) = 3.670$, $p < 0.001$, Cohen's $f^2 = 0.005$) (Figure 3B).

Figure 3. Group differences of subgenual anterior cingulate cortex (sgACC) functional connectivity (FC) profiles are related to TMS treatment efficacy, demonstrating clinical significance. (A) Two-sample *t*-test maps of MDD-related sgACC FC abnormalities with global signal regression (GSR) implemented. (B) Two clusters showed significant case-control differences in sgACC-FC. (C) The TMS targets were extracted from the TRD-TMS dataset [37]. The sizes of the spheres indicate the magnitudes in Hamilton Depression Rating Scale (HAMD) reductions, with the group difference map rendered on the surface. (D) The magnitudes of the group differences in the FC between TMS targets and sgACC were positively related to clinical improvements in the TRD dataset.

sgACC-FC group differences correlate with TMS treatment outcomes

Using the TRD dataset, we examined the relationship between group differences (*t*-values)

and clinical outcomes (HAMD reductions) to test the clinical relevance of the group

difference maps. Group differences were positively correlated with HAMD score reductions

(r(23) = 0.448, p = 0.025, Figure 3C-D), suggesting that group-level difference maps may be

useful for enhancing the outcomes of TMS for treating MDD by improving target localization.

sgACC anticorrelation peaks in MDD and HCs

Our results highlighted the case-control differences in sgACC-FC profiles. Since the prior

sgACC group target (MNI coordinates: x = −42, y = 44, z = 30) had been based on a cohort of

young, healthy adults [37], we sought to examine potential differences in anticorrelation peaks

extracted from the mean sgACC FC maps of MDD and HC groups (Figure 4B-C) in the DIRECT

dataset. We found that the anticorrelation peak of MDD patients (MNI coordinates: x = -40, y

= 50, z = 22) differed from that of HCs (MNI coordinates: x = -42, y = 38, z = 32), probably due

to abnormal FCs within the left DLPFC in MDD patients. The anticorrelation peak extracted

from the DIRECT HCs was closer to the previously reported locus [37] (Figure 4).

Figure 4. Distribution of dual regression (DR) based individualized TMS targets guided by different

subgenual anterior cingulate cortex (sgACC)-FC statistical maps. The peak of the sgACC anticorrelation

in MDD patients differed from that in HCs. (A) The peak sgACC anticorrelation location in the average

sgACC-FC map of the MDD group in the DIRECT dataset. (B) The peak sgACC anticorrelation location in

the average sgACC-FC map of the HC group in the DIRECT dataset. (C) The sgACC-FC case-control group

difference map and the peak in the left dorsal lateral prefrontal cortex (DLPFC) in the DIRECT dataset.

(D-F) DR-based individualized TMS targets guided by the mean HC sgACC-FC map, the mean MDD

sgACC-FC map, and the group difference map, respectively. The sizes of the spheres indicate the

magnitudes of Hamilton Depression Rating Scale (HAMD) reductions.

Individualized targets suggest higher clinical efficacy than group targets

To integrate the reliable group-level statistical map information based on large samples with

individual spontaneous brain activity, we utilized a DR-based approach to calculate

individualized optimal targets for each participant based on three group-level statistic maps.

The individualized target locations are illustrated in Figure 4. The clinical relevance of the

targets was determined by the correlation between target offset and clinical improvement in

the CUD dataset. All the individualized targets (Figure 5D-F) outperformed their

corresponding group-level targets (Figure 5A-C). Among the three individualized targets, the

targets based on group-level difference maps (r(14) = -0.560, p = 0.029) outperformed the

individualized targets based on the MDD group-average template (r(14) = -0.375, p = 0.115)

or the HC group-average template (r(14) = -0.372, p = 0.117).

457
Figure 5. The individualized targets exhibited greater clinical efficacy compared to the group targets.

Clinical efficacy was characterized by computing the correlations between the offset distances of TMS
targets and the clinical improvements observed in the CUD-TMS dataset. The TMS targeting offset
distance was defined as the Euclidean distance between the actual rTMS stimulation coordinates and
the group/individualized targets. Clinical improvement was defined as the maximum HAMD reduction
during rTMS treatment, adjusted for age, sex, education, and head motion. (A-C) The clinical efficacy
of the group targets. (D-F) The clinical efficacy of the individualized targets. $*p < 0.05$.
465

Discussion

In the present study, we leveraged a large multi-site fMRI sample (1660 MDD patients and
1341 HCs) and two openly shared TMS datasets to delineate abnormalities in sgACC FC in
MDD and explore their potential impact on the localization of TMS targets. Specifically, with
GSR implemented, we found enhanced FCs between sgACC and left DLPFC, bilateral
supplementary motor areas and inferior parietal lobes, thalamus and visual areas, and
decreased FCs between sgACC and left anterior insula, left superior temporal lobe, and
bilateral temporal poles in MDD patients. Crucially, we showed that the clinical outcomes of
TMS treatments were related to the magnitude of the case-control differences in the FCs
between sgACC and TMS targets. Furthermore, the thresholded group difference maps
showed outstanding performance when identifying individualized TMS targets. Additionally,
the DR-based approach to identify individualized TMS targets showed better clinical efficacy
than TMS targets based on group sgACC-FC profiles.

MDD-related FC abnormalities of sgACC

Our results add to a growing literature documenting functional network abnormalities
involving the sgACC in MDD[8,10-12,16,18,59-61]. Nevertheless, notable discrepancies in the type of
abnormality (enhanced/reduced) and specific brain regions showing altered sgACC FCs have
been reported. Considering the small effect sizes (Cohen's $f^2 < 0.01$) of MDD-related
sgACC-FC abnormalities, the limited sample sizes in previous studies entail a high risk of false
positive findings [20,62,63]. Another potential source of heterogeneity in previous findings may
be whether or not they applied GSR (See the supplementary materials for detailed

discussion). With an unprecedented sample size, our results provide among the most robust

evidence to date. Specifically, MDD patients showed enhanced sgACC-thalamus FC and

decreased sgACC-limbic network FC regardless of whether GSR was implemented. These

results are consistent with previous studies in adolescents [18] and adults [57] with MDD.

Previous studies have reported abnormal FCs between sgACC and limbic areas and some

subcortical regions, such as the amygdala [8,19] and parahippocampus regions [12]. Together,

these results are consistent with a model highlighting sgACC as a critical hub in an "extended

medial network," which also encompasses limbic, thalamic, and striatal regions and plays a

key role in the pathophysiology of MDD [64,65]. This "extended medial network" overlaps

substantially with the DMN. Indeed, decreased FCs between sgACC and DMN regions, such

as the medial prefrontal cortex, precuneus, temporal gyrus, and parahippocampus regions,

were revealed in MDD relative to HCs when GSR was not implemented. Such abnormalities

have been previously reported [9,11,14,33], which led to the hypothesis that abnormally

enhanced FC between sgACC and DMN are the network underpinnings of rumination [66].

However, contrary to the aforementioned hypothesis, we found reduced, instead of

enhanced FC between sgACC and DMN. Similarly, in the first phase of DIRECT, we

demonstrated that MDD was characterized by reduced FC within DMN [22,23]. We note that

both the first phase and the current second phase of the DIRECT data are solely comprised of

Chinese samples, while most studies that have reported enhanced sgACC-DMN FCs have

been in Caucasian samples. Different prevalence rates [67,68], heterogeneous symptoms [69], and

different risk alleles [70] have been reported in Caucasian and Eastern Asian groups.

Accordingly, we cannot exclude racial differences contributing to this discrepancy.

Clinical relevance of abnormal sgACC-DLPFC anticorrelation in MDD

Once the group difference maps of sgACC FC profiles were delineated, we further explored

the impact of such an abnormality, especially in DLPFC, on identifying TMS targets. Most

clinical trials have focused on applying TMS to the left DLPFC based on the hypothesis that

high-frequency rTMS will enhance hypoactivity during depressive episodes [7,71]. The DLPFC is

anatomically extensive, comprising BA 46 and BA 9 [72]. However, which DLPFC sub-field is the

best target for TMS remains unclear. The current FDA-approved protocol (i.e., the "5 cm"

method) leads to large interindividual variation in stimulation sites, which may contribute to

the heterogeneity in the effect sizes of antidepressant responses in prior trials [73,74].

However, leveraging anatomical landmarks to target the boundary between BA 9 and BA 46

precisely has not consistently outperformed the "5 cm" method or the F3 Beam method [75,76].

Considering the unsatisfactory effect of TMS target localization based on brain anatomical

parcellation, group-level normative sgACC anticorrelation peaks based on healthy population

datasets have been frequently used as TMS targets in recent years [35,37,40]. However, in the

present study, we found that the locations of such anticorrelated peaks differ substantially

between MDD and HC samples when measured in a large clinical cohort. Therefore, it might

be problematic to identify TMS targets based solely on sgACC-FC profiles in healthy or

depressed individuals. Intriguingly, we found that the magnitudes of case-control differences

in TMS targets' FCs to sgACC were positively related to the clinical improvements after

receiving rTMS to the left DLPFC. Such correlation implies that the case-control differences in

the FC between sgACC and the left DLPFC might bear important information that could be

leveraged to guide the identification of reliable, individualized TMS targets.

A DR-based individualized approach may improve the clinical efficacy of TMS targets

In the current study, we developed a DR-based approach to individualize the TMS targets

derived from group-level statistical maps. The proposed approach takes advantage of the

high signal-to-noise ratio and reliability of large sample statistical maps while integrating

individual spontaneous brain activity of individuals with MDD. Most existing individualized

TMS target localization algorithms are based on calculating sgACC FCs using densely sampled

MRI images from single subjects [77]. However, individual MRI images tend to be noisy and

unreliable [78,79], especially in the sgACC region. Air in the sinuses often introduces

susceptibility artifacts due to the different magnetic properties of air and brain tissue. Signal

loss and geometric distortion are common in areas close to air-filled sinuses, such as the

inferior frontal cortex, including the sgACC [80]. The seed map approach has been proposed to

alleviate such difficulties due to the subpar image quality of the sgACC region [40,43].

Nevertheless, the sgACC is still the major contributor to the seed time series extracted for

the seed map, and the final targeting approach still relies on calculating FC using MRI images

from a single subject. A cutting-edge MDD TMS therapy combining FC-guided target

localization, high dose, and intermittent theta-burst stimulation (iTBS) was reported to be

highly effective in a randomized, double-blinded, sham-controlled clinical trial [44,45]. However,

the targeting algorithm relied on hierarchical agglomerative clustering in the sgACC area

which has a low signal-to-noise ratio. However, the final target was still determined

according to individual-level sgACC-DLPFC FCs. For the proposed DR-based approach, the

entire regression computation procedure at the individual level is conducted in the DLPFC

region, which is less affected by susceptibility artifacts. In this way, the superior signal quality

of DLPFC and the effective and reliable properties of the dual regression algorithm enhance

the accuracy of target localization.

Encouragingly, the DR-based individualization targets enhanced the clinical significance of

corresponding group-level targets, regardless of which template was used. This result

supports the generalizability and extensibility of the algorithm, offering the potential for TMS

targeting based on other circuits and biomarkers. Considering the majority of existing TMS

research still relies on the traditional 5 cm or Beam F3 methods for targeting, one approach

based on our data to improve targeting would be to simply shift the target to a more anterior

and lateral position. However, such a simple shift was not supported by the present study.

Instead, 25% of targets individualized using the group difference map were more medial than

the original targets, and 50% of targets individualized by the group difference map were

more posterior than the original targets. Therefore, the DR-based algorithm does not simply

set a more anterior and lateral coordinate in the BA46 area as the individualized optimization

target for subjects. Among the group-level templates used in the DR-based individualized

algorithm, the sgACC-FC group difference map performed the best, instead of the commonly

used average sgACC based on healthy individuals. This may reflect the abnormal posterior

shift of the MDD sgACC anticorrelation peak we found, and it emphasizes the immense

clinical value of examining spontaneous brain activity differences between MDD and HC in a

large sample. Previous studies either included only a small quantity of MDD functional MRI

data [45,46,81] or developed targeting algorithms based on large-scale HC samples [48]. In this

paper, we utilized an unprecedented amount of functional MRI data from both MDD and HC,

obtained a reliable group-level difference map of sgACC-FC, and successfully validated its

potential in treating MDD with TMS.

Limitations

Several limitations need to be considered. First, we noted that the inconsistency between

our results and the broader literature could partly be due to racial differences of the samples

(i.e., Eastern Asian vs. Caucasian). Consortiums that intend to pool existing neuroimaging

data worldwide, such as the Enhancing NeuroImaging Genetics through Meta-Analysis

(ENIGMA) Major Depressive Disorder (MDD) consortium [24], have accumulated large-scale

data primarily from Western countries and published several high-impact studies delineating

MDD-related anatomical abnormalities [82,83]. Planned collaborations between the DIRECT and

the ENIGMA-MDD consortiums may help address potential cultural, genetic, and

environmental mechanisms in more diverse groups of MDD patients. Second, we performed

volume-based preprocessing to facilitate comparison with previous literature. Surface-based

preprocessing strategies have provided more accurate and detailed representations of

cortical and subcortical structures [84]. Recent research has begun to explore surface-based

rTMS target identification algorithms and has shown promising clinical relevance [38,39,85].

Future research should consider using surface-based case-control difference maps to further

refine ways to identify TMS targets. Given the established involvement of the left DLPFC in

the existing TMS treatment protocols, the present study restricted the investigation within

this area. We noted that other brain regions (e.g., angular gyrus and supplementary motor

area) showing significant case-control differences worth further research and may serve as

potential targets for neuromodulation [86,87].

Identifying a reliable personalized TMS target solely based on an individual's R-fMRI data

(around 8 mins of fMRI scan in clinical practice) is challenging. Due to the poor replicability of

FC [88], existing individual-level network parcellation algorithms need a large quantity of fMRI

images (usually more than one hour of scanning time)[89]. The present study utilized two

openly-shared TMS samples to validate the efficacy of the individualized algorithm. However,

prospective, double-blind clinical trials are warranted to compare the treatment outcome

across different rTMS targeting algorithms (e.g., traditional anatomical landmark-based

targeting, group-level sgACC-FC targeting, and DR-based individualized targeting methods).

In addition, mounting evidence has suggested that MDD patients may benefit from rTMS

over alternative targets such as dorsomedial prefrontal cortex (DMPFC) [90]. The DR-based

individualized positioning algorithm can be readily transferred to other neural circuits or

other brain imaging-derived feature maps (e.g., ICA, functional gradient, normative

modeling). The clinical efficacy of these alternative targets is worthy of future investigations.

Conclusion

In summary, we leveraged a large sample of MDD patients to fully delineate group

differences in sgACC-FC maps between MDD patients and HCs. We next demonstrated the

impact of such case-control differences on group TMS targets based on sgACC-FC profiles by

showing that the magnitudes of case-control differences in the FC between sgACC and TMS

targets were positively associated with clinical outcomes. The peak sgACC anticorrelation

locations were different in MDD patients as compared to HCs. Moreover, we developed a

DR-based approach to identify individualized TMS targets and demonstrated that this

approach may improved clinical efficacy compared to group targets based on sgACC-FC

profiles.

Contributors

C-GY, Y-FZ, X-NZ, XC, and BL conceived and designed the study. C-GY, Y-FZ, J-PZ, and X-NZ

coordinated the collaboration. JQ, LK, T-MS, TL, K-RZ, Z-NL, L-PC, JY, X-PW, Y-GY, C-YW, C-MX,

G-RX, Y-SL, Y-QY, XW, YW, X-FX, Y-QC, Q-YG, W-BG, J-PZ, YH, H-NW, B-JL, WZ, and J-PL

supervised data collection. X-RW, QH, Y-KW, HY, A-XZ, Y-CL, J-SC, P-FS, X-YL, FL, C-CH, X-LC,

F-NJ, J-JZ, X-LJ, G-MC, Z-SC, T-LC, X-XS, TC, B-JL, M-LY, Z-PX and BL organized the data. XC and

C-GY analyzed and interpreted the data. BL and C-GY designed the dual regression-based

TMS targeting algorithm. XC, BL, and C-GY drafted the manuscript. All authors revised the

manuscript for important intellectual content and approved the final submitted version. XC,

BL, and C-GY accessed and verified the data. XC, BL, and C-GY had full access to all the data in

the study. C-GY had final responsibility for the decision to submit for publication. All authors

were responsible for the final decision to submit for publication and have seen and approved

the manuscript.

Conflict of interest

Inclusion and Ethics

Local researchers were included throughout the research process. Research protocols have

been approved by local ethics review committees. No potential risk was involved in the

current research and all local researchers have discussed and approved the research protocol.

All participants provided written informed consent, and local institutional review boards

approved each study from all included cohorts. The analysis plan of the current study has

been reviewed and approved by the Institutional Review Board of Institute of Psychology,

Chinese Academy of Sciences (No. H21102).

Data sharing

According to the success of the data sharing model of DIRECT Phase I data (REST-meta-MDD,

http://rfmri.org/REST-meta-MDD), DIRECT Phase II data will also have 2 sharing stages. 1)

Stage 1: coordinated sharing, upon the publication of this announcing manuscript. To reduce

conflict of the researchers, the consortium will review and coordinate the proposals

submitted by interested researchers. The interested researchers first send a letter of intent

to rfmrilab@gmail.com. Then the consortium will send all the approved proposals to the

applicant. The applicant should submit a new innovative proposal while avoiding conflict

with approved proposals. This proposal would be reviewed and approved by the consortium

if no conflict. Once approved, this proposal would enter the pool of approved proposals and

prevent future conflict. 2) Stage 2: unrestricted sharing, after January 1, 2026. The

researchers can perform any analyses of interest while not violating ethics. All codes have

been made openly available

(https://github.com/Chaogan-Yan/PaperScripts/tree/master/Chen_2023).

Acknowledgments

This work was funded by the Sci-Tech Innovation 2030 - Major Project of Brain Science and

Brain-inspired Intelligence Technology (No. 2021ZD0200600), the National Key R&D Program

of China (No. 2017YFC1309902), the National Natural Science Foundation of China (No.

82122035, No. 81671774 and No. 81630031), the 13th Five-year Informatization Plan of

Chinese Academy of Sciences (No. XXH13505), the Key Research Program of the Chinese

Academy of Sciences (No. ZDBS-SSW-JSC006), Beijing Nova Program of Science and

Technology (No. Z191100001119104), the Scientific Foundation of Institute of Psychology,

Chinese Academy of Sciences (No. E2CX4425YZ and No. Y9CX422005), the China

Postdoctoral Science Foundation (No. 2019M660847), the China National Postdoctoral

Program for Innovative Talents (No. BX20200360), the Special Research Assistant Program of

the Chinese Academy of Sciences (No. E2CX0624), and the Scientific Foundation of Institute

of Psychology, Chinese Academy of Sciences (grant number: E2CX4425YZ). The support

provided by China Scholarship Council (CSC, No. 202104910248) during a visit of Xiao Chen

to the Centre for Addiction and Mental Health is acknowledged. In addition, we would like to

acknowledge the valuable insights provided by Prof. B.T. Thomas Yeo, whose comments

helped to shape our understanding of the data and refine our analysis.

References

1 Malhi, G. S. & Mann, J. J. Depression. *Lancet (London, England)* **392**, 2299-2312
(2018). https://doi.org:10.1016/s0140-6736(18)31948-2

2 Drevets, W. C., Price, J. L. & Furey, M. L. Brain structural and functional abnormalities
in mood disorders: implications for neurocircuitry models of depression. *Brain*
structure & function **213**, 93-118 (2008). https://doi.org:10.1007/s00429-008-0189-x

3 Mayberg, H. S. Modulating dysfunctional limbic-cortical circuits in depression:
towards development of brain-based algorithms for diagnosis and optimised
treatment. *British medical bulletin* **65**, 193-207 (2003).

4 Mayberg, H. S. *et al.* Deep brain stimulation for treatment-resistant depression.
Neuron **45**, 651-660 (2005). https://doi.org:10.1016/j.neuron.2005.02.014

5 Disner, S. G., Beevers, C. G., Haigh, E. A. P. & Beck, A. T. Neural mechanisms of the
cognitive model of depression. *Nat Rev Neurosci* **12**, 467-477 (2011).

6 Cash, R. F. H. *et al.* Using Brain Imaging to Improve Spatial Targeting of Transcranial
Magnetic Stimulation for Depression. *Biol Psychiatry* (2020).
https://doi.org:10.1016/j.biopsych.2020.05.033

7 Fitzgerald, P. B. Targeting repetitive transcranial magnetic stimulation in depression:
do we really know what we are stimulating and how best to do it? *Brain Stimul* **14**,
730-736 (2021). https://doi.org:10.1016/j.brs.2021.04.018

8 Connolly, C. G. *et al.* Resting-State Functional Connectivity of Subgenual Anterior
Cingulate Cortex in Depressed Adolescents. *Biological Psychiatry* **74**, 898-907 (2013).
https://doi.org:http://dx.doi.org/10.1016/j.biopsych.2013.05.036

9 Greicius, M. D. *et al.* Resting-state functional connectivity in major depression:
abnormally increased contributions from subgenual cingulate cortex and thalamus.
Biol Psychiatry **62**, 429-437 (2007). https://doi.org:10.1016/j.biopsych.2006.09.020

10 Wu, H. *et al.* Changed Hub and Corresponding Functional Connectivity of Subgenual
Anterior Cingulate Cortex in Major Depressive Disorder. *Front Neuroanat* **10**, 120
(2016). https://doi.org:10.3389/fnana.2016.00120

11 Davey, C. G., Harrison, B. J., Yucel, M. & Allen, N. B. Regionally specific alterations in

functional connectivity of the anterior cingulate cortex in major depressive disorder.
Psychol Med **42**, 2071-2081 (2012). https://doi.org:10.1017/s0033291712000323

12   Cheng, B. *et al.* Functional connectivity patterns of the subgenual anterior cingulate
cortex in first-episode refractory major depressive disorder. *Brain imaging and*
behavior **15**, 2397-2405 (2021). https://doi.org:10.1007/s11682-020-00436-x

13   Sheline, Y. I., Price, J. L., Yan, Z. & Mintun, M. A. Resting-state functional MRI in
depression unmasks increased connectivity between networks via the dorsal nexus.
Proc Natl Acad Sci U S A **107**, 11020-11025 (2010).
https://doi.org:10.1073/pnas.1000446107

14   Berman, M. G. *et al.* Depression, rumination and the default network. *Social*
cognitive and affective neuroscience **6**, 548-555 (2011).
https://doi.org:10.1093/scan/nsq080

15   Wu, X. *et al.* Dysfunction of the cingulo-opercular network in first-episode
medication-naive patients with major depressive disorder. *J Affect Disord* **200**,
275-283 (2016). https://doi.org:10.1016/j.jad.2016.04.046

16   Sudheimer, K. *et al.* Decreased hypothalamic functional connectivity with subgenual
cortex in psychotic major depression. *Neuropsychopharmacology* **40**, 849-860 (2015).
https://doi.org:10.1038/npp.2014.259

17   Davey, C. G., Yücel, M., Allen, N. B. & Harrison, B. J. Task-related deactivation and
functional connectivity of the subgenual cingulate cortex in major depressive
disorder. *Frontiers in psychiatry* **3**, 14 (2012).
https://doi.org:10.3389/fpsyt.2012.00014

18   Gaffrey, M. S. *et al.* Subgenual cingulate connectivity in children with a history of
preschool-depression. *Neuroreport* **21**, 1182-1188 (2010).
https://doi.org:10.1097/WNR.0b013e32834127eb

19   Ho, T. C. *et al.* Functional connectivity of negative emotional processing in adolescent
depression. *J Affect Disord* **155**, 65-74 (2014).
https://doi.org:10.1016/j.jad.2013.10.025

20   Button, K. S. *et al.* Power failure: why small sample size undermines the reliability of
neuroscience. *Nat Rev Neurosci* **14**, 365-376 (2013).
https://doi.org:10.1038/nrn3475

21   Algermissen, J. & Mehler, D. M. A. May the power be with you: are there highly
powered studies in neuroscience, and how can we get more of them? *J Neurophysiol*
119, 2114-2117 (2018). https://doi.org:10.1152/jn.00765.2017

22   Chen, X. *et al.* The DIRECT consortium and the REST-meta-MDD project: towards
neuroimaging biomarkers of major depressive disorder. *Psychoradiology* **2**, 32-42
(2022). https://doi.org:10.1093/psyrad/kkac005

23   Yan, C. G. *et al.* Reduced default mode network functional connectivity in patients
with recurrent major depressive disorder. *Proc Natl Acad Sci U S A* (2019).
https://doi.org:10.1073/pnas.1900390116

24   Schmaal, L. *et al.* ENIGMA MDD: seven years of global neuroimaging studies of major
depression through worldwide data sharing. *Transl Psychiatry* **10**, 172 (2020).
https://doi.org:10.1038/s41398-020-0842-6

25   Yang, H. *et al.* Disrupted intrinsic functional brain topology in patients with major
depressive disorder. *Mol Psychiatry* (2021).
https://doi.org:10.1038/s41380-021-01247-2

26   Long, Y. *et al.* Altered resting-state dynamic functional brain networks in major
depressive disorder: Findings from the REST-meta-MDD consortium. *Neuroimage*
Clin **26**, 102163 (2020). https://doi.org:10.1016/j.nicl.2020.102163

27   Ding, Y. D. *et al.* Disrupted hemispheric connectivity specialization in patients with
major depressive disorder: Evidence from the REST-meta-MDD Project. *J Affect*
Disord **284**, 217-228 (2021). https://doi.org:10.1016/j.jad.2021.02.030

28   Deng, K. *et al.* Impaired robust interhemispheric function integration of depressive
brain from REST-meta-MDD database in China. *Bipolar disorders* (2021).
https://doi.org:10.1111/bdi.13139

29   Liang, S. *et al.* Biotypes of major depressive disorder: Neuroimaging evidence from
resting-state default mode network patterns. *Neuroimage Clin* **28**, 102514 (2020).
https://doi.org:10.1016/j.nicl.2020.102514

30   Liu, P. H. *et al.* Brain structural alterations in MDD patients with gastrointestinal
symptoms: Evidence from the REST-meta-MDD project. *Progress in*
neuro-psychopharmacology & biological psychiatry **111**, 110386 (2021).
https://doi.org:10.1016/j.pnpbp.2021.110386

31   Yan, C.-G., Wang, X.-D. & Lu, B. DPABISurf: data processing & analysis for brain
imaging on surface. *Science Bulletin* (2021).
https://doi.org:https://doi.org/10.1016/j.scib.2021.09.016

32   Ge, R., Downar, J., Blumberger, D. M., Daskalakis, Z. J. & Vila-Rodriguez, F. Functional
connectivity of the anterior cingulate cortex predicts treatment outcome for rTMS in
treatment-resistant depression at 3-month follow-up. *Brain Stimul* **13**, 206-214
(2020). https://doi.org:10.1016/j.brs.2019.10.012

33   Cash, R. F. H. *et al.* A multivariate neuroimaging biomarker of individual outcome to
transcranial magnetic stimulation in depression. *Hum Brain Mapp* **40**, 4618-4629
(2019). https://doi.org:10.1002/hbm.24725

34   Philip, N. S. *et al.* Network Mechanisms of Clinical Response to Transcranial Magnetic
Stimulation in Posttraumatic Stress Disorder and Major Depressive Disorder. *Biol*
Psychiatry **83**, 263-272 (2018). https://doi.org:10.1016/j.biopsych.2017.07.021

35   Fox, M. D., Buckner, R. L., White, M. P., Greicius, M. D. & Pascual-Leone, A. Efficacy of
transcranial magnetic stimulation targets for depression is related to intrinsic
functional connectivity with the subgenual cingulate. *Biol Psychiatry* **72**, 595-603
(2012). https://doi.org:10.1016/j.biopsych.2012.04.028

36   Elbau, I. G. *et al.* Functional Connectivity Mapping for rTMS Target Selection in
Depression. *Am J Psychiatry* **180**, 230-240 (2023).
https://doi.org:10.1176/appi.ajp.20220306

37   Weigand, A. *et al.* Prospective Validation That Subgenual Connectivity Predicts
Antidepressant Efficacy of Transcranial Magnetic Stimulation Sites. *Biol Psychiatry* **84**,
28-37 (2018). https://doi.org:10.1016/j.biopsych.2017.10.028

38   Cash, R. F. H., Cocchi, L., Lv, J., Fitzgerald, P. B. & Zalesky, A. Functional Magnetic

801   Resonance Imaging-Guided Personalization of Transcranial Magnetic Stimulation
802   Treatment for Depression. *JAMA psychiatry* **78**, 337-339 (2021).
803   https://doi.org:10.1001/jamapsychiatry.2020.3794

804   39   Siddiqi, S. H., Weigand, A., Pascual-Leone, A. & Fox, M. D. Identification of
805   Personalized Transcranial Magnetic Stimulation Targets Based on Subgenual
806   Cingulate Connectivity: An Independent Replication. *Biol Psychiatry* **90**, e55-e56
807   (2021). https://doi.org:10.1016/j.biopsych.2021.02.015

808   40   Fox, M. D., Liu, H. & Pascual-Leone, A. Identification of reproducible individualized
809   targets for treatment of depression with TMS based on intrinsic connectivity.
810   *Neuroimage* **66**, 151-160 (2013). https://doi.org:10.1016/j.neuroimage.2012.10.082

811   41   Gordon, E. M. & Nelson, S. M. Three types of individual variation in brain networks
812   revealed by single-subject functional connectivity analyses. *Current Opinion in*
813   *Behavioral Sciences* **40**, 79-86 (2021).

814   42   Doucet, G. E., Lee, W. H. & Frangou, S. Evaluation of the spatial variability in the
815   major resting-state networks across human brain functional atlases. *Hum Brain*
816   *Mapp* **40**, 4577-4587 (2019). https://doi.org:10.1002/hbm.24722

817   43   Cash, R. F. H. *et al.* Personalized connectivity-guided DLPFC-TMS for depression:
818   Advancing computational feasibility, precision and reproducibility. *Hum Brain Mapp*
819   (2021). https://doi.org:10.1002/hbm.25330

820   44   Cole, E. J. *et al.* Stanford Neuromodulation Therapy (SNT): A Double-Blind
821   Randomized Controlled Trial. *Am J Psychiatry*, appiajp202120101429 (2021).
822   https://doi.org:10.1176/appi.ajp.2021.20101429

823   45   Cole, E. J. *et al.* Stanford Accelerated Intelligent Neuromodulation Therapy for
824   Treatment-Resistant Depression. *Am J Psychiatry* **177**, 716-726 (2020).
825   https://doi.org:10.1176/appi.ajp.2019.19070720

826   46   Williams, N. R. *et al.* High-dose spaced theta-burst TMS as a rapid-acting
827   antidepressant in highly refractory depression. *Brain* **141**, e18 (2018).
828   https://doi.org:10.1093/brain/awx379

829   47   Cash, R. F. H. *et al.* Subgenual Functional Connectivity Predicts Antidepressant
830   Treatment Response to Transcranial Magnetic Stimulation: Independent Validation
831   and Evaluation of Personalization. *Biological psychiatry* **86**, e5-e7 (2019).
832   https://doi.org:10.1016/j.biopsych.2018.12.002

833   48   Cash, R. F. H. *et al.* Personalized connectivity-guided DLPFC-TMS for depression:
834   Advancing computational feasibility, precision and reproducibility. *Hum Brain Mapp*
835   **42**, 4155-4172 (2021). https://doi.org:10.1002/hbm.25330

836   49   Ojemann, J. G. *et al.* Anatomic localization and quantitative analysis of gradient
837   refocused echo-planar fMRI susceptibility artifacts. *Neuroimage* **6**, 156-167 (1997).
838   https://doi.org:10.1006/nimg.1997.0289

839   50   Beckmann, C. F., Mackay, C. E., Filippini, N. & Smith, S. M. Group comparison of
840   resting-state FMRI data using multi-subject ICA and dual regression. *NeuroImage* **47**
841   (2009). https://doi.org:10.1016/s1053-8119(09)71511-3

842   51   Zuo, X. N. *et al.* Reliable intrinsic connectivity networks: test-retest evaluation using
843   ICA and dual regression approach. *Neuroimage* **49**, 2163-2177 (2010).

https://doi.org:10.1016/j.neuroimage.2009.10.080

52   Garza-Villarreal, E. A. *et al.* Clinical and Functional Connectivity Outcomes of 5-Hz
Repetitive Transcranial Magnetic Stimulation as an Add-on Treatment in Cocaine Use
Disorder: A Double-Blind Randomized Controlled Trial. *Biological psychiatry.*
Cognitive neuroscience and neuroimaging **6**, 745-757 (2021).
https://doi.org:10.1016/j.bpsc.2021.01.003

53   Team, R. C. *R: A language and environment for statistical computing.* (R Foundation
for Statistical Computing, 2013).

54   Champely, S. *et al.* pwr: Basic functions for power analysis. (2017).

55   Yan, C.-G., Wang, X.-D., Zuo, X.-N. & Zang, Y.-F. DPABI: Data Processing & Analysis for
(Resting-State) Brain Imaging. *Neuroinformatics* **14**, 339-351 (2016).
https://doi.org:10.1007/s12021-016-9299-4

56   Yan, C.-G. & Zang, Y.-F. DPARSF: A MATLAB Toolbox for "Pipeline" Data Analysis of
Resting-State fMRI. *Frontiers in systems neuroscience* **4**, 13 (2010).
https://doi.org:10.3389/fnsys.2010.00013

57   Liston, C. *et al.* Default mode network mechanisms of transcranial magnetic
stimulation in depression. *Biol Psychiatry* **76**, 517-526 (2014).
https://doi.org:10.1016/j.biopsych.2014.01.023

58   Johnson, W. E., Li, C. & Rabinovic, A. Adjusting batch effects in microarray expression
data using empirical Bayes methods. *Biostatistics* **8**, 118-127 (2007).
https://doi.org:10.1093/biostatistics/kxj037

59   Greicius, M. Resting-state functional connectivity in neuropsychiatric disorders.
Current opinion in neurology **21**, 424-430 (2008).
https://doi.org:10.1097/WCO.0b013e328306f2c5

60   Cullen, K. R. *et al.* A preliminary study of functional connectivity in comorbid
adolescent depression. *Neuroscience Letters* **460**, 227-231 (2009).
https://doi.org:http://dx.doi.org/10.1016/j.neulet.2009.05.022

61   Strikwerda-Brown, C. *et al.* Mapping the relationship between subgenual cingulate
cortex functional connectivity and depressive symptoms across adolescence. *Social*
cognitive and affective neuroscience **10**, 961-968 (2015).
https://doi.org:10.1093/scan/nsu143

62   Ioannidis, J. P. A. Why Most Published Research Findings Are False. *PLOS Medicine* **2**,
e124 (2005). https://doi.org:10.1371/journal.pmed.0020124

63   Chen, X., Lu, B. & Yan, C. G. Reproducibility of R-fMRI metrics on the impact of
different strategies for multiple comparison correction and sample sizes. *Hum Brain*
Mapp **39**, 300-318 (2018). https://doi.org:10.1002/hbm.23843

64   Price, J. L. & Drevets, W. C. Neurocircuitry of mood disorders.
Neuropsychopharmacology **35**, 192-216 (2010).
https://doi.org:10.1038/npp.2009.104

65   Price, J. L. & Drevets, W. C. Neural circuits underlying the pathophysiology of mood
disorders. *Trends Cogn Sci* **16**, 61-71 (2012).
https://doi.org:10.1016/j.tics.2011.12.011

66   Hamilton, J. P., Farmer, M., Fogelman, P. & Gotlib, I. H. Depressive Rumination, the

Default-Mode Network, and the Dark Matter of Clinical Neuroscience. *Biol Psychiatry*
78, 224-230 (2015). https://doi.org:10.1016/j.biopsych.2015.02.020

67 Parker, G., Chan, B. & Hadzi-Pavlovic, D. Lower rates of depression in westernised
Chinese in the US. *J Affect Disord* **104**, 175-178 (2007).
https://doi.org:10.1016/j.jad.2007.02.020

68 Parker, G., Gladstone, G. & Chee, K. T. Depression in the planet's largest ethnic group:
the Chinese. *Am J Psychiatry* **158**, 857-864 (2001).
https://doi.org:10.1176/appi.ajp.158.6.857

69 Qiu, P., Caine, E. D., Hou, F., Cerulli, C. & Wittink, M. N. Depression as seen through
the eyes of rural Chinese women: Implications for help-seeking and the future of
mental health care in China. *J Affect Disord* **227**, 38-47 (2018).
https://doi.org:10.1016/j.jad.2017.10.016

70 Kiyohara, C. & Yoshimasu, K. Association between major depressive disorder and a
functional polymorphism of the 5-hydroxytryptamine (serotonin) transporter gene: a
meta-analysis. *Psychiatr Genet* **20**, 49-58 (2010).
https://doi.org:10.1097/YPG.0b013e328335112b

71 Fitzgerald, P. B. *et al.* An analysis of functional neuroimaging studies of dorsolateral
prefrontal cortical activity in depression. *Psychiatry Res* **148**, 33-45 (2006).
https://doi.org:10.1016/j.pscychresns.2006.04.006

72 Rajkowska, G. & Goldman-Rakic, P. S. Cytoarchitectonic definition of prefrontal areas
in the normal human cortex: II. Variability in locations of areas 9 and 46 and
relationship to the Talairach Coordinate System. *Cerebral cortex (New York, N.Y. :*
1991) **5**, 323-337 (1995). https://doi.org:10.1093/cercor/5.4.323

73 Johnson, K. A. *et al.* Prefrontal rTMS for treating depression: location and intensity
results from the OPT-TMS multi-site clinical trial. *Brain Stimul* **6**, 108-117 (2013).
https://doi.org:10.1016/j.brs.2012.02.003

74 Perera, T. *et al.* The Clinical TMS Society Consensus Review and Treatment
Recommendations for TMS Therapy for Major Depressive Disorder. *Brain Stimul* **9**,
336-346 (2016). https://doi.org:10.1016/j.brs.2016.03.010

75 Fitzgerald, P. B. *et al.* A randomized trial of rTMS targeted with MRI based
neuro-navigation in treatment-resistant depression. *Neuropsychopharmacology* **34**,
1255-1262 (2009). https://doi.org:10.1038/npp.2008.233

76 Hebel, T. *et al.* A direct comparison of neuronavigated and non-neuronavigated
intermittent theta burst stimulation in the treatment of depression. *Brain Stimul* **14**,
335-343 (2021). https://doi.org:10.1016/j.brs.2021.01.013

77 Modak, A. & Fitzgerald, P. B. Personalising transcranial magnetic stimulation for
depression using neuroimaging: A systematic review. *The World Journal of Biological*
Psychiatry **22**, 647-669 (2021). https://doi.org:10.1080/15622975.2021.1907710

78 Dubois, J. & Adolphs, R. Building a Science of Individual Differences from fMRI.
Trends in Cognitive Sciences **20**, 425-443 (2016).
https://doi.org:http://dx.doi.org/10.1016/j.tics.2016.03.014

79 Zuo, X. N. & Xing, X. X. Test-retest reliabilities of resting-state FMRI measurements in
human brain functional connectomics: a systems neuroscience perspective. *Neurosci*

Biobehav Rev **45**, 100-118 (2014). https://doi.org:10.1016/j.neubiorev.2014.05.009

80   Zald, D. H. & Rauch, S. *The orbitofrontal cortex.*   (Oxford University Press, USA,
2006).

81   Siddiqi, S. H. *et al.* Repetitive Transcranial Magnetic Stimulation with Resting-State
Network Targeting for Treatment-Resistant Depression in Traumatic Brain Injury: A
Randomized, Controlled, Double-Blinded Pilot Study. *J Neurotrauma* **36**, 1361-1374
(2019). https://doi.org:10.1089/neu.2018.5889

82   Schmaal, L. *et al.* Cortical abnormalities in adults and adolescents with major
depression based on brain scans from 20 cohorts worldwide in the ENIGMA Major
Depressive Disorder Working Group. *Mol Psychiatry* (2016).
https://doi.org:10.1038/mp.2016.60

83   Schmaal, L. *et al.* Subcortical brain alterations in major depressive disorder: findings
from the ENIGMA Major Depressive Disorder working group. *Mol Psychiatry* **21**,
806-812 (2016). https://doi.org:10.1038/mp.2015.69

84   Esteban, O. *et al.* fMRIPrep: a robust preprocessing pipeline for functional MRI. *Nat*
Methods **16**, 111-116 (2019). https://doi.org:10.1038/s41592-018-0235-4

85   Lynch, C. J. *et al.* Automated optimization of TMS coil placement for personalized
functional network engagement. *Neuron* **110**, 3263-3277.e3264 (2022).
https://doi.org:10.1016/j.neuron.2022.08.012

86   Downar, J. & Daskalakis, Z. J. New targets for rTMS in depression: a review of
convergent evidence. *Brain Stimul* **6**, 231-240 (2013).
https://doi.org:10.1016/j.brs.2012.08.006

87   Siddiqi, S. H., Khosravani, S., Rolston, J. D. & Fox, M. D. The future of brain
circuit-targeted therapeutics. *Neuropsychopharmacology* (2023).
https://doi.org:10.1038/s41386-023-01670-9

88   Zuo, X.-N. & Xing, X.-X. Test-retest reliabilities of resting-state FMRI measurements in
human brain functional connectomics: A systems neuroscience perspective.
Neuroscience & Biobehavioral Reviews **45**, 100-118 (2014).
https://doi.org:http://dx.doi.org/10.1016/j.neubiorev.2014.05.009

89   Wang, D. *et al.* Parcellating cortical functional networks in individuals. *Nat Neurosci*
18, 1853-1860 (2015). https://doi.org:10.1038/nn.4164

90   Dunlop, K. *et al.* Dorsomedial prefrontal cortex repetitive transcranial magnetic
stimulation for treatment-refractory major depressive disorder: A three-arm, blinded,
randomized controlled trial. *Brain Stimul* **13**, 337-340 (2020).
https://doi.org:10.1016/j.brs.2019.10.020

 Table 1. Demographic data for all samples included in the DIRECT II project

Site	Age (HC)[1]	Age (MDD)[1]	% Male (HC)[2]	% Female (HC)[2]	% Male (MDD)[2]	% Female (MDD)[2]	Education (HC, years)[1]	Education (MDD, years)[1]	N (HC)	N (MDD)
1	39.33 (15.80)	38.67 (13.50)	87 (35.22%)	160 (64.78%)	98 (35.13%)	181 (64.87%)	13.05 (3.89)	12.03 (3.62)	247	279
2	20.90 (2.89)	19.02 (3.79)	24 (38.71%)	38 (61.29%)	37 (27.41%)	98 (72.59%)	13.73 (1.80)	11.97 (2.55)	62	135
3	26.50 (11.22)	34.40 (10.56)	5 (25.00%)	15 (75.00%)	5 (25.00%)	15 (75.00%)	15.40 (1.47)	14.90 (2.97)	20	20
4	29.89 (9.88)	30.87 (10.39)	9 (33.33%)	18 (66.67%)	13 (41.94%)	18 (58.06%)	13.04 (3.80)	12.65 (4.22)	27	31
5	36.79 (9.07)	33.59 (9.85)	32 (56.14%)	25 (43.86%)	60 (46.88%)	68 (53.13%)	14.37 (2.64)	13.07 (3.43)	57	128
6	23.75 (5.15)	30.90 (9.29)	45 (44.55%)	56 (55.45%)	31 (52.54%)	28 (47.46%)	13.88 (2.31)	12.03 (3.06)	101	59
7	19.80 (3.85)	20.00 (6.51)	8 (26.67%)	22 (73.33%)	5 (23.81%)	16 (76.19%)	11.90 (2.55)	10.81 (2.86)	30	21
8	31.93 (9.66)	32.84 (8.95)	18 (42.86%)	24 (57.14%)	15 (34.88%)	28 (65.12%)	15.40 (1.86)	15.79 (1.82)	42	43
9	36.12 (13.85)	45.11 (14.32)	10 (40.00%)	15 (60.00%)	19 (40.43%)	28 (59.57%)	13.56 (4.01)	10.83 (4.52)	25	47
10	36.49 (12.64)	34.00 (11.81)	30 (44.78%)	37 (55.22%)	36 (45.00%)	44 (55.00%)	14.58 (2.30)	13.25 (3.08)	67	80
11	36.62 (12.11)	38.29 (12.75)	30 (44.12%)	38 (55.88%)	55 (48.67%)	58 (51.33%)	12.96 (3.57)	10.96 (3.48)	68	113
12	29.59 (5.00)	30.47 (7.20)	15 (46.88%)	17 (53.13%)	21 (33.87%)	41 (66.13%)	14.59 (2.82)	13.73 (3.39)	32	62
13	35.45 (13.76)	38.00 (14.56)	9 (31.03%)	20 (68.97%)	13 (40.63%)	19 (59.38%)	11.28 (4.31)	12.03 (3.87)	29	32
14	38.10 (11.96)	43.50 (11.28)	65 (45.14%)	79 (54.86%)	47 (34.56%)	89 (65.44%)	13.24 (4.20)	9.07 (3.60)	144	136
15	21.51 (4.58)	22.06 (4.70)	23 (65.71%)	12 (34.29%)	7 (20.59%)	27 (79.41%)	12.57 (1.42)	11.85 (2.39)	35	34

16	25.73 (7.38)	26.07 (7.66)	44 (54.32%)	37 (45.68%)	37 (44.58%)	46 (55.42%)	15.53 (2.81)	13.81 (2.60)	81	83
17	34.73 (9.66)	35.21 (9.43)	19 (38.78%)	30 (61.22%)	15 (35.71%)	27 (64.29%)	14.31 (4.21)	12.50 (4.09)	49	42
18	32.19 (11.83)	35.25 (13.15)	13 (40.63%)	19 (59.38%)	12 (50.00%)	12 (50.00%)	12.53 (3.03)	13.42 (4.47)	32	24
19	30.94 (7.32)	27.03 (11.04)	16 (50.00%)	16 (50.00%)	5 (15.63%)	27 (84.38%)	13.94 (2.84)	12.69 (3.18)	32	32
20	31.40 (10.99)	27.80 (8.99)	5 (25.00%)	15 (75.00%)	4 (20.00%)	16 (80.00%)	16.10 (3.21)	13.60 (4.36)	20	20
21	32.80 (11.06)	32.52 (11.44)	31 (55.36%)	25 (44.64%)	17 (28.33%)	43 (71.67%)	15.88 (4.59)	12.20 (4.12)	56	60
22	34.67 (13.54)	35.74 (10.02)	8 (33.33%)	16 (66.67%)	10 (28.57%)	25 (71.43%)	14.21 (2.86)	13.20 (3.75)	24	35
23	13.57 (2.18)	13.91 (1.48)	14 (50.00%)	14 (50.00%)	8 (13.79%)	50 (86.21%)	7.57 (2.18)	7.91 (1.48)	28	58

967   1 Mean (SD)

968   2 N (%)