Effectiveness of heat tolerance rice cultivars in preserving grain appearance quality under high temperatures - A meta-analysis

Hitomi Wakatsuki*, Takahiro Takimoto*, Yasushi Ishigooka**, Motoki Nishimori*, Mototaka Sakata***, Naoya Saida***, Kosuke Akagi***, David Makowski****, Toshihiro Hasegawa*

* Institute for Agro-Environmental Sciences, NARO, Tsukuba, 305-8604, Japan

** Hokkaido Agricultural Research Center, NARO, 082-0081, Japan

***Kochi Prefectural Agricultural Research Center, Nankoku, 783-0023, Japan

**** Applied Mathematics and Computer Science (MIA 518), INRAE AgroParisTech, Université Paris-Saclay, 75231, Paris, France

Email addresses

Hitomi Wakatsuki, kawatsuki55@affrc.go.jp
Takahiro Takimoto, t.takimoto@naro.affrc.go.jp
Yasushi Ishigooka, isigo@naro.affrc.go.jp
Motoki Nishimori, mnishi@naro.affrc.go.jp
Mototaka Sakata, mototaka_sakata@ken2.pref.kochi.lg.jp
Naoya Saida, naoya_saida@ken4.pref.kochi.lg.jp
Kosuke Akagi, kousuke_akagi@ken4.pref.kochi.lg.jp
David Makowski, david.makowski@inrae.fr

Corresponding Autor
Toshihiro Hasegawa
Institute for Agro-Environmental Sciences
National Agriculture and Food Research Organization (NARO)
3-1-3 Kannondai, Tsukuba, Ibaraki 305-8604, Japan
Tel: +81 29 838 8204
E-mail: thase@affrc.go.jp

Abstract

Background: Climate change, particularly rising temperatures, negatively affects rice grain quality, increasing chalky grain percentage (CG) and hampering rice grade and price. Heat-tolerant cultivars have been bred and released since the 2000s, but the effectiveness of heat tolerance in reducing the occurrence of CG has yet to be quantified.

Objectives: This study aimed to measure the effectiveness of breeding for better heat tolerance in reducing the negative impact of high temperatures on rice quality.

Methods: Through a systematic literature search, we developed a dataset including 1297 field observations covering 48 cultivars from five different heat tolerant ranks (HTRs) at 44 sites across Japan. A linear mixed-effect model (LME) and a random forest model (RF) were fitted to the data to analyze the effect of HTR and climatic factors such as the cumulative mean air temperature above 26 °C (TaHD), mean solar radiation, and mean relative humidity for 20 days after heading on CG.

Results: The LME model explained 63 % of the variation with a 14% RMSE. The RF partial dependence plot revealed that the logit-transformed CG response to climate factors was linear, supporting the assumption of LME. The statistical analysis showed that CG increased as a function of TaHD ($P < 0.001$), with significant differences among HTRs ($P < 0.001$). The strongest effect of TaHD was obtained for the lowest HTR and was found to decrease with increasing HTR. CG also increased with higher relative humidity ($P < 0.001$) and solar radiation ($P < 0.01$). Based on our modeling, we estimated that as TaHD increased from 20 to 80 °Cd (equivalent to a mean temperature increase from 27 °C to 30 °C), CG increased by 66 % points (difference in CG) for cultivars with the lowest HTR, 45 % points for cultivars with an intermediate HTR, and 19 % points for cultivars with the highest HTR. Raising HTR by just one step (from intermediate to moderately tolerant) is projected to increase the proportion of first-grade rice at a grain-filling temperature of 27 °C, but tolerance levels need to be improved further in case of stronger warming.

Conclusions: The effect of high temperatures on CG was highly dependent on the cultivar's HTR.

Improvements in HTR effectively reduce the negative impacts of high temperatures on rice grain

quality.

Significance: Heat-tolerant cultivars are projected to suppress the prevalence of CG more than

threefold compared with heat-sensitive cultivars when grain-filling temperature increases from 27

to 30 °C.

Keywords

64 Climate change adaptation, Chalky grain, High-temperature, Oryza sativa, Systematic literature

65 search

1. Introduction

Ongoing climate change, particularly rising temperatures, negatively affects rice grain yield and

quality (Krishnan et al.,2011). Since rice (*Oryza sativa*) is mostly consumed as whole grain, the quality

of rice involves a few unique traits different from other cereals, including milling, eating and cooking

properties, grain appearance, and nutritional quality (Neerja and Renu,2020). Among various quality

traits, appearance quality, especially grain chalkiness, has been of primary concern in major rice-

production areas in the world because it affects the milling, eating, and cooking properties and is

strongly affected by climate change (Cheng et al.,2005; Fitzgerald et al.,2009; Chun et al.,2010;

Lanning et al.,2011; Ali et al.,2019). Not only does the occurrence of chalky grain (CG) lower the

market price, but milling losses caused by fragile CG also reduce head rice yield and create direct

income reduction (Lyman et al.,2013). A few simulation studies estimated increasing economic losses

due to the further degradation of grain quality resulting from increased temperatures under climate

change by the mid-century in Japan (Masutomi et al.,2019) and in Europe (Cappelli and

Bregaglio,2021).

The chalkiness of grain is due to the presence of an opaque part in the grain endosperm, and

is often classified into milky-white, basal-white, and white-back/belly, depending on the locations

where opaque parts appear(Nagato and Ebata,1965). It looks opaque because light reflects

irregularly in endosperms with larger air spaces formed by loosely packed starch granules, often

caused by insufficient starch accumulation induced by high temperatures during the grain-filling

period (Morita,2008). The occurrence of CG increases as the temperature during the grain-filling

period increases. In Japan, as extremely high temperatures have become more frequent since the

beginning of the 1990s (Japan Meteorological Agency,2015), CG has been increasingly reported

across the country by the Ministry of Agriculture, Forestry, and Fisheries of Japan (MAFF,2021). To

reduce the heat-induced damage on appearance quality, genetic improvement of heat tolerance is

considered effective, and heat-tolerant cultivars have been bred and released since the 2000s

(Ishimaru et al.,2016). The cultivation area of heat-tolerant cultivars increased from 2.3 % in 2010 to

11 % in 2020, and it is expected to expand more (MAFF,2021).

However, heat tolerance levels differed even among 'heat-tolerant cultivars'. In 2018, MAFF

established nine heat tolerance ranks (HTRs) to enhance heat tolerance breeding. In addition,

assigned several check cultivars to each HTR as part of the Plant Variety Protection and Seed Act (No.

83 of 1998)(MAFF,1998; Kaji et al.,2016; Tamura et al.,2018). Rice breeders can now have a clear

breeding target and determine the HTRs of new rice cultivars by comparing their proportions of CG in

high-temperature settings (mean temperature for 20 days after heading of around 27 °C) with those

of check cultivars. HTRs could potentially be used to provide quantitative levels of heat tolerance of

rice cultivars at different temperature levels, but the effects of HTRs on CG are yet to be quantified. A

lack of quantified effectiveness of different HTRs in response to temperatures limits our ability to

develop quantitative breeding targets in specific regions and at specific global warming levels.Many

experimental studies have investigated the effects of high temperatures on grain appearance quality

of various cultivars across Japan for different purposes, including variety selection and management

improvement (See a list of references in Supplementary Materials). A few studies have quantified the

effects of air temperature and solar radiation on the occurrence of CG in specific conditions (Okada

et al.,2011; Masutomi et al.,2015; Takimoto et al.,2019). Masutomi et al. (2022) conducted a

parameter sensitivity test to demonstrate the importance of CG dependence on temperature in the

impacts of climate change on rice grain quality. It remains unclear, however, how much actual

cultivars with high HTRs can alleviate the negative impacts of ongoing and projected climate change,

which is necessary to develop concrete breeding targets in different regions and global warming

levels.

Previous modelling studies often used parameters for individual cultivars, referred to as

"genetic coefficients" (e.g., Mavromatis et al.,2001), but lack of data availability frequently limits

model applicability for a large number of cultivars. An option would be to use the same set of

coefficients for all cultivars belonging to the same HTR groups, thus reducing the need for cultivar-

specific coefficients. Quantifying the heat tolerance for HTRs has at least two advantages over

quantifying it for each cultivar. First, the effectiveness of increasing heat tolerance levels can be

easily quantified in response to global warming levels, leading to the development of breeding

targets for specific areas and temperatures. Second, each parameter can be derived using more

observations covering a wide range of environments, improving the robustness of parameters and

estimates. To utilize HTR as a quantitative measure, we need to examine if cultivars in the same HTR

behave similarly in response to major climatic variables and if high HTRs have clear advantages over

lower HTRs with increasing temperature levels. To answer these questions, we need a large dataset

covering cultivars with various HTRs obtained from widely different environments. In this study, we

conducted a systematic literature and database search to develop a nationwide dataset on rice grain

appearance quality of cultivars of different HTRs. We then performed a meta-analysis to model the

effects of temperature and other weather conditions on the CG of various HTRs. The results are used

to examine the effectiveness of cultivar improvement in reducing the negative impact of

temperature on rice quality.

2. Material and Methods

2.1. Selection of studies

135   We conducted a systematic literature search from three different sources to obtain CG data of rice

136   cultivars grown under field conditions (Figure1).

139
Figure 1. A PRISMA diagram depicting data collection and selection processes. Nc is the number of
cultivars, N is the number of studies (articles) found through the literature search, and n is the
number of observations.

The first data source is the Rice Cultivar and Characteristics (RCC) database developed by the

Institute of Crop Science, NARO (https://ineweb.narcc.affrc.go.jp/), containing breeding progress,

morphological and ecological characteristics, and appearance and taste quality characteristics of

more than 500 varieties registered by MAFF. In this study, we focused on 63 cultivars： 39 cultivars

selected as standard cultivars of different heat tolerance ranks under the Plant Variety Protection

and Seed Act (No. 83 of 1998)(MAFF,1998), and 24 cultivars released as heat-tolerant cultivars

(Supplementary Table1). Using 63 cultivars' names as search terms, we searched experimental data

in the RCC database and selected relevant data according to the following criteria. i) Experiments

were conducted under open field conditions without warming treatments. Pot experiments and

glasshouse experiments were excluded. ii) Experimental results should include CG, heading dates,

and precise experimental field locations. In this study, we defined CG as the total % of milky-white,

basal-white, belly-white, and back-white grains. We identified 427 observations of 32 cultivars to

meet our criteria from the database.

The second data source consists of four databases containing peer-reviewed scientific papers

and bulletins of national and prefectural agricultural research stations in Japan:

i) J-STAGE (https://www.jstage.jst.go.jp/browse/-char/en)

ii) CiNii Research (https://cir.nii.ac.jp/?lang=en)

iii) SCOPUS

iv) AgriKnowledge, a database for agricultural science and technology of Japan

(https://agriknowledge.affrc.go.jp/).

In December 2021, we searched J-STAGE and CiNii Research, using "Kouon-tojuku" in Japanese,

meaning "grain filling under high temperatures", as a search term, and obtained 248 and 424 results,

respectively. For the SCOPUS search, the following equation was used;

Search equation: PUBYEAR > 2001 AND (TITLE-ABS((("heat stress tolerance" OR "heat tolerance" OR

"heat stress") AND ("genotyp*" OR "cultivar" OR "grain quality" OR "nutrient quality"))) OR

AUTHKEY((("heat stress tolerance" OR "heat tolerance" OR "heat stress") AND ("genotyp*" OR

"cultivar" OR "grain quality" OR "nutrient quality")))) AND (TITLE-ABS({greenhouse gas} OR {global

warming} OR {climate change} OR {climatic change} OR {climate variability} OR {climate warming}) OR

AUTHKEY({greenhouse gas} OR {global warming} OR {climate change} OR {climatic change} OR

{climate variability} OR {climate warming})) AND NOT ((TITLE-ABS(MRV) OR AUTHKEY(MRV)))

MRV stands for Measurement, Reporting, and Verification, the process for quantifying greenhouse

gas emissions, and was used with "NOT" to exclude papers that focused on climate change mitigation.

The SCOPUS search resulted in 459 articles. In June 2022, we conducted an additional search in

AgriKnowledge using the 63 cultivars' names as a search string and collected 650 articles. From a

total of 1781 papers obtained from the four databases, we manually selected studies using the same

two criteria described earlier. This manual selection screened 1691 papers and returned 90 studies.

After removing duplicates, 50 studies (articles) were selected containing 811 field observations of

128 cultivars.

The third data source was the Performance Tests for Recommended Varieties of Rice data of

Kochi Prefecture from 2013 to 2020, provided by the Kochi Prefectural Agricultural Research Center,

containing 1094 sets of field observations of 261 cultivars. In all three sources combined, the dataset

consists of 2332 observations of 381 cultivars collected between 1994 and 2020 in 46 experimental

sites covering 33 prefectures (Figure 2).

Figure2. Geographical distribution of the 46 experimental sites in the rice quality dataset in Japan

2.2 Data extraction

Variables	Unit	Definition
Prefecture		Prefecture of experimental site
Municipality		Municipality of experimental site
Institution/Station		Name of institution and or experimental station
Latitude	°N	Latitude of experimental site
Longitude	°E	Longitude of experimental site
Altitude	m	Altitude of experimental site
Soil group		Soil classification of experimental sites
Year		Year of experiment
Cultivar		Name of cultivar
Heading DOY		Heading day of year
HTR		Heat Tolerance Rank (HTR) from 3 (susceptible) to 7 (tolerant)
Tave	℃	Average temperature of 20 days after heading
TaHD	℃ d	Cumulative mean air temperature above 26 ℃ for 20 days after heading
SR	MJ m^{-2} d^{-1}	Average solar radiation for 20 days after heading
RH	%	Average relative humidity for 20 days after heading
CG	%	% of chalky grains, a total % of milky-white, basal-white and belly/back-white grain

Milky-white	%	% of milky-white grain type CG
Basal-white	%	% of basal-white grain type CG
Belly/back-white	%	% of belly/back-white grain type CG
Reference		Source of the data records

Table1. List of variables obtained from the systematic literature search

We extracted attributes such as prefecture, location of experimental sites, latitude and

longitude (when available), soil group, cultivar name, heading date, and grain quality for 2332

observations (Table 1). Soil group to soil classification groups for each experimental site based on

literature values or classified using the Soil Inventory website. (https://soil-

inventory.rad.naro.go.jp/figure.html)(NARO). In addition to the variables listed in Table 1, the dataset

contains sieve size for grain screening and CG measurement methods, such as "manual visual

inspection" and "mechanical grain discriminator". The standard errors and numbers of replicates

were not available in the selected studies and could thus not be extracted. Of the 321 cultivars in our

dataset, 48 have been classified in one of the existing heat tolerance ranks, while the rest have not

been classified. Of the 48 cultivars with known HTRs (Table 2), 33 were listed in the MAFF guideline,

ten were rated in reference to the standard cultivars, and their HTRs were reported in the RCC

database. HTRs of the other five cultivars were obtained from the literature (Wada et al.,2010;

NARO,2011; Mizobuchi et al.,2018; NARO,2018) (Table 2). A total of 1302 observations of the 48

classified cultivars were registered in our dataset. However, data in the zero nitrogen fertilizer plots

(n = 5) were excluded from the following statistical analysis because nitrogen deficiency was

reported to increase CG substantially (Wakamatsu et al., 2008) and hinders cultivar comparisons. The

48 cultivars registered in our dataset belonged to HTRs ranging from 3 (susceptible) to 7 (tolerant)

(Table 2). Currently, cultivars with HTRs higher than 7 or lower than 3 have not been registered.

Of the 1297 observations, 245 contained CG by type, such as milky-white, basal-white, belly-

white, or back-white grains. This subset of data, referred to as "CG-type subset data", inclluded 30

cultivars covering 5 HTRs and was used to examine the environmental responses of each CG type

separately.

218   Weather data were extracted from the gridded daily meteorological dataset with a 1-km

resolution developed by NARO (Ishigooka et al.,2017). We averaged daily mean temperature, solar

radiation (SR), and relative humidity (RH) for 20 days after heading, a critical period affecting the

occurrence of CG (Morita,2008). Where shading treatments were imposed, the incident SR was

corrected using the shading ratio reported in each study. We derived TaHD as a heat stress index for

rice quality, defined as the cumulative mean air temperature above the threshold temperature (T_b)

for 20 days after heading, following earlier research (Usui et al.,2016; Ishigooka et al.,2017; Nishimori

et al.,2020) as follows;

$\text{TaHD} = \sum_{i=1}^{20} \max(Ta_i - Tb, 0)$

Ta_i is the daily mean temperature on the i-th day from the heading date (Table 1). Tb was set at 26 °C

in our study based on a preliminary test that showed that 26 °C was appropriate across different

HTRs (Supplementary Figure 1), in agreement with previous studies (Usui et al.,2016; Ishigooka et

al.,2017; Nishimori et al.,2020).

Table 2. Number of observations (n) and names of cultivars for each heat tolerance rank (HTR)
group

HTR	n	Cultivars
3	296	Akanezora[1], Akisayaka, Aoinokaze, Hinohikari, Hatsuboshi, Matsuribare, Sainokagayaki, Satojiman
4	83	Kinuhikari, Koganebare, Koisomeshi[3], Sinrei, Tachiharuka, **Yumiazusa**[2]
5	540	Akitakomachi, **Chihominori, Emidawara**, Haenuki, Harumoni[4], Hitomebore, **Inahokkori, Kankinokaze**, Koshihikari, Mutsuhomare, Nikomaru, Nipponbare
6	159	**Akiharuka**, Akisakari, Fukei227, Hanaechizen,Koganemasari, **Koinoyokan**, Kokoromachi, Mineharuka, **Nijinokirameki**, , Nishihikari, Shifukunominori, Tochiginohoshi, **Tsukiakari,** Tsuyahime, **Tsuyakirari**
7	224	Eminokizuna, Fusaotome, Genkitsukushi[5], Natsuhikari[6], Natsuhonoka, Otentosodachi, Yosakoibijin[6]

	Total	1302

1) Cultivars in plain text are standard cultivars designated to each HTR in the MAFF guideline
(MAFF,1998).
2) Cultivars in bold text are recently released cultivars registered in the RCC database with their HTR
classified following the MAFF guideline.
3) HTR based on NARO (2018)
4) HTR based on NARO (2011)
5) HTR based on Wada et al. (2010).
6) HTR based on Mizobuchi et al. (2018)
242

2.3 Statistical analysis

All analyses were conducted using R (R Core Team,2022). We fitted a parametric linear mixed effects

(LME) model and a non-parametric machine learning model (random forest, RF) to the whole dataset

od 1297 observations to quantify the effects of weather variables (TaHD, mean RH, and SR) and HTR

on CG. The following logit transformation was applied to stabilize the variance of CG proportion data

(Yandell,1997), logit (CG) = ln (CG/(100-CG)). The variables TaHD, mean RH, and SR were treated as

numerical variables and HTR as a factor. The use of two very different types of models (LME and RF)

allowed us to test the robustness of the results to the model assumptions.

The following LME model was applied to examine the effects of HTRs and environmental

factors on CG;

$$logit\left(CG_{ijk}\right) = \beta_1(HTR_j) + \beta_2(HTR_j) \times TaHD_i + \beta_3 SR_i + \beta_4 RH_i + u_i + \varepsilon_{ijk} \quad (1)$$

where i, j, and k are the indices of the reference-site (i.e., unique combination of article reference

and site name), cultivar, and sub-treatment number for each site-cultivar, $\beta_1, ..., \beta_4$ are fixed

parameters, u_i is a random reference-site effect, ε_{ijk} is a random residual term. The sub-treatment

includes different nitrogen applications, planting patterns, or previous crops. Only TaHD × HTR

interaction was included in this model because our preliminary tests indicated interactions between

other environmental variables and HTR were not significant. Another preliminary test revealed that

the inclusion of soil classification (group) did not improve the model goodness of fit, and so it was not

included in the model. The model was fitted using the lme4 R package (Bates et al.,2015). The

significance of the effects of explanatory variables was tested using ANOVA, and differences among

HTRs were tested using the marginal means of CG estimated using the emmeans package

(Lenth,2022). We also used the metafor package (Viechtbauer,2010) to illustrate CG responses to

climatic factors with 95 % confidence intervals computed using bootstrapping.

We also used an RF model to determine the effects of HTR and climatic factors on CG

without making a linear assumption and without the need to select any specific interaction (all

interactions are taken into account in RF). This model was implemented using the randomForest

package with the following settings (mtry = 2, ntree = 500) (Liaw and Wiener,2001). CG was also

logit-transformed before training RF. The importance of variables in the RF model was evaluated with

the mean increase in mean square error (MSE) (computed by random permutation of the model

input values), and the partial dependence plot function was used as an analysis tool to detect

influential explanatory variables and then show how the average RF model responses are influenced

by each predictor.

Both LME and RF models were fitted/trained with 80% of the total observations selected

randomly while keeping the same proportions of HTR categories as in the entire dataset using the

"sample" function in R. The fitted/trained models were then evaluated by computing the root mean

square error (RMSE) with the test data, which accounted for 20% of all observations.

To examine the effects of HTR and environmental factors on different types of CG, the same

LME model (Eq. (1)) was fitted to CG by type recorded in the sub-datasets, leading to three separate

models. CG data by type were logit-transformed, but data with CG = 0% (n = 5) were excluded

because they could not be logit-transformed. As a result, a subset of 240 observations was used for

this analysis. For each CG type, an LME model was trained using 80% of all observations randomly

selected from the subset, following the method described earlier. The root mean square error

(RMSE) was calculated using 20% of the remaining data. We also fitted an LME model to the whole

CG data (sum of three types) in the same manner, and we compared the model accuracy of CG

estimated in two ways to determine whether the predictions of CG by type were more accurate than

those obtained from a single model fitted to the aggregated CG.

3. Results
3.1 Grain quality observed in rice cultivars with different heat tolerance

The dataset used in our analysis contained 1297 CG observations for 48 cultivars of five different

HTRs, obtained at 44 sites (196 site × year combinations) across Japan. HTR5 (intermediate tolerance

level) recorded the most observations, accounting for 41% of the dataset, followed by HTR3

(susceptible, 23%) (Table 2). HTR4 (moderately susceptible) had the fewest observations but still had

over 80 observations. CG spanned from 0 to 100 % (Supplementary Figure 2a), with a skewed

distribution to the right. The overall mean CG was 20.7 %, 5.4 % points(percentage-points) greater

than the median (15.3 %) (Table 3). Logit-transformed CG data, however, showed a bell-shaped

symmetric distribution (Supplementary Figure 2b). The average values of CG obtained for each HTR

class ranged from 12 to 29%, and declined as HTR increased (Table 3).

The logit-transformed CG data also exhibited a symmetric bell-shaped distribution when

analyzed for each type of CG separately (Supplementary Fig. 4). The mean CG obtained for each HTR

ranged from 9 to 28%, similar to the whole dataset. All three CG types (milky-white, basal-white, and

belly/back white) decreased with increasing HTR (Supplementary Figure 5).

Air temperature, relative humidity (RH), and solar radiation (SR) averaged over 20 days after

heading varied greatly among 196 site-years. The mean air temperature ranged from 20.8 °C to 30 °C

with a median of 26.5 °C, implying that more than half of the observations were exposed to

temperatures above the threshold temperature of 26°C (Supplementary Figure 2c). Mean SR and RH

ranged from 7.4 to 23.8 MJ m^{-2} d^{-1} and from 64.2 % to 94.7 %, respectively (Supplementary Figures

2d and 2e). Weather conditions were similar among HTRs (Table 3, Supplementary Figure 2c-e). CG

increased almost linearly with TaHD in all HTRs, but the increase in CG values was greater in low HTR

levels than in high HTR levels (Figure 3).

Table3. Mean measured CG (%), mean and standard error and weather conditions (averaged over 20
days after heading) for each HTR

HTR	Mean CG (%) +SE			Median CG (%)	Mean Temperature (°C)	TaHD(°C d)	Mean SR (MJ m² d⁻¹)	Mean RH (%)
3	29.4	±	6.4	24.5	26.5	24.1	16.1	78.1
4	26.4	±	6.1	22.4	26.9	28.9	16.7	76.3
5	20.6	±	5.2	16.5	26.3	21.5	16.7	80.2
6	14.6	±	4.2	11.8	26.5	23.3	16.2	80.0
7	11.8	±	3.7	8.2	26.0	18.9	16.0	82.2
Total	20.7	±	4.2	15.6	26.3	21.2	16.6	80.6

319 **Figure3**. Boxplot of measured CG (%) in relationship to TaHD (°C d, representing the sum of
320 temperature above 26°C, 20 days after heading) for each HTR category. Each box indicates the
321 interquartile range (IQR) and the middle line in the box represents the median. The upper- and
322 lower-end of whiskers are median 1.5 ×IQR ± median. Open circles are values outside the
323 1.5 ×IQR.

3.2 Model comparison

We first applied the LME model (Eq.1) to test if the effect of temperature on CG differed among HTRs

and if other environmental factors had a significant effect on CG. Overall, the model explained 63%

of the CG when accounting for the reference-site effects, and 33% without (i.e., based on climate-

and HTR-related inputs only). The RMSE of LME with the test data set was about 14% without

accounting for reference-site effects (Figure 4).

The LME results revealed a strong effect of TaHD on CG (Table 4, $P < 0.001$), and a significant

interaction between TaHD and HTRs ($P < 0.001$). The TaHD effect (i.e. coefficient) was greatest in

HTR3 and decreased progressively as HTRs increased. Intercepts also differed significantly among

HTRs, being smaller in HTR5, 6, and 7 than in HTR3 ($P < 0.001$). The effects of SR and RH were both

significantly positive ($P < 0.005$ and $P < 0.001$, respectively), with RH having a larger influence than SR

(Table 4). The marginal mean CG values revealed that increasing HTRs from 4 to 5 could significantly

reduce CG by 7.7 % points and that increasing HTRs from 5 to 6 would decrease CG even further, by

5.0 % points, when the other environmental factors (SR and RH) were set to their mean values.

We also tested the RF model using the same explanatory variables and found that it

explained 49% of the CG variation of the train data set. The RMSE with the test data set was about

14%, comparable to that of LME (Figure 4). The variable importance analysis of the RF model showed

that TaHD was the most influential variable, followed by HTR, RH, and SR, in close agreement with

the LME model (Table 4, Supplementary figure 3).

The LME model applied to the CG by type showed that all three CG types increased with

TaHD, but that the TaHD effect tended to become progressively smaller with increasing HTRs, as

evidenced by the significant interaction between HTR and TaHD (Supplementary Table 2). The effect

of SR was not significant, while that of RH was significant except for basal-white (P<0.001 for milky-

white and belly/back-white). Still, the estimates of coefficients were all positive in agreement with

those for the whole CG (Supplementary Figure 6). We compared the accuracy of CG prediction

obtained from the model fitted for the aggregated CG with that from the models fitted separately for

three CG types and found that both predictions gave similar RMSE (15.4 % and 16.5 %, respectively)

(Supplementary Figure 7).

Table 4. ANOVA results of the rice LME model, estimated parameters, and estimated marginal means
for each HTR.

	Variables	Mean Square	DFn[1]	DFd[2]	F value	P[3]
ANOVA	TaHD	105.6	1	1117	176.1	0.000***
	HTR	15.06	4	1257	24.99	0.000***
	SR	4.534	1	811	7.558	0.006**
	RH	25.70	1	1137	42.85	0.000***
	TaHD x HTR	4.325	4	1236	7.209	0.000***
	HTR	Estimate[4]	SE	df	t value	P
Intercepts						
	HTR3	-7.314	0.855	987	-8.557	0.000***
	HTR4-HTR3	0.025	0.193	1282	0.127	0.899
	HTR5-HTR3	-0.501	0.102	1270	-4.922	0.000***
	HTR6-HTR3	-0.906	0.131	1248	-6.891	0.000***
	HTR7-HTR3	-1.020	0.114	1242	-8.933	0.000***
Coefficients for covariates						
TaHD	HTR3	0.046	0.003	1245	13.56	0.000***
	HTR4-HTR3	-0.012	0.005	1242	-2.204	0.028*
	HTR5-HTR3	-0.011	0.003	1245	-3.351	0.001**
	HTR6-HTR3	-0.014	0.004	1228	-3.352	0.001**
	HTR7-HTR3	-0.020	0.004	1227	-5.156	0.000***
SR		0.042	0.015	811	2.749	0.006**
RH		0.057	0.009	1137	6.546	0.000***

	Estimated marginal mean CG (%)[5]	Confidence limits (%)	
		Lower	Upper
HTR 3	26.5 a	22.8	30.8
HTR 4	22.3 a	17.8	27.3
HTR 5	14.6 b	12.3	17.2
HTR 6	9.6 c	7.9	11.7
HTR 7	7.7 d	6.2	9.3

1) Degree of freedom for the numerator

2) Degree of freedom fo the denominator estimated using the Kenward-Roger method in the lme4 R
package (Bates et al., 2015).

3) ***, P< 0.001; **, P < 0.01; and * , P<0.05 .

4) Estimated parameters in Eq. (1) based on the logit-transformed CG.

5) Estimated marginal means based on the untransformed CG. Values followed by the same
alphabetic letters are not statistically different at P=0.05 using Tukey's method. Note that the
difference between HTR6 and 7 was found significant, whereas that between HTR 3 and 4 was not,
despite it being greater in the latter comparison. This was because the pairwise comparison was
made using the logit-transformed data; the difference was greater in the HTR6-7 comparison than in
the HTR3-4.

Figure 4. Relationships between observed and predicted values from the LME (a in logit scale, d in %) and the RF model (b in logit scale, e in %), and between the predicted values of the two models (c in logit scale, f in %).

3.3 Responses of CG to temperature, solar radiation, and humidity

Both models predicted similar responses of Logit (CG) to TaHD, SR, and RH (Supplementary Figure 3).

The partial dependence plots of the RF model indicated that Logit(CG) responded almost linearly to

all climatic variables studied, confirming that the linear assumption of the model LME was plausible.

Since the RF and LME models showed similar goodness of fit (Section 3.2) and responses to climatic

factors, we chose the LME model to describe the responses of CG to TaHD, RH, and SR for the

different HTRs, due to its simplicity and interpretability.

CG increased as a function of TaHD for all HTRs, but the response was much weaker for high

HTRs than for low HTRs (Figure 5). When SR and RH were fixed to their mean observed values,

increasing TaHD from 20 to 80 °C d was projected to increase CG by 66 % points for HTR3, 45 %

points for HTR5, and 19 % points for HTR7, confirming the effectiveness of heat-tolerant cultivars in

reducing the negative impacts of high temperatures. The projected responses also showed that CG

increased with both increasing RH and SR (Figure 5) to a greater extent for low HTRs than for high

HTRs. When TaHD and SR were held constant, increasing RH from 70 to 90% was projected to

increase CG by 22 % points for HTR3, 14 % points for HTR5, and 8 % points for HTR7. The effect of

HTR was smaller for SR; increasing SR from 10 to 22 MJ m^{-2} was estimated to raise CG by 9.8 % points

for HTR3, 6 % points for HTR5, and only 4 % points for HTR7 (Figure 5)

Figure 5. Response of CG (%) to TaHD (a), RH (b), and SR (c) estimated with the LME model for each
heat tolerance rank (HTR). Error bars show 95% confidence intervals. Response to each weather
variable was drawn by fixing other two environmental variables at their average values in the
dataset: SR = 16 MJ m^{-2} d^{-1}, RH = 80%, and TaHD = 22 °C d.

4. Discussion

Using a systematic literature search strategy, we collected 2,332 field trials of 321 cultivars conducted at 46 experimental stations across Japan to develop a quantitative effectiveness measure against heat damage on rice quality. The dataset covers all regions except Hokkaido, where temperatures are low and heat damage has yet to be reported. The dataset contains established national-level standardized cultivars with contrasting levels of heat tolerance. All the data were obtained in open fields, without additional warming, in conditions close to real farms. The dataset also includes key climatic variables and geographical information for each observation, facilitating the analysis of quality-climate relationships.

We tested two different models; a parametric model based on mixed-effect regression (LME) and a non-parametric model based on the random forest (RF) algorithm. The LME model assumes a linear relationship between the response and explanatory variables, while the RF model makes no linear assumption but is more difficult to interpret (Xie et al.,2021). Our results demonstrated that the two models agreed on their capacity to predict the CG variation in the test data (RMSE = 14 %), the importance of explanatory variables (TaHD > HTR > RH > SR), and the linear responses of CG to weather-related variables. Both models were able to detect the negative effect of high temperature on quality, which was found to be highly dependent on cultivars' HTR. Each HTR group includes 6–15 cultivars (Table 2), bred in different locations and years from various backgrounds. They were selected as reference cultivars for each HTR based on their performance under high grain-filling temperatures (mean temperatures for 20 days after heading are typically around 27–28 °C). (Kaji et al.,2016; Tamura et al.,2018). The proposed models demonstrated that increasing HTR could reduce the occurrence of CG at high temperatures (>26 °C), and we found that a shift of HTR by one rank could decrease CG by more than 5%.

Grading of rice grain quality in Japan is determined based on several criteria, the most important one being the percentage of undamaged grains as described in Article 3 of the Agricultural

Products Inspection Law, where the first-grade rice must have at least 70% of undamaged grains.

Chalky grains are considered immature and one of the major causes of lowering the rice grade,

particularly under high temperatures during the grain filling period. Earlier, Nishimori et al. (2020)

showed that, from a nationwide survey in Japan, the percentage of first-grade rice sharply dropped

by 3-4 % points each % point rise in CG when CG exceeded 5 %. When CG exceeded 25 %, the first-

grade rice % was nearly zero and was replaced by second or third-grade rice. Reducing CG by raising

HTR can help to maintain appearance quality (Table 4) and, thus, quality grade in hotter climates. For

instance, at a mean grain-filling temperature of 27 °C (0.6 °C higher than the average of the entire

dataset, Table 3, equivalent to TaHD = 20 °C d), CG was estimated to be 24.5 % (95% CI ranging from

21.2 to 27.75%) for HTR3 (susceptible) (Figure 5), implying that it is already difficult for HRT3 cultivars

to produce first-grade rice. At the same grain-filling temperature, CG was calculated to be 13.6 %

(95% CI, 11.4 ~ 15.9 %) with HTR5 (intermediate) and 9.0 % (7.5~10.9) with HTR6 (moderately

tolerant) (Figure 5), indicating that raising HTR by just one step could substantially contribute to

attaining first-grade rice. When the air temperature further rises, however, the adaptation limit is

reached, even with HTR6 or 7. For instance, at a grain-filling temperature of 28.5 °C (TaHD = 50 °C d),

CG was projected to reach 20% (16.2~25.4) for HTR6 and 14% (11.1~18.1) for HTR7, indicating that

the reduction of first-grade rice is inevitable, calling for further improvement in heat tolerance.

Our results confirmed that grain-filling temperature is the dominant climatic factor affecting

CG. The variable TaHD (cumulative temperature above the threshold value 26°C) was found to be a

key factor, as reported elsewhere (Nishimori et al.,2020). Some researchers used a different base

temperature for different genotypes, assuming that heat tolerance would raise the threshold

temperature (Takimoto et al.,2019; Masutomi et al.,2022). We also examined this possibility by

varying the base temperature for each HTR, but found that the best model fit was obtained with a

base temperature of about 26 °C regardless of HTRs (Supplementary Figure 1). We, therefore, fixed

the base temperature at 26 °C for all HTRs. It is possible that the intercepts of the LME model, which

also differed significantly among HTRs, accounted for the CG variation near the threshold

temperature.

Two other climatic factors, Relative Humidity (RH) and Solar Radiation (SR), were also found

to have significant effects on CG according to the LME model. CG increased with both RH and SR, but

the effect of RH was stronger, with the F value being 43 for RH and 7.6 for SR (Table 4). To our

knowledge, only a few have studies reported the effects of RH on CG; one of them used a growth

chamber to impose RH treatments, and the other statistically analyzed climatic factors on CG of field-

grown rice in wet and dry seasons (Wakamatsu et al.,2009; Zhao and Fitzgerald,2013), both

demonstrating that CG increased with RH in line with the current findings. The effect of SR on CG has

been frequently reported (Deng et al.,2018; Ishimaru et al.,2018), with evidence that a specific type

of chalk, milky-white grains, which depends on assimilate supply (Kobata et al.,2004; Tsukaguchi and

Iida,2008), increased under low SR (Ishimaru et al.,2018). Some modelling studies have already

considered the effects of SR (Okada et al.,2011; Yoshida et al.,2016; Takimoto et al.,2019),

accounting for the rise in CG when solar radiation is limited. Our analysis showed, however, that high

SR had a positive impact on CG although the effect of SR was much smaller than that of TaHD or RH

(Table 4, Figure 5). Our additional analysis based on a subset data for three types of CG revealed that

the dependence of CG on SR was either positive or close to 0 in all CG types (Supplementary Table2

and Figure 6). This could be explained in part by the fact that an important mechanism by which SR

increases CG despite its positive effects on assimilate supply is greater incident radiation, which

raises panicle temperatures. High RH could also raise panicle temperatures by reducing

transpirational cooling (Yoshimoto et al.,2022). The fact that both RH and SR increased CG suggests

that the indirect impacts of these climatic factors on CG via panicle temperatures override a

beneficial effect of SR through increased assimilate supply, which could reduce the occurrence of

milky-white grains. It is important to note that increases in CG with rising RH and SR decreased with

higher HTRs, implying that the effects of RH and SR were heat-mediated (Figure 5).

Our analysis by CG type revealed that HTR was effective in reducing all three types of CG

(milky-white, basal-white, and belly/back-white grains). The analysis also confirmed the significant

TaHD x HTR interactions, with a stronger interaction for basal-white and belly/back-white grains

(Supplementary Table 2 and Figure 6) for their sensitivity to high temperatures (Wakamatsu et

al.,2007). Our model comparison did not show any advantage in distinguishing different types of CG,

and thus supports the use of the whole CG as the response variable (Supplementary Figure 7).

Our models can play at least two roles in genetic improvements in heat tolerance. First,

when combined with climate scenarios, the model can be used to backcast quantitative breeding

targets that are temporally and spatially explicit. Breeding targets can also be defined based on

global warming levels. Because the lead time in breeding is often long, the breeding target—heat

tolerance level in the current context—must be developed well before variety release. Current

temperatures during the grain-filling period vary across regions. For instance, our dataset showed

grain-filling temperatures averaged for each prefecture in the recent decade, ranged from below

25 °C to above 28 °C. They are projected to rise over time as global warming proceeds, strongly

suggesting that the breeding target for HTR needs to be tailored for each region and time. The

current models can estimate CG from major climatic factors for different tolerance levels and can

help design locally specific targets. The models are built on the Japanese rice dataset but might be

applied to other countries/regions where standard cultivars are already established. Second, the

models may be used to derive objective and quantitative heat tolerance ranks for new cultivars

based on field trials and their environmental conditions. Heat tolerance ranks of new cultivars are

currently determined based on relative cultivar performances, established comparing candidate lines

with standard cultivars. The proposed models may potentially assist expert judgment by providing

quantitative tolerance levels from multiple field data sets, although further testing of the model is

needed for this purpose. Heat tolerance is required for other characteristics, particularly heat-

induced spikelet sterility, which can substantially affect cereal yield (Ye et al.,2021). However,

because the growth processes and temperature sensitivity for spikelet sterility are different from

those for grain quality in threshold values and sensitive stages (Sanchez et al.,2014), modelling and

enhancing heat tolerance for high-temperature-induced spikelet sterility requires different strategies

that should be addressed separately from the present study.

In principle, the proposed models could be used to estimate the effects of weather- and

cultivar-related inputs, but many other factors are involved in the CG variation. The LME model

explained 63 % of the variations when accounting for the reference-site effects. Other factors may

include some management factors such as nitrogen application (Wakamatsu et al.,2008; Tanaka et

al.,2010), planting density (Morita et al.,2012), and water management (Chiba et al.,2017). In this

study, most data were from cultivar trials on experimental stations where standard levels of

fertilizers were applied and sufficient irrigation was possible, which might have decreased the CG

variation compared to farmers' fields. Experimental sites had different soils, but adding soil groups

did not improve the prediction accuracy (data not shown). Nevertheless, diverse fertilizer

applications, soil fertility, and irrigation methods could exist in actual farmers' fields. Model

adjustments, incorporating more covariates related to soil or management practices, may be

necessary for the model to be widely applicable in the future.

Because the model is empirical, it cannot be extrapolated. The maximum TaHD in the dataset

was 80 °C d, which is equivalent to the average grain-filling temperature of 30 °C. According to the

climate projections based on various greenhouse gas emissions, the mean air temperature in Japan

may rise by 0.6 ~ 3.0 °C and by 1.1 ~ 6.7 °C from the 1981-2000 period by mid- and late-century,

respectively (Ishigooka et al.,2017), suggesting that grain-filling temperatures could exceed 30 °C,

particularly in currently warm regions. In addition, the current model does not account for

atmospheric CO_2 concentration or wind speed. Notably, the effect of CO_2 is relevant in the context of

climate change because elevated CO_2 exacerbates the heat-induced degradation of rice appearance

quality (Yang et al.,2007; Usui et al.,2016), which could dramatically influence the projected impacts

of climate change on grain appearance quality (Ishigooka et al.,2021). There are a few reports that

show heat-tolerant cultivars effectively sustain appearance quality under elevated CO_2 levels (Usui et al.,2014; Hasegawa et al.,2019), but this needs to be examined more extensively. Further expansion of the rice CG dataset and model improvement will be needed.

Conclusions

We aimed to quantify the effects of improving cultivars' heat tolerance ranks (HTRs) to alleviate heat damage on rice grain quality using recently established national-level standardized cultivar heat tolerance ranks in Japan. A linear mixed effect model and a random forest model were developed and tested with a dataset containing 1,297 field trials of 48 cultivars from 44 experimental stations across Japan, covering all regions where high temperatures are a source of concerns. The two models agreed on their capacity to predict the variation in chalky grain percentage (CG) in test data (RMSE = 14 or 15%), the importance of explanatory variables, and linear responses to climatic variables. The proposed models demonstrated that increasing HTR could reduce the occurrence of CG, especially under high temperature conditions. Increasing the HTR by a single step (from intermediate to moderately tolerant) can significantly increase the chances of obtaining first-grade quality rice under grain filling temperatures of up to 27°C, but tolerance levels must be improved if more extreme temperatures are expected. The proposed models can be used to backcast quantitative, temporally, and spatially explicit breeding targets under climate change.

Acknowledgements

This study was partly supported by the Environment Research and Technology Development Fund (JPMEERF20S11820) of the Environmental Restoration and Conservation Agency of Japan; JST Grant Number JPMJPF2013 supported this work; the Joint Linkage Call supported by NARO and INRAE; and the project CLIMAE of INRAE.

References

Ali, F., Waters, D.L.E., Ovenden, B., Bundock, P., Raymond, C.A., Rose, T.J., 2019. Heat stress during grain fill reduces head rice yield through genotype dependant increased husk biomass and grain breakage. Journal of Cereal Science 90. doi:10.1016/j.jcs.2019.102820

Bates, D., Mächler, M., Bolker, B., Walker, S., 2015. Fitting Linear Mixed-Effects Models Usinglme4. Journal of Statistical Software 67. doi:10.18637/jss.v067.i01

Cappelli, G.A., Bregaglio, S., 2021. Model‐based evaluation of climate change impacts on rice grain quality in the main European rice district. Food and Energy Security 10. doi:10.1002/fes3.307

Cheng, F.M., Zhong, L.J., Wang, F., Zhang, G.P., 2005. Differences in cooking and eating properties between chalky and translucent parts in rice grains. Food Chemistry 90, 39-46. doi:10.1016/j.foodchem.2004.03.018

Chiba, M., Terao, T., Watanabe, H., Matsumura, O., Takahashi, Y., 2017. Improvement in Rice Grain Quality by Deep-flood Irrigation and its Underlying Mechanisms. Japan Agricultural Research Quarterly: JARQ 51, 107-116. doi:10.6090/jarq.51.107

Chun, A., Song, J., Kim, K.-J., Lee, H.-J., 2010. Quality of head and chalky rice and deterioration of eating quality by chalky rice. Journal of Crop Science and Biotechnology 12, 239-244. doi:10.1007/s12892-009-0142-4

Deng, F., Wang, L., Pu, S.-L., Mei, X.-F., Li, S.-X., Li, Q.-P., Ren, W.-J., 2018. Shading stress increases chalkiness by postponing caryopsis development and disturbing starch characteristics of rice grains. Agricultural and Forest Meteorology 263, 49-58. doi:10.1016/j.agrformet.2018.08.006

Fitzgerald, M.A., McCouch, S.R., Hall, R.D., 2009. Not just a grain of rice: the quest for quality. Trends in Plant Science 14, 133-139. doi:10.1016/j.tplants.2008.12.004

Hasegawa, T., Sakai, H., Tokida, T., Usui, Y., Nakamura, H., Wakatsuki, H., Chen, C.P., Ikawa, H., Zhang, G., Nakano, H., Matsushima, M.Y., Hayashi, K., 2019. A High-Yielding Rice Cultivar "Takanari" Shows No N Constraints on CO2 Fertilization. Front Plant Sci 10, 361. doi:10.3389/fpls.2019.00361

Ishigooka, Y., Fukui, S., Hasegawa, T., Kuwagata, T., Nishimori, M., Kondo, M., 2017. Large-scale evaluation of the effects of adaptation to climate change by shifting transplanting date on rice production and quality in Japan. Journal of Agricultural Meteorology 73, 156-173. doi:10.2480/agrmet.D-16-00024

Ishigooka, Y., Hasegawa, T., Kuwagata, T., Nishimori, M., Wakatsuki, H., 2021. Revision of estimates of climate change impacts on rice yield and quality in Japan by considering the combined effects of temperature and CO2 concentration. Journal of Agricultural Meteorology 77, 139-149. doi:10.2480/agrmet.D-20-00038

Ishimaru, T., Hirabayashi, H., Sasaki, K., Ye, C., Kobayashi, A., 2016. Breeding efforts to mitigate damage by heat stress to spikelet sterility and grain quality. Plant Production Science 19, 12-21. doi:10.1080/1343943X.2015.1128113

Ishimaru, T., Nakayama, Y., Aoki, N., Ohsumi, A., Suzuki, K., Umemoto, T., Yoshinaga, S., Kondo, M., 2018. High temperature and low solar radiation during ripening differentially affect the composition of milky-white grains in rice (Oryza sativa L.). Plant Production Science 21, 370-379. doi:10.1080/1343943X.2018.1520047

Japan Meteorological Agency, 2015. Climate Change Monitoring Report 2014. Japan Meteorological Agency, , Tokyo, Japan

Kaji, R., Ohta, H., Fukushima, A., Tsuda, N., Moriyama, S., Kon, C., Endo, T., Nakagomi, Y., Sato, H., Kawamoto, T., Kato, K., Goto, H., Abe, Y., Sasaki, S., Yoshida, N., Otera, M., 2016. Selection of Standard Rice Varieties for Evaluating Grain Quality under High Temperature during Ripening Stage

for Tohoku Region. Bulletin of TOHOKU Agricultural Research Center 118, 49-55.
doi:10.24514/00001292

Kobata, T., Uemuki, N., Inamura, T., Kagata, H., 2004. Shortage of Assimilate Supply to Grain
Increases the Proportion of Milky White Rice Kernels under High Temperatures. Japanese Journal
of Crop Science 73, 315-322. doi:10.1626/jcs.73.315

Krishnan, P., Ramakrishnan, B., Reddy, K.R., Reddy, V.R., 2011. Chapter three - High-Temperature
Effects on Rice Growth, Yield, and Grain Quality. In: Sparks, D.L. (Ed.), Advances in Agronomy.
Academic Press, pp. 87-206.

Lanning, S.B., Siebenmorgen, T.J., Counce, P.A., Ambardekara, A.A., Mauromoustakos, A., 2011.
Extreme nighttime air temperatures in 2010 impact rice chalkiness and milling quality. Field Crop
Res 124, 132-136. doi:10.1016/j.fcr.2011.06.012

Lenth, R.V., 2022. emmeans: Estimated Marginal Means, aka Least-Squares Means.

Liaw, A., Wiener, M., 2001. Classification and Regression by RandomForest. Forest 23.

Lyman, N.B., Jagadish, K.S., Nalley, L.L., Dixon, B.L., Siebenmorgen, T., 2013. Neglecting rice milling
yield and quality underestimates economic losses from high-temperature stress. PLoS One 8,
e72157. doi:10.1371/journal.pone.0072157

MAFF, 1998. The guideline for variety registration, "Rice". In: MAFF (Ed.), The plant variety protection
and seed act

MAFF, 2021. Climate change impacts report 2020. Ministry of Agriculture, Forestry and Fisheries
(MAFF), .

Masutomi, Y., Arakawa, M., Minoda, T., Yonekura, T., Shimada, T., 2015. Critical air temperature and
sensitivity of the incidence of chalky rice kernels for the rice cultivar "Sai-no-kagayaki". Agricultural
and Forest Meteorology 203, 11-16. doi:10.1016/j.agrformet.2014.11.016

Masutomi, Y., Takimoto, T., Manabe, T., Imai, Y., Tamura, M., Kobayashi, K., 2022. Breeding targets
for heat-tolerant rice varieties in Japan in a warming climate. Mitigation and Adaptation Strategies
for Global Change 28, 2. doi:10.1007/s11027-022-10027-4

Masutomi, Y., Takimoto, T., Shimamura, M., Manabe, T., Arakawa, M., Shibota, N., Ooto, A., Azuma,
S., Imai, Y., Tamura, M., 2019. Rice grain quality degradation and economic loss due to global
warming in Japan. Environmental Research Communications 1. doi:10.1088/2515-7620/ab52e7

Mavromatis, T., Boote, K.J., Jones, J.W., Irmak, A., Shinde, D., Hoogenboom, G., 2001. Developing
Genetic Coefficients for Crop Simulation Models with Data from Crop Performance Trials. Crop
Science 41, 40-51. doi:10.2135/cropsci2001.41140x

Mizobuchi, M., Takata, S., Sakata, M., Kameshima, M., Yoshida, Y., 2018. "Kouiku 76 gou", A new
extremerly early rice line with superior eating quality and grain appearance under high
temperatures during ripening. Bulletin of Kochi Agricultural Research Center 27, 55-64.

Morita, K., Matsushima, T., Yamaguchi, T., Saito, A., Furuhata, M., 2012. Evaluation of the Dense
Planting in Late Transplanting Aimed to Avoid High Temperatures During the Ripening Period in
Rice Cultivar Koshihikari. Japanese Journal of Crop Science 81, 349-356. doi:10.1626/jcs.81.349

Morita, S., 2008. Prospect for Developing Measures to Prevent High-Temperature Damage to Rice
Grain Ripening. Japanese Journal of Crop Science 77, 1-12. doi:10.1626/jcs.77.1

Nagato, K., Ebata, M., 1965. Effects of High Temperature during Ripening Period on the Development
and the Quality of Rice Kernels. Japanese journal of crop science 34, 59-66. doi:10.1626/jcs.34.59

NARO, Japan Soil inventry.

NARO, 2011. A new rice cultivar "Harumoni". Press relsease.

NARO, 2018. A new rice cultivar "Koisomeshi". Press release.

Neerja, S., Renu, K., 2020. Rice Grain Quality: Current Developments and Future Prospects. In: Farooq,
S. (Ed.), Recent Advances in Grain Crops Research.

Nishimori, M., Ishigooka, Y., Wakatsuki, H., Kuwagata, T., Hasegawa, T., Yoshida, H., Takimoto, T.,
Kondo, M., 2020. Rice quality analysis using the Crop Survey database of Japan for climate impact
assessment. Climate in Biosphere 20, 1-8. doi:10.2480/cib.J-20-054

Okada, M., Iizumi, T., Hayashi, Y., Yokozawa, M., 2011. Modeling the multiple effects of temperature
648 and radiation on rice quality. Environmental Research Letters 6, 034031. doi:10.1088/1748-
649 9326/6/3/034031

650 R Core Team, 2022. R: A Language and Environment for Statistical Computing. R Foundation for
651 Statistical Computing, Vienna, Austria.

652 Sanchez, B., Rasmussen, A., Porter, J.R., 2014. Temperatures and the growth and development of
653 maize and rice: a review. Glob Chang Biol 20, 408-417. doi:10.1111/gcb.12389

654 Takimoto, T., Masutomi, Y., Tamura, M., Nitta, Y., Tanaka, K., 2019. The effect of air temperature and
655 solar radiation on the occurrence of chalky rice grains in rice cultivars "Koshihikari" and
656 "Akitakomachi". Journal of Agricultural Meteorology 75, 203-210. doi:10.2480/agrmet.D-18-00039

657 Tamura, K., Kataoka, T., Nakanishi, A., Sato, H., Tamura, Y., Sakai, M., Yamaguchi, O., Wada, T.,
658 Tsubone, M., Tatara, I., Tokuda, S., Yoshida, K., Koga, J., Nakayama, M., Fujii, Y., Mitsukawa, M.,
659 Shimizu, Y., Hasegawa, W., Shiraishi, M., Nagayoshi, Y., Matsuura, S., Wakamatsu, K.-i., Sato, M.,
660 Sonoda, J., Takeuchi, Y., 2018. Standard Rice Varieties with High Temperature Tolerance at the
661 Ripening Stage in the Warm Region of Japan. Japanese Journal of Crop Science 87, 209-214.
662 doi:10.1626/jcs.87.209

663 Tanaka, K., Miyazaki, M., Uchikawa, O., Araki, M., 2010. Effects of the Nitrogen Nutrient Condition
664 and Nitrogen Application on Kernel Quality of Rice. Japanese Journal of Crop Science 79, 450-459.
665 doi:10.1626/jcs.79.450

666 Tsukaguchi, T., Iida, Y., 2008. Effects of Assimilate Supply and High Temperature during Grain-Filling
667 Period on the Occurrence of Various Types of Chalky Kernels in Rice Plants (Oryza sativa L.). Plant
668 Production Science 11, 203-210. doi:10.1626/pps.11.203

669 Usui, Y., Sakai, H., Tokida, T., Nakamura, H., Nakagawa, H., Hasegawa, T., 2014. Heat-tolerant rice
670 cultivars retain grain appearance quality under free-air CO_2 enrichment. Rice (N Y) 7, 6.
671 doi:10.1186/s12284-014-0006-5

672 Usui, Y., Sakai, H., Tokida, T., Nakamura, H., Nakagawa, H., Hasegawa, T., 2016. Rice grain yield and
673 quality responses to free-air CO_2 enrichment combined with soil and water warming. Global
674 Change Biology 22, 1256-1270. doi:10.1111/gcb.13128

675 Viechtbauer, W., 2010. Conducting Meta-Analyses in R with the metafor Package. Journal of
676 Statistical Software 36, 1 - 48. doi:10.18637/jss.v036.i03

677 Wada, T., Tsubone, M., Inoue, T., Ogata, T., Hamachi, Y., Matsue, Y., Oosato, K.F., Yasunaga, T.,
678 Kawamura, Y., Ishizuka, A., 2010. "Genkitsukushi", a new rice cultivar with tolerance to high
679 temperature during maturing period and high eating quality. Bulletin of Fukuoka Agricultural
680 Research Center 29, 1-9.

681 Wakamatsu, K.-i., Sasaki, O., Tanaka, A., 2009. Effects of the Amount of Insolation and Humidity
682 during the Ripening Period on the Grain Quality of Brown Rice in Warm Regions of Japan. Japanese
683 Journal of Crop Science 78, 476-482. doi:10.1626/jcs.78.476

684 Wakamatsu, K.-i., Sasaki, O., Uezono, I., Tanaka, A., 2007. Effects of High Air Temperature during the
685 Ripening Period on the Grain Quality of Rice in Warm Regions of Japan. Japanese Journal of Crop
686 Science 76, 71-78. doi:10.1626/jcs.76.71

687 Wakamatsu, K.-i., Sasaki, O., Uezono, I., Tanaka, A., 2008. Effect of the Amount of Nitrogen
688 Application on Occurrence of White-back Kernels during Ripening of Rice under High-temperature
689 Conditions. Japanese Journal of Crop Science 77, 424-433. doi:10.1626/jcs.77.424

690 Xie, X., Wu, T., Zhu, M., Jiang, G., Xu, Y., Wang, X., Pu, L., 2021. Comparison of random forest and
691 multiple linear regression models for estimation of soil extracellular enzyme activities in
692 agricultural reclaimed coastal saline land. Ecological Indicators 120, 106925.
693 doi:10.1016/j.ecolind.2020.106925

694 Yandell, B.S., 1997. Practical Data Analysis for Designed Experiments. Chapman and Hall/CRC, FL, USA.

695 Yang, L., Wang, Y., Dong, G., Gu, H., Huang, J., Zhu, J., Yang, H., Liu, G., Han, Y., 2007. The impact of
696 free-air CO_2 enrichment (FACE) and nitrogen supply on grain quality of rice. Field Crop Res 102,
697 128-140. doi:10.1016/j.fcr.2007.03.006

Ye, C., Li, X., Redoña, E., Ishimaru, T., Jagadish, K., 2021. Genetics and Breeding of Heat Tolerance in
Rice BT - Rice Improvement: Physiological, Molecular Breeding and Genetic Perspectives. In: Ali, J.,
Wani, S.H. (Eds.). Springer International Publishing, Cham, pp. 203-220.
Yoshida, H., Takehisa, K., Kojima, T., Ohno, H., Sasaki, K., Nakagawa, H., 2016. Modeling the effects of
N application on growth, yield and plant properties associated with the occurrence of chalky grains
of rice. Plant Production Science 19, 30-42. doi:10.1080/1343943X.2015.1128111
Yoshimoto, M., Fukuoka, M., Tsujimoto, Y., Matsui, T., Kobayasi, K., Saito, K., van Oort, P.A.J., Inusah,
B.I.Y., Vijayalakshmi, C., Vijayalakshmi, D., Weerakoon, W.M.W., Silva, L.C., Myint, T.T., Phyo, Z.C.,
Tian, X., Lur, H.-S., Yang, C.-M., Tarpley, L., Manigbas, N.L., Hasegawa, T., 2022. Monitoring canopy
micrometeorology in diverse climates to improve the prediction of heat-induced spikelet sterility
in rice under climate change. Agricultural and Forest Meteorology 316, 108860.
doi:10.1016/j.agrformet.2022.108860
Zhao, X., Fitzgerald, M., 2013. Climate Change: Implications for the Yield of Edible Rice. PLOS ONE 8,
e66218. doi:10.1371/journal.pone.0066218

712