A $MSTN^{Del273C}$ mutation with *FGF5* knockout sheep by CRISPR/Cas9 promotes skeletal muscle myofiber hyperplasia

Ming-Ming Chen[1, †], Yue Zhao[1, †], Xue-Ling Xu[1], Xiao-Sheng Zhang[2], Jin-Long Zhang[2], Su-Jun Wu[1], Zhi-Mei Liu[1], Yi-Ming Yuan[1], Xiao-Fei Guo[2], Shi-Yu Qi[1], Guang Yi[1], Shu-Qi Wang[1], Huang-Xiang Li[1], Ao-Wu Wu[1], Guo-Shi Liu[1], Kun Yu[1], Shoulong Deng[3], Hong-Bing Han[1], Feng-Hua Lv[1, *], Yan Li[4, *], Zheng-Xing Lian[1, *]

[1] Beijing Key Laboratory for Animal Genetic Improvement, National Engineering Laboratory for Animal Breeding, Key Laboratory of Animal Genetics and Breeding of the Ministry of Agriculture, College of Animal Science and Technology, China Agricultural University, Beijing 100193, China

[2] Institute of Animal Husbandry and Veterinary Medicine, Tianjin Academy of Agricultural Sciences, Tianjin 300381, China

[3] NHC Key Laboratory of Human Disease Comparative Medicine, Institute of Laboratory Animal Sciences, Chinese Academy of Medical Sciences and Comparative Medicine Center, Peking Union Medical College, Beijing, China

[4] Laboratory Animal Center of the Academy of Military Medical Sciences, Beijing 100071, China

† These authors contributed equally to this work.
* Correspondence: lianzhx@cau.edu.cn (Zheng-Xing Lian), scauly@cau.edu.cn (Yan Li); lvfenghua@cau.edu.cn (Feng-Hua Lv)

Abstract

Mutations in the well-known Myostatin (*MSTN*) produce a "double-muscle" phenotype, which makes it commercially invaluable for improving livestock meat production and providing high-quality protein for humans. However, mutations at different loci of the *MSTN* often produce a variety of different phenotypes. In the current study, we increased the delivery ratio of Cas9 mRNA to sgRNA from the traditional 1:2 to 1:10, which improves the efficiency of the homozygous mutation of biallelic gene. Here, a $MSTN^{Del273C}$ mutation with *FGF5* knockout sheep, in which the *MSTN* and *FGF5* dual-gene biallelic homozygous mutations were produced via the deletion of 3-base pairs of AGC in the third exon of *MSTN*, resulting in cysteine-depleted at amino acid position 273, and the *FGF5* double allele mutation led to inactivation of *FGF5* gene. The $MSTN^{Del273C}$ mutation with *FGF5* knockout sheep highlights a dominant "double-muscle" phenotype, which can be stably inherited. Both F0 and F1 generation mutants highlight the excellent trait of high-yield meat with a smaller cross-sectional area and higher number of muscle fibers per unit area. Mechanistically, the $MSTN^{Del273C}$ mutation with *FGF5* knockout mediated the activation of *FOSL1* via the MEK-ERK-FOSL1 axis. The activated *FOSL1* promotes skeletal muscle satellite cell proliferation and inhibits myogenic differentiation by inhibiting the expression of MyoD1, and resulting in smaller myotubes. In addition, activated ERK1/2 may inhibit the secondary fusion of myotubes by Ca^{2+}-dependent CaMKII activation pathway, leading to myoblasts fusion to form smaller myotubes.

Keywords: *MSTN*; *FGF5*; dual-gene biallelic mutation; *FOSL1*; myogenesis

1 Introduction

Myostatin (*MSTN*) has been well-known as a negative regulator of muscle growth and development. Its mutation produces a "double-muscle" phenotype, which shows its inestimable commercial value in improving meat production of livestock and poultry, and providing high-quality protein for humans (Fan *et al.*, 2022; Chen *et al.*, 2021b). Due to its role in promoting muscle atrophy and cachexia, *MSTN* has been recognized as a promising therapeutic target to offset the loss of muscle mass (Lee, 2021; Baig *et al.*, 2022; Wijaya *et al.*, 2022).

MSTN is highly conserved in mammals, and mutations in the *MSTN* gene, either artificially or naturally, will result in increased skeletal muscle weight and produce a "double-muscle" phenotype, which has been reported in many species, including cattle, sheep, and pigs, rabbits, and humans (Grisolia *et al.*, 2009; Dilger *et al.*, 2010; Kambadur *et al.*, 1997). However, mutations at different loci of the *MSTN* often produce variety of different phenotypes, and its molecular mechanism of skeletal muscle growth and development remains controversial (Hanset and Michaux, 1985; Grobet *et al.*, 1997; Wegner *et al.*, 2000; Kambadur *et al.*, 1997; Marchitelli *et al.*, 2003). More than 77 natural mutation sites of *MSTN* have been reported in various sheep breeds, most of these mutations were found to be located in the non-coding regions, and did not affect *MSTN* activity (Kijas *et al.*, 2007; Sjakste *et al.*, 2011; Han *et al.*, 2013; Dehnavi *et al.*, 2012). In addition to introns, it is still possible that mutations in regulatory regions and exons may not affect the sheep phenotypes (Pothuraju *et al.*, 2015; Kijas *et al.*, 2007; Boman and Vage, 2009; Boman *et al.*, 2009).

Fibroblast growth factor 5 (*FGF5*) belongs to the fibroblast growth factor (FGF) family and is a secretory signaling protein. *FGF5* played an inhibitory effect on mouse hair growth (Hebert *et al.*, 1994), and its natural mutation can lead to a significant increase in hair growth in angora mice (Sundberg *et al.*, 1997). Subsequent studies have also successively confirmed the inhibitory effect of *FGF5* on mammalian hair growth and is recognized to be a negative regulator of hair growth (Kehler *et al.*, 2007; Dierks *et al.*, 2013; Yoshizawa *et al.*, 2015; Legrand *et al.*, 2014; Higgins *et al.*, 2014).

81   In this study, to increase both meat and wool production, we first produced the

82 *MSTN* and *FGF5* dual-gene biallelic homozygous mutations sheep by the increased

83 delivery ratio of Cas9 mRNA to sgRNA targeting *MSTN* and *FGF5*. The *MSTN*^{Del273C}

delivery ratio of Cas9 mRNA to sgRNA targeting *MSTN* and *FGF5*. The $MSTN^{Del273C}$

mutation with *FGF5* knockout sheep highlights a dominant "double-muscle"

phenotype by decreasing the muscle fiber cross-sectional area and increasing the

number of muscle fibers per unit area. Then, we used the *MSTN* and *FGF5* dual-gene

biallelic homozygous mutations sheep to unravel the molecular mechanism of the

"double-muscle" phenotype and myofiber hyperplasia.

89 2 Materials and Methods

90 2.1 Production of Cas9 mRNA and sgRNA

The Cas9 and U6-sgRNA co-expression vector backbones pX330 were

purchased from Addgene (plasmid ID: 42230 and 48138). Sheep MSTN and FGF5

sgRNAs were designed using CRISPR Design Tool

(http://tools.genome-engineering.org). The MSTN sgRNA

(GACATCTTTGTAGGAGTACAGCAA) and FGF5 sgRNA

(AGGTTCCCCTTTCCGCACCT) were used in this study. Two complementary guide

sequence oligos were synthesized, annealed, and cloned into the pX330 backbone

vector to form the functional co-expression plasmids. T7 promoter was linked to the 5'

ends of the Cas9 coding region and MSTN/FGF5 sgRNA template by PCR

amplification from the pX330-MSTN/FGF5 plasmid constructed as described above.

Then, these PCR products, as transcription templates, were purified using E.Z.N.A.

Cycle Pure Kit (Omega Bio-Tek). MSTN sgRNA and FGF5 sgRNA were prepared by

in vitro transcription (IVT) using the MEGAshortscript T7 Kit (Life Technologies).

Cas9 mRNA was transcribed with the m7G(5')ppp(5') G cap on its 5' terminal and

poly (A) tail on its 3' terminal using the mMESSAGE mMACHINE T7 Ultra Kit

(Life Technologies). Both the Cas9 mRNA and the sgRNAs were purified using the

MEGAclear Kit (Life Technologies) and eluted in RNase-free water.

108 2.2 Microinjection and embryo transfer

The procedure for the efficient production of pronuclear embryos has been

described previously (Li *et al.*, 2016). Briefly, Cas9 mRNA (1000 ng/μL) and

sgRNAs (200 ng/μL) were mixed and injected into the transferable embryos in which

the zona pellucida was clear, cytoplasm was uniform, and pronucleus was visible

using a FemtoJet microinjector (Eppendorf). Following microinjection, three to five

embryos were transplanted into the oviduct of each recipient within 1 h after starting

the laparotomy operation. Pregnancy was confirmed by transabdominal ultrasound

scanning on the 60th day after embryo transfer.

2.3 Tissue sample collection and preparation

Gluteus medius and longissimus dorsi were harvested from WT and *MSTN*^Del273C

mutation with *FGF5* knockout (MF$^{-/-}$) sheep, and three WT sheep and four MF$^{-/-}$ F1

generation sheep (half-sib) were used for feeding and slaughter. All sheep were

female and are slaughtered at 12-month-old. All samples were immediately frozen in

liquid nitrogen and then stored at $-80°C$ until analysis. All sheep are raised by the

national feeding standard NT/T815-2004. All procedures performed for this study

were consistent with the National Research Council Guide for the Care and Use of

Laboratory Animals. All experimental animal protocols in this study were approved

and performed following the requirements of the Animal Care and Use Committee at

China Agricultural University (AW02012202-1-3). All surgeries were performed

under sodium pentobarbital anesthesia, and all efforts were made to minimize any

suffering experienced by the animals used in this study.

2.4 H&E staining and morphological analysis of muscle fibers

Fresh muscle tissue samples were fixed, dehydrated, embedded and frozen

sectioned, respectively. Next, they were sequentially stained with hematoxylin and

eosin, then dehydrated with gradient ethanol and transparentized with xylene, and

finally sealed with neutral resin to make tissue sections. The images from at least five

random fields were captured with an inverted microscope. The Image J software was

used to segment, count and calculate the area of each muscle fiber cell. The number of

fiber cell in a fixed area size was calculated to estimate the number of fiber cell per

unit area.

2.5 Cell isolation, culture, and transfection

Sheep skeletal muscle satellite cells were isolated and cultured as previously

described (Chen *et al.*, 2021a). In brief, the muscle tissues of the hind limbs from

3-month-old sheep fetuses were cut into small pieces, digested sequentially with 0.2%

collagenase type II (Gibco, Grand Island, NY) and 0.25% trypsin (Gbico, Grand

Island, NY). The cell suspension was successively filtered through 100, 200 and 400

mesh cell sieves. After this, the cells were resuspended in growth medium (GM)

containing DMEM/F12 (Gbico, Grand Island, NY) with 20% fetal bovine serum (FBS,

Gibco) and 1% penicillin-streptomycin liquid (Gbico, Grand Island, NY), and

cultured for 2-3 times with differential adhesion. To induce differentiation, the cells

were cultured to 70% confluence in GM, and followed by an exchange to

differentiation medium (DM) containing DMEM high glucose (Gbico, Grand Island,

NY) with 2% horse serum (HS, Gibco) and 1% penicillin-streptomycin. To produce

viral solution for over-expression of the target gene, it was subcloned into the XbaI

and BamHI sites of the lentiviral vector by seamless cloning, and the primer

sequences of gene cloning were listed in Table S1. HEK 293T cells were

co-transfected with the envelope plasmid pMD2.G, the packaging plasmid psPAX2

and the target plasmid at a mass ratio of 1:2:4 to produce the virus. The siRNA were

synthesized by Guangzhou RiboBio Co., Ltd, and the sequences were listed in Table

S2. Then, the cells were infected with packaged lentivirus or transfected with siRNA

using Lipofectamine 3000 (Invitrogen, USA) when they were cultured to 60%-70%

confluence.

2.6 Total RNA isolation and real-time quantitative PCR (RT- qPCR)

The total RNA of tissues and cells was isolated using TRIzol reagent (Sangon

Biotech, Shanghai, China) following the manufacturer's protocol. In short, after

tissues or cells were lysed, chloroform was added to separate the organic and

inorganic phases, followed by precipitation with isopropanol and ethanol in turn, and

finally, the RNA was dissolved in DEPC water. Then, the first strand cDNA was

prepared using PrimeScript II 1st Strand cDNA Synthesis Kit (Takara, Beijing, China).

qPCR was performed using 2× SYBR Green qPCR Mix (Low ROX) (Aidlab

Biotechnologies, Beijing, Chian) in a Stratagene Mx3000P (Agilent Technologies,

SUA). With GAPDH mRNA as endogenous control, the relative expression level of

genes was calculated by the $2^{-\Delta\Delta Ct}$ method. All primers used were listed in Table S3.

2.7 Western blot

Tissue or cell samples were lysed in RIPA buffer (Solarbio, Beijing, China)

supplemented with protease and phosphatase inhibitor cocktail (Beyotime, Beijing,

China) for total protein extraction. Then, equal amounts of tissue or cell lysate were

resolved by 10% SDS-PAGE and transferred onto PVDF membranes (Millipore,

USA). The membranes were blocked with 5% BSA for 1h, incubated with primary

antibody at 4℃ overnight, then incubated with secondary antibody for 1h before

detection. The fold change of protein was normalized to GAPDH for quantitative

analysis by ImageJ software. The antibodies information was listed in Table S4.

2.8 5-Ethynyl-2′-deoxyuridine (EdU) assay

At 24 h after transfection, sheep skeletal muscle satellite cells were incubated at

37℃ for 2 h in 96-well plates with 50 µM EdU (RiboBio, Guangzhou, China). Then,

fixed the cells in 4% paraformaldehyde for 30 min and neutralized using 2 mg/mL

glycine solution. The Apollo® staining solution which contains EdU was added and

incubated at room temperature for 30 min in the dark to label the DNA in the

synthesis stage, the nuclear was then counterstained with DAPI. The number of EdU

positive cells was counted from the images of five random fields obtained with an

inverted fluorescence microscope at a magnification of 100×. EdU labeling index was

expressed as the number of EdU-positive cell nuclei/total cell nuclei.

2.9 Cell counting kit-8 (CCK-8) and cell cycle detection

Skeletal muscle satellite cells were seeded in 96-well plates and cultured for

appropriate time according to different experimental treatments. Then, 10 µL CCK-8

solution was added to each well and incubated at 37°C in a 5% CO_2 incubator for 2 h,

and then the absorbance at 450 nm was measured with a microplate reader. The

cultured skeletal muscle satellite cells were digested with trypsin, centrifuged at 1000

g for 5 min to collect the cell pellet, washed once with ice-cold PBS, and then 1 mL of

ice-cold 70% ethanol was added to fix the cells overnight at 4°C. The next day, the

cells were washed with ice-cold PBS again, and the cells were incubated with 0.5 mL

PI staining solution at 37°C for 30 min and collected by flow cytometry at low speed.

2.10 Immunofluorescence staining

Sheep skeletal muscle cells were fixed in 4% paraformaldehyde for 30 min,

permeabilized in 0.1% Triton X-100 for 20 min and blocked with 5% normal goat

serum for 30 min at room temperature, and then incubated with primary antibody at 4℃

overnight. Next, the fluorescent secondary antibody was added and incubated at 37℃

for 1 h in the dark, and the nuclear was then counterstained with DAPI. The

immunofluorescence images from five random fields were captured with an inverted

fluorescence microscope.

2.11 Chromatin Immunoprecipitation (ChIP)

The cells were fixed with 1% formaldehyde, then the cells were collected with a

cell scraper and resuspended in cell lysis buffer (10 mM HEPES, 0.5% NP-40, 1.5

mM MgCl2, 10 mM KCl, pH 7.9) containing protease inhibitor cocktail (Beyotime,

Beijing, China) and incubated on ice to release the cytoplasm. Next, cell pellets were

collected and resuspended in nuclear lysis buffer (50 mM Tris, 10 mM EDTA, 0.3%

SDS, pH 8.0) containing protease inhibitor cocktail. After the DNA was fragmented

by ultrasonication, the supernatant was collected. The samples were diluted ChIP

dilution buffer (0.01% SDS, 1.1% Triton X- 100, 1.2 mM EDTA, 16.7 mM Tris-HCl

pH 8.0, 167 mM NaCl), then 5 μg primary antibody was added and incubated

overnight at 4°C with rotation. The next day, protein A/G magnetic beads were added

to each sample and incubated at 4°C with rotation for 2h. Then, the magnetic beads

were respectively washed once with low-salt wash buffer (0.1% SDS, 1% Triton

X-100, 2 mM EDTA, 20 mM Tris-HCl pH 8.0, 150 mM NaCl), high-salt wash buffer

(0.1% SDS, 1% Triton X-100, 2 mM EDTA, 500mM NaCl, 20 mM Tris-HCl pH 8.0),

LiCl wash buffer (0.25 M LiCl, 1% NP-40, 1% sodium deoxycholate, 1 mM EDTA,

10 mM Tris-HCl pH 8.0), and TE buffer (10 mM Tris-HCl pH 8.0, 1 mM EDTA) at

4°C. Next, ChIP elution buffer (1% SDS, 100 mM NaHCO$_3$) containing proteinase K

was added to each sample, then incubated at 62°C overnight, and the DNA was finally

purified by a purification column.

2.12 RNA-seq

The rRNA was removed from each total RNA sample of the gluteus medius to

construct a strand-specific transcriptome sequencing library, and the Illumina

Novaseq 6000 sequencing platform was used to perform high-throughput sequencing

with a paired-end read length of 150 bp. Raw data were transformed into clean reads

by removing reads containing adapter, ploy-N and low-quality reads from raw data.

At the same time, Q20, Q30, GC-content, and sequence duplication levels of the clean

data were calculated. The genome index was constructed using Hisat2 software and

the clean reads were mapped to the sheep reference genome (*Oar Rambouillet v1.0*),

the featureCounts software was used for expression quantification, and DESeq2

software was used for differential expression analysis based on *P*-value < 0.05 and |

log2 Fold Change | > 1.

3 Results

3.1 Elevated molar ratio of Cas9/sgRNA can efficiently generate biallelic

homozygous mutant sheep

The sgRNAs for targeting were designed in the third exon of the MSTN and

FGF5 genes, respectively (Figure 1A, B). Both MSTN and FGF5 PCR products could

be cleaved by T7E1 and the fragment sizes were also as expected, and grayscale

analysis showed that the editing efficiency was 14.6% and 11.4%, respectively

(Figure 1C), which indicates that the designed sgRNAs can achieve more efficient

gene targeting. The microinjection was performed according to the molar ratio of

Cas9 mRNA:sgRNAs (1:2, 1:10, and 1:15), respectively. The number of embryos

injected, recipients of nuclear transfer, pregnancy, and alive lambs per group were

listed in Table S5. The subsequent gene mutation detection showed that a total of 3

lambs were mutated in the MSTN and FGF5 genes at a Cas9 mRNA:sgRNAs

injection molar ratio of 1:2, with a gene editing mutation rate of 14.3% (4/28).

However, all 3 lambs were chimeric, that is, there were both mutant and wild-type after editing, and the biallelic mutation rate was 0% (Table S5 ,Figure 1D). Increasing the injection molar ratio of Cas9 mRNA:sgRNAs to 1:10 resulted in mutations in MSTN and FGF5 genes of two lambs with a gene editing mutation rate of 18.2 % (4/22), which contributed to a significant ($P<0.05$) increase in the biallelic mutation rate (Table S5, Figure 1D). While the injection molar ratio of Cas9 mRNA: sgRNAs was continuously increased to 1:15, one lamb had a mutation, which was a biallelic mutation of *MSTN* gene, and the gene editing mutation rate was 7.14% (1/14) (Table S5, Figure 1D). These results indicate that increasing the delivery molar ratio of Cas9 mRNA to sgRNA from 1:2 to 1:10 can greatly improve the efficiency of biallelic mutation in sheep.

3.2 The *MSTN*^{Del273C} mutation with *FGF5* knockout sheep highlights a dominant "double-muscle" phenotype and muscle fiber hyperplasia

Among gene-edited sheep, a sheep with biallelic deletion of *MSTN* and biallelic mutation of *FGF5* aroused our great interest. Specifically, gene editing caused a deletion of 3-base pairs of AGC in the third exon of *MSTN* (Figure 1E, F), resulting in the deletion of cysteine at amino acid position 273 (*MSTN*^{Del273C}) (Figure 1 E, F), which is highlighted by the "double-muscle" phenotype (Figure 2A, 2B). At the same time, a biallelic mutation in *FGF5* caused the knockout of *FGF5* gene and increased the density and length of hairs (Zhang *et al.*, 2020). Compared to WT sheep, the fiber cell number per unit area of gluteus medius and longissimus dorsi in MF$^{-/-}$ sheep was significantly ($P<0.01$) increased (Figure 2C, D), and the cross-sectional area were smaller (Figure 2C, E-F). Similarly, the cross-sectional area of gluteus medius muscle fibers in the offspring generation MF$^{+/-}$ sheep was also smaller (Figure 2G-H), and the number of muscle fiber cells per unit area was significantly increased ($P<0.0001$) (Figure 2G, I); the percentage of smaller muscle fiber area in MF$^{+/-}$ sheep was significantly increased ($P<0.05$) (Figure 2G, J), these results was consistent with that in MF$^{-/-}$ sheep. Interestingly, the FGF5 knockout alone had no significant ($P>0.05$) effect on muscle fiber size (Figure S1A-D). Although the mRNA expression levels of *MSTN* and *FGF5* were significantly ($P<0.05$) reduced in MF$^{+/-}$ sheep during early embryonic development (3-month-old) (Figure S2A), there was no significant

differences in *MSTN* mRNA and protein expression levels at 12-month-old after birth (Figure S2B-D). In addition, there was no significant differenc in the proportion of centrally nucleated myofibres (P>0.05) (Figure S2E), nor were there aberrant expression of some genes related to muscular dystrophy and muscle atrophy (Figure S2F). These results indicate that *MSTN*^{Del273C} mutation with *FGF5* knockout produces muscle fiber hyperplasia instead of muscular dystrophy or muscle atrophy.

Further, although the muscle weight of different parts in WT and MF$^{+/-}$ sheep has no significant difference (Table S6), the proportion of hind leg meat was significantly (P<0.05) increased by 21.2% (Table S7), and the proportion of gluteus medius in the carcass of MF$^{+/-}$ sheep was significantly (P<0.01) increased by 26.3% compared to WT sheep (Figure 2K). In addition, there were no significant (P>0.05) differences in pH, color, drip loss, cooking loss, shearing force, and amino acid content of the longissimus dorsi between WT and MF$^{+/-}$ sheep (Table S8-10). All these results demonstrated that the *MSTN*^{Del273C} mutation with *FGF5* knockout sheep had well-developed hip muscles with smaller muscle fibers, which do not affect meat quality, and this phenotype may be dominated by *MSTN* gene.

3.3 The *MSTN*^{Del273C} mutation with *FGF5* knockout promotes skeletal muscle satellite cells proliferation and inhibits myogenic differentiation

The proliferation and differentiation of skeletal muscle satellite cells is a key step in muscle formation and development. The CCK-8 and EdU cell proliferation experiments showed that the proliferative rate of MF$^{+/-}$ cells were highly significantly (P<0.01) elevated (Figure 3A) with a significant (P<0.05) increase in the rate of EdU-positive cells (Figure 3B, C) compared to WT cells. In addition, cell cycle detection showed a significant (P<0.01) reduce in the proportion of G1 phase and a significant increase (P<0.05) in the proportion of S phase in MF$^{+/-}$ cells (Figure 3D, E). Meanwhile, the mRNA expression levels of the cell cycle marker genes CyclinB1, CDK4, Cyclin A1, Cyclin E1, and CDK2 were significantly increased (P<0.05) (Figure 3F). These results suggest that the *MSTN*^{Del273C} mutation with *FGF5* knockout may promote cell proliferation by accelerating the cell cycle from G0/G1 phase to S phase.

The mRNA level of MyHC and the protein levels of MyoD1, MyoG, and MyHC

(Figure 3G-I) were dramatically decreased ($P<0.05$) after induced differentiation 2

days in MF$^{+/-}$ cells, suggesting that the $MSTN^{Del273C}$ mutation with $FGF5$ inhibit

myogenic differentiation. Meanwhile, the immunofluorescence staining of MyoG and

MyHC in myotubes showed that myotube fusion index (Figure 3J, K), number of

myotubes (Figure 3J, L), and number of nuclei per myotube (Figure 3J, M) were all

highly significantly ($P<0.01$) reduced after inducing differentiation for 2 days of

MF$^{+/-}$ cells compared to WT cells, as was the myotube diameter at the maximum

measured (Figure 3J, N). In addtion, the differentiation capacity and fusion ability of

MF$^{+/-}$ cells were consistently significantly lower than WT cells during the ongoing

differentiation process, as was the diameter of fused myotubes (Figure S3A-J). The

reduced expression of myogenic differentiation markers further confirmed that the

$MSTN^{Del273C}$ mutation with $FGF5$ knockout consistently inhibits myogenic

differentiation of skeletal muscle satellite cells (Figure S3K-N). Taken together, our

results elucidated that the $MSTN^{Del273C}$ mutation with $FGF5$ knockout promotes

proliferation and inhibits myogenic differentiation of skeletal muscle satellite cells,

and induces a smaller myotube diameter of myotubes, which may explain the muscle

fiber hyperplasia phenotype and the decreased cross-sectional area of muscle fibers in

MF$^{-/-}$ and MF$^{+/-}$ sheep.

335   3.4 The $MSTN^{Del273C}$ mutation with $FGF5$ knockout contribute to muscle

336   phenotype via MEK-ERK-FOSL1 axis

To elucidate the potential mechanism of the $MSTN^{Del273C}$ mutation with $FGF5$

knockout result in smaller muscle fiber cross-sectional area and myotube diameter,

the RNA-seq was performed in gluteus medius. GO and KEGG enrichment analysis

indicating that differentially expressed genes (DEGs) were significantly closely

related to cell proliferation, myogenic differentiation, and muscle development; and

significantly enriched in MAPK signaling pathway (Figure 4A-B). Among DEGs,

FOSL1 has aroused our interest. Compared to WT sheep, the FSOL1 mRNA level

was significantly ($P<0.05$) lower in both gluteus medius and longissimus dorsi in

MF$^{+/-}$ sheep (Figure 4C), but it was significantly ($P<0.05$) elevated in MF$^{+/-}$ cells at

GM and DM2 (Figure 4D). More strikingly, FOSL1 mRNA level was strongly

(P<0.01) decreased after induced differentiation (Figure 5E), and its expression

diminished continuously with the differentiation progress. These results suggested

that FOSL1 may play a crucial role in the proliferation and myogenic differentiation

of skeletal muscle satellite cells.

FOSL1 is a member of the AP-1 family, and another member of this family,

c-Fos, inhibits myogenesis and MyoD1 expression by directly binding to the MyoD1

promoter region. Therefore, we speculated that FOSL1 might have similar functions

to c-Fos. Subsequent protein-protein interaction (PPI) analysis further suggested that

there was a potential interaction between FOSL1 and MyoD1 (Figure 5F). In addition,

the c-Fos mRNA level was highly significantly (P<0.01) reduced in MF$^{+/-}$ myoblasts

compared to WT cells, whereas the MyoD1 mRNA level was dramatically (P<0.01)

increased (Figure 5G). We futher found that two bZIP recognition sites in the MyoD1

promoter region had the most significant binding potential to FOSL1 (Figure 5H-J).

Subsequently, ChIP-qPCR confirmed that FOSL1 directly binds to these two bZIP

recognition sites in the MyoD1 promoter region (Figure 5K-L). The dual luciferase

report experiment confirmed that transcription factor FOSL1 can significantly

(P<0.05) inhibit MyoD1 promoter activity (Figure 4M). These results indicated that

FOSL1 plays an important role in the transcriptional regulation of MyoD1.

As mentioned above, FOSL1 may be involved in the proliferation and myogenic

differentiation of skeletal muscle satellite cells. Here, the protein levels of FOSL1 and

c-Fos were significantly (P<0.05) reduced in MF$^{+/-}$ cells at GM compared to WT cells

(Figure 4N-O), and accompanied by a significant (P<0.05) increase in p-FOSL1

protein levels (Figure 4N-O), whereas FOSL1 protein levels were significantly

(P<0.05) diminished and c-Fos protein levels were highly significantly (P<0.01)

elevated in MF$^{+/-}$ cells after induced differentiation (Figure 4P-Q), these results

further support the key role of FOSL1 on myogenesis. As demonstrated previously,

enrichment analysis significantly enriched the MAPK signaling pathway. Compared

to WT cells, ERK1/2 protein level was extremely significantly (P<0.01) decreased,

and accompanied by a significant (P<0.05) increase in p-ERK1/2 protein levels

(Figure 4N-O). After induced differentiation, although both MEK1/2 and ERK1/2

protein levels were dramatically (P<0.01) inhibited (Figure 4P-Q).

Considering the possible serum regulation of MSTN, we examined the effects of
MSTN mutations on its receptors and downstream target genes, and observed that
both MSTN receptors were significantly up-regulated (Figure S4A), whereas the
expression of downstream Smand and Jun families was also inhibited to a varying
degree (Figure S4B-C). Furthermore, serum from $MF^{+/-}$ sheep promoted the
proliferation of skeletal muscle satellite cells (Figure S4D). $MSTN^{Del273C}$ mutation
with *FGF5* knockout promoted FOSL1 expression using WT sheep serum (Figure
S4E), which was similar to the results of FBS culture and HS induction. The serum
from $MF^{+/-}$ sheep strongly stimulated FOSL1 expression and the inhibition of MyoD1
(Figure S4F). These results suggest that serum regulation cannot be ignored after
$MSTN^{Del273C}$ mutation with *FGF5* knockout.

In summary, $MSTN^{Del273C}$ mutation with *FGF5* knockout may regulate the
expression and activity of FOSL1 via the MEK1/2-ERK1/2-FOSL1 axis to affect the
proliferation and myogenic differentiation of skeletal muscle satellite cells, and
further contribute to muscle phenotype.

3.5 FOSL1 expression and activity control the proliferation and myogenic
differentiation of skeletal muscle satellite cells

To investigate the role of FOSL1 on the proliferation and myogenic
differentiation of skeletal muscle satellite cells, we successfully constructed *FOSL1*
gain-of-function model (Figure 5A). We found that overexpression of *FOSL1*
significantly ($P<0.05$) promotes cell proliferation (Figure 5B-E), suppressed c-Fos
mRNA level (Figure 5F) ($P<0.05$), and inhibited MyoD1 mRNA ($P<0.01$) and protein
($P<0.05$) levels (Figure 5F-H). These results are consistent with what we observed in
$MF^{+/-}$ cells at GM, suggesting a potential inhibitory effect of FOSL1 protein on
MyoD1. In addition, the protein levels of MyoD1, MyoG and MyHC were all
significantly decreased ($P<0.05$), proving that the myogenic differentiation was
inhibited after FOSL1 was overexpressed (Figure 5I-J). Subsequently,
immunofluorescence staining further confirmed the significant ($P<0.05$) inhibitory
effect on cell differentiation (Figure 5K-L). Also, the number of myotubes, the
number of nuclei per myotube, and the myotube diameter were all significantly
decreased ($P<0.05$) (Figure 5K, M-O). Complementarily, we also constructed *FOSL1*

loss-of-function model (Figure 6A). In contrast to what observed in FOSL1
overexpression, interference with FOSL1 inhibited skeletal muscle satellite cell
proliferation (Figure 6B-E), elevated c-Fos and MyoD1 mRNA levels (Figure 6F) and
significantly ($P<0.05$) increased MyoD1 protein level (Figure 6G-H). With myogenic
differentiation markers were significantly up-regulated (Figure 6I-J), the myotube
fusion index, the number of myotubes, the number of nuclei per myotube, and the
myotube diameter were all significantly increased (Figure 6K-O). These results
further demonstrated that elevated FOSL1 level inhibits myogenic differentiation and
produces smaller myotubes.

To further ascertain this insight, the tert-butylhydroquinone (TBHQ), which can
strongly activate ERK1/2 and increase p-ERK1/2 protein expression level, was used
to activate ERK1/2 and act as an indirect activator of FOSL1. As expected, the
addition of 20 μM TBHQ significantly ($P<0.01$) inhibited the myogenic
differentiation of skeletal muscle satellite cells (Figure 7A-B). And, the number of
myotubes was significantly increased ($P<0.05$) (Figure 7A, C), while the number of
nuclei per myotube was significantly ($P<0.05$) decreased and produced a smaller
myotube diameter ($P<0.05$) (Figure 7A, D-E). In addition, PD98509, which can
inhibit ERK1/2 and dose-dependent reduce p-ERK1/2 protein expression level, was
used for complementary experiments. We observed opposite results to TBHQ
treatment, including increase in myotube fusion index, the number of nuclei per
myotube, and the myotube diameter (Figure 7F-J). Taken together, our results shed
light that the $MSTN^{Del273C}$ mutation with $FGF5$ knockout mediated the activation of
FOSL1 via MEK-ERK-FOSL1 axis, further promotes skeletal muscle satellite cell
proliferation, and inhibits myogenic differentiation by inhibiting the transcription of
MyoD1, and resulting in smaller myotubes.

3.6 The $MSTN^{Del273C}$ mutation with $FGF5$ knockout inhibit calcium-dependent
transcription signal pathway to regulate secondary fusion of myotubes

As mentioned previously, DEGs identified by RNA-seq were significantly
enriched in biological processes such as muscle contraction and cardiac muscle
contraction. In muscle contraction, calcium ions (Ca^{2+}) play a crucial role, triggering
the process of muscle contraction and relaxation by binding to proteins in muscle

fibers. Here, the calcium-calmodulin dependent protein kinase II (CaMKII) α/δ

protein level was significantly ($P<0.05$) reduced in $MF^{+/-}$ cells compared to WT cells

(Figure 8A-B), and the intracellular Ca^{2+} concentration was also significantly ($P<0.05$)

reduced (Figure 8C-D). The high-throughput Ca^{2+} channel RYR is responsible for

rapid and massive release Ca^{2+} from the endoplasmic reticulum into cytoplasm. We

observed a significant ($P<0.05$) decrease in RYR1 and/or RYR3 mRNA levels in

$MF^{+/-}$ sheep skeletal muscle satellite cells and myotubes cells (Figure 8E-F). In

addition, intracellular Ca^{2+} concentration correlated with myoblasts fusion, whereas

the mRNA levels of MYMK and MYMX, which control myoblasts fusion, were

significantly ($P<0.01$) reduced in $MF^{+/-}$ cells (Figure 8E-F). These results suggest that

the decrease Ca^{2+} levels and inhibition of myoblasts fusion genes may be potential

triggers for the decrease of myotube diameter and myofiber cross-sectional area in

$MF^{+/-}$ sheep.

In a word, our results shed light that the $MSTN^{Del273C}$ mutation with *FGF5*

knockout mediated the activation of FOSL1 via MEK-ERK-FOSL1 axis (Figure 9).

The activated FOSL1 promotes skeletal muscle satellite cell proliferation and inhibits

myogenic differentiation by inhibiting the expression of MyoD1, and resulting in

fusion to form smaller myotubes (Figure 9). In addition, activated ERK1/2 may

inhibit the secondary fusion of myotubes by Ca^{2+}-dependent CaMKII activation

pathway, leading to myoblasts fusion to form smaller myotubes (Figure 9).

4 Discussion

4.1 Optimized Cas9 mRNA and sgRNA delivery ratio improves the efficiency of

dual-gene biallelic homozygous mutations

The strategy for producing gene knockout animals by CRISPR/Cas9 gene editing

system is usually to introduce the Cas9 mRNA and the sgRNA of the target gene into

their prokaryotic embryos by microinjection. However, this "one-step" method often

results in a "mosaic" of gene-edited offspring (Wan *et al.*, 2015). Such chimeric

mutants have now been reported in gene knockout mice (Wang *et al.*, 2013), rats (Bao

et al., 2015), monkeys (Niu *et al.*, 2014), pigs (Hai *et al.*, 2014), sheep (Hongbing

HAN, 2014), goats (Wang *et al.*, 2015), rabbits (Lv *et al.*, 2016), and humans (Wang

and Yang, 2019) prepared by a "one-step" method using the CRISPR/Cas9 system.

For studies involved in genetic phenotypes, chimeric gene knockout animals require

further cross-breeding to obtain animals with a complete knockout of the target gene.

Once required to generate multiple gene knockout animals, this time-consuming and

laborious operation will become extremely difficult. Although many studies have

been devoted to eliminating this widespread chimeric mutation (Sato *et al.*, 2015;

Sung *et al.*, 2014; Kotani *et al.*, 2015; Chen *et al.*, 2015; Zhou *et al.*, 2014; Tu *et al.*,

2017; Wang *et al.*, 2015), however, these optimizations did not bring about a

significant improvement in the production efficiency of biallelic knockout animals.

Here, we increased the delivery ratio of Cas9 mRNA to sgRNA from 1:2 to 1:10,

which improve the efficiency of the homozygous mutation of the biallelic gene. This

unprecedented optimization method not only improved the overall gene knockout

efficiency, but also the obtained gene-edited offspring were all dual-gene biallelic

mutation. However, it is necessary to point out that although there are statistical

differences, due to the limited number of sheep we actually produced and used for

evaluation, the strategy to improve the efficiency of the homozygous mutation of

biallelic gene by increasing the Cas9 and mRNA delivery ratio needs to be further

comfirm in future studies.

4.2 Phenotypes produced by *MSTN* mutations are mutation site-dependent

As mentioned previously, although *MSTN* mutations have been found to produce

a "double-muscle" phenotype in multiple species, the microscopic phenotypes are

different, and this difference is closely related to the mutation site and species types.

In mice, the number of skeletal muscle fibers with *MSTN* gene knockout significantly

increased by 86% (McPherron *et al.*, 1997). A missense mutant *MSTN* only increased

the number of mouse muscle fibers, while dominant negative *MSTN* resulted in

increased muscle fiber cross-sectional area in mice, but not the number of muscle

fibers (Nishi *et al.*, 2002; Zhu *et al.*, 2000). In addition, the use of *MSTN* neutralizing

antibody on adult rats also resulted in an increased muscle fiber cross-sectional area

(Haidet *et al.*, 2008). In cattle, natural MSTN mutant Belgian Blue cattle had an

increased number of muscle fibers and reduced muscle fiber diameter (Wegner *et al.*,

2000). The muscle fiber cross-sectional area of longissimus dorsi and gluteus medius

in sheep was significantly increased after a 4bp deletion of the first exon of *MSTN* (Zhiliang *et al.*, 2004). In pigs, both the *MSTN* gene-edited Meishan and Hubei pigs showed a phenotype with increased muscle fiber density (Qian *et al.*, 2015; Xu *et al.*, 2013). Here, we prepared $MSTN^{Del273C}$ mutation with *FGF5* knockout sheep with 3-base pairs of AGC in the third exon of *MSTN*, which caused the deletion of cysteine at amino acid position 273. Its macroscopic phenotype is similar to that of the MSTN-edited sheep with the first exon knocked out 4-base pairs. Both of them showed an abnormally developed "double-muscle" phenotype of hip muscle, but the $MSTN^{Del273C}$ mutation with *FGF5* knockout sheep highlights a muscle fiber hyperplasia phenotype.

4.3 FOSL1 recognizes and binds to the MyoD1 promoter and inhibits its expression

As previously described, AP-1 family members play key roles in skeletal muscle cell proliferation, differentiation, and muscle development. In this study, AP-1 family member *FOSL1* was significantly reduced in $MF^{+/-}$ sheep, and its expression were drastically reduced during myogenic differentiation, which was consistent with the decrease of *FOSL1* expression during C2C12 differentiation (Tobin *et al.*, 2016). In addition, FOSL2, another AP-1 family member, can also inhibit myoblast differentiation (Alli *et al.*, 2013a), which may support the inhibitory effect of FOSL1 on myogenic differentiation. Therefore, *FOSL1* was recognized as a potential gatekeeper. It has been shown that *FOSL1* heterodimerizes with other transcription factors, such as the members of the bZIP family, and these dimers are either disabling the transcriptional activator complex or saving the interacting proteins from degradation in proteasomes (Sobolev *et al.*, 2022). Moreover, c-Fos, a member of the AP-1 family, has been shown to inhibit MyoD1 expression and myogenesis by directly binding to the MyoD1 promoter region (Li *et al.*, 1992). Therefore, we speculate that *FOSL1* may have similar functions to c-Fos. PPI analysis suggested a potential interaction between *FOSL1* and MyoD1. Subsequently, we confirmed that FOSL1 directly binds to two bZIP recognition sites in the MyoD1 promoter region, and inhibits MyoD1 promoter activity. Meanwhile, the overexpression and interference of FOSL1 further confirmed the inhibitory effect of FOSL1 on the

expression of MyoD1. In a word, these results fully support our hypothesis that

FOSL1 binds the MyoD1 promoter and inhibits its expression.

4.4 The *MSTN*^{Del273C} mutation with *FGF5* knockout contribute to muscle

phenotype via MEK-ERK-FOSL1 axis

The nonclassical pathway of MSTN involves PI3K/Akt/mTOR signaling

pathway and MAPK signaling pathway, which mainly includes ERKs, JNKs and p38

MAPK (Huang *et al.*, 2007; Gui *et al.*, 2012). All of those pathways are involved in

the signal transduction pathway of MSTN and mediate the transcription of MRFs

(Myogenin, Myf5, MyoD), MuRF-1 and Atrogin-1, to regulate myogenic

differentiation and skeletal muscle quality (Chen *et al.*, 2021b). *MSTN* induces muscle

fiber hypertrophy prior to satellite cell activation (Wang and McPherron, 2012) and

inhibits IGF-I-induced increase in myotube diameter through Akt signaling pathway

(Morissette *et al.*, 2009). In our study, the MSTN^{Del273C} mutation with FGF5 knockout

resulted in the inhibition of myogenic differentiation of skeletal muscle satellite cells,

and the number of myotubes and the myotube size were significantly reduced.

As previously described, the DEGs of gluteus medius RNA-seq were

significantly enriched in the MAPK signaling pathway. In fact, the MAPK signaling

pathway has been proven to be closely related to muscle development and myoblast

differentiation (Xie *et al.*, 2018; Segales *et al.*, 2016). For example, ERK1/2 promotes

myoblast proliferation in response to various growth factors (Campbell *et al.*, 1995),

inhibits signaling pathways that activate ERK1/2, or isolates ERK1/2 in the cytoplasm,

leading to cell cycle exit and cell differentiation (Jones *et al.*, 2001; Michailovici *et al.*,

2014). The RXR activity in myoblasts promotes myogenesis by regulating MyoD

expression and acting as a MyoG cofactor (Zhu *et al.*, 2009). Inhibition of MEK1/2

activates satellite cell differentiation in primary muscle fibers (Alli *et al.*, 2013b), and

also induces myogenic differentiation and excessive fusion (Eigler *et al.*, 2021). A

recent study on glioma showed that FOSL1 can be activated by the

Ras-MEK1/2-ERK1/2 axis in MAPK signaling pathway (Marques *et al.*, 2021).

Similarly, the activated MEK1/2-ERK1/2 axis in aged skeletal muscle also activates

FOSL1 and increases the abundance of FOSL1 and the trans-activation capacity of the

Fos-Jun heterodimer (Mathes *et al.*, 2021). In our study, the *MSTN*^{Del273C} mutation

with *FGF5* knockout regulates FOSL1 expression and activity through MEK1/2-ERK1/2-FOSL1 axis and activated FOSL1 further inhibits myogenic differentiation of skeletal muscle satellite cells, resulting in smaller myotube diameter. However, despite the high expression of p-FOSL1 in MF$^{+/-}$ myoblasts, it did not significantly inhibit the transcription of MyoD1, which may be related to a dramatic enhance in c-Fos, or there might be other parallel signaling pathways regulating MyoD1 after *MSTN*^{Del273C} mutation with *FGF5* knockout.

Furthermore, it has been demonstrated that the inhibition of MEK1/2 using MEK1/2-specific inhibitor PD184352 can significantly down-regulate FOSL1 expression (Mathes *et al.*, 2021). In our study, the indirect activation of FOSL1 by TBHQ can inhibit the myogenic differentiation of sheep skeletal muscle satellite cells, leading to reduced myoblast fusion capacity and smaller myotube diameter. In contrast, inhibition of the MEK1/2 pathway by PD98059 to suppress FOSL1 activity produced the opposite effect. Taken together, these results shed light on the potential mechanisms by which *MSTN*^{Del273C} mutation with *FGF5* knockout leads to increased myofiber numbers and decreased fiber cross-sectional area.

4.5 The *MSTN*^{Del273C} mutation with *FGF5* knockout may affect myoblasts fusion through ERK-mediated calcium dependent transcriptional signaling pathway

Ca^{2+} is recognized as a regulator of mammalian muscle fusion (Eigler *et al.*, 2021; Constantin *et al.*, 1996). The transient depletion of Ca^{2+} in the endoplasmic reticulum is associated with myoblast differentiation and fusion (Jones *et al.*, 2001). However, the signaling cascade leading to Ca^{2+}-mediated myoblast fusion remains unclear. Intracellular Ca^{2+} level is regulated by various Ca^{2+} voltage-gated channels, including but not limited to ryanodine receptors (RYR), which is the main Ca^{2+} release channel of the sarcoplasmic reticulum. CaMKII is a member of the calcium/calmodulin-dependent serine/threonine kinase family. CaMKII δ, CaMKII γ, and CaMKII β are the main isoforms expressed in skeletal muscle (Bayer *et al.*, 1996). Recent studies have shown that elevated intracellular Ca^{2+} level is crucial for myoblasts fusion and that Ca^{2+} signaling in newly formed myotubes occurs prior to the rapid growth stage of myotubes, indicating that Ca^{2+} released from the endoplasmic reticulum in early myotubes may promote secondary fusion of myoblasts

and myotube expansion (Eigler *et al.*, 2021). Further studies showed that
Ca^{2+}-dependent activation of CaMKII is essential for myotubes expansion and may
mediate myoblast-myotube fusion by regulating MYMK and Rac1, but it is not
necessary for myoblast-myoblast fusion (Eigler *et al.*, 2021).

In our study, the *MSTN*^Del273C mutation with *FGF5* knockout resulted in a
significant reduction of Ca^{2+} level and CaMKII α/δ protein level, and led to a decrease
in myotube fusion capacity. Therefore, our results support that the increased
p-ERK1/2 level promotes cell proliferation and inhibits myogenic differentiation in
MF$^{+/-}$ sheep skeletal muscle satellite cells. Meanwhile, activated ERK1/2 further
inhibited RYR activity by suppressing the phosphorylation of RXR, thereby reducing
the release of endoplasmic reticulum Ca^{2+} and potentially inhibiting the secondary
fusion of myotubes by Ca^{2+}-dependent CaMKII activation pathway, and further
mediating myofiber hyperplasia.

607 5 Conclusion

In this study, we found that increasing the delivery ratio of Cas9 mRNA to
sgRNA can improve the efficiency of the homozygous mutation of the biallelic gene.
Based on this, we generated a *MSTN*^Del273C mutation with *FGF5* knockout sheep, a
dual-gene biallelic homozygous mutant, which highlights a dominant "double-muscle"
phenotype. Both F0 and F1 generation mutants highlight the excellent trait of
high-yield meat and the more number of muscle fibers per unit area. Our results
suggested the *MSTN*^Del273C mutation with *FGF5* knockout mediated the activation of
FOSL1 via MEK-ERK-FOSL1 axis, further promotes skeletal muscle satellite cell
proliferation, and inhibits myogenic differentiation by inhibiting the expression of
MyoD1, and resulting in smaller myotubes. In addition, activated ERK1/2 may inhibit
the secondary fusion of myotubes by Ca^{2+}-dependent CaMKII activation pathway,
leading to myoblasts fusion to form smaller myotubes. This supports the myofiber
hyperplasia that more number of muscle fibers and smaller cross sectional area,
caused by the MSTN^Del273C mutation with FGF5 knockout.

Data availability statement

The raw sequence data reported in this paper have been deposited in the Genome Sequence Archive (Genomics, Proteomics & Bioinformatics 2021) in National Genomics Data Center (Nucleic Acids Res 2022), China National Center for Bioinformation / Beijing Institute of Genomics, Chinese Academy of Sciences (GSA: CRA008539) that are publicly accessible at https://ngdc.cncb.ac.cn/gsa.

Ethics statement

All experiments were performed in accordance with relevant guidelines and adhere to the ARRIVE guidelines (https://arriveguidelines.org/) for the reporting of animal experiments. All sheep are raised in accordance with the national feeding standard NT/T815-2004. All procedures performed were consistent with the National Research Council Guide for the Care and Use of Laboratory Animals. All experimental animal protocols were approved and performed in accordance with the requirements of the Animal Care and Use Committee at China Agricultural University (AW02012202-1-3).

Competing financial interests

The authors declare that there are no competing financial interests.

Author contributions

MMC performed the majority of experiments, data analysis, and drafted the manuscript. YZ performed a part of experiments and revised the manuscript. XLX, SJW and ZML helped with data analysis. XSZ, JLZ and XFG were responsible for the management of the feeding plant, slaughtering, and collecting samples. YMY helped to process some biological information data. SYQ, GY, SQW, HXL and AWW helped to collect and organize original data. GSL led the prokaryotic injection and embryo transfer. YL prepared the gene editing sheep. KY and HBH participated in project management. KY, FHL and ZXL conceived the project, revised manuscript and final approval of manuscript. All authors read and approved the final manuscript.

Acknowledgments

650   This work was supported by Major Agricultural Biological Breeding Project

651   (2022ZD04014), National Key Research and Development Program-Key Projects

652   (2021YFF1000704, 2021YFD1200900), Natural Science Foundation of China

653   (32072722), and National Transgenic Creature Breeding Grand Project

654   (2016zx08008-003).

  Figure Legends

Figure 1 Efficient generation of sheep carrying biallelic mutations in dual gene

via the CRISPR/Cas9 system

(A) Schematic of sgRNAs specific to exon 3 of the sheep MSTN locus. The crRNA

sequences are highlighted in blue typeface and the PAM in red. (B) Schematic of

sgRNAs specific to exon 3 of the sheep FGF5 locus. The crRNA sequences are

highlighted in blue typeface and the PAM in red. (C) T7EI assay for sgRNAs of

MSTN and FGF5 in sheep fetal fibroblasts. The cleavage bands are marked with an

red asterisk (*) and the indel frequencies were calculated using the expected

fragments. (D) Summary of the generation of sheep carrying biallelic mutations in

dual gene via zygote injection of Cas9 mRNA/sgRNAs. Biallelic mutation rate was

statistically analyzed using chi square test. *$P < 0.05$. (E) Analysis of genome

sequence and amino acid sequence of MSTN-modified sheep. The location of sgRNA

and PAM are highlighted in blue and red, respectively. The deletions are indicated by

a dashed line (-). (F) Analysis of genome sequence and amino acid sequence of

FGF5-modified sheep. The location of sgRNA and PAM are highlighted in blue and

red, respectively. The deletions are indicated by a dashed line (-).

Figure 2 The *MSTN*^Del273C mutation with *FGF5* knockout sheep highlights a dominant "double-muscle" phenotype and muscle fiber hyperplasia

(A) The 6-month-old WT and MF^-/- sheep. The genome-edited sheep displayed an obvious "double-muscle" phenotype compared with the WT. (B) The CT scanning image of the brisket and hip of WT and MF^-/- sheep. (C) HE sections of gluteus medius and longissimus dorsi of WT and MF^-/- sheep. Scale bar 200 μm. (D) Quantification of muscle fibre cell number of per unit area in WT (n=3) and MF^-/- (n=1) sheep. All data points were shown. (E-F) Quantification of muscle fibre cell area of gluteus medius and longissimus dorsi in WT (n=3) and MF^-/- (n=1) sheep. All

data points were shown. (G) HE sections of gluteus medius in WT and MF$^{+/-}$ sheep.

Scale bar 100 μm. (H) Quantification of muscle fibre cell area of gluteus medius in

WT (n=3) and MF$^{+/-}$ (n=4) sheep. (I) Quantification of muscle fibre cell number of

per unit area in WT (n=3) and MF$^{+/-}$ (n=4) sheep. (J) The percentage of

cross-sectional area of different size muscle fibers. (K) The proportion of different

muscles in carcass in WT (n=3) and MF$^{+/-}$ (n=4) sheep. Data: mean ± SEM. Unpaired

student's t-test were used for statistical analysis after the equal variance test,

otherwise the t-test with Welch's correction were performed. *$P < 0.05$, **$P < 0.01$,

***$P<0.001$, and ****$P<0.0001$.

Figure 3 The *MSTN*^{Del273C} mutation with *FGF5* knockout promote proliferation

and inhibit differentiation of skeletal muscle satellite cells

(A) The number of cells was detected by CCK-8 at 12h, 24h, 36h, 48h, and 60h in

GM (n=7-8 per group). (B-C) EdU assay showed that the number of EdU positive

cells and EdU labeling index were significantly increased in MF$^{+/-}$ cells (n=3). Scale

bar 130 μm. All data points were shown. (D-E) PI staining to detect cell cycle and

showed a significant reduce in the proportion of G1 phase and a significant increase in

the proportion of S phase in MF$^{+/-}$ cells (n=4). (F) The mRNA expression levels of

cell cycle marker genes and cell proliferation marker genes (n=3). (G) The mRNA

expression levels of myogenic differentiation marker genes MyoG, MyoD1, and

MyHC (n=3). (H-I) The protein expression levels of myogenic differentiation marker

genes MyoG, MyoD1, and MyHC (n=3). (J) The MyoG and MyHC

immunofluorescence staining of myotubes in DM2. Scale bar 130 μm. (K) The

myotube fusion index, which was represented by the number of cell nuclei in

myotubes/total cell nuclei (n=3). All data points were shown. (L) The number of

myotubes, which was the number of all myotubes in the field of view (n=3). All data

points were shown. (M) The number of nuclei per myotube (n=3). All data points

were shown. (N) The myotube diameter (n=3). To reflect the myotube diameter as

accurately as possible, the vertical line at the thinnest position of the myotube is taken

as the minimum measured (Min), the mid-perpendicular line of the long myotube axis

is taken as the middle measured (Middle), and the vertical line at the widest position

of the myotube is taken as the maximum measured (Max). All data points were shown.

Data: mean ± SEM. Unpaired student's t-test and chi square test were used for

statistical analysis. All student's t-test were performed after the equal variance test,

otherwise the t-test with Welch's correction were used. *$P < 0.05$, **$P < 0.01$, and

***$P<0.001$.

 Figure 4 The *MSTN*^{Del273C} mutation with *FGF5* knockout contributes to muscle

725 phenotype via MEK-ERK-FOSL1 axis

(A) Go enrichment analysis of DEGs. Among them, the top 20 entries with significant

enrichment are listed in biological process (BP). CC, cellular component; MF,

molecular function. (B) KEGG enrichment analysis of DEGs. (C) The mRNA

expression level of FOSL1 both at gluteus medius and longissimus dorsi in WT (n=3)

and $MF^{+/-}$ (n=4) sheep. (D) The mRNA expression level of FOSL1 both at GM and

DM2 in WT and $MF^{+/-}$ cells (n=3). (E) The expression level of FOSL1 mRNA during

myogenic differentiation (n=3). (F) The protein-protein interaction (PPI) analysis of

FOSL1, c-Fos and MyoD1. (G) The mRNA expression level of c-Fos and MyoD1 at

GM in WT and $MF^{+/-}$ myoblasts (n=3). (H) Schematic diagram of MyoD1 gene body,

promoter region and binding sites. (I-J) FOSL1 recognition motif in the MyoD1

promoter region. (K) FOSL1 ChIP-qPCR of motif 1 recognition region (n=3). (L)

FOSL1 ChIP-qPCR of motif 2 recognition region (n=3). (M) Dual luciferase assay for

the effect of FOSL1 on MyoD1 promoter activity (n=4). (N) Western blot of FOSL1,

c-Fos, and key kinases of MAPK signaling pathways at GM. (O) Quantification of

protein expression of FOSL1, c-Fos, and key kinases of MAPK signaling pathways at

GM (n=3). (P) Western blot of FOSL1, c-Fos, and key kinases of MAPK signaling

pathways at DM2. (Q) Quantification of protein expression of FOSL1, c-Fos, and key

kinases of MAPK signaling pathways at DM2 (n=3). Data: mean ± SEM. Unpaired

student's t-test was used for statistical analysis. All student's t-test were performed

after the equal variance test, otherwise the t-test with Welch's correction were used.

$*P < 0.05$, $**P < 0.01$, and $***P<0.001$.

Figure 5 The overexpression of *FOSL1* promotes proliferation and inhibits

differentiation of skeletal muscle satellite cells

(A) The mRNA expression level of FOSL1 at GM and DM2 after lentivirus infection

(n=3). (B) The number of cells detected by CCK-8 at 0h, 12h, 24h, 36h, 48h, and 60h

after infection with lentivirus (n=4-6). (C-D) EdU assay showed that the number of

EdU positive cells and EdU labeling index were significantly increased after infection

with lentivirus (n=3). Scale bar 130 μm. All data points were shown. (E) The mRNA

expression levels of cell cycle marker genes and cell proliferation marker genes (n=3).

(F) The mRNA expression levels of c-Fos and MyoD1 at GM after overexpression of

FOSL1 (n=3). (G-H) The protein expression levels of MyoD1 at GM after

overexpression of FOSL1 (n=3). (I-J) The protein expression levels of myogenic

differentiation marker genes MyoD1, MyoG and MyHC at DM2 after overexpression

of FOSL1 (n=3). (K) The MyoG and MyHC immunofluorescence staining of

myotubes at DM2 after overexpression of FOSL1. Scale bar 130 μm. (L-O) The

myotube fusion index, number of myotubes, number of nuclei per myotube and the

myotube diameter at DM2 after overexpression of FOSL1 (n=3). All data points were

shown. Data: mean ± SEM. Unpaired student's t-test and chi square test were used for

statistical analysis. All student's t-test were performed after the equal variance test,

otherwise the t-test with Welch's correction were used. *$P < 0.05$, **$P < 0.01$, and

***$P<0.001$.

Figure 6 The inhibition of *FOSL1* suppresses proliferation and promotes differentiation of skeletal muscle satellite cells

(A) The mRNA expression level of FOSL1 at GM and DM2 after inhibiting FOSL1 (n=3). (B) The number of cells detected by CCK-8 at 0h, 12h, 24h, 36h, 48h, and 60h after inhibiting FOSL1 (n=4-6). (C-D) EdU assay showed that the number of EdU positive cells and EdU labeling index were significantly decreased after inhibiting

FOSL1 (n=3). Scale bar 130 μm. All data points were shown. (E) The mRNA

expression levels of cell cycle marker genes and cell proliferation marker genes (n=3).

(F) The mRNA expression levels of c-Fos and MyoD1 at GM after inhibiting FOSL1

(n=3). (G-H) The protein expression levels of MyoD1 at GM after inhibiting FOSL1

(n=3). (I-J) The protein expression levels of myogenic differentiation marker genes

MyoD1, MyoG and MyHC at DM2 after inhibiting FOSL1 (n=3). (K) The MyoG and

MyHC immunofluorescence staining of myotubes at DM2 after inhibiting FOSL1.

Scale bar 130 μm. (L-O) The myotube fusion index, number of myotubes, number of

nuclei per myotube and the myotube diameter at DM2 after overexpression of FOSL1

(n=3). All data points were shown. Data: mean ± SEM. Unpaired student's t-test and

chi square test were used for statistical analysis. All student's t-test were performed

after the equal variance test, otherwise the t-test with Welch's correction were used.

$*P < 0.05$, $**P < 0.01$, $***P < 0.001$, and $****P < 0.0001$.

Figure 7 FOSL1 activity is a key regulator of myogenic differentiation and
muscle myofiber hyperplasia

(A) Immunofluorescence staining of myogenic differentiation markers MyoG and
MyHC in sheep skeletal muscle satellite cells at DM2 after addition of 20 μM TBHQ.
Scale bar 130 μm or 50 μm. (B-E) The myotube fusion index, number of myotubes,
number of nuclei per myotube and the myotube diameter at DM2 after addition of 20
μM TBHQ (n=3). All data points were shown. (F) Immunofluorescence staining of
myogenic differentiation markers MyoG and MyHC in sheep skeletal muscle satellite
cells at DM2 after addition of 1 μM PD98059. Scale bar 130 μm or 50 μm. (G-J) The
myotube fusion index, number of myotubes, number of nuclei per myotube and the

myotube diameter at DM2 after addition of 1 μM PD98059 (n=3). All data points

were shown. Data: mean ± SEM. Unpaired student's t-test and chi square test were

used for statistical analysis. All student's t-test were performed after the equal

variance test, otherwise the t-test with Welch's correction were used. *$P < 0.05$, **$P <$

0.01, and ***$P<0.001$.

Figure 8 The *MSTN*^{Del273C} mutation with *FGF5* knockout inhibit

calcium-dependent transcription signal pathway

(A-B) The protein expression level of CaMKII α/δ between WT and MF$^{+/-}$ cells at

GM (n=3). (C) Distribution of intracellular Ca^{2+} signals between WT and MF$^{+/-}$ cells

at GM. (D) Average intracellular Ca^{2+} fluorescence intensity between WT and MF$^{+/-}$

cells at GM (n=4). (E) The mRNA expression levels of Ca^{2+} channels and myoblast

fusion-related genes at GM (n=3). (F) The mRNA expression levels of Ca^{2+} channels

and myoblast fusion-related genes at DM2 (n=3). Data: mean ± SEM. Unpaired

student's t-test was used for statistical analysis. All student's t-test were performed

after the equal variance test, otherwise the t-test with Welch's correction were used.

$*P < 0.05$, $**P < 0.01$.

Figure 9 Schematic illustration of the regulation of muscle phenotypes by
***MSTN*^{Del273C} mutation with *FGF5* knockout**

The *MSTN*^{Del273C} mutation with *FGF5* knockout mediated the activation of
FOSL1 via MEK-ERK-FOSL1 axis. The activated FOSL1 promotes skeletal muscle
satellite cell proliferation and inhibits myogenic differentiation by inhibiting the
expression of MyoD1, and resulting in fusion to form smaller myotubes. In addition,
activated ERK1/2 may inhibit the secondary fusion of myotubes by Ca^{2+}-dependent
CaMKII activation pathway, leading to myoblasts fusion to form smaller myotubes.

Figure S1 *FGF5* mutation does not affect muscle fiber size

(A) HE sections of gluteus medius in WT and *FGF5*[+/-] sheep. Scale bar 100 μm. (B)

Quantification of muscle fibre cell area of gluteus medius in WT and *FGF5*[+/-] sheep.

(C) Quantification of muscle fibre cell number of per unit area in WT and *FGF5*[+/-]

sheep. (D) The percentage of cross-sectional area of different size muscle fibers. Data:

mean ± SEM. Unpaired student's t-test was used for statistical analysis. All student's

t-test were performed after the equal variance test, otherwise the t-test with Welch's

correction were used.

Figure S2 The *MSTN*^{Del273C} mutation with *FGF5* knockout has no potential effect

on MSTN expression and muscular dystrophy

(A) The mRNA expression levels of *MSTN* and *FGF5* of gluteus medius in WT (n=4)

and MF$^{+/-}$ (n=4) sheep at 3-month-old. (B) *MSTN* mRNA expression level of gluteus

medius in WT (n=3) and MF$^{+/-}$ (n=4) sheep. (C-D) MSTN protein expression level of

gluteus medius in WT (n=3) and MF$^{+/-}$ (n=4) sheep. (E) The proportion of centrally

nucleated myofibres between WT (n=3) and MF$^{+/-}$ (n=4) sheep. (F) The mRNA

expression of muscular dystrophy related genes between WT and MF$^{+/-}$ sheep (n=4).

Data: mean ± SEM. Unpaired student's t-test was used for statistical analysis. All

student's t-test were performed after the equal variance test, otherwise the t-test with

Welch's correction were used. *$P < 0.05$, **$P < 0.01$, and ***$P < 0.001$.

Figure S3 The myogenic differentiation ability of MF$^{+/-}$ cells was continuously

inhibited

(A) The MyoG and MyHC immunofluorescence staining of myotubes at DM4. Scale

bar 130 μm. (B-E) The myotube fusion index, number of myotubes, number of nuclei

per myotube and the myotube diameter at DM4 (n=3). (F) The MyoG and MyHC

immunofluorescence staining of myotubes at DM6. Scale bar 130 μm. (G-J) The

myotube fusion index, number of myotubes, number of nuclei per myotube and the

myotube diameter at DM6 (n=3). (K-N) The protein expression levels of myogenic

differentiation markers during myogenic differentiation between WT and MF$^{+/-}$ cells

(n=2-3). Data: mean ± SEM. Unpaired student's t-test and chi square test were used

for statistical analysis. All student's t-test were performed after the equal variance test,

otherwise the t-test with Welch's correction were used. *P < 0.05, **P < 0.01,

***P<0.001, and ****P<0.0001.

Figure S4 The effects of MSTN signaling pathway and MF$^{+/-}$ serum on

proliferation and differentiation of skeletal muscle satellite cells

(A) The mRNA expression levels of type I and type II receptors of MSTN between

WT (n=3) and MF$^{+/-}$ (n=4) sheep gluteus medius. (B) The mRNA expression levels of

the Smad family downstream of MSTN between WT (n=3) and MF$^{+/-}$ (n=4) sheep

gluteus medius. (C) The mRNA expression levels of the Jun family downstream of

MSTN between WT (n=3) and MF$^{+/-}$ (n=4) sheep gluteus medius. (D) The number of

cells detected by CCK-8 at 0h, 24h, and 48h after culturing cells with serum from WT

sheep and MF$^{+/-}$ sheep (n=4-5). (E) The mRNA expression levels of FOSL1 and

myogenic differentiation markers between WT and MF$^{+/-}$ sheep skeletal muscle

satellite cells cultured and induced differentiation in sheep serum (n=4). (F) The

mRNA expression levels of FOSL1 and myogenic differentiation markers in skeletal

muscle satellite cells cultured and induced differentiation by serum from WT and

MF$^{+/-}$ sheep (n=4). Data: mean ± SEM. Unpaired student's t-test and chi square test

were used for statistical analysis. All student's t-test were performed after the equal

variance test, otherwise the t-test with Welch's correction were used. *P < 0.05, **P <

0.01, and ***P<0.001.

References

Alli, N. S., Yang, E. C., Miyake, T., Aziz, A., Collins-Hooper, H., Patel, K. & McDermott, J. C. (2013a). Signal-dependent fra-2 regulation in skeletal muscle reserve and satellite cells. *Cell Death Dis* 4: e692.

Alli, N. S., Yang, E. C., Miyake, T., Aziz, A., Collins-Hooper, H., Patel, K. & McDermott, J. C. (2013b). Signal-dependent fra-2 regulation in skeletal muscle reserve and satellite cells. *Cell Death Dis* 4(6): e692.

Baig, M. H., Ahmad, K., Moon, J. S., Park, S. Y., Ho Lim, J., Chun, H. J., Qadri, A. F., Hwang, Y. C., Jan, A. T., Ahmad, S. S., Ali, S., Shaikh, S., Lee, E. J. & Choi, I. (2022). Myostatin and its Regulation: A Comprehensive Review of Myostatin Inhibiting Strategies. *Front Physiol* 13: 876078.

Bao, D., Ma, Y., Zhang, X., Guan, F., Chen, W., Gao, K., Qin, C. & Zhang, L. (2015). Preliminary Characterization of a Leptin Receptor Knockout Rat Created by CRISPR/Cas9 System. *Sci Rep* 5: 15942.

Bayer, K. U., Lohler, J. & Harbers, K. (1996). An alternative, nonkinase product of the brain-specifically expressed Ca2+/calmodulin-dependent kinase II alpha isoform gene in skeletal muscle. *Mol Cell Biol* 16(1): 29-36.

Boman, I. A., Klemetsdal, G., Blichfeldt, T., Nafstad, O. & Vage, D. I. (2009). A frameshift mutation in the coding region of the myostatin gene (MSTN) affects carcass conformation and fatness in Norwegian White Sheep (Ovis aries). *Anim Genet* 40(4): 418-422.

Boman, I. A. & Vage, D. I. (2009). An insertion in the coding region of the myostatin (MSTN) gene affects carcass conformation and fatness in the Norwegian Spaelsau (Ovis aries). *BMC Res Notes* 2: 98.

Campbell, J. S., Wenderoth, M. P., Hauschka, S. D. & Krebs, E. G. (1995). Differential activation of mitogen-activated protein kinase in response to basic fibroblast growth factor in skeletal muscle cells. *Proc Natl Acad Sci U S A* 92(3): 870-874.

Chen, M., Zhang, L., Guo, Y., Liu, X., Song, Y., Li, X., Ding, X. & Guo, H. (2021a). A novel lncRNA promotes myogenesis of bovine skeletal muscle satellite cells via PFN1-RhoA/Rac1. *J Cell Mol Med*.

Chen, M. M., Zhao, Y. P., Zhao, Y., Deng, S. L. & Yu, K. (2021b). Regulation of Myostatin on the Growth and Development of Skeletal Muscle. *Front Cell Dev Biol* 9: 785712.

Chen, Y., Zheng, Y., Kang, Y., Yang, W., Niu, Y., Guo, X., Tu, Z., Si, C., Wang, H., Xing, R., Pu, X., Yang, S. H., Li, S., Ji, W. & Li, X. J. (2015). Functional disruption of the dystrophin gene in rhesus monkey using CRISPR/Cas9. *Hum Mol Genet* 24(13): 3764-3774.

Constantin, B., Cognard, C. & Raymond, G. (1996). Myoblast fusion requires cytosolic calcium elevation but not activation of voltage-dependent calcium channels. *Cell Calcium* 19(5): 365-374.

Dehnavi, E., Ahani Azari, M., Hasani, S., Nassiry, M. R., Mohajer, M., Khan Ahmadi, A., Shahmohamadi, L. & Yousefi, S. (2012). Polymorphism of Myostatin Gene in Intron 1 and 2 and Exon 3, and Their Associations with Yearling Weight, Using PCR-RFLP and PCR-SSCP Techniques in Zel Sheep. *Biotechnol Res Int* 2012: 472307.

Dierks, C., Momke, S., Philipp, U. & Distl, O. (2013). Allelic heterogeneity of FGF5 mutations causes the long-hair phenotype in dogs. *Anim Genet* 44(4): 425-431.

Dilger, A. C., Gabriel, S. R., Kutzler, L. W., McKeith, F. K. & Killefer, J. (2010). The myostatin null mutation and clenbuterol administration elicit additive effects in mice. *Animal* 4(3): 466-471.

Eigler, T., Zarfati, G., Amzallag, E., Sinha, S., Segev, N., Zabary, Y., Zaritsky, A., Shakked, A., Umansky, K. B., Schejter, E. D., Millay, D. P., Tzahor, E. & Avinoam, O. (2021). ERK1/2 inhibition promotes robust myotube growth via CaMKII activation resulting in myoblast-to-myotube fusion. *Dev Cell* 56(24): 3349-3363 e3346.

Fan, Z., Liu, Z., Xu, K., Wu, T., Ruan, J., Zheng, X., Bao, S., Mu, Y., Sonstegard, T. &
Li, K. (2022). Long-term, multidomain analyses to identify the breed and
allelic effects in MSTN-edited pigs to overcome lameness and sustainably
improve nutritional meat production. *Sci China Life Sci* 65(2): 362-375.
Grisolia, A. B., D'Angelo, G. T., Porto Neto, L. R., Siqueira, F. & Garcia, J. F. (2009).
Myostatin (GDF8) single nucleotide polymorphisms in Nellore cattle. *Genet*
Mol Res 8(3): 822-830.
Grobet, L., Martin, L. J., Poncelet, D., Pirottin, D., Brouwers, B., Riquet, J.,
Schoeberlein, A., Dunner, S., Menissier, F., Massabanda, J., Fries, R., Hanset,
R. & Georges, M. (1997). A deletion in the bovine myostatin gene causes the
double-muscled phenotype in cattle. *Nat Genet* 17(1): 71-74.
Gui, T., Sun, Y., Shimokado, A. & Muragaki, Y. (2012). The Roles of
Mitogen-Activated Protein Kinase Pathways in TGF-beta-Induced
Epithelial-Mesenchymal Transition. *J Signal Transduct* 2012: 289243.
Hai, T., Teng, F., Guo, R., Li, W. & Zhou, Q. (2014). One-step generation of knockout
pigs by zygote injection of CRISPR/Cas system. *Cell Res* 24(3): 372-375.
Haidet, A. M., Rizo, L., Handy, C., Umapathi, P., Eagle, A., Shilling, C., Boue, D.,
Martin, P. T., Sahenk, Z., Mendell, J. R. & Kaspar, B. K. (2008). Long-term
enhancement of skeletal muscle mass and strength by single gene
administration of myostatin inhibitors. *Proc Natl Acad Sci U S A* 105(11):
4318-4322.
Han, J., Forrest, R. H. & Hickford, J. G. (2013). Genetic variations in the myostatin
gene (MSTN) in New Zealand sheep breeds. *Mol Biol Rep* 40(11): 6379-6384.
Hanset, R. & Michaux, C. (1985). On the genetic determinism of muscular
hypertrophy in the Belgian White and Blue cattle breed. I. Experimental data.
Genet Sel Evol (1983) 17(3): 359-368.
Hebert, J. M., Rosenquist, T., Gotz, J. & Martin, G. R. (1994). FGF5 as a regulator of
the hair growth cycle: evidence from targeted and spontaneous mutations. *Cell*
78(6): 1017-1025.
Higgins, C. A., Petukhova, L., Harel, S., Ho, Y. Y., Drill, E., Shapiro, L., Wajid, M. &
Christiano, A. M. (2014). FGF5 is a crucial regulator of hair length in humans.
Proc Natl Acad Sci U S A 111(29): 10648-10653.
Hongbing HAN, Y. M., Tao WANG,Ling LIAN,Xiuzhi TIAN,Rui HU,Shoulong
DENG,Kongpan LI,Feng WANG,Ning LI,Guoshi LIU,Yaofeng
ZHAO,Zhengxing LIAN (2014). One-step generation of myostatin gene
knockout sheep via the CRISPR/Cas9 system. *Front. Agr. Sci. Eng.* 1(1): 2-5.
Huang, Z., Chen, D., Zhang, K., Yu, B., Chen, X. & Meng, J. (2007). Regulation of
myostatin signaling by c-Jun N-terminal kinase in C2C12 cells. *Cell Signal*
19(11): 2286-2295.
Jones, N. C., Fedorov, Y. V., Rosenthal, R. S. & Olwin, B. B. (2001). ERK1/2 is
required for myoblast proliferation but is dispensable for muscle gene
expression and cell fusion. *J Cell Physiol* 186(1): 104-115.
Kambadur, R., Sharma, M., Smith, T. P. & Bass, J. J. (1997). Mutations in myostatin
(GDF8) in double-muscled Belgian Blue and Piedmontese cattle. *Genome Res*
7(9): 910-916.
Kehler, J. S., David, V. A., Schaffer, A. A., Bajema, K., Eizirik, E., Ryugo, D. K.,
Hannah, S. S., O'Brien, S. J. & Menotti-Raymond, M. (2007). Four
independent mutations in the feline fibroblast growth factor 5 gene determine
the long-haired phenotype in domestic cats. *J Hered* 98(6): 555-566.
Kijas, J. W., McCulloch, R., Edwards, J. E., Oddy, V. H., Lee, S. H. & van der Werf, J.
(2007). Evidence for multiple alleles effecting muscling and fatness at the
ovine GDF8 locus. *BMC Genet* 8: 80.
Kotani, H., Taimatsu, K., Ohga, R., Ota, S. & Kawahara, A. (2015). Efficient Multiple
Genome Modifications Induced by the crRNAs, tracrRNA and Cas9 Protein
Complex in Zebrafish. *PLoS One* 10(5): e0128319.
Lee, S. J. (2021). Targeting the myostatin signaling pathway to treat muscle loss and
metabolic dysfunction. *J Clin Invest* 131(9).

Legrand, R., Tiret, L. & Abitbol, M. (2014). Two recessive mutations in FGF5 are associated with the long-hair phenotype in donkeys. *Genet Sel Evol* 46: 65.

Li, L., Chambard, J. C., Karin, M. & Olson, E. N. (1992). Fos and Jun repress transcriptional activation by myogenin and MyoD: the amino terminus of Jun can mediate repression. *Genes Dev* 6(4): 676-689.

Li, Y., Lian, D., Deng, S., Zhang, X., Zhang, J., Li, W., Bai, H., Wang, Z., Wu, H., Fu, J., Han, H., Feng, J., Liu, G., Lian, L. & Lian, Z. (2016). Efficient production of pronuclear embryos in breeding and nonbreeding season for generating transgenic sheep overexpressing TLR4. *J Anim Sci Biotechnol* 7: 38.

Lv, Q., Yuan, L., Deng, J., Chen, M., Wang, Y., Zeng, J., Li, Z. & Lai, L. (2016). Efficient Generation of Myostatin Gene Mutated Rabbit by CRISPR/Cas9. *Sci Rep* 6: 25029.

Marchitelli, C., Savarese, M. C., Crisa, A., Nardone, A., Marsan, P. A. & Valentini, A. (2003). Double muscling in Marchigiana beef breed is caused by a stop codon in the third exon of myostatin gene. *Mamm Genome* 14(6): 392-395.

Marques, C., Unterkircher, T., Kroon, P., Oldrini, B., Izzo, A., Dramaretska, Y., Ferrarese, R., Kling, E., Schnell, O., Nelander, S., Wagner, E. F., Bakiri, L., Gargiulo, G., Carro, M. S. & Squatrito, M. (2021). NF1 regulates mesenchymal glioblastoma plasticity and aggressiveness through the AP-1 transcription factor FOSL1. *Elife* 10.

Mathes, S., Fahrner, A., Ghoshdastider, U., Rudiger, H. A., Leunig, M., Wolfrum, C. & Krutzfeldt, J. (2021). FGF-2-dependent signaling activated in aged human skeletal muscle promotes intramuscular adipogenesis. *Proc Natl Acad Sci U S A* 118(37).

McPherron, A. C., Lawler, A. M. & Lee, S. J. (1997). Regulation of skeletal muscle mass in mice by a new TGF-beta superfamily member. *Nature* 387(6628): 83-90.

Michailovici, I., Harrington, H. A., Azogui, H. H., Yahalom-Ronen, Y., Plotnikov, A., Ching, S., Stumpf, M. P., Klein, O. D., Seger, R. & Tzahor, E. (2014). Nuclear to cytoplasmic shuttling of ERK promotes differentiation of muscle stem/progenitor cells. *Development* 141(13): 2611-2620.

Morissette, M. R., Cook, S. A., Buranasombati, C., Rosenberg, M. A. & Rosenzweig, A. (2009). Myostatin inhibits IGF-I-induced myotube hypertrophy through Akt. *Am J Physiol Cell Physiol* 297(5): C1124-1132.

Nishi, M., Yasue, A., Nishimatu, S., Nohno, T., Yamaoka, T., Itakura, M., Moriyama, K., Ohuchi, H. & Noji, S. (2002). A missense mutant myostatin causes hyperplasia without hypertrophy in the mouse muscle. *Biochem Biophys Res Commun* 293(1): 247-251.

Niu, Y., Shen, B., Cui, Y., Chen, Y., Wang, J., Wang, L., Kang, Y., Zhao, X., Si, W., Li, W., Xiang, A. P., Zhou, J., Guo, X., Bi, Y., Si, C., Hu, B., Dong, G., Wang, H., Zhou, Z., Li, T., Tan, T., Pu, X., Wang, F., Ji, S., Zhou, Q., Huang, X., Ji, W. & Sha, J. (2014). Generation of gene-modified cynomolgus monkey via Cas9/RNA-mediated gene targeting in one-cell embryos. *Cell* 156(4): 836-843.

Pothuraju, M., Mishra, S. K., Kumar, S. N., Mohamed, N. F., Kataria, R. S., Yadav, D. K. & Arora, R. (2015). Polymorphism in the Coding Region Sequence of Gdf8 Gene in Indian Sheep. *Genetika* 51(11): 1297-1300.

Qian, L., Tang, M., Yang, J., Wang, Q., Cai, C., Jiang, S., Li, H., Jiang, K., Gao, P., Ma, D., Chen, Y., An, X., Li, K. & Cui, W. (2015). Targeted mutations in myostatin by zinc-finger nucleases result in double-muscled phenotype in Meishan pigs. *Sci Rep* 5: 14435.

Sato, M., Koriyama, M., Watanabe, S., Ohtsuka, M., Sakurai, T., Inada, E., Saitoh, I., Nakamura, S. & Miyoshi, K. (2015). Direct Injection of CRISPR/Cas9-Related mRNA into Cytoplasm of Parthenogenetically Activated Porcine Oocytes Causes Frequent Mosaicism for Indel Mutations. *Int J Mol Sci* 16(8): 17838-17856.

Segales, J., Perdiguero, E. & Munoz-Canoves, P. (2016). Regulation of Muscle Stem Cell Functions: A Focus on the p38 MAPK Signaling Pathway. *Front Cell Dev Biol* 4: 91.

Sjakste, T., Paramonova, N., Grislis, Z., Trapina, I. & Kairisa, D. (2011). Analysis of the single-nucleotide polymorphism in the 5'UTR and part of intron I of the sheep MSTN gene. *DNA Cell Biol* 30(7): 433-444.

Sobolev, V. V., Khashukoeva, A. Z., Evina, O. E., Geppe, N. A., Chebysheva, S. N., Korsunskaya, I. M., Tchepourina, E. & Mezentsev, A. (2022). Role of the Transcription Factor FOSL1 in Organ Development and Tumorigenesis. *Int J Mol Sci* 23(3).

Sundberg, J. P., Rourk, M. H., Boggess, D., Hogan, M. E., Sundberg, B. A. & Bertolino, A. P. (1997). Angora mouse mutation: altered hair cycle, follicular dystrophy, phenotypic maintenance of skin grafts, and changes in keratin expression. *Vet Pathol* 34(3): 171-179.

Sung, Y. H., Kim, J. M., Kim, H. T., Lee, J., Jeon, J., Jin, Y., Choi, J. H., Ban, Y. H., Ha, S. J., Kim, C. H., Lee, H. W. & Kim, J. S. (2014). Highly efficient gene knockout in mice and zebrafish with RNA-guided endonucleases. *Genome Res* 24(1): 125-131.

Tobin, S. W., Yang, D., Girgis, J., Farahzad, A., Blais, A. & McDermott, J. C. (2016). Regulation of Hspb7 by MEF2 and AP-1: implications for Hspb7 in muscle atrophy. *J Cell Sci* 129(21): 4076-4090.

Tu, Z., Yang, W., Yan, S., Yin, A., Gao, J., Liu, X., Zheng, Y., Zheng, J., Li, Z., Yang, S., Li, S., Guo, X. & Li, X. J. (2017). Promoting Cas9 degradation reduces mosaic mutations in non-human primate embryos. *Sci Rep* 7: 42081.

Wan, H., Feng, C., Teng, F., Yang, S., Hu, B., Niu, Y., Xiang, A. P., Fang, W., Ji, W., Li, W., Zhao, X. & Zhou, Q. (2015). One-step generation of p53 gene biallelic mutant Cynomolgus monkey via the CRISPR/Cas system. *Cell Res* 25(2): 258-261.

Wang, H. & Yang, H. (2019). Gene-edited babies: What went wrong and what could go wrong. *PLoS Biol* 17(4): e3000224.

Wang, H., Yang, H., Shivalila, C. S., Dawlaty, M. M., Cheng, A. W., Zhang, F. & Jaenisch, R. (2013). One-step generation of mice carrying mutations in multiple genes by CRISPR/Cas-mediated genome engineering. *Cell* 153(4): 910-918.

Wang, Q. & McPherron, A. C. (2012). Myostatin inhibition induces muscle fibre hypertrophy prior to satellite cell activation. *J Physiol* 590(9): 2151-2165.

Wang, X., Yu, H., Lei, A., Zhou, J., Zeng, W., Zhu, H., Dong, Z., Niu, Y., Shi, B., Cai, B., Liu, J., Huang, S., Yan, H., Zhao, X., Zhou, G., He, X., Chen, X., Yang, Y., Jiang, Y., Shi, L., Tian, X., Wang, Y., Ma, B., Huang, X., Qu, L. & Chen, Y. (2015). Generation of gene-modified goats targeting MSTN and FGF5 via zygote injection of CRISPR/Cas9 system. *Sci Rep* 5: 13878.

Wegner, J., Albrecht, E., Fiedler, I., Teuscher, F., Papstein, H. J. & Ender, K. (2000). Growth- and breed-related changes of muscle fiber characteristics in cattle. *J Anim Sci* 78(6): 1485-1496.

Wijaya, Y. T., Setiawan, T., Sari, I. N., Park, K., Lee, C. H., Cho, K. W., Lee, Y. K., Lim, J. Y., Yoon, J. K., Lee, S. H. & Kwon, H. Y. (2022). Ginsenoside Rd ameliorates muscle wasting by suppressing the signal transducer and activator of transcription 3 pathway. *J Cachexia Sarcopenia Muscle*.

Xie, S. J., Li, J. H., Chen, H. F., Tan, Y. Y., Liu, S. R., Zhang, Y., Xu, H., Yang, J. H., Liu, S., Zheng, L. L., Huang, M. B., Guo, Y. H., Zhang, Q., Zhou, H. & Qu, L. H. (2018). Inhibition of the JNK/MAPK signaling pathway by myogenesis-associated miRNAs is required for skeletal muscle development. *Cell Death Differ* 25(9): 1581-1597.

Xu, T. S., Gu, L. H., Zhang, X. H., Ye, B. G., Liu, X. L. & Hou, S. S. (2013). Characterization of myostatin gene (MSTN) of Pekin duck and the association of its polymorphism with breast muscle traits. *Genet Mol Res* 12(3): 3166-3177.

Yoshizawa, Y., Wada, K., Shimoi, G., Kameyama, Y., Wakabayashi, Y., Fukuta, K. &
Hashizume, R. (2015). A 1-bp deletion in Fgf5 causes male-dominant long
hair in the Syrian hamster. *Mamm Genome* 26(11-12): 630-637.
Zhang, R., Li, Y., Jia, K., Xu, X., Li, Y., Zhao, Y., Zhang, X., Zhang, J., Liu, G., Deng,
S. & Lian, Z. (2020). Crosstalk between androgen and Wnt/beta-catenin leads
to changes of wool density in FGF5-knockout sheep. *Cell Death Dis* 11(5):
407.
Zhiliang, G., Dahai, Z., Ning, L., Hui, L., Xuemei, D. & Changxin, W. (2004). The
single nucleotide polymorphisms of the chicken myostatin gene are associated
with skeletal muscle and adipose growth. *Sci China C Life Sci* 47(1): 25-30.
Zhou, J., Wang, J., Shen, B., Chen, L., Su, Y., Yang, J., Zhang, W., Tian, X. & Huang,
X. (2014). Dual sgRNAs facilitate CRISPR/Cas9-mediated mouse genome
targeting. *FEBS J* 281(7): 1717-1725.
Zhu, G. H., Huang, J., Bi, Y., Su, Y., Tang, Y., He, B. C., He, Y., Luo, J., Wang, Y.,
Chen, L., Zuo, G. W., Jiang, W., Luo, Q., Shen, J., Liu, B., Zhang, W. L., Shi,
Q., Zhang, B. Q., Kang, Q., Zhu, J., Tian, J., Luu, H. H., Haydon, R. C., Chen,
Y. & He, T. C. (2009). Activation of RXR and RAR signaling promotes
myogenic differentiation of myoblastic C2C12 cells. *Differentiation* 78(4):
195-204.
Zhu, X., Hadhazy, M., Wehling, M., Tidball, J. G. & McNally, E. M. (2000).
Dominant negative myostatin produces hypertrophy without hyperplasia in
muscle. *FEBS Lett* 474(1): 71-75.

1140