Recent evolutionary origin and localized diversity hotspots of mammalian coronaviruses.

Renan Maestri[a,b,1], Benoît Perez-Lamarque[a,c,1], Anna Zhukova[d], Hélène Morlon[a]

[a]Institut de Biologie de l'École Normale Supérieure (IBENS), École Normale Supérieure, CNRS, INSERM, Université PSL, Paris, France

[b]Departamento de Ecologia, Instituto de Biociências, Universidade Federal do Rio Grande do Sul, Porto Alegre, RS, Brazil

[c]Institut de Systématique, Évolution, Biodiversité (ISYEB), Muséum national d'histoire naturelle, CNRS, Sorbonne Université, EPHE, UA, Paris, France

[d]Institut Pasteur, Université Paris Cité, Bioinformatics and Biostatistics Hub, Paris, France

[1]R.M. and B.P.-L. contributed equally to this work.

*Corresponding author: Renan Maestri

Email: renanmaestri@gmail.com

Author Contributions: R.M. and H.M. conceived the study; R.M, B.P-L., A.Z. and H.M. designed and performed the research; B.P-L. and R.M. analyzed the data; R.M, B.P-L., A.Z. and H.M. wrote the paper.

Competing Interest Statement: The authors declare no competing interest.

Classification: Biological Sciences. Evolution.

Keywords: Paste the keywords here. There should be at least three and no more than five.

coronavirus evolution | diversity of coronaviruses | preferential host switches | parasite diversification

Abstract

Several coronaviruses infect humans, with three, including the SARS-CoV2, causing diseases. While coronaviruses are especially prone to induce pandemics, we know little about their evolutionary history, host-to-host transmissions, and biogeography, which impedes the prediction of future transmission scenarios. One of the difficulties lies in dating the origination of the family, a particularly challenging task for RNA viruses in general. Previous cophylogenetic tests of virus-host associations, including in the Coronaviridae family, have suggested a virus-host codiversification history stretching many millions of years. Here, we establish a framework for robustly testing scenarios of ancient origination and codiversification *versus* recent origination and diversification by host switches. Applied to coronaviruses and their mammalian hosts, our results support a scenario of recent origination of coronaviruses in bats and diversification by host switches, with preferential host switches within mammalian orders. Hotspots of coronavirus diversity, concentrated in East Asia and Europe, are consistent with this scenario of relatively recent origination and localized host switches. Spillovers from bats to other species are rare, but have the highest probability to be towards humans than to any other mammal species, implicating humans as the evolutionary intermediate host. The high host-switching rates within orders, as well as between humans, domesticated mammals, and non-flying wild mammals, indicates the potential for rapid additional spreading of coronaviruses across the world. Our results suggest that the evolutionary history of extant mammalian coronaviruses is recent, and that cases of long-term virus–host codiversification have been largely over-estimated.

Main Text

Introduction

Coronaviruses are RNA-viruses of the family Coronaviridae, comprising positive-sense and single-stranded viruses that have the largest genomes among nidoviruses (1, 2). As several other RNA viruses, they may cause diseases in humans and other animals (3). Depending on the taxonomic arrangement, seven (4–6) or eight (7) species of coronaviruses infect humans, three of which being pathogenic: the SARS-CoV (8, 9), the MERS-CoV (10), and the SARS-CoV2 (11). The latter is at the origin of the recent COVID-19 pandemic that infected more than 620 million people and caused the death of more than six and a half million (12). Coronaviruses' high frequency of recombination (13), broad host range, and high mutation rates (7) make them especially prone to causing yet future diseases. Nevertheless, their evolutionary history and biogeography are very poorly understood. Resolving the evolutionary origins of Coronaviridae, understanding how they diversified, and characterizing their geographic diversity patterns would facilitate attempts to predict future zoonoses (7, 14, 15).

Coronaviruses infect mammals, birds, and fish (2), although they predominate in mammalian species (16–20). A consensus exists on the taxonomic segregation of four genera within Coronaviridae: Orthocoronavirinae, namely Alpha-, Beta-, Gamma- and Deltacoronavirus (2, 21). Alpha- and Betacoronaviruses are found exclusively in mammals, while Delta- and Gammacoronaviruses infect mostly birds but also mammals to a lesser extent (20, 22, 23). Coronaviruses are most numerous and genetically diversified in mammals (2, 23), in particular bats, suggesting a mammalian origin in bats (2, 20, 23, 24), although this remains to be tested.

The timing of origination of the Coronaviridae family is debated, with results that vary by several orders of magnitude. Woo et al (23) found a recent origin, around 10 thousand years ago. This dating was obtained by sequencing the well-conserved RNA-dependent RNA-polymerase (RdRp) genome region of representatives of all four coronavirus genera, and fitting to these

sequences a neutral nucleotide-based substitution model with an uncorrelated log-normal relaxed
clock (25) calibrated with serial samples. This calibration provided a mean substitution rate
estimate of 1.3 x 10-4 substitutions per site per year. Wertheim et al. (26) used this estimate and
the same genome region (RdRp), but with a codon-based substitution model accounting for the
effect of selection. Indeed, purifying selection can lead to an underestimation of viral origins when
not accounted for (26, 27). They found an ancient origin, around 293 (95% confidence interval,
190 to 489) million years ago (26). More recently, Hayman & Knox (28) obtained similar results,
but using the splitting times of hosts as constraints, therefore assuming a priori that coronaviruses
codiversified with their hosts.
84
More generally, dating the phylogenies of RNA virus families is a difficult task (29). While
for some of them dated calibration points can be used, based on orthologous copies of
endogenous virus elements (EVEs) present in the genomes of related mammalian species with
known times of divergence (30), in many others, including in the Coronaviridae, such elements
have not been found (31). Despite the difficulty in dating viral families, it has been proposed, from
cophylogenetic analyses investigating the congruence of the host and viral phylogenetic trees
(32), that vertebrate-associated RNA viruses have codiversified with their hosts over hundreds of
millions of years (31, 33). Indeed, RNA virus phylogenies tend to mirror that of their hosts; for
example, closely-related coronaviruses infect closely-related mammals (e.g. (28)). However, a
major caveat is that such cophylogenetic signals can emerge when viruses diversify by host
switches preferentially occurring among closely-related hosts, in the absence of any cospeciation
event (34). Event-based cophylogenetic methods can in principle identify cospeciation and host
switches events (32, 34), but their behavior in the presence of diversification by preferential host
switches is not well understood. Under a perfect codiversification scenario, host and symbiont
phylogenies would be identical. Events of host switches, duplications and losses induce
mismatches, and cophylogenetic methods aim to identify parsimonious sets of events that allow
"reconciling" the two phylogenies (34, 35). However, most of these methods rely entirely on tree
topology (and not branching times), such that time-inconsistent host switches between non-
contemporary host lineages are allowed during the reconciliation. In the presence of preferential
host switches, these methods may thus favor biologically unrealistic reconciliations that involve
cospeciation events and 'back-in-time' host switches to reconciliations that involve more frequent
contemporary host switches. This would have remained unnoticed, unless users of the methods
specifically looked at the time consistency of the inferred host switches, which is usually not done.
108
Here, we establish a framework for testing scenarios of ancient origination and
codiversification versus recent origination and diversification by host switches that combines
probabilistic cophylogenetic models and biogeographic analyses (Fig 1). We then apply this
framework to the Coronaviridae-mammals association. We assemble a dataset of all mammalian
hosts of coronaviruses and a complete association matrix between host species and species-like
Operational Taxonomic Units (sOTUs) of coronaviruses, as well as geographic repartition of
Coronaviridae and their mammalian hosts. We construct a new Coronaviridae tree based on a
recent proposition for the use of a well-conserved region of their RNA genome (36, 37). Under the
ancient origination scenario (Fig 1A), long-term vertical transmission of Coronaviridae within
mammalian lineages could lead to events of mammal-coronavirus cospeciations. Coronaviruses'
diversification would then be modulated by both cospeciations and horizontal host switches from
one mammalian lineage to another (26, 31). The most recent common ancestor of coronaviruses
could even have infected the most recent common ancestor of mammals and birds (26). Under
the recent origination scenario (Fig 1B), codiversification with hosts is virtually impossible, and
coronaviruses' diversification would then be largely dominated by recent host switches.
Expectations for the output of reconciliation and biogeographic analyses under these different
scenarios, as well as a scenario of random associations, are explicated in Fig 1. We identify the
likely origination of coronaviruses in the mammalian tree, quantify the frequency of cospeciation
and host-switching events, and locate these host switches, therefore identifying 'reservoirs' of
Coronaviridae and potential transmission routes across mammals.

Results

By screening the 46 sOTUs of Coronaviridae identified by Edgar et al. (36) in public databases,
we found 35 that were associated with mammalian hosts. Our trees of these 35 sOTUs support a
well-defined split between Alphacoronaviruses and the other genera, regardless of the
phylogenetic method used (Fig 2; *SI Appendix*, Fig. S1). Overall, Alphacoronaviruses form a
monophyletic clade, Delta- and Gammacoronoviruses form sister clades, with the main
uncertainty being on the placement of their ancestor in relation to Beta (i.e. as a sister to a
monophyletic Beta-clade (Fig. 2) or within the Beta-clade (*SI Appendix*, Fig. S1).

We found that mammalian hosts of coronaviruses belong to 31 families and 10 orders of
mammals, and are widely distributed throughout the mammalian phylogeny (SI Appendix, Fig.
S2). Most mammalian hosts are bats (Chiroptera - 55 species), followed by rodents (Rodentia -
22 species), artiodactyls (Artiodactyla - 15 species), carnivores (Carnivora - 11 species), and
primates (Primates - 5 species). Five other orders have at least one representative species:
Eulipotyphla (4), Lagomorpha (1), Perissodactyla (1), Pholidota (1), and Sirenia (1). The number
of mammalian hosts per coronavirus' sOTU varies across the Coronaviridae tree, ranging from 1
to 22 species, with an average of 4.94 (Fig. 2). Of the 35 sOTUs, 23 are found in at least one bat
species and 17, mostly in alphacoronaviruses, are found exclusively in bats (Fig. 2). Eight sOTUs
are found in humans, six of which, including the three pathogenetic sOTUs, are
betacoronaviruses. Betacoronaviruses infect a larger average number of hosts and a larger
diversity of non-bat species than Alphacoronaviruses. Twenty-two coronaviruses occur in more
than one species; of those, 11 are found in multiple orders (Fig. 2; *SI Appendix*, Fig. S3) and 11
in multiple species of a single order (*SI Appendix*, Fig. S3).

We first tested whether closely-related coronaviruses tend to infect closely-related
mammals. A negative answer to this question would suggest that the diversification of
Coronaviridae is independent of mammalian history, excluding the scenarios of codiversification
or diversification per preferential host switches (Fig. 1). To the contrary, we found a significant
phylogenetic signal for the overall association between coronaviruses and mammals (Mantel test:
r= 0.38; *P*= 0.0001) and vice-versa (r= 0.29; *P*= 0.0001), after accounting for the confounding
phylogenetic signal in the number of partners (38). Mantel tests across sub-clades of both
phylogenies revealed that this overall phylogenetic signal is linked to phylogenetic signal in the
deep nodes of the Coronaviridae and mammal phylogenies rather than at shallow phylogenetic
scales (*SI Appendix*, Fig. S4). This pattern could arise from ancient codiversification followed by
un-preferential host switches, or from recent host switches preferentially occurring between hosts
from the same high-level taxonomic grouping (such as mammalian orders). We also found that
closely related coronaviruses tend to infect a similar number of hosts (r= 0.29; *P*=0.002), while
closely related mammals do not tend to host a similar number of distinct coronaviruses (r= 0.04,
P=0.1), suggesting that coronaviruses' specificity towards hosts is evolutionarily conserved while
hosts' susceptibility to coronaviruses is not.

To further investigate the hypotheses of ancient codiversification versus recent host
switches, we used a probabilistic cophylogenetic model, the amalgamated likelihood estimation
(ALE - (39)), that reconciliates the host and symbiont phylogenies using events of cospeciations,
host switches, duplications, or losses, while accounting for phylogenetic uncertainty in the
symbiont phylogenies and undersampling of the host species (35, 39, 40). The main version of
ALE we used is an "undated" version that accounts for topology but not branch lengths, as the
dated version did not perform well on our data (see Methods). Time-inconsistent host switches
are thus allowed during the reconciliation. If the scenario of ancient diversification holds, we
expect to find reconciliations requiring more cospeciations and fewer host switches than expected
under a scenario of independent evolution (hereafter referred to as 'significant reconciliation'),

and few time-inconsistent switches (Fig. 1A). Under the alternative scenario of recent origination
and diversification by preferential host switches, we also expect to infer a significant reconciliation,
but with many time-inconsistent switches, as the algorithm tends to explain the cophylogenetic
signal in the interactions by cospeciation events (Fig. 1B). We indeed found a significant
reconciliation between the Coronaviridae and the mammalian trees, confirming the non-
independence of their evolution, which we evaluated by randomly shuffling mammal species
across the full tree or within biogeographic regions (*SI Appendix*, Fig. S7). ALE reconciliations
inferred average numbers of 145 cospeciations, 65 losses, 0 duplication, and 92 host switches.
Without investigating the time-consistency of the host switches, we would conclude that there are
almost 1.5 more diversification events of Coronaviridae that are related to ancient
codiversification rather than host switches. However, on average 20% of the inferred host
switches are time-inconsistent, including "back-in-time" host switches of >50 Myr (*SI Appendix*,
Fig. S8), which suggests instead that extent Coronaviridae originated recently and diversified by
frequent preferential host switches. 64% of the reconciliations found an origination of
coronaviruses within bats, in particular within the Pteropodidae family (Fig. 2A-C). We no longer
found an origination in bats when randomly shuffling the dataset (*SI Appendix*, Fig. S5,S6),
suggesting that this result is not artifactual. We checked the interpretation of our results by
simulating the two scenarios of (i) ancient origination in the ancestors of bats followed by
codiversification and (ii) recent origination in an extant bat species and a subsequent
diversification by preferential host switches. On the first set of simulations, ALE correctly inferred
an origination in bats and a few time-inconsistent switches (*SI Appendix*, Fig. S12). On the
second set, ALE correctly inferred an origination in bats, although with lower confidence, and a
large fraction (~20%) of time-inconsistent host switches, similar to what we observed for
Coronaviridae. These results therefore indicate a scenario of recent origination of coronaviruses
in bats followed by diversification by preferential host switches.
207
To investigate this scenario in more detail, we gradually applied a tree transformation to
the mammalian phylogeny, which excludes the possibility of an ancient origination happening
earlier than a given time. We found that we had to impose a very recent time of origination
(younger than 5 Myr) to obtain few time-inconsistent switches (*SI Appendix*, Table S1). We thus
carried out our follow-up analyses with a mammals' tree transformation (star phylogeny) that
assumes an origination in an extant mammalian lineage, such that coronavirus diversification is
explained entirely by host switches between extant mammalian species. Simulations validated
this approach in terms of properly inferring originations and identifying preferential host switches
(*SI Appendix*, Fig. S13). Applied to the data, the approach inferred a high probability of origination
in bats (56%, Fig, 2B-C, *SI Appendix*, Fig. S6) and a scenario of diversification by preferential
host switches: 68% of the inferred host switches happened within mammal orders (Fig 2D, *SI
Appendix*, Fig. S10), whereas we would expect on average only 28% of within-order host
switches if happening at random. We also inferred more-than-expected host switches between
closely related mammal orders (e.g. between Artiodactyla and Perissodactyla) and between the
order containing humans (Primates) and those of their domesticated animals, such as
Artiodactyla and Carnivora (*SI Appendix*, Fig. S10, Table S2). In contrast, host switches were five
times less numerous than expected by chance between bats and other orders (10.7%, against
50.2% on average if host switches were randomly distributed, Fig. 2D), in particular Artiodactyla
and Rodentia (*SI Appendix*, Fig. S11, Table S2). When occurring, host switches from bats often
occurred toward humans (1.9 host switches per reconciliation on average) or toward urban-living
and/or domesticated animals, such as rats, camels, or pigs (>1 host switch on average; *SI
Appendix*, Table S3). Host switches to humans occurred mostly from domesticated mammals
(camels, pigs, dogs), the house shrew and the house mouse, then followed by Asian palm civets,
and lastly by bats and other rodents (*SI Appendix*, Table S4). Results were consistent when
subsampling the dataset to have an equal sampling effort per host species, suggesting that our
results are not artifactually explained by the enhanced monitoring of coronaviruses in humans or
domesticated animals (*SI Appendix*, Supplementary Information Text). Finally, we found that
some sOTUs, in particular from Betacoronaviruses (e.g, u24667 and u175, both with humans

among their hosts), have experienced frequent host switches, whereas others have not (e.g.
u165, which is restricted to pigs). In particular, u944 (SARS-Cov-2) has experienced an
intermediate number of host switches compared to other coronaviruses (*SI Appendix*, Fig. S9).
239
We found qualitatively similar results when applying ALE on different sub-parts of the
palmprint region, suggesting that the potential occurrence of recombination does not bias our
conclusions (*SI Appendix*, Table S5). The percentage of originations inferred to occur in bats
decreased in the analyses on the first sub-part, probably because using such a short fragment
(75 aa-long) does not allow robust reconciliations. We also obtained consistent results using a
reconciliation method based on maximum parsimony (eMPRess) instead of maximum likelihood
(ALE). Whatever the costs that we set for the different reconciliation events, eMPRess estimated
significant reconciliations (p-values<0.01). For instance, when favoring host switches, we inferred
a recent origination in bats in 54% of the reconciliations and observed on average 32
cospeciations (s.d.   3), 2 losses (s.d.   1), 0.1 duplication (s.d.   0.3), and 140 host switches
(s.d.   3) including several "back-in-time" host switches of >30 Myr. eMPRess therefore also
supports a scenario of recent origination in bats and diversification by preferential host switches
(Fig. 1B). Without investigating the time-consistency of the host switches, we would have wrongly
concluded that almost one fourth of the diversification of Coronaviridae is related to ancient
cospeciation events.
255
An additional piece of evidence for a recent origination scenario comes from the
geographical distribution of coronaviruses, with a hotspot of diversity in Eurasia that has not
colonized the whole world (Fig. 3A, Fig. 1B). The coronavirus' hotspot is more strongly influenced
by the diversity of alphacoronaviruses than of betacoronaviruses (*SI Appendix*, Fig. S14). The
higher host switches rates and broader host range of betacoronaviruses is reflected in a more
widespread geographic distribution, with less pronounced hotspots when compared to
alphacoronaviruses (*SI Appendix*, Fig. S14). Mammalian hosts of coronaviruses have a hotspot
of species diversity concentrated in East Asia (Fig. 3C). The richness of coronaviruses presents a
similar pattern, but with two comparable hotspots of species diversity in East Asia and Southern
Europe (Fig. 3A), suggesting that the European hotspot is composed by fewer host species,
together carrying as diverse a set of coronaviruses as the Asian hotspot. Other regions with a
relatively high richness of coronaviruses and their hosts include parts of the African continent.
The Americas and Australia have relatively low richness of coronaviruses and their hosts.
Phylogenetic diversity of both hosts and coronaviruses (Fig. 3C,D) depict a similar pattern but
with phylogenetic diversity more evenly distributed across most world regions, including the
Americas.
272

273 **Discussion**
274
Together, our results suggest that the common ancestor of extant mammalian coronaviruses
originated recently in a bat species, and that coronaviruses diversification occurred via
preferential host switches rather than through codiversification with mammals. Although we
cannot unequivocally reject that ancestors of present-day coronaviruses were not present several
million years ago, we demonstrate that Coronaviridae is a highly dynamic clade in which
diversification operates through host switches at a much faster pace than that of their hosts.
sOTUs are rapidly replaced by newly-generated ones, with little role for codiversification with the
hosts. The high diversity and endemicity of coronaviruses among bats has led others to anticipate
that bats might be implicated in the origin of coronaviruses (2, 20, 23, 24), although definitive
proof was lacking. We provided evidence for that hypothesis using a probabilistic cophylogenetic
model after sampling the entire diversity of coronaviruses across mammals. Independent
evidence for coronavirus recent host switches among different species exists in the literature (41,
42). The envisioned scenario suggests a timing of origination for extant Coronaviridae that is
much more recent than the hundreds of millions of years ago suggested by (26). This is not

surprising given the difficulties in estimating divergence times and inferring branch lengths for
viral phylogenies (26, 27, 29), and provided that the dating of Wertheim et al. (26) relied on a
substitution rate estimated from data with limited temporal signal (~50 serially sampled
contemporary sequences of a short gene fragment, (23)).
293
Our results contradict previous suggestions that codiversification with vertebrate hosts
played an important role in Coronaviridae diversification (26, 28, 31). They also suggest that
previously reported cases of long-term codiversification in vertebrate RNA viruses have been
largely over-estimated, as many of them may instead be cases of diversification by host switches
occurring preferentially among closely-related hosts. Indeed, these two scenarios both generate
cophylogenetic signal in host-symbionts associations, such that cophylogenetic signal alone is
not evidence for long-term codiversification (34). In addition, under a scenario of recent
origination and preferential host switches, event-based cophylogenetic methods tend to
artifactually favor biologically unrealistic scenarios with codiversification and back-in-time host
switches, as we have shown here. As the time-consistency of host switches is typically not
investigated, this has remained unnoticed, and evidence for codiversification has been taken for
real. Ideally, cophylogenetic reconciliation methods would not allow such time-inconsistent host
switches. However, imposing time constraints in methods based on parsimony is NP-hard (43),
and the 'dated' version of ALE is not well adapted when recent host switch events dominate
evolutionary history. We have found two ways to get around the problem, by interpreting time-
inconsistent host switches as evidence for recent preferential host switches, and by gradually
transforming the host tree to avoid large back-in-time switches, however future efforts should
focus on developing time-consistent cophylogenetic methods. This would allow more robust and
precise inferences of host-virus (and more generally host-symbiont) evolutionary history.
313
During their evolution, coronavirus' host switches occurred more frequently within than
between mammalian orders. This suggests that mammalian characteristics shared between
relatives (e.g., genetic, behavioral, ecological), and the frequency of encounters among hosts
play important roles in determining coronavirus' host switches. Additionally, between-order host
switches occurred more frequently among non-flying mammals and among orders containing
humans and urban and domesticated mammals, suggesting that contact frequency alone is likely
a key characteristic in host switches. Accordingly, amongst the most-likely host switches towards
humans were those coming from mammals suspected to be involved in the transfer of specific
coronavirus sOTUs likely through contact, for instance, camels in the case of MERS-CoV (41),
Asian palm civets with SARS-CoV (16, 17) and the house mouse with SARS-CoV2 (42).
Importantly, we found that host switches from bats to other mammalian species were rare during
the evolutionary history of Coronaviridae, even though coronaviruses originated and are more
diverse within bats. These pieces of evidences suggest that bats are a closed reservoir of the
Coronaviridae diversity.
328
Spillovers from bats to non-bat species, when they occurred, were found more likely to be
towards humans than to any other mammalian species, suggesting humans may have acted as
evolutionary intermediate hosts amongst mammals. From an ecological perspective, the large
abundance and widespread geographic distribution of humans, together with our habits of forcing
contact with other species, including bats, make it unsurprising that humans, among all mammal
species, have acted as intermediate hosts of ancestral forms of coronaviruses. Interestingly, for
some individual species of coronaviruses, such as the SARS-CoV2 and other SARS-like
coronaviruses, the dominant hypothesized scenario is that precursor forms spread from a bat to
another intermediate mammalian host before infecting humans (20, 44). Our molecular marker
lacks the intra-OTU resolution necessary to make species-level predictions, but our results
suggest that more ancient coronaviruses host switches may have occurred in the other direction:
from bats to humans to non-bat mammals. Many human activities lend credit to the human-as-
evolutionarily-intermediate-host-hypothesis, including human excursions to bat caves (45),
hunting (46), and habitat destruction and modification (47), all of which increase the contact

between bats and humans and their domesticated animals (47). Conservation of bats' natural
habitats, away from human contact, could help avoiding further spreads of coronaviruses among
humans.
346
Insights of past and future host switches are gained from coronavirus geographic
distribution. Coronaviruses are found worldwide and their hotspots of diversity are concentrated in
East Asia and Southern Europe, where they likely originated. Previous assessments of the
diversity of bat hosts of betacoronaviruses suggested similar hotspots but with a distribution of
coronaviruses more concentrated in the hotspots (7, 14, 15) than the more pervasive pattern we
found using all mammalian hosts. Moreover, the distribution of coronaviruses is less concentrated
in the hotspots when phylogenetic metrics of diversity are included, suggesting that species
richness alone is masking the global evolutionary potential of these viruses (48). Coronaviruses'
likely recent origination in bats, high within-order transmission rates, and their capacity to switch
between mammal orders in some cases suggest the potential for future fast spreading and
increase in the number of species across most world regions. Among alphacoronaviruses, the
spread is more likely to remain concentrated within bats, while betacoronaviruses have a higher
potential for among-orders spreading and infection of new mammalian hosts. The
betacoronaviruses lineages already detected in humans have especially high host generalism
and transmission rates, indicating that these lineages should be particularly monitored to avoid
future pandemics.
363
Finally, a few important limitations of our analyses deserve to be mentioned.
Recombination is an important mechanism of viral evolution (49), and approaches more
adequately designed to investigate the role of recombination are needed. The fact that different
subparts of the palmprint region lead to similar results indicates that recombination acting on the
palmprint region is unlikely to bias our conclusions. However, looking at other genomic regions
would allow gaining a more complete understanding of the role of recombination in coronavirus
evolution. Lastly, because the palmprint region is a conserved region, we could not reconstruct
the recent evolutionary history of coronaviruses (i.e. the within sOTU transmission dynamic):
combining the palmprint region with a fast-evolving region(s) would enable more precise
estimates of the recent routes of coronaviruses' transmission, including that of SARS-CoV-2.
374
Understanding the evolutionary origins and diversification of viruses is crucial to any
attempt of predicting new transmission routes, yet the relative frequencies of virus–host
cospeciation versus cross-species transmission in the evolution of vertebrate RNA viruses
remains uncertain (31). We found that coronaviruses originated in bats where they are more
diverse nowadays, and later diversified in other mammal orders through preferential host
switches. Spillovers from bats were rare but likely human-induced, suggesting humans are the
intermediate evolutionary bridge that facilitated the spread of coronaviruses across mammals.
Host switches between primates and artiodactyls, perissodactyls, and carnivorans happen at high
rates and we can thus expect a spread of coronaviruses amongst new mammalian hosts and
outside of their current diversity hotspots in East Asia and Europe, as well as future
coronaviruses-related pandemics. Our results suggest reducing human-bat contact, for example
by conserving bat habitats, as a mitigation strategy. They also suggest that cases of long-term
virus–host codiversification, reported on the basis of cophylogenetic tests, have been largely
over-estimated.
389

390 **Materials and Methods**
391
Operational Taxonomic Units for Coronaviridae

We used the 46 described species-like Operational Taxonomic Units (sOTUs) for Coronaviridae
delimited using 'palmprint' sequences by (36, 37). The palmprint is a conserved amino acid (aa)

sub-sequence (150 aa in Coronaviridae) of central importance in the viral RdRp (36), selected for
its homology across the large majority of sequences, allowing estimation of sequence divergence
and phylogenetic trees (37). sOTUs were identified by Edgar et al. (36) after clustering palmprint
sequences at 90% amino acid identity; and released through the Serratus project. Their approach
is equivalent to the species delimitation proposed by the International Committee on the
Taxonomy of Viruses for Coronaviridae ((2); *SI Appendix*, Supplementary Information Text),
which suggests 90% similarity of amino acid sequences for conserved domains (2, 37), and are,
therefore, ideal for species tree construction. Under the 'palmprint framework' a centroid definition
of species is applied to characterize a new OTU when a threshold of 90% amino acid identity is
surpassed, serving as a useful taxonomic barcode (37). We downloaded the palmprint amino acid
sequences of Coronaviridae sOTUs from the Serratus project (https://serratus.io/; (36)) on April
13 of 2022.
407

Mammalian hosts of Coronaviridae

All 46 sOTUs of Coronaviridae with a full palmprint and associated data in the NCBI database
were screened for the identification of its hosts. From those, 35 sOTUs were associated with
mammalian hosts and were kept for downstream analyses. Serratus' associated metadata was
used to identify GenBank accession codes linked to each sOTU. The complete set of 90,540
associated GenBank accession codes was screened to obtain the host information for each
sOTU (on NCBI, Features>source>/host=). All the host species with a full Linnean name were
kept as such. Accession codes with hosts leading to a generic level information were further
inspected to identify the associated publication and determine the complete species name.
Dubious cases or accession codes without publications had their hosts disregarded. Common
names or high-level host information (e.g., host="bats") were generally eliminated except in a few
cases where a domesticated species was found to be the host (i.e., host="dog","canine" were
Canis lupus; host="cat","feline" were Felis catus; host="pig","piglet","newborn piglet","sucking
piglet","porcine","swine" were Sus scrofa). A final dataset of 116 mammalian hosts associated
with the 35 sOTUs was assembled and used in downstream analyses. A matrix with the
association between Coronaviridae sOTUs and mammalian species is available in *SI Appendix*,
Dataset S1.
425

Coronaviridae phylogenetic trees

We constructed a Coronaviridae tree using the palmprint amino acid sequence information of the
35 sOTUs. We aligned the amino acid sequences with MAFFT (50) and trimmed them with trimAl
(51). The final alignment contained 150 amino acid positions. We used two main phylogenetic
software, BEAST2 (52) and PhyloBayes (53), both to assess the robustness of the tree to the
phylogenetic method, and because they have different advantages (e.g. rooting and time
calibration are performed in BEAST2 while PhyloBayes outputs are adapted to the
cophylogenetic algorithm we used). We visualized phylogenetic trees using R (54).
In order to run BEAST2, we generated an input file using BEAUti with the following
parameters: a WAG model with 4 classes of rates and invariant sites, a birth-death prior, and a
relaxed log-normal clock. BEAST2 sampled a posterior distribution of ultrametric trees using
Markov chain Monte Carlo (MCMC) with 4 independent chains each composed of 100,000,000
steps sampled every 10,000 generations. We checked the convergence of the 4 chains using
Tracer (55). We used LogCombiner to merge the results setting a 25% burn-in and TreeAnnotator
to obtain a Maximum Clade Credibility (MCC) tree with median branch lengths. PhyloBayes was
run using an LG model, 4 classes of rates, and a chain composed of 4,000 steps with a 25%
burn-in.
To further assess the robustness of the BEAST2 tree rooting, we estimated the root
position on a 46-sOTU maximum likelihood tree (from the Serratus project - (36)) assuming a

strict molecular clock and an ultrametric tree. We used an ultrametric setting as temporary
information from the tip dates (ranging between 1999 and 2022, a neglectable difference with
respect to the root age of dozens of thousands or even millions of years) was not sufficient to
infer the mutation rate (we assessed the temporal signal with TempEst, (56)). We performed
rooting and time-scaling with LSD2 (v2.3, (57)), assuming a tree of unknown scale (e.g. fixing all
the tips dates to 1 and the root date to 0) with outlier removal and root search on all branches.
LSD2 detected no outliers and positioned the root on the same branch as in the BEAST2 MCC
tree (between alpha and betacoronaviruses).
453

Mammalian phylogenetic tree

455 We obtained a phylogenetic hypothesis for mammals from the consensus DNA-only tree of (58),
456 one of the most complete and updated phylogenies for mammals. We downloaded the node-
457 dated tree for 4,098 mammals, constructed based on a 31-gene supermatrix, from the VertLife
458 website (http://vertlife.org/data/mammals/). We used a pruned version of the tree with the 116
459 mammalian hosts of Coronaviridae in all analyses in this paper.
460

Phylogenetic signal in the association between coronaviruses and mammals

To assess whether closely related coronaviruses interact with similar mammals, and vice-versa,
i.e. presence of phylogenetic signal in the association, we used Mantel tests following (38).
Mantel tests were constructed by taking the Pearson correlation between phylogenetic distances
and ecological distances. Phylogenetic distances of coronaviruses were computed on the
BEAST2 MCC phylogeny. Ecological distances were calculated based on the interaction network
matrix containing the association between coronavirus' sOTUs and mammals, accounting for the
evolutionary relationships among interaction partners using UniFrac distances (59). Firstly, we
conducted Mantel tests permuting the identity of species but keeping the number of partners per
species constant; this allows for assessing the effect of species identity while controlling for the
confounding effect of the number of partners. Then, we evaluated the phylogenetic signal in the
number of partners alone. Lastly, we calculated clade-specific Mantel tests for sub-networks
containing at least 10 species (38) to evaluate whether phylogenetic signal was stronger for
specific subclades of mammals or coronaviruses. Ten thousand permutations were used in each
analysis to assess significance. Analyses were conducted using the phylosignal_network and
phylosignal_sub_network functions in the R package RPANDA (60).
477

Coronaviridae origination and host switches

We used the amalgamated likelihood estimation (ALE - (39)) to reconciliate the mammal and
coronaviruses evolutionary history using events of cospeciations, host switches, duplications, and
losses. Originally designed in the context of gene tree – species tree reconciliations (39), ALE
has also been particularly useful in the context of host-symbiont cophylogenetic analyses as it
considers both phylogenetic uncertainty of the symbiont evolutionary history and undersampling
of host species (35, 61, 62). ALE indeed assumes that host switches may imply an unsampled or
extinct intermediate host lineage (40). We ran ALE with the posterior distribution of phylogenetic
trees of coronaviruses generated with PhyloBayes to estimate the maximum likelihood rates of
host switches, duplications, and losses of the coronaviruses. We first tried running the "dated"
version of ALE, which accounts for the order of branching events in the host phylogeny, therefore
only allowing for time-consistent host switches (i.e. host switches that happen between two
contemporary host lineages). However, this led to unrealistic parameter estimates (such as very
high loss rates) and ALE was not able to output possible reconciliations, suggesting that the
mammalian and Coronaviridae trees are too incongruent to be reconciled with only time-

consistent host switches. We therefore used the "undated" version of ALE that only exploits the
topology of both the host and the symbiont tree and thus does not constrain the host switches to
be time-consistent. ALE generated a total of 5,000 reconciliations, from which we extracted the
mean number of cospeciations, host switches, duplications, and losses. We also reported the
likely origination of coronaviruses in mammals (i.e. the branch in the mammal phylogeny that was
first infected by coronaviruses) by computing, for each branch of the mammalian tree, the
frequency of reconciliations (among the 5,000) that supported an origination in that branch. If a
reconciliation requires more cospeciation events and fewer host switch events, than expected
under a null scenario of independent evolution, this indicates that the evolution of the symbiont
was not independent of that of the host, and in this case, we talk about a "significant
reconciliation" (63). We evaluated the significance of the reconciliation by comparing the
estimated number of cospeciation and host switch events to null expectations obtained with ALE
by shuffling the mammal host species across the mammal tree, both randomly or within major
biogeographic regions according to the proposal of regions by (64) for mammals (six
biogeographic regions: North American, South American, African, Eurasian, Oriental, and
Australian). We considered a reconciliation to be significant if the observed number of
cospeciations was higher than 95% of the null expectations and if the number of host switches
was lower than 95% of the null expectations (35). The likeliness of a host switch between two
mammal lineages is measured as the frequency of the reconciliations in which it occurs. Finally,
we reported the ratio of time-inconsistent host switches by focusing on "back-in-time" switches,
from a donor mammal lineage to an older receiver mammal lineage that never coexisted.

Because ALE estimated a large proportion of time-inconsistent host switches (see
Results), we first tested the scenario of a more recent origination by collapsing all mammalian
nodes anterior to X Myr into a polytomy at the root of the phylogeny (with X varying from 55 Myr
to 5 Myr), such that the coronavirus origination and host switches inferred by ALE could not
involve mammal lineages older than X Myr. Second, we investigated the scenario of
diversification by pure preferential host switches of the coronaviruses among extant mammals. To
do so, we ran ALE on a star mammalian phylogenetic tree. In this context, ALE could no longer
infer cospeciations, and only fit events of host switches, duplications, or losses. When inferring a
likely host switch between two specific mammalian lineages on a star phylogeny, there are often
as many reconciliations suggesting one directionality of the host switch (i.e. from one of the
lineages to the other) as the other. We then only kept host switches present in at least 10% of the
reconciliations and looked at the ratio between the number of host switches that were estimated
within versus between mammal orders. We compared this ratio to a null expectation obtained by
randomly shuffling the host mammal species.

Recombination is frequent in viruses and the palmprint region may be recombined, such
that different fragments of the palmprint region may have different evolutionary histories,
potentially biasing our inference. To test whether the results we obtained on the whole 150-amino
acid palmprint region were not impacted by recombination, we replicated the ALE analyses on
two sub-regions: the first part (positions 1-75) and the last part (positions 76-150).

Finally, we repeated our cophylogenetic analyses using eMPRess (43), another event-
based cophylogenetic approach that reconciliates host-symbiont evolutionary histories using
maximum parsimony. eMPRess is a recent improved version of the popular Jane approach (32);
it differs from Jane especially by not only relying on a heuristic and therefore guarantying that the
solution truly corresponds to the maximum parsimony reconciliation(s) (43). However, contrary to
Jane, eMPRess does not allow the same symbiont species to be present in different host species,
and does not offer the possibility to constrain host switches to occur only among lineages from
pre-specified time periods. eMPRess requires specifying cost values for the events of host
switches (t), duplications (d), and losses (l). We tested two sets of cost values: (1) cost values
that disadvantage host switches (d=6, t=6, l=1) and (2) uniform cost values that favor host
switches (d=1, t=1, l=1). As with ALE, we evaluated the significance of the reconciliations using

permutations. We ran eMPRess analyses on a set of 50 trees randomly sampled from the
posterior distribution of PhyloBayes.
549

Simulation analyses

By running the undated version of ALE either on the mammal phylogeny or a star phylogeny, we
proposed a framework to evaluate whether the cophylogenetic pattern is due to a history of
ancient codiversification (i.e. a mix of cospeciations, host switches, duplications, and losses; Fig.
1A) or to a scenario where the coronaviruses diversify more recently by preferential host switches
((34); Fig. 1B). To validate the interpretation of our ALE results, we performed simulations under
the two alternative scenarios of codiversification and diversification by preferential host switches.
For the scenario of codiversification, we assumed that coronaviruses originated in the ancestors
of bats and that they subsequently codiversified with the mammals by experiencing events of
cospeciations, host switches, duplications, and losses. We used the function *sim_microbiota* in
the R-package HOME to obtain the corresponding coronavirus sequences and coronavirus-
mammals associations (65) (*SI Appendix*, Supplementary Information Text). For the scenario of
coronaviruses diversification by preferential host switches, we used a birth-death model (pbtree
function in the R-package phytools) to simulate a phylogenetic tree of the coronaviruses: in our
model, each coronavirus lineage is associated with a single host species, a birth event
corresponds to a host switch (at rate 50), while a death event corresponds to a loss of a
coronavirus in a host lineage (at rate 5). We started the diversification by assuming a single
coronavirus infection in *Eidolon helvum* (a bat host of external lineages within Betacoroviruses,
u25738 and u27845). Then, following de (66) and (67), we modeled preferential host switches by
assuming that for a host switch from a given donor mammal species, each potential receiver
species has a probability proportional to exp(-0.035*d) where d is the phylogenetic distance
between the donor and receiver species. Finally, we simulated DNA sequences of the
coronavirus sequences using the function *simulate_alignment* in HOME. For each type of
simulation, we generated 50 simulated datasets of mammal-coronavirus associations. For each
dataset, we ran PhyloBayes and ALE on both the mammalian phylogeny and the star phylogeny.
575

Geographic distribution of Coronaviridae

577 We downloaded geographic range maps for each mammalian host species, with the exception of
578 *Homo sapiens*, from the Map of Life website (https://mol.org/species/); see (68). These maps
579 follow the taxonomy of the Mammal Diversity Database (69) supplemented with the Handbook of
580 the Mammals of the World (HWM) database and the Alien Checklist database for invasive
581 species (68).
582

We created a world map with hexagonal, equal-area grid cells of 220km on which we
mapped host and coronavirus species diversity, using the Mollweide world projection to
accurately represent areas. At large spatial scales, cells with ~220km resolution return more
reliable diversity estimates than smaller cells (70). We considered that a host species was
present in any given cell if its range covered at least 30% of the cell area to avoid overestimating
diversity. We calculated host species diversity as a simple sum of the species occurring in any
given cell, and host phylogenetic diversity as Faith's phylogenetic diversity index (PD - (71)) for
each cell. We mapped Coronaviridae diversity using the host-filling method (72): we constructed
a range map for each Coronaviridae sOTU by overlapping the range maps of all its hosts. We
consider the host filling method appropriate in this case because coronaviruses are obligatory
parasites that can only live inside hosts. Next, we calculated Coronaviridae sOTU diversity by
summing range maps overlapping on each cell, and Coronaviridae phylogenetic diversity as
Faith's PD (71). We created these maps in R (54) using the packages epm (73), sf (74), and ape
(75).

Acknowledgments

This work was performed using HPC resources from GENCI-IDRIS (Grants 2021-A0100312405 and 2022- AD010313735). The authors thank B. Boussau and the Morlon lab for helpful discussions. RM thanks Campus France and the One Health/Make Our Planet Again Program for a fellowship to conduct this research.

References

1. K. P. Alekseev, et al., Bovine-Like Coronaviruses Isolated from Four Species of Captive Wild Ruminants Are Homologous to Bovine Coronaviruses, Based on Complete Genomic Sequences. J Virol 82, 12422–12431 (2008).
2. R. J. de Groot, et al., Family: Coronaviridae. ICTV Ninth Report. https://ictv.global/report_9th/RNApos/Nidovirales/Coronaviridae (2022).
3. A. M. Q. King, Adams M.J., Carstens E.B., Lefkowitz E.J., Virus taxonomy: classification and nomenclature of viruses: Ninth Report of the International Committee on Taxonomy of Viruses (Elsevier, 2012).
4. J. C. Leao, et al., Coronaviridae—Old friends, new enemy! Oral Dis 28, 858–866 (2022).
5. M. Wardeh, M. Baylis, M. S. C. Blagrove, Predicting mammalian hosts in which novel coronaviruses can be generated. Nat Commun 12 (2021).
6. C. M. Zmasek, E. J. Lefkowitz, A. Niewiadomska, R. H. Scheuermann, Genomic evolution of the Coronaviridae family. Virology 570, 123–133 (2022).
7. D. J. Becker, et al., Optimising predictive models to prioritise viral discovery in zoonotic reservoirs. Lancet Microbe 3, e625–e637 (2022).
8. C. Drosten, et al., Identification of a Novel Coronavirus in Patients with Severe Acute Respiratory Syndrome. New England Journal of Medicine 348, 1967–1976 (2003).
9. J. Peiris, et al., Coronavirus as a possible cause of severe acute respiratory syndrome. The Lancet 361, 1319–1325 (2003).
10. A. M. Zaki, S. van Boheemen, T. M. Bestebroer, A. D. M. E. Osterhaus, R. A. M. Fouchier, Isolation of a Novel Coronavirus from a Man with Pneumonia in Saudi Arabia. New England Journal of Medicine 367, 1814–1820 (2012).
11. P. Zhou, et al., A pneumonia outbreak associated with a new coronavirus of probable bat origin. Nature 579, 270–273 (2020).
12. H. Ritchie, et al., Coronavirus Pandemic (COVID-19). Published online at OurWorldInData.org (2022).
13. P. C. Y. Woo, Y. Huang, S. K. P. Lau, K. Y. Yuen, Coronavirus genomics and bioinformatics analysis. Viruses 2, 1805–1820 (2010).
14. S. J. Anthony, et al., Global patterns in coronavirus diversity. Virus Evol 3 (2017).
15. N. F. R. Munoz, et al., The coevolutionary mosaic of bat betacoronavirus emer-gence risk. EcoEvoRXiv (2022).
16. Y. Guan, et al., Isolation and characterization of viruses related to the SARS coronavirus from animals in Southern China. Science (1979) 302, 276–278 (2003).
17. C. SMEC, Molecular Evolution of the SARS Coronavirus During the Course of the SARS Epidemic in China. Science (1979) 303, 1666–1669 (2004).
18. R. L. Graham, R. S. Baric, Recombination, Reservoirs, and the Modular Spike: Mechanisms of Coronavirus Cross-Species Transmission. J Virol 84, 3134–3146 (2010).
19. R. J. de Groot, et al., Commentary: Middle East Respiratory Syndrome Coronavirus (MERS-CoV): Announcement of the Coronavirus Study Group. J Virol 87, 7790–7792 (2013).

20.   V. M. Corman, D. Muth, D. Niemeyer, C. Drosten, "Hosts and Sources of Endemic
Human Coronaviruses" in Advances in Virus Research, (Academic Press Inc., 2018), pp.
163–188.
21.   E. Mavrodiev, M. L. Tursky, S. Vincent, D. M. Williams, L. Schroder, On Classication and
Taxonomy of Coronaviruses (Riboviria, Nidovirales, Coronaviridae) with Special Focus on
Severe Acute Respiratory Syndrome-Related Coronavirus 2 (SARS-CoV-2) (2021)
https:/doi.org/10.21203/rs.3.rs-282371/v1.
22.   P. C. Y. Woo, S. K. P. Lau, Y. Huang, K. Y. Yuen, Coronavirus diversity, phylogeny and
interspecies jumping. Exp Biol Med 234, 1117–1127 (2009).
23.   P. C. Y. Woo, et al., Discovery of Seven Novel Mammalian and Avian Coronaviruses in
the Genus Deltacoronavirus Supports Bat Coronaviruses as the Gene Source of
Alphacoronavirus and Betacoronavirus and Avian Coronaviruses as the Gene Source of
Gammacoronavirus and Deltacoronavirus. J Virol 86, 3995–4008 (2012).
24.   D. Vijaykrishna, et al., Evolutionary Insights into the Ecology of Coronaviruses. J Virol 81,
4012–4020 (2007).
25.   A. J. Drummond, S. Y. W. Ho, M. J. Phillips, A. Rambaut, Relaxed Phylogenetics and
Dating with Confidence. PLoS Biol 4, e88 (2006).
26.   J. O. Wertheim, D. K. W. Chu, J. S. M. Peiris, S. L. Kosakovsky Pond, L. L. M. Poon, A
Case for the Ancient Origin of Coronaviruses. J Virol 87, 7039–7045 (2013).
27.   J. O. Wertheim, S. L. Kosakovsky Pond, Purifying selection can obscure the ancient age
of viral lineages. Mol Biol Evol 28, 3355–3365 (2011).
28.   D. T. S. Hayman, M. A. Knox, Estimating the age of the subfamily Orthocoronavirinae
using host divergence times as calibration ages at two internal nodes. Virology 563, 20–
27 (2021).
29.   S. Duchêne, E. C. Holmes, S. Y. W. Ho, Analyses of evolutionary dynamics in viruses
are hindered by a time-dependent bias in rate estimates. Proceedings of the Royal
Society B: Biological Sciences 281, 20140732 (2014).
30.   A. Katzourakis, R. J. Gifford, Endogenous Viral Elements in Animal Genomes. PLoS
Genet 6, e1001191 (2010).
31.   M. Shi, et al., The evolutionary history of vertebrate RNA viruses. Nature 556, 197–202
(2018).
32.   C. Conow, D. Fielder, Y. Ovadia, R. Libeskind-Hadas, Jane: A new tool for the
cophylogeny reconstruction problem. Algorithms for Molecular Biology 5 (2010).
33.   Y.-Z. Zhang, W.-C. Wu, M. Shi, E. C. Holmes, The diversity, evolution and origins of
vertebrate RNA viruses. Curr Opin Virol 31, 9–16 (2018).
34.   D. M. de Vienne, et al., Cospeciation vs host-shift speciation: methods for testing,
evidence from natural associations and relation to coevolution. New Phytologist 198, 347–
385 (2013).
35.   B. Perez-Lamarque, H. Morlon, Comparing different computational approaches for
detecting long-term vertical transmission in host-associated microbiota in Molecular
Ecology, (John Wiley and Sons Inc, 2022) https:/doi.org/10.1111/mec.16681.
36.   R. C. Edgar, et al., Petabase-scale sequence alignment catalyses viral discovery. Nature
602, 142–147 (2022).
37.   A. Babaian, R. Edgar, Ribovirus classification by a polymerase barcode sequence. PeerJ
10 (2022).
38.   B. Perez-Lamarque, et al., Do closely related species interact with similar partners?
Testing for phylogenetic signal in bipartite interaction networks. Peer Community Journal
2, XX (2022).
39.   G. J. Szöllosi, W. Rosikiewicz, B. Boussau, E. Tannier, V. Daubin, Efficient exploration of
the space of reconciled gene trees. Syst Biol 62, 901–912 (2013).
40.   G. J. Szöllosi, E. Tannier, N. Lartillot, V. Daubin, Lateral gene transfer from the dead.
Syst Biol 62, 386–397 (2013).
41.   G. Dudas, L. M. Carvalho, A. Rambaut, T. Bedford, MERS-CoV spillover at the camel-
human interface. Elife 7 (2018).

42.   C. Wei, et al., Evidence for a mouse origin of the SARS-CoV-2 Omicron variant. Journal
of Genetics and Genomics 48, 1111–1121 (2021).

43.   S. Santichaivekin, et al., eMPRess: a systematic cophylogeny reconciliation tool.
Bioinformatics 37, 2481–2482 (2021).

44.   C. C. S. Tan, et al., Transmission of SARS-CoV-2 from humans to animals and potential
host adaptation. Nat Commun 13 (2022).

45.   N. M. Furey, P. A. Racey, "Conservation Ecology of Cave Bats" in Bats in the
Anthropocene: Conservation of Bats in a Changing World, (Springer International
Publishing, 2016), pp. 463–500.

46.   T. Mildenstein, I. Tanshi, P. A. Racey, "Exploitation of Bats for Bushmeat and Medicine"
in Bats in the Anthropocene: Conservation of Bats in a Changing World, (Springer
International Publishing, 2016), pp. 325–375.

47.   I. Smith, L. F. Wang, Bats and their virome: An important source of emerging viruses
capable of infecting humans. Curr Opin Virol 3, 84–91 (2013).

48.   S. Leopardi, et al., Interplay between co-divergence and cross-species transmission in
the evolutionary history of bat coronaviruses. Infection, Genetics and Evolution 58, 279–
289 (2018).

49.   M. Pérez-Losada, M. Arenas, J. C. Galán, F. Palero, F. González-Candelas,
Recombination in viruses: Mechanisms, methods of study, and evolutionary
consequences. Infection, Genetics and Evolution 30, 296–307 (2015).

50.   K. Katoh, D. M. Standley, MAFFT Multiple Sequence Alignment Software Version 7:
Improvements in Performance and Usability. Mol Biol Evol 30, 772–780 (2013).

51.   S. Capella-Gutiérrez, J. M. Silla-Martínez, T. Gabaldón, trimAl: a tool for automated
alignment trimming in large-scale phylogenetic analyses. Bioinformatics 25, 1972–1973
(2009).

52.   R. Bouckaert, et al., BEAST 2: A Software Platform for Bayesian Evolutionary Analysis.
PLoS Comput Biol 10, e1003537 (2014).

53.   N. Lartillot, H. Philippe, A Bayesian Mixture Model for Across-Site Heterogeneities in the
Amino-Acid Replacement Process. Mol Biol Evol 21, 1095–1109 (2004).

54.   R Core Team, R: a language and environment for statistical computing (2018).

55.   A. Rambaut, A. J. Drummond, D. Xie, G. Baele, M. A. Suchard, Posterior Summarization
in Bayesian Phylogenetics Using Tracer 1.7. Syst Biol 67, 901–904 (2018).

56.   A. Rambaut, T. T. Lam, L. Max Carvalho, O. G. Pybus, Exploring the temporal structure
of heterochronous sequences using TempEst (formerly Path-O-Gen). Virus Evol 2,
vew007 (2016).

57.   T.-H. To, M. Jung, S. Lycett, O. Gascuel, Fast Dating Using Least-Squares Criteria and
Algorithms. Syst Biol 65, 82–97 (2016).

58.   N. S. Upham, J. A. Esselstyn, W. Jetz, Inferring the mammal tree: Species-level sets of
phylogenies for questions in ecology, evolution, and conservation. PLoS Biol 17 (2019).

59.   J. Chen, et al., Associating microbiome composition with environmental covariates using
generalized UniFrac distances. Bioinformatics 28, 2106–2113 (2012).

60.   H. Morlon, et al., RPANDA: an R package for macroevolutionary analyses on
phylogenetic trees. Methods Ecol Evol 7, 589–597 (2016).

61.   M. Groussin, et al., Unraveling the processes shaping mammalian gut microbiomes over
evolutionary time. Nat Commun 8 (2017).

62.   M. Bailly-Bechet, et al., How Long Does Wolbachia Remain on Board? Mol Biol Evol 34,
1183–1193 (2017).

63.   R. G. Dorrell, et al., Phylogenomic fingerprinting of tempo and functions of horizontal
gene transfer within ochrophytes https:/doi.org/10.1073/pnas.2009974118/-
/DCSupplemental.

64.   C. Barry Cox, The biogeographic regions reconsidered. J Biogeogr 28, 511–523 (2001).

65.   B. Perez-Lamarque, H. Morlon, Characterizing symbiont inheritance during host–
microbiota evolution: Application to the great apes gut microbiota. Mol Ecol Resour 19,
1659–1671 (2019).

66.   D. M., de Vienne, T. Giraud, J. A. Shykoff, When can host shifts produce congruent host
and parasite phylogenies? A simulation approach. J Evol Biol 20, 1428–1438 (2007).
67.   B. Perez-Lamarque, H. Krehenwinkel, R. G. Gillespie, H. Morlon, Limited evidence for
microbial transmission in the phylosymbiosis between Hawaiian spiders and their
microbiota. mSystems 7, e01104-21 (2022).
68.   C. J. Marsh, et al., Expert range maps of global mammal distributions harmonised to
three taxonomic authorities. J Biogeogr 49, 979–992 (2022).
69.   C. J. Burgin, J. P. Colella, P. L. Kahn, N. S. Upham, How many species of mammals are
there? J Mammal 99, 1-14 (2018).
70.   A. H. Hurlbert, W. Jetz, Species richness, hotspots, and the scale dependence of range
maps in ecology and conservation. PNAS 104, 13384-13389 (2007).
71.   D. P. Faith, Conservation evaluation and phylogenetic diversity. Biol Cons 61, 1-10
(1992).
72.   P. Pappalardo, et al., Comparing methods for mapping global parasite diversity. Global
Ecol Biogeogr 29, 182-193 (2019).
73.   P. O. Title, D. L. Swiderski, M. L. Zelditch, EcoPhyloMapper: an R package for integrating
geographical ranges, phylogeny and morphology. Methods Ecol Evol 13, 1912-1922
(2022).
74.   E. Pebesma, Simple features for R: standardized support for spatial vector data. The R
Journal 10, 439-446 (2018).
75.   E. Paradis, K. Schliep, ape 5.0: an environment for modern phylogenetics and
evolutionary analyses in R. Bioinformatics 35, 526-528 (2019).
775

Figures and Tables

Figure 1. A framework for testing scenarios of virus-host evolution, illustrated with the example of Coronaviridae and their mammalian hosts: In (A), a scenario of ancient origination and codiversification; in (B) a scenario of recent origination and diversification by preferential host switches; and in (C) a scenario of independent evolution. For each scenario, we indicate the associated predictions in the grey boxes. Contrary to scenario C, both scenarios A and B are expected to generate a cophylogenetic signal, *i.e.* closely-related coronaviruses tend to infect closely-related mammals, resulting in significant reconciliations when using topology-based probabilistic cophylogenetic methods, such as the undated version of ALE (Szollozi et al 2013), Jane (Conow et al 2010), or eMPRess (Sanitchaivekin et al, 2021). However, we expect scenario B to be distinguishable from scenario A in terms of the time consistency of host-switching events. Under scenario B, cophylogenetic methods wrongly estimate a combination of cospeciations and "back-in-time" host switches (see Methods & Results). We also expect different biogeographic patterns under the different scenarios.

Figure 2. Species-level relationships among coronaviruses and their associated mammalian hosts. A consensus Maximum Clade Credibility phylogenetic tree of coronaviruses is shown on the left. The tree was constructed with BEAST2 based on 150-aa palmprint amino acid sequences of the RdRp gene. sOTUs of Coronaviridae followed the definition of the Serratus project. The putative location of four genera of coronaviruses, Beta, Gamma, Delta, and Alphacoronaviruses, is shown. Bar scale is in units of aa substitution. On the right, a barplot gives the number of total mammalian host species and the number of host species by main mammalian order. Ancestral states on the left were obtained for illustrative purposes with the make.simmap function of the phytools R package (Revell 2012). Mammal silhouettes taken from open-to-use sources in phylopic.org, detailed credits given in SI Appendix Table S6.

804
805

Figure 3. The origination of coronaviruses in mammals is estimated among bats, which
tend to form a closed reservoir. (A) Phylogenetic tree of the mammals with branches colored
as the percentage of ALE reconciliations which inferred this branch or its ancestral lineages as
the origination of coronaviruses in mammals. Red branches are likely originations, whereas blue
branches are unlikely. (B) Boxplots recapitulating the probability of inferred origination per branch
in bats *versus* other mammal orders, with ALE applied on the original mammal tree (left panel) or
on the mammal tree transformed into a star phylogeny (right panel), therefore assuming an
origination in extant species. (C) Distributions of the percentages of host switches occurring
within mammalian orders (left panel) and between-orders involving bats (right panel). Observed
values (in orange) are compared to null expectations if host switches were happening at random
(in grey). Mammal silhouettes taken from open-to-use sources in phylopic.org, detailed credits
given in SI Appendix Table S6.

821
Figure 4. Maps of the diversity of coronaviruses and their mammal hosts. In A) the richness of species of coronaviruses; geographic range maps of coronaviruses were constructed after applying the host-filling method on the geographic range maps of mammalian hosts of coronaviruses. In B) Faith's (1992) phylogenetic diversity of coronaviruses, calculated using the phylogenetic tree of coronaviruses (see main text). In C) and D), the richness and phylogenetic diversity of mammal hosts of coronaviruses, respectively. All maps are on the Mollweide projection.

829
830