Impaired potency of neutralizing antibodies against cell-cell fusion mediated by SARS-CoV-2

Qian Wang[a,#], Andre Yanchen Yeh [a,b,#], Yicheng Guo[a], Hiroshi Mohri[a], Jian Yu[a], David D. Ho[a,c,d], Lihong Liu[a, d,*]

[a]*Aaron Diamond AIDS Research Center, Columbia University Vagelos College of Physicians and Surgeons, New York, USA;* [b]*School of Medicine, National Taiwan University, Taipei, Taiwan;* [c]*Department of Microbiology and Immunology, Columbia University Vagelos College of Physicians and Surgeons, New York, NY, USA;* [d]*Division of Infectious Diseases, Department of Medicine, Columbia University Vagelos College of Physicians and Surgeons, New York, NY, USA.*

[#]These authors contributed equally

[*]Correspondence: ll3411@cumc.columbia.edu (L.L.)

Abstract

The SARS-CoV-2 Omicron subvariants have dominated the pandemic due to their high
transmissibility and immune evasion conferred by the spike mutations. The Omicron
subvariants can spread by cell-free virus infection and cell-cell fusion, the latter of
which is more effective but has not been extensively investigated. In this study, we
developed a simple and high-throughput assay that provides a rapid readout to quantify
cell-cell fusion mediated by the SARS-CoV-2 spike proteins without using live or
pseudotyped virus. This assay can be used to identify variants of concern and to screen
for prophylactic and therapeutic agents. We further evaluated a panel of monoclonal
antibodies (mAbs) and vaccinee sera against D614G and Omicron subvariants, finding
that cell-cell fusion is substantially more resistant to mAb and serum inhibition than
cell-free virus infection. Such results have important implications for the development
of vaccines and antiviral antibody drugs against cell-cell fusion induced by SARS-CoV-
2 spikes.

Keywords: SARS-CoV-2, Omicron subvariants, spike protein, cell-cell fusion,
monoclonal antibody, polyclonal serum, neutralization, antibody evasion

Main text

Severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) is a novel beta-
coronavirus that causes the global coronavirus disease 2019 (COVID-19) pandemic.
The spike protein on the viral particles of SARS-CoV-2 mediates cell entry by binding
to the human angiotensin-converting enzyme 2 (hACE2) receptor, which triggers the
fusion of viral and cellular membranes (1). The spike protein is also the target for
neutralizing antibodies, which are critical for the host's immunity to control viral
infection (2, 3).

Previous studies have shown that the SARS-CoV-2 spike protein is capable of
mediating both cell-free virus infection and cell-cell transmission. Compared to cell-free
virus infection, cell-cell transmission is a more efficient mode of spreading due to the
direct transfer of viral particles between cells facilitated by tight cell-cell contact (4-7).
Cell-cell transmission has also been observed in other coronaviruses such as the
infectious bronchitis virus (IBV) and SARS-CoV (8). Like these coronaviruses, SARS-
CoV-2 cell-cell transmission involves S1/S2 cleavage, cell-cell fusion, and syncytia
formation (6). Notably, SARS-CoV-2 has been shown to exhibit higher cell-cell
transmission activity than SARS-CoV (4).

Cell-free virus infection of Omicron subvariants, with additional spike mutations, is
more resistant to antibody neutralization relative to the ancestral strain D614G (9).
Whether cell-cell fusion mediated by Omicron subvariants has such antibody evasion
properties, however, is not fully understood. Herein, we developed a simple, rapid, and
high-throughput assay to quantitatively analyse cell-cell fusion. We further showed that
cell-cell fusion induced by SARS-CoV-2 spike protein is largely resistant to the
inhibition by monoclonal antibodies (mAbs) and vaccinee sera, as compared to the cell-
free virus infection.

The experimental design of the cell-cell fusion assay is illustrated in **Figure 1A.** The
TZM-bl-hACE2 target cell line was generated from parental TZM-bl cell line through
stable transduction of hACE2 using a lentiviral system, followed by successive rounds
of fluorescence-activated cell sorting (FACS) and selection of transduced cells with
puromycin. Cell surface staining data revealed that TZM-bl-hACE2 cells displayed
elevated levels of hACE2 compared to TZM-bl cells and Vero-E6 cells (**Figure S1A**),
which are commonly used as target cells in pseudovirus assays for human
immunodeficiency virus (HIV)-1 and SARS-CoV-2, respectively. The donor 293T cells
were transiently transfected with HIV-1 Trans-Activator of Transcription (Tat) and
spike genes, before being co-cultured with the TZM-bl-hACE2 target cells at a ratio of
2:1. Cell-cell fusion could be evaluated both qualitatively, via observation with a light
microscope (**Figure S1B**), and quantitatively, through a luciferase reporter assay
(**Figure S1C**). Cell-cell fusion mediated by the D614G and BA.1 variants was assessed
at various time points after co-culture, and it was observed that the fusion activity of
both variants increased within the first 16-24 hours of co-culture (**Figure S1C**).
However, the fusion activity of D614G decreased after 24 hours, while that of BA.1

remained stable during the tested period after 16 hours of co-culture. Therefore, a co-
culture time of 16 hours was chosen for subsequent experiments to ensure comparable
levels of cell-cell fusion for both variants. Additionally, it was observed that D614G
spike mediated slightly higher cell-cell fusion activity than BA.1 spike, despite the fact
that BA.1 spike has higher affinity for ACE2 than D614G spike (10). To investigate this
discrepancy, the expression and processing of the spike protein in the donor cells
(293T) were examined (**Figure S2**). The results showed that the cleavage efficiency of
BA.1 and other Omicron subvariants was substantially lower than that of D614G,
resulting in lower levels of mature S1/S2 trimers in expressing cells, which may have
contributed to the observed reduction in cell-cell fusion activity for BA.1 compared to
D614G.

We initially assessed the susceptibility of TZM-bl-hACE2 cells to SARS-CoV-2
infection using vesicular stomatitis virus (VSV)-pseudotypes. The results showed a
dose-dependent infection of TZM-bl-hACE2 cells by the wild-type SARS-CoV-2
D614G, while TZM-bl cells were not infected, suggesting that hACE2 receptors were
necessary for virus entry (**Figure S3A**). Furthermore, we tested the infectivity of
multiple Omicron subvariants and found that they were all capable of infecting TZM-bl-
hACE2 cells, although with varying levels of infectivity (**Figure S3B**). Through these
evaluations, a rapid and reliable platform was developed and validated, allowing for the
comparison of SARS-CoV-2 variant spikes in mediating cell-free virus infection and
cell-cell fusion, as well as for testing the inhibitory effects of mAbs and vaccinee sera
on these processes.

We then assessed the efficacy of mAbs and human ACE2 (hACE2) in blocking cell-free
pseudovirus infection and cell-cell fusion mediated by the spikes of SARS-CoV-2
variants. The mAbs used in this study included CAB-A17 (11), Omi-3 (12), XGv347
(13), XGv282 (13), LY-CoV1404 (14), and COV2-2130 (15), which could neutralize
SARS-CoV-2 by recognizing the receptor binding domain (RBD) on the spike and
preventing hACE2 binding. **Figure 1B and Figure S4** display the structural modeling
of the binding of these RBD-directed mAbs and hACE2 to the RBD of D614G spike.
The neutralization activity of aforementioned mAbs was evaluated with VSV-
pseudotyped SARS-CoV-2 on Vero-E6 cells (**Figure S5**) and TZM-bl-hACE2 cells
(**Figures 1C and S6A**). Our results indicated that all mAbs effectively inhibited cell-
free infection of D614G, with the 50% inhibitory concentrations (IC_{50}) less than 0.3
μg/mL on TZM-bl-hACE2 cells and less than 0.1 μg/mL on Vero-E6 cells. Although
the neutralizing efficacy of some mAbs was significantly impaired by Omicron, such as
XGv347 against BA.4/5 and COV2-2130 against BA.1, most mAbs remained potent
against the cell-free virus infection of multiple Omicron subvariants. The IC_{50} values on
TZM-bl-hACE2 cells were higher than those on Vero-E6 cells, likely because TZM-bl-
hACE2 cells had a higher level of hACE2 expression, which hinders the inhibition of
virus entry (**Figure S1A**). As consistent with the pseudovirus neutralization data, we
found that mAbs CAB-A17, Omi-3, XGv347, XGv282, and LY-CoV1404 also
effectively neutralized authentic BA.1 on Vero-E6 cells with IC_{50} ranging from 0.007 to

0.217 µg/mL (**Figure S7**). But in contrast, our results showed that none of the mAbs
tested could potently and broadly inhibit cell-cell fusion mediated by SARS-CoV-2
spikes (**Figures 1D and S6B**). They retained weak potency against some of the
Omicron subvariants we tested. Among the variants tested, cell-cell fusion induced by
D614G was the most resistant to mAb inhibition, followed by BA.4/5, while BA.2 was
the most sensitive. Overall, our results demonstrated that cell-to-cell fusion mediated by
SARS-CoV-2 is more resistant to mAb inhibition than cell-free virus infection.

We further profiled the inhibition efficiency of serum samples in cell-free virus
infection and cell-cell fusion mediated by SARS-CoV-2 spikes. Sera were collected
from a study cohort in which each subject had received three doses of mRNA-based
SARS-CoV-2 vaccines. The clinical information of this study cohort is provided in
Table S1. In both Vero-E6 cells and TZM-bl-hACE2 cells, VSV-pseudotyped SARS-
CoV-2 variants were sensitive to serum neutralization, with the 50% inhibitory dose
(ID_{50}) titers ranging from 525 to 2074 on Vero-E6 cells, and from 403 to 1017 on TZM-
bl-hACE2 cells (**Figures 1E and S8**). The ID_{50} titers on the TZM-bl-hACE2 cells were
slightly lower than those on the Vero-E6 cells, which is consistent with what was
observed in the pseudovirus neutralization assay to mAbs. However, in the cell-cell
fusion inhibition assay on TZM-bl-hACE2 cells, the ID50 titers to serum inhibition
were more than 3-fold lower than those obtained from the cell-free virus neutralization
assay on the same cell line. Our data suggest that cell-cell fusion induced by SARS-
CoV-2 spike protein is more resistant to inhibition by vaccinee sera, compared to cell-
free virus infection mediated by the spike protein on viral particles. Among the
Omicron subvariants we tested, the cell-cell fusion mediated by BA.4/5 showed the
strongest evasion to vaccinee sera.

It was previously shown that cell-cell transmission of SARS-CoV-2 pre-Omicron
variants is more efficient and more refractory to blockade by mAbs and convalescent
sera than cell-free infection (4). In this study, we further demonstrate that the cell-cell
fusion mediated by various Omicron subvariants is also resistant to inhibition by mAbs
(**Figure 1D**) and vaccinee sera (**Figure 1E**). Future prophylactic or therapeutic agents
should consider strategies to suppress the cell-cell fusion of SARS-CoV-2.

It should be noted that all experiments in this study were performed *in vitro*, and it is
possible that the results may differ from *in vivo* outcomes. In addition, only humoral
immunity against cell-cell transmission of SARS-CoV-2 was evaluated, and the role of
cellular immunity requires further investigation. Lastly, cell-cell transmission was not
measured directly, but indirectly through a cell-cell fusion assay. If SARS-CoV-2
triggers intense cell-cell fusion, giant syncytia may form and result in cell death, which
in turn lessens cell-cell transmission.

In summary, we designed and validated a rapid and reliable cell-cell fusion assay for
SARS-CoV-2. Using this assay, we demonstrated that mAbs and vaccinee sera could
not effectively inhibit the cell-cell fusion mediated by SARS-CoV-2 spikes, in contrast
to cell-free infection. Such results suggest that developing vaccines and antiviral

antibody drugs that target the cell-cell fusion of SARS-CoV-2 may be a promising
approach.

Acknowledgements

157 We thank C. Lu and the Columbia Center for Translational Immunology (CCTI) Flow
158 Cytometry Core for their help with cell sorting.

Disclosure statement

160 L.L., and D.D.H. are inventors on patent applications (WO2021236998) or provisional
161 patent applications (63/271,627) filed by Columbia University. D.D.H. is a co-founder
162 of TaiMed Biologics and RenBio, consultant to WuXi Biologics and Brii Biosciences,
163 and board director for Vicarious Surgical.

Funding

165 This study was supported financially by the SARS-CoV-2 Assessment of Viral
166 Evolution Program, NIAID, NIH (Subcontract No. 0258-A709-4609 under Federal
167 Contract No. 75N93021C00014).

ORCID

Qian Wang https://orcid.org/0000-0001-9549-3140
Andre Yanchen Yeh https://orcid.org/0000-0002-5761-528X
Yicheng Guo https://orcid.org/0000-0001-5357-660X
Hiroshi Mohri https://orcid.org/0000-0003-3101-7625
Jian Yu https://orcid.org/0000-0001-8604-4538
Lihong Liu https://orcid.org/0000-0003-3128-8513
David D. Ho https://orcid.org/0000-0003-1627-149X

C

Pesudovirus inhibition on TZM-bl-hACE2 cells

IC50 (µg/mL)	D614G	BA.1	BA.2	BA.2.12.1	BA.4/5	BA.2.75
hACE2-Fc	1.097	0.206	0.160	0.196	0.222	0.168
CAB-A17	0.042	0.051	0.069	0.043	0.043	1.977
Omi-3	0.030	0.026	0.048	0.050	0.106	0.122
XGv347	0.009	0.005	0.005	0.005	>10	0.001
XGv282	0.0005	1.877	0.001	0.003	0.002	0.242
LY-CoV1404	0.005	0.004	0.002	0.002	0.003	0.020
COV2-2130	0.013	>10	0.016	0.035	0.604	0.498

D

Cell-cell fusion inhibition on TZM-bl-hACE2 cells

IC50 (µg/mL)	D614G	BA.1	BA.2	BA.2.12.1	BA.4/5	BA.2.75
hACE2-Fc	>50	>50	>50	>50	>50	>50
CAB-A17	>50	>50	>50	>50	>50	>50
Omi-3	>50	24.64	30.66	>50	>50	5.069
XGv347	>50	8.702	14.18	10.82	>50	0.782
XGv282	>50	>50	0.494	42.89	33.46	>50
LY-CoV1404	>50	20.71	4.931	7.463	13.30	>50
COV2-2130	>50	>50	5.738	>50	>50	>50

Figure 1. Resistance of cell-cell fusion induced by SARS-CoV-2 to monoclonal antibody and polyclonal serum neutralisation. (**A**) Schematic illustration of the cell-cell fusion assay. Briefly, 293T cells that were transiently transfected with the HIV-1 trans-activator of transcription (Tat) gene and SARS-CoV-2 spike gene were used as the donor cells. The target cells were TZM-bl-hACE2 cells that contained the firefly luciferase (luc) gene regulated by the long terminal repeat (LTR) promoter. Upon co-culture, spike protein on donor cells bound to the hACE2 receptors on target cells and initiated membrane fusion. In the syncytia, Tat proteins activated the LTR promoter, causing luc gene expression, which was quantified by luciferase reporter assay to evaluate the cell-cell fusion activity. (**B**) Structural modelling of RBD, hACE2, and mAbs. Different colors indicate different classes of mAbs, as grouped by their epitopes. (**C**) Pseudovirus neutralization IC_{50} titres for mAbs against D614G and Omicron subvariants on TZM-bl cells. (**D**) Cell-cell fusion inhibition IC_{50} titres for mAbs against D614G and Omicron subvariants. (**E**) Pseudovirus neutralization and cell-cell fusion inhibition ID_{50} titres for sera against D614G and Omicron subvariants. Each point represents a serum sample, with square and circle showing samples from individuals who received 3 doses of COVID-19 mRNA vaccines with and without breakthrough infection, respectively. Data are shown as geometric mean ± SD. Horizontal solid bar indicates the geometric mean of ID_{50} titres. Horizontal dotted line denotes the limit of

195   detection (LOD) at 100 or 30. Data shown in **C-E** are representative of one in three

196   independent experiments.

References

1. Shang J, Wan Y, Luo C, Ye G, Geng Q, Auerbach A, et al. Cell entry mechanisms of SARS-CoV-2. Proc Natl Acad Sci U S A. 2020 May 26;117(21):11727-34.

2. Liu L, Wang P, Nair MS, Yu J, Rapp M, Wang Q, et al. Potent neutralizing antibodies against multiple epitopes on SARS-CoV-2 spike. Nature. 2020 Aug;584(7821):450-6.

3. Rogers TF, Zhao F, Huang D, Beutler N, Burns A, He WT, et al. Isolation of potent SARS-CoV-2 neutralizing antibodies and protection from disease in a small animal model. Science. 2020 Aug 21;369(6506):956-63.

4. Zeng C, Evans JP, King T, Zheng YM, Oltz EM, Whelan SPJ, et al. SARS-CoV-2 spreads through cell-to-cell transmission. Proc Natl Acad Sci U S A. 2022 Jan 4;119(1).

5. Zhao M, Su PY, Castro DA, Tripler TN, Hu Y, Cook M, et al. Rapid, reliable, and reproducible cell fusion assay to quantify SARS-Cov-2 spike interaction with hACE2. PLoS Pathog. 2021 Jun;17(6):e1009683.

6. Rajah MM, Bernier A, Buchrieser J, Schwartz O. The Mechanism and Consequences of SARS-CoV-2 Spike-Mediated Fusion and Syncytia Formation. J Mol Biol. 2022 Mar 30;434(6):167280.

7. Papa G, Mallery DL, Albecka A, Welch LG, Cattin-Ortola J, Luptak J, et al. Furin cleavage of SARS-CoV-2 Spike promotes but is not essential for infection and cell-cell fusion. PLoS Pathog. 2021 Jan;17(1):e1009246.

8. Belouzard S, Millet JK, Licitra BN, Whittaker GR. Mechanisms of coronavirus cell entry mediated by the viral spike protein. Viruses. 2012 Jun;4(6):1011-33.

9. Wang Q, Iketani S, Li Z, Liu L, Guo Y, Huang Y, et al. Alarming antibody evasion properties of rising SARS-CoV-2 BQ and XBB subvariants. Cell. 2023 Jan 19;186(2):279-86 e8.

10. Li L, Liao H, Meng Y, Li W, Han P, Liu K, et al. Structural basis of human ACE2 higher binding affinity to currently circulating Omicron SARS-CoV-2 sub-variants BA.2 and BA.1.1. Cell. 2022 Aug 4;185(16):2952-60 e10.

11. Sheward DJ, Pushparaj P, Das H, Kim C, Kim S, Hanke L, et al. Structural basis of Omicron neutralization by affinity-matured public antibodies. bioRxiv. 2022.

12. Nutalai R, Zhou D, Tuekprakhon A, Ginn HM, Supasa P, Liu C, et al. Potent cross-reactive antibodies following Omicron breakthrough in vaccinees. Cell. 2022 Jun 9;185(12):2116-31 e18.

13. Wang K, Jia Z, Bao L, Wang L, Cao L, Chi H, et al. Memory B cell repertoire from triple vaccinees against diverse SARS-CoV-2 variants. Nature. 2022 Mar;603(7903):919-25.

14. Westendorf K, Zentelis S, Wang L, Foster D, Vaillancourt P, Wiggin M, et al. LY-CoV1404 (bebtelovimab) potently neutralizes SARS-CoV-2 variants. Cell Rep. 2022 May 17;39(7):110812.

15. Zost SJ, Gilchuk P, Case JB, Binshtein E, Chen RE, Nkolola JP, et al. Potently neutralizing and protective human antibodies against SARS-CoV-2. Nature. 2020 Aug;584(7821):443-9.