CXCR4 blockade alleviates pulmonary and cardiac outcomes in early COPD

Isabelle Dupin,[1,2,3] Pauline Henrot,[1,2,4] Elise Maurat,[1,2] Reshed Abohalaka,[1,2,5] Sébastien

Chaigne,[1,2,4] Dounia El Hamrani,[1,2], Edmée Eyraud,[1,2], Renaud Prevel,[1,2,4] Pauline Esteves,[1,2]

Maryline Campagnac,[1,2] Marielle Dubreuil,[1,2] Guillaume Cardouat,[1,2] Clément Bouchet,[1,2] Olga

Ousova,[1,2], Jean-William Dupuy,[1,2] Thomas Trian,[1,2] Matthieu Thumerel,[1,2,4] Hugues

Bégueret,[1,2,4] Pierre-Olivier Girodet,[1,2,4] Roger Marthan,[1,2,4] Maeva Zysman,[1,2,4] Véronique

Freund-Michel,[1,2], Patrick Berger,[1,2,4]

[1] Univ. Bordeaux, Centre de Recherche Cardio-thoracique de Bordeaux, INSERM U1045, IHU

Liryc, CIC 1401, Proteomics Facility, F-33600 Pessac, France

[2] INSERM, Centre de Recherche Cardio-thoracique de Bordeaux, U1045, CIC 1401, F-33600

Pessac, France

[3] Institut universitaire de France (IUF)

[4] CHU Bordeaux, Service d'exploration fonctionnelle respiratoire, Service de réanimation,

Service de pneumologie, Service de chirurgie thoracique, Service de cardiologie-

électrophysiologie et stimulation cardiaque, Service d'anatomopathologie, CIC-P 1401, F-33600

Pessac, France

[5] Krefting Research Centre, Department of Internal Medicine and Clinical Nutrition, Institute of

Medicine, Sahlgrenska Academy, University of Gothenburg - Gothenburg, Sweden,

Corresponding author: Prof. Isabelle Dupin
Centre de Recherche Cardio-thoracique de Bordeaux, INSERM U1045
PTIB. Hôpital Xavier Arnozan. Avenue du Haut Lévêque. 33600 Pessac, FRANCE
Tel : +33 5 47 30 27 51

e-mail : isabelle.dupin@u-bordeaux.fr

ABSTRACT

Chronic obstructive pulmonary disease (COPD) is a prevalent respiratory disease lacking effective treatment. Focusing on early COPD should help to discover disease modifying therapies. We aimed to examine the role of the CXCL12/CXCR4 axis in early COPD from both human samples and murine models. Blood samples and lung tissues of early COPD patients and controls were obtained in order to analyse CXCL12 and CXCR4 levels. To generate an early COPD model, ten-week-old male C57BL/6J mice were exposed to cigarette smoke (CS) for 10 weeks and intranasal instillations of polyinosinic–polycytidylic acid (poly(I:C)) for the last 5 weeks to mimic exacerbations. CXCR4 expressing cells number was increased in the blood of patients with COPD, as well as in the blood of exposed mice. Lung CXCL12 expression was higher in both early COPD patients and exposed mice. Exposed mice presented mild airway obstruction, peri-bronchial fibrosis and right heart thickening. The density of fibrocytes expressing CXCR4 was increased in the bronchial submucosa of these mice. Conditional inactivation of CXCR4 at adult stage as well as pharmacological inhibition of CXCR4 with plerixafor injections improved lung function, reduced inflammation, and protected against CS and poly-(I:C)-induced airway and cardiac remodeling. CXCR4$^{-/-}$ and plerixafor-treated mice also had less CXCR4-expressing circulating cells and a lower density of peri-bronchial fibrocytes. We demonstrate that targeting CXCR4 has beneficial effects in an animal model of early COPD and provide a framework to translate these preclinical findings to clinical settings in a drug repurposing approach.

Key words: COPD pathology, macrophage biology, cytokine biology

Clinical relevance

We demonstrate that CXCL12/CXCR4 axis plays an important role in the pathogenesis of early

COPD. Inhibition of this axis improves lung function and cardiac tissue remodeling, supporting

the future use of CXCR4 inhibitors to slow down the progression of the disease.

Abbreviations

BAL   Broncho-alveolar lavage

bFGF   basic Fibroblast Growth Factor

BSM   Bronchial smooth muscle

CS   Cigarette smoke

CT   Computed tomographic

ECG   Electrocardiogram

FEV1   Forced expiratory volume in 1 second

FEV0.05   Forced expiratory volume in 0.05 second

FVC   Forced vital capacity

IFN   Interferon

LV + S   Left ventricle plus septum

MRI   Magnetic Resonance Imaging

Poly-(I:C)   Polyinosinic–polycytidylic acid

RA   Room air

RV   Right ventricle

RVSP   Right ventricular systolic pressure

VEGF   Vascular Endothelial Growth Factor

INTRODUCTION

Chronic obstructive pulmonary disease (COPD) is a common disease, characterized by persistent respiratory symptoms and airflow limitation (1). Major risk factors for COPD include chronic exposure to noxious particles, mainly cigarette smoke (CS), and airway infections (2). Early COPD has been recently defined as patients aged <50 years with a smoking exposure >10 pack-years, with a ratio between forced expiratory volume in 1 second (FEV1) and forced vital capacity (FVC) less than 70% or less than the lower limit of normal (LLN), compatible computed tomographic (CT) abnormalities (*i.e*., visual emphysema, air trapping, or bronchial thickening graded mild or worse), and/or evidence of accelerated FEV1 decline of >60 ml/years (3). Twenty-four percent of these early COPD patients developed clinical COPD 10 years later (4). Therapeutic interventions in early COPD patients could help to tackle the disease before reaching irreversible tissue damage (5). However, it is extremely difficult to enrol those patients into trials and collect their bronchial samples, since they are rarely diagnosed, demonstrating the urgent need to develop animal models of early COPD.

Exposure of lung structural and immune cells to CS and infectious agents results in the release of various inflammatory substances, including chemokines (6). The chemokine receptor CXCR4 and its associated ligand CXCL12 appear to be attractive therapeutic targets for early COPD since (i) they are implicated in the migration of inflammatory cells from the blood (7) to the COPD lungs (8), (ii) their expression and function are controlled by other cytokines (9), oxygen concentration (10) and microbial agents (11, 12), which are disrupted in COPD, and (iii) pharmacological blockade of CXCR4 reduces emphysema development in a long-term COPD murine model (13). CXCR4 is notably expressed by fibrocytes, a rare population of circulating fibroblast-like cells (14), that are assumed to play a crucial role in COPD (7, 8).

In the present study, we thus hypothesized that the CXCL12/CXCR4 axis plays a role in the

pathological processes leading to COPD development during early adulthood and, hence, that

inhibiting CXCR4 would protect from CS-induced airflow limitation and tissue remodeling. In

early COPD patients' samples, as well as in murine model of early COPD, both CXCR4-

expressing cells in the blood and CXCL12 in the lung were overexpressed. Using either

transgenic mice deficient for CXCR4 or the CXCR4 antagonist plerixafor, we showed that lungs

were protected against CS-induced both lung function alteration and fibrocytes accumulation, and

that the right heart was resistant to cardiac remodeling.

Some of these findings have been previously reported in abstract form (15, 16).

METHODS

Complete details concerning methods appear in an online data supplement.

Study Population

Human lung tissues were obtained from both the "Fibrochir" study (NCT01692444) (8) and

"TUBE" (i.e., TissUs Bronchiques et PulmonairEs, sponsored by the University hospital of

Bordeaux) biological collection (Table S1). Blood samples of COPD patients were obtained from

"COBRA" (i.e., COhort of BRonchial obstruction and Asthma; sponsored by INSERM) (Table

S2) and. All subjects gave their written informed consent to participate to the studies. The studies

received approval from the local or national ethics committees.

Dataset transcriptomic analysis

Transcriptomes obtained on human blood and lung tissues were downloaded from the NCBI

Gene Expression Omnibus (GEO) database (http://www.ncbi.nlm.nih.gov/geo/) using datasets

under the accession codes of GSE100153 and GSE76925, respectively.

Mouse model of early COPD

Male C57BL/6J mice were obtained from Janvier (St Berthevin, France). CXCR4$^{lox/lox}$ and

Cre2ERT2 mice were obtained from The Jackson Laboratory (17) and the "Institut Clinique de

la Souris" (18), respectively. All animal studies were performed according to European directives

for the protection of vertebrate animals. Agreement was obtained from French authorities

(numbers A33-063-907 and A33-318-3) and all the protocols were approved by the local ethics

committee.

Ten-week-old mice were exposed for 10 weeks to either room air (RA) or CS from non-filtered

research cigarettes (2R4; University of Kentucky, Lexington, KY) using the smoking apparatus

(Anitech, Paris, France), as described previously by Almolki *et al*. (19). After 5 weeks of CS

exposure, mice were anesthetized and 50 μg of poly(I:C) diluted in PBS, mimicking responses by

RNA virus infections (20), or its vehicle control, were administered twice per week for 5 weeks

via nasal aspiration. Electrocardiograms (ECG) and Magnetic Resonance Imaging (MRI) were

performed during the 10[th] week of the protocol. After lung function measurement, right

ventricular systolic pressure (RVSP) was determined invasively. Following bronchoalveolar

lavage (BAL), mice were sacrificed and lung, heart and blood were collected. The Fulton index

was calculated as the ratio between the right ventricle (RV) and the left ventricle plus septum

(LV + S).

CXCR4 genetic deletion was induced by tamoxifen. CXCR4 pharmacological inhibition was

induced by subcutaneous injection of 1 mg/kg plerixafor (Sigma-Aldrich, Saint-Quentin-

Fallavier, France), 5 times/week during the last 5 weeks of the exposition protocol.

Experimental procedures in both human and mice

CXCR4-expressing cells were quantified in the blood and in the lungs by flow cytometry and

immunohistochemistry, as described previously (8). CXCL12 and CXCR4 levels were evaluated

by ELISA, western blotting and RT-PCR. Proteomics analyses were performed using mass

spectrometry, as described previously (21).

Statistical Analysis

Statistical significance, defined as $P < 0.05$, was assessed by t-tests and MANOVA for variables

with parametric distribution, and by Kruskal-Wallis with multiple comparison z tests, Mann-

Whitney tests, Wilcoxon tests and Spearman correlation coefficients for variables with non-

parametric distribution.

RESULTS

CXCL12 and CXCR4 expression is increased in lung and blood, respectively, in patients

with COPD

First, we determined whether the expression of CXCL12 and CXCR4 was altered in patients with

COPD by interrogating RNAseq data from lung of control subjects and COPD patients

(GSE76925 data set, http://www.ncbi.nlm.nih.gov/geo/). We found that CXCL12 expression was

significantly increased in the lung of patients with COPD (GSE76925 data set, figure 1A). In a

separate cohort of patients with and without early COPD (table S1), CXCL12 was predominantly

localized to bronchial epithelial cells and peri-bronchial infiltrating cells, which are likely

immune cells (figure 1B). The CXCL12 surface immunostaining was significantly increased in

lungs of patients with early COPD in comparison to control lungs (figure 1B-C). CXCR4 level

was not significantly altered in the lungs of patients with COPD compared to those of control

smokers at the mRNA level (GSE76925 data set, figure 1D), which was corroborated by a similar

result at the protein level by quantification of immunostaining (figure 1E) in patients with and

without early COPD. In the blood, CXCR4 was significantly increased, at the mRNA level, in

patients with COPD compared to control subjects (figure S1A). In a separate cohort of moderate

to mild COPD patients (GOLD I to II, as defined by (1), table S2), the percentage of CXCR4-

expressing cells was also increased in the blood of moderate COPD patients in comparison with

mild COPD patients (figure S1B-C).

Figure 1: Characterization of CXCR4 and CXCL12 expression in lung of patients with or without COPD. (A, D) CXCL12 (A) and CXCR4 (D) mRNA expression in human lungs. Data are derived from a publicly available GSE-set (GSE76925). Control subjects n=40; patients with COPD n=111. (B) Representative staining of CXCL12 (magenta) and nucleus (white) in peri-bronchial areas of human lungs. Lower panels: higher magnification of the images surrounded by white squares in the upper panels. Images obtained in control *vs* early COPD lungs were acquired with the same exposure time and gain. The white arrowheads and arrows indicate respectively CXCL12-expressing bronchial cells and immune-type cells. (C, E) Quantification of the percentage of positive CXCL12 (C) or CXCR4 (E) surface (normalized by the total area) (n=5 control subjects, n=6 patients with early COPD). Medians are represented as horizontal lines. (A, C, D, E) *: P <0.05, ***: P<0.001, t-test or Mann-Whitney test.

Chronic CS and poly(I:C) exposure induces functional obstruction, lung inflammation and both lung and heart tissue remodeling in mice

To model early COPD, we used mice at the beginning of adulthood (*i.e.*, 10-week old) (22), which were exposed to a combination of CS exposure and intranasal instillations of poly(I:C) in order to mirror the effects of smoking and exacerbations (23). The protocol was based on a previous study (20), where mice were exposed during 5 weeks to cigarette smoke, with poly-IC instillations during the 2 last weeks, based on the intricate relationships between innate immune response and COPD pathogenesis. We reproduced a similar protocol and found no modification of lung function at 5 weeks (figure S2A-D), prompting us to extend to 10 weeks the duration of

the protocol. Ten weeks of CS exposure with 10 repeated instillations of poly(I:C) during the last

5 weeks significantly decreased the FEV0.05/FVC ratio (figure 2A-C). Other lung function

parameters, in particular respiratory system compliance and tissue elastance, were not modified

by CS and poly(I:C) exposure (figure S2E-J). At the end of the protocol, the weight gain of the

exposed mice was significantly different from that of the control mice (figure S3A-B). CS and

poly(I:C) exposure increased total cell count in the BAL fluid, as well as the absolute neutrophil,

lymphocyte and macrophage numbers, 1 day after the last poly(I:C) instillation (figure 2D-F).

Although lower, the increase in neutrophils and lymphocytes persisted 4 days after the last

poly(I:C) instillation (figure 2D-F). CS and poly(I:C) synergistically enhanced airflow

obstruction and BAL inflammation (figure S4). We also analyzed lung proteomics obtained 1 day

after the last poly-IC instillation. Of the 137 pathways that were significantly different in early

COPD   vs   control   mice,   largest   upregulated   genes   were   those   of   the

hypercytokinemia/hyperchemokinemia box and of the pathogenesis of influenza and interferon

(IFN) signaling (figure 2G-H). In contrast, inhibited pathways were the wound healing and the

xenobiotic metabolism PXR signaling pathways (figure 2G-H). Exposed mice had a moderate but

significant increase in peri-bronchial fibrosis compared to control mice (figure 2I-J), mimicking

the thickening of distal airway tissue in COPD patients (24).

Then, we investigated the impact of CS and poly(I:C) exposure on cardiac function. There was no

evidence of heart failure in our model (figure S5A-B). Moreover, there was no sign of heart

rhythm disorder as assessed by ECG monitoring (figure S5C and figure S6A-F). Using MRI, we

also found that CS and poly(I:C)-exposition did not alter RV or LV volumes (figure S5D-F).

However, we detected a significant increase of RV wall thickness in exposed mice, indicating RV

hypertrophy (figure 2K-L). This finding was confirmed by the measurement of the Fulton index

(figure 2M). Of note, the RV systolic pressure (RVSP) did not differ between early COPD and

control mice (figure 2N).

Figure 2: Evaluation of lung function, inflammation, bronchial and heart remodeling in experimental COPD. (A) Mice are exposed either to room air (RA) and challenged with PBS, or exposed to cigarette smoke (CS) and challenged with poly(I:C), during 10 weeks. Mice are sacrificed at day 68 or 71 (*i.e.*, 1 day or 4 days after the last poly(I:C) instillation). (B) Average expiratory flow-volume curves of RA+PBS-exposed mice (black curve) and CS+poly(I:C)-exposed mice (grey curve). Lower and upper error bars represent standard deviations for CS+poly(I:C) and RA+PBS-exposed mice, respectively. (C) FEV0.05/FVC (FEV0.05: Forced Expiratory Volume during 0.05 s, FVC: Forced Vital Capacity) of RA+PBS-exposed mice (black circles, n=40) and CS+poly(I:C)-exposed mice (grey squares, n=41). Data represent individual mice. (D) Total cells count in bronchoalveolar lavage (BAL). Kruskal-Wallis test followed by Dunn's post-tests. (E-F), BAL differential cell recovery 1 day (d+1, RA+PBS: n=24,

CS+poly(I:C): n=22) and 4 days (d+4, RA+PBS: n=26, CS+poly(I:C): n=31) after the last

poly(I:C) instillation. (G) Top Canonical Ingenuity Pathways significantly altered in

CS+poly(I:C) (n=5) *vs* RA+PBS (n=5)-exposed lungs (n=5) obtained 1 day after the last poly-IC

instillation, ranked by Z-score (negative and positive for the pathways represented respectively

on the top and bottom part of the graph), obtained by Gene Set Enrichment Analysis. (H)

Heatmaps of differentially regulated proteins in CS+poly(I:C) vs RA+PBS-exposed lungs, from

the pathways "hypercytokinemia/hyperchemokinemia in the pathogenesis of influenza" (left),

"wound healing signaling pathway" (middle) and "xenobiotic metabolism PXR signaling

pathway" (right). The colour scale indicates the log_2 fold changes of abundance for each protein.

(I) Representative Masson's trichrome staining to assess peri-bronchial fibrosis. (J) Standardized

fibrosis, defined as (peri-bronchial fibrosis ("PF", %) - mean PFcontrol)/standard deviation

PFcontrol). The peri-bronchial fibrosis ("PF", percentage) is defined by the ratio between the area

of segmented pixels in the area of analysis divided by the total peri-bronchial area. RA+PBS:

n=35, CS+poly(I:C): n=34. (K) Representative short-axis cine magnetic resonance images of

heart at end-systolic stage (left panels) and end-diastolic stage (middle panels) in control mice

(top panels) and CS and poly(I:C)-exposed mice (bottom panels). Right panels: higher

magnification of RV wall. (L) Wall thickness. RA+PBS: n=6, CS+poly(I:C): n=6. (M) Fulton

index, defined as (right ventricle (RV)/ left ventricle plus septum (LV + S). RA+PBS: n=26,

CS+poly(I:C): n=24. (N) Right ventricular systolic pressure (RVSP). RA+PBS: n=12,

CS+poly(I:C): n=12. (C, E, F, I, L, M, N) Unpaired t test or Mann-Whitney test. *: P<0.05, **:

P<0.01, ***: P<0.001.

CXCR4 and CXCL12 expression is increased in blood and lung, respectively, in

experimental COPD

The percentage of CXCR4-expressing cells was significantly increased in the blood of CS and

poly(I:C)-exposed mice compared to control mice, 1 day after the last poly(I:C) instillation

(figure 3A-B), indicating that experimental COPD replicated the increased levels of CXCR4 in

the blood of COPD patients. This effect appeared to be due to a synergy between CS and

poly(I:C) (figure S7). The level of CXCR4$^+$ circulating cells decreased and became similar

between control and exposed mice 4 days after the last poly(I:C) instillation (figure 3B). No

significant modification of CXCL12 plasma concentration was evidenced in mice (figure 3C).

The percentage of CXCR4-expressing cells was not significantly altered in whole lung

homogenates in experimental early COPD (figure 3D-E). In lung homogenates, CS and

poly(I:C)-exposure significantly increased CXCL12 protein levels compared to control mice, 1

and 4 days after the last poly(I:C) instillation (figure 3F). In good agreement with our findings in

patients' lungs, CXCL12 was expressed by bronchial epithelial cells and peri-bronchial

infiltrating cells (figure 3G), with a higher expression in lungs of animals exposed to CS and

poly-(I:C) in comparison to control lungs (figure 3H).

Figure 3: Characterization of CXCR4 and CXCL12 expression in the blood and lung in
experimental COPD. Mice are exposed either to room air (RA) and challenged with PBS, or
exposed to cigarette smoke (CS) and challenged with poly(I:C), during 10 weeks. They are
sacrificed at day 68 or 71 (i.e. 1 day or 4 days after the last poly(I:C) instillation). (A, D) Upper
left panels: dot plots represent representative side scatter (SSC, y-axis)-forward scatter (FSC, x-
axis) graphs of circulating (A) and lung (D) cells. DAPI⁻ cells, and DAPI⁻ CXCR4⁺ cells are
shown respectively in grey and pink. Upper right panels and bottom panels: histograms represent
representative cell count (y-axis) versus Phycoerythrin (PE) fluorescence (x-axis) in DAPI⁻
circulating (A) and lung (D) cells. Percentages of CXCR4⁺ cells in the DAPI⁻ cells are shown in
pink. (B, E) Levels of circulating (B) and lung (E) CXCR4⁺ cells 1 day (d+1) and 4 days (d+4)
after the last poly(I:C) instillation. (C, F) Plasma (C) and lung (F) CXCL12 concentration at d+1
and d+4. For lung CXCL12 concentration, the value is normalized to total protein concentration.
(G) Representative staining of CXCL12 (magenta) and nucleus (white) in peri-bronchial areas of
mouse lungs. Lower panels: higher magnification of the images surrounded by white squares in
the upper panels. Images obtained in RA and PBS *vs* CS and poly(I:C)-exposed have been
acquired with the same exposure time and gain. The white arrowheads and arrows indicate
respectively CXCL12-expressing bronchial cells and immune-type cells. (H) Quantification of
the percentage of positive CXCL12 surface (normalized by the total area). (B, C, E, F, H)
Medians are represented as horizontal lines. *: P <0.05, **: P<0.01, Mann-Whitney test.

291
292
293
294

Fibrocytes and CXCR4$^+$ fibrocytes are increased in lungs of experimental early COPD

To quantify fibrocytes in control and experimental COPD lungs, single cell suspension was

stained with CD45 and FSP1 as well as CXCR4. We defined fibrocytes as cells expressing CD45

and a high level of FSP1 ("CD45$^+$ FSP1^{high}") (25, 26), which is also in agreement with previous

studies performed in human lung (8, 27). The percentage of fibrocytes was significantly increased

in CS and poly(I:C)-exposed lungs 1 day after the last poly(I:C) instillation (figure 4A-C). The

average percentage of leukocytes (CD45$^+$ cells) in control and exposed lungs was 69.2 % and

74.0 %, respectively (p=0.07, figure 4D). In control and experimental early COPD, a vast

majority of lung fibrocytes expressed CXCR4 (96.1 and 96.6 %, respectively, figure 4B) and the

percentage of CXCR4$^+$ fibrocytes was significantly increased in the lungs of exposed mice

(figure 4C). Four days after the last poly(I:C) instillation, the total level of fibrocytes decreased

and became similar between control and experimental early COPD lungs (figure 4C). Since the

flow cytometry approach limits our analysis to the whole lung, we co-stained FSP1 with CD45 in

murine lungs, 4 days after the last poly(I:C) instillation (figure 4F) using immunohistochemistry.

Similarly to that previously reported in human lung (8), the density of peri-bronchial fibrocytes

was higher in CS and poly(I:C)-exposed murine lungs than in control lungs (figure 4G).

Moreover, we also observed *in situ* CXCR4-expressing fibrocytes in peri-bronchial areas of

human lung samples ("Fibrochir" study (8), figure 4H). Of note, fibrocytes were found in right

and left ventricles, but their densities were unchanged by CS and poly(I:C) exposure (figure S8).

Figure 4: Characterization of lung fibrocyte level and localization in experimental COPD and in patients with or without COPD. (A-G) Mice are exposed either to room air (RA) and challenged with PBS, or exposed to cigarette smoke (CS) and challenged with poly(I:C), during 10 weeks. Mice are sacrificed 1 day ("d+1") or 4 days ("d+4") after the last poly(I:C) instillation. a) Representative flow cytometry contour plots in each condition. The x-axis is for CD45-Allophycocyanin (APC) fluorescence and the y-axis, FSP1-Fluorescein-5-isothiocyanate (FITC) fluorescence. Percentages of CD45$^+$ FSP1^{high} cells are shown in green. (B) Representative cell count (y-axis) versus CXCR4-Phycoerythrin (PE) fluorescence (x-axis) in CD45$^+$ FSP1^{high}-cell subsets in mouse lungs. (C-E) Levels of lung CD45$^+$ FSP1^{high} cells (C), CD45$^+$ cells (D), CD45$^+$ FSP1^{high} CXCR4$^+$ cells (E), in RA+PBS-exposed mice (black circles) and CS+poly(I:C)-exposed mice (grey squares), at d+1 and d+4. (F) Representative staining of CD45 (green) and FSP1 (red) in peri-bronchial areas (delimited in pink) of lung mice at d+4. The white arrows indicate fibrocytes, defined as CD45$^+$ FSP1$^+$ cells. (G) Quantification of fibrocyte density (normalized by peri-bronchial area). (H) Representative staining of CD45 (green) and FSP1 (red) in peri-bronchial area. The white arrowheads indicate fibrocytes, defined as CD45$^+$ FSP1$^+$ cells. (F)

Representative staining of CD45 (green), FSP1 (red), nuclei (DAPI, blue) and CXCR4 (white) in
peri-bronchial areas of lung samples of control subjects and patients with COPD. The white
arrows indicate CXCR4-expressing fibrocytes, defined as $CD45^+$ $FSP1^+$ $CXCR4^+$ cells. (F, H)
Left and lower panels: higher magnification of the images surrounded by white squares in the
right upper panels. (C-E, G) Medians are represented as horizontal lines. *: P <0.05, **: P<0.01,
Mann-Whitney test.
336

CXCR4 deletion protects against functional airway obstruction, lung and heart tissue

remodeling, and inhibits fibrocyte recruitment into the lungs in experimental early COPD

As mice genetically deficient for CXCR4 die perinatally (28), we generated conditional CXCR4

mutants using floxed allele (17) and a Cre2ERT2 strain (18), thereby obtaining mutants harboring

a specific inactivation of CXCR4 at early adulthood after tamoxifen treatment. Intraperitoneal

tamoxifen administration, for 5 consecutive days, in $CXCR4^{lox/lox}$/Cre2ERT2 mice was not

sufficient to induce a large CXCR4 downregulation, particularly, in the lungs (figure S9A-B).

Therefore, we combined intraperitoneal tamoxifen administration and feeding mice with

tamoxifen-containing food to obtain a more robust decrease in CXCR4 expression in the lung and

bone marrow (figure S9C-E). As tamoxifen-containing food reduced weight gain in mice (figure

S3C-D), we used $CXCR4^{lox/lox}$ mice as control mice, and gave all transgenic mice tamoxifen-

containing food.

$CXCR4^{lox/lox}$/Cre2ERT2 and $CXCR4^{lox/lox}$ mice were exposed to CS and poly(I:C) for 10 weeks

(figure 5A). As expected, CXCR4 expression decreases in the lung and blood of

$CXCR4^{lox/lox}$/Cre2ERT2 mice (figure S10). Moreover, FEV0.05/FVC ratio was higher in

$CXCR4^{-/-}$ mice compared to $CXCR4^{lox/lox}$ mice, when exposed to CS and poly(I:C) (figure 5B).

In the BAL fluid, CXCR4 deletion induced a significant decrease of macrophage and increase of

neutrophil percentages, 1 day after the last poly(I:C) instillation (figure 5C-E). $CXCR4^{-/-}$ mice

had a significant reduction in collagen deposition around small airways, compared to CS and

poly(I:C)-exposed CXCR4$^{+/+}$ mice (figure 5F-G), confirming an important role of CXCR4 in CS-

induced bronchial remodeling.

We then assessed the impact of CXCR4 deletion on right heart remodeling. CXCR4 deletion did

not affect electrophysiological and hemodynamic (figure S6G-L and figure S11) but it

significantly reduced the RV wall thickness as assessed by heart MRI *in vivo* and the Fulton

index *ex vivo* in exposed mice (figure 5H-J). RVSP value was unchanged by CXCR4 deletion

(figure 5K).

Figure 5: Evaluation of lung function, inflammation, and bronchial and heart remodeling upon conditional knockout of CXCR4 in experimental COPD. (A) $CXCR4^{lox/lox}$ (grey squares) and $CXCR4^{lox/lox}$/Cre2ERT2 (pink squares) mice are exposed to cigarette smoke (CS) and challenged with poly(I:C). All the mice were administered with 1 mg intraperitoneal tamoxifen for 5 consecutive days at the beginning of the COPD protocol, and then fed by tamoxifen-containing food during the remaining protocol. Mice are sacrificed at day 68 or 71 (1 day and 4 days after the last poly(I:C) instillation). (B) FEV0.05/FVC in the different conditions. Data represent individual mice. $CXCR4^{lox/lox}$: n=27, $CXCR4^{lox/lox}$/Cre2ERT2: n=32. (C) Total cells count in bronchoalveolar lavage (BAL). Kruskal-Wallis test followed by Dunn's post-tests. (D-E) BAL relative (D) and absolute (E) cell composition respectively 1 day and 4 days after the last poly(I:C) instillation. d+1: $CXCR4^{lox/lox}$: n=18, $CXCR4^{lox/lox}$/Cre2ERT2: n=19; d+4: $CXCR4^{lox/lox}$: n=21, $CXCR4^{lox/lox}$/Cre2ERT2: n=23. (F) Representative Masson's trichrome staining to assess peri-bronchial fibrosis. (G) Standardized fibrosis, defined as (peri-bronchial fibrosis ("PF", %) - mean PFcontrol)/standard deviation PFcontrol). The peri-bronchial fibrosis ("PF", percentage) is defined by the ratio between the area of segmented pixels in the area of analysis divided by the total peri-bronchial area. $CXCR4^{lox/lox}$: n=14, $CXCR4^{lox/lox}$/Cre2ERT2: n=18. (H) Representative short-axis cine magnetic resonance images at end-systolic stage (left panels) and end-diastolic stage (middle panels) in CS and poly(IC)-exposed $CXCR4^{lox/lox}$ mice (top panels) and CS and poly(IC)-exposed $CXCR4^{lox/lox}$/Cre2ERT2 mice (bottom panels). Right panels: higher magnification of RV wall. i) Wall thickness. $CXCR4^{lox/lox}$: n=4, $CXCR4^{lox/lox}$/Cre2ERT2: n=4. (J) Fulton index, defined as (right ventricle (RV)/ left ventricle plus septum (LV + S). $CXCR4^{lox/lox}$: n=46, $CXCR4^{lox/lox}$/Cre2ERT2: n=53. (K) Right ventricular systolic pressure (RVSP). $CXCR4^{lox/lox}$: n=29, $CXCR4^{lox/lox}$/Cre2ERT2: n=36. (B, D, E, G, J, K) *: $P<0.05$, **: $P<0.01$, ***: $P<0.001$. Unpaired t test or Mann-Whitney test.

Moreover, we observed reduced flow cytometry levels of $CXCR4^+$ fibrocytes in CS and poly(I:C)-exposed whole lungs of $CXCR4^{-/-}$ mice compared with $CXCR4^{+/+}$ exposed mice, 1 day after the last poly(I:C) instillation (figure 6A-D). Four days after the last poly(I:C) instillation, the density of fibrocytes was also significantly decreased around the small airways (figure 6F-G).

Figure 6: Characterization of fibrocyte level and peri-bronchial density in lung upon conditional knockout of CXCR4 in experimental COPD. CXCR4$^{lox/lox}$ (grey squares) and CXCR4$^{lox/lox}$/Cre2ERT2 (pink squares) mice are exposed to cigarette smoke (CS) and challenged with poly(I:C). Mice are sacrificed at day 68 or 71 (1 day and 4 days after the last poly(I:C) instillation). (A) Representative flow cytometry contour plots in each condition. The x-axis is for CD45-Allophycocyanin (APC) fluorescence and the y-axis, FSP1-Fluorescein-5-isothiocyanate (FITC) fluorescence. Percentages of CD45$^+$ FSP1^{high} cells are shown in green. (B) Representative cell count (y-axis) versus CXCR4-Phycoerythrin (PE) fluorescence (x-axis) in CD45$^+$ FSP1^{high}-cell subsets in mouse lungs. (C-E) Levels of lung CD45$^+$ FSP1^{high} cells (C), CD45$^+$ FSP1^{high} CXCR4$^+$ cells (D), CD45$^+$ cells (E) 1 day after the last poly(I:C) instillation in each condition. (F) Quantification of fibrocyte density (normalized by peri-bronchial area) 4 days after the last poly(I:C) instillation in each condition. (G) Representative staining of CD45 (red) and FSP1 (green) in peri-bronchial area (delimited in pink) in each condition. The white arrows indicate fibrocytes, defined as CD45$^+$ FSP1$^+$ cells. (C-F) Medians are represented as horizontal lines. *: P <0.05, Unpaired t test or Mann-Whitney test.

Pharmacological inhibition of CXCR4 protects against functional obstruction and inhibits

fibrocyte recruitment into the lungs in experimental COPD

We then assessed the efficacy of a CXCR4 antagonist (*i.e.* AMD3100, plerixafor) in the early

COPD murine model (figure 7A). Mice weight was not affected by daily subcutaneous plerixafor

treatment during the second half of the protocol (figure S3E-F). Plerixafor significantly reduced

CXCR4 expression 1 day after the last poly(I:C) instillation in circulating cells, but not in lung

cells (figure S12). In exposed-mice treated with plerixafor, the FEV0.05/FVC ratio was

significantly higher than vehicle-treated exposed mice (figure 7B). The total number of cells in

the BAL fluid and its composition was unchanged by plerixafor treatment (figure 7C-E). The

peri-bronchial fibrosis was significantly decreased in the lungs of plerixafor-treated exposed mice

(figure 7F-G). To identify pathways that were differentially regulated by the early COPD

protocol and plerixafor treatment, we performed proteome analysis on lungs obtained 1 day after

the last poly-IC instillation. Comparison analysis revealed a significant inhibition of pathways of

inflammation, cell invasion, movement, adhesion, and synthesis of reactive oxygen species

(ROS) in relation to plerixafor treatment (figure 7H). These changes were driven by down-

regulation of a core set of proteins such as thioredoxin interacting protein (TXNIP), matrix

metalloproteinase-8 (MMP-8) and vimentin (figure S13).

Treatment with plerixafor also reduced the development of RV hypertrophy in exposed mice

(figure 7I). The RVSP, as well as heart electrophysiological properties remained unchanged by

CXCR4 inhibition (figure 7J, figure S6M-R).

Flow cytometry 1 day after the last poly(I:C) instillation showed a significant decrease of the

numbers of both fibrocytes and CXCR4$^+$ fibrocytes in the lungs of exposed mice by plerixafor

(figure S14A-E). Four days after the last poly(I:C) instillation, the density of peri-bronchial

fibrocytes identified by immunohistochemistry was significantly reduced by plerixafor (figure

S14F-G).

Figure 7: Evaluation of lung function, inflammation, bronchial and heart remodeling upon pharmacological blockage of CXCR4 in experimental COPD. (A) Mice are exposed to cigarette smoke (CS), challenged with poly(I:C) and treated with plerixafor ("pleri", 1 mg/kg, purple squares) or its vehicle (grey squares). Mice are sacrificed at day 68 or 71 (1 day and 4 days after the last poly(I:C) instillation). (B) FEV0.05/FVC in the different conditions. Data represent individual mice. Vehicle: n=30, pleri: n=27. (C) Total cells count in bronchoalveolar lavage (BAL). Kruskal-Wallis test followed by Dunn's post-tests. (D-E) BAL relative (D) and absolute (E) cell composition respectively 1 day and 4 days after the last poly(I:C) instillation. d+1: Vehicle: n=13, Pleri: n=14; d+4: Vehicle: n=21, Pleri: n=26. (F) Representative Masson's trichrome staining to assess peri-bronchial fibrosis. (G) Standardized fibrosis, defined as (peri-bronchial fibrosis ("PF", %) - mean PFcontrol)/standard deviation PFcontrol. The peri-bronchial fibrosis ("PF", percentage) is defined by the ratio between the area of segmented pixels in the

area of analysis divided by the total peri-bronchial area. Vehicle: n=27, pleri: n=26. (H)
Heatmaps of significantly differentially regulated pathways identified by Ingenuity Pathway
analysis (IPA; Qiagen), obtained by proteomics comparison of CS+poly(I:C) (n=5) *vs* RA+PBS
(n=5)-exposed lungs (n=5) (all treated by vehicle), and plerixafor (n=5) vs vehicle (n=5)-treated
lungs (all exposed to CS+poly(I:C)). The colour scale indicates the Z-score value. (I) Fulton
index, defined as (right ventricle (RV)/ left ventricle plus septum (LV + S). Vehicle: n=31, pleri:
n=34. (J) Right ventricular systolic pressure (RVSP). Vehicle: n=11, pleri: n=15. (B, D, E, G, I,
J) *: P<0.05, **: P<0.01, ***: P<0.001. Unpaired t test or Mann-Whitney test.
458

DISCUSSION

Taken together, our data demonstrate that the CXCL12/CXCR4 axis plays a major role in the

pathogenesis of lung and heart tissue remodeling of early COPD. Both genetic invalidation and

pharmacological inhibition of CXCR4 in our experimental early COPD murine model reduced

both the level of CXCR4-positive cells in the peripheral circulation and fibrocyte recruitment into

the lungs, along with a proteomic signature consistent with decreased inflammation. As a

consequence, CXCR4 inhibition prevented airflow limitation and both lung and cardiac

remodeling. Our findings are likely to be translatable to humans as we showed that CXCL12 was

increased in the lung of early COPD patients and CXCR4 in the blood of COPD patients.

We designed the present dedicated murine model of early COPD to examine pathophysiological

processes at the onset of the disease. Instead of deciphering the relative contribution of CS and

poly-(I:C), this study aims at developing and using an early COPD model, based on the

following: first, 10-week old mice exposed to CS for 10 weeks would correspond to young

human adults chronically exposed to tobacco smoke for several years (22), and poly-(I:C)

administration mimics exacerbations occurring in early COPD at a frequency similar to the

COPD population (23). Moreover, poly-(I:C) allowed to mimic respiratory viral infections (20),

which are more frequent in smokers (29). Second, as in humans, exposed mice exhibited limited

but significant airflow limitation together with both neutrophilic/lymphocytic airway

inflammation in the BAL, allowing our design to model the bronchial component of COPD (30).

Lung histological observations evidenced peribronchial fibrosis, but no obvious signs of alveolar

destructions. Neither respiratory system compliance nor tissue elastance was different between

control and COPD animals. By contrast, the vast majority of murine COPD models are based on

long-term CS exposure (up to 6 months) either mimicking emphysema, or reproducing severe

airway obstruction together with emphysema-like changes (31).

The present model also includes features of early RV hypertrophy (32), thus confirming again the

relevance of this animal model of early COPD. Cardiovascular diseases are indeed frequently

seen in current smokers (33) and patients with COPD (34). Several studies have shown that

morphological changes of the right ventricle occur in the initial stages of COPD. Indeed, right

ventricular hypertrophy has been described as an early sign of adaptation of the right ventricle to

pressure overload in COPD, even in normoxemia or in mild hypoxemia conditions, and initially

without affecting right ventricular function (35). Development of right ventricular alterations has

also been evidenced very early when airway resistance was increased in preschool-aged children

with obstructive respiratory diseases (*i.e*., suffering from chronic coughing and wheezing) (36).

Although right ventricular hypertrophy is often described as a consequence of increased

pulmonary arterial pressures and development of pulmonary hypertension (PH), right ventricular

remodeling has been shown to occur in early stages of COPD before PH development (32, 35,

37). Our mouse model therefore reproduces such early right ventricular alterations regardless of

PH development, with significant right ventricular hypertrophy, but without modifications of

right ventricular systolic pressure, volume or function. As right ventricular systolic pressure

measurements performed in our mice COPD model might not completely reflect PAP,

subclinical/minor elevations of the PAP in our model of experimental COPD cannot be ruled out.

This hypothesis is in line with the findings of another COPD mice model, where elevations of

pressure and higher modification of cardiac remodeling have been found, presumably because of

longer time of CS exposure (6 months) (38). More likely, cardiac remodeling could emerge

secondary to alternative mechanisms including lung hyperinflation, systemic inflammation, and

endothelial dysfunction.

An important result of the present study is the increase of CXCL12 expression in the lungs of

early COPD mice and patients, along with an increase in $CXCR4^+$-cells in the blood of early

COPD mice as well as in COPD patients. The increased level of CXCR4 in both experimental

COPD and patients with COPD suggests a therapeutic opportunity to target CXCR4 for COPD

treatment. Treating mice with the CXCR4 antagonist plerixafor prevented the increase of

$CXCR4^+$-cells in the blood, attenuated tissue fibrocytes recruitment and lung function

degradation. The beneficial effect of plerixafor on lung function and remodeling in exposed mice,

exhibiting increased CXCR4 solely in the blood, was similar with that observed in $CXCR4^{-/-}$

mice. This suggests that such beneficial effects occurred through CXCR4 reduction in the blood

rather than in the lungs, which could encourage targeting CXCR4 systemically rather than locally

in the lung.

As CXCR4 expression is strongly associated with chemotactic responsiveness (10), CXCR4 up-

regulation is likely to result in a dynamic change of cell recruitment into the lungs. In support of

this hypothesis, we demonstrated an increased infiltration of fibrocytes into the lungs, that

accumulate into peribronchial areas at steady state in CS and poly(I:C)-exposed mice. The

elevated lung CXCL12 level in those mice and in patients with early COPD is also likely to

participate in fibrocyte chemotaxis from the general circulation to lung tissue, as well as in

fibrocyte increase in peribronchial areas, as the binding of CXCL12 on its receptor CXCR4

protects against cell death (39). Pharmacological and genetic targeting of CXCR4 reduced

CXCR4-expressing fibrocyte infiltration into the lungs during exacerbations-like events and

fibrocyte accumulation around bronchi at the steady state in COPD mice. These results are in

agreement with findings obtained in mice treated with another CXCR4 antagonist or with

CXCL12 neutralizing antibody, in the context of pulmonary fibrosis, where reduction of

fibrocyte accumulation has been found concomitantly of an improvement of lung outcomes (14,

40). Altogether, this indicates that fibrocytes could play a major role in the development of

peribronchial fibrosis, either directly through excessive repair processes, or indirectly by

paracrine activation of extracellular matrix-producing cells such as fibroblasts and myofibroblasts

(41). This could be partially mediated through MMP-8 production, that fibrocyte are able to

secrete (42), and that we evidenced as down-regulated by plerixafor treatment along with

fibrocyte level reduction. Additionally, and importantly, these fibrocytes could also perpetuate

chronic inflammation, through interaction with CD8$^+$ T cells. Such cells are indeed implicated in

COPD lung pathogenesis (43) and we have recently shown that fibrocytes directly interact with

CD8+ T cells in the lungs, leading to increased proliferation and activation (44).

Several other cell types might be implicated in plerixafor-mediated beneficial effects in the lungs,

including neutrophils. CXCL12 not only acts as a chemoattractant for neutrophils but also

suppresses neutrophils cellular death, allowing other functions for neutrophils such as the

production of neutrophil extracellular traps (NETs) (45). CXCR4-expressing neutrophils

accumulate in inflamed lungs and they are more prone to produce NETs (46), that may cause

tissue damage and contribute to bronchial obstruction. Neutrophils might also control the

secondary migration of other cells such as CD8$^+$ T cells by leaving CXCL12-containing trails

behind during their migration (47). This lead is worthy of investigation and could be the topic of

future studies.

Targeting CXCR4 with daily injections of plerixafor during the 5 last weeks of the protocol as

well as genetic deletion of CXCR4 decreased smoke induced-right heart remodeling. This

extends the findings of another study showing that neutralizing CXCL12 or antagonizing CXCR4

attenuates right ventricular hypertrophy in a rat model of severe pulmonary arterial hypertension

(48), and this suggests a general role of the CXCL12-CXCR4 axis in the development of

pulmonary hypertension. In the latter study (48), the beneficial effect of CXCL12 neutralization

has been attributed to a decrease of vascular pericyte coverage and macrophage infiltration into

the lungs, that deserves further investigation in the COPD model. As cardiac comorbidities have

a huge impact on the natural history of COPD, with a poor prognosis of COPD patients with

pulmonary artery pressure elevations (49), it reinforces the therapeutic interest of CXCR4 as a

target for COPD treatment.

Our findings obtained mainly in mice are likely to apply in humans for the following reasons: (i)

the expression patterns of CXCL12 are strikingly similar in murine and human lungs, (ii) the

CXCL12 and CXCR4 genes originate early in the phylogenetic tree and are highly conserved

among vertebrate species, especially between mice and humans (50–52). In the same page, the

degree of homology between human and murine CXCR4 and CXCL12 is 90% and 99%,

respectively (53). Last, the fact that plerixafor, a drug already used in clinics for CXCR4

antagonization (in the context of hematopoietic stem cells mobilization in the peripheral blood),

shows efficacy in our mouse model, further supports the translation of our results to humans.

Collectively, our data show that the CXCL12-CXCR4 axis plays important roles in the

pathogenesis of COPD and that inhibitors of the CXCL12-CXCR4 axis could open new

therapeutic options in this devastating disease, by improving lung function and decreasing cardiac

comorbidities. However, inhibiting the CXCL12-CXCR4 axis may also increase the severity of

exacerbations due to its central role in immune response. Therefore, further studies are required

to precise the balance between benefits and risks of targeting CXCR4 in COPD.

ACKNOWLEDGMENTS

We thank the study participants and the staff of the Thoracic Surgery, Radiology, Pathology, Respiratory, Lung Function Testing departments from the University Hospital of Bordeaux (Bordeaux, France), Fabienne Estella for technical assistance with mice protocol, Isabelle Goasdoue, Isabelle Bernis, Natacha Robert, Virginie Niel, and Marine Servat from the clinical investigation center for technical assistance, and Atika Zouine and Vincent Pitard for technical assistance at the Flow cytometry facility (CNRS UMS 3427, INSERM US 005, Univ. Bordeaux, F-33000 Bordeaux, France), Anne-Aurélie Raymond for help with proteomic analysis at the Oncoprot Platform (TBMCore US005). Christel Poujol, Sébastien Marais and Fabrice Cordelières for help with imaging and image analysis et the Bordeaux Imaging Centre (BIC; Bordeaux, France). Microscopy was performed at BIC, a service unit of the CNRS-INSERM and Bordeaux University, a member of the national BioImaging infrastructure of France supported by the French National Research Agency (ANR-10-INBS-04).

The project was supported by a grant from the "Fondation de l'Université de Bordeaux" (Fonds pour les maladies chroniques nécessitant une assistance médico-technique FGLMR/AVAD), a grant from « Aquitaine Science Transfert » (PB) and an unrestricted grant from AstraZeneca, the "Investments in the Future" program managed by the "Agence Nationale de la Recherche" (ANR, grant reference ANR-10-IAHU-04). The COBRA cohort was funded by AstraZeneca, Chiesi, Glaxo-SmithKline, Novartis, Chiesi, Roche and Legs Poix fondation.

Data sharing

Qualified researchers can request access to all data through the corresponding author.

Contributors

- conception and design (ID, PB)

- data acquisition (ID, PH, EM, RA, SC, DEH, EE, PE, MC, MD, GC, CB, OO, JWD,

TT, VFM)

- data analysis (ID, PH, EM, RA, SC, DEH, RP, MD, GC, CB, JWD, VFM, PB)

- data interpretation (ID, PH, EM, RA, SC, DEH, EE, RP, PE, MD, GC, CB, TT, RM,

MZ, VFM, PB)

- resources (PH, MC, MD, MT, HB, POG, MZ, PB)

- drafting the manuscript (ID, PH, PB)

- revision and final approval of the version to be published, agreement to be accountable

for all aspects of the work in ensuring that questions related to the accuracy or integrity

of any part of the work are appropriately investigated and resolved (All)

Declaration of interests

ID has 2 patents delivered (i) (EP N°3050574 *i.e.*, Use of plerixafor for treating and/or preventing

acute exacerbations of chronic obstructive pulmonary disease); (ii) (EP N°20173595.8 *i.e.*, New

compositions and methods of treating COVID-19 Disease). ID report a grant from the

"Fondation Bordeaux Université," with funding from "Assistance Ventilatoire à Domicile"

(AVAD) and "Fédération Girondine de Lutte contre les Maladies Respiratoires" (FGLMR).

PH reports non-financial support from AVAD, Chiesi and GlaxoSmithKline, outside the

submitted work and a grant from the "Fondation Bordeaux Université," with funding from

AVAD and FGLMR.

POG reports grants, personal fees and non-financial support from AstraZeneca, personal fees and non-financial support from Chiesi, personal fees and non-financial support from GlaxoSmithKline, personal fees and non-financial support from Novartis, personal fees and non-financial support from Sanofi, outside the submitted work. POG has 2 patents delivered (i) (EP N°3050574 *i.e*., Use of plerixafor for treating and/or preventing acute exacerbations of chronic obstructive pulmonary disease); (ii) (EP N°20173595.8 *i.e*., New compositions and methods of treating COVID-19 Disease).

MZ reports personal fees from AstraZeneca, Boehringer Ingelheim, CSL Behring, Novartis, Chiesi, GlaxoSmithKline and non-financial support Lilly outside the submitted work and a grant from the "Fondation Bordeaux Université," with funding from AVAD and FGLMR.

PB is the medical coordinator of the French national cohort (*i.e*., COBRA), which received grants from AstraZeneca, GlaxoSmithKine, and Chiesi. Moreover, PB reports grants and personal fees from Novartis, personal fees and non-financial support from Chiesi, grants, personal fees and non-financial support from Boehringer Ingelheim, grants, personal fees and non-financial support from AstraZeneca, personal fees and non-financial support from GSK, personal fees and non-financial support from Sanofi, outside the submitted work; in addition, PB has 4 patents delivered (i) (EP N°3050574 *i.e*., Use of plerixafor for treating and/or preventing acute exacerbations of chronic obstructive pulmonary disease); (ii) (EP N°20173595.8 *i.e*., New compositions and methods of treating COVID-19 Disease); (iii) (N° WO2017203064A1 *i.e*., MRI Method for the geometrical characterization of pulmonary airways); (iv) (N° WO2021018439A1 *i.e*., Method for generating a biomarker system). All other authors declare they have no competing interests.

REFERENCES

1. Global Initiative for Chronic Obstructive Lung Disease. *Global Initiative for Chronic Obstructive Lung*

Disease - GOLD at <https://goldcopd.org/>.

2. Mannino DM, Buist AS. Global burden of COPD: risk factors, prevalence, and future trends. *Lancet*

2007;370:765–773.

3. Martinez FJ, Han MK, Allinson JP, Barr RG, Boucher RC, Calverley PMA, *et al.* At the Root: Defining

and Halting Progression of Early Chronic Obstructive Pulmonary Disease. *Am J Respir Crit Care Med*

2018;197:1540–1551.

4. Çolak Y, Afzal S, Nordestgaard BG, Lange P, Vestbo J. Importance of Early COPD in Young Adults for

Development of Clinical COPD: Findings from the Copenhagen General Population Study. *Am J Respir*

Crit Care Med 2021;203:1245–1256.

5. Martinez FJ, Agusti A, Celli BR, Han MK, Allinson JP, Bhatt SP, *et al.* Treatment Trials in Young

Patients with Chronic Obstructive Pulmonary Disease and Pre-Chronic Obstructive Pulmonary

Disease Patients: Time to Move Forward. *Am J Respir Crit Care Med* 2022;205:275–287.

6. Henrot P, Prevel R, Berger P, Dupin I. Chemokines in COPD: From Implication to Therapeutic Use. *Int*

J Mol Sci 2019;20:.

7. Dupin I, Allard B, Ozier A, Maurat E, Ousova O, Delbrel E, *et al.* Blood fibrocytes are recruited during

acute exacerbations of chronic obstructive pulmonary disease through a CXCR4-dependent pathway.

J Allergy Clin Immunol 2016;137:1036-1042.e7.

8. Dupin I, Thumerel M, Maurat E, Coste F, Eyraud E, Begueret H, *et al.* Fibrocyte accumulation in the

airway walls of COPD patients. *Eur Respir J* 2019;54:.

9. Busillo JM, Benovic JL. Regulation of CXCR4 signaling. *Biochimica et Biophysica Acta (BBA) -*

Biomembranes 2007;1768:952–963.

10. Schioppa T, Uranchimeg B, Saccani A, Biswas SK, Doni A, Rapisarda A, *et al.* Regulation of the

Chemokine Receptor CXCR4 by Hypoxia. *Journal of Experimental Medicine* 2003;198:1391–1402.

11. Tian X, Xie G, Xiao H, Ding F, Bao W, Zhang M. CXCR4 knockdown prevents inflammatory cytokine

expression in macrophages by suppressing activation of MAPK and NF-κB signaling pathways. *Cell &*

Bioscience 2019;9:55.

12. Petty JM, Sueblinvong V, Lenox CC, Jones CC, Cosgrove GP, Cool CD, *et al.* Pulmonary stromal-

derived factor-1 expression and effect on neutrophil recruitment during acute lung injury. *J Immunol*

2007;178:8148–8157.

13. Barwinska D, Oueini H, Poirier C, Albrecht ME, Bogatcheva NV, Justice MJ, *et al.* AMD3100

ameliorates cigarette smoke-induced emphysema-like manifestations in mice. *Am J Physiol Lung Cell*

Mol Physiol 2018;315:L382–L386.

14. Phillips RJ, Burdick MD, Hong K, Lutz MA, Murray LA, Xue YY, *et al.* Circulating fibrocytes traffic to the

lungs in response to CXCL12 and mediate fibrosis. *J Clin Invest* 2004;114:438–446.

15. Dupin I, Henrot P, Maurat E, Eyraud E, Esteves P, Zysman M, *et al.* Targeting CXCR4 as a therapeutic

strategy to improve outcomes in a mouse model of early chronic obstructive pulmonary disease

(COPD). *ERJ Open Research* 2022;8:.

16. Dupin I, Dubreuil M, Freund-Michel V, Maurat E, Campagnac M, Ousova O, *et al.* Targeting CXCR4 as

a therapeutic strategy to improve outcomes in a mouse model of early chronic obstructive

pulmonary disease (COPD). *ERJ Open Research* 2019;5:.

17. Tokoyoda K, Egawa T, Sugiyama T, Choi B-I, Nagasawa T. Cellular Niches Controlling B Lymphocyte

Behavior within Bone Marrow during Development. *Immunity* 2004;20:707–718.

18. Jullien N, Goddard I, Selmi-Ruby S, Fina J-L, Cremer H, Herman J-P. Use of ERT2-iCre-ERT2 for

conditional transgenesis. *Genesis* 2008;46:193–199.

19. Almolki A, Guenegou A, Golda S, Boyer L, Benallaoua M, Amara N, *et al*. Heme oxygenase-1 prevents

airway mucus hypersecretion induced by cigarette smoke in rodents and humans. *Am J Pathol*

2008;173:981–992.

20. Kang M-J, Lee CG, Lee J-Y, Dela Cruz CS, Chen ZJ, Enelow R, *et al*. Cigarette smoke selectively

enhances viral PAMP- and virus-induced pulmonary innate immune and remodeling responses in

mice. *J Clin Invest* 2008;118:2771–2784.

21. Esteves P, Blanc L, Celle A, Dupin I, Maurat E, Amoedo N, *et al*. Crucial role of fatty acid oxidation in

asthmatic bronchial smooth muscle remodelling. *Eur Respir J* 2021;58:2004252.

22. Dutta S, Sengupta P. Men and mice: Relating their ages. *Life Sciences* 2016;152:244–248.

23. Cosío BG, Casanova C, Soler-Cataluña JJ, Soriano JB, García-Río F, de Lucas P, *et al*. Unravelling young

COPD and pre-COPD in the general population. *ERJ Open Res* 2023;9:00334–02022.

24. Hogg JC, McDonough JE, Sanchez PG, Cooper JD, Coxson HO, Elliott WM, *et al*. Micro–Computed

Tomography Measurements of Peripheral Lung Pathology in Chronic Obstructive Pulmonary Disease.

Proc Am Thorac Soc 2009;6:546–549.

25. Sangaletti S, Tripodo C, Cappetti B, Casalini P, Chiodoni C, Piconese S, *et al*. SPARC oppositely

regulates inflammation and fibrosis in bleomycin-induced lung damage. *Am J Pathol* 2011;179:3000–

3010.

26. Aono Y, Kishi M, Yokota Y, Azuma M, Kinoshita K, Takezaki A, *et al*. Role of platelet-derived growth

factor/platelet-derived growth factor receptor axis in the trafficking of circulating fibrocytes in

pulmonary fibrosis. *Am J Respir Cell Mol Biol* 2014;51:793–801.

27. Mitsuhashi A, Goto H, Saijo A, Trung VT, Aono Y, Ogino H, *et al*. Fibrocyte-like cells mediate acquired

resistance to anti-angiogenic therapy with bevacizumab. *Nat Commun* 2015;6:8792.

28. Tachibana K, Hirota S, Iizasa H, Yoshida H, Kawabata K, Kataoka Y, *et al*. The chemokine receptor

CXCR4 is essential for vascularization of the gastrointestinal tract. *Nature* 1998;393:591–594.

29. Arcavi L, Benowitz NL. Cigarette Smoking and Infection. *Archives of Internal Medicine*

2004;164:2206–2216.

30. Higham A, Quinn AM, Cançado JED, Singh D. The pathology of small airways disease in COPD:

historical aspects and future directions. *Respir Res* 2019;20:49.

31. Tanner L, Single AB. Animal Models Reflecting Chronic Obstructive Pulmonary Disease and Related

Respiratory Disorders: Translating Pre-Clinical Data into Clinical Relevance. *JIN* 2020;12:203–225.

32. Hilde JM, Skjørten I, Grøtta OJ, Hansteen V, Melsom MN, Hisdal J, *et al.* Right ventricular dysfunction

and remodeling in chronic obstructive pulmonary disease without pulmonary hypertension. *J Am*

Coll Cardiol 2013;62:1103–1111.

33. Banks E, Joshy G, Korda RJ, Stavreski B, Soga K, Egger S, *et al.* Tobacco smoking and risk of 36

cardiovascular disease subtypes: fatal and non-fatal outcomes in a large prospective Australian

study. *BMC Medicine* 2019;17:128.

34. Chen W, Thomas J, Sadatsafavi M, FitzGerald JM. Risk of cardiovascular comorbidity in patients with

chronic obstructive pulmonary disease: a systematic review and meta-analysis. *Lancet Respir Med*

2015;3:631–639.

35. Vonk-Noordegraaf A, Marcus JT, Holverda S, Roseboom B, Postmus PE. Early changes of cardiac

structure and function in COPD patients with mild hypoxemia. *Chest* 2005;127:1898–1903.

36. Manti S, Parisi GF, Giacchi V, Sciacca P, Tardino L, Cuppari C, *et al.* Pilot study shows right ventricular

diastolic function impairment in young children with obstructive respiratory disease. *Acta*

Paediatrica 2019;108:740–744.

37. Vonk-Noordegraaf A, Haddad F, Chin KM, Forfia PR, Kawut SM, Lumens J, *et al.* Right heart

adaptation to pulmonary arterial hypertension: physiology and pathobiology. *J Am Coll Cardiol*

2013;62:D22-33.

38. Weissmann N, Lobo B, Pichl A, Parajuli N, Seimetz M, Puig-Pey R, *et al.* Stimulation of soluble

guanylate cyclase prevents cigarette smoke-induced pulmonary hypertension and emphysema. *Am J*

Respir Crit Care Med 2014;189:1359–1373.

39. Suzuki Y, Rahman M, Mitsuya H. Diverse transcriptional response of CD4+ T cells to stromal cell-

derived factor SDF-1: cell survival promotion and priming effects of SDF-1 on CD4+ T cells. *J Immunol*

2001;167:3064–3073.

40. Griffiths K, Habiel DM, Jaffar J, Binder U, Darby WG, Hosking CG, *et al.* Anti-fibrotic Effects of CXCR4-

Targeting i-body AD-114 in Preclinical Models of Pulmonary Fibrosis. *Sci Rep* 2018;8:3212.

41. Dupin I, Contin-Bordes C, Berger P. Fibrocytes in Asthma and Chronic Obstructive Pulmonary

Disease: Variations on the Same Theme. *Am J Respir Cell Mol Biol* 2018;58:288–298.

42. García-de-Alba C, Becerril C, Ruiz V, González Y, Reyes S, García-Alvarez J, *et al.* Expression of matrix

metalloproteases by fibrocytes: possible role in migration and homing. *Am J Respir Crit Care Med*

2010;182:1144–1152.

43. Roos-Engstrand E, Ekstrand-Hammarström B, Pourazar J, Behndig AF, Bucht A, Blomberg A. Influence

of smoking cessation on airway T lymphocyte subsets in COPD. *COPD* 2009;6:112–120.

44. Eyraud E, Maurat E, Sac-Epée J-M, Henrot P, Zysman M, Esteves P, *et al.* Short-range interactions

between fibrocytes and CD8+ T cells in COPD bronchial inflammatory response. *Elife*

2023;12:RP85875.

45. Yamada M, Kubo H, Kobayashi S, Ishizawa K, He M, Suzuki T, *et al.* The increase in surface CXCR4

expression on lung extravascular neutrophils and its effects on neutrophils during endotoxin-induced

lung injury. *Cell Mol Immunol* 2011;8:305–314.

46. Radermecker C, Sabatel C, Vanwinge C, Ruscitti C, Maréchal P, Perin F, *et al.* Locally instructed

CXCR4hi neutrophils trigger environment-driven allergic asthma through the release of neutrophil

extracellular traps. *Nat Immunol* 2019;20:1444–1455.

47. Lim K, Hyun Y-M, Lambert-Emo K, Capece T, Bae S, Miller R, *et al.* Neutrophil trails guide influenza-

specific CD8[+] T cells in the airways. *Science* 2015;349:aaa4352.

48. Bordenave J, Thuillet R, Tu L, Phan C, Cumont A, Marsol C, *et al.* Neutralization of CXCL12 attenuates

established pulmonary hypertension in rats. *Cardiovasc Res* 2019;doi:10.1093/cvr/cvz153.

49. Oswald-Mammosser M, Weitzenblum E, Quoix E, Moser G, Chaouat A, Charpentier C, *et al.*

Prognostic factors in COPD patients receiving long-term oxygen therapy. Importance of pulmonary

artery pressure. *Chest* 1995;107:1193–1198.

50. Nomiyama H, Osada N, Yoshie O. A family tree of vertebrate chemokine receptors for a unified

nomenclature. *Developmental & Comparative Immunology* 2011;35:705–715.

51. Zlotnik A, Yoshie O. Chemokines: a new classification system and their role in immunity. *Immunity*

2000;12:121–127.

52. Aleotti A, Goulty M, Lewis C, Giorgini F, Feuda R. The origin, evolution, and molecular diversity of the

chemokine system. *Life Science Alliance* 2024;7:.

53. Nagasawa T, Tachibana K, Kishimoto T. A novel CXC chemokine PBSF/SDF-1 and its receptor CXCR4:

their functions in development, hematopoiesis and HIV infection. *Seminars in Immunology*

1998;10:179–185.