CXCR4 blockade alleviates pulmonary and cardiac outcomes in young COPD

Isabelle Dupin,[1,2] Pauline Henrot,[1,2,3] Elise Maurat,[1,2] Reshed Abohalaka,[1,2] Sébastien Chaigne,[1,2,3] Dounia El Hamrani,[1,2], Edmée Eyraud,[1,2], Renaud Prevel,[1,2,3] Pauline Esteves,[1,2] Maryline Campagnac,[1,2] Marielle Dubreuil,[1,2] Guillaume Cardouat,[1,2] Clément Bouchet,[1,2] Olga Ousova,[1,2], Jean-William Dupuy,[1,2] Thomas Trian,[1,2] Matthieu Thumerel,[1,2,3] Hugues Bégueret,[1,2,3] Pierre-Olivier Girodet,[1,2,3] Roger Marthan,[1,2,3] Maeva Zysman,[2,3] Véronique Freund-Michel,[1,2], Patrick Berger,[1,2,3]

[1] Univ. Bordeaux, Centre de Recherche Cardio-thoracique de Bordeaux, INSERM U1045, IHU Liryc, CIC 1401, Proteomics Facility, F-33600 Pessac, France

[2] INSERM, Centre de Recherche Cardio-thoracique de Bordeaux, U1045, CIC 1401, F-33600 Pessac, France

[3] CHU Bordeaux, Service d'exploration fonctionnelle respiratoire, Service de réanimation, Service de pneumologie, Service de chirurgie thoracique, Service de cardiologie-électrophysiologie et stimulation cardiaque, Service d'anatomopathologie, CIC-P 1401, F-33600 Pessac, France

Corresponding author: Prof. Isabelle Dupin
Centre de Recherche Cardio-thoracique de Bordeaux, INSERM U1045
PTIB. Hôpital Xavier Arnozan. Avenue du Haut Lévêque. 33600 Pessac, FRANCE
Tel : +33 5 47 30 27 51
e-mail : isabelle.dupin@u-bordeaux.fr

ABSTRACT

Background: Chronic obstructive pulmonary disease (COPD) is a prevalent respiratory disease lacking effective treatment. Focusing on young COPD should help to discover disease modifying therapies. We aimed to examine the role of the CXCL12/CXCR4 axis in young COPD from both human samples and murine models.

Methods: Blood samples and lung tissues of young COPD patients and controls were obtained in order to analyse CXCL12 and CXCR4 levels. To generate a young COPD model, ten-week-old mice were exposed to cigarette smoke (CS) for 10 weeks and intranasal instillations of polyinosinic–polycytidylic acid (poly(I:C)) for the last 5 weeks to mimic exacerbations.

Results: CXCR4 expressing cells number was increased in the blood of patients with COPD, as well as in the blood of exposed mice. Lung CXCL12 expression was higher in both young COPD patients and exposed mice. Exposed mice presented mild airway obstruction, peri-bronchial fibrosis and right heart thickening. The density of fibrocytes expressing CXCR4 was increased in the bronchial submucosa of exposed mice. Conditional inactivation of CXCR4 at adult stage as well as pharmacological inhibition of CXCR4 with plerixafor injections improved lung function, reduced inflammation and protected against CS and poly-(I:C)-induced airway and cardiac remodelling. CXCR4$^{-/-}$ and plerixafor-treated mice also had less CXCR4-expressing circulating cells and a lower density of peri-bronchial fibrocytes.

Conclusion: We demonstrate that targeting CXCR4 has beneficial effects in an animal model of young COPD and provide a framework to translate these preclinical findings to clinical settings in a drug repurposing approach.

Abstract word count: 244 words /250

What is already known on this topic

Whereas the CXCL12/CXCR4 axis has already been identified in COPD pathophysiology,

preclinical evidences supporting a beneficial role of CXCR4 antagonists for COPD treatment are

lacking.

What this study adds

CXCL12 and CXCR4 are upregulated in the lung and in the blood of patients with young COPD,

respectively. Genetic and pharmacological inhibition of CXCR4 in experimental young COPD

mice model reduces the number of CXCR4-expressing cells in the peripheral circulation and

fibrocyte recruitment into the lungs, along with a proteomic signature consistent with a decrease

of inflammation. Overall, it improves lung function and cardiac tissue remodelling.

How this study might affect research, practice or policy

CXCR4 inhibitors may be therapeutically exploited to slow down the progression of young

COPD.

Key words: fibrocytes, bronchial remodelling, inflammation, obstruction, animal model

Abbreviations

BAL   Broncho-alveolar lavage

bFGF   basic Fibroblast Growth Factor

BSM   Bronchial smooth muscle

CS   Cigarette smoke

CT   Computed tomographic

ECG   Electrocardiogram

FEV1   Forced expiratory volume in 1 second

FEV0.05   Forced expiratory volume in 0.05 second

FVC   Forced vital capacity

IFN   Interferon

LV + S   Left ventricle plus septum

MRI   Magnetic Resonance Imaging

Poly-(I:C)   Polyinosinic–polycytidylic acid

RA   Room air

RV   Right ventricle

RVSP   Right ventricular systolic pressure

VEGF   Vascular Endothelial Growth Factor

INTRODUCTION

Chronic obstructive pulmonary disease (COPD) is a common disease, characterized by persistent

respiratory symptoms and airflow limitation [1]. Major risk factors for COPD include chronic

exposure to noxious particles, mainly cigarette smoke (CS), and airway infections [2]. Young

COPD has been recently defined as patients aged <50 years, with a smoking exposure >10 pack-

years, and a ratio between forced expiratory volume in 1 second (FEV1) and forced vital capacity

(FVC) less than 70% [1,3]. Therapeutic interventions in young COPD patients could help to

tackle the disease before reaching irreversible tissue damage [3]. However, it is extremely

difficult to enrol those patients into trials and collect their bronchial samples, since they are rarely

diagnosed, demonstrating the urgent need to develop animal models of young COPD.

Exposure of lung structural and immune cells to CS and infectious agents results in the release of

various inflammatory substances, including chemokines [4]. The chemokine receptor CXCR4

and its associated ligand CXCL12 appear to be attractive therapeutic targets for young COPD

since (i) they are implicated in the migration of inflammatory cells in COPD lungs [5], (ii) their

expression and function are controlled by other cytokines [6], oxygen concentration [7] and

microbial agents [8,9], which are disrupted in COPD, and (iii) pharmacological blockade of

CXCR4 reduces emphysema development in a long-term COPD murine model [10]. CXCR4 is

notably expressed by fibrocytes, a rare population of circulating fibroblast-like cells [11], that are

assumed to play a crucial role in COPD [5,12].

In the present study, we thus hypothesized that the CXCL12/CXCR4 axis plays a role in the

pathological processes leading to COPD development during early adulthood and, hence, that

inhibiting CXCR4 would protect from CS-induced airflow limitation and tissue remodelling. In a

young COPD patients' samples, as well as in murine model of young COPD, both CXCR4-

expressing cells in the blood and CXCL12 in the lung were upregulated. Using transgenic mice

deficient for CXCR4 or the CXCR4 antagonist plerixafor, we showed that lungs were protected

against CS-induced both lung function alteration and fibrocytes accumulation, and that the right

heart was resistant to cardiac remodelling.

Some of these findings have been previously reported in abstract form [13,14].

MATERIALS AND METHODS

Detailed methods are described in the online supplementary material.

Study Population

Human lung tissues were obtained from both the "Fibrochir" study (NCT01692444) [12] and

"TUBE" (*i.e*., TissUs Bronchiques et PulmonairEs, sponsored by the University hospital of

Bordeaux) biological collection (table S1). According to the French law and the MR004

regulation, patients received an information form, allowing them to refuse the use of their

surgical samples for research. Blood samples of COPD patients were obtained from the

"COBRA" cohort (table S2). Research was conducted according to the principles of the World

Medical Association Declaration of Helsinki. The studies received approval from the local and

national ethics committees.

Mouse model of young COPD

All animal studies were performed according to European directives for the protection of

vertebrate animals and to the Declaration of Helsinki conventions for the use and care of animals.

Agreement was obtained from French authorities (number A33-063-907) and all the protocols

were approved by the local ethics committee.

Ten-week-old mice were exposed for 10 weeks (whole-body) to either room air (RA) or CS, as

described by Almolki *et al*. [15]. After 5 weeks of CS exposure, 50 μg of poly(I:C) or its vehicle

control was administered twice per week for 5 weeks via nasal aspiration.

CXCR4 genetic deletion was induced by tamoxifen. CXCR4 pharmacological inhibition was

induced by subcutaneous injection of 1 mg/kg plerixafor, 5 times/week during the last 5 weeks of

the protocol.

Statistical Analysis

Statistical significance, defined as $P < 0.05$, was assessed by t-tests and ANOVA for variables

with parametric distribution, and by Kruskal-Wallis with multiple comparison z tests, Mann-

Whitney tests, Wilcoxon tests and Spearman correlation coefficients for variables with non-

parametric distribution.

RESULTS

CXCL12 and CXCR4 expression is increased in lung and blood, respectively, in patients

with COPD

First, we determined whether the expression of CXCL12 and CXCR4 was altered in patients with

COPD by interrogating RNAseq data from lung of control subjects and COPD patients

(GSE76925 data set, http://www.ncbi.nlm.nih.gov/geo/). We found that CXCL12 expression was

significantly increased in the lung of patients with COPD (GSE76925 data set, figure 1A). In a

separate cohort of patients with and without young COPD (table S1), CXCL12 was

predominantly localized to bronchial epithelial cells and peri-bronchial infiltrating cells, which

are likely immune cells (figure 1B). The CXCL12 surface immunostaining was significantly

increased in lungs of patients with young COPD in comparison to control lungs (figure 1B-C).

CXCR4 level was not significantly altered in the lungs of patients with COPD compared to those

of control smokers at the mRNA level (GSE76925 data set, figure 1D), which was corroborated

by a similar result at the protein level by quantification of immunostaining (figure 1E) in patients

with and without young COPD. In the blood, CXCR4 was significantly increased, at the mRNA

level, in patients with COPD compared to control subjects (figure S1A). In a separate cohort of

moderate to mild COPD patients (GOLD I to II, as defined by [1], table S2), the percentage of

CXCR4-expressing cells was also increased in the blood of moderate COPD patients in

comparison with mild COPD patients (figure S1B-C).

Chronic CS and poly(I:C) exposure induces functional obstruction, lung inflammation and

both lung and heart tissue remodelling in mice

To model young COPD, we used mice at the beginning of adulthood (*i.e*., 10-week old) [16],

which were exposed to a combination of CS exposure and intranasal instillations of poly(I:C) in

order to mirror the effects of smoking and exacerbations [17]. CS exposure for 5 weeks combined

with 4 doses of poly(I:C) were not sufficient to trigger functional airway obstruction (figure S2A-

D). In contrast, 10 weeks of CS exposure with 10 repeated instillations of poly(I:C) during the

last 5 weeks, significantly decreased the FEV0.05/FVC ratio (figure 2A-C). Other lung function

parameters, in particular respiratory system compliance and tissue elastance, were not modified

by CS and poly(I:C) exposure (figure S2E-J). At the end of the protocol, the weight gain of the

exposed mice was significantly different from that of the control mice (figure S3A-B). CS and

poly(I:C) exposure increased total cell count in the BAL fluid, as well as the absolute neutrophil,

lymphocyte and macrophage numbers, 1 day after the last poly(I:C) instillation (figure 2D-F).

Although lower, the increase in neutrophils and lymphocytes persisted 4 days after the last

poly(I:C) instillation (figure 2D-F). CS and poly(I:C) synergistically enhanced airflow

obstruction and BAL inflammation (figure S4). We also analysed lung proteomics obtained 1 day

after the last poly-IC instillation. Of the 137 pathways that were significantly different in young

COPD vs control mice, largest upregulated genes were those of the hypercytokinemia/

hyperchemokinemia box and of the pathogenesis of influenza and interferon (IFN) signalling

(figure 2G-H). In contrast, inhibited pathways were the wound healing and the xenobiotic

metabolism PXR signalling pathways (figure 2G-H). Exposed mice had a moderate but

significant increase in peri-bronchial fibrosis compared to control mice (figure 2I-J), mimicking

the thickening of distal airway tissue in COPD patients [18].

Then, we investigated the impact of CS and poly(I:C) exposure on cardiac function. There was no

evidence of heart failure in our model (figure S5A-B). Moreover, there was no sign of heart

rhythm disorder as assessed by ECG monitoring (figure S5C and figure S6A-F). Using MRI, we

also found that CS and poly(I:C)-exposition did not alter RV or LV volumes (figure S5D-F).

However, we detected a significant increase of RV wall thickness in exposed mice, indicating RV

hypertrophy (figure 2K-L). This finding was confirmed by the measurement of the Fulton index

(figure 2M). Of note, the RV systolic pressure (RVSP) did not differ between young COPD and

control mice (figure 2N).

CXCR4 and CXCL12 expression is increased in blood and lung, respectively, in

experimental COPD

The percentage of CXCR4-expressing cells was significantly increased in the blood of CS and

poly(I:C)-exposed mice compared to control mice, 1 day after the last poly(I:C) instillation

(figure 3A-B), indicating that experimental COPD replicated the increased levels of CXCR4 in

the blood of COPD patients. This effect appeared to be due to a synergy between CS and

poly(I:C) (figure S7). The level of CXCR4$^+$ circulating cells decreased and became similar

between control and exposed mice 4 days after the last poly(I:C) instillation (figure 3B). No

significant modification of CXCL12 plasma concentration was evidenced in mice (figure 3C).

The percentage of CXCR4-expressing cells was not significantly altered in whole lung

homogenates in experimental young COPD (figure 3D-E). In lung homogenates, CS and

poly(I:C)-exposure significantly increased CXCL12 protein levels compared to control mice, 1

and 4 days after the last poly(I:C) instillation (figure 3F). In good agreement with our findings in

patients' lungs, CXCL12 was expressed by bronchial epithelial cells and peri-bronchial

infiltrating cells (figure 3G), with a higher expression in lungs of animals exposed to CS and

poly-(I:C) in comparison to control lungs (figure 3H).

Fibrocytes and CXCR4$^+$ fibrocytes are increased in lungs of experimental young COPD

The percentage of fibrocytes, defined as cells expressing CD45 and a high level of FSP1

("CD45$^+$ FSP1^{high}") [19], was significantly increased in CS and poly(I:C)-exposed lungs, 1 day

after the last poly(I:C) instillation (figure 4A-C). The average percentage of leukocytes (CD45$^+$

cells) in control and exposed lungs was 69.2 % and 74.0 %, respectively (p=0.07, figure 4D). In

control and experimental young COPD, a vast majority of lung fibrocytes expressed CXCR4

(96.1 and 96.6 %, respectively, figure 4B) and the percentage of CXCR4$^+$ fibrocytes was

significantly increased in the lungs of exposed mice (figure 4C). Four days after the last poly(I:C)

instillation, the total level of fibrocytes decreased and became similar between control and

experimental young COPD lungs (figure 4C). Since the flow cytometry approach limits our

analysis to the whole lung, we co-stained FSP1 with CD45 in murine lungs, 4 days after the last

poly(I:C) instillation (figure 4F) using immunohistochemistry. Similarly to that previously

reported in human lung [12], the density of peri-bronchial fibrocytes was higher in CS and

poly(I:C)-exposed murine lungs than in control lungs (figure 4G). Moreover, we also observed *in*

situ CXCR4-expressing fibrocytes in peri-bronchial areas of human lung samples ("Fibrochir"

study [12], figure 4H). Of note, fibrocytes were found in right and left ventricles, but their

densities were unchanged by CS and poly(I:C) exposure (figure S8).

CXCR4 deletion protects against functional airway obstruction, lung and heart tissue

remodelling, and inhibits fibrocyte recruitment into the lungs in experimental young COPD

As mice genetically deficient for CXCR4 die perinatally [20], we generated conditional CXCR4

mutants using floxed allele [21] and a Cre2ERT2 strain [22], thereby obtaining mutants

harbouring a specific inactivation of CXCR4 at young adulthood after tamoxifen treatment.

Intraperitoneal tamoxifen administration, for 5 consecutive days, in CXCR4$^{lox/lox}$/Cre2ERT2 mice

was not sufficient to induce a large CXCR4 downregulation, in particular in the lungs (figure

S9A-B). Therefore, we combined intraperitoneal tamoxifen administration and feeding mice with

tamoxifen-containing food to obtain a more robust decrease in CXCR4 expression in the lung and

bone marrow (figure S9C-E). As tamoxifen-containing food reduced weight gain in mice (figure

S3C-D), we used CXCR4$^{lox/lox}$ mice as control mice, and gave all transgenic mice tamoxifen-

containing food.

CXCR4$^{lox/lox}$/Cre2ERT2 and CXCR4$^{lox/lox}$ mice were exposed to CS and poly(I:C) for 10 weeks

(figure 5A). As expected, CXCR4 expression decreases in the lung and blood of

CXCR4$^{lox/lox}$/Cre2ERT2 mice (figure S10). Moreover, FEV0.05/FVC ratio was higher in

CXCR4$^{-/-}$ mice compared to CXCR4$^{lox/lox}$ mice, when exposed to CS and poly(I:C) (figure 5B).

In the BAL fluid, CXCR4 deletion induced a significant decrease of macrophage and increase of

neutrophil percentages, 1 day after the last poly(I:C) instillation (figure 5C-E). CXCR4$^{-/-}$ mice

had a significant reduction in collagen deposition around small airways, compared to CS and

poly(I:C)-exposed CXCR4$^{+/+}$ mice (figure 5F-G), confirming an important role of CXCR4 in CS-

induced bronchial remodelling.

We then assessed the impact of CXCR4 deletion on right heart remodelling. CXCR4 deletion did

not affect electrophysiological and hemodynamic (figure S6G-L and figure S11) but it

significantly reduced the RV wall thickness as assessed by heart MRI *in vivo* and the Fulton

index *ex vivo* in exposed mice (figure 5H-J). RVSP value was unchanged by CXCR4 deletion

(figure 5K).

Moreover, we observed reduced flow cytometry levels of CXCR4^{+} fibrocytes in CS and

poly(I:C)-exposed whole lungs of CXCR4$^{-/-}$ mice compared with CXCR4$^{+/+}$ exposed mice, 1 day

after the last poly(I:C) instillation (figure 6A-D). Four days after the last poly(I:C) instillation, the

density of fibrocytes was also significantly decreased around the small airways (figure 6F-G).

Pharmacological inhibition of CXCR4 protects against functional obstruction and inhibits

fibrocyte recruitment into the lungs in experimental COPD

We then assessed the efficacy of a CXCR4 antagonist (*i.e*. AMD3100, plerixafor) in the young

COPD murine model (figure 7A). Mice weight was not affected by daily subcutaneous plerixafor

treatment during the second half of the protocol (figure S3E-F). Plerixafor significantly reduced

CXCR4 expression 1 day after the last poly(I:C) instillation in circulating cells, but not in lung

cells (figure S12). In exposed-mice treated with plerixafor, the FEV0.05/FVC ratio was

significantly higher than vehicle-treated exposed mice (figure 7B). The total number of cells in

the BAL fluid and its composition was unchanged by plerixafor treatment (figure 7C-E). The

peri-bronchial fibrosis was significantly decreased in the lungs of plerixafor-treated exposed mice

(figure 7F-G). To identify pathways that were differentially regulated by the young COPD

protocol and plerixafor treatment, we performed proteome analysis on lungs obtained 1 day after

the last poly-IC instillation. Comparison analysis revealed a significant inhibition of pathways of

inflammation, cell invasion, movement, adhesion and synthesis of reactive oxygen species (ROS)

in relation to plerixafor treatment (figure 7H). These changes were driven by down-regulation of

a core set of proteins such as thioredoxin interacting protein (TXNIP), matrix metalloproteinase-8

(MMP-8) and vimentin (figure S13).

Treatment with plerixafor also reduced the development of RV hypertrophy in exposed mice

(figure 7I). The RVSP, as well as heart electrophysiological properties remained unchanged by

CXCR4 inhibition (figure 7J, figure S6M-R).

Flow cytometry 1 day after the last poly(I:C) instillation showed a significant decrease of the

numbers of both fibrocytes and CXCR4^{+} fibrocytes in the lungs of exposed mice by plerixafor

(figure S14A-E). Four days after the last poly(I:C) instillation, the density of peri-bronchial

fibrocytes identified by immunohistochemistry was significantly reduced by plerixafor (figure

S14F-G).

DISCUSSION

Taken together, our data demonstrate that the CXCL12/CXCR4 axis plays a major role in the

pathogenesis of lung and heart remodelling of young COPD. We showed that CXCR4$^+$-cells

were increased in the blood of patients with a mild COPD whereas CXCL12 was increased in the

lung of young COPD patients. Both genetic silencing and pharmacological inhibition of CXCR4

in our experimental young COPD model reduced both the level of CXCR4-expressing cells in the

peripheral circulation and fibrocyte recruitment into the lungs, along with a proteomic signature

consistent with decreased inflammation. As a consequence, CXCR4 inhibition prevented airflow

limitation and both lung and cardiac remodelling.

We designed the present dedicated murine model of young COPD to examine pathophysiological

processes at the onset of the disease. Instead of deciphering the relative contribution of CS and

poly-(I:C), this study aims at developing and using a young COPD model, based on the

following: first, 10-week old mice exposed to CS for 10 weeks would correspond to young

human adults chronically exposed to tobacco smoke for several years [16], and poly-(I:C)

administration mimics exacerbations occurring in young COPD a frequency similar to the COPD

population [17]. Second, as in humans, exposed mice exhibited limited but significant airflow

limitation together with both neutrophilic/lymphocytic airway inflammation and distal airway

remodelling, without obvious signs of alveolar destruction, allowing our design to model the

bronchial component of COPD [23]. By contrast, the vast majority of murine COPD models are

based on long-term CS exposure (up to 6 months) either mimicking emphysema, or reproducing

severe airway obstruction together with emphysema-like changes [24]. Moreover, we used a

combination of CS exposure with poly-(I:C) to mimic respiratory viral infections [25], which are

more frequent in smokers [26] and young COPD [17]. The present model also includes features

of early RV hypertrophy [27], without established pulmonary hypertension, as previously

identified in a long term murine model [28], thus confirming again the relevance of this animal

model of young COPD. Cardiovascular diseases, are indeed frequently seen in current smokers

[29] and patients with COPD [30], including young COPD [31].

A main result of the present study is the increase of CXCL12 expression in the lungs of young

COPD mice and patients, along with an increase in CXCR4[+]-cells in the blood of young COPD

mice as well as in COPD patients. Treating mice with the CXCR4 antagonist plerixafor prevented

the increase of CXCR4[+]-cells, attenuated tissue fibrocytes recruitment and lung function

degradation. Targeting CXCR4 with daily injections of plerixafor during the 5 last weeks of the

protocol as well as genetic deletion of CXCR4 also decreased CS and poly-(I:C) induced-right

heart remodelling. In this connection, neutralizing CXCL12 or antagonizing CXCR4 has been

previously shown to attenuate RV hypertrophy in a rat model of severe pulmonary hypertension

[32]. The beneficial effect of plerixafor on lung function and remodelling in exposed mice,

exhibiting increased CXCR4 solely in the blood, was similar with that observed in CXCR4[-/-]

mice. This suggests that such beneficial effects occurred through CXCR4 reduction in the blood

rather than in the lungs. These results are in agreement with findings obtained in mice treated

with another CXCR4 antagonist or with CXCL12 neutralizing antibodies, in the context of

pulmonary fibrosis, where reduction of fibrocytes accumulation concomitantly improved lung

outcomes [11,33]. The reduction of tissue fibrocytes could thus contribute to lower chronic

inflammation [34,35]. Finally, plerixafor-induced increased mobilization of hematopoietic

progenitor cells from the bone marrow may also participate in the improvement of lung function

and remodelling [10]. Several mechanisms of up-regulation of blood CXCR4[+]-cells may be

proposed, including effects mediated by mild/intermittent hypoxia [7], the Vascular Endothelial

Growth Factor (VEGF) and the basic Fibroblast Growth Factor (bFGF), described in endothelial

cells [36], or IL-33, described in fibrocytes [37,38].

ACKNOWLEDGMENTS

We thank the study participants and the staff of the Thoracic Surgery, Radiology, Pathology, Respiratory, Lung Function Testing departments from the University Hospital of Bordeaux (Bordeaux, France), Fabienne Estella for technical assistance with mice protocol, Isabelle Goasdoue, Isabelle Bernis, Natacha Robert, Virginie Niel, and Marine Servat from the clinical investigation center for technical assistance, and Atika Zouine and Vincent Pitard for technical assistance at the Flow cytometry facility (CNRS UMS 3427, INSERM US 005, Univ. Bordeaux, F-33000 Bordeaux, France), Anne-Aurélie Raymond for help with proteomic analysis at the Oncoprot Platform (TBMCore US005). Christel Poujol, Sébastien Marais and Fabrice Cordelières for help with imaging and image analysis et the Bordeaux Imaging Centre (BIC; Bordeaux, France). Microscopy was performed at BIC, a service unit of the CNRS-INSERM and Bordeaux University, a member of the national BioImaging infrastructure of France supported by the French National Research Agency (ANR-10-INBS-04).

Contributors

- conception and design (ID, PB)

- data acquisition (ID, PH, EM, RA, SC, DEH, EE, PE, MC, MD, GC, CB, OO, JWD, TT, VFM)

- data analysis (ID, PH, EM, RA, SC, DEH, RP, MD, GC, CB, JWD, VFM, PB)

- data interpretation (ID, PH, EM, RA, SC, DEH, EE, RP, PE, MD, GC, CB, TT, RM, MZ, VFM, PB)

- resources (PH, MC, MD, MT, HB, POG, MZ, PB)

- drafting the manuscript (ID, PH, PB)

• revision and final approval of the version to be published, agreement to be accountable

for all aspects of the work in ensuring that questions related to the accuracy or integrity

of any part of the work are appropriately investigated and resolved (All)

Funding

The project was supported by a grant from the "Fondation de l'Université de Bordeaux" (Fonds

pour les maladies chroniques nécessitant une assistance médico-technique FGLMR/AVAD), a

grant from « Aquitaine Science Transfert » (PB) and an unrestricted grant from AstraZeneca, the

"Investments in the Future" program managed by the "Agence Nationale de la Recherche"

(ANR, grant reference ANR-10-IAHU-04). The COBRA cohort was funded by AstraZeneca,

Chiesi, Glaxo-SmithKline, Novartis, Chiesi,Roche and Legs Poix fondation.

Competing interests

ID has 2 patents delivered (i) (EP N°3050574 *i.e.*, Use of plerixafor for treating and/or preventing

acute exacerbations of chronic obstructive pulmonary disease); (ii) (EP N°20173595.8 *i.e.*, New

compositions and methods of treating COVID-19 Disease). ID report a grant from the

"Fondation Bordeaux Université," with funding from "Assistance Ventilatoire à Domicile"

(AVAD) and "Fédération Girondine de Lutte contre les Maladies Respiratoires" (FGLMR).

PH reports non-financial support from AVAD, Chiesi and GlaxoSmithKline, outside the

submitted work and a grant from the "Fondation Bordeaux Université," with funding from

AVAD and FGLMR.

POG reports grants, personal fees and non-financial support from AstraZeneca, personal fees and

non-financial support from Chiesi, personal fees and non-financial support from

GlaxoSmithKline, personal fees and non-financial support from Novartis, personal fees and non-

financial support from Sanofi, outside the submitted work. POG has 2 patents delivered (i) (EP

N°3050574 *i.e*., Use of plerixafor for treating and/or preventing acute exacerbations of chronic

obstructive pulmonary disease); (ii) (EP N°20173595.8 *i.e*., New compositions and methods of

treating COVID-19 Disease).

MZ reports personal fees from AstraZeneca, Boehringer Ingelheim, CSL Behring, Novartis,

Chiesi, GlaxoSmithKline and non-financial support Lilly outside the submitted work and a grant

from the "Fondation Bordeaux Université," with funding from AVAD and FGLMR.

PB is the medical coordinator of the French national cohort (*i.e*., COBRA), which received grants

from AstraZeneca, GlaxoSmithKine, and Chiesi. Moreover, PB reports grants and personal fees

from Novartis, personal fees and non-financial support from Chiesi, grants, personal fees and

non-financial support from Boehringer Ingelheim, grants, personal fees and non-financial

support from AstraZeneca, personal fees and non-financial support from GSK, personal fees and

non-financial support from Sanofi, personal fees from Menarini, personal fees from TEVA,

outside the submitted work; in addition, PB has 4 patents delivered (i) (EP N°3050574 *i.e*., Use

of plerixafor for treating and/or preventing acute exacerbations of chronic obstructive pulmonary

disease); (ii) (EP N°20173595.8 *i.e*., New compositions and methods of treating COVID-19

Disease); (iii) (N° WO2017203064A1 *i.e*., MRI Method for the geometrical characterization of

pulmonary airways); (iv) (N° WO2021018439A1 *i.e*., Method for generating a biomarker

system). All other authors declare they have no competing interests.

REFERENCES

Global Initiative for Chronic Obstructive Lung Disease. Global Initiative for Chronic Obstructive Lung Disease - GOLD. https://goldcopd.org/ (accessed 6 February 2023)

Mannino DM, Buist AS. Global burden of COPD: risk factors, prevalence, and future trends. *Lancet*. 2007;370:765–73.

Martinez FJ, Agusti A, Celli BR, *et al.* Treatment Trials in Young Patients with Chronic Obstructive Pulmonary Disease and Pre-Chronic Obstructive Pulmonary Disease Patients: Time to Move Forward. *Am J Respir Crit Care Med*. 2022;205:275–87.

Henrot P, Prevel R, Berger P, *et al.* Chemokines in COPD: From Implication to Therapeutic Use. *Int J Mol Sci*. 2019;20. doi: 10.3390/ijms20112785

Dupin I, Allard B, Ozier A, *et al.* Blood fibrocytes are recruited during acute exacerbations of chronic obstructive pulmonary disease through a CXCR4-dependent pathway. *J Allergy Clin Immunol*. 2016;137:1036-1042.e7.

Busillo JM, Benovic JL. Regulation of CXCR4 signaling. *Biochimica et Biophysica Acta (BBA) - Biomembranes*. 2007;1768:952–63.

Schioppa T, Uranchimeg B, Saccani A, *et al.* Regulation of the Chemokine Receptor CXCR4 by Hypoxia. *Journal of Experimental Medicine*. 2003;198:1391–402.

Tian X, Xie G, Xiao H, *et al.* CXCR4 knockdown prevents inflammatory cytokine expression in macrophages by suppressing activation of MAPK and NF-κB signaling pathways. *Cell & Bioscience*. 2019;9:55.

Petty JM, Sueblinvong V, Lenox CC, *et al.* Pulmonary stromal-derived factor-1 expression and effect on neutrophil recruitment during acute lung injury. *J Immunol*. 2007;178:8148–57.

Barwinska D, Oueini H, Poirier C, *et al.* AMD3100 ameliorates cigarette smoke-induced emphysema-like manifestations in mice. *Am J Physiol Lung Cell Mol Physiol*. 2018;315:L382–6. doi: 10.1152/ajplung.00185.2018

Phillips RJ, Burdick MD, Hong K, *et al.* Circulating fibrocytes traffic to the lungs in response to CXCL12 and mediate fibrosis. *J Clin Invest*. 2004;114:438–46. doi: 10.1172/JCI20997

Dupin I, Thumerel M, Maurat E, *et al.* Fibrocyte accumulation in the airway walls of COPD patients. *Eur Respir J*. 2019;54. doi: 10.1183/13993003.02173-2018

Dupin I, Henrot P, Maurat E, *et al.* Targeting CXCR4 as a therapeutic strategy to improve outcomes in a mouse model of early chronic obstructive pulmonary disease (COPD). *ERJ Open Research*. 2022;8. doi: 10.1183/23120541.LSC-2022.9

14 Dupin I, Dubreuil M, Freund-Michel V, *et al*. Targeting CXCR4 as a therapeutic strategy to
improve outcomes in a mouse model of early chronic obstructive pulmonary disease (COPD).
ERJ Open Research. 2019;5. doi: 10.1183/23120541.lungscienceconference-2019.PP225

15 Almolki A, Guenegou A, Golda S, *et al*. Heme oxygenase-1 prevents airway mucus
hypersecretion induced by cigarette smoke in rodents and humans. *Am J Pathol*.
2008;173:981–92.

16 Dutta S, Sengupta P. Men and mice: Relating their ages. *Life Sciences*. 2016;152:244–8.

17 Cosío BG, Casanova C, Soler-Cataluña JJ, *et al*. Unravelling young COPD and pre-COPD in
the general population. *ERJ Open Res*. 2023;9:00334–2022.

18 Hogg JC, McDonough JE, Sanchez PG, *et al*. Micro–Computed Tomography Measurements
of Peripheral Lung Pathology in Chronic Obstructive Pulmonary Disease. *Proc Am Thorac
Soc*. 2009;6:546–9.

19 Sangaletti S, Tripodo C, Cappetti B, *et al*. SPARC oppositely regulates inflammation and
fibrosis in bleomycin-induced lung damage. *Am J Pathol*. 2011;179:3000–10.

20 Tachibana K, Hirota S, Iizasa H, *et al*. The chemokine receptor CXCR4 is essential for
vascularization of the gastrointestinal tract. *Nature*. 1998;393:591–4.

21 Tokoyoda K, Egawa T, Sugiyama T, *et al*. Cellular Niches Controlling B Lymphocyte
Behavior within Bone Marrow during Development. *Immunity*. 2004;20:707–18.

22 Jullien N, Goddard I, Selmi-Ruby S, *et al*. Use of ERT2-iCre-ERT2 for conditional
transgenesis. *Genesis*. 2008;46:193–9.

23 Higham A, Quinn AM, Cançado JED, *et al*. The pathology of small airways disease in
COPD: historical aspects and future directions. *Respir Res*. 2019;20:49.

24 Tanner L, Single AB. Animal Models Reflecting Chronic Obstructive Pulmonary Disease
and Related Respiratory Disorders: Translating Pre-Clinical Data into Clinical Relevance.
JIN. 2020;12:203–25.

25 Kang M-J, Lee CG, Lee J-Y, *et al*. Cigarette smoke selectively enhances viral PAMP- and
virus-induced pulmonary innate immune and remodeling responses in mice. *J Clin Invest*.
2008;118:2771–84.

26 Arcavi L, Benowitz NL. Cigarette Smoking and Infection. *Archives of Internal Medicine*.
2004;164:2206–16.

27 Hilde JM, Skjørten I, Grøtta OJ, *et al*. Right ventricular dysfunction and remodeling in
chronic obstructive pulmonary disease without pulmonary hypertension. *J Am Coll Cardiol*.
2013;62:1103–11.

28 Weissmann N, Lobo B, Pichl A, *et al.* Stimulation of soluble guanylate cyclase prevents
cigarette smoke-induced pulmonary hypertension and emphysema. *Am J Respir Crit Care*
Med. 2014;189:1359–73.

29 Banks E, Joshy G, Korda RJ, *et al.* Tobacco smoking and risk of 36 cardiovascular disease
subtypes: fatal and non-fatal outcomes in a large prospective Australian study. *BMC*
Medicine. 2019;17:128.

30 Chen W, Thomas J, Sadatsafavi M, *et al.* Risk of cardiovascular comorbidity in patients with
chronic obstructive pulmonary disease: a systematic review and meta-analysis. *Lancet Respir*
Med. 2015;3:631–9. doi: 10.1016/S2213-2600(15)00241-6

31 Divo MJ, Marin JM, Casanova C, *et al.* Comorbidities and mortality risk in adults younger
than 50 years of age with chronic obstructive pulmonary disease. *Respiratory Research*.
2022;23:267.

32 Bordenave J, Thuillet R, Tu L, *et al.* Neutralization of CXCL12 attenuates established
pulmonary hypertension in rats. *Cardiovasc Res*. Published Online First: 7 June 2019. doi:
10.1093/cvr/cvz153

33 Griffiths K, Habiel DM, Jaffar J, *et al.* Anti-fibrotic Effects of CXCR4-Targeting i-body AD-
114 in Preclinical Models of Pulmonary Fibrosis. *Sci Rep*. 2018;8:3212.

34 Eyraud E, Maurat E, Sac-Epée J-M, *et al.* Short-range interactions between fibrocytes and
CD8+ T cells in COPD bronchial inflammatory response. *eLife*. 2023;12:RP85875.

35 Dupin I, Eyraud E, Maurat É, *et al.* Probabilistic cellular automata modelling of intercellular
interactions in airways: complex pattern formation in patients with chronic obstructive
pulmonary disease. *Journal of Theoretical Biology*. 2023;564:111448.

36 Salcedo R, Wasserman K, Young HA, *et al.* Vascular endothelial growth factor and basic
fibroblast growth factor induce expression of CXCR4 on human endothelial cells: In vivo
neovascularization induced by stromal-derived factor-1alpha. *Am J Pathol*. 1999;154:1125–
35.

37 Wang C-H, Weng C-M, Huang T-T, *et al.* Anti-IgE therapy inhibits chemotaxis, proliferation
and transformation of circulating fibrocytes in patients with severe allergic asthma.
Respirology. Published Online First: 9 June 2021. doi: 10.1111/resp.14096

38 Berger P, Dupin I. Unravelling the effects of omalizumab on fibrocytes. *Respirology*.
2021;26:825–7.

Figure legends

Figure 1: Characterization of CXCR4 and CXCL12 expression in lung in patients with or

without COPD. (A, D) CXCL12 (A) and CXCR4 (D) mRNA expression in human lungs. Data

are derived from a publicly available GSE-set (GSE76925). Control subjects n=40; patients with

COPD n=111. (B) Representative staining of CXCL12 (magenta) and nucleus (white) in peri-

bronchial areas of human lungs. Lower panels: higher magnification of the images surrounded by

white squares in the upper panels. Images obtained in control *vs* young COPD lungs were

acquired with the same exposure time and gain. The white arrowheads and arrows indicate

respectively CXCL12-expressing bronchial cells and immune-type cells. (C, E) Quantification of

the percentage of positive CXCL12 (C) or CXCR4 (E) surface (normalized by the total area)

(n=5 control subjects, n=6 patients with young COPD). Medians are represented as horizontal

lines. (A, C, D, E) *: P <0.05, ***: P<0.001, t-test or Mann-Whitney test.

Figure 2: Evaluation of lung function, inflammation, bronchial and heart remodelling in experimental COPD. (A) Mice are exposed either to room air (RA) and challenged with PBS, or exposed to cigarette smoke (CS) and challenged with poly(I:C), during 10 weeks. They are sacrificed at day 68 or 71 (i.e. 1 day or 4 days after the last poly(I:C) instillation). (B) Average expiratory flow-volume curves of RA+PBS-exposed mice (black curve) and CS+poly(I:C)-exposed mice (grey curve). Lower and upper error bars represent standard deviations for CS+poly(I:C) and RA+PBS-exposed mice, respectively. (C) FEV0.05/FVC (FEV0.05: Forced

Expiratory Volume during 0.05 s, FVC: Forced Vital Capacity) of RA+PBS-exposed mice (black

circles) and CS+poly(I:C)-exposed mice (grey squares). Data represent individual mice. (D) Total

cell count in bronchoalveolar lavage (BAL). Kruskal-Wallis test followed by Dunn's post-tests.

(E-F), BAL differential cell recovery 1 day (d+1) and 4 days (d+4) after the last poly(I:C)

instillation. (G) Top Canonical Ingenuity Pathways significantly altered in CS+poly(I:C) (n=5) *vs*

RA+PBS (n=5)-exposed lungs (n=5) obtained 1 day after the last poly-IC instillation, ranked by

Z-score (negative and positive for the pathways represented respectively on the top and bottom

part of the graph), obtained by Gene Set Enrichment Analysis. (H) Heatmaps of differentially

regulated proteins in CS+poly(I:C) vs RA+PBS-exposed lungs, from the pathways

"hypercytokinemia/hyperchemokinemia in the pathogenesis of influenza" (left), "wound healing

signalling pathway" (middle) and "xenobiotic metabolism PXR signalling pathway" (right). The

colour scale indicates the $\log_2$ fold changes of abundance for each protein. (I) Representative

Masson's trichrome stainings to assess peri-bronchial fibrosis. (J) Standardized fibrosis, defined

as (peri-bronchial fibrosis ("PF", %) - mean PFcontrol)/standard deviation PFcontrol). The peri-

bronchial fibrosis ("PF", percentage) is defined by the ratio between the area of segmented pixels

in the area of analysis divided by the total peri-bronchial area. (K) Representative short-axis cine

magnetic resonance images of heart at end-systolic stage (left panels) and end-diastolic stage

(middle panels) in control mice (top panels) and CS and poly(I:C)-exposed mice (bottom panels).

Right panels: higher magnification of RV wall. (L) Wall thickness. (M) Fulton index, defined as

(right ventricle (RV)/ left ventricle plus septum (LV + S). (N) Right ventricular systolic pressure

(RVSP). (C, E, F, I, L, M, N) Unpaired t test or Mann-Whitney test. *: P<0.05, **: P<0.01, ***:

P<0.001.

Figure 3: Characterization of CXCR4 and CXCL12 expression in the blood and lung in experimental COPD. Mice are exposed either to room air (RA) and challenged with PBS, or exposed to cigarette smoke (CS) and challenged with poly(I:C), during 10 weeks. They are sacrificed at day 68 or 71 (i.e. 1 day or 4 days after the last poly(I:C) instillation). (A, D) Upper left panels: dot plots represent representative side scatter (SSC, y-axis)-forward scatter (FSC, x-axis) graphs of circulating (A) and lung (D) cells. DAPI⁻ cells, and DAPI⁻ CXCR4⁺ cells are shown respectively in grey and pink. Upper right panels and bottom panels: histograms represent representative cell count (y-axis) versus Phycoerythrin (PE) fluorescence (x-axis) in DAPI⁻ circulating (A) and lung (D) cells. Percentages of CXCR4⁺ cells in the DAPI⁻ cells are shown in pink. (B, E) Levels of circulating (B) and lung (E) CXCR4⁺ cells 1 day (d+1) and 4 days (d+4) after the last poly(I:C) instillation. (C, F) Plasma (C) and lung (F) CXCL12 concentration at d+1 and d+4. For lung CXCL12 concentration, the value is normalized to total protein concentration. (G) Representative stainings of CXCL12 (magenta) and nucleus (white) in peri-bronchial areas of mouse lungs. Lower panels: higher magnification of the images surrounded by white squares in

557   the upper panels. Images obtained in RA and PBS *vs* CS and poly(I:C)-exposed have been

558   acquired with the same exposure time and gain. The white arrowheads and arrows indicate

559   respectively CXCL12-expressing bronchial cells and immune-type cells. (H) Quantification of

560   the percentage of positive CXCL12 surface (normalized by the total area). (B, C, E, F, H)

561   Medians are represented as horizontal lines. *: P <0.05, **: P<0.01, Mann-Whitney test.

Figure 4: Characterization of lung fibrocyte level and localization in experimental COPD and in patients with or without COPD. (A-G) Mice are exposed either to room air (RA) and challenged with PBS, or exposed to cigarette smoke (CS) and challenged with poly(I:C), during 10 weeks. They are sacrificed 1 day ("d+1") or 4 days ("d+4") after the last poly(I:C) instillation. a) Representative flow cytometry contour plots in each condition. The x-axis is for CD45-

Allophycocyanin (APC) fluorescence and the y-axis, FSP1-Fluorescein-5-isothiocyanate (FITC)

fluorescence. Percentages of $CD45^+$ $FSP1^{high}$ cells are shown in green. (B) Representative cell

count (y-axis) versus CXCR4-Phycoerythrin (PE) fluorescence (x-axis) in $CD45^+$ $FSP1^{high}$-cell

subsets in mouse lungs. (C-E) Levels of lung $CD45^+$ $FSP1^{high}$ cells (C), $CD45^+$ cells (D), $CD45^+$

$FSP1^{high}$ $CXCR4^+$ cells (E), in RA+PBS-exposed mice (black circles) and CS+poly(I:C)-exposed

mice (grey squares), at d+1 and d+4. (F) Representative stainings of CD45 (green) and FSP1

(red) in peri-bronchial areas (delimited in pink) of lung mice at d+4. The white arrows indicate

fibrocytes, defined as $CD45^+$ $FSP1^+$ cells. (G) Quantification of fibrocyte density (normalized by

peri-bronchial area). (H) Representative stainings of CD45 (green) and FSP1 (red) in peri-

bronchial area. The white arrowheads indicate fibrocytes, defined as $CD45^+$ $FSP1^+$ cells. (F)

Representative stainings of CD45 (green), FSP1 (red), nuclei (DAPI, blue) and CXCR4 (white)

in peri-bronchial areas of lung samples of control subjects and patients with COPD. The white

arrows indicate CXCR4-expressing fibrocytes, defined as $CD45^+$ $FSP1^+$ $CXCR4^+$ cells. (F, H)

Left and lower panels: higher magnification of the images surrounded by white squares in the

right upper panels. (C-E, G) Medians are represented as horizontal lines. *: P <0.05, **: P<0.01,

Mann-Whitney test.

Figure 5: Evaluation of lung function, inflammation and bronchial and heart remodelling upon conditional knockout of CXCR4 in experimental COPD. (A) CXCR4$^{lox/lox}$ (grey squares) and CXCR4$^{lox/lox}$/Cre2ERT2 (pink squares) mice are exposed to cigarette smoke (CS) and challenged with poly(I:C). All the mice were administered with 1 mg intraperitoneal tamoxifen for 5 consecutive days at the beginning of the COPD protocol, and then fed by tamoxifen-containing food during the remaining protocol. They are sacrificed at day 68 or 71 (1 day and 4 days after the last poly(I:C) instillation). (B) FEV0.05/FVC in the different conditions. Data represent individual mice. (C) Total cell count in bronchoalveolar lavage (BAL). Kruskal-

Wallis test followed by Dunn's post-tests. (D-E) BAL relative (D) and absolute (E) cell

composition respectively 1 day and 4 days after the last poly(I:C) instillation. (F) Representative

Masson's trichrome stainings to assess peri-bronchial fibrosis. (G) Standardized fibrosis, defined

as (peri-bronchial fibrosis ("PF", %) - mean PFcontrol)/standard deviation PFcontrol). The peri-

bronchial fibrosis ("PF", percentage) is defined by the ratio between the area of segmented pixels

in the area of analysis divided by the total peri-bronchial area. (H) Representative short-axis cine

magnetic resonance images at end-systolic stage (left panels) and end-diastolic stage (middle

panels) in CS and poly(IC)-exposed $CXCR4^{lox/lox}$ mice (top panels) and CS and poly(IC)-exposed

$CXCR4^{lox/lox}$/Cre2ERT2 mice (bottom panels). Right panels: higher magnification of RV wall. i)

Wall thickness. (J) Fulton index, defined as (right ventricle (RV)/ left ventricle plus septum (LV

+ S). (K) Right ventricular systolic pressure (RVSP). (B, D, E, G, J, K) *: $P<0.05$, **: $P<0.01$,

***: $P<0.001$. Unpaired t test or Mann-Whitney test.

Figure 6: Characterization of fibrocyte level and peri-bronchial density in lung upon conditional knockout of CXCR4 in experimental COPD. CXCR4$^{lox/lox}$ (grey squares) and CXCR4$^{lox/lox}$/Cre2ERT2 (pink squares) mice are exposed to cigarette smoke (CS) and challenged with poly(I:C). They are sacrificed at day 68 or 71 (1 day and 4 days after the last poly(I:C) instillation). (A) Representative flow cytometry contour plots in each condition. The x-axis is for CD45-Allophycocyanin (APC) fluorescence and the y-axis, FSP1-Fluorescein-5-isothiocyanate (FITC) fluorescence. Percentages of CD45$^+$ FSP1^{high} cells are shown in green. (B) Representative cell count (y-axis) versus CXCR4-Phycoerythrin (PE) fluorescence (x-axis) in CD45$^+$ FSP1^{high}-cell subsets in mouse lungs. (C-E) Levels of lung CD45$^+$ FSP1^{high} cells (C), CD45$^+$ FSP1^{high} CXCR4$^+$ cells (D), CD45$^+$ cells (E) 1 day after the last poly(I:C) instillation in each condition. (F)

616   Quantification of fibrocyte density (normalized by peri-bronchial area) 4 days after the last

617   poly(I:C) instillation in each condition. (G) Representative stainings of CD45 (red) and FSP1

618   (green) in peri-bronchial area (delimited in pink) in each condition. The white arrows indicate

619   fibrocytes, defined as CD45$^+$ FSP1$^+$ cells. (C-F) Medians are represented as horizontal lines. *: P

620   <0.05, Unpaired t test or Mann-Whitney test.

Figure 7: Evaluation of lung function, inflammation and bronchial and heart remodelling upon pharmacological blockage of CXCR4 in experimental COPD. (A) Mice are exposed to cigarette smoke (CS), challenged with poly(I:C) and treated with plerixafor ("pleri", 1 mg/kg, purple squares) or its vehicle (grey squares). They are sacrificed at day 68 or 71 (1 day and 4 days after the last poly(I:C) instillation). (B) FEV0.05/FVC in the different conditions. Data represent individual mice. (C) Total cell count in bronchoalveolar lavage (BAL). Kruskal-Wallis test followed by Dunn's post-tests. (D-E) BAL relative (D) and absolute (E) cell composition respectively 1 day and 4 days after the last poly(I:C) instillation. (F) Representative Masson's trichrome stainings to assess peri-bronchial fibrosis. (G) Standardized fibrosis, defined as (peri-

bronchial fibrosis ("PF", %) - mean PFcontrol)/standard deviation PFcontrol). The peri-bronchial

fibrosis ("PF", percentage) is defined by the ratio between the area of segmented pixels in the

area of analysis divided by the total peri-bronchial area. (H) Heatmaps of significantly

differentially regulated pathways identified by Ingenuity Pathway analysis (IPA; Qiagen),

obtained by proteomics comparison of CS+poly(I:C) (n=5) *vs* RA+PBS (n=5)-exposed lungs

(n=5) (all treated by vehicle), and plerixafor (n=5) vs vehicle (n=5)-treated lungs (all exposed to

CS+poly(I:C)). The colour scale indicates the Z-score value. (I) Fulton index, defined as (right

ventricle (RV)/ left ventricle plus septum (LV + S). (J) Right ventricular systolic pressure

(RVSP). (B, D, E, G, I, J) *: P<0.05, **: P<0.01, ***: P<0.001. Unpaired t test or Mann-Whitney

test.