Loss of MC1R signaling implicates TBX3 in pheomelanogenesis and melanoma predisposition

H. Matthew Berns[1,2], Dawn E. Watkins-Chow[1], Sizhu Lu[2], Pakavarin Louphrasitthiphol[2,3], Tongwu Zhang[4], Kevin M. Brown[4], Pedro Moura-Alves[2,5,6], Colin R. Goding[2], William J. Pavan[1,*]

[1]Genetic Disease Research Branch, National Human Genome Research Institute, National Institutes of Health, Bethesda, MD, 20892, USA.

[2]Ludwig Institute for Cancer Research, Nuffield Department of Clinical Medicine, University of Oxford, Headington, Oxford, OX3 7DQ, UK.

[3]Department of Gastrointestinal and Hepato-Biliary-Pancreatic Surgery, Faculty of Medicine, University of Tsukuba, Tsukuba, Japan.

[4]Division of Cancer Epidemiology and Genetics, National Cancer Institute, Bethesda, MD 20892, USA.

[5]i3S-Instituto de Investigação e Inovação em Saúde, Universidade do Porto, 4200-135 Porto, PT.

[6]IBMC-Instituto de Biologia Molecular e Celular, Universidade do Porto, 4200-135 Porto, PT.

*Corresponding Author: William J. Pavan

Email: bpavan@mail.nih.gov

Author Contributions: Contribution: W.J.P., D.E.W.-C, H.M.B, P.M.A., and C.R.G. designed the research study; H.M.B., D.E.W.-C, S.L., P.L., K.M.B., T.Z. acquired the data; H.M.B., D.E.W.-C, K.M.B., T.Z. analyzed data; H.M.B, D.E.W.-C, C.R.G., and W.J.P. wrote the manuscript; H.M.B., D.E.W.-C., S.L., P.L., C.R.G., P.M.A., and W.J.P. edited the manuscript; and all authors approved the final manuscript.

Financial Disclosure Statement:

This research was supported by the National Human Genome Research Institute (NHGRI)

Intramural Research Program at the National Institutes of Health (NIH) (1ZIAHG000068-15), the

Intramural Research Program of the National Cancer Institute (NCI) at the National Institutes of

Health (NIH) (1ZIACP010201), and the Ludwig Institute for Cancer Research. H.M.B. is

supported by an NHGRI Intramural Research Training Award, the NIH Oxford-Cambridge

Scholars Program, and the NIH Medical Research Scholars Program, a public-private partnership

supported jointly by the NIH and contributions to the Foundation for the NIH from the Doris Duke

Charitable Foundation, Genentech, the American Association for Dental Research, and the

Colgate-Palmolive Company. Additionally, P.M.A. would like to thank H2020-WIDESPREAD-

2018-951921- ImmunoHUB for financial support. The funders had no role in study design, data

collection and analysis, decision to publish, or preparation of the manuscript.

Competing Interest Statement: Authors have no competing interests to declare.

Keywords: melanoma; melanocytes; melanocortin 1 receptor (MC1R); TBX3; red hair color.

Data Availability: Original data created for the study are or will be available in a persistent

repository upon publication. TBX3 ChIP is deposited in Gene Expression Omnibus (GEO) as

GSE223256; Currently compiling pertinent information for submission of bulk and single-cell

RNA-seq datasets to GEO.

Abstract

The human Red Hair Color (RHC) trait is caused by increased pheomelanin (red-yellow) and reduced eumelanin (black-brown) pigment in skin and hair due to diminished melanocortin 1 receptor (MC1R) function. In addition, individuals harboring the RHC trait are predisposed to melanoma development. While *MC1R* variants have been established as causative of RHC and are a well-defined risk factor for melanoma, it remains unclear mechanistically why decreased MC1R signaling alters pigmentation and increases melanoma susceptibility. Here, we use single-cell RNA-sequencing (scRNA-seq) of melanocytes isolated from RHC mouse models to reveal a Pheomelanin Gene Signature (PGS) comprising genes implicated in melanogenesis and oncogenic transformation. We show that TBX3, a well-known anti-senescence transcription factor implicated in melanoma progression, is part of the PGS and binds both E-box and T-box elements to regulate genes associated with melanogenesis and senescence bypass. Our results provide key insights into mechanisms by which MC1R signaling regulates pigmentation and how individuals with the RHC phenotype are predisposed to melanoma.

Introduction

Cancer predisposition is usually defined by inherited genetic mutations which increase an individual's risk of developing the disease. Identification of the genes and molecular pathways underlying predisposition is essential for both cancer treatment and prevention. Melanoma, the most lethal form of cutaneous cancer, arises from melanin-producing melanocytes and represents an excellent model to study cancer risk factors. Melanoma is an increasingly common skin cancer caused by high levels of sun exposure and subsequent damage by ultraviolet (UV) light. All stages of the disease are amenable to study and the genetic basis of disease initiation is well-defined. Specifically, cutaneous melanoma usually arises from acquisition of activated pro-oncogenes (e.g., BRAF, NRAS) combined with senescence bypass through inactivation of cell

cycle regulators (e.g., CDKN2A/INK4A)[1–4]. However, in addition to genes falling into these two

categories, additional genetic loci have been implicated in melanoma predisposition.

Understanding the molecular basis for melanoma predisposition is a key issue which has the

potential to inform melanoma prevention strategies.

Red Hair Color (RHC) is a well-described melanoma predisposition phenotype.

Individuals with the RHC phenotype, possessing red hair, pale skin, freckles, and an inability to

tan, show an increased risk of developing melanoma compared to those with a non-RHC

disposition[5–7]. The RHC trait is mostly explained by coding variants of reduced function in the

melanocortin 1 receptor (MC1R) gene which encodes a G-protein-coupled receptor (GPCR)

located on the cell surface of melanocytes. MC1R signaling regulates pigment/melanin

production[7–9]; therefore, diminished MC1R function alters the ratio of cellular eumelanin and

pheomelanin that largely determines skin, hair, and eye color. Decreased MC1R signaling results

in the loss of photoprotective eumelanin synthesis and in turn allows for almost exclusive

production of pheomelanin. High levels of pheomelanin in RHC individuals exacerbate DNA

damage through the amplification of free radicals which is worsened when cells are exposed to

UV[10,11], and pheomelanin itself can contribute to melanomagenesis in the absence of UV

radiation[11]. These observations may partly explain why individuals with genetic RHC variants in

MC1R exhibit an increased somatic mutational burden in melanocytes[12].

Although the influence of *MC1R* RHC variants on melanin composition may play a role in

melanoma predisposition, *MC1R* mutation status may also have an impact on melanoma initiation

unrelated to its role in pigmentation. For example, individuals with *MC1R* variants fail to activate

PTEN upon UV exposure, resulting in increased PI3K/AKT signaling and oncogenic

transformation[13]. However, whether MC1R signaling plays additional roles in the prevention of

melanoma initiation remains unknown. In part, this may be because, by definition, a molecular

mechanism underlying melanoma predisposition must already be operating prior to the

acquisition of the ultimate trigger for melanoma initiation, such as acquiring activating BRAF or

NRAS mutations.

MC1R signaling shows high evolutionary conservation among mammals, thus murine

species provide an excellent model to explore this pathway. This includes mice with active MC1R

signaling (nonagouti; *a/a*), and suppressed MC1R signaling (recessive yellow, *Mc1r^{e/e}*, and lethal

yellow, *A^{y}/a*). Nonagouti mice have a dark coat color that results from a loss-of-function (LOF)

allele of the nonagouti gene (*a*), which encodes the mouse homolog of agouti signaling protein

(ASIP). In the absence of this important MC1R antagonist, nonagouti melanocytes predominantly

synthesize eumelanin. In contrast, recessive yellow (*Mc1r^{e/e}*) and lethal yellow (*A^{y}/a*) mice provide

two models for RHC, each with melanocytes that synthesize primarily pheomelanin due to distinct

molecular mechanisms which cause reduced MC1R signaling. Recessive yellow mice carry a

LOF allele in the MC1R receptor while the similar yellow coat color of lethal yellow mice is caused

by constitutive overexpression of the MC1R antagonist, agouti signaling protein[14–16].

110   Given the role of the RHC trait in melanoma risk, a better understanding of the genetic

regulatory network underlying aberrant MC1R signaling is needed. Previously, April et al[17]

identified several novel genes linked to pheomelanogenesis in mouse models using bulk RNA-

sequencing. Since this study used whole skin layers, of which melanocytes comprise only a small

fraction of the total cell population, gene expression from other cell types such as keratinocytes,

basal cells, and dendritic cells potentially masked the detection of melanocyte-intrinsic changes.

Haltaufderhyde and Oancea[18] performed bulk RNA-sequencing on human epidermal

melanocytes to assess global gene expression changes between light and dark colored cells.

However, the genotypes were unknown, and melanocytes were not exclusively producing one

form of melanin over the other. While these previous studies established an important foundation

for our understanding of pheomelanogenesis, additional, high-resolution genomic studies are

needed to fully evaluate the intricate molecular networks regulating eumelanogenesis,

pheomelanogenesis, and melanoma development.

123   To better understand the gene regulatory network associated with the RHC trait, we used

single-cell RNA-sequencing (scRNA-seq) to compare melanocytes with both active and

suppressed MC1R signaling. We defined a set of gene candidates, the Pheomelanin Gene

Signature (PGS), implicated in regulating pheomelanogenesis and melanomagenesis. We further

explored the chromatin occupancy of TBX3, a PGS- and melanoma-associated T-box family

member. Our results provide key insights into how RHC trait variants may increase melanoma

risk via altered regulation of PGS candidate genes downstream of MC1R.

Results

Single-cell RNA-sequencing of primary melanocytes identifies a

Pheomelanin Gene Signature (PGS)

To better understand global gene expression changes downstream of MC1R signaling, we used

single-cell RNA-sequencing (scRNA-seq) to define the transcriptomic profiles of primary mouse

melanocytes which predominantly synthesize either pheomelanin (lethal yellow, A^y; or recessive

yellow, $Mc1r^e$) or eumelanin (nonagouti, a). The dorsal skin was removed from post-natal day 3

(P3) animals along anatomically defined margins consisting of four incisions (Fig 1A): two

transverse cuts, with one between each axilla, and the other between the anterior pelvises, and

bilateral incisions at the midline from the axilla to the anterior pelvis. This tissue was then

enzymatically digested into a single-cell solution. Melanocytes comprise ~5% of cells within the

skin; therefore, after dissection and digestion of tissue, a fluorescent-activated cell sorting (FACS)

approach was used to enrich the pool of melanocytes for sequencing[19]. Positively selected cells

were defined as c-KIT$^+$ and CD45$^-$ (Fig 1A,B). To eliminate selection bias among potential

subpopulations of melanocytes that may vary in c-KIT levels, a lower intensity threshold for c-KIT

positivity was used, yielding a population enriched for melanocytes but expected to also contain

other cell types. Isolated cells were processed for library construction and RNA-sequencing via

the 10x Genomics Chromium pipeline. After the implementation of quality control measures (see

Methods), sequencing data was obtained from a total of 11,058 cells from all samples. The

corresponding data underwent dimensional reduction via non-biased principal component (PCA)

and graph-based clustering analyses for graphical display by Uniform Manifold Approximation

and Projection (UMAP) (Fig 1C,D). We classified and labeled captured cell types by overlaying

published murine skin cell gene signatures[20] (Fig 1C,D; Table S1; Fig S1-S2). Cells classified as

melanocytes included 4 of the 16 graph-based clusters (GBC) identified in cells isolated from a

nonagouti and lethal yellow pair of samples, and 1 of 12 GBCs identified in cells isolated from a

nonagouti and recessive yellow pair of samples. Subsequently, for each of the 2 pairwise

experiments, all non-melanocyte cells were filtered out and only gene expression from the cells

classified as melanocytes were used for further analysis.

Because the mitotic stage has a significant and influential effect on gene expression, all

melanocytes outside the G_0/G_1 phase was removed using the same gene expression overlay

approach with previously published cell cycle marker genes[21,22] (Fig S3A, S3B). This defined the

non-cycling melanocytes that were used for differential gene expression analysis comparison.

Subsequent graph-based clustering analysis on the melanocyte population showed that both

RHC mouse models displayed a bifurcation between the pheomelanin melanocytes and the

eumelanin melanocytes captured in parallel, as displayed by UMAP (Fig 1C,D). Consistent with

this, clustering of melanocytes by genotype was unique to melanocytes and not observed in the

other cell types, suggesting that the effects of MC1R signaling loss are indeed restricted to

melanocytes[23].

Using this melanocyte population, differentially expressed genes (DEGs) (ANOVA; FDR

< 0.05 and fold-change (FC) ±2) were identified in pair-wise experiments to compare lethal yellow

to nonagouti melanocytes (Fig 1C) as well as recessive yellow to nonagouti melanocytes (Fig

1D). The differential lists from each experiment were intersected to identify shared genes which

have consistent FC direction when comparing the nonagouti (*a/a*) melanocytes to the lethal

yellow (*A^y/a*) or recessive yellow (*Mc1r$^{e/e}$*) melanocytes. These comparisons resulted in a set of

216 genes which we have identified as the Pheomelanin Gene Signature (PGS) (Table S2). We

next asked to what extent these genes are associated with pigmentation by querying the PGS

against a compiled list of genes annotated as causative for pigmentation phenotypes[24] to assess

the degree of overlap. Interestingly, only 5% of the PGS genes appeared in the previously

annotated pigmentation gene list (Table 1).

We next used PANTHER Gene Ontology (GO) and Ingenuity® Pathway Analysis (IPA) to

identify significantly over-represented processes and pathways in the PGS. GO analysis revealed

a prevalence of significant terms (FDR < 0.05) related to 'morphogenesis', 'melanin biosynthesis',

and 'neurogenesis' / 'neurotransmitter regulation' (Fig 2A). Terms related to 'melanin processes'

had equally high fold-enrichment; for example, 4 out of the 15 genes associated with 'melanin

biosynthetic process' are present in the PGS (*Tyrp1*, *Mc1r*, *Oca2*, and *Pmel*). IPA highlighted

canonical pathways related to 'Neuroinflammation, 'Synaptogenesis' and 'Neuron receptor

signaling' (Fig 2B). These terms are primarily driven by a host of receptors and ligands within the

gamma-aminobutyric acid (GABA), N-methyl-D-aspartate (NMDA)/Glutamate, Ephrin, and

Semaphorin families.

Predicted Transcriptional Regulators of the Pheomelanin Gene

Signature include TBX3

To better identify the gene regulatory network regulating pheomelanin synthesis, we examined

differentially expressed transcription factors, as well as predicted molecular upstream regulators

of gene regulatory networks. The PGS contains 22 annotated transcription and co-transcription

factors[25] (Fig 3A). Many of these have established relationships to melanocyte and pigment cell

biology, including neural crest development (*Hmga2*[26], *Id4*[27], *Mef2c*[28], *Tbx3*[29] and *Tfap2b*[30]),

melanocyte differentiation (*Mef2c*[31]), melanogenesis (*Id4*[32], *Irf4*[33], and *Tbx3*[34,35]), oxidative stress

protection (*Nr4a*[36]), and melanoma (*Hmga2*[37], *Irf4*[38], *Nfactc2*[39], *Nr4a*[40], *Tbx3*[41–45] and *Zeb1*[46–48]).

Of note, the *microphthalmia-associated transcription factor* (*Mitf*), a master melanocyte regulatory

transcription factor[49], was not identified as transcriptionally altered. Collectively, these data

suggest that pheomelanin synthesis is not simply a result of eumelanin synthesis inactivation, but

rather could involve multiple complex pathways downstream of MC1R signaling loss.

IPA was used to predict upstream regulators which may be driving expression of the PGS

(Fig 3B). The top 10 predicted regulators include three molecules -- SMAD4, GDF2, and SMAD1

-- that are all related to TGFβ/BMP signaling. Importantly, TGFβ signaling can regulate melanin

synthesis, with activation of the pathway resulting in hypopigmentation[50]. We reasoned that those

transcriptional targets downstream of TGFβ/BMP signaling merited further analysis. Among these

was *Tbx3*, a T-box family member activated by the TGFβ signaling pathway[51]. *Tbx3* showed

significantly increased expression in both the lethal yellow and recessive yellow samples (Fig 3A).

TBX3 has previously been implicated in melanoma invasion[44], but also has a key role as an anti-

senescence transcription factor[1–4,45,52]. Up-regulation of TBX3 and consequent suppression of

senescence in melanocytes lacking MC1R prior to acquisition of activating mutations in BRAF or

NRAS would therefore provide a potential mechanism by which redheads would exhibit a

predisposition to melanoma. In addition, TBX3 has previously been implicated in pigmentation, of

Dun horses for example[53], however, how TBX3 mediates these effects in unclear. Therefore,

TBX3 was selected for further exploration with the goal of confirming TBX3 as a candidate

transcription factor for direct regulation of key anti-senescence and melanogenesis target genes

downstream from MC1R.

221   TBX3 binds at T-box and E-box motifs

To assess the functional targets of TBX3 in melanocytes, Chromatin immunoprecipitation

sequencing (ChIP-seq) was used to identify global targets of TBX3 binding. For this we used

doxycycline-inducible TBX3 tagged with an HA epitope previously used to produce high-quality

ChIP-seq datasets[54–56] (Fig S4A). In our studies, the level of induction of TBX3 by doxycycline

was adjusted to ensure physiological levels of TBX3-HA expression in the human melanoma cell

line, 501mel (Fig S4B). We performed two replicate experiments and utilized MACS2 for peak

calling (q-value cutoff ≤ 0.01). To identify common peaks between the two replicates, a filter was

applied requiring an 80% overlap of coordinates between the 2 replicates, resulting in the

identification of 20,449 common peaks (Fig S4C).

To determine the biological significance of these peaks, we used Genomic Regions

Enrichment Annotation Tool (GREAT)[57] to identify 10,000 peaks associated with 7,109 candidate

TBX3 gene targets (Methods; Table S3). We next assessed the binding distribution of these

10,000 peaks relative to gene transcriptional start sites (TSS). Promoter region binding within

±5kb of the TSS comprised only 12.68% of peaks, and the vast majority were situated ±20kb from

the TSS, suggesting TBX3 binds distal enhancer elements (Fig 4A,B). GO enrichment for the

7,109 candidate TBX3 gene targets included multiple terms related to pigmentation/melanogenic

pathways, as well as multiple terms associated with EMT, of which TBX3 is a well-known

regulator (Fig 4C). Phenotype association analyses showed that pigmentation-related phenotypes

were the most abundant for both human and mouse with a particular emphasis on

hypopigmentation/abnormal melanocyte activity (Fig 4D).

We next used MEME-ChIP to assess motif enrichment within the TBX3 peaks. *De novo*

motif analysis revealed that the top 10 sequences (JASPAR CORE vertebrates non-redundant

v2), ranked by e-value, were not only associated with the expected T-box related elements, but

also with E-box related motifs (Fig 5A). Interestingly, the palindromic canonical E-box sequence

CACGTG is bound by basic-Helix-Loop-Helix (bHLH) and bHLHHzip transcription factors

including MITF and its related family member. This sequence shares a 3bp GTG with the well-

described core of the T-box element (AGGTGTGA) bound by T-box family members [55–57].

Furthermore, TBX3 represses *CDH1* at a CAGGTGT site which shares 5bp with the canonical E-

box[44]. Interestingly, a TBX3 peak does not appear at this site in our ChIP dataset; however, TBX3

and MITF share two peaks within intron 2, and another peak 1.6kb upstream of the promoter

region (Fig S4D). Collectively, T-box and E-box motifs were found in the majority of TBX3-bound

input sequences (8,320/10,000; 83.20%) with the E-box motifs occurring more frequently than the

T-box motifs (69.17% vs 45.92%, respectively) (Fig 5B).

Importantly, as the E-box motif is the primary recognition sequence associated with

MITF[49], we wanted to interrogate the degree of shared genome-wide binding sites between TBX3

and MITF. To assess for common occupancy of TBX3 and MITF we intersected our 20,449 TBX3

peaks with 12,137 called peaks from a previously published human MITF ChIP dataset in 501mel

cells[58]. This yielded 4,758 common peaks which were assessed with MEME-ChIP for motif

enrichment. The top 10 sequences (JASPAR CORE vertebrates non-redundant v2), ranked by e-

value, were all E-box related motifs (Fig 5C) centrally located within the TBX3 bound peaks (Fig

5D). The frequency of TBX3 binding at centrally located E-box motifs also known to be occupied

by MITF suggests that TBX3 and MITF may compete for binding at a set of shared regulatory

elements.

To query the set of TBX3 and MITF dual candidate target genes, we utilized the same

GREAT parameters as our TBX3 analysis to identify 7,095 MITF candidate target genes

associated with peaks from the MITF ChIP dataset[59] (Table S4). We then identified the mouse

homologs for both the TBX3 and MITF target gene sets to allow comparison to the PGS gene list

derived from *in vivo* mouse experiments (Table S5,S6). The intersection of TBX3 (6,499/7,109)

and MITF (6,661/7,094) gene target lists identified a shared set of 4,072 genes, representing

62.6% of the TBX3 genome-wide target list (Fig 6A). Of these, 59 are found on the PGS (59/216;

27.3%), suggesting that a portion of the PGS is directly regulated by both TBX3 and MITF (Fig

6B/Table S7). Importantly, TBX3 and MITF shared binding sites at important pigmentation genes,

with four of these having well-described MITF regulatory elements *IRF4, MC1R, OCA2,* and

TYRP1[59–62] (Fig 6C). Collectively, these data suggest that TBX3 can directly bind at E-box motifs

associated with PGS genes, thus regulating pheomelanogenesis via modulation of MITF-

mediated gene regulation.

TBX3 functions as a transcriptional activator and repressor

Our bioinformatic analyses strongly suggest that TBX3 can bind near PGS genes, including

important genes related to melanogenesis. Traditionally, TBX3 has been shown to function as a

transcriptional repressor; however, it is known to include an activator domain[52,63]. In MC1R-

inhibited melanocytes, individual genes contained in the PGS exhibit either increased or

decreased expression relative to eumelanin-producing nonagouti melanocytes. Therefore, we

wanted to quantitatively analyze the extent to which TBX3 directly influences PGS expression

levels. Bulk RNA-sequencing in human melanoma cells was performed following the transient

reduction of TBX3, to determine (1) if the putative target genes were regulated by TBX3 and (2) if

TBX3 repressed or activated these targets.

We performed transient knock-down of *TBX3* in 501mels for 24h, in triplicate, with two

distinct siRNAs (TBX3_13.6 and TBX3_13.8) and a non-coding control. At 24 hours we collected

lysates, harvested RNA, and then submitted samples for bulk RNA-seq. We then performed

differential analysis and identified significant DEGs (padj < 0.05) that were responsive to TBX3

depletion (Table S8). We next intersected DEGs from the two small interfering RNAs and

removed genes in which FC occurred in contradictory directions, resulting in 1,420 positive FC

genes (+FC) and 1,340 negative FC genes (-FC) (Fig 7A). To identify candidate genes regulated

by TBX3, we intersected this list with our TBX3 ChIP target genes, which identified a total of

1,091 targets (623 +FC, 468 -FC) that show both responsiveness to TBX3 depletion as well as

TBX3 binding near their respective genomic loci (Table S9).

300   TBX3 regulates PGS and cell cycle genes

We next performed Gene Set Enrichment Analysis (GSEA) on the candidate direct TBX3 targets

to identify significant (FDR < 0.05) KEGG terms associated with the +FC and -FC genes. Terms

for the +FC list include those related to TBX3's known role in melanoma, such as 'TGFb signaling

pathway', 'focal adhesion', and 'ECM receptor interaction' (Fig 7B). Importantly, 'melanogenesis'

was one of the most significant terms. Enriched terms for -FC genes include those consistent with

TBX3's pro-proliferative and anti-senescent functions, such as 'cell cycle', MAPK signaling

pathway', and 'p53 signaling pathway' (Fig 7B). Specifically, this includes cell cycle progression

genes *CDK1* and *CDK2* (Table S9). Lastly, we wanted to assess which PGS targets were

specifically bound and regulated in our *in vitro* system. We first converted the 1,091 +/- FC genes

bound and regulated by TBX3 to their known mouse homologs, yielding 1,041 genes. These were

intersected with the 98 genes comprising TBX3 targets of the PGS, of which 23 were present (Fig

7C). Collectively, these data suggest that TBX3 can regulate melanogenesis as both a

transcriptional activator and repressor of PGS genes.

Discussion

Coding variants in *MC1R* are associated with the RHC trait. However, the gene regulatory

networks (GRNs) downstream of MC1R remain poorly understood. Through a high-resolution,

single-cell RNA-sequencing approach we have identified a Pheomelanin Gene Signature in

melanocytes comprising candidate genes which may regulate both pheomelanogenesis and

melanomagenesis. One such candidate is *Tbx3* which we show can bind and regulate PGS

genes. Interestingly, we have shown that the transcriptional regulatory activities of TBX3 include

activation and repression of candidate genes. Furthermore, this function is not restricted to DNA

binding via the T-box motif, rather it includes binding the canonical CACGTG E-box, which is the

main recognition sequence of the melanocyte master regulatory transcription factor MITF.

The vast majority of genes comprising the PGS have no previously established ties to

pigmentation phenotypes (205/216; 94.9%). A common approach to begin investigation of a novel

gene list involves identifying over-enriched gene families and pathways. One such PGS-enriched

family encompasses neurogenesis and neurotransmitter regulation and is primarily driven by

receptors and ligands within the gamma-aminobutyric acid (GABA), N-methyl-D-aspartate

(NMDA)/Glutamate, Ephrin, and Semaphorin families. Mature melanocytes show morphological

similarities with neurons in that they are highly dendritic, which enables contact with multiple

keratinocytes for melanin transfer. Furthermore, normal human melanocytes have GO enrichment

for 'neuron ensheathment and synapse'[22]. Interestingly, melanocytes that predominantly produce

pheomelanin in the presence of ASIP lack dendricity[64]. Likewise, inhibition of glutamate signaling

through NDMA receptors has been shown to inhibit melanocyte filipodia attachment to

keratinocytes as well as reduce melanosome transfer efficacy[65]. A recent study has shown that

GABA signaling between keratinocytes and melanoma cells occurs through neuronal like

synapses, and further shows that this communication promotes melanoma initation[66].

Interestingly, in this context, keratinocytes showed increased GABA receptor expression while

melanoma cells did not. Collectively, increased expression of these gene families may reflect

both intrinsic and extrinsic signaling to promote melanocyte development, dendrite formation, and

melanosome transfer abilities.

Previous work has shown that the benzodiazepine Diazepam, which works to increase

the action of GABA, increases both melanogenesis and melanocyte dendricity[67]. Likewise, NMDA

receptors appear to promote melanin synthesis and tyrosinase activity in the cuttlefish, *Sepia*

officinalis[67,68]. In addition, glutamate receptor genes, especially *Grm1*, are expressed in human

melanomas[69], and ectopic expression of *mGluR1* in melanocytes leads to oncogenic

transformation and metastatic spread with 100% penetrance[69]. Ephrins play an essential role in

regulating biological processes such as cell-cell attachment, cell shape, and motility[70]. Along with

Semaphorins, they are expressed in neural crest cells (which give rise to melanocytes) and are of

vital importance in regulating neural crest migration[71]. Genes within these families also show

differential expression in melanocytes treated with ASIP, suggesting that this treatment results in

dedifferentiation towards a neural crest-like state[72]. Our data further validate this idea and

suggest that MC1R acts downstream of ASIP to effect these changes, as loss of MC1R in

primary melanocytes drives similar gene expression changes.

Transcription factors play an essential role in the regulation of gene expression. The most

well-characterized of these in melanocytes is *Mitf.* This gene encodes a master regulatory

transcription factor implicit in melanocyte specificity, proliferation, survival, migration,

differentiation, and melanogenesis, and whose expression has been reported to lie downstream

from MC1R signaling[73,74]. *Mitf* does not exhibit differential expression in our study, nor do many of

its known post-transcriptional/translational modifiers, such as MAPKs, *Gsk3b*, *p300*, and *Stat3,*

although changes in the activity of these signaling molecules may not be apparent in the gene

expression profile. Similarly, other well-characterized transcription factors in melanocyte biology,

including *Sox10*, *Pax3*, *Lef1*, and *Creb,* which can all regulate *Mitf,* also exhibit no differential

expression. This supports that the primary transcriptional changes driving activation of

pheomelanin, and repression of eumelanin, synthesis downstream of MC1R signaling are not

mediated through reduced *Mitf* transcription, but instead through transcriptional regulation of

other factors that modulate the DNA binding or post-transcriptional regulation of MITF.

One of the primary candidate genes we performed further analysis on is the T-box

transcription factor family member, *Tbx3*. This candidate stood out for the following reasons: (1) it

is activated by TGFβ/BMP signaling[51,63,75], (2) it has high fold-enrichment in melanocytes from

both pheomelanin-producing mouse genotypes (Fig 3A), (3) it is found across multiple terms from

our GO analysis, (4) it has known functional roles in developmental states of melanocytes[29,34,35]

(5) it regulates developmental processes across multiple tissue/cell types, including heart[76–78],

lungs[79], limbs[80,81], inner ear[82], and neural crest[29,76], and (6) is highly expressed in a multitude of

cancers, especially melanoma, where its primary roles appear to be in metastasis by activation of

EMT, at least in part through down-regulation of E-cadherin, as well as suppression of

senescence[41,43–45,83,84].

The differential expression of *Tbx3* in eumelanin- and pheomelanin-expressing

melanocytes, along with genome-wide binding of TBX3 to T- and E-box elements in melanoma

cells, suggested that TBX3 may directly regulate MITF targets downstream of MC1R signaling.

Through ChIP and bulk RNA-seq we've shown that TBX3 can bind and regulate genes within the

PGS. Although TBX3 is often viewed as a transcriptional repressor, it has been shown to activate

gene expression of putative target genes[52]. Likewise, our data strongly suggests that TBX3 can

both activate or repress target genes. These targets include genes with known associations to

pigmentation phenotypes. An intriguing example of TBX3-mediated activation is *Slc7a11*. This

gene encodes the cystine/glutamate exchanger (xCT) which imports cystine into the cell where it

is reduced into two molecules of cysteine, a necessary component for pheomelanin synthesis[85,86].

The increased pool of cysteine provides an environment that is favorable for pheomelanin

synthesis. Increased expression of *Slc7a11* by TBX3 suggests an active process downstream of

MC1R to favor pheomelanin synthesis, distinct from the repression of eumelanin synthesis. The

importance of cysteine goes beyond pheomelanin synthesis, as it is an essential factor of the

glutathione system and oxidative stress regulation as well as ferroptosis inhibiton[86,87]. Following

TBX3 knockdown we observed a significant reduction of *SLC7A11* expression, with associated

enrichment of pathways associated with 'glutathione metabolism' and 'apoptosis'. Given the

cytotoxicity of pheomelanin, regulation of cysteine levels within melanocytes is critical and

presents a paradox where cysteine is essential for regulating oxidative stress; however, too much

can exacerbate cell stress through pheomelanin accumulation.

TBX3 is implicated in senescence by-pass partly through its ability to repress cell cycle

checkpoint genes such as CDKN2B[52]. Our *TBX3* knockdown data support this, with *CDKN2B*

showing significantly increased expression. Furthermore, our data also support a role for TBX3 in

senescence by-pass through activation of cell cycle progression genes, where the loss of *TBX3*

yields decreased expression of *CDK1* and *CDK2* (Table S9). Willmer et al[88] have shown that

TBX3 is important in S-phase transition in part due to post-translational phosphorylation via

CDK2[88]. More recent work from this group suggests a more complex relationship, whereby TBX3

itself promotes *CDK2* expression[89]. Our data further support this model and suggest a further way

in which TBX3 can contribute to oncogenic transformation by facilitating senescence bypass.

These observations may pave the way in explaining why redheads with diminished MC1R

function may be predisposed to melanoma.

The T-box family members comprise five sub-groups, with TBX3 as a member of the

TBX2 sub-group. TBX2 and TBX3 share 95% homology within the DNA binding domain[63]. TBX2,

which canonically binds the T-box, can also bind the E-box motif, and functions as both an

activator and repressor[56]. Our data support the idea that TBX3 also binds to, and regulates gene

expression, via the E-box motif. However, unlike TBX2, the type of motif bound does not seem to

influence its action (Fig S5A). E-box site recognition of TBX3 can possibly be explained with

protein structural studies. As mentioned previously, the canonical E-box and T-box elements

share a 3bp (GTG) region. This corresponds to base pairs 3-5 of the T-box (AGGTGTGA).

Consequently, this 3bp core significantly affects TBX3's ability to bind DNA, whereas the flanking

base pairs are deemed less crucial[90]. As a monomer, TBX3 can also bind to either the major or

minor groove[90]. Therefore, while bound in the minor grove, it is possible that TBX3 may interact

with other transcription factors bound in the major grove, including MITF. The degree to which

TBX3 modulates gene expression is most likely dependent on several factors, including, cell type,

differentiation state, co-factor availability, chromatin accessibility, and biological state (e.g.,

methylated, acetylated). Variation across these factors could explain why more putative genes

targets regulated by TBX3 in human melanoma cells *in vitro* are not represented in the *in vivo*

healthy murine melanocyte PGS.

In this study, we aimed to profile the transcriptome in melanocytes with aberrant MC1R

signaling. To this end we have defined a set of genes (PGS) that merit further investigation

towards better understanding the complex GRNs downstream of MC1R signaling. This includes

the melanoma-associated transcription factor, *Tbx3*. Furthermore, we believe the Pheomelanin

Gene Signature provides important candidate genes that merit further follow-up regarding the

regulation of melanocyte development, melanogenesis, and melanomagenesis.

434 Materials and Methods

435 Single-Cell RNA-Sequencing and Differential Expression

436 Analysis

FACS-isolated melanocytes were captured in a solution of 1xPBS + 0.04% FBS. Cell viability and

concentration were analyzed using a Luna-FX7 cell counter (Logos Biosystems). Gel Bead-in-

Emulsions were prepared by loading 5,000 cells/sample onto a Chromium Chip G (10x Genomics

#1000127) and run using the Chromium controller (10x Genomics #1000204). cDNA libraries

were generated with Chromium Single Cell 3' GEM, Library and Gel Bead Kit V3.1 (10x

Genomics #1000128). Samples were indexed and pairs for each experiment were pooled for

sequencing using the NextSeq 500/550 Hi Output KT v2.5 (Illumina #20024907) on an Illumina

NextSeq550. Barcoded reads were aligned to the mm39 genome using STAR aligner.

Downstream processing of the data was performed with Partek Flow™ Genomic Analysis

Software.

Single cell QA/QC included steps to filter out cells which did not meet the following parameters

defined in the two pairwise experiments (lethal yellow / recessive yellow) as: barcode count 600-

32,000/3,000-45,000, number of genes detected 500-6000/500-7000, and % ribosomal counts

12-43%/12-39%. Of the 31,017 annotated murine genes, 16,793 (54.14%)/20,060 (64.7%)

passed an exclusion cutoff where genes were excluded if they had zero counts in at least 99.9%

of the cells. The remaining read counts were normalized using a $\log_2$-based method on Counts

Per Million (CPM). Normalized counts underwent dimensional reduction via a Principal

Components Analysis (PCA) on the first 100 principal components and unsupervised, non-

biased, Graph-based clustering through a Louvain algorithm on the first 20 principal components.

Graph-based clusters underwent further dimensional reduction for visualization on a Uniform

Manifold Approximation and Projection diagram. Postnatal mouse skin cell types including dermal

fibroblasts, dermal papilla, transit amplifying cells, matrix cells, and epidermal cells were identified

by utilizing published signature gene lists[20]. The most highly expressed genes (top 10%) from the

signature gene lists were overlayed on the UMAP and the graph-based clusters which coincided

with expression overlay were assigned to each cell type. All non-melanocyte cell types were

subsequently filtered out, leaving only clusters classified as melanocytes for downstream

analysis. The melanocyte only population underwent a further iteration of PCA, graph-based

clustering, and UMAP display. To eliminate the variable of cellular heterogeneity as influenced by

the mitotic state, we selected for cells within the G_0 phase using a gene expression overlay

established by Hsiao et al[21]. The melanocytes within the G_0 phase underwent a further iteration of

PCA, graph-based clustering, and UMAP analysis. Differential gene expression lists of

melanocytes were generated using the Partek ANOVA differential expression analysis. Genes

considered statistically significant had a False-discovery rate of <0.05 and a fold-change of ±2.

Gene ontology and pathway analysis

Genes comprising the Pheomelanin Gene Signature were uploaded to the Gene Ontology

Resource for GO Enrichment Analysis via PANTHER for Overrepresentation Tests (GO Ontology

database DOI:10.5281/zenodo.6399963; Released 2022-03-22). The PGS was run against a

reference list comprising all *Mus musculus* genes in the database. Fisher's Exact test was

performed with a False Discovery Rate correction for GO Biological Process Complete, GO Cell

Component Complete, and GO Molecular Function Complete with significant results being those

with an FDR < 0.05. To identify predicted upstream regulators, we utilized QIAGEN's Ingenuity®

Pathway Analysis (73620684 (Release Date: 2022-03-12). Testing was done via Expression Core

Analysis on the observed experimental fold change of the PGS for both pairwise experiments.

Gene Set Enrichment Analysis

483　Gene lists comprising the significant DE +FC and -FC genes regulated by TBX3 were uploaded

484　to the Molecular Signatures Database (MSigDB) for gene set enrichment analysis utilizing the

485　Kyoto Encyclopedia of Genes and Genomes (KEGG). For the +FC list, the top 500 genes by fold-

486　change for TBX3_13.8 were used as the maximum input value is 500 genes. Each list was tested

487　independently with significant terms being those with an FDR q-value < 0.05.

TBX3 ChIP Analysis

Prior to mapping to the human reference genome (hg38) with Bowtie2 (v.2.3.5)[91], quality of the

raw sequencing data was evaluated using FastQC (v.0.11.7), with adapter contamination

removed using CutAdapt (v.2.8). Peak calling was performed using MACS2 (v.2.1.2)[92] with a 0.01

q-value threshold. Bowtie2-generated SAM files were compressed to BAM files, indexed using

SAMtools (v.1.9), and sequentially converted to bigWig files using USCStools (v.373). The BED

files of both replicates were intersected using the UCSC Table Browser to pull out common peaks

from replicate 2 (40,974) that had ≥80% overlap with replicate 1 (32,479), which yielded a group

of 20,449 common peaks. For simplicity of downstream analysis, the 20,449 peak coordinates

used were derived from the BED position parameters of replicate 1.

The Genomic Regions Enrichment of Annotations Tool (GREAT v4.0.4)[57] was used to analyze

the functional significance of cis-regulatory regions of the TBX3 bound ChIP peaks. Each

annotated gene was assigned a basal regulatory domain of 5kb upstream, and 1kb downstream,

of the transcriptional start site. The regulatory domain of that gene was then extended in both

directions to the nearest gene's basal domain but with no more than a 100kb extension in one

direction[57]. Discovery of enriched binding motifs was done using the MEME-ChIP[93] DNA binding

motif analysis tool. 250bps were added up and downstream of the chromosomal coordinates

corresponding to the absolute summit of all overlapping peaks. A FASTA file was generated

using the usegalaxy.org bedtools application 'GetFastaBed'[94]. This was run against the default

settings in MEME-ChIP, except for maximum width for 'What width motifs should MEME and

STREME find?' which was increased to 20 bases. Peak distributions were obtained from GREAT

and ChIPseeker[95], via usegalaxy.org. The histogram was produced by GREAT, while the venn

diagram was created with ChIPseeker.

513   Bulk RNAseq of *in vitro* 501mel Human Melanoma Cells

501mel cells (140,000/well) were plated in a 12-well plate (1.2mL/well). dsiRNAs (NC1, TBX3

13.6, TBX 13.8; Integrated DNA Technologies) were diluted in 300uL of Opti-MEM (final

concertation of 20nM). Lipofectamine RNAiMAX (6uL/100uL; Invitrogen 13778075) was then

diluted in 300uL of Opti-MEM. Diluted dsiRNA and Lipofectamine were mixed and set to incubate

at RT for 5 mins. 200ul of dsiRNA-Lipofectamine mix was added to each well. At 24h, cells were

lysed for RNA collection via (Zymo Research R2053). RNA purity was assessed via tapestation

with all samples having a RIN purity of ≥ 9.0. RNA samples were submitted to Omega

Bioservices (Norcross, GA, USA) for bulk RNAseq. RNA purity was again assessed via an Agilent

tapestation with all samples having a RIN ≥ 9.0. Library construction was completed using

NEBNext® Ultra™ II Directional RNA Library Prep Kit for Illumina®. Samples were sequenced

using NovaSeq 6000 (Illumina) PE150 with a ~40M (20M in each direction) read throughput for

each sample. Phred Mean Quality scores for all samples was ≥ 38.86. Data were analyzed by

ROSALIND® (https://rosalind.bio/), with a HyperScale architecture developed by ROSALIND, Inc.

(San Diego, CA).

529   Figure Construction

530   Figures were assembled and created with BioRender.com. UMAPs were generated with Partek

531   Flow. Venn diagrams were produced using DeepVenn. Binding distribution, Fig 3A, was modified

532   from original GREAT output, while Fig 3B was modified from the ChIP seeker output from

533   usegalaxy.org. Motif distribution for Fig 5D is modified from the MEME CentriMo tool.

535   Acknowledgments

536   We are grateful to Dr. Jonathan Zippin for providing the C57BL/6J-Mc1r^e/J mice and helpful

537   scientific discussion. We further extend thanks to the NHGRI Microarray (Dr. Abdel G. Elkahloun

538   and Bayu Sisay) and Flow Cytometry (Stacie Anderson and Martha Kirby) cores. Lastly, we

539   greatly appreciate the help of Dr. Laura Baxter in providing helpful editing and feedback.

References

1.) Ha, L., Merlino, G. & Sviderskaya, E. V. Melanomagenesis: Overcoming the barrier of melanocyte senescence. *Cell Cycle* **7**, 1944–1948 (2008).

2.) Bennett, D. C. & Medrano, E. E. Molecular Regulation of Melanocyte Senescence. *Pigm Cell Res* **15**, 242–250 (2002).

3.) Vredeveld, L. C. W. *et al.* Abrogation of BRAFV600E-induced senescence by PI3K pathway activation contributes to melanomagenesis. *Gene Dev* **26**, 1055–1069 (2012).

4.) Wagstaff, W. *et al.* Melanoma --- Molecular genetics, metastasis, targeted therapies, immunotherapies, and therapeutic resistance. *Genes Dis* **9**, 1608–1623 (2022).

5.) Rhodes, A. R., Weinstock, M. A., Fitzpatrick, T. B., Mihm, M. C. & Sober, A. J. Risk Factors for Cutaneous Melanoma: A Practical Method of Recognizing Predisposed Individuals. *Jama* **258**, 3146–3154 (1987).

6.) Rayner, J. E. *et al.* Phenotypic and genotypic analysis of amelanotic and hypomelanotic melanoma patients. *J Eur Acad Dermatol* **33**, 1076–1083 (2019).

7.) Schiöth, H. B. *et al.* Loss of Function Mutations of the Human Melanocortin 1 Receptor Are Common and Are Associated with Red Hair. *Biochem Bioph Res Co* **260**, 488–491 (1999).

8.) Pavan, W. J. & Sturm, R. A. The Genetics of Human Skin and Hair Pigmentation. *Annu Rev Genom Hum G* **20**, 1–32 (2019).

9.) Mort, R. L., Jackson, I. J. & Patton, E. E. The melanocyte lineage in development and disease. *Development* **142**, 620–632 (2015).

10.) Napolitano, A., Panzella, L., Monfrecola, G. & d'Ischia, M. Pheomelanin-induced oxidative stress: bright and dark chemistry bridging red hair phenotype and melanoma. *Pigm Cell Melanoma R* **27**, 721–733 (2014).

11.) Mitra, D. *et al.* An ultraviolet-radiation-independent pathway to melanoma carcinogenesis in the red hair/fair skin background. *Nature* **491**, 449–453 (2012).

12.)Robles-Espinoza, C. D. *et al.* Germline MC1R status influences somatic mutation burden

in melanoma. *Nat Commun* **7**, 12064 (2016).

13.)Cao, J. *et al.* MC1R Is a Potent Regulator of PTEN after UV Exposure in Melanocytes.

Mol Cell **51**, 409–422 (2013).

14.)HAUSCHKA, T. S., JACOBS, B. B. & HOLDRIDGE, B. A. Recessive Yellow and its

Interaction With Belted in the Mouse. *J Hered* **59**, 339–341 (1968).

15.)Michaud, E. J., Bultman, S. J., Stubbs, L. J. & Woychik, R. P. The embryonic lethality of

homozygous lethal yellow mice (Ay/Ay) is associated with the disruption of a novel RNA-

binding protein. *Gene Dev* **7**, 1203–1213 (1993).

16.)Michaud, E. J. *et al.* A molecular model for the genetic and phenotypic characteristics of

the mouse lethal yellow (Ay) mutation. *Proc National Acad Sci* **91**, 2562–2566 (1994).

17.)April, C. S. & Barsh, G. S. Skin layer-specific transcriptional profiles in normal and

recessive yellow (Mc1re/Mc1re) mice. *Pigm Cell Res* **19**, 194–205 (2006).

18.)Haltaufderhyde, K. D. & Oancea, E. Genome-wide transcriptome analysis of human

epidermal melanocytes. *Genomics* **104**, 482–489 (2014).

19.)Harris, M. L. *et al.* A direct link between MITF, innate immunity, and hair graying. *Plos*

Biol **16**, e2003648 (2018).

20.)Rezza, A. *et al.* Signaling Networks among Stem Cell Precursors, Transit-Amplifying

Progenitors, and their Niche in Developing Hair Follicles. *Cell Reports* **14**, 3001–3018

(2016).

21.)Hsiao, C. J. *et al.* Characterizing and inferring quantitative cell cycle phase in single-cell

RNA-seq data analysis. *Genome Res* **30**, 611–621 (2020).

22.)Belote, R. L. *et al.* Human melanocyte development and melanoma dedifferentiation at

single-cell resolution. *Nat Cell Biol* 1–13 (2021) doi:10.1038/s41556-021-00740-8.

23.)Guida, S., Guida, G. & Goding, C. R. MC1R Functions, Expression, and Implications for

Targeted Therapy. *J Invest Dermatol* (2021) doi:10.1016/j.jid.2021.06.018.

24.)Baxter, L. L., Watkins-Chow, D. E., Pavan, W. J. & Loftus, S. K. A curated gene list for

expanding the horizons of pigmentation biology. *Pigm Cell Melanoma R* **32**, 348–358

(2019).

25.)Hu, H. *et al.* AnimalTFDB 3.0: a comprehensive resource for annotation and prediction of

animal transcription factors. *Nucleic Acids Res* **47**, gky822 (2018).

26.)Macrì, S. *et al.* Hmga2 is required for neural crest cell specification in Xenopus laevis.

Dev Biol **411**, 25–37 (2016).

27.)Kee, Y. & Bronner-Fraser, M. Id4 expression and its relationship to other Id genes during

avian embryonic development. *Mech Develop* **109**, 341–345 (2001).

28.)Hu, J. *et al.* Endothelin signaling activates Mef2c expression in the neural crest through a

MEF2C-dependent positive-feedback transcriptional pathway. *Development* **142**, 2775–

2780 (2015).

29.)López, S. H. *et al.* Loss of Tbx3 in murine neural crest reduces enteric glia and causes

cleft palate, but does not influence heart development or bowel transit. *Dev Biol* **444**,

S337–S351 (2018).

30.)Seberg, H. E., Otterloo, E. V. & Cornell, R. A. Beyond MITF: Multiple transcription factors

directly regulate the cellular phenotype in melanocytes and melanoma. *Pigm Cell*

Melanoma R **30**, 454–466 (2017).

31.)Agarwal, P. *et al.* The MADS box transcription factor MEF2C regulates melanocyte

development and is a direct transcriptional target and partner of SOX10. *Development*

138, 2555–2565 (2011).

32.)DiVito, K. A., Trabosh, V. A., Chen, Y., Simbulan-Rosenthal, C. M. & Rosenthal, D. S.

Inhibitor of differentiation-4 (Id4) stimulates pigmentation in melanoma leading to

histiocyte infiltration. *Exp Dermatol* **24**, 101–107 (2015).

33.)Chhabra, Y. *et al.* Genetic variation in IRF4 expression modulates growth characteristics,

tyrosinase expression and interferon-gamma response in melanocytic cells. *Pigm Cell*

Melanoma R **31**, 51–63 (2018).

34.) Wang, C. *et al.* Donkey genomes provide new insights into domestication and selection

for coat color. *Nat Commun* **11**, 6014 (2020).

35.) Imsland, F. *et al.* Regulatory mutations in TBX3 disrupt asymmetric hair pigmentation that

underlies Dun camouflage color in horses. *Nat Genet* **48**, 152–158 (2016).

36.) Yin, K., Sturm, R. A. & Smith, A. G. MC1R and NR4A receptors in cellular stress and

DNA repair: implications for UVR protection. *Exp Dermatol* **23**, 449–452 (2014).

37.) Moon, H. *et al.* Melanocyte Stem Cell Activation and Translocation Initiate Cutaneous

Melanoma in Response to UV Exposure. *Cell Stem Cell* **21**, 665-678.e6 (2017).

38.) Zhang, T. *et al.* Cell-type–specific eQTL of primary melanocytes facilitates identification

of melanoma susceptibility genes. *Genome Res* **28**, 1621–1635 (2018).

39.) Perotti, V. *et al.* NFATc2 is an intrinsic regulator of melanoma dedifferentiation.

Oncogene **35**, 2862–2872 (2016).

40.) Smith, A. G., Lim, W., Pearen, M., Muscat, G. E. O. & Sturm, R. A. Regulation of NR4A

nuclear receptor expression by oncogenic BRAF in melanoma cells. *Pigm Cell Melanoma*

R **24**, 551–563 (2011).

41.) Boyd, S. C. *et al.* Oncogenic B-RAFV600E Signaling Induces the T-Box3 Transcriptional

Repressor to Repress E-Cadherin and Enhance Melanoma Cell Invasion. *J Invest*

Dermatol **133**, 1269–1277 (2013).

42.) Lu, J., Li, X.-P., Dong, Q., Kung, H. & He, M.-L. TBX2 and TBX3: The special value for

anticancer drug targets. *Biochimica Et Biophysica Acta Bba - Rev Cancer* **1806**, 268–274

(2010).

43.) Peres, J. *et al.* TBX3 promotes melanoma migration by transcriptional activation of ID1

which prevents activation of E-cadherin by MITF. *J Invest Dermatol* (2021)

doi:10.1016/j.jid.2021.02.740.

44.) Rodriguez, M., Aladowicz, E., Lanfrancone, L. & Goding, C. R. Tbx3 Represses E-

Cadherin Expression and Enhances Melanoma Invasiveness. *Cancer Res* **68**, 7872–

7881 (2008).

45.) Peres, J. *et al.* The Highly Homologous T-Box Transcription Factors, TBX2 and TBX3,

Have Distinct Roles in the Oncogenic Process. *Genes Cancer* **1**, 272–282 (2010).

46.) Perrot, C. Y. *et al.* GLI2 cooperates with ZEB1 for transcriptional repression of CDH1

expression in human melanoma cells. *Pigm Cell Melanoma R* **26**, 861–873 (2013).

47.) Heppt, M. V. *et al.* MSX1-Induced Neural Crest-Like Reprogramming Promotes

Melanoma Progression. *J Invest Dermatol* **138**, 141–149 (2018).

48.) Chen, Y. *et al.* ZEB1 Regulates Multiple Oncogenic Components Involved in Uveal

Melanoma Progression. *Sci Rep-uk* **7**, 45 (2017).

49.) Goding, C. R. & Arnheiter, H. MITF—the first 25 years. *Gene Dev* **33**, 983–1007 (2019).

50.) Murakami, M., Matsuzaki, F. & Funaba, M. Regulation of melanin synthesis by the TGF-β

family in B16 melanoma cells. *Mol Biol Rep* **36**, 1247–1250 (2009).

51.) Li, J., Weinberg, M. S., Zerbini, L. & Prince, S. The oncogenic TBX3 is a downstream

target and mediator of the TGF-β1 signaling pathway. *Mol Biol Cell* **24**, 3569–3576

(2013).

52.) Laurette, P. *et al.* Transcription factor MITF and remodeller BRG1 define chromatin

organisation at regulatory elements in melanoma cells. *Elife* **4**, e06857 (2015).

53.) Louphrasitthiphol, P. *et al.* Tuning Transcription Factor Availability through Acetylation-

Mediated Genomic Redistribution. *Mol Cell* **79**, 472-487.e10 (2020).

54.) Lu, S. *et al.* TBX2 controls a proproliferative gene expression program in melanoma.

Gene Dev **35**, 1657–1677 (2021).

55.) McLean, C. Y. *et al.* GREAT improves functional interpretation of cis-regulatory regions.

Nat Biotechnol **28**, 495–501 (2010).

56.) Strub, T. *et al.* Essential role of microphthalmia transcription factor for DNA replication,

mitosis and genomic stability in melanoma. *Oncogene* **30**, 2319–2332 (2011).

57.) Praetorius, C. *et al.* A Polymorphism in IRF4 Affects Human Pigmentation through a

Tyrosinase-Dependent MITF/TFAP2A Pathway. *Cell* **155**, 1022–1033 (2013).

58.) Aoki, H. & Moro, O. Involvement of microphthalmia-associated transcription factor (MITF)

in expression of human melanocortin-1 receptor (MC1R). *Life Sci* **71**, 2171–2179 (2002).

59.) Visser, M., Kayser, M. & Palstra, R.-J. HERC2 rs12913832 modulates human

pigmentation by attenuating chromatin-loop formation between a long-range enhancer

and the OCA2 promoter. *Genome Res* **22**, 446–455 (2012).

60.) Gautron, A. *et al.* Human TYRP1: two functions for a single gene? *Pigm Cell Melanoma*

R (2020) doi:10.1111/pcmr.12951.

61.) Washkowitz, A. J., Gavrilov, S., Begum, S. & Papaioannou, V. E. Diverse functional

networks of Tbx3 in development and disease. *Wiley Interdiscip Rev Syst Biology*

Medicine **4**, 273–283 (2012).

62.) Khan, S. F. *et al.* *Gene* **726**, 144223 (2019).

63.) Pape, E. L., Wakamatsu, K., Ito, S., Wolber, R. & Hearing, V. J. Regulation of

eumelanin/pheomelanin synthesis and visible pigmentation in melanocytes by ligands of

the melanocortin 1 receptor. *Pigm Cell Melanoma R* **21**, 477–486 (2008).

64.) Ni, J. *et al.* The effect of the NMDA receptor-dependent signaling pathway on cell

morphology and melanosome transfer in melanocytes. *J Dermatol Sci* **84**, 296–304

(2016).

65.) Tagore, M. *et al.* Electrical activity between skin cells regulates melanoma initiation.

Biorxiv 2021.12.19.473393 (2021) doi:10.1101/2021.12.19.473393.

66.) Enkhtaivan, E. & Lee, C. H. Role of Amine Neurotransmitters and Their Receptors in Skin

Pigmentation: Therapeutic Implication. *Int J Mol Sci* **22**, 8071 (2021).

67.) Palumbo, A. Melanogenesis in the Ink Gland of Sepia officinalis. *Pigm Cell Res* **16**, 517–

522 (2003).

68.) Eddy, K. & Chen, S. Glutamatergic Signaling a Therapeutic Vulnerability in Melanoma.

Cancers **13**, 3874 (2021).

69.) Janes, P. W. *et al.* EphA3 biology and cancer. *Growth Factors* **32**, 176–189 (2014).

70.) Theveneau, E. & Mayor, R. Neural crest migration: interplay between chemorepellents,

chemoattractants, contact inhibition, epithelial–mesenchymal transition, and collective

cell migration. *Wiley Interdiscip Rev Dev Biology* **1**, 435–445 (2012).

71.) Pape, E. L. *et al.* Microarray analysis sheds light on the dedifferentiating role of agouti

signal protein in murine melanocytes via the Mc1r. *Proc National Acad Sci* **106**, 1802–

1807 (2009).

72.) Arnheiter, H. The discovery of the microphthalmia locus and its gene, Mitf. *Pigm Cell*

Melanoma R **23**, 729–735 (2010).

73.) Levy, C., Khaled, M. & Fisher, D. E. MITF: master regulator of melanocyte development

and melanoma oncogene. *Trends Mol Med* **12**, 406–414 (2006).

74.) Li, J. *et al.* The Anti-proliferative Function of the TGF-β1 Signaling Pathway Involves the

Repression of the Oncogenic TBX2 by Its Homologue TBX3*. *J Biol Chem* **289**, 35633–

35643 (2014).

75.) Mesbah, K. *et al.* Identification of a Tbx1/Tbx2/Tbx3 genetic pathway governing

pharyngeal and arterial pole morphogenesis. *Hum Mol Genet* **21**, 1217–1229 (2012).

76.) Mohan, R. A. *et al.* T-box transcription factor 3 governs a transcriptional program for the

function of the mouse atrioventricular conduction system. *Proc National Acad Sci* **117**,

18617–18626 (2020).

77.) Singh, R. *et al.* Tbx2 and Tbx3 induce atrioventricular myocardial development and

endocardial cushion formation. *Cell Mol Life Sci* **69**, 1377–1389 (2012).

78.) Lüdtke, T. H. *et al.* Tbx2 and Tbx3 Act Downstream of Shh to Maintain Canonical Wnt

Signaling during Branching Morphogenesis of the Murine Lung. *Dev Cell* **39**, 239–253

(2016).

79.) Sheeba, C. J. & Logan, M. P. O. Chapter Twelve The Roles of T-Box Genes in

Vertebrate Limb Development. *Curr Top Dev Biol* **122**, 355–381 (2017).

80.) Frank, D. U., Emechebe, U., Thomas, K. R. & Moon, A. M. Mouse Tbx3 Mutants Suggest

724   Novel Molecular Mechanisms for Ulnar-Mammary Syndrome. *Plos One* **8**, e67841

725   (2013).

726   81.) Kaiser, M. *et al.* Regulation of otocyst patterning by Tbx2 and Tbx3 is required for inner

727   ear morphogenesis in the mouse. *Development* **148**, dev.195651 (2021).

728   82.) Dong, L. *et al.* Novel HDAC5-interacting motifs of Tbx3 are essential for the suppression

729   of E-cadherin expression and for the promotion of metastasis in hepatocellular

730   carcinoma. *Signal Transduct Target Ther* **3**, 22 (2018).

731   83.) Peres, J. & Prince, S. The T-box transcription factor, TBX3, is sufficient to promote

732   melanoma formation and invasion. *Mol Cancer* **12**, 117 (2013).

733   84.) Hoekstra, H. E. Genetics, development and evolution of adaptive pigmentation in

734   vertebrates. *Heredity* **97**, 222–234 (2006).

735   85.) Chintala, S. *et al.* Slc7a11 gene controls production of pheomelanin pigment and

736   proliferation of cultured cells. *P Natl Acad Sci Usa* **102**, 10964–10969 (2005).

737   86.) Poltorack, C. D. & Dixon, S. J. Understanding the role of cysteine in ferroptosis: progress

738   & paradoxes. *Febs J* **289**, 374–385 (2022).

739   87.) Willmer, T., Peres, J., Mowla, S., Abrahams, A. & Prince, S. The T-Box factor TBX3 is

740   important in S-phase and is regulated by c-Myc and cyclin A-CDK2. *Cell Cycle* **14**, 3173–

741   3183 (2015).

742   88.) Damerell, V. *et al.* The c-Myc/TBX3 Axis Promotes Cellular Transformation of Sarcoma-

743   Initiating Cells. *Frontiers Oncol* **11**, 801691 (2022).

744   89.) Coll, M., Seidman, J. G. & Müller, C. W. Structure of the DNA-Bound T-Box Domain of

745   Human TBX3, a Transcription Factor Responsible for Ulnar-Mammary Syndrome.

746   *Structure* **10**, 343–356 (2002).

747   90.) Langmead, B. & Salzberg, S. L. Fast gapped-read alignment with Bowtie 2. *Nat Methods*

748   **9**, 357–359 (2012).

749  91.) Zhang, Y. *et al.* Model-based Analysis of ChIP-Seq (MACS). *Genome Biol* **9**, R137

750    (2008).

751  92.) Bailey, T. L., Johnson, J., Grant, C. E. & Noble, W. S. The MEME Suite. *Nucleic Acids*

752    *Res* **43**, W39–W49 (2015).

753  93.) Quinlan, A. R. & Hall, I. M. BEDTools: a flexible suite of utilities for comparing genomic

754    features. *Bioinformatics* **26**, 841–842 (2010).

755  94.) Yu, G., Wang, L.-G. & He, Q.-Y. ChIPseeker: an R/Bioconductor package for ChIP peak

756    annotation, comparison and visualization. *Bioinformatics* **31**, 2382–2383 (2015).

Figures and Tables

759
Figure 1. Overview of single cell sequencing of primary murine melanocytes. (A) Schematic

of tissue harvest and digestion for melanocyte isolation from dorsal skin (a = nonagouti; A^y =

lethal yellow, $Mc1r^e$ = recessive yellow). (B) Representative FACS enrichment of c-KIT$^+$/CD45$^-$

cells isolated from the lethal yellow sample (red). (C-D) Uniform Manifold Approximation and

Projection (UMAP) plots of the sequenced cells that passed quality control for each pairwise

experiment. Among the cells isolated from nonagouti and lethal yellow paired samples (6,413

cells in C) and nonagouti and recessive yellow paired samples (4645 cells in D), skin cell types

were identified and labeled as follows: Melanocytes (Mc), Dermal fibroblasts (DF), Dermal papilla

(DP), Epidermis (Epi), Hair follicle and Outer root sheath (HF + ORS), and Transit amplifying and

Matrix cells (TAC + Mx). Right panels show UMAP plots of the 2256 cells classified as non-

cycling melanocytes. The nonagouti melanocytes (black) segregate from the recessive yellow

and lethal yellow (gold) melanocytes consistent with a eumelanin and pheomelanin producing

melanocytes having distinct transcriptional profiles.

774
Figure 2. Enriched processes and pathways present in the PGS. (A) Top 15 significant GO

terms ranked by fold-enrichment (FDR < 0.05). (B) Top 15 significant pathway terms from IPA

ranked by significance (p-value <0.05).

Figure 3. Identifying PGS regulatory factors. (A) List of all differentially expressed

transcription/co-transcription factors found within the PGS and ranked by FDR from the

nonagouti/lethal yellow pairwise differential analysis. (B) Top 10 predicted activated (green) and

inhibited (red) regulators of the PGS gene list ranked by Z-score.

Figure 4. ChIP-seq analysis implicates TBX3 as a regulator of melanogenesis. (A) Position

of called TBX3 peaks relative to the TSS. (B) ChIPseeker plot identifying genomic regions in

which TBX3 is bound. (C) Top 15 GO terms of genes associated with TBX3 ChIP target genes

ranked by -$\log_{10}$ of the FDR. (D) Top 5 predicted phenotype associations (human and mouse)

from TBX3 ChIP gene by -$\log_{10}$ of the FDR.

Figure 5. TBX3 binds MITF targets at E-boxes. (A) The top 10 motifs enriched within TBX3 ChIP peaks show over-representation of core E-box (red) and T-box (purple) sequences (B) The majority of TBX3 ChIP peaks contain either a T-box motif (14%), E-box motif (27%), or both (32%). (C) Top motifs of shared MITF and TBX3 peak sites (E-box = red; T-box = purple). (D) CentriMo plot showing top 9 centrally bound motifs.

Figure 6. TBX3 binds the E-box of pigmentation gene targets. (A) Intersection showing the number of murine genes (4072) associated with ChIP peaks in both our TBX3 dataset and a published MITF dataset. (B) Intersection of PGS genes identified as targets of both TBX3 and MITF. (C) UCSC browser view of TBX3 and MITF ChIP peaks associated with pigmentation genes. Browser tracks are as follows: TBX3 = replicated binding sites in our dataset; MITF = ChIP peaks from Strub et al; cCRE=encode annotated candidate cis-regulatory elements. The location of verified MITF regulatory elements containing E-box motifs are indicated (*) (refs in text).

Figure 7. TBX3 binds to and regulates PGS genes. (A) Heatmap depicting relative gene

expression of 2780 DEG (Adjusted p-value < 0.05) from the TBX3 knockdown samples against a

non-coding control. (B) Top KEGG terms ranked by -Log$_{10}$ of the p-value for upregulated (green)

and downregulated (red) genes. (C) Intersection of DE TBX3 gene targets against PGS TBX3

targets.

Table 1. Pigmentation phenotype associated genes found within the Pheomelanin Gene

Signature (PGS) with FDR and FC values for each pairwise experiment. In each pairwise

experiment a negative FC indicates reduced expression in the pheomelanin producing

melanocytes (lethal yellow, recessive yellow) compared to the eumelanin producing melanocytes

isolated from nonagouti littermates.

	Lethal yellow		Recessive yellow	
Gene	FDR step up	Fold change	FDR step up	Fold change
Tyrp1	0.00E+00	-19.31	1.12E-67	-83.90
Pmel	4.82E-46	-2.33	5.82E-16	-2.55
Mef2c	8.98E-40	2.69	1.93E-07	2.33
Oca2	1.77E-25	-3.16	9.28E-05	-2.47
Slc7a11	5.38E-23	7.72	3.79E-03	3.92
Irf4	1.32E-19	-3.02	7.19E-03	-2.61
Mc1r	2.58E-11	2.94	3.64E-03	3.63
Gas7	1.37E-07	-2.28	1.05E-02	-8.57
Sdc2	1.70E-07	2.39	8.94E-13	4.29
Pcbd1	1.25E-03	2.12	2.37E-02	2.34
Masp1	9.76E-03	3.11	1.73E-02	2.69

Supporting Information

S1 Fig. Classification Scheme for nonagouti and lethal yellow cells. (A) Uniform Manifold Approximation and Projection (UMAP) plot of lethal yellow and nonagouti cells classified by skin cell type: Melanocytes (Mc), Dermal fibroblasts (DF), Dermal papilla (DP), Epidermis (Epi), Hair follicle and Outer root sheath (HF + ORS), and Transit amplifying and Matrix cells (TAC + Mx). (B) UMAP of cells classified by graph-based cluster. (C) UMAPs depicting heatmap of cell type scoring.

S2 Fig. Classification Scheme for nonagouti and recessive yellow cells. (A) Uniform Manifold Approximation and Projection (UMAP) plot of the recessive yellow and nonagouti cells classified by skin cell type: Melanocytes (Mc), Dermal fibroblasts (DF), Dermal papilla (DP), Epidermis (Epi), Hair follicle and Outer root sheath (HF + ORS), and Transit amplifying and Matrix cells (TAC + Mx). (B) UMAP of cells classified by graph-based cluster. (C) UMAPs depicting heatmap of cell type scoring.

S3 Fig. Melanocytes cell cycle. (A) Total melanocytes from lethal yellow and nonagouti pairwise experiment (left), cells within the cell cycle highlighted in blue (right). (B) Total melanocytes from recessive yellow and nonagouti pairwise experiment (left), cells within the cell cycle highlighted in blue (right).

S4 Fig. Inducible TBX3 cell line validation. (A) schematic of piggyBAC transposase system with Tet-on inducible expression of TBX3. Dox, doxycycline; TRE, tetracycline response element; rtTA, reverse tetracycline-controlled transactivator. (B) Western blot of polyclonal 501mel-iTBX3-3xHA-PuroR inducible cells treated with varying amounts of doxycycline for 24h. (C) Venn diagram depicting shared peaks between both replicate ChIP experiments. (D) UCSC browser view of TBX3 and MITF ChIP peaks associated with E-cadherin. Browser tracks are as follows:

TBX3 = replicated binding sites in our dataset; MITF = ChIP peaks from Strub et al;

cCRE=encode annotated candidate cis-regulatory elements.

S5 Fig. Motif enrichment by fold-change direction in TBX3 KD studies. (A) Top 10 motifs,

ranked by e-value, of TBX3 ChIP peaks associated with putative target genes with increased

expression (left) or decreased expression (right) upon TBX3 knock-down.

S1-S5 Figure File. Berns et al_Supplementary Figures S1-S5.doc.

S1 Table. Gene Signature of Murine Skin Cells from Rezza et al.

S2 Table. Pheomelanin Gene Signature.

S3 Table. GREAT Output for TBX3 ChIP Dataset.

S4 Table. GREAT Output for MITF ChIP Dataset.

S5 Table. Human to Mouse Conversion of TBX3 Target Genes.

S6 Table. Human to Mouse Conversion of MITF Target Genes.

S7 Table. Gene Target Overlap.

S8 Table. Significant Differentially Expressed Genes from TBX3 Knockdown.

S9 Table. Significant Differentially Expressed Genes from TBX3 Knockdown which are

TBX3 Targets Identified by ChIP.

S1-S9 Tables File. Berns et al_Supplementary Tables S1-S9.xls.