TITLE

What happens to the inhibitory control functions of the right inferior frontal

cortex when this area is dominant for language?

AUTHORS

Esteban Villar-Rodríguez[1]

Cristina Cano-Melle[1]

Lidón Marin-Marin[1]

Maria Antònia Parcet[1]

César Avila*[1]

[1]Neuropsychology and Functional Neuroimaging; Jaume I University; Castellón

de la Plana, E-12071; Spain.

***Corresponding author:** email: avila@psb.uji.es – phone: +34 964387662

ABSTRACT

A low number of individuals show an atypical brain control of language

functions that differs from the typical lateralization in the left cerebral

hemisphere. In these cases, the neural distribution of other cognitive functions

is not fully understood. Although there is a bias towards a mirrored brain

organization consistent with the Causal hypothesis, some individuals are found

to be exceptions to this rule. However, no study has focused on what happens

to the homologous language areas in the right frontal inferior cortex. Using an

fMRI-adapted stop-signal task in a healthy non right-handed sample (50 typically

lateralized and 36 atypically lateralized for language production), our results

show that atypical lateralization is associated with a mirrored brain organization

of the inhibitory control network in the left hemisphere: inferior frontal cortex,

presupplementary motor area, and subthalamic nucleus. However, the

individual analyses revealed a large number of cases with a noteworthy overlap

in the inferior frontal gyrus, which shared both inhibitory and language

functions. Further analyses showed that atypical lateralization was associated

with stronger functional and structural interhemispheric connectivity.

Importantly, we did not find task performance differences as a function of

lateralization, but there was an association between atypical dominance in the

inferior frontal cortex and higher scores on schizotypy and autistic spectrum

traits, as well as worse performance on a reading accuracy test. Together, these

results partially support the Causal hypothesis of hemispheric specialization and

provide further evidence of the link between atypical hemispheric lateralization

and increased interhemispheric transfer through the corpus callosum.

KEYWORDS: Lateralization, hemispheric specialization, language, inhibitory

control, inferior frontal cortex, left-handers, corpus callosum, schizotypy,

autistic spectrum disorder, dyslexia.

INTRODUCTION

One of the oldest findings in human neuroscience is that the brain control of

language is lateralized to the left hemisphere[1–3]. However, not all individuals

present a typical organization of this function. Some left-handers (22-24%) show

an atypical lateralization of language – right or ambilateral[4–6]. Also, this atypical

organization is observed in some neurodevelopmental disorders such as

schizophrenia, dyslexia, or autistic spectrum[7–9].

How these atypical individuals organize the rest of their lateralized functions has

been an extensively researched question. Historically, two different hypotheses

have been proposed to explain the hemispheric specialization of the brain. The

Causal hypothesis[10,11] argues that rapid activities requiring the implementation

of sequences of cognitive processes are better performed from a single,

unilateral control. As a result, these activities are innately programmed to be

lateralized, and, according to this model, some functions will be performed

better if they are controlled by different hemispheres. The Statistical

hypothesis[12], on the other hand, postulates that each cognitive function

lateralizes independently from the others, making the probability of finding an

atypical hemispheric dominance merely a statistical phenomenon. The evidence

supporting each theory is mixed. Some studies have obtained data favoring the

Statistical hypothesis when analyzing language, visuospatial attention, and face

processing[13–15]. Most investigations, however, have demonstrated a mirrored

organization of praxis[16,17], visuospatial attention[14,17,18], face recognition[14,17,19], and

emotional prosody[17] in most individuals with atypical lateralization of language.

Nonetheless, it should be noted that this reversed pattern is not observed in all

atypical participants, and all the studies found specific cases in favor of both

theories[17].

All these studies have employed a production task to investigate language

lateralization in the inferior frontal cortex (IFC), classifying left-handed

participants into typical and atypical groups. The rest of the lateralized cognitive

functions have been examined using several tasks, but they all mainly depended

on posterior parts of the brain. This may have limited the results, given that they

were not dealing with directly homotopic functions, which would have been the

most sensitive scenario to the proposed advantages of hemispheric

specialization[20,21]. To directly address the language production function, which is

typically dependent on the left IFC, we would have to use a cognitive task that is

typically dependent on the homotopic right IFC. According to previous

literature, this function would be inhibitory control. Extensive results support

the notion that stopping an ongoing response involves a rightward network that

includes the IFC, presupplementary motor area (preSMA), and subthalamic

nucleus (STN)[22–25]. Thus, inhibitory control and language production together are

the ideal candidates to investigate the Causal vs. Statistical disjunctive. But

surprisingly, no investigation to date has analyzed this.

In this study, we explored the relationship between the lateralization of

language production and inhibitory control. To do so, we divided non-right-

handed participants into typically lateralized and atypically lateralized for

language using the verb generation task. These two groups then completed the

stop-signal task. We hypothesized that the atypical group would show a

mirrored brain organization during the stop-signal task. Additionally, we

investigated if a relation existed between the lateralization of both functions

and inter-hemispheric structural-functional connectivity[26], as well as with

behavioral markers of certain clinical conditions that have been related with

atypical lateralization[7–9].

METHOD

1. Participants

Eighty-six participants were included in the present study. They were selected

following a functional magnetic resonance imaging (fMRI) language

lateralization assessment via a verb generation task. Hence, 50 were typically

lateralized for language – left-dominant – (mean ± SD age = 22.4 ± 3.6 years; 24

male, 26 female) and 36 were atypically lateralized (mean ± SD age = 23.4 ± 4

years; 16 male, 20 female).

All the participants were non-right-handed, according to the Edinburgh

Handedness Inventory[27,28]. There were no significant between-group differences

in age (t_{84} = 1.28; P = 0.2), sex (χ^2 =0.11; P = 0.74), or general intelligence by

WAIS-IV matrix reasoning subtest[29] (t_{84} = 1.8; P = 0.08). The participants had no

history of any neurological or psychiatric disorders or head injury with loss of

consciousness. Written informed consent was obtained from all participants

following a protocol approved by Universitat Jaume I. All methods were carried

out in accordance with approved guidelines and regulations.

2. Experimental Design

Participants were recruited via multiple advertisements across Castellón and

Valencia universities (bulletin boards, mass emailing, etc.) asking for the

collaboration of left-handers in an fMRI brain study. Persons older than 36 years

or younger than 16 years were discarded. Valid participants were cited for a first

fMRI session, in which they completed the verb generation task and the resting-

state acquisition. During this session, we used the BrainWave software (GE

HealthCare Technologies Inc.) to visualize real-time data and roughly categorize

participants as potentially typical or potentially atypical. 43 participants were

found potentially atypical for language lateralization, and were invited to a

second fMRI session, along with 43 potentially typical participants. During this

second session, participants completed the stop-signal task (scanner) and a

reading skill test (out of scanner). Also, data regarding schizotypy personality

and autistic spectrum was gathered via self-questionnaires. Note that

classification of participants as potentially typical or potentially atypical was

used for screening purposes, and it did not completely match the final

assessment.

2.1 Verb generation Task

Expressive language function was measured via fMRI during a computerized

verb generation task[30,31] that consists of a 2-block design paradigm with

intercalating activation and control blocks. In the activation blocks, participants

are asked to overtly say the first verb that comes to mind when visually

presented with a concrete noun. In the control blocks, they have to read

visually-presented letter pairs aloud. This task lasted for 6 minutes, with a block

duration of 30 seconds (6 activation blocks and 6 control blocks), a stimulus

duration of 1500 ms, and a blank inter-stimulus interval of 1500 ms. Before

entering the scanner, participants practiced with a different version of the task

for 1 minute. Stimuli were presented using MRI-compatible goggles (VisuaStim

Digital, Resonance Technology Inc.), and responses were recorded with a noise-

cancelling microphone (FOMRI III+, Optoacoustics Ltd.) to verify that each

participant was engaged correctly in the task.

2.2 Stop-signal Task

Response inhibition was measured via fMRI during a computerized stop-signal

task adapted from another study[23]. The task consists of an event-related design

paradigm with 'Go' and 'Stop' trials. In the 'Go' trials, participants are asked to

manually answer by pressing a button if the visually presented string is a word

(has a meaning) or a pseudoword (mimics typical word structure but has no

meaning). Words require an index button press, whereas pseudowords require

a thumb button press. However, on the 'Stop' trials, the string is followed by a

'beep' noise that signals to the participants to inhibit their response and refrain

from pressing any button. The instructions emphasized the importance of going

correctly and stopping correctly. However, participants were asked to respond

as quickly and accurately as possible and avoid withholding their responses in

anticipation of a possible 'beep'.

This task took 13 minutes and 28 seconds, and it was divided into two runs that

lasted 6 minutes and 14 seconds each, separated by a 1-minute rest. Each run

consisted of 135 trials, of which 32 (23.7%) were 'Stop' trials. All the trials

randomly presented words or pseudowords. The trial structure consisted of a

fixation crosshair (500 ms), a word or pseudoword (1000 ms, during which the

'beep' may or may not be presented at some point), and a blank inter-stimulus

interval (ranging from 500 to 4000 ms, sampled from an exponential distribution

truncated at 4000 ms, with mean of 1000 ms). The Stop Signal Delay or SSD (the

amount of time after the onset of the word or pseudoword when the 'beep' is

presented during 'Stop' trials) changed dynamically during the task after each

'Stop' trial, depending on whether inhibition was successful (+25 ms) or

unsuccessful (−25 ms), with a minimum of 100 ms and a maximum of 800 ms.

The lower the SSD, the easier it is to inhibit the response, and vice versa. Hence,

a dynamic SSD normalizes the task's difficulty across all participants based on

their performance, aiming at a 50% successful inhibition rate. The SSD used for

the first 'Stop' trial was estimated for each participant based on their practice

session before entering the scanner. This practice session lasted 6 minutes and

used a different set of words and pseudowords. Stimuli were presented using

MRI-compatible goggles and headset (VisuaStim Digital, Resonance Technology

Inc.), and responses were recorded via an MRI-compatible response-grip

(ResponseGrips, NordicNeuroLab). The 'beep' volume was kept at a comfortable

level and was constant across participants.

2.3 Resting-state

Functional connectivity was measured via fMRI during a resting-state session[32].

In this paradigm, participants were presented with a fixation crosshair and

instructed to just lie in the scanner with their eyes open and try not to sleep or

think about anything in particular. This session lasted for 7 minutes. Seven

participants did not complete this session due to time constraints, and they

were subsequently removed from the functional connectivity analyses.

2.4 Behavioral measures

Individual inhibition speed was estimated by calculating the Stop-Signal

Reaction Time (SSRT)[33]. SSRT was computed as the difference between the

median reaction time (RT) on correct 'go' trials and the mean SSD on the stop-

signal task. It should be noted that technical problems involving the MRI-

compatible response-grip invalidated the SSD data of four participants, and so

their SSRT was estimated using the SSD data from the practice session instead.

Reading skill was evaluated using the word reading subtest of the PROLEC-SE-

R[34], a battery that assesses reading processes in Spanish. Participants had to

read four lists of words consisting, respectively, of short familiar words, long

familiar words, short unfamiliar words, and long unfamiliar words. Responses

were recorded with a microphone, and their accuracy and speed were

subsequently measured by computing length (performance during short words

– performance during long words) and familiarity (performance during familiar

words – performance during unfamiliar words) indicators.

Schizotypal traits were explored with the SPQ[35], a preclinical self-report

questionnaire modeled after DSM-III criteria[36]. This questionnaire evaluates

schizotypy based on the three-dimensional model: disorganized traits, cognitive-

perceptual traits, and interpersonal traits. For two participants, these data were

not collected due to time constraints.

Autistic spectrum traits were explored with the AQ[37], a preclinical self-report

questionnaire modeled after DSM-IV criteria[38]. This questionnaire evaluates

autistic spectrum based on the "triad" of impaired communication, impaired

social skills, and a restricted and repetitive way of acting[39], plus two sub-scales

for imagination and attention to detail. For two participants, these data were

not collected due to time constraints.

3. Image acquisition and processing

Images were acquired on a 3T General Electric Signa Architect magnetic

resonance imaging (MRI) scanner. A 3D structural MRI was acquired for each

subject using a T1-weighted magnetization-prepared rapid gradient-echo

sequence (TR/TE = 8.5/3.3 ms; matrix = 512 × 512 × 384; voxel size = 0.47 × 0.47

× 0.5). For the fMRI, a gradient-echo T2*-weighted echo-planar imaging

sequence was used in the acquisition of 150 functional volumes on the verb

generation task (TR/TE = 2500/30 ms; matrix = 64 × 64 × 30; voxel size = 3.75 ×

3.75 × 4), and a different sequence was used in the acquisition of 374 functional

volumes during the stop-signal task and 210 functional volumes in the resting-

state (TR/TE = 2000/30 ms; matrix = 64 × 64 × 27; voxel size = 3.75 × 3.75 × 4.5).

Task functional images were processed using the Statistical Parametric Mapping

software package (SPM12; Wellcome Trust Centre for Neuroimaging, London,

UK). Preprocessing followed the default pipeline and included: (a) alignment of

each participant's fMRI data to the AC-PC plane by using the anatomical image;

(b) head motion correction, where the functional images were realigned and

resliced to fit the mean functional image; (c) co-registration of the anatomical

image to the mean functional image; (d) re-segmentation of the transformed

anatomical image using a symmetric tissue probability map; (e) spatial

normalization of the functional images to the MNI (Montreal Neurological

Institute, Montreal, Canada) space with 3 mm^3 resolution; and (f) spatial

smoothing (FWHM = 4 mm). The general linear model for the verb generation

task was defined for each participant by contrasting *activation > control blocks*.

The general linear model for the stop-signal task was defined for each

participant by contrasting *correct 'stop' > correct 'go' trials.* For both tasks, the

BOLD (Blood-Oxygen-Level-Dependent) signal was estimated by convolving the

task's block/trial onsets with the canonical hemodynamic response function. Six

motion realignment parameters extracted from head motion preprocessing

were included as covariates of no interest, and a high-pass filter (128 s) was

applied to the contrast images to eliminate low-frequency components. After

that, mean BOLD values were extracted from the stop-signal contrast images for

the main components of the inhibitory control network: pars opercularis, pars

triangularis, preSMA, and STN. Pars opercularis and pars triangularis of the IFC

were defined following the criteria of the Harvard-Oxford atlas[40–43]. For the

PreSMA region, we used the SMA region, as defined by the same atlas, but

including only voxels anterior to MNI Y = 0[44]. STN was defined by a 10mm box

centered at MNI coordinates 10, −15, −5[22,45]. Lastly, we computed whole-brain

voxel-wise functional asymmetry maps from the stop-signal contrast images. To

do so, stop-signal contrast images were flipped at midline, inverting the right

and left hemispheres, and subsequently subtracted from the original unflipped

contrast images[46].

Resting-state functional images were processed using the Data Processing

Assistant for the Resting-State toolbox (DPARSFA)[47], which is based on SPM and

the Data Processing & Analysis of Brain Imaging toolbox (DPABI)[48]. Preprocessing

steps included: (a) slice-timing correction for interleaved acquisitions; (b) head

motion correction (no participant had a head motion of more than 2 mm

maximum displacement in any direction or 2° of any angular motion throughout

the scan); (c) co-registration of the anatomical image with the mean functional

image; (d) new segmentation to DARTEL; (e) removal of nuisance variance

through linear regression: six parameters from the head motion correction,

white matter signal, cerebrospinal fluid signal, and global mean signal; (f) spatial

normalization to the MNI (3 mm^3); (g) spatial smoothing (FWHM = 4 mm); (h)

removal of the linear trend in the time series; (i) band-pass temporal filtering

(0.01–0.1); (j) normalization to a symmetric template; (k) Voxel-Mirrored

Homotopic Computation (VMHC)[49], calculated as the Pearson correlation

coefficient of every voxel with its hemispheric counterpart; and (l) normalization

of all voxel-wise time courses to Fisher *z* values. Finally, mean VMHC values

were extracted for the pars opercularis and pars triangularis regions, as defined

by the Harvard-Oxford atlas[40–43].

Structural images were processed via voxel-based morphometry (VBM) using

the CAT12 toolbox[50], which is based on SPM. Preprocessing steps followed the

recommended pipeline and included: (a) segmentation into grey matter, white

matter, and cerebrospinal fluid; (b) registration to the ICBM standard template;

(c) modulated normalization of grey matter and white matter segments to the

MNI template; (d) spatial smoothing (FWHM = 6 mm); and (e) extraction of

Regions of Interest (ROIs) values from native space. Three ROIs were delimited

by the voxels mapped as genu, body and splenium of the corpus callosum

according to the Mori atlas[51]. Total intracranial volume was also extracted for

use as covariate of no interest.

4. Individual assessment of functional lateralization

Individual functional lateralization was assessed by calculating the Laterality

Index (LI) on the unflipped contrast images[30,31]. We used the bootstrap method

implemented in the LI-toolbox[52], based on SPM. LI is a proportion of the brain

activation between the two hemispheres, thus giving us information about the

direction and degree of hemispheric specialization during a particular function

in a single individual. LI ranges from +1 (totally leftward function) to −1 (totally

rightward function). For both the verb generation task and the stop-signal task,

we explored the LI of the inferior frontal region roughly corresponding to the

classic Broca's area: pars opercularis and pars triangularis of the inferior frontal

gyrus, according to the Harvard-Oxford atlas[40–43]. This region is critical for both

language production[3] and response inhibition[24], depending on the hemisphere.

During the verb generation task, language production was classified as typically

lateralized (LI higher than +0.4) or atypically lateralized (LI lower than +0.4). We

used +0.4 as a cut-off point (contrary to the more traditional +0.2), based on

previous findings that emphasized the importance of lateralization strength

when grouping individuals[6,53].

5. Statistical analyses

We performed different analyses to investigate the hypothesis that the atypical

group would show a mirrored brain organization during the stop-signal task.

First, a series of ROI analyses were employed to compare the functional

asymmetry of the main components of the inhibitory control network (during

the stop-signal task) between typically and atypically lateralized groups. Hence,

four repeated-measures MANOVAs (one for the mean BOLD value inside each

ROI: pars opercularis, pars triangularis, preSMA, and STN) were calculated,

including Hemisphere (left/right) as within-subject factor and Group

(typical/atypical) as between-subject factor. All *p*-values were FDR-corrected for

four comparisons. This analysis was complemented with whole-brain functional

asymmetry contrast maps from the stop-signal task that were compared in

typically and atypically lateralized groups based on language production. To do

so, we performed a whole-brain two-sample *t*-test (voxel-wise $P < 0.001$; FWE

cluster-corrected at $P < 0.05$) via SPM12. Next, a Pearson's correlation was

carried out between the LI from the verb generation task and the LI from the

stop-signal task used in the individual lateralization classification ($P < 0.05$, 1-

tailed). Finally, we explored the functional overlap between both tasks in the

subgroup of participants that presented an ambilateral organization of at least

one function (n = 48). To do so, we overlayed two one-sample *t*-tests restricted

to the IFC: one for the stop-signal task (voxel-wise $P < 0.001$; FWE cluster-

corrected at $P < 0.05$), and one for the verb generation task (voxel-wise $P < 0.05$;

uncorrected). Functional overlap was then expressed as the proportion of

overlapped voxels for every function and hemisphere.

Using the LIs from the verb generation and stop-signal tasks, we tested if a

linear relationship existed between the functional lateralization of an individual,

and several behavioral tests and neuroanatomical measures. Partial Spearman's

correlations were computed between the LI of both tasks and: stop-signal task

performance (RT, accuracy, and SSRT), SPQ derived-scores, AQ derived-scores,

reading test performance (length and familiarity effects on speed and accuracy),

functional connectivity measures (VMHC), and white matter volume of the

callosal genu. Age and general intelligence were included as covariates in all

analyses. Total intracranial volume was also included as covariate in the

volumetric analysis.

RESULTS

1. Inhibitory control shifts its lateralization according to language production

Inhibitory control components during the stop-signal task were defined by four

Regions Of Interest (ROIs) and subsequently analyzed by computing MANOVAs,

including Hemisphere as within-subject factor and Group (typical/atypical) as

between-subject factor. Results showed that all the tested structures presented

a statistically significant Hemisphere × Group interaction in their BOLD signal

during the 'correct stop > correct go' condition (pars opercularis $F_{1,84}$ = 25.59, P

< 0.001; pars triangularis $F_{1,84}$ = 19.95, P < 0.001; preSMA $F_{1,84}$ = 7.63, P = 0.007;

STN $F_{1,84}$ = 14.24, P < 0.001; FDR-corrected for 4 comparisons; see **Fig 1a**). In the

atypical group, all the regions seemed to have a shift towards leftward

lateralization of brain function.

A voxel-wise whole-brain analysis of the functional asymmetry maps during the

stop-signal task was performed to complement the ROI results. Two-sample *t*-

test comparisons of the typical and atypical groups during the 'correct stop >

correct go' condition showed a differential asymmetry pattern in all the ROI-

corresponding areas, while also revealing differences in other cortical and

subcortical regions, including the angular gyrus, middle temporal gyrus, and

caudate (**Fig 1b**). Specifically, the typical group presented a rightward activity in

these regions, whereas the atypical group exhibited a leftward shift (**Fig 1c**).

Finally, the Pearson's correlation between the frontal LIs (pars opercularis + pars

triangularis) from the verb generation task and the stop-signal task revealed a

strong inverse correlation ($r_{84} = -.58$; one-tailed $P < 0.001$; $R^2 = .34$) (**Fig 2**). Out

of 50 typically lateralized for language, 33 (66%) presented a typical right

lateralization for inhibitory control, and 17 (34%) showed an atypical (left –

ambilateral) lateralization. Among the 36 individuals from the atypical group,

only 8 (22.2%) presented a typical lateralization during the stop-signal task,

whereas 28 (77.8%) showed an atypical lateralization. Importantly, no

participant presented a leftward + leftward or rightward + rightward

segregation pattern for language production and response inhibition. Strong

segregation of both functions into different hemispheres was observed in 38

participants (44.2%). The remaining 48 individuals (55.8%) presented an

ambilateral representation of one or both functions, with an important level of

spatial overlap in their activations (see **Fig 1d**). In this ambilateral group,

language production presented an overlap in 32.3% of its voxels in the left

hemisphere, and in 58.6% of its voxels in the right hemisphere. Inhibitory

control showed an overlap in 57.2% of its voxels in the left hemisphere, and in

22.8% of its voxels of the right hemisphere.

2. Laterality Indexes correlate to interhemispheric connectivity and preclinical

markers

Correlational analyses revealed certain links between the functional

lateralization of both tasks and behavioral, neuroanatomical and connectivity

measures (**Table 1**).

First, we studied how the structural and functional connectivity between both

IFC behaved in relation to LI. Volume in the genu, body and splenium of the

corpus callosum was inversely related with language LI (respectively: $\rho_{83} = -.25$,

$P = 0.02$; $\rho_{83} = -.26$, $P = 0.02$; and $\rho_{83} = -.25$, $P = 0.02$). However, only the callosal

genu was found related with inhibition LI ($\rho_{83} = .25$, $P = 0.02$). That is, the

volume of the callosal genu increased as functional organization of the IFC

became more atypical, extending this effect to the whole corpus callosum when

considering exclusively language lateralization. On the same line, ROI

interhemispheric functional connectivity analyses at rest revealed that VMHC of

the pars triangularis also increased as a function of inhibition LI (ρ_{77} = .26, P =

0.02), but a similar relation failed to be found for language LI (ρ_{77} = −.12, P =

0.3). No statistically significant linear relationships were found when exploring

the VHMC of the pars opercularis.

Behaviorally, performance (RT, SSRT and accuracy) during the scanner stop-

signal task did not correlate with language and inhibition LIs. Regarding the

reading test, accuracy for word length and word familiarity were found

correlated with both LIs (length and language: ρ_{84} = −.24, P = 0.03; length and

inhibition: ρ_{84} = .21, P = 0.04; familiarity and language: ρ_{84} = −.23, P = 0.03;

familiarity and inhibition: ρ_{84} = .3, P = 0.01). That is, an atypical organization was

related with a higher rate of errors when reading long words and unfamiliar

words. No significant correlations were found when exploring reading speed.

SPQ score was found significantly related with inhibition LI (ρ_{82} = .25, P = 0.02),

but it did not reach statistical significance when paired with language LI (ρ_{84} =

−.19, P = 0.08). AQ, on the other hand, presented an association with language

LI (ρ_{84} = −.25, P = 0.02) but not with inhibition LI (ρ_{84} = .08, P = 0.5).

DISCUSSION

The present study employed the stop-signal task to investigate brain areas involved in the inhibitory control function in two different groups of individuals: typically and atypically lateralized for language production. As expected, the atypical participants showed, as a group, a mirrored brain organization of the inhibitory control function compared to the typical group. This leftward organization affected the entire inhibitory control network, including the IFC, preSMA, and STN. However, some participants manifested a clear overlap in the control of language and inhibitory functions. Our results also demonstrate that atypical organization of language production is associated with an increased white matter volume of the corpus callosum, and that atypical lateralization of inhibitory control is related with a higher interhemispheric functional coupling of the IFC. Behaviorally, atypical lateralization was not associated with performance on the task, but an association existed with worse reading accuracy, and more schizotypy and autistic traits.

Our first relevant result is that the group of individuals with atypical lateralization of language presented a mirrored brain organization during the stop-signal task compared to the typically lateralized group. Remarkably, the hemispheric asymmetry differences obtained affected the entire inhibitory

control brain network[25]. The inferior frontal cortex, or IFC, associated with the

initiation of the stop command and the main cortical area of this system,

showed a clear lateralization effect, with the typical group almost exclusively

using the right part and the atypical group mostly activating the left part. The

presupplementary motor area, or pre-SMA, related to the implementation of

the stop command, was more bilaterally involved in both groups, although

atypicals activated the left part more and typicals activated the right part more.

Importantly, the hemispheric reversal of organization also affected the

subcortical structures participating in this network, namely the STN and

thalamus. These structures displayed a lateralization pattern similar to the

preSMA, with a more pronounced bilateral engagement but higher involvement

of the right part in typicals and of the left part in atypicals. Thus, this is the first

demonstration that an entire cerebral network controlling a cognitive function

shows a completely flipped organization in an atypical population for language,

which supports the Causal hypothesis of lateralization[10,11]. Moreover, this result

reveals that hemispheric lateralization goes beyond cerebral cortices, and even

interhemispheric connectivity, through the corpus callosum, given that it also

involves frontal-subcortical circuits. Additionally, the specificity of this result to

the areas of the proposed network is another element that supports the

relevance of this system in inhibitory control.

It should be noted that this pattern of atypical brain specialization was observed

in the absence of a relation between functional lateralization and performance

differences during the 'go' or 'stop' conditions. Even though the 'go' condition

involves a lexical decision requiring language processing, and the 'stop'

condition consists of inhibiting a linguistic process, the functional organization

of the IFC during the tasks did not affect the response speed (RT and SSRT) or

the 'go' accuracy in either case. Thus, brain organization for language and

inhibition, as initially defined in our study, and cognitive efficiency are not

directly related, based on our data. This finding is consistent with previous

reports showing that cognitive deficits appear only in specific cognitive

domains[17,54].

Importantly, despite observing a complimentary relationship between language

production and inhibitory control at the group level, not all the individuals

presented a strong and segregated distribution of these two functions. The vast

majority of hemispherically segregated individuals showed the typical

segregation of language in the left hemisphere and inhibitory control in the right

hemisphere, whereas only a low percentage displayed the opposite segregated

organization. The rest, in agreement with similar reports on other functions[19],

lacked a clear hemispheric lateralization of language production, inhibitory

control, or both. In all cases, this implied some level of overlapping or sharing of

the same area of the inferior frontal cortex. Both the low ratio of atypical

segregation and the high ratio of bilateral inhibitory control could be explained

by the fact that typical right hemispheric functions (such as inhibition) seem to

be more bilaterally represented than typical left functions (such as language)[55].

Some conclusions can be extracted from this: (1) Although infrequent, a strong

atypical segregation of the two functions is possible; (2) Our data have not

revealed any case with a segregation of both frontal cognitive functions in the

same hemisphere; and (3) A large percentage of participants with a strong left

lateralization of language presented a bilateral control of inhibition, showing

that a typical dominance for language does not imply a strongly lateralized

organization of other cognitive functions. The existence of exceptions to the

group pattern of mirrored brain organization requires us to use caution when

interpreting these data as supporting the causal hypothesis. Although the

negative correlation between the lateralization indexes during the verb

generation task and the stop-signal task is strong at the group level, some

individual data support the statistical hypothesis[12] instead. Given that we found

no cases of both cognitive functions completely sharing a hemisphere (see **Fig**

2), we propose that causal-supporting vs. statistical-supporting results reflects

two different pathways to cognitive control. On the one hand, the segregated

pathway (in line with the Causal hypothesis) develops due to the cerebral bias

towards lateralizing certain cognitive functions in a single hemisphere. On the

other hand, the integrated pathway (in line with the Statistical hypothesis),

appears when ontogenetic development towards strong lateralized control fails

in some way.

The corpus callosum is the main cerebral structure in interhemispheric

connectivity, with the genu area being responsible for the connectivity between

the two inferior frontal gyri[56]. Phylogenetic data have demonstrated a negative

association between the corpus callosum volume and hemispheric

specialization, suggesting that the ontogenetic development of the brain is

supported by a decrease in interhemispheric connectivity to potentiate and

establish intrahemispheric connectivity and hemispheric specialization[57]. There

is evidence of this mechanism in language development[58] and in agenesis of the

corpus callosum, which has been associated with lower lateralization of

language and worse language performance[59,60]. In line with this proposal, we

have replicated the relation between callosal enlargement and atypical

functional organization found in a previous study[53]. Moreover, in our data, this

structural correlation was accompanied by an association between atypical

lateralization of inhibitory control and strength of interhemispheric functional

connectivity in the pars triangularis of the IFC.

As hypothesized, when considering the functional lateralization during both

tasks, atypical organizations were found linearly related with higher scores on a

schizotypy test (only for inhibitory control), higher scores on an autistic

spectrum test (only for language production), and a higher rate of reading errors

(both language production and inhibitory control). The first result is consistent

with previous structural, functional, and behavioral data that have shown

reduced language lateralization in schizophrenia patients[8,61]. In the case of

schizotypy, the evidence for reduced lateralization of cognitive functions is

mixed and may respond to differences in the methodology employed[62].

However, consistently with the present results, this condition has been

previously associated to impaired behavioral and neural processing during the

stop-signal task[63], and to an abnormal frontal functional asymmetry[64]. The

second result relates atypical language lateralization with the presence of

autistic traits, a result consistent with data obtained in autism[65]. The third result

shows that atypical organization was related with a higher rate of reading errors

when considering word length and word familiarity. The length effect has been

considered a pathognomonic symptom of reading disorders such as

developmental dyslexia[66] and pure alexia[67]. These disorders have also been

characterized by a weak language lateralization[7]. Thus, presented results

support the hypothesis that atypical hemispheric specialization is related with

worse cognitive performance in the linguistic domain, and even preclinical traits

of some neurodevelopmental disorders among healthy population.

DATA AVAILABILITY

All data presented in this study are included in the following public dataset:

https://doi.org/10.6084/m9.figshare.22047299.

ACKNOWLEDGMENTS

This work was supported by the Spanish State Research Agency (PID2019-

108198GB-I00) and the Universitat Jaume I (UJI-B2021-11)). E.V.-R. was

supported by a predoctoral graduate program grant from the Spanish Ministry

of Education, Culture and Sports (FPU18/00687). We also thank all of our

participants for their collaboration in this study, as well as the radiographers at

the clinic ASCIRES Castellón for their technical support during data acquisition.

AUTHOR CONTRIBUTIONS

Conceptualization, C.A. and M.-A.P.; Methodology, E.V.-R. and C.A.; Formal

Analysis, E.V.-R.; Investigation, E.V.-R., C.C.-M., and L.M.-M.; Data Curation, E.V.-

R. and C.C.-M., Writing, E.V.-R., C.A. and M.-A.P.; Supervision, C.A.; Project

Administration, C.A.

DECLARATION OF INTERESTS

The authors declare no competing interests.

REFERENCES

1.   Broca, P. (1861). Remarks on the Seat of the Faculty of Articulated Language, Following an
Observation of Aphemia (Loss of Speech). Bulletin de La Société Anatomique, 330–357.

2.   Wernicke, C. (1874). The Aphasic Symptom-Complex. Arch Neurol *3*, 2–80.

3.   Price, C.J. (2012). A review and synthesis of the first 20 years of PET and fMRI studies of
heard speech, spoken language and reading. Neuroimage *62*, 816–847.
10.1016/j.neuroimage.2012.04.062.

4.   Pujol, J., Deus, J., Losilla, J.M., and Capdevila, a (1999). Cerebral lateralization of language in
normal left-handed people studied by functional MRI. Neurology *52*, 1038–1043.
10.1212/WNL.52.5.1038.

5.   Szaflarski, J.P., Binder, J.R., Possing, E.T., McKiernan, K.A., Ward, B.D., and Hammeke, T.A.
(2002). Language lateralization in left-handed and ambidextrous people: fMRI data.
Neurology *59*, 238–244.

6.   Mazoyer, B., Zago, L., Jobard, G., Crivello, F., Joliot, M., Perchey, G., Mellet, E., Petit, L., and
Tzourio-Mazoyer, N. (2014). Gaussian mixture modeling of hemispheric lateralization for
language in a large sample of healthy individuals balanced for handedness. PLoS One *9*, 9–
14. 10.1371/journal.pone.0101165.

7.   Bishop, D.V.M. (2013). Cerebral asymmetry and language development: Cause, correlate,
or consequence? Science (1979) *340*. 10.1126/science.1230531.

8.   Sommer, I.E., Ramsey, N.F., and Kahn, R.S. (2001). Language lateralization in schizophrenia,
an fMRI study. Schizophr Res *52*, 57–67.

9.   Eyler, L.T., Pierce, K., and Courchesne, E. (2012). A failure of left temporal cortex to
specialize for language is an early emerging and fundamental property of autism. Brain *135*,
949. 10.1093/BRAIN/AWR364.

10.   Kosslyn, S.M. (1987). Seeing and Imagining in the Cerebral Hemispheres: A Computational
Approach. Psychol Rev *94*, 148–175. 10.1037/0033-295X.94.2.148.

11.   Hellige, J.B. (1990). Hemispheric Asymmetry. Annu Rev Psychol *41*, 55–80.
10.1146/annurev.ps.41.020190.000415.

12.   Bryden, M.P., Hécaen, H., and DeAgostini, M. (1983). Patterns of cerebral organization.
Brain Lang *20*, 249–262. 10.1016/0093-934X(83)90044-5.

13.   Whitehouse, A.J.O., and Bishop, D.V.M. (2009). Hemispheric division of function is the
result of independent probabilistic biases. Neuropsychologia *47*, 1938–1943.
10.1016/j.neuropsychologia.2009.03.005.

14.   Badzakova-Trajkov, G., Häberling, I.S., Roberts, R.P., and Corballis, M.C. (2010). Cerebral
Asymmetries: Complementary and Independent Processes. PLoS One *5*, e9682.
10.1371/journal.pone.0009682.

15.   Rosch, R.E., Bishop, D.V.M., and Badcock, N.A. (2012). Lateralised visual attention is
unrelated to language lateralisation, and not influenced by task difficulty - a functional
transcranial Doppler study. Neuropsychologia *50*, 810–815.
10.1016/j.neuropsychologia.2012.01.015.

16.   Vingerhoets, G., Alderweireldt, A.-S., Vandemaele, P., Cai, Q., Van der Haegen, L.,
Brysbaert, M., and Achten, E. (2013). Praxis and language are linked: Evidence from co-
lateralization in individuals with atypical language dominance. Cortex *49*, 172–183.
10.1016/J.CORTEX.2011.11.003.

17.   Gerrits, R., Verhelst, H., Vingerhoets, G., and Vingerhoets, G. (2020). Mirrored brain
organization: Statistical anomaly or reversal of hemispheric functional segregation bias?
Proc Natl Acad Sci U S A *117*, 14057–14065.
10.1073/PNAS.2002981117/SUPPL_FILE/PNAS.2002981117.SAPP.PDF.

18.   Cai, Q., Van Der Haegen, L., and Brysbaert, M. (2013). Complementary hemispheric
specialization for language production and visuospatial attention. Proc Natl Acad Sci U S A
110. 10.1073/pnas.1212956110.

19.   Gerrits, R., Van der Haegen, L., Brysbaert, M., and Vingerhoets, G. (2019). Laterality for
recognizing written words and faces in the fusiform gyrus covaries with language
dominance. Cortex *117*, 196–204. 10.1016/J.CORTEX.2019.03.010.

20.   Rogers, L.J. (2000). Evolution of hemispheric specialization: Advantages and disadvantages.
Brain Lang *73*, 236–253. 10.1006/BRLN.2000.2305.

21. Vallortigara, G., and Rogers, L.J. (2020). A function for the bicameral mind. Cortex *124*,
274–285. 10.1016/J.CORTEX.2019.11.018.

22. Aron, A.R., and Poldrack, R.A. (2006). Cortical and Subcortical Contributions to Stop Signal
Response Inhibition: Role of the Subthalamic Nucleus. 10.1523/JNEUROSCI.4682-05.2006.

23. Xue, G., Aron, A.R., and Poldrack, R.A. (2008). Common Neural Substrates for Inhibition of
Spoken and Manual Responses. Cerebral Cortex *18*, 1923–1932.
10.1093/CERCOR/BHM220.

24. Aron, A.R., Robbins, T.W., and Poldrack, R.A. (2014). Inhibition and the right inferior frontal
cortex: one decade on. Trends Cogn Sci *18*, 177–185. 10.1016/j.tics.2013.12.003.

25. Hannah, R., and Aron, A.R. (2021). Towards real-world generalizability of a circuit for
action-stopping. Nat Rev Neurosci *22*, 538–552. 10.1038/S41583-021-00485-1.

26. Tzourio-Mazoyer, N., Perrone-Bertolotti, M., Jobard, G., Mazoyer, B., and Baciu, M. (2017).
Multi-factorial modulation of hemispheric specialization and plasticity for language in
healthy and pathological conditions: A review. Cortex *86*, 314–339.
10.1016/j.cortex.2016.05.013.

27. Bryden, M.P. (1977). Measuring handedness with questionnaires. Neuropsychologia *15*,
617–624. 10.1016/0028-3932(77)90067-7.

28. Oldfield, R.C. (1971). The assessment and analysis of handedness: The Edinburgh inventory.
Neuropsychologia *9*, 97–113. 10.1016/0028-3932(71)90067-4.

29. Wechsler, D. (2012). Wais-IV: Escala de inteligencia de Wechsler para adultos IV (Madrid:
Pearson).

30. Sanjuán, A., Bustamante, J.C., Forn, C., Ventura-Campos, N., Barrós-Loscertales, A.,
Martínez, J.C., Villanueva, V., and Ávila, C. (2010). Comparison of two fMRI tasks for the
evaluation of the expressive language function. Neuroradiology *52*, 407–415.
10.1007/s00234-010-0667-8.

31. Villar-Rodríguez, E., Palomar-García, M., Hernández, M., Adrián-Ventura, J., Olcina-
Sempere, G., Parcet, M., and Ávila, C. (2020). Left-handed musicians show a higher
probability of atypical cerebral dominance for language. Hum Brain Mapp *41*, 2048–2058.
10.1002/hbm.24929.

32. Biswal, B., Zerrin Yetkin, F., Haughton, V.M., and Hyde, J.S. (1995). Functional connectivity
in the motor cortex of resting human brain using echo-planar mri. Magn Reson Med *34*,
537–541. 10.1002/MRM.1910340409.

33. Verbruggen, F., Aron, A.R., Band, G.P.H., Beste, C., Bissett, P.G., Brockett, A.T., Brown, J.W.,
Chamberlain, S.R., Chambers, C.D., Colonius, H., et al. (2019). A consensus guide to
capturing the ability to inhibit actions and impulsive behaviors in the stop-signal task. Elife
8. 10.7554/ELIFE.46323.

34. Cuetos Vega, F., Ramos Sánchez, J.L., and Arribas, D. (2016). PROLEC-SE-R : batería de
evaluación de los procesos lectores en secundaria y bachillerato.

35. Raine, A., and Raine, A. (1991). The SPQ: A Scale for the Assessment of Schizotypal
Personality Based on DSM-III-R Criteria. *17*.

36. American Psychiatric Association (1980). Diagnostic and Statistical Manual of Mental
Disorders, 3rd ed 3rd ed. 10.1176/ajp.138.1.128.

37. Baron-Cohen, S., Wheelwright, S., Skinner, R., Martin, J., and Clubley, E. (2001). The autism-
spectrum quotient (AQ): evidence from Asperger syndrome/high-functioning autism, males
and females, scientists and mathematicians. J Autism Dev Disord *31*, 5–17.
10.1023/A:1005653411471.

38. American Psychiatric Association (1994). Diagnostic and Statistical Manual of Mental
Disorders, 4th ed.

39. Rutter, M. (1978). Diagnosis and definition of childhood autism. J Autism Child Schizophr *8*,
139–161. 10.1007/BF01537863.

40. Frazier, J.A., Chiu, S., Breeze, J.L., Makris, N., Lange, N., Kennedy, D.N., Herbert, M.R., Bent,
E.K., Koneru, V.K., Dieterich, M.E., et al. (2005). Structural brain magnetic resonance
imaging of limbic and thalamic volumes in pediatric bipolar disorder. Am J Psychiatry *162*,
1256–1265. 10.1176/APPI.AJP.162.7.1256.

41. Desikan, R.S., Ségonne, F., Fischl, B., Quinn, B.T., Dickerson, B.C., Blacker, D., Buckner, R.L.,
Dale, A.M., Maguire, R.P., Hyman, B.T., et al. (2006). An automated labeling system for
subdividing the human cerebral cortex on MRI scans into gyral based regions of interest.
Neuroimage *31*, 968–980. 10.1016/J.NEUROIMAGE.2006.01.021.

42. Makris, N., Goldstein, J.M., Kennedy, D., Hodge, S.M., Caviness, V.S., Faraone, S. V., Tsuang,
M.T., and Seidman, L.J. (2006). Decreased volume of left and total anterior insular lobule in
schizophrenia. Schizophr Res *83*, 155–171. 10.1016/J.SCHRES.2005.11.020.

43. Goldstein, J.M., Seidman, L.J., Makris, N., Ahern, T., O'Brien, L.M., Caviness, V.S., Kennedy,
D.N., Faraone, S. V., and Tsuang, M.T. (2007). Hypothalamic Abnormalities in
Schizophrenia: Sex Effects and Genetic Vulnerability. Biol Psychiatry *61*, 935–945.
10.1016/J.BIOPSYCH.2006.06.027.

44. Ruan, J., Bludau, S., Palomero-Gallagher, N., Caspers, S., Mohlberg, H., Eickhoff, S.B., Seitz,
R.J., and Amunts, K. (2018). Cytoarchitecture, probability maps, and functions of the human
supplementary and pre-supplementary motor areas. Brain Struct Funct *223*, 4169–4186.
10.1007/S00429-018-1738-6.

45. Lucerna, S., Salpietro, F.M., Alafaci, C., and Tomasello, F. (2002). In Vivo Atlas of Deep Brain
Structures. In Vivo Atlas of Deep Brain Structures. 10.1007/978-3-642-56381-2.

46. Kurth, F., Gaser, C., and Luders, E. (2015). A 12-step user guide for analyzing voxel-wise gray
matter asymmetries in statistical parametric mapping (SPM). Nat Protoc *10*, 293–304.
10.1038/nprot.2015.014.

47.   Chao-Gan, Y., and Yu-Feng, Z. (2010). DPARSF: A MATLAB Toolbox for "Pipeline" Data
Analysis of Resting-State fMRI. Front Syst Neurosci *4*, 13. 10.3389/fnsys.2010.00013.

48.   Yan, C.-G., Wang, X.-D., Zuo, X.-N., and Zang, Y.-F. (2016). DPABI: Data Processing &
Analysis for (Resting-State) Brain Imaging. Neuroinformatics *14*, 339–351. 10.1007/s12021-
016-9299-4.

49.   Zuo, X.N., Kelly, C., Di Martino, A., Mennes, M., Margulies, D.S., Bangaru, S., Grzadzinski, R.,
Evans, A.C., Zang, Y.F., Castellanos, F.X., et al. (2010). Growing together and growing apart:
Regional and sex differences in the lifespan developmental trajectories of functional
homotopy. Journal of Neuroscience *30*, 15034–15043. 10.1523/JNEUROSCI.2612-10.2010.

50.   Gaser, C., and Kurth, F. (2016). Manual Computational Anatomy Toolbox - CAT12. 1–43.

51.   Oishi, K., Faria, A., Jiang, H., Li, X., Akhter, K., Zhang, J., Hsu, J.T., Miller, M.I., Van Zijl,
P.C.M., Albert, M., et al. (2009). Atlas-based whole brain white matter analysis using large
deformation diffeomorphic metric mapping: Application to normal elderly and Alzheimer's
disease participants. Neuroimage *46*, 486–499.

52.   Wilke, M., and Lidzba, K. (2007). LI-tool: A new toolbox to assess lateralization in functional
MR-data. J Neurosci Methods *163*, 128–136. 10.1016/j.jneumeth.2007.01.026.

53.   Labache, L., Mazoyer, B., Joliot, M., Crivello, F., Hesling, I., and Tzourio-Mazoyer, N. (2020).
Typical and atypical language brain organization based on intrinsic connectivity and
multitask functional asymmetries. Elife *9*, 1–31. 10.7554/eLife.58722.

54.   Mellet, E., Zago, L., Jobard, G., Crivello, F., Petit, L., Joliot, M., Mazoyer, B., and Tzourio-
Mazoyer, N. (2014). Weak language lateralization affects both verbal and spatial skills: An
fMRI study in 297 subjects. Neuropsychologia *65*, 56–62.
10.1016/j.neuropsychologia.2014.10.010.

55.   Gotts, S.J., Jo, H.J., Wallace, G.L., Saad, Z.S., Cox, R.W., and Martin, A. (2013). Two distinct
forms of functional lateralization in the human brain. Proc Natl Acad Sci U S A *110*, E3435.
10.1073/PNAS.1302581110/-/DCSUPPLEMENTAL.

56.   Hofer, S., and Frahm, J. (2006). Topography of the human corpus callosum revisited—
Comprehensive fiber tractography using diffusion tensor magnetic resonance imaging.
Neuroimage *32*, 989–994. 10.1016/J.NEUROIMAGE.2006.05.044.

57.   Tzourio-Mazoyer, N. (2016). Intra- and Inter-hemispheric Connectivity Supporting
Hemispheric Specialization. Research and Perspectives in Neurosciences, 129–146.
10.1007/978-3-319-27777-6_9.

58.   Tzourio-Mazoyer, N., and Seghier, M.L. (2016). The neural bases of hemispheric
specialization. Neuropsychologia *93*, 319–324. 10.1016/j.neuropsychologia.2016.10.010.

59.   Hinkley, L.B.N., Marco, E.J., Brown, E.G., Bukshpun, P., Gold, J., Hill, S., Findlay, A.M.,
Jeremy, R.J., Wakahiro, M.L., James Barkovich, A., et al. (2016). The Contribution of the
Corpus Callosum to Language Lateralization. J Neurosci *36*, 4522–4533.
10.1523/JNEUROSCI.3850-14.2016.

60. Bartha-Doering, L., Schwartz, E., Kollndorfer, K., Fischmeister, F.P.S., Novak, A., Langs, G.,
Werneck, H., Prayer, D., Seidl, R., and Kasprian, G. (2021). Effect of corpus callosum
agenesis on the language network in children and adolescents. Brain Struct Funct *226*, 701.
10.1007/S00429-020-02203-6.

61. Li, X., Branch, C.A., Ardekani, B.A., Bertisch, H., Hicks, C., and DeLisi, L.E. (2007). fMRI study
of language activation in schizophrenia, schizoaffective disorder and in individuals
genetically at high risk. Schizophr Res *96*, 14–24. 10.1016/j.schres.2007.07.013.

62. Park, H.R.P., and Waldie, K.E. (2017). Associations between schizotypy and cerebral
laterality. Laterality *22*, 195–218. 10.1080/1357650X.2016.1154066.

63. Jia, L. xia, Qin, X. jing, Cui, J. fang, Zheng, Q., Yang, T. xiao, Wang, Y., and Chan, R.C.K.
(2021). An ERP study on proactive and reactive response inhibition in individuals with
schizotypy. Sci Rep *11*. 10.1038/s41598-021-87735-5.

64. Le, T.P., Lucas, H.D., Schwartz, E.K., Mitchell, K.R., and Cohen, A.S. (2020). Frontal alpha
asymmetry in schizotypy: electrophysiological evidence for motivational dysfunction.
https://doi.org/10.1080/13546805.2020.1813096 *25*, 371–386.
10.1080/13546805.2020.1813096.

65. Jouravlev, O., Kell, A.J.E., Mineroff, Z., Haskins, A.J., Ayyash, D., Kanwisher, N., and
Fedorenko, E. (2020). Reduced Language Lateralization in Autism and the Broader Autism
Phenotype as Assessed with Robust Individual-Subjects Analyses. Autism Res *13*, 1746–
1761. 10.1002/AUR.2393.

66. Provazza, S., Giofrè, D., Adams, A.M., and Roberts, D.J. (2019). The Clock Counts – Length
Effects in English Dyslexic Readers. Front Psychol *10*, 2495.
10.3389/FPSYG.2019.02495/BIBTEX.

67. Roberts, D.J., Woollams, A.M., Kim, E., Beeson, P.M., Rapcsak, S.Z., and Lambon Ralph, M.A.
(2013). Efficient Visual Object and Word Recognition Relies on High Spatial Frequency
Coding in the Left Posterior Fusiform Gyrus: Evidence from a Case-Series of Patients with
Ventral Occipito-Temporal Cortex Damage. Cerebral Cortex *23*, 2568–2580.
10.1093/CERCOR/BHS224.

FIGURES

Fig 1. Hemispheric lateralization of inhibitory control according to language lateralization.

(A) ROI analysis of the main components of the inhibitory control network. Graphs depict adjusted mean BOLD signal during successful inhibitions on the stop-signal task ('stop > go' contrast) for both hemispheres and both groups. All four structures showed significant Hemisphere × Group interactions. Error bars represent one standard error. pOper = pars opercularis, pTri = pars triangularis, preSMA = presupplementary motor area, and STN = subthalamic nucleus. **(B)** Voxel-wise whole-brain analysis of functional asymmetry. Significance maps (voxel-wise $P < 0.001$; FWE cluster-corrected at $P < 0.05$; color bar represents t

value) are displayed in three-dimensional reconstructions plus coronal and

transversal slices using MNI space. (**C**) Mean BOLD values of the significant

regions found in the voxel-wise whole-brain analysis. Graphic depicts BOLD

values for every region, hemisphere, and group. IFC = inferior frontal cortex,

preSMA = presupplementary motor area, AngG = angular gyrus, MTG = middle

temporal gyrus, Thal = thalamus, STN = subthalamic nucleus, Caud = caudate.

(D) Functional overlap between language production and inhibitory control in

the IFC of ambilateral participants. Conjunction maps for inhibition (voxel-wise

$P < 0.001$; FWE cluster-corrected at $P < 0.05$) and language (voxel-wise $P < 0.05$;

uncorrected) are displayed in coronal, sagittal and transversal slices using MNI

space. VGT = verb generation task, SST = stop-signal task.

Fig 2. Correlation between the LIs of the verb generation and stop-signal tasks.

$r = -.583$, one-tailed $P < 0.001$, $R^2 = .339$. Negative values indicate rightward lateralization, whereas positive values indicate leftward lateralization. Segregated and ambilateral phenotypes are also depicted according to the background color. The green area corresponds to segregated individuals (both

functions strongly lateralized), and the red area corresponds to integrated

individuals (at least one function ambilaterally controlled). Numbers inside each

quadrant denote the number of individuals (n) contained in it. Each individual

datapoint is symbolized according to its functional organization: ○ = typical

segregation; ● = reversed segregation; △ = ambilateral inhibition; ▽ =

ambilateral language; ◇ = ambilateral language and inhibition.

TABLES

	Verb generation task LI		Stop-signal task LI	
	Spearman's ρ	P	Spearman's ρ	P
Callosal genu volume	−.249	.021*	.251	.02*
Callosal body volume	−.258	.017*	.132	.224
Callosal splenium volume	−.249	.021*	.118	.279
Pars Opercularis VMHC	−.062	.584	.148	.192
Pars Triangularis VMHC	−.119	.295	.258	.022*
'Go' reaction time	.125	.251	−.041	.71
'Go' accuracy	−.076	.489	.039	.724
SSRT	.173	.111	−.038	.726
Reading length accuracy	−.237	.028*	.214	.047*
Reading familiarity accuracy	−.232	.032*	.301	.005**
Reading length time	.054	.622	−.184	.068
Reading familiarity time	.091	.403	−.198	.091
SPQ	−.194	.078	.247	.023*
AQ	−.247	.023*	.075	.498

766 **Table 1. Spearman's partial correlations between task LIs and**

767 **neuroanatomical plus behavioral variables.**

768 General intelligence and age were included as covariates of no interest. Callosal

769 volume correlations were additionally corrected for total intracranial volume.