Impact of Socioeconomic Status on Amygdala and Hippocampus Subdivisions in Children and Adolescents

Jamie L Hanson[1]*, Dorthea J Adkins[1]*, Brendon M Nacewicz[2] & Kelly R Barry[3]

1: University of Pittsburgh, 2: University of Wisconsin-Madison, 3: University of Houston

(*denotes equal contribution of authorship*)

Author Notes
**Corresponding Author:** Jamie Hanson, Ph.D., *Learning Research & Development Center University of Pittsburgh, Murdoch Building 3420 Forbes Ave. Rm. 639 Pittsburgh, PA 15260,* Telephone: 412-383-3250, Email: jamie.hanson@pitt.eduad

*Keywords*: socioeconomic status; neurodevelopment; neurobiology; amygdala; hippocampus; sex differences

**ABSTRACT**

Socioeconomic status (SES) in childhood can impact behavioral and brain development. Past work has consistently focused on the amygdala and hippocampus, two brain areas critical for emotion, memory, and learning. While there are SES differences in total amygdala and hippocampal volumes, there are many unanswered questions in this domain connected to neurobiological specificity and whether these effects vary by participant age or sex. To address these gaps, we combined multiple large neuroimaging datasets of children and adolescents with information about neurobiology and SES (N=2,765). We examined subdivisions of the amygdala and hippocampus, derived from Freesurfer, using linear mixed effects models. Higher SES was associated with larger volumes across all three amygdala subdivisions examined, the superficial cortical division, basolateral complex, and centromedial region. Within the hippocampus, SES was specifically related to volumes in the head, with no significant associations for the body or tail. Contrary to our hypotheses, we found no significant interactions between SES and participant age or sex after correcting for multiple comparisons, suggesting these associations were relatively consistent across the developmental period examined (ages 5-18) and similar for males and females. These results fill in important gaps regarding the neurobiological specificity of SES effects, demonstrating associations across functionally distinct subdivisions of these critical brain structures.

**INTRODUCTION**

Socioeconomic status (SES) in childhood has been associated with multiple negative physical and mental health outcomes, with several meta-analyses noting these links (Peverill et al., 2021; Piotrowska et al., 2015). These disparities, in part, reflect structural inequalities in how resources, opportunities, and risks are distributed across socioeconomic strata, including unequal access to quality education, healthcare, safe housing, and economic security (Adler & Newman, 2002; Braveman & Gottlieb, 2014). The mechanisms underlying these relations, however, are poorly understood. An emerging approach leverages precise quantitation of neurobiology to understand SES-gradients of health (Hair et al., 2015, 2022; Hanson & Hackman, 2012). Neuroscientific investigations may allow a more elemental focus, as the brain determines behavioral and physiological responses (McEwen, 2007). This focus may be particularly valuable given the protracted nature of brain development and that the brain is shaped by experiences early in life (Noble & Giebler, 2020).

A growing body of research has found neurobiological alterations in samples exposed to poverty or lower SES conditions (Johnson et al., 2016; Noble & Giebler, 2020). Notably, childhood poverty has been implicated in structural differences across multiple brain regions, with differences in hippocampal and amygdala structure being commonly reported. The link between childhood poverty and smaller hippocampal volumes has been replicated by at least eight research groups (Ellwood-Lowe et al., 2018; Hanson et al., 2011; Luby et al., 2013; McDermott et al., 2019; Noble et al., 2012, 2015; Raffington et al., 2019; Weissman et al., 2023; Yu et al., 2018). However, studies examining the directional impact of childhood poverty on amygdala structure have produced a landscape of heterogeneous results, with reports of larger and smaller amygdalae (Hanson & Nacewicz, 2021). Given these areas' connections to important socioemotional functions and learning, understanding how poverty may shape these regions could shed light onto the mechanisms of SES-related disparities. The amygdala is a central neural hub for vigilance and processing negative emotions (Adolphs, 2010; LeDoux, 2007). The hippocampus plays a critical role in memory representations and using previously acquired information in service of goal-directed behavior (Murty et al., 2016; Shohamy & Turk-Browne, 2013). As such, these brain areas are critical for emotion and behavioral responding.

While there are SES differences in amygdala and hippocampal volumes, there are many unanswered questions in this domain connected to neurobiological specificity, and for whom these effects may be more pronounced. First, related to neurobiology, while research often treats the amygdala and hippocampus as unitary structures, they are complex and heterogeneous. Different amygdala nuclei have unique connectivity profiles, patterns of developmental changes, and behavioral correlates (Klein-Flügge et al., 2022; LeDoux, 2007; Swanson & Petrovich, 1998). Similarly, the hippocampus is composed of functionally distinct subregions, with differential connectivity and cytoarchitectonics (Gruber & Ranganath, 2019; Strange et al., 2014; Vogel et al., 2020). The posterior hippocampus has been linked more to cognitive functions, while more anterior regions relate to stress and affect (Fanselow & Dong, 2010; Poppenk & Moscovitch, 2011). Second, there may be potential sociodemographic subgroups where the effects are more pronounced; specifically, sex and age may both moderate the impact of SES on neurobiology. Motivated in part by sex disparities in many neuropsychiatric disorders (Hartung & Lefler, 2019), there has been a growing emphasis on sex as a biological variable. After stress exposure, sex differences in neuronal firing, dendritic spines, neurogenesis, and fMRI responsivity have been found in both the amygdala and the hippocampus (Stevens & Hamann, 2012; van Eijk et al., 2020; Yagi & Galea, 2019). These sex differences may be due to different neuroendocrine processes, responses from the environment, or sex chromosome-specific neuroprogramming (Bale & Epperson, 2017; Shansky

& Woolley, 2016). Sex may also be related to differential responses to stress exposure, like those associated with lower SES (Bath, 2020). Related to age, there are non-linear trajectories for brain development, with many structures increasing in volume in childhood, and then showing lower volumes in adolescence and adulthood (Lenroot & Giedd, 2006). The impacts of stress on neurobiology may vary with age and development (Hanson & Nacewicz, 2021). Stress may increase volumes in certain regions early in development, but then relate to "excitotoxic burnout" and smaller volumes later in time. As such, it will be critical to explore connections between neurobiology, age, sex, and SES.

Connected to neurobiological specificity, there is a growing body of past work examining amygdala and hippocampal subdivisions after childhood adversity. For the hippocampus, past projects have commonly reported smaller volumes in Cornu Ammonis (CA) 1 for those exposed to high levels of adversity (Lee et al., 2018; Margolis et al., 2022; Teicher et al., 2012; Yuan et al., 2020); however, results are not perfectly uniform with other studies only finding differences in CA3, and not CA1 (Brody et al., 2017; Malhi et al., 2019). With the amygdala, less work has been completed. Two studies reported smaller basolateral amygdala volumes after exposure to adversity (Nogovitsyn et al., 2022; Oshri et al., 2019), but these projects also reported stress was sometimes related to differences in accessory basal, central-medial, and paralaminar subdivisions. Related to SES, limited work has examined if there are potential alterations in volumes of amygdala and hippocampal subdivisions. Three past studies have found anterior hippocampal volumes (i.e., Dentate gyrus; CA1) were positively related to different operationalizations of SES, including parental education (Merz et al., 2019), family household income (Decker et al., 2020), and socioeconomic conditions in a census tract (Botdorf et al., 2022). Notably, no work to date has examined amygdala subdivisions in relation to SES.

Attempting to overcome these limitations, here, we combined multiple, large neuroimaging datasets of children and adolescents with information about neurobiology and SES (N=2,765). To improve neurobiological specificity, we examined subdivisions of the amygdala and hippocampus. We aimed to richly probe the main effects of SES on these smaller regions, as well as examine potential sex- or age-specific impacts of SES on these volumes. In keeping with past reports of smaller volumes in lower SES youth, we predicted lower SES would be related to smaller volumes in amygdala and hippocampal subdivisions. Related to past human and non-human research in stress-exposed groups (Hanson & Nacewicz, 2021), we also predicted that lower SES would be related to smaller volumes in the head of the hippocampus, as well as the basolateral and central amygdala. Finally, given potential neuroprotective effects of estrogen (McEwen, 2002, 2020) and developmental trajectories of brain development (Aoki et al., 2017), we predicted relations between SES and volumes would be stronger for older participants, especially boys.

## METHOD

**Participants.** Participants between 5-17 years of age were drawn from four large neuroimaging projects: The National Consortium on Alcohol and Neuro-Development in Adolescence (NCANDA; (Brown et al., 2015), the Healthy Brain Network (HBN; (Alexander et al., 2017), the Pediatric Imaging, Neurocognition, and Genetics (PING study) (Jernigan et al., 2016), and the Human Connectome Project in Development (HCP-D; (Somerville et al., 2018). Richer sample descriptions are available in our supplemental materials. Across these studies, the total number of participants with usable data was N=2765 (44% Female; Mean Age= 11.9, Age SD= 3.5; Age Range = 5.04-17.99). Descriptive information for the combined sample is shown below in Table 1. Information for each project is noted in our Supplement (Table S1). A histogram of participant age by study is shown in Figure 1.

*— Insert Table 1 Approximately Here—*

*— Insert Figure 1 Approximately Here—*

**MRI Data Acquisition and Imaging Processing.** High-resolution T1-weighted structural images were acquired with varying parameters across each project. The majority of scans came from 3T scanners, with the exception of a 1.5T scanner used at one site. Scans resolution varied in in-plane resolution from 0.8 to 1.2mm. Information about MRI parameters are noted in our supplemental materials. These MRI scans were processed in Freesurfer 7.1, deployed via Brainlife.io (Hayashi et al., 2024). Freesurfer is a widely documented morphometric processing tool suite, http://surfer.nmr.mgh.harvard.edu/ (Dale et al., 1999; Fischl et al., 2002). Based on hand-tracing on high-definition, ex-vivo T1-weighted 7T scans, Freesurfer can output 12 subfields and 9 amygdala subnuclei (see (Iglesias et al., 2015; Saygin et al., 2017) for additional details). With the hippocampus, we focused on segmentation that divided this region into the head, body, and tail. This was motivated by: 1) work finding that hippocampal organization and connectivity varies on its longitudinal axis (i.e., head/body/tail) (Genon et al., 2021; Vogel et al., 2020); and 2) commentary suggesting that segmentation of the longitudinal axis of the hippocampus is more appropriate, than much smaller (and potentially less reliably) segmented subfields, i.e., dentate gyrus, and subiculum (Kahhale et al., 2023; Wisse et al., 2021). With the amygdala, subnuclei volumes were grouped into three divisions based on cytoarchitecture, connectivity, and functional organization (Amunts et al., 2005; Saygin et al., 2017): (1) a basolateral complex (lateral, basal, and accessory basal nuclei), comprising glutamatergic principal neurons that serve as the primary input station for cortical and thalamic sensory information and support associative learning (Paton et al., 2006; Sah et al., 2003); (2) a centromedial division (central and medial nuclei) consists predominantly of GABAergic output neurons that project to hypothalamic and brainstem autonomic centers to coordinate neuroendocrine and behavioral stress responses (Ciocchi et al., 2010); (3) a superficial cortical division (cortical nucleus, anterior amygdaloid area, and corticoamygdaloid transition area) represents transitional cortical-like regions with prominent olfactory and limbic connectivity (Kemppainen et al., 2002; Majak & Pitkänen, 2003). The paralaminar nucleus was excluded from analyses due to ongoing debates about subgroups (deCampo & Fudge, 2012; Kedo et al., 2018), as well as segmentation uncertainty and inconsistent detection across participants (Kahhale et al., 2023). For the 3 hippocampal and the 3 amygdala subdivision groups, we calculated the total volume by summing volumes from the left and right hemispheres of each of these hippocampal or amygdala subdivisions.

*— Insert Figure 2 Approximately Here—*

To exclude particularly high-motion scans and limit the impact of image quality on subcortical segmentation, we generated a quantitative metric of image quality combining noise-to-contrast ratio, coefficient of joint variation, inhomogeneity-to-contrast ratio, and root-mean-squared voxel resolution (Gaser et al., 2024). Scans with an image quality grade < 0.8 were excluded from analyses, a threshold determined to be effective in previous projects for ensuring adequate volumetric and morphometric measurements (Adkins & Hanson, 2025; Bacas et al., 2023; Gilmore et al., 2021; Hanson et al., 2024). The image quality metric was then included as a covariate in all statistical models to further account for remaining variability in scan quality among retained scans. This was motivated by our work finding that T1-weighted image quality is related to Freesurfer outputs (Gilmore et al., 2021). Additional details about this metric, image processing in Freesurfer, and MRI Data Acquisition are noted in our supplemental materials.

**Measures of SES.** We used a multimethod determination of SES, using metrics of both caregiver education and household income. For caregiver education, caregivers reported on how much school they completed (e.g., obtained a high school diploma; some college; graduate degree); this was converted into numbers of years (i.e., high school diploma = 12 years; some college = 14 years) and then took the highest value by either caregiver. For household income, caregivers selected an income range (i.e., $30,000-39,999/year; $50,000-$99,999/year), or reported a continuous value. In keeping with recommendations from demographers (Parker & Fenwick, 1983), we took the midpoint for reported ranges and then log-transformed this value (i.e., $30,000-39,999=$34,999.5; log($34,999.5)=4.54406184). For continuous income reports, we also log-transformed these values. Then, we created an SES composite by taking an average of z-scored, log-transformed income and z-scored, maximum parental education.

**Analytic plan**. To test relations between SES and volumes of the hippocampus and amygdala, we employed linear mixed effects models (LMEMs) using the *lme4* R package (Bates et al., 2015). In all models, we included a random effect for study site, and examined the independent (fixed effect) variables of participant age (in years), participant sex (binary-coded), estimated Total Intracranial Volume (eTIV), image quality (CAT12 rating), our SES composite, and interactions between age, sex, and SES. These models include two- and three-way interactions of these variables (i.e., SES x Age x Sex; SES x Sex) and were completed for 6 regions of interest (3 hippocampal subdivisions [head, body, and tail]; 3 amygdala subdivisions [accessory basal, central amygdala, basolateral complex, and medial cortical]). We corrected these different steps of analyses for multiple comparisons using False Discovery Rate approaches (Benjamini & Hochberg, 1995). For any models yielding significant interactions, we planned to probe these effects using simple slopes analysis to examine the association between SES and regional volumes at different levels of the moderating variable (e.g., testing the SES-volume relationship separately for males and females, or at ±1 SD of age). Of note, we did not apply ComBat harmonization, a common approach to dealing with neuroimaging data from different sites. This choice was motivated by recent work finding that harmonization methods obscured some of the true biological effect in structural data and structural MRI data "was harmed by harmonization" (Richter et al., 2022). The mixed-effects modeling approach allowed us to statistically control for systematic site-level differences while maximally preserving individual-level variation in brain structure. In our supplement, we also completed analyses: 1) focused on volumes and household income or education (as individual/separate variables); and 2) for smaller parcellations of the hippocampus and amygdala output by Freesurfer.

## RESULTS

*Main Effects of SES on Amygdala and Hippocampal Subregions*
SES was associated with volumes of all amygdala subdivisions and the hippocampal head. For the amygdala, higher SES was related to larger volumes in the superficial cluster (*β=0.07, p<0.001*), basolateral complex (*β=0.05, p=0.014*), and centromedial region (*β=0.05, p=0.028*). Among hippocampal subfields, SES was only related to volumes in the head of the hippocampus (*β=0.07, p=0.002*), with no significant associations observed for the body or tail (p's >.7). After correcting for multiple comparisons, all significant associations remained robust (*superficial: p-fdr=0.005; hippocampal head: p-fdr=0.006; basolateral: p-fdr=0.028; centromedial: p-fdr=0.042*).

*— Insert Figure 3 Approximately Here—*

*Two-Way Interaction: SES x Age*

We examined whether the relationship between SES and regional brain volumes varied as a function of participant age. After correcting for multiple comparisons, no significant SES × Age interactions emerged for any of the amygdala or hippocampal subdivisions (all p-fdr>0.38). The strongest effect was observed in the hippocampal tail (β=0.04, p=0.095), suggesting a trend toward stronger positive SES associations with increasing age, though this did not reach statistical significance. Overall, the association between SES and subcortical volumes did not differ significantly across the age range examined in this sample.

*Two-Way Interaction: SES × Sex*
To determine whether sex moderated the relationship between SES and regional brain volumes, we tested for SES × Sex interactions across all subdivisions. No significant interactions were observed for any amygdala or hippocampal subdivision after FDR correction (all p-fdr>0.70). The largest effect was in the amygdala superficial cluster (β=-0.03, p=0.233), though this was far from significant. These null findings indicate that the association between SES and subcortical volumes was similar for males and females in this sample.

*Three-Way Interaction: SES × Sex × Age*
Finally, we investigated the three-way interaction between SES, sex, and age to test whether sex-specific SES effects varied across development. No three-way interactions reached statistical significance after FDR correction (all p-fdr>0.19). While the hippocampal tail (β=-0.06, p=0.060) and amygdala centromedial region (β=-0.06, p=0.064) showed marginal trends, these did not survive correction for multiple comparisons. Overall, there was no evidence that the relationship between SES and subcortical volumes varied as a function of both sex and age.

*Main Effects of Age and Sex on Subcortical Volumes*
Independent of SES and consistent with prior developmental neuroimaging research, we observed robust main effects of age and sex on subcortical volumes. Age demonstrated significant positive linear associations with volumes across all amygdala and hippocampal subdivisions (*all p-fdr<0.001*), with the strongest effects in the hippocampal body (β=0.12) and tail (β=0.11). Additionally, significant quadratic age effects were observed for all regions (*all p-fdr<0.001*), indicating inverted-U developmental trajectories with volume increases during childhood followed by decreases in later adolescence. Main effects of sex were also pronounced, with females showing larger volumes than males across all three amygdala subdivisions (basolateral: β=0.10, p-fdr<0.001; centromedial: β=0.11, p-fdr<0.001; superficial: β=0.08, p-fdr=0.003) and the hippocampal head (β=0.08, p-fdr=0.002). Sex effects were not significant for the hippocampal body or tail after FDR correction (p-fdr>0.10). Figure 4 displays the pattern of standardized coefficients for all main effects and interactions across the examined subdivisions.

*— Insert Figure 4 Approximately Here—*

**DISCUSSION**

This study investigated associations between SES and volumetric variations in the hippocampus and amygdala. We parcellated the hippocampus and amygdala into smaller subdivisions, aiming to increase neurobiological specificity. With the amygdala, we found that SES was positively associated with volumes across all three subdivisions investigated-- the basolateral complex, centromedial region, and superficial cortical division. For the hippocampus, effects were localized to the head of the hippocampus, with higher SES being associated with

larger volumes in that subdivision. We tested whether these associations varied by participant age or sex through two-way and three-way interaction models. However, we did not find significant moderation effects after correcting for multiple comparisons. Our findings are partially in line with our a priori predictions. We predicted that subdivisions of the hippocampus and amygdala would be related to SES, but we also believed that relations would vary by sex and age; however, interaction analyses did not support these hypotheses in this large sample.

It is critical to note that these SES-related neurobiological differences likely reflect structural and systemic inequalities rather than inherent biological differences between socioeconomic groups. The volumetric variations we observe are likely downstream consequences of differential access to resources, opportunities, and environmental quality-- disparities that are fundamentally rooted in structural inequalities including unequal distribution of wealth, residential segregation by income, educational funding disparities, limited access to quality healthcare, and neighborhood-level disadvantage (Braveman & Gottlieb, 2014; Chetty et al., 2014; Jones et al., 2025; Miller et al., 2025, 2026). These structural barriers constrain opportunities for upward mobility and create conditions where lower-income families face chronic stress, resource scarcity, and reduced access to enriching environments-- factors that may shape the neurodevelopmental patterns we observe.

With the hippocampus, our results are similar to previous projects that have found relations between SES and volumetric differences in the whole hippocampus (Hair et al., 2015; Hanson et al., 2011, 2015; Luby et al., 2013; Noble et al., 2012, 2015). With subfields and specific subdivisions, our results are mostly in line with the small number of publications focused on this question. However, each of these papers varies in the parcellation of the hippocampus and the SES variable examined. Specifically, Merz and colleagues found lower parental education was related to smaller volumes in the dentate gyrus and CA1 subfields of the hippocampus (Merz et al., 2019). The CA1 subfield is squarely in the anterior (head) portion of the hippocampus, while their dentate gyrus subfield can span the head and body of the hippocampus. Botdorf and coworkers found smaller anterior and posterior volumes in the hippocampus with greater area deprivation index, a census-based index of SES (Botdorf et al., 2022). Decker et al. found that lower household income was related to smaller anterior hippocampal volumes (Decker et al., 2020). Examined collectively, there is clear evidence that facets of SES are related to volumes in the head of the hippocampus.

Regarding the amygdala, there is a raft of inconsistencies in past work focused on amygdala volumes, SES, and stress exposure (see (Hanson & Nacewicz, 2021). Multiple groups have reported smaller (whole) amygdala volumes in lower SES youth, but results have not been perfectly uniform. There has been no published work to date focused on SES and amygdala subnuclei. In adult samples exposed to childhood adversity, smaller volumes of basal and accessory basal portions of the amygdala have been reported (Nogovitsyn et al., 2022; Oshri et al., 2019). Oshri and colleagues also found high levels of adversity were related to smaller volumes in central-medial portions of the amygdala. Our findings extend these relations to multiple amygdala subdivisions in a large pediatric sample, demonstrating that SES-related volumetric differences are evident across functionally distinct amygdala regions including the basolateral complex (which serves as the primary sensory input station), the centromedial region (which coordinates autonomic stress responses), and the superficial cortical division (which maintains prominent olfactory and limbic connectivity).

Related to age- and sex-specific effects, we tested whether the association between SES and subcortical volumes varied as a function of participant age, sex, or their combination. Despite a reasonably large sample (N=2,765), we found no significant two-way or three-way interactions

after FDR correction. This suggests that the relations between SES and volumes were similar for males and females and across the age range. We should, however, interpret these null with care, as interaction analyses are often underpowered even in large samples (Baranger et al., 2023; Castillo et al., 2025; McClelland & Judd, 1993). Past work has provided theoretical rationale for expecting age- and sex-specific effects, but subtle moderation effects could exist below our detection threshold. Regarding sex-specific effects, limited work has specifically considered relations between SES and these volumes in males versus females, with inconsistent findings across studies focused on (Kim et al., 2019; Rakesh et al., 2021). Preclinical work examining "sex as a biological variable" suggests that stress exposure often causes more significant neurobiological alterations in males compared to females (Shansky & Woolley, 2016), with chronic stress causing dendritic atrophy and spine elimination in male rodents (McEwen, 2001), but only slight changes in female rats (Galea et al., 1997). With age, Merz et al. found lower SES was significantly associated with smaller amygdala volumes in adolescent participants, but there were no significant associations for younger age participants (Merz et al., 2018). Previously, we also noted that in adversity exposed samples larger volumes in the amygdala were seen for child samples, but small volumes were reported in adults (Hanson & Nacewicz, 2021). Additionally, some evidence suggests timing matters: adversity exposure before 8 years of age was more likely to impact hippocampal volumes in males, while adversity after 9 years of age impacted females more significantly (Teicher et al., 2018). Interactive effects may exist, but our sample design and analytical approaches may have made detection more challenging. Future work, especially longitudinal studies, will be critical to providing clarity about potential age- and sex-specific neurobiological impacts of SES.

Given our results, it will be crucial to think about how these neurobiological alterations may influence future behavior. We did not explore relations between volume and behavior, as specific behavioral measures greatly varied by project. In thinking about relations between SES and the head of the hippocampus, a growing literature suggests that this hippocampal subdivision may be more related to emotion and stress responding (Herman et al., 1998; Kheirbek et al., 2013; Strange et al., 2014). Lesions to this region in rodents (Henke, 1990), as well as variations in human functional connectivity of this area (Gryglewski et al., 2019; Hanson, Gillmore, et al., 2019; Hubachek et al., 2021) have been linked to alterations in stress responding and emotional behavior. Thinking about the amygdala, we find multiple subdivisions are related to SES and these different divisions have been implicated in different aspects of fear responsivity and reward valuation (Phelps & LeDoux, 2005). The basolateral complex serves in sensory gating and associative learning, while the centromedial division relays emotional information to coordinate autonomic responses. Collectively, these alterations could create attentional biases toward negative valenced stimuli, potentially contributing to challenges in emotion regulation (Barry et al., 2022; Hanson, Albert, et al., 2019; Oshri et al., 2019). Preclinical work supports these broad presumptions, as mice with smaller basolateral volumes show significantly greater levels of conditioned freezing compared to those with larger volumes (Yang et al., 2008).These different behavioral processes will be important to interrogate in future work connecting SES and medial temporal lobe neurobiology.

Considering the strengths of our work, the current study benefited from a large sample size (N=2,765), well-validated methods, and an SES composite considering both parental education and income. However, we must consider potential issues with the work. First, our work was cross-sectional in nature, based on a single MRI scan. Volumetric differences could "equalize" over time; this may be particularly true of the hippocampus, where research has demonstrated reversibility in volumetric differences if given a "stress-free" period (Vyas et al., 2002). In future work, we hope to assess other structural and functional properties of the amygdala and hippocampus through the use of longitudinal functional MRI and magnetic resonance

spectroscopy (Nacewicz et al., 2012). Second, automated methods, like FreeSurfer, may be less accurate than hand-delineation of these subdivisions (Hanson et al., 2012). Quantification of these areas by hand is not feasible in such a large sample, but there may be novel automated approaches that more accurately quantify small amygdala and hippocampal subdivisions (DeKraker et al., 2022; Liu et al., 2020). Third, our null findings for age and sex interactions should be interpreted with appropriate caution. Interaction analyses require substantially larger sample sizes than main effect analyses to achieve adequate power (McClelland & Judd, 1993), and subtle moderation effects could exist below our detection threshold despite our large sample. Finally, it will be important to think mechanistically about how lower SES may be impacting neurobiology. Lower SES encompasses a host of challenges likely to impact development, including higher levels of stress, food insecurity, residential instability, community violence, and structural disadvantage (Evans, 2004; Hanson & Hackman, 2012; Masarik & Conger, 2017). This fits with past work finding relations between hippocampal volumes and rich measures of stress exposure (Hanson et al., 2015). Similarly, environmental stimulation is another proximal factor through which SES may impact these volumes, especially the hippocampus. Economically marginalized families may have lower levels of cognitive stimulation in the home, including fewer toys and educational resources (Evans, 2004; Masarik & Conger, 2017). Environmental enrichment and stimulation can impact hippocampal structure, including dendritic branching, neurogenesis, and synaptic density (for review, see (Ohline & Abraham, 2019; Van Praag et al., 2000). Probing different dimensions of experience common to poverty (e.g., stress exposure; environmental stimulation) to understand the patterns reported here will be critical moving forward (Palacios-Barrios et al., 2021; Palacios-Barrios & Hanson, 2019). In our supplement, we examined income and parental education independently and these are important sets of results to also consider. Thoroughly isolating major drivers of neurobiological and behavioral differences could be particularly powerful, especially if this information can be translated into effective interventions to lessen different SES-related disparities.

These limitations notwithstanding, our results provide data about neurobiological alterations seen in relation to SES. Few investigations have examined subdivisions of the amygdala and hippocampus in relation to SES, and our study represents the first to examine these associations with statistical interactions testing age and sex moderation in a large pediatric sample. Such neurobiological differences may connect to SES-gradients in health and well-being. These patterns also suggest how structural inequalities, in education, healthcare access, housing stability, and economic opportunity, may shape brain development, highlighting the critical need for policy interventions that address these systemic barriers. Additional research is needed to clarify the complex relations among early poverty exposure and long-term mental health difficulties; our data are, however, a needed step in the ability to understand the impact of SES on neurobiology critical for emotion, memory, and learning.

**Author contributions**: Jamie Hanson conceived of the project, outlined approaches to statistical analyses, and wrote the majority of the manuscript. Dorthea Adkins wrote portions of the manuscript and constructed statistical models. Brendon Nacewicz guided the team on conceptual and methodological elements of amygdala neurobiology and provided feedback on draft writing. Kelly Barry aided with data cleaning, coding, and organization.

**Funding**: This work was supported by R03HD095048 to JLH.

**Competing Interests**: None to report.

**REFERENCES**

Adkins, D. J., & Hanson, J. L. (2025). Racial and Ethnic Disparities in Brain Age Algorithm Performance: Investigating Bias Across Six Popular Methods. *medRxiv*, 2025–09.

Adler, N. E., & Newman, K. (2002). Socioeconomic disparities in health: Pathways and policies. *Health Affairs*, *21*(2), 60–76.

Adolphs, R. (2010). What does the amygdala contribute to social cognition? *Annals of the New York Academy of Sciences*, *1191*(1), Article 1.

Alexander, L. M., Escalera, J., Ai, L., Andreotti, C., Febre, K., Mangone, A., Vega-Potler, N., Langer, N., Alexander, A., Kovacs, M., & others. (2017). An open resource for transdiagnostic research in pediatric mental health and learning disorders. *Scientific Data*, *4*(1), 1–26.

Amunts, K., Kedo, O., Kindler, M., Pieperhoff, P., Mohlberg, H., Shah, N., Habel, U., Schneider, F., & Zilles, K. (2005). Cytoarchitectonic mapping of the human amygdala, hippocampal region and entorhinal cortex: Intersubject variability and probability maps. *Anatomy and Embryology*, *210*(5), 343–352.

Aoki, C., Romeo, R. D., & Smith, S. S. (2017). Adolescence as a critical period for developmental plasticity. *Brain Research*.

Bacas, E., Kahhalé, I., Raamana, P. R., Pablo, J. B., Anand, A. S., & Hanson, J. L. (2023). Probing multiple algorithms to calculate brain age: Examining reliability, relations with demographics, and predictive power. *Human Brain Mapping*, *44*(9), 3481–3492.

Bale, T. L., & Epperson, C. N. (2017). Sex as a biological variable: Who, what, when, why, and how. *Neuropsychopharmacology*, *42*(2), 386–396.

Baranger, D. A., Finsaas, M. C., Goldstein, B. L., Vize, C. E., Lynam, D. R., & Olino, T. M. (2023). Tutorial: Power analyses for interaction effects in cross-sectional regressions. *Advances in Methods and Practices in Psychological Science*, *6*(3), 25152459231187531.

Barry, K. R., Hanson, J. L., Calma-Birling, D., Lansford, J. E., Bates, J. E., & Dodge, K. A. (2022). Developmental connections between socioeconomic status, self-regulation, and adult externalizing problems. *Developmental Science*, e13260.

Bates, D., Mächler, M., Bolker, B., & Walker, S. (2015). Fitting Linear Mixed-Effects Models Using lme4. *Journal of Statistical Software*, *67*(1), 1–48. https://doi.org/10.18637/jss.v067.i01

Bath, K. G. (2020). Synthesizing views to understand sex differences in response to early life adversity. *Trends in Neurosciences*, *43*(5), 300–310.

Benjamini, Y., & Hochberg, Y. (1995). Controlling the False Discovery Rate: A Practical and Powerful Approach to Multiple Testing. *Journal of the Royal Statistical Society: Series B (Methodological)*, *57*(1), 289–300. https://doi.org/10.1111/j.2517-6161.1995.tb02031.x

Botdorf, M., Dunstan, J., Sorcher, L., Dougherty, L. R., & Riggins, T. (2022). Socioeconomic disadvantage and episodic memory ability in the ABCD sample: Contributions of hippocampal subregion and subfield volumes. *Developmental Cognitive Neuroscience*, *57*, 101138.

Braveman, P., & Gottlieb, L. (2014). The social determinants of health: It's time to consider the causes of the causes. *Public Health Reports*, *129*(1_suppl2), 19–31.

Brody, G. H., Gray, J. C., Yu, T., Barton, A. W., Beach, S. R., Galván, A., MacKillop, J., Windle, M., Chen, E., Miller, G. E., & others. (2017). Protective prevention effects on the association of poverty with brain development. *JAMA Pediatrics*, *171*(1), 46–52.

Brown, S. A., Brumback, T., Tomlinson, K., Cummins, K., Thompson, W. K., Nagel, B. J., Bellis, M. D. D., Hooper, S. R., Clark, D. B., Chung, T., Hasler, B. P., Colrain, I. M., Baker, F. C., Prouty, D., Pfefferbaum, A., Sullivan, E. V., Pohl, K. M., Rohlfing, T., Nichols, B. N., … Tapert, S. F. (2015). The national consortium on alcohol and neuro-development in

adolescence (NCANDA): A multisite study of adolescent development and substance use. *Journal of Studies on Alcohol and Drugs*, *76*(6), 895–908. https://doi.org/10.15288/jsad.2015.76.895

Castillo, A., Miller, J., Vize, C., Baranger, D., & Lynam, D. (2025). When do interaction/moderation effects stabilize in linear regression? *Manuscript Submitted for Publication*.

Chetty, R., Hendren, N., Kline, P., & Saez, E. (2014). Where is the land of opportunity? The geography of intergenerational mobility in the United States. *The Quarterly Journal of Economics*, *129*(4), 1553–1623.

Ciocchi, S., Herry, C., Grenier, F., Wolff, S. B., Letzkus, J. J., Vlachos, I., Ehrlich, I., Sprengel, R., Deisseroth, K., Stadler, M. B., & others. (2010). Encoding of conditioned fear in central amygdala inhibitory circuits. *Nature*, *468*(7321), 277–282.

Dale, A. M., Fischl, B., & Sereno, M. I. (1999). Cortical surface-based analysis: I. Segmentation and surface reconstruction. *Neuroimage*, *9*(2), 179–194.

deCampo, D. M., & Fudge, J. L. (2012). Where and what is the paralaminar nucleus? A review on a unique and frequently overlooked area of the primate amygdala. *Neuroscience & Biobehavioral Reviews*, *36*(1), 520–535.

Decker, A. L., Duncan, K., Finn, A. S., & Mabbott, D. J. (2020). Children's family income is associated with cognitive function and volume of anterior not posterior hippocampus. *Nature Communications*, *11*(1), 1–11.

DeKraker, J., Haast, R. A., Yousif, M. D., Karat, B., Lau, J. C., Köhler, S., & Khan, A. R. (2022). Automated hippocampal unfolding for morphometry and subfield segmentation with HippUnfold. *Elife*, *11*, e77945.

Ellwood-Lowe, M. E., Humphreys, K. L., Ordaz, S. J., Camacho, M. C., Sacchet, M. D., & Gotlib, I. H. (2018). Time-varying effects of income on hippocampal volume trajectories in adolescent girls. *Developmental Cognitive Neuroscience*, *30*, 41–50.

Evans, G. W. (2004). The environment of childhood poverty. *American Psychologist*, *59*(2), Article 2.

Fanselow, M. S., & Dong, H.-W. (2010). Are the dorsal and ventral hippocampus functionally distinct structures? *Neuron*, *65*(1), 7–19.

Fischl, B., Salat, D. H., Busa, E., Albert, M., Dieterich, M., Haselgrove, C., Kouwe, A. V. D., Killiany, R., Kennedy, D., Klaveness, S., Montillo, A., Makris, N., Rosen, B., & Dale, A. M. (2002). Whole brain segmentation: Automated labeling of neuroanatomical structures in the human brain. *Neuron*, *33*(3), 341–355. https://doi.org/10.1016/S0896-6273(02)00569-X

Galea, L., McEwen, B., Tanapat, P., Deak, T., Spencer, R., & Dhabhar, F. (1997). Sex differences in dendritic atrophy of CA3 pyramidal neurons in response to chronic restraint stress. *Neuroscience*, *81*(3), 689–697.

Gaser, C., Dahnke, R., Thompson, P. M., Kurth, F., Luders, E., Initiative, A. D. N., & others. (2024). CAT: a computational anatomy toolbox for the analysis of structural MRI data. *Gigascience*, *13*, giae049.

Genon, S., Bernhardt, B. C., La Joie, R., Amunts, K., & Eickhoff, S. B. (2021). The many dimensions of human hippocampal organization and (dys)function. *Trends in Neurosciences*, *44*(12), 977–989.

Gilmore, A. D., Buser, N. J., & Hanson, J. L. (2021). Variations in structural MRI quality significantly impact commonly used measures of brain anatomy. *Brain Informatics*, *8*(1), 1–15.

Gruber, M. J., & Ranganath, C. (2019). How curiosity enhances hippocampus-dependent memory: The prediction, appraisal, curiosity, and exploration (PACE) framework. *Trends in Cognitive Sciences*, *23*(12), 1014–1025.

Gryglewski, G., Baldinger-Melich, P., Seiger, R., Godbersen, G. M., Michenthaler, P., Klöbl, M., Spurny, B., Kautzky, A., Vanicek, T., Kasper, S., & others. (2019). Structural changes in amygdala nuclei, hippocampal subfields and cortical thickness following electroconvulsive therapy in treatment-resistant depression: Longitudinal analysis. *The British Journal of Psychiatry*, *214*(3), 159–167.

Hair, N. L., Hanson, J. L., Wolfe, B. L., & Pollak, S. D. (2015). Association of child poverty, brain development, and academic achievement. *JAMA Pediatrics*, *169*(9). https://doi.org/10.1001/jamapediatrics.2015.1475

Hair, N. L., Hanson, J. L., Wolfe, B. L., & Pollak, S. D. (2022). Low household income and neurodevelopment from infancy through adolescence. *PloS One*, *17*(1), Article 1.

Hanson, J. L., Adkins, D. J., Bacas, E., & Zhou, P. (2024). Examining the reliability of brain age algorithms under varying degrees of participant motion. *Brain Informatics*, *11*(1), 9.

Hanson, J. L., Albert, W. D., Skinner, A. T., Shen, S. H., Dodge, K. A., & Lansford, J. E. (2019). Resting state coupling between the amygdala and ventromedial prefrontal cortex is related to household income in childhood and indexes future psychological vulnerability to stress. *Development and Psychopathology, 31*(3), 1053–1066.

Hanson, J. L., Chandra, A., Wolfe, B. L., & Pollak, S. D. (2011). Association between Income and the Hippocampus. *PloS One, 6*(5), e18712.

Hanson, J. L., Gillmore, A. D., Yu, T., Holmes, C. J., Hallowell, E. S., Barton, A. W., Beach, S. R., Galván, A., MacKillop, J., Windle, M., & others. (2019). A family focused intervention influences hippocampal-prefrontal connectivity through gains in self-regulation. *Child Development*, *90*(4), 1389–1401.

Hanson, J. L., & Hackman, D. A. (2012). Cognitive neuroscience and SES disparities. *Biological Consequences of Socioeconomic Inequalities, New York, NY: Russell Sage Foundation*.

Hanson, J. L., & Nacewicz, B. M. (2021). Amygdala allostasis and early life adversity: Considering excitotoxicity and inescapability in the sequelae of stress. *Frontiers in Human Neuroscience*.

Hanson, J. L., Nacewicz, B. M., Sutterer, M. J., Cayo, A. A., Schaefer, S. M., Rudolph, K. D., Shirtcliff, E. A., Pollak, S. D., & Davidson, R. J. (2015). Behavioral problems after early life stress: Contributions of the hippocampus and amygdala. *Biological Psychiatry, 77*(4), 314–323.

Hanson, J. L., Suh, J. W., Nacewicz, B. M., Sutterer, M. J., Cayo, A. A., Stodola, D. E., Burghy, C. A., Wang, H., Avants, B. B., Yushkevich, P. A., & others. (2012). Robust automated amygdala segmentation via multi-atlas diffeomorphic registration. *Frontiers in Neuroscience*, *6*, 166.

Hartung, C. M., & Lefler, E. K. (2019). Sex and gender in psychopathology: DSM–5 and beyond. *Psychological Bulletin*, *145*(4), 390.

Hayashi, S., Caron, B. A., Heinsfeld, A. S., Vinci-Booher, S., McPherson, B., Bullock, D. N., Bertò, G., Niso, G., Hanekamp, S., Levitas, D., & others. (2024). Brainlife. Io: A decentralized and open-source cloud platform to support neuroscience research. *Nature Methods, 21*(5), 809–813.

Henke, P. G. (1990). Hippocampal pathway to the amygdala and stress ulcer development. *Brain Research Bulletin*, *25*(5), 691–695.

Herman, J., Dolgas, C., & Carlson, S. (1998). Ventral subiculum regulates hypothalamo–pituitary–adrenocortical and behavioural responses to cognitive stressors. *Neuroscience*, *86*(2), 449–459.

Hubachek, S., Botdorf, M., Riggins, T., Leong, H.-C., Klein, D. N., & Dougherty, L. R. (2021). Hippocampal subregion volume in high-risk offspring is associated with increases in depressive symptoms across the transition to adolescence. *Journal of Affective Disorders*, *281*, 358–366.

Iglesias, J. E., Augustinack, J. C., Nguyen, K., Player, C. M., Player, A., Wright, M., Roy, N., Frosch, M. P., McKee, A. C., Wald, L. L., Fischl, B., & Van Leemput, K. (2015). A computational atlas of the hippocampal formation using ex vivo, ultra-high resolution MRI: Application to adaptive segmentation of in vivo MRI. *Neuroimage*, *115*, 117-137.

Jernigan, T. L., Brown, T. T., Hagler Jr, D. J., Akshoomoff, N., Bartsch, H., Newman, E., Thompson, W. K., Bloss, C. S., Murray, S. S., Schork, N., & others. (2016). The pediatric imaging, neurocognition, and genetics (PING) data repository. *Neuroimage*, *124*, 1149–1154.

Johnson, S. B., Riis, J. L., & Noble, K. G. (2016). State of the art review: Poverty and the developing brain. *Pediatrics*, *137*(4).

Jones, E. J., Miller, P., Gibson-Davis, C. M., Hanson, J. L., & Votruba-Drzal, E. (2025). Family wealth and adolescent physical health. *Health Psychology Review*, *19*(2), 315–343.

Kahhale, I., Buser, N. J., Madan, C. R., & Hanson, J. L. (2023). Quantifying numerical and spatial reliability of hippocampal and amygdala subdivisions in FreeSurfer. *Brain Informatics*, *10*(1), 9.

Kedo, O., Zilles, K., Palomero-Gallagher, N., Schleicher, A., Mohlberg, H., Bludau, S., & Amunts, K. (2018). Receptor-driven, multimodal mapping of the human amygdala. *Brain Structure and Function*, *223*(4), 1637–1666.

Kemppainen, S., Jolkkonen, E., & Pitkänen, A. (2002). Projections from the posterior cortical nucleus of the amygdala to the hippocampal formation and parahippocampal region in rat. *Hippocampus*, *12*(6), 735–755.

Kheirbek, M. A., Drew, L. J., Burghardt, N. S., Costantini, D. O., Tannenholz, L., Ahmari, S. E., Zeng, H., Fenton, A. A., & Hen, R. (2013). Differential control of learning and anxiety along the dorsoventral axis of the dentate gyrus. *Neuron*, *77*(5), 955–968.

Kim, D.-J., Davis, E. P., Sandman, C. A., Glynn, L., Sporns, O., O'Donnell, B. F., & Hetrick, W. P. (2019). Childhood poverty and the organization of structural brain connectome. *NeuroImage*, *184*, 409–416.

Klein-Flügge, M. C., Jensen, D. E., Takagi, Y., Priestley, L., Verhagen, L., Smith, S. M., & Rushworth, M. F. (2022). Relationship between nuclei-specific amygdala connectivity and mental health dimensions in humans. *Nature Human Behaviour*, 1–18.

LeDoux, J. (2007). The amygdala. *Current Biology*, *17*(20), R868–R874.

Lee, S. W., Yoo, J. H., Kim, K. W., Kim, D., Park, H., Choi, J., Teicher, M. H., & Jeong, B. (2018). Hippocampal subfields volume reduction in high schoolers with previous verbal abuse experiences. *Clinical Psychopharmacology and Neuroscience*.

Lenroot, R. K., & Giedd, J. N. (2006). Brain development in children and adolescents: Insights from anatomical magnetic resonance imaging. *Neuroscience & Behavioral Reviews*, *30*(6), 718–729.

Liu, Y., Nacewicz, B. M., Zhao, G., Adluru, N., Kirk, G. R., Ferrazzano, P. A., Styner, M. A., & Alexander, A. L. (2020). A 3D fully convolutional neural network with top-down attention-guided refinement for accurate and robust automatic segmentation of amygdala and its subnuclei. *Frontiers in Neuroscience*, *14*, 260.

Luby, J., Belden, A., Botteron, K., Marrus, N., Harms, M. P., Babb, C., Nishino, T., & Barch, D. (2013). The effects of poverty on childhood brain development: The mediating effect of caregiving and stressful life events. *JAMA Pediatrics*, *167*(12), 1135–1142.

Majak, K., & Pitkänen, A. (2003). Projections from the periamygdaloid cortex to the amygdaloid complex, the hippocampal formation, and the parahippocampal region: A PHA-L study in the rat. *Hippocampus*, *13*(8), 922–942.

Malhi, G. S., Das, P., Outhred, T., Irwin, L., Gessler, D., Bwabi, Z., Bryant, R., & Mannie, Z. (2019). The effects of childhood trauma on adolescent hippocampal subfields. *Australian & New Zealand Journal of Psychiatry*, *53*(5), 447–457.

Margolis, A. E., Cohen, J. W., Ramphal, B., Thomas, L., Rauh, V., Herbstman, J., & Pagliaccio, D. (2022). Prenatal Exposure to Air Pollution and Early Life Stress Effects on Hippocampal Subregional Volumes and Associations with Visual-Spatial Reasoning. *Biological Psychiatry Global Open Science*.

Masarik, A. S., & Conger, R. D. (2017). Stress and child development: A review of the Family Stress Model. *Current Opinion in Psychology*, *13*, 85–90.

McClelland, G. H., & Judd, C. M. (1993). Statistical difficulties of detecting interactions and moderator effects. *Psychological Bulletin*, *114*(2), 376.

McDermott, C. L., Seidlitz, J., Nadig, A., Liu, S., Clasen, L. S., Blumenthal, J. D., Reardon, P. K., Lalonde, F., Greenstein, D., Patel, R., & others. (2019). Longitudinally mapping childhood socioeconomic status associations with cortical and subcortical morphology. *Journal of Neuroscience*, *39*(8), 1365–1373.

McEwen, B. S. (2001). Plasticity of the hippocampus: Adaptation to chronic stress and allostatic load. *Annals of the New York Academy of Sciences*, *933*(1), 265–277.

McEwen, B. S. (2002). Sex, stress and the hippocampus: Allostasis, allostatic load and the aging process. *Neurobiology of Aging*, *23*(5), Article 5.

McEwen, B. S. (2007). Physiology and neurobiology of stress and adaptation: Central role of the brain. *Physiological Reviews*, *87*(3), 873–904.

McEwen, B. S. (2020). Hormones and behavior and the integration of brain-body science. *Hormones and Behavior*, *119*, 104619.

Merz, E. C., Desai, P. M., Maskus, E. A., Melvin, S. A., Rehman, R., Torres, S. D., Meyer, J., He, X., & Noble, K. G. (2019). Socioeconomic disparities in chronic physiologic stress are associated with brain structure in children. *Biological Psychiatry*, *86*(12), 921–929.

Merz, E. C., Tottenham, N., & Noble, K. G. (2018). Socioeconomic Status, Amygdala Volume, and Internalizing Symptoms in Children and Adolescents. *Journal of Clinical Child & Adolescent Psychology*, *47*(2), Article 2. https://doi.org/10.1080/15374416.2017.1326122

Miller, P., Blatt, L., Hunter-Rue, D., Barry, K. R., Jamal-Orozco, N., Hanson, J. L., & Votruba-Drzal, E. (2025). Economic hardship and adolescent behavioral outcomes: Within-and between-family associations. *Development and Psychopathology*, *37*(1), 107–124.

Miller, P., Podvysotska, T., Jones, E., Hanson, J. L., & Votruba-Drzal, E. (2026). How Family Assets and Debts Relate to Children's Achievement and Behavior Problems Across Development. *Social Development*, *35*(1), e70033.

Murty, V. P., Calabro, F., & Luna, B. (2016). The role of experience in adolescent cognitive development: Integration of executive, memory, and mesolimbic systems. *Neuroscience & Biobehavioral Reviews*, *70*, 46–58.

Nacewicz, B. M., Angelos, L., Dalton, K. M., Fischer, R., Anderle, M. J., Alexander, A. L., & Davidson, R. J. (2012). Reliable non-invasive measurement of human neurochemistry using proton spectroscopy with an anatomically defined amygdala-specific voxel. *Neuroimage*, *59*(3), 2548–2559.

Noble, K. G., & Giebler, M. A. (2020). The neuroscience of socioeconomic inequality. *Current Opinion in Behavioral Sciences*, *36*, 23–28.

Noble, K. G., Houston, S. M., Brito, N. H., Bartsch, H., Kan, E., Kuperman, J. M., Akshoomoff, N., Amaral, D. G., Bloss, C. S., Libiger, O., Schork, N. J., Murray, S. S., Casey, B. J., Chang, L., Ernst, T. M., Frazier, J. A., Gruen, J. R., Kennedy, D. N., Zijl, P. V., … Sowell, E. R. (2015). Family income, parental education and brain structure in children and adolescents. *Nature Neuroscience*, *18*, 773–778.

Noble, K. G., Houston, S. M., Kan, E., & Sowell, E. R. (2012). Neural correlates of socioeconomic status in the developing human brain. *Developmental Science*, *15*(4), Article 4.

Nogovitsyn, N., Addington, J., Souza, R., Placsko, T. J., Stowkowy, J., Wang, J., Goldstein, B. I., Bray, S., Lebel, C., Taylor, V. H., & others. (2022). Childhood trauma and amygdala nuclei volumes in youth at risk for mental illness. *Psychological Medicine*, *52*(6), 1192–1199.

Ohline, S., & Abraham, W. (2019). Environmental enrichment effects on synaptic and cellular physiology of hippocampal neurons. *Neuropharmacology*, *145*, 3–12.

Oshri, A., Gray, J. C., Owens, M. M., Liu, S., Duprey, E. B., Sweet, L. H., & MacKillop, J. (2019). Adverse childhood experiences and amygdalar reduction: High-resolution segmentation reveals associations with subnuclei and psychiatric outcomes. *Child Maltreatment*, *24*(4), 400–410.

Palacios-Barrios, E. E., & Hanson, J. L. (2019). Poverty and self-regulation: Connecting psychosocial processes, neurobiology, and the risk for psychopathology. *Comprehensive Psychiatry*, *90*, 52–64.

Palacios-Barrios, E. E., Hanson, J. L., Barry, K. R., Albert, W. D., White, S. F., Skinner, A. T., Dodge, K. A., & Lansford, J. E. (2021). Lower neural value signaling in the prefrontal cortex is related to childhood family income and depressive symptomatology during adolescence. *Developmental Cognitive Neuroscience*, *48*, 100920.

Parker, R. N., & Fenwick, R. (1983). The Pareto curve and its utility for open-ended income distributions in survey research. *Social Forces*, *61*(3), 872–885.

Paton, J. J., Belova, M. A., Morrison, S. E., & Salzman, C. D. (2006). The primate amygdala represents the positive and negative value of visual stimuli during learning. *Nature*, *439*(7078), 865–870.

Peverill, M., Dirks, M. A., Narvaja, T., Herts, K. L., Comer, J. S., & McLaughlin, K. A. (2021). Socioeconomic status and child psychopathology in the United States: A meta-analysis of population-based studies. *Clinical Psychology Review*, *83*, 101933. https://doi.org/10.1016/j.cpr.2020.101933

Phelps, E. A., & LeDoux, J. E. (2005). Contributions of the amygdala to emotion processing: From animal models to human behavior. *Neuron*, *48*(2), 175–187.

Piotrowska, P. J., Stride, C. B., Croft, S. E., & Rowe, R. (2015). Socioeconomic status and antisocial behaviour among children and adolescents: A systematic review and meta-analysis. *Clinical Psychology Review*, *35*, 47–55.

Poppenk, J., & Moscovitch, M. (2011). A hippocampal marker of recollection memory ability among healthy young adults: Contributions of posterior and anterior segments. *Neuron*, *72*(6), 931–937.

Raffington, L., Czamara, D., Mohn, J. J., Falck, J., Schmoll, V., Heim, C., Binder, E. B., & Shing, Y. L. (2019). Stable longitudinal associations of family income with children's hippocampal volume and memory persist after controlling for polygenic scores of educational attainment. *Developmental Cognitive Neuroscience*, *40*, 100720.

Rakesh, D., Seguin, C., Zalesky, A., Cropley, V., & Whittle, S. (2021). Associations between neighborhood disadvantage, resting-state functional connectivity, and behavior in the adolescent brain cognitive development study: The moderating role of positive family and school environments. *Biological Psychiatry: Cognitive Neuroscience and Neuroimaging*, *6*(9), 877–886.

Richter, S., Winzeck, S., Correia, M. M., Kornaropoulos, E. N., Manktelow, A., Outtrim, J., Chatfield, D., Posti, J. P., Tenovuo, O., Williams, G. B., & others. (2022). Validation of cross-sectional and longitudinal ComBat harmonization methods for magnetic resonance imaging data on a travelling subject cohort. *Neuroimage: Reports*, *2*(4), 100136.

Sah, P., Faber, E. L., Lopez de Armentia, M., & Power, J. (2003). The amygdaloid complex: Anatomy and physiology. *Physiological Reviews*, *83*(3), 803–834.

Saygin, Z. M., Kliemann, D., Iglesias, J. E., van der Kouwe, A. J., Boyd, E., Reuter, M., Stevens, A., Van Leemput, K., McKee, A., Frosch, M. P., & others. (2017). High-resolution

magnetic resonance imaging reveals nuclei of the human amygdala: Manual segmentation to automatic atlas. *Neuroimage*, *155*, 370–382.

Shansky, R. M., & Woolley, C. S. (2016). Considering sex as a biological variable will be valuable for neuroscience research. *Journal of Neuroscience*, *36*(47), 11817–11822.

Shohamy, D., & Turk-Browne, N. B. (2013). Mechanisms for widespread hippocampal involvement in cognition. *Journal of Experimental Psychology: General*, *142*(4), Article 4.

Somerville, L. H., Bookheimer, S. Y., Buckner, R. L., Burgess, G. C., Curtiss, S. W., Dapretto, M., Elam, J. S., Gaffrey, M. S., Harms, M. P., Hodge, C., & others. (2018). The Lifespan Human Connectome Project in Development: A large-scale study of brain connectivity development in 5–21 year olds. *Neuroimage*, *183*, 456–468.

Stevens, J. S., & Hamann, S. (2012). Sex differences in brain activation to emotional stimuli: A meta-analysis of neuroimaging studies. *Neuropsychologia*, *50*(7), 1578–1593.

Strange, B. A., Witter, M. P., Lein, E. S., & Moser, E. I. (2014). Functional organization of the hippocampal longitudinal axis. *Nature Reviews Neuroscience*, *15*(10), 655–669.

Swanson, L. W., & Petrovich, G. D. (1998). What is the amygdala? *Trends in Neurosciences*, *21*(8), 323–331.

Teicher, M. H., Anderson, C. M., Ohashi, K., Khan, A., McGreenery, C. E., Bolger, E. A., Rohan, M. L., & Vitaliano, G. D. (2018). Differential effects of childhood neglect and abuse during sensitive exposure periods on male and female hippocampus. *Neuroimage*, *169*, 443–452.

Teicher, M. H., Anderson, C. M., & Polcari, A. (2012). Childhood maltreatment is associated with reduced volume in the hippocampal subfields CA3, dentate gyrus, and subiculum. *Proceedings of the National Academy of Sciences*, *109*(9), E563–E572.

van Eijk, L., Hansell, N. K., Strike, L. T., Couvy-Duchesne, B., de Zubicaray, G. I., Thompson, P. M., McMahon, K. L., Zietsch, B. P., & Wright, M. J. (2020). Region-specific sex differences in the hippocampus. *Neuroimage*, *215*, 116781.

Van Praag, H., Kempermann, G., & Gage, F. H. (2000). Neural consequences of enviromental enrichment. *Nature Reviews Neuroscience*, *1*(3), 191–198.

Vogel, J. W., La Joie, R., Grothe, M. J., Diaz-Papkovich, A., Doyle, A., Vachon-Presseau, E., Lepage, C., Vos de Wael, R., Thomas, R. A., & Iturria-Medina, Y. (2020). A molecular gradient along the longitudinal axis of the human hippocampus informs large-scale behavioral systems. *Nature Communications*, *11*(1), 1–17.

Vyas, A., Mitra, R., Rao, B. S., & Chattarji, S. (2002). Chronic stress induces contrasting patterns of dendritic remodeling in hippocampal and amygdaloid neurons. *Journal of Neuroscience*, *22*(15), 6810–6818.

Weissman, D. G., Hatzenbuehler, M. L., Cikara, M., Barch, D. M., & McLaughlin, K. A. (2023). State-level macro-economic factors moderate the association of low income with brain structure and mental health in US children. *Nature Communications*, *14*(1), 2085.

Wisse, L. E., Chételat, G., Daugherty, A. M., de Flores, R., la Joie, R., Mueller, S. G., Stark, C. E., Wang, L., Yushkevich, P. A., Berron, D., & others. (2021). Hippocampal subfield volumetry from structural isotropic 1 mm3 MRI scans: A note of caution. *Human Brain Mapping*, *42*(2), 539–550.

Yagi, S., & Galea, L. A. (2019). Sex differences in hippocampal cognition and neurogenesis. *Neuropsychopharmacology*, *44*(1), 200–213.

Yang, R. J., Mozhui, K., Karlsson, R.-M., Cameron, H. A., Williams, R. W., & Holmes, A. (2008). Variation in mouse basolateral amygdala volume is associated with differences in stress reactivity and fear learning. *Neuropsychopharmacology*, *33*(11), 2595–2604.

Yu, Q., Daugherty, A. M., Anderson, D. M., Nishimura, M., Brush, D., Hardwick, A., Lacey, W., Raz, S., & Ofen, N. (2018). Socioeconomic status and hippocampal volume in children and young adults. *Developmental Science*, *21*(3), e12561.

Yuan, M., Rubin-Falcone, H., Lin, X., Rizk, M. M., Miller, J. M., Sublette, M. E., Oquendo, M. A., Burke, A., Ogden, R. T., & Mann, J. J. (2020). Smaller left hippocampal subfield CA1 volume is associated with reported childhood physical and/or sexual abuse in major depression: A pilot study. *Journal of Affective Disorders*, *272*, 348–354.

Table 1.

| Characteristic | N = 2765 |
| --- | --- |
| **Age (years), M (SD)** | 11.91 (3.45) |
| **Sex, n (%)** | |
| F | 1,228 (44.4%) |
| M | 1,537 (55.6%) |
| **Household Income (USD), M (SD)** | 123,641 (99,112) |
| **Maximum Parental Education (years), M (SD)** | 17.49 (2.95) |

Demographic information for the full sample, including the means and standard deviations for age (in years), income (in US dollars, USD), and highest parental education (in years), as well as the distribution of sex in this sample.

Figure 1.

 Stacked histogram depicting the number of participants at each age, for each study, between the ages of 5 and 18. The top set of bins represents the numbers of participants in the HBN study at each age (colored in blue); stacked below are the bins for the HCPD (purple), NCANDA (green), and HBN (orange) studies, respectively. The dashed and dotted lines display the mean ages for each study (with each color corresponding to a different study; blue = HBN; purple = HCPD; green = NCANDA; orange = PING).

Figure 2.

Depiction of our amygdala and hippocampal regions of interest. The left side of this figure shows our 3 amygdala subnuclei groups overlaid on an average T1-weighted scan. The right side of this figure shows our 3 hippocampal subdivisions also overlaid on an average T1-weighted scan.

Figure 3.

Scatterplots depicting main effects of socioeconomic status on amygdala and hippocampal subdivisions. Each panel displays the relationship between standardized SES composite scores (x-axis) and regional brain volumes in mm³ (y-axis) for six subcortical subdivisions: amygdala basolateral complex, centromedial division, and superficial cortical division (top row), and hippocampal head, body, and tail (bottom row). Orange and blue regression lines (amygdala = orange; blue = hippocampus) indicate FDR-significant positive associations between SES and regional volumes (q < 0.05). Gray lines indicate non-significant associations. All models controlled for age, sex, estimated total intracranial volume, image quality, and study site.

Figure 4.

Comprehensive effects heatmap of socioeconomic status, sex, and age associations with subcortical volumes. Heatmap displays standardized regression coefficients (β) for main effects (SES, Sex, Age, Age²), two-way interactions (SES×Sex, SES×Age, Sex×Age), and three-way interaction (SES×Sex×Age) across amygdala subdivisions (top three rows: basolateral complex, centromedial division, superficial cortical division) and hippocampal subdivisions (bottom three rows: head, body, tail). Red shading indicates positive associations; blue shading indicates negative associations; white indicates no association. Double asterisks (**) denote effects surviving False Discovery Rate correction (q < 0.05); single asterisks (*) indicate uncorrected significance (p < 0.05). All models controlled for estimated total intracranial volume (eTIV), image quality (CAT12 rating), and study site (modeled as a random effect).