Spatial Transcriptomic Cell-type Deconvolution Using Graph Neural Networks

Yawei Li[1,2], Yuan Luo[1,2,*]

[1] *Department of Preventive Medicine, Northwestern University, Feinberg School of Medicine, Chicago, IL 60611, USA*

[2] *Center for Collaborative AI in Healthcare, Northwestern University, Feinberg School of Medicine, Chicago, IL 60611, USA*

[*] Corresponding author. Email: yuan.luo@northwestern.edu (Yuan Luo)

Abstract

Spatially resolved transcriptomics performs high-throughput measurement of transcriptomes while preserving spatial information about the cellular organizations. However, many spatially resolved transcriptomic technologies can only distinguish spots consisting of a mixture of cells instead of working at single-cell resolution. Here, we present STdGCN, a graph neural network model designed for cell type deconvolution of spatial transcriptomic (ST) data that can leverage abundant single-cell RNA sequencing (scRNA-seq) data as reference. STdGCN is the first model incorporating the expression profiles from single cell data as well as the spatial localization information from the ST data for cell type deconvolution. Extensive benchmarking experiments on multiple ST datasets showed that STdGCN outperformed 14 published state-of-the-art models. Applied to a human breast cancer Visium dataset, STdGCN discerned spatial

23 distributions between stroma, lymphocytes and cancer cells for tumor microenvironment

24 dissection. In a human heart ST dataset, STdGCN detected the changes of potential endothelial-

25 cardiomyocyte communications during tissue development.

Introduction

The development of spatially resolved transcriptomic technologies has improved our

understanding of the spatial organization and gene expression heterogeneity within tissues [1-4].

Spatial transcriptomics (ST) technologies enable the measurement of transcriptomes while

maintaining the spatial context of the tissue, offering valuable insights into disease pathology

research [5-7]. However, ST technologies still face certain limitations due to the trade-off between

cell resolution and throughput. For instance, many fluorescence in situ hybridization (FISH)

based technologies are unable to capture the full scope of the transcriptome within a single cell

[8]. This poses computational challenges when attempting to infer the missing gene expression

information from FISH data. In addition, many sequencing-based methods used in ST often

lack sufficient resolution at the single cell level, raising computational challenges for

deconvolving cell types detected within the same spatial spot/location. In contrast to ST

technologies, single-cell RNA sequencing (scRNA-seq) is capable of detecting many thousands

of gene expressions within individual cells. As a result, scRNA-seq can be effectively leveraged

as a complementary tool to address certain limitations of ST technologies. For example,

restoring cell type decomposition within a spot/location in ST data bears similarities to using

scRNA-seq to deconvolve bulk RNA-seq data. Indeed, several models have been developed for

bulk RNA-seq data deconvolution [9-14], and in theory, these models could be adapted to handle

ST data. However, the cell numbers within a spot/location in ST data are considerably lower

than those in bulk RNA-seq data. Consequently, applying a bulk RNA-seq deconvolution

method to such a small sample size usually introduces noise from unrelated cell types, and has

been demonstrated by several benchmark analyses [15-17].

In recent years, several models have been developed for deconvolving ST data, employing

either statistics-based or machine learning-based algorithms [15-28]. Statistics-based algorithms

typically assume a distribution , usually Poisson [18] or negative binomial [15, 17, 22, 26], for the gene

transcript count. These methods first employ the scRNA-seq data to calculate the distribution

coefficients for genes in each cell type. Cell type proportions are then inferred using maximum-

likelihood estimation or maximum a posteriori estimation. On the other hand, machine learning

based algorithms usually integrate the predicted ST data with scRNA-seq data to directly

predict cell type proportions through neural network systems. These models eliminate the need

for an additional step to learn cell type gene signatures from scRNA-seq data [20, 21]. However, a

common limitation of these models is that they deconvolve the spots in a given ST dataset

independently, without considering the neighborhood connections between spots. Indeed, it has

been demonstrated that ST data can be divided into distinct spatial domains [29, 30]. By

incorporating the spatial coordinate information between spots, the ST cell type decomposition

learned from scRNA-seq can calibrated, leading to increased predictive accuracy.

In this study, we present the Spatial Transcriptomics deconvolution using Graph

Convolutional Networks (STdGCN) as a novel approach for ST data cell-type deconvolution.

Our model leverages graph convolutional networks (GCN), a widely used graph-based deep

learning model. Notably, STdGCN integrates the expression profiles from scRNA-seq data with

spatial localization information from the ST data for cell type deconvolution. By integrating

expression and spatial information, STdGCN achieves accurate predictions of cell type

proportions in ST data. To evaluate the performance of STdGCN, we conducted benchmarking

against 14 state-of-the-art models. The results demonstrate that STdGCN consistently

outperforms the benchmarked models across various ST platforms. This highlights the

superiority of STdGCN in cell-type deconvolution for ST data.

Results

STdGCN description. STdGCN is designed for cell type deconvolution in ST data by using

scRNA-seq data as a reference (**Fig. 1**). The underlying hypothesis is that both ST data and

scRNA-seq data share common cell types, and their cell type gene transcript signatures exhibit

similarities. In short, the normalized gene expression of a spot can be considered a combination

of different cell types with varying proportions. The initial step of STdGCN involves

identifying cell-type marker genes and generating pseudo-spots using the scRNA-seq data (**Fig.**

1A). Subsequently, it builds two link graphs to establish the GCN pipeline (**Fig. 1B~1C**). The

first link graph, known as the expression graph, is a hybrid graph comprising three sub-graphs,

a pseudo-spots internal graph, a real-spots internal graph, and a real-to-pseudo-spots graph.

Each subgraph is constructed based on mutual nearest neighbors (MNN) using the expressed

similarity between spots. The second link graph, the spatial graph, is built based on the

Euclidean distance between real-spots in ST data. During the execution of STdGCN, the input

feature matrix is forward propagated through the expression GCN layers and the spatial GCN

layers, respectively. The outputs of the two GCN layers, namely Exp-feature and Spa-feature,

are then concatenated column-wise into a single matrix. This concatenated matrix is

subsequently fed into fully-connected layers to predict the cell type proportions for each spot.

To train the model, we divide the pseudo-spots into a training dataset and a validation dataset.

Only pseudo-spots in the training dataset are used for back propagation, whereas the validation

dataset serves the purpose of early-stopping. Using this approach, the cell type proportions of

real-spots can also be updated through the GCN pipeline, benefiting the learning of the pseudo-

spots (**Fig. 1D**). The details are described in **Methods**.

Benchmarks of STdGCN in different datasets. To assess the performance of STdGCN, we

conducted a comprehensive comparison with 14 state-of-the-art deconvolution models,

including stereoscope [15], RCTD [18], SPOTlight [16], SpatialDWLS [19], Cell2location [26], DSTG [20],

CellDART [21], DestVI [22], STdeconvolve [27], STRIDE [23], Tangram [24], BayesPrism [25], AdRoit [17]

and SpatialDecon [28]. We selected four different ST platforms with single cell level cell type

annotations: seqFISH [31], seqFISH+ [30], MERFISH [32] and slide-seq [33]. Synthetic multi-cellular

ST datasets were generated using these platforms, allowing us to evaluate the performance of

the models under controlled conditions (**Fig. 2**, **Methods**). To quantitatively assess accuracy of

the 15 models, we employed two metrics: Jensen-Shannon divergence (JSD) and root-mean-

square error (RMSE). Based on the two values, we ranked the models accordingly. Since each

dataset consisted of multiple slices, the performance of a model for a given dataset was

determined by calculating the mean rank across all slices within that dataset.

Table 1 presents the performance of the 15 models. The rankings showed that STdGCN

achieved the lowest average JSD and RMSE across all four ST datasets. It tied for first place

with RCTD in JSD of the MERFISH dataset. Specifically, in the seqFISH and seqFISH+

datasets, STdGCN achieved the top ranking in both JSD and RMSE across all slices. In the case

of the Slide-seq dataset, STdGCN secured the first rank among four out of the six slices.

Furthermore, in the MERFISH dataset, STdGCN demonstrated the best performance among

66% of the slices in RMSE and 42% of the slices in JSD, respectively.

To further evaluate the performance of the models, we investigated their effectiveness in

spots with different cell numbers. We divided the spots into two categories: a smaller group

consisting of spots with five or fewer cells and a larger group comprising spots with more than

five cells. We then benchmarked the models in each of the two groups across all datasets. For

most models, the average JSD and RMSE of the larger group were lower than those of the

smaller group (P-value < 0.05, Wilcoxon test) (**Fig. 3A~3D**, **Extended Data Fig. 1**), indicating

that spots with smaller cell number pose greater challenges for accurate deconvolution. Despite

this difficulty, when comparing the relative performance between two groups, the average ranks

of most methods remained consistent (**Table S1**). Remarkably, STdGCN consistently

outperformed other models in both groups, except for the smaller group of the MERFISH

dataset in terms of JSD metrics.

Cell-cell communication plays a vital role in multicellular organisms, and it is influenced

by physical location within the cellular microenvironment. Spatially adjacent cells are more

likely to interact with each other. Thus, accurate deconvolution of cell types enables the

identification of co-localization and spatial interactions. In light of this, we scrutinized the

performance of the benchmarked models when dealing with different number of cell types

within a single spot. We divided the spots into four groups based on the number of cell types:

singleton (one cell type), doubleton (two cell types), tripleton (three cell types), and multiton

(four or more cell types). Of the four groups, STdGCN consistently demonstrated strong

performance (**Fig. 3E~3H**, **Extended Data Fig. 2**, **Table S2**), and in most circumstances,

STdGCN achieved the lowest JSD and RMSE scores.

Lastly, we assessed the models' capability for predicting diverse cell types. For this

analysis, we employed a binary Jensen-Shannon divergence (JSD) to measure the discrepancy

between the predicted proportions and the actual proportions by summarizing the proportions

of the remaining cell types within each spot (**Methods**). To measure the RMSE of a cell type,

we exclusively calculated the RMSE between predicted proportions and the real proportions

across all spots corresponding to that particular cell type within a given slice. In terms of the

average ranks of the 15 models, STdGCN outperformed other models for most cell types (**Fig.**

3I~3L, **Extended Data Fig. 3**). It is noteworthy that based on the average ranks of all cell types,

STdGCN demonstrated a greater specialization in predicting cell types with higher frequencies

(**Table S3~S4**).

Developing human heart spatial transcriptomic dataset. In order to show the utility of

STdGCN, we first employed STdGCN on a developing human heart [34] to assess its performance

on a real-word multi-cellular ST dataset. This study used a combination of spatial

transcriptomics [35, 36], scRNA-seq and *in situ* sequencing (ISS), to capture transcriptomic profiles

in human embryonic heart tissues (**Fig. 4A~4D**). Leveraging the scRNA-seq data as the

reference, we employed STdGCN to infer the cell-type composition within the spatial

transcriptomics dataset. Remarkably, the cell type distribution predicted by STdGCN exhibited

a high degree of concordance with the ISS dataset (**Fig. 4A, Extended Data Fig. 4**). STdGCN

accurately identified the spatial localization patterns of various cell types within the heart tissue:

epicardial cells were predominantly located on the outer surface of the heart tissue, atrial

cardiomyocytes were primarily situated within the atrium, and smooth muscle cells were

located in the aorta and pulmonary artery area (**Fig. 4E~4G**). These predicted cell types were

consistent with existing medical knowledge [34], providing compelling evidence for the efficacy

of STdGCN in capturing the spatial organization of cell types within complex tissue

environments.

In a previous review study [37], various models were compared in their ability to deconvolve

the aforementioned dataset. The finding revealed that most models struggled to distinguish

between *Myoz2*-enriched cardiomyocytes and ventricular cardiomyocytes (Figure 3 in Chen et

al. [37]) (**Fig. 4H~4I**). It has been established that that the *Myoz2*-enriched cardiomyocytes are a

subpopulation of cardiomyocytes in the healthy heart, characterized by the expression of

MYOZ2 [34, 38]. Notably, both the UMAP plots of the ISS dataset and the scRNA-seq dataset

demonstrated that the expression level of *Myoz2*-enriched cardiomyocytes closely resembled

that of ventricular cardiomyocytes and atrial cardiomyocytes (**Fig. 4B and 4D**).

We investigated their respective cell type marker genes (**Methods**) and found that 41.6%

of the Myoz2-enriched cardiomyocyte marker genes were also shared with atrial cardiomyocyte

marker genes, while 100% of the Myoz2-enriched cardiomyocyte marker genes overlapped

with ventricular cardiomyocyte marker genes (**Fig. 4J**). Notably, *MYOZ2*, which has been

identified as the most significant marker gene for Myoz2-enriched cardiomyocytes (average

log-fold change = 1.81, expressed in 99.0% *Myoz2*-enriched ventricular cardiomyocytes, FDR

$\approx 1.08 \times 10^{-89}$) [34, 38], also serves as a marker gene for ventricular cardiomyocytes (average log-

fold change = 1.30, expressed in 92.7% ventricular cardiomyocytes, FDR ≈ 0) (**Methods**). The

considerable overlap of marker genes between ventricular cardiomyocyte and *Myoz2*-enriched

cardiomyocyte elucidates why the majority of models fail to effectively differentiate between

the two cell types. Despite the challenge, we compared STdGCN with eight previously well-

performed models. The results showcased that STdGCN outperformed other models,

successfully distinguish between the two cell types (**Fig. 4H~4I, Extended Data Fig. 5**).

Studies have emphasized the crucial role of communication between endothelial cells and

cardiomyocytes in cardiac development. Endothelial cells not only facilitate the transmission

of oxygenated blood supply to cardiomyocytes, but also provide local protective signals that

promote cardiomyocyte organization and survival [39]. To investigate the extent of endothelial-

cardiomyocyte communication within the analyzed dataset, we explore the cell type co-

localization score (**Methods**) across three developmental stages of heart tissue sections

(PCW4.5-5, PCW6.5, PCW9) (**Fig. 4K**). The heatmap shows a higher co-localization score

between endothelial cells (including "Endothelium / pericytes / adventitia" and "Capillary

endothelium" in the dataset) and cardiomyocytes (including "Myoz2-enriched

cardiomyocytes", "ventricular cardiomyocytes", "atrial cardiomyocytes" in the dataset)

compared to the overall cell type level. This finding aligns with previous research and

underscores the significance of endothelial-cardiomyocyte interactions in the context of this

dataset.

Developing human breast cancer transcriptomic dataset. To assess the versatility of

STdGCN, we then applied it to investigate the spatial organization of a human breast cancer ST

dataset [40]. This dataset involved the analysis of breast tumor samples using both scRNA-seq

(**Fig. 5A~5B**) and 10X Genomics Visium ST sequencing. By utilizing the scRNA-seq data as a

reference, we obtained the cell type proportions corresponding to each spot in the 10X

Genomics Visium dataset (**Fig. S1~S6**).

By using STdGCN, we were able to quantitatively dissect the tumor microenvironment

(TME) for these tissues in high resolution. For example, previous studies have highlighted the

significance of tumor-infiltrating lymphocytes (TILs) as a strong prognostic marker for patient

survival. In the case of locally developed breast cancer treated with neoadjuvant chemotherapy,

the presence of TILs is a predictor of the response [41-43]. Leveraging the deconvolution results

predicted by STdGCN, we compared the cell type co-localization score (**Methods**) between

estrogen receptor-positive (ER+) samples and triple-negative breast cancer (TNBC) samples.

The co-localization heatmap revealed higher TILs in ER+ samples than TNBC samples,

suggesting a more favorable prognosis for ER+ samples (**Fig. 5C**). Additionally, the

interactions between stromal cells, cancer cells and lymphocytes have been reported to be

important for maintaining the TME [44]. Cancer cells that interact with stromal cells can create

favorable physical or molecular signaling environments that promote tumor progression [45],

while certain stromal cells, such as cancer-associated fibroblasts (CAFs) can recruit immune

cells to the tumor tissues and influence their behavior towards cancer cells [46]. Through STdGCN,

we were able to analyze cell-cell interactions in spatial locations. For instance, we compared

the TME of an ER+ patient (CID4535) with that of a TNBC patient (CID4465) (**Fig. 5D~5L**).

Despite similar overall proportions of cell types between the two patients (**Fig. 5D**), the spatial

structure and co-localization heatmap revealed substantial differences in their respective (**Fig.**

5E~5L), indicating distinct microenvironmental characteristics.

Discussion

ST profiles have significantly enhanced our understanding of tissue architecture at the cellular

level. However, many existing ST profiling technologies lack sufficient single cell resolution.

To this end, we introduce STdGCN, a graph-based deep learning framework that leverages cell

type profiles learned from single-cell RNA-seq to effectively deconvolve the cell type mixtures

in ST data. In order to evaluate the performance of STdGCN, we conducted a comprehensive

benchmarking study involving four ST datasets and compared its performance against 14 state-

of-the-art ST deconvolution methods [15-27]. The models were ranked based on their performance

metrics. Across all four benchmarked datasets, STdGCN demonstrated the most superior

performance in terms of both JSD and RMSE (**Table 1**).

To improve the performance of cell type deconvolution, distinct designs were adapted

based on the characteristics of ST datasets. First, STdGCN stands out from previous models by

incorporating spatial structure information from ST data and expression profiles from scRNA-

seq for cell type deconvolution. This innovative approach capitalizes on the inherent similarity

between spots within a spatial domain, allowing for precise calibration of predictions based on

the expression profiles derived from scRNA-seq data. Next, to maximize the heterogeneity of

the pseudo-spot pool, we developed a sampling-by-cell-type method to simulate pseudo-spots.

This approach does not rely on cell-type distributions from scRNA-seq by assigning equal

weights for each cell type. This effectively addresses the challenge of rare cell types being

unlikely to be randomly selected using traditional methods. In addition, we imposed a

maximum limit on the number of cell types within a single pseudo-spot to accurately mimic the

characteristics observed in real-word ST data.

The modular design of STdGCN provides users with a high degree of flexibility. While

default parameters are provided, STdGCN allows users to fine-tune the model according to the

specific characteristics of their ST platforms. For example, in the preprocessing module, users

have the option to input either a raw count matrix or a normalized expression matrix from both

ST data and scRNA-seq data. Additionally, users can utilize their own genes of interest to

customize the deconvolution process. The pseudo-spot simulation module offers users the

flexibility to select the desired cell number and cell-type number within a spot, adapting to the

specific characteristics of their ST platform. The number of simulated spots is also adjustable,

allowing users to strike a balance between improved performance (i.e. more simulated spot

number) and computational efficiency and memory usage. In the GCN module, users have the

freedom to customize various aspects of the model, such as the weights of the link graphs, the

depth of expression/spatial GCN layers, the depth of the fully-connected neural network layers,

and the dimensionality of the hidden layers. Lastly, STdGCN provides output in both csv table

and the anndata format [47]. This enables users to seamlessly integrate the results with numerous

downstream analysis tools, such as Scanpy, which are widely accessible and user-friendly.

Despite the aforementioned merits, STdGCN still has limitations. One limitation is that

the predictive performance of STdGCN varies across cell types. In particular, the relative

performance of STdGCN in cell types with lower proportions is not as good as in the cell types

with higher proportions (**Table S3~S4**). This indicates that there is room for improvement in

the current pseudo-spot generation method and the construction model of the link graph. Thus,

our future work will focus on ameliorating the effectiveness of utilizing reference data.

In conclusion, STdGCN stands as the pioneering model that incorporates the spatial

coordinates from ST data and cell type profiles learned from single-cell RNA-seq data for the

purpose of deconvolving local cell type proportions within complex tissues. By leveraging this

innovative approach, STdGCN has the potential to greatly enhance our understanding of tissue

spatial architecture and provide invaluable support for downstream analyses at the cellular level.

With its ability to unravel the intricate composition of cell types within ST datasets, STdGCN

opens new avenues for exploring the complexities of tissue organization and advancing our

knowledge of biological systems.

Competing Interests

The authors declare that they have no conflict of interests.

Authors' contributions

Y.Luo and Y.Li designed the research; Y.Li collected and analyzed data, and contributed to the

theory; Y.Luo and Y.Li drafted the manuscript. All authors have read, edited and approved the

final manuscript.

Acknowledgements

We are grateful to Xin Wu, Zexian Zeng, and Yufeng He for the comments and suggestions

during the preparation of the manuscript. This study is supported in part by NIH grant

U01TR003528, 1R01LM013337.

References

1.   Eng, C.L. et al. Transcriptome-scale super-resolved imaging in tissues by RNA

seqFISH. *Nature* **568**, 235-239 (2019).

2.   Merritt, C.R. et al. Multiplex digital spatial profiling of proteins and RNA in fixed

tissue. *Nat Biotechnol* **38**, 586-599 (2020).

3.   Rodriques, S.G. et al. Slide-seq: A scalable technology for measuring genome-wide

expression at high spatial resolution. *Science* **363**, 1463-1467 (2019).

4.   Wang, X. et al. Three-dimensional intact-tissue sequencing of single-cell

transcriptional states. *Science* **361** (2018).

5.   Berglund, E. et al. Spatial maps of prostate cancer transcriptomes reveal an unexplored

landscape of heterogeneity. *Nat Commun* **9**, 2419 (2018).

6.   Moncada, R. et al. Integrating microarray-based spatial transcriptomics and single-cell

RNA-seq reveals tissue architecture in pancreatic ductal adenocarcinomas. *Nat*

Biotechnol **38**, 333-342 (2020).

7.   Thrane, K., Eriksson, H., Maaskola, J., Hansson, J. & Lundeberg, J. Spatially Resolved

Transcriptomics Enables Dissection of Genetic Heterogeneity in Stage III Cutaneous

Malignant Melanoma. *Cancer Res* **78**, 5970-5979 (2018).

8.   Stuart, T. & Satija, R. Integrative single-cell analysis. *Nat Rev Genet* **20**, 257-272

(2019).

9.   Du, R., Carey, V. & Weiss, S.T. deconvSeq: deconvolution of cell mixture distribution

in sequencing data. *Bioinformatics* **35**, 5095-5102 (2019).

10.   Tsoucas, D. et al. Accurate estimation of cell-type composition from gene expression

data. *Nat Commun* **10**, 2975 (2019).

11.   Wang, X., Park, J., Susztak, K., Zhang, N.R. & Li, M. Bulk tissue cell type

deconvolution with multi-subject single-cell expression reference. *Nat Commun* **10**,

380 (2019).

12.   Newman, A.M. et al. Determining cell type abundance and expression from bulk tissues

309   with digital cytometry. *Nat Biotechnol* **37**, 773-782 (2019).

310   13.   Gong, T. & Szustakowski, J.D. DeconRNASeq: a statistical framework for

311   deconvolution of heterogeneous tissue samples based on mRNA-Seq data.

312   *Bioinformatics* **29**, 1083-1085 (2013).

313   14.   Dong, M. et al. SCDC: bulk gene expression deconvolution by multiple single-cell

314   RNA sequencing references. *Brief Bioinform* **22**, 416-427 (2021).

315   15.   Andersson, A. et al. Single-cell and spatial transcriptomics enables probabilistic

316   inference of cell type topography. *Commun Biol* **3**, 565 (2020).

317   16.   Elosua-Bayes, M., Nieto, P., Mereu, E., Gut, I. & Heyn, H. SPOTlight: seeded NMF

318   regression to deconvolute spatial transcriptomics spots with single-cell transcriptomes.

319   *Nucleic Acids Res* **49**, e50 (2021).

320   17.   Yang, T. et al. AdRoit is an accurate and robust method to infer complex transcriptome

321   composition. *Commun Biol* **4**, 1218 (2021).

322   18.   Cable, D.M. et al. Robust decomposition of cell type mixtures in spatial transcriptomics.

323   *Nat Biotechnol* **40**, 517-526 (2022).

324   19.   Dong, R. & Yuan, G.C. SpatialDWLS: accurate deconvolution of spatial transcriptomic

325   data. *Genome Biol* **22**, 145 (2021).

326   20.   Song, Q. & Su, J. DSTG: deconvoluting spatial transcriptomics data through graph-

327   based artificial intelligence. *Brief Bioinform* **22** (2021).

328   21.   Bae, S. et al. CellDART: cell type inference by domain adaptation of single-cell and

329   spatial transcriptomic data. *Nucleic Acids Res* **50**, e57 (2022).

22.   Lopez, R. et al. DestVI identifies continuums of cell types in spatial transcriptomics

data. *Nat Biotechnol* (2022).

23.   Sun, D., Liu, Z., Li, T., Wu, Q. & Wang, C. STRIDE: accurately decomposing and

integrating spatial transcriptomics using single-cell RNA sequencing. *Nucleic Acids*

Res **50**, e42 (2022).

24.   Biancalani, T. et al. Deep learning and alignment of spatially resolved single-cell

transcriptomes with Tangram. *Nat Methods* **18**, 1352-1362 (2021).

25.   Chu, T., Wang, Z., Pe'er, D. & Danko, C.G. Cell type and gene expression

deconvolution with BayesPrism enables Bayesian integrative analysis across bulk and

single-cell RNA sequencing in oncology. *Nat Cancer* **3**, 505-517 (2022).

26.   Kleshchevnikov, V. et al. Cell2location maps fine-grained cell types in spatial

transcriptomics. *Nat Biotechnol* **40**, 661-671 (2022).

27.   Miller, B.F., Huang, F., Atta, L., Sahoo, A. & Fan, J. Reference-free cell type

deconvolution of multi-cellular pixel-resolution spatially resolved transcriptomics data.

Nat Commun **13**, 2339 (2022).

28.   Danaher, P. et al. Advances in mixed cell deconvolution enable quantification of cell

types in spatial transcriptomic data. *Nature Communications* **13** (2022).

29.   Zhao, E. et al. Spatial transcriptomics at subspot resolution with BayesSpace. *Nat*

Biotechnol **39**, 1375-1384 (2021).

30.   Zhu, Q., Shah, S., Dries, R., Cai, L. & Yuan, G.C. Identification of spatially associated

subpopulations by combining scRNAseq and sequential fluorescence in situ

hybridization data. *Nat Biotechnol* (2018).

31.   Lohoff, T. et al. Integration of spatial and single-cell transcriptomic data elucidates

353   mouse organogenesis. *Nat Biotechnol* **40**, 74-85 (2022).

354   32.   Moffitt, J.R. et al. Molecular, spatial, and functional single-cell profiling of the

355   hypothalamic preoptic region. *Science* **362** (2018).

356   33.   Chen, H. et al. Dissecting mammalian spermatogenesis using spatial transcriptomics.

357   *Cell Rep* **37**, 109915 (2021).

358   34.   Asp, M. et al. A Spatiotemporal Organ-Wide Gene Expression and Cell Atlas of the

359   Developing Human Heart. *Cell* **179**, 1647-1660 e1619 (2019).

360   35.   Stahl, P.L. et al. Visualization and analysis of gene expression in tissue sections by

361   spatial transcriptomics. *Science* **353**, 78-82 (2016).

362   36.   Salmen, F. et al. Barcoded solid-phase RNA capture for Spatial Transcriptomics

363   profiling in mammalian tissue sections. *Nat Protoc* **13**, 2501-2534 (2018).

364   37.   Chen, J. et al. A comprehensive comparison on cell-type composition inference for

365   spatial transcriptomics data. *Brief Bioinform* **23** (2022).

366   38.   Gladka, M.M. et al. Single-Cell Sequencing of the Healthy and Diseased Heart Reveals

367   Cytoskeleton-Associated Protein 4 as a New Modulator of Fibroblasts Activation.

368   *Circulation* **138**, 166-180 (2018).

369   39.   Hsieh, P.C., Davis, M.E., Lisowski, L.K. & Lee, R.T. Endothelial-cardiomyocyte

370   interactions in cardiac development and repair. *Annu Rev Physiol* **68**, 51-66 (2006).

371   40.   Wu, S.Z. et al. A single-cell and spatially resolved atlas of human breast cancers. *Nat

372   Genet* **53**, 1334-1347 (2021).

41.   Loi, S. et al. Tumor-Infiltrating Lymphocytes and Prognosis: A Pooled Individual

Patient Analysis of Early-Stage Triple-Negative Breast Cancers. *J Clin Oncol* **37**, 559-

569 (2019).

42.   Stanton, S.E. & Disis, M.L. Clinical significance of tumor-infiltrating lymphocytes in

breast cancer. *J Immunother Cancer* **4**, 59 (2016).

43.   Zgura, A., Galesa, L., Bratila, E. & Anghel, R. Relationship between tumor infiltrating

lymphocytes and progression in breast cancer. *Maedica (Bucur)* **13**, 317-320 (2018).

44.   Seager, R.J., Hajal, C., Spill, F., Kamm, R.D. & Zaman, M.H. Dynamic interplay

between tumour, stroma and immune system can drive or prevent tumour progression.

Converg Sci Phys Oncol **3** (2017).

45.   Pietras, K. & Ostman, A. Hallmarks of cancer: interactions with the tumor stroma. *Exp*

Cell Res **316**, 1324-1331 (2010).

46.   Kalluri, R. The biology and function of fibroblasts in cancer. *Nat Rev Cancer* **16**, 582-

598 (2016).

47.   Wolf, F.A., Angerer, P. & Theis, F.J. SCANPY: large-scale single-cell gene expression

data analysis. *Genome Biol* **19**, 15 (2018).

48.   Li, H. et al. A comprehensive benchmarking with practical guidelines for cellular

deconvolution of spatial transcriptomics. *Nat Commun* **14**, 1548 (2023).

49.   Dries, R. et al. Giotto: a toolbox for integrative analysis and visualization of spatial

expression data. *Genome Biol* **22**, 78 (2021).

50.   Haghverdi, L., Lun, A.T.L., Morgan, M.D. & Marioni, J.C. Batch effects in single-cell

RNA-sequencing data are corrected by matching mutual nearest neighbors. *Nat*

Biotechnol **36**, 421-427 (2018).

51.   Kipf, T.N. & Welling, M. Semi-Supervised Classification with Graph Convolutional

397   Networks. arXiv:1609.02907 (2016).

Methods

Benchmark data selection and preprocess. To curate the datasets for benchmark, we

conducted a thorough review of two published comparison studies on ST deconvolution [37, 48],

as well as examined the models benchmarked in this study. After strict comparison process, we

selected four datasets from four ST platforms, including seqFISH, MERFISH, seqFISH+ and

slide-seq (**Table S5**). These datasets were chosen because they offer single cell-level resolution.

Moreover, the selected datasets come with cell type annotations provided by the authors,

enabling us to obtain the ground truth after simulating low-resolution spots. In addition, the

selected datasets include both sequencing-based technologies and imaging-based technologies,

with gene sizes ranging from hundreds to genome wide (>20,000) and the spot size ranging

from tens to thousands. This diverse range of characteristics allows for a comprehensive

evaluation of the models in various scenarios. Importantly, all four datasets contain multiple

slices, which facilitates testing the robustness of the models across different samples. To create

synthetic spots for analysis, we resampled the datasets by dividing each slice into multiple

square pixel areas. Cells within each square pixel area were merged to form a synthetic spot.

As the ST platforms used provided single-cell-level data, we had access to the exact cell type

proportions for each synthetic spot. The single-cell level data was also used as the internal

single-cell reference in our study.

The seqFISH+ dataset was obtained from Zhu et al. [30] (access via Giotto package [49]:

getSpatialDataset(dataset = "seqfish_SS_cortex", method = "wget"), which encompasses the

detection of mRNAs for 10,000 genes corresponding to a section of the mouse somatosensory

(SS) region. We divided the SS region into two slices based on the author's annotation,

including a cortex slice and a sub-ventricular zone (SVZ) slice. For the cortex slice, we selected

400×400 square pixel area as a synthetic spot, whereas for the SVZ slice we chose 200×200

square pixel area. Any spot with cell number below two was discarded. There were 109 cortex

synthetic spots and 59 SVZ synthetic spots after resampling, respectively.

The   seqFISH   dataset   was   obtained   from   Lohoff   et   al. [31]

(https://content.cruk.cam.ac.uk/jmlab/SpatialMouseAtlas2020/). This study detected mRNAs

for 351 target genes in tissue sections of three mouse embryos at the 8–12 somite stage, with

two slices obtained for each embryo. In terms of the cell types annotated by the authors from

single-cell transcriptome atlases, we removed the cell types with very small proportions ($< 3\%$)

for cell type deconvolution. The coordinates of the cells were stored in columns

"x_global_affine" and "y_global_affine" within the file "metadata.Rds". For each slice, we

selected the 0.1×0.1 square "global_affine" area as a synthetic spot. Any spot with cell number

less than two was discarded. The number of predictive synthetic spots obtained for each slice

after applying the filtering criteria ranged from 1279 to 2350.

The   Slide-seq   dataset   was   downloaded   from   Chen   et   al. [33]

(https://www.dropbox.com/s/ygzpj0d0oh67br0/Testis_Slideseq_Data.zip?dl=0).   This   study

captured the spatial gene expression patterns in mouse testis slices, including three diabetes

mouse testis slices and three wild type mouse testis slices. The coordinates of the cells were

stored in columns "x" and "y" in the csv files starting with "BeadLocationsForR". For each

slice, we selected a synthetic spot size of 80×80 square pixel area. Any spot with cell number

less than two was discarded. The final number of predictive synthetic spots for each slice after

filtering ranged from 2046 to 2615.

The MERFISH dataset was downloaded from Chen et al. [32] (https://datadryad.org/stash/dataset/doi:10.5061/dryad.8t8s248). This study detected mRNAs for a set of 155 target genes in the mouse preoptic region. Approximately 1.1 million cells were measured across 181 slices obtained from 36 mice. Cell types with proportions smaller than 3% were removed for cell type deconvolution. The coordinates of the cells were stored in columns "xcoord" and "ycoord" in the file named "Moffitt_and_Bambah-Mukku_et_al_merfish_all_cells.csv". For each slice, we selected a synthetic spot size of 50×50 square pixel area. Any spot with cell number less than two was discarded. The final number of predictive synthetic spots for each slice after filtering ranged from 938 to 1247.

Preprocess of the human heart dataset. The human heart spatial transcriptomics dataset was obtained from Asp et al. [34]. This dataset contains three sequencing platforms: spatial transcriptomics [35, 36], scRNA-seq, and ISS. The tissue sections were collected from human embryonic cardiac samples at three developmental stages in the first trimester: 4.5–5, 6.5, and 9 post-conception weeks (PCW). Herein, we employed STdGCN to infer the cell-type composition of the spatial transcriptomics data. This dataset consists of 19 slices, with four slices from 4.5-5 PCW, nine slices from 6.5 PCW, and six slices from 9 PCW. During data deconvolution, we considered only the spots located within the tissues. The coordinates of the spots can be obtained from their corresponding spot names. We used the scRNA-seq dataset provided in this study as the reference data. It is important to note that the spatial transcriptomics dataset lacks single-cell resolution, making it impossible to obtain realistic cell-type compositions for the spots. Additionally, this study offers the differentially expressed gene

(DEG) analysis results based on the scRNA-Seq data (Table S3 in Asp et al. [34]). To filter cell-

type marker genes for Myoz2-enriched cardiomyocytes, ventricular cardiomyocytes, atrial

cardiomyocytes, we retained genes with FDR (marked as "'p_val_adj'") < 0.01, average log-

fold change (marked as "'avg_logFC'") > 1, fraction of cells expressing the genes in the cell

type (marked as "pct.1") > 0.7, and fraction of cells expressing the genes in the remaining cell

types (marked as "pct.2") < 0.3 from the downloaded DEG analysis table.

Preprocess of the human breast cancer dataset. The human breast cancer 10X Genomics

Visium dataset was obtained from Wu et al. [40]. The dataset consists of data from six breast

cancer patients, including two ER$^+$ samples and four TNBC samples. When performing data

deconvolution, we considered only the spots located within the tissues. The coordinates of the

spots can be found in the file "tissue_positions_list.csv". Additionally, in the same study, the

authors processed scRNA-seq from 26 breast tumor samples. For the purpose of deconvolution

in our study, we employed this scRNA-seq data as the reference dataset.

Benchmark metrics. To assess the effectiveness of cell type deconvolution algorithms across

the synthetic real-wrod datasets, we selected two metrics: JSD and RMSE. These metrics were

chosen to quantify the disparities between the predicted cell type proportions and the actual

proportions within each spot.

The JSD is a symmetric form of the Kullback-Leibler (KL) divergence. Denote P and Q

are the probability distributions representing the predicted and ground-truth results, respectively.

The JSD can be expressed as:

$$487 \qquad \text{JSD(P||Q)} = \frac{1}{2} D_{KL}(P||\frac{1}{2}(P+Q)) + \frac{1}{2} D_{KL}(Q||\frac{1}{2}(P+Q)) \qquad (1)$$

488 where D_{KL} is the Kullback-Leibler (KL) divergence, which can be expressed as:

$$489 \qquad D_{KL}(P||Q) = \sum_{x \in X} P(x) \log \frac{P(x)}{Q(x)} \qquad (2)$$

Typically, the JSD is a nonnegative number, and a smaller JSD represents a higher similarity

between the predicted result and the ground-truth, indicating a better performance of the

corresponding method in accurately estimating the cell type proportions.

To assess the performance of the models specifically for cell type A, denote the predicted

and ground-truth proportions of a spot are P(A) and Q(A). For the purpose of analysis, we

considered the remaining cell types as a group, collectively referred to as the non-A type. Their

proportions are denoted as 1-P(A) and 1-Q(A), respectively. The equation (2) will be expressed

as:

$$498 \qquad D_{KL}(P||Q) = P(A) \log \frac{P(A)}{Q(A)} + (1 - P(A)) \log \frac{(1 - P(A))}{(1 - Q(A))} \qquad (3)$$

499 Based on equation (3), we can eventually get the binary JSD between predicted proportions

500 compared to the ground truth of the cell type A.

501 We used the following equation to calculate RMSE:

$$502 \qquad \text{RMSE(P||Q)} = \sqrt{\frac{\sum_{i=1}^{K}(P(x_i) - Q(x_i))^2}{K}} \qquad (4)$$

503 When evaluating the performance of a model for deconvolving a spot, the variable K represents

504 the number of cell types. Conversely, when assessing the performance of the models for a

505 specific cell type, K represents the number of spots within a slice. RMSE is a nonnegative value,

506 and a smaller RMSE represents a lower dissimilarity between the predicted result and the

507 ground truth.

Cell type co-localization score. Cell–cell interactions often occur via receptor–ligand pairs.

Thus, identifying cell type co-localization provides potential cell-cell interactions within a

tissue. To quantify the degree of co-localization between two cell types within a spatial

transcriptomics (ST) slice, we have developed a cell type co-localization score denoted as

$(S(A, B))$. $S(A, B)$ can be calculated using the following functions:

$$S(A, B) = \frac{\sum_{all\ spots} S_A = 1\ and\ S_B = 1}{\sum_{all\ spots} S_A = 1\ or\ S_B = 1}$$

where $S_A = 1$ denotes that the location has cell type A. The value of $S(A, B)$ ranges between

0 and 1. A higher $S(A, B)$ a higher probability of co-localization between the two cell types

within a spot. Considering that the proportions of all cell types within a spot predicted by

STdGCN are greater than 0, To ensure meaningful co-localization analysis, we introduce a

threshold for $S(A, B)$ calculations. When the predicted proportion of cell type A within a spot

is below the threshold, we set $S_A = 0$. We used a threshold of 0.15 for the 10X Genomics

Visium human breast cancer dataset, and 0.1 for the developing human heart spatial

transcriptomic. This choice was made based on the characteristics of the dataset and the

platform used.

Implementation of STdGCN

Single-cell data preprocessing and feature selection. The first step in the processing pipeline

of STdGCN is to identify the cell type marker genes from scRNA-seq data. The input of the

single-cell expression matrix is flexible and can be used for either the raw count matrix or the

normalized matrix. For the raw count matrix, we used the Scanpy package in Python to

preprocess the data. We used the command "scanpy.pp.normalize_total" to normalize counts to

a standardized value of 10,000 per cell. Subsequently, we logarithmized the data matrix using

the command "sc.pp.log1p", followed by "scanpy.pp.scale" to scale data to achieve unit

variance and a mean of zero. During the preprocessing phase, we have incorporated options to

allow users to make decisions regarding the filtering of highly variable genes and the regression

of mitochondrial genes. These options provide flexibility based on specific analysis

requirements. Following preprocessing step, we used the principal component analysis (PCA)

(sc.tl.pca) for dimensionality reduction. To select cell type marker genes for each cell type, we

utilized logistic regression (sc.tl.rank_genes_groups(method= 'logreg')). The resulting gene

features were obtained by aggregating the top "n" cell type marker genes, where "n" represents

the desired number of markers to be considered. To streamline the preprocessing and marker

gene selection steps, we have integrated them into a single command named

"find_marker_genes." This integration enhances the ease of use and efficiency of the STdGCN

pipeline.

Pseudo-spots generation. The generation of pseudo-spots involves randomly selecting a

mixture of cells from scRNA-seq data. In our experience, the heterogeneity of the pseudo-spot

pool is a key factor in constructing an effective GCN link graph. Conventional methods for

pseudo-spot generation may result in an unbalanced pseudo-spot pool and limited heterogeneity,

particularly due to the unbalanced proportions of cell types in reference single-cell data. To

address this issue, we designed a sampling-by-cell-type method for obtaining a more balanced

pseudo-spot generation. Simulating pseudo-spots requires specifying three user-defined hyper-

parameters: N_{up}, the upper limit of cell numbers in a pseudo-spot; N_{low}, the lower limit of cell

numbers in a pseudo-spot; and K_{up}, the upper limit of the number of cell types in a pseudo-spot.

The use of K_{up} is influenced by the fact that the number of cell types within a spot is usually

limited in real-word ST datasets, unlike bulk sequencing data. For each pseudo-spot simulation,

we first generated a random number between 1 and K_{up} to determine the number of cell types

(k) within a pseudo-spot. Subsequently, we randomly selected k cell-types from the entire cell

types (K_w) to form the cell type set for the corresponding pseudo-spot. We next randomly

selected a number (N_s) between N_{low} and N_{up} to determine the cell number in this pseudo-spot.

We then randomly picked up N_s cells from the selected cell-type set to construct the pseudo-

spot. The expression level of the pseudo-spot was determined by the summation of the counts

of the randomly selected cells. To streamline the entire process of pseudo-spot generation, we

have implemented a single command called "pseudo_spot_generation" within the STdGCN

framework.

Real-spots and pseudo-spots preprocessing and integration. The preprocessing steps for

real-spots and pseudo-spots were performed in a similar manner to the reference single-cell

data. After preprocessing, we retained the previously selected marker genes for downstream

analyses. To facilitate data integration, we employed PCA for dimensionality reduction,

allowing the mapping of real- and pseudo- spots into a common latent space. The integrated

dimensionality reduction matrix was used as the input for constructing the subsequent link

graph.

Link graph construction. The STdGCN system includes two link graphs, an expression graph

and a spatial graph. The expression graph captures the expression similarity between spots. It

is a hybrid graph composed of three sub-graphs, a pseudo-spots internal graph, a real-spots

internal graph, and a real-to-pseudo-spot graph. We employed the "cosine" distance to measure

the distances between spots and the MNN [50] algorithm to establish links between spots. In MNN

model, a link between two spots (a and b) if and only if spot a belongs to one of the nearest

neighbors of spot b, and spot b is also one of the nearest neighbors of spot a. For the real-to-

pseudo-spots link graph, we applied the integrated dimensionality reduction matrix of the two

datasets as the input feature, with the parameter "nearest neighbors = 20" per cell. For the two

internal link graphs, data integration was not required, we thus directly used the normalized

expression matrix of the real-spot/pseudo-spot as the input feature. The parameter "nearest

neighbors" was set to 10 for the real-spots internal graph and 20 for pseudo-spots internal graph.

The adjacency matrix of the hybrid expression graph (A^{exp}) is a binary matrix. $A_{ab}^{exp} = 1$

represents a link between two spots and vice versa.

The spatial graph (A^{sp}) represents the spatial distances between real-spots. We calculated

the Euclidean distance between a given spot and all other spots (D_{ab}) using their relative spatial

coordinates. In our approach, we defined the distance between two neighboring spots as one

unit (=1). To construct the spatial graph, we established links between spots that have distances

smaller than a predefined radius (D_{upper}). The adjacency matrix of the spatial graph is defined

as follows:

594
$$A_{ab} = \begin{cases} 0 & when\ D_{ab} \geq D_{upper} \\ 1/D_{ab} & when\ D_{ab} < D_{upper} \end{cases}$$

Unlike the expression link graph, which is represented as a binary matrix, the spatial graph is a

weighted link graph based on the spatial distance between spots.

Model description. The STdGCN is built on the GCN [51], a semi-supervised learning algorithm

designed for graph-structured data. GCN is based on an efficient variant of convolutional neural

networks that directly operate on graphs. Denoting V and E are the sets of nodes and edges

in the graph $G = (V, E)$, the objective of GCN is to learn the graph representations by

combining node embeddings with those of their neighboring nodes. The model takes two

components as input: a feature matrix (X) representing the embeddings of all graph nodes,

and an adjacency matrix (A) depicting the structural connections in the link graph. Each GCN

layer can be written as a non-linear function

$$H^{(l+1)} = f(H^l, \tilde{A}) = \sigma(\tilde{A} H^l W^l)$$

where H^l is the output for the l-th layer with $H^l = X$, W^l is the weight matrix for the l-th

layer that needs to be learned during the training, $\sigma(\cdot)$ is the activation function, and $\tilde{A} =$

$D^{-1/2}(A/\lambda + I)D^{1/2}$ is the symmetric normalized adjacency matrix of A. The $\lambda = 20$ is a

stabilizing factor, as we have observed that incorporating this factor can increase the accuracy

and robustness of STdGCN. Here, I represents the identity matrix and D is the diagonal

degree matrix of $(A/\lambda + I)$.

The STdGCN constructs a comprehensive and diverse graph, containing both pseudo-spots

and real-spots as individual nodes, enabling the adaptation of real-spots from pseudo-spots. The

objective of STdGCN is to predict the cell-type proportions for the real-spots. Therefore, the

label for each spot is a vector that includes proportions for all cell-types ($Y_{prop} \in$

$\mathbb{R}^{(n_r + n_p) \times n_{type}}$), where n_r and n_p denote the real- and pseudo- spot numbers and n_{type}

denote the cell-type number. We used the expression level for the real- and pseudo spots as the

input matrix ($X \in \mathbb{R}^{(n_r+n_p) \times g}$), where g represents the selected gene number. The input matrix

was then forward propagated to two two-layer GCNs using the two previously constructed link

graphs, an expression graph ($A^{exp} \in \mathbb{R}^{(n_r+n_p) \times (n_r+n_p)}$) and a spatial graph ($A^{sp} \in$

$\mathbb{R}^{(n_r+n_p) \times (n_r+n_p)}$), with the function

623
$$\begin{cases} X_{exp} = ELU(\tilde{A}^{exp}ELU(\tilde{A}^{exp}XW_{exp}^{(0)})W_{exp}^{(1)}) \\ X_{sp} = ELU(\tilde{A}^{sp}ELU(\tilde{A}^{sp}XW_{sp}^{(0)})W_{sp}^{(1)}) \end{cases}$$

624   to obtain an expression embedding ($X_{exp} \in \mathbb{R}^{(n_r+n_p) \times m}$) and a spatial embedding ($X_{sp} \in$

625    $\mathbb{R}^{(n_r+n_p) \times m}$). Where m is the dimension of the embedding space and $ELU(\cdot)$ is the activation

626   function that can be expressed as

627
$$ELU(x) = \begin{cases} x, & if\,x > 0 \\ e^x - 1, & if\,x \le 0 \end{cases}.$$

628   We next concatenated the two GCN output embedding matrix ($X_{con} = [X_{exp},\, X_{sp}] \in$

629    $\mathbb{R}^{(n_r+n_p) \times 2m}$) as the input feature and propagated it through two additional two-layer fully

630   connected neural networks using the following function

631
$$\tilde{Y}_{prop} = softmax(b^{(1)} + a^1 ELU(b^{(0)} + a^{(0)}X_{con}))$$

632   where $a^{(l)}$ and $b^{(l)}$ indicate the weight matrix and additive bias for the l-th layer respectively,

633    $\tilde{Y}_{prop} \in \mathbb{R}^{(n_r+n_p) \times n_{type}}$ is the prediction of cell-type proportions, and $softmax(\cdot)$ is the

634   activation function that can be expressed as

635
$$softmax(x_i) = \frac{x_i}{\sum_j x_j}.$$

We used the KL divergence with L2 penalty as the loss function during the training of STdGCN.

Parameter settings. Herein, we used a two-layer GCN architecture for the implementation of

STdGCN, as we found that utilizing two GCN layers yielded superior performance compared

to employing a single layer or multiple layers. To train STdGCN, we adopted stochastic gradient

descent (SGD) with a maximum of 3,000 epochs (Python command: torch.optim.SGD

(model.parameters(), lr=0.2, momentum=0.9, weight_decay=0.0001, dampening=0,

nesterov=True)). We implemented an early stopping mechanism where training would cease if

the validation loss did not decrease for 20 consecutive epochs. We used a self-adaption method

called orch.optim.lr_scheduler.ReduceLROnPlateau to adjust the learning rate during training.

The reduced learning rate factor was 0.1 and the learning rate was reduced after 5 epochs

without improvement.

Code Availability:

STdGCN and all script files used in the analysis in this manuscript can be downloaded from

GitHub at https://github.com/luoyuanlab/stdgcn.

Data Availability:

All datasets used in this study are publicly available. The seqFISH dataset of the mouse

embryos are available for download at

https://content.cruk.cam.ac.uk/jmlab/SpatialMouseAtlas2020/. The seqFISH+ dataset of the

mouse somatosensory region are available using the Giotto command getSpatialDataset(dataset

= "seqfish_SS_cortex", method = "wget"). The Slide-seq data of the mouse testis are available

for download at https://www.dropbox.com/s/ygzpj0d0oh67br0/Testis_Slideseq_Data.zip?dl=0.

The MERFISH data of the mouse testis are available for download at

https://datadryad.org/stash/dataset/doi:10.5061/dryad.8t8s248. The 10X Genomics Visium data

of the human breast cancer are available for download at

https://doi.org/10.5281/zenodo.4739739 and https://doi.org/10.5281/zenodo.3957257. The

spatial transcriptomic data of the developing human heart are available for download at

https://www.spatialresearch.org/resources-published-datasets/doi-10-1016-j-cell-2019-11-

025/.

Table 1. The average ranks of the 15 benchmarked models across the four datasets.

	Model	seqFISH	seqFISH+	Slide-seq	MERFISH
	STdGCN	1.00	1.00	1.33	1.83
	cell2location	8.17	3.50	9.17	14.07
	CellDART	9.17	7.50	8.50	5.07
	DestVI	3.83	15.00	15.00	11.87
	DSTG	15.00	13.00	14.00	14.80
	RCTD	2.33	4.00	3.67	1.83
	spatialDWLS	4.83	6.00	6.17	4.10
JSD	SPOTlight	13.33	12.50	11.50	11.82
	STdeconvolve	12.17	13.00	12.00	7.61
	stereoscope	5.17	7.50	7.33	10.96
	STRIDE	13.50	10.50	9.50	9.57
	Tangram	10.00	10.50	6.83	7.30
	Adroit	6.33	3.00	3.50	3.07
	BayesPrism	10.67	4.00	1.83	10.24
	SpatialDecon	4.50	9.00	9.67	5.87
	STdGCN	1.00	1.00	1.33	1.40
	cell2location	6.83	2.50	7.50	12.89
	CellDART	9.33	7.50	11.17	4.19
	DestVI	3.17	15.00	14.17	12.08
	DSTG	15.00	12.50	14.33	14.46
	RCTD	2.67	4.00	4.17	3.06
	spatialDWLS	7.50	6.50	6.17	5.48
RMSE	SPOTlight	13.83	12.50	9.67	10.94
	STdeconvolve	12.50	13.50	12.33	8.18
	stereoscope	5.00	7.50	7.33	13.45
	STRIDE	12.67	9.50	8.50	7.98
	Tangram	10.33	11.50	7.67	8.50
	Adroit	5.50	3.00	2.83	2.35
	BayesPrism	10.00	4.50	2.00	9.85
	SpatialDecon	4.67	9.00	10.83	5.21

667   Note: JSD, Jensen–Shannon divergence; RMSE, root mean square error

Figure 1 Schematic view of the STdGCN framework. STdGCN is a cell type deconvolution

method designed for spatial transcriptomics data, using scRNA-seq data as the reference. The

workflow of STdGCN involves several key steps. Firstly, STdGCN employs the scRNA-seq

reference data to identify cell-type marker genes and generate a pseudo-spot pool **(A)**. It then

builds two link graphs: a spatial graph **(B)** and an expression graph **(C)**. The expression graph

is a hybrid graph composed of three sub-graphs, a pseudo-spot internal graph, a real-spot

internal graph, and a real-to-pseudo-spot graph. These sub-graphs are formed using mutual

677   nearest neighbors (MNN) based on expression similarity. Based on the two link graphs, a GCN-

678   based model is utilized to propagate information from both real- and pseudo- spots. The output

679   of STdGCN is the predicted cell-type proportions for each spot **(D)**. Additionally, STdGCN

680   provides visualizations of the predicted results for the real-spot dataset in two different formats.

Figure 2 Four representative multi-cellular synthetic benchmark ST slices corresponding

to the four ST platforms. The scatter plots display the cell atlas of the mouse embryos

sequenced with seqFISH (**A**), mouse preoptic region sequenced with MERFISH (**C**), mouse

somatosensory (SS) region sequenced with seqFISH+ (**E**), and mouse testis region sequenced

with Slide-seq (**G**). We resampled these datasets by dividing each slice into multiple square

pixel areas (dash lines). Cells within each square pixel area were then merged into synthetic

spots. Pie plots illustrate the cell-type proportions for each synthetic spot after resampling

process (**B, D, F, and H**).

Figure 3 Assessment of STdGCN and benchmarked methods for estimation of cell type

proportions. (**A, B, C, and D**) The bar plots compare the performance of models with different

numbers of cells within a spot. We divided the spots into two groups: a smaller group (spots

with ≤5 cells) and a larger group (spots with >5 cells). Bar plots display the Jensen-Shannon

divergence (JSD) of the benchmarked methods for the two groups, as well as the total spots

across the four datasets. (**E, F, G, and H**) Comparison of the models in spots with different

number of cell type mixtures. We divided the spots into four groups based on the number of

cell types within a spot: singleton (one cell type), doubleton (two cell types), tripleton (three

cell types), and multiton (four or more cell types). Bar plots display the JSD of the benchmarked

methods for the four groups, as well as the total spots across the four datasets. (**I, J, K, and L**)

Bar plots display the JSD of STdGCN and benchmarked methods for distinguishing diverse

cell types across the four datasets.

Figure 4 Performance of STdGCN on the developing human heart spatial transcriptomic

dataset. (**A**) The spatial structure and cell type distribution of cells identified by in situ

sequencing (ISS) in a PCW6.5 heart tissue section. (**B**) UMAP plot of the cells as in (**A**). (**C**)

The pie plot displays the predicted cell-type proportions in each spot of the PCW6.5-6 heart

tissue section. (**D**) UMAP plot of the cells from the scRNA-seq dataset. (**E, F and G**) The

scatter plot displays the predicted proportions of epicardium cells (**E**), atrial cardiomyocytes

(**G**), smooth muscle cells from the PCW6.5-7 heart tissue section and their annotated results in

ISS data. (**H and I**) The scatter plots display the annotated results of Myoz2-enriched

cardiomyocytes (**H**) and Ventricular cardiomyocytes (**I**) in ISS data and the predicted

proportions from STdGCN, Adroit and RCTD, respectively. (**J**) The Venn diagram displays

the overlapping of cell-type marker genes between ventricular cardiomyocytes, Myoz2-

enriched cardiomyocytes, and atrial Cardiomyocytes after filtering. (**K**) The heatmap displays

the co-localization score for all deconvolved cell types in the three stages (PCW4.5-5, PCW6.5,

PCW9) of heart tissue sections. Red arrows are cardiomyocytes and blue arrows are endothelial

cells. (**M**) The distributions of ventricular cardiomyocytes in the ISS dataset.

Figure 5 Performance of STdGCN on the human breast cancer 10X Genomics Visium dataset. (**A and B**) UMAP plots of the cells from the scRNA-seq dataset divided by patient and cell type. (**C**) The heatmap displays the co-localization score (**Methods**) for normal cells, cancer cells, lymphocytes (integration of B-cells, T-cells and plasmablasts), stroma (integration of cancer-associated fibroblasts, perivascular-like and endothelial cells), and myeloid. (**D**) Mean predicted cell type proportions for the patient CID4465 (TNBC) and CID4535 (ER+). (**E,**

729   **F and G**) The scatter plot displays the predicted result for the patient CID4535. (**H**) The

730   heatmap displays the co-localization score for the patient CID4535. (**I, J and K**) The scatter

731   plot displays the predicted result for the patient CID4465. (**L**) The heatmap displays the co-

732   localization score for the patient CID4465.

Extended Data Figure 1 Assessment of STdGCN and benchmarked methods in spots with

different cell numbers. To compare the performance of models with different numbers of cells

within a spot, we divided the spots into two groups: a smaller group (spots with ≤5 cells) and a

738 larger group (spots with >5 cells). Bar plots display the root mean square error (RMSE) of the

739 benchmarked methods for the two groups, as well as the total spots across the four datasets.

Extended Data Figure 2 Assessment of STdGCN and benchmarked methods in spots with

different number of cell types. We divided the spots into four groups based on the number of

cell types within a spot: singleton (one cell type), doubleton (two cell types), tripleton (three

744   cell types), and multiton (four or more cell types). Bar plots display the root mean square error

745   (RMSE) of the benchmarked methods for the four groups, as well as the total spots across the

746   four datasets.

Extended Data Figure 3 Assessment of STdGCN and benchmarked methods in different

cell types. Bar plots display the root mean square error (RMSE) of STdGCN and benchmarked

methods for distinguishing diverse cell types across the four datasets.

Extended Data Figure 4 Pie plots of the predicted cell type proportions of the developing

human heart spatial transcriptomic dataset from STdGCN.

Extended Data Figure 5 Performance of STdGCN and benchmark models in distinguish ventricular cardiomyocytes and Myoz2-enriched cardiomyocytes on the developing human heart spatial transcriptomic dataset.

761