Differential contribution of nitrifying prokaryotes to groundwater nitrification

Markus Krüger[1], Narendrakumar Chaudhari[1,2], Bo Thamdrup[3], Will Overholt[1], Laura Bristow[3], Martin Taubert[1], Kirsten Küsel[1,2], Nico Jehmlich[4], Martin von Bergen[2,4,5], Martina Herrmann[1,2*]

[1]Aquatic Geomicrobiology, Institute of Biodiversity, Friedrich Schiller University, Jena, Germany.

[2]German Center for Integrative Biodiversity Research (iDiv) Halle-Jena-Leipzig, Leipzig, Germany.

[3]Nordcee, Department of Biology, University of Southern Denmark, Odense, Denmark. [4]Department of Molecular Systems Biology, Helmholtz Centre for Environmental Research – UFZ, Leipzig, Germany.

[5]Faculty of Biosciences, Pharmacy and Psychology, Institute of Biochemistry, University of Leipzig, Germany.

* Corresponding author: martina.herrmann@uni-jena.de

Abstract

The ecophysiology of complete ammonia-oxidizing bacteria (CMX) of the genus *Nitrospira* and their widespread occurrence in groundwater suggests that CMX bacteria have a competitive advantage over ammonia-oxidizing bacteria (AOB) and archaea (AOA) in these environments. However, the specific contribution of their activity to nitrification processes has remained unclear. We aimed to disentangle the contribution of CMX, AOA and AOB to nitrification and to identify the environmental drivers of their niche differentiation at different levels of ammonium and oxygen in oligotrophic carbonate rock aquifers. CMX *amoA* genes accounted on average for 16 to 75% of the total groundwater *amoA* genes detected. Nitrification rates were positively correlated to CMX clade A associated phylotypes and AOB affiliated with *Nitrosomonas ureae*. Short-term incubations amended with the nitrification inhibitors allylthiourea and chlorate suggested that AOB contributed a large fraction to overall ammonia oxidation, while metaproteomics analysis confirmed an active role of CMX in both ammonia and nitrite oxidation. Ecophysiological niche differentiation of CMX clades A and B, AOB and AOA was linked to their requirements for ammonium, oxygen tolerance, and metabolic versatility. Our results demonstrate that despite numerical predominance of CMX, the first step of nitrification in oligotrophic groundwater appears to be primarily governed by AOB. Higher growth yields at lower ammonia turnover rates and energy derived from nitrite oxidation most likely enable CMX to maintain consistently high populations.

Introduction

Nitrification is a globally relevant process of the nitrogen cycle mediated by a polyphyletic group of microorganisms [1]. The discovery of complete ammonia oxidizers (comammox bacteria, CMX) challenged the paradigm of a strict division of labor in the process of nitrification [2]. While ammonia oxidation is carried out by ammonia-oxidizing bacteria (AOB) and archaea (AOA) [3] and nitrite oxidation by nitrite-oxidizing bacteria (NOB) [2], CMX of the genus *Nitrospira* can perform complete oxidation of ammonia (NH_3) to nitrate (NO_3^-) in one cell [4, 5]. CMX have been frequently reported from engineered environments such as drinking water or wastewater treatment facilities [6–8] and from natural environments like soils [9, 10], surface freshwater [11, 12] and groundwater [13, 14]. However, our knowledge about the role and relevance of CMX for nitrification activity in natural environments is still scarce.

In most environments, CMX will compete with canonical nitrifiers for either ammonia (NH_3) or nitrite (NO_2^-) or both, raising the question of the actual contribution of CMX to nitrification processes and of the factors that control their co-occurrence with AOB, AOA, and NOB. Recent investigations consider affinity to NH_3, growth rates, growth yields, and metabolic versatility as key factors that determine the niche differentiation and competition between CMX, AOB, and AOA [2, 15]. Since CMX exhibit higher affinity to NH_3 and oxygen than terrestrial AOB [15, 16] along with higher growth yields compared to AOB and AOA [2], CMX might have a competitive advantage over AOB in environments that are limited in NH_3 and ammonium (NH_4^+) such as oligotrophic groundwater. Particular lineages of AOB and AOA exhibit adaptation to NH_4^+ limited conditions [17–20] and have previously been reported from groundwater environments [21–23]. As such, oligotrophic groundwater represents an excellent microbial habitat to study the co-occurrence of CMX, AOB and AOA and their contribution to nitrification activity.

Based on the phylogeny of the ammonia monooxygenase (Amo), the key enzyme of ammonia oxidation, CMX *Nitrospira* are divided into two clades, A and B [2, 4]. Metagenomic studies uncovering the phylogeny and metabolic features of CMX proposed an ecophysiological niche

differentiation between the two clades in natural habitats [4, 21, 24–26]. Those findings are

linked to genomic variability in nitrogen uptake and alternative energy metabolism [25, 26],

and several studies suggest a more prominent role of clade A in nitrification. So far, CMX

enrichments and isolates comprise only clade A representatives [27]. [13]C stable isotope

probing confirmed growth of CMX clade A with NH_4^+ during incubations of wetland coast

sediments [28], and CMX clade A were highly abundant in salt marsh sediments showing

strong potential for nitrification [29]. However, the ecophysiological niche differentiation

between the clades has rarely been studied in the context of their co-occurrence and

competition with canonical nitrifiers.

Previous studies from karstic carbonate rock aquifers of the Hainich Critical Zone Exploratory

in central Germany pointed to strong links between *in situ* nitrification and

chemolithoautotrophic carbon fixation [30, 31]. In addition, genome resolved metagenomics

revealed an important role of CMX *Nitrospira* in the nitrogen and carbon cycle in this

oligotrophic habitat [21]. The aquifers are characterized by a high spatial heterogeneity of NH_4^+

and NO_3^- concentrations across an oxygen gradient, linked to differences in groundwater flow

regime and distance to groundwater recharge areas [28, 29, 30]. These settings are ideal for

the investigation of the ecophysiological niche differentiation of CMX clade A and B, AOB and

AOA.

We combined [15]N-based nitrification rate measurements with selective nitrification inhibitors

and genome-resolved metagenomics and metaproteomics to assess the contributions of CMX

Nitrospira, AOB and AOA to overall groundwater nitrification in these aquifers and to follow

the composition of the groundwater nitrifier populations across the hydrochemical gradients.

We hypothesized that (i) CMX act as key ammonia oxidizers in these oligotrophic groundwater

habitats and account for a large fraction of overall nitrification activity. We further hypothesized

that (ii) distribution patterns of ammonia oxidizers across gradients of NH_4^+ and oxygen

availability are linked to their ecophysiological niche differentiation, indicated by differences in

encoded metabolic potentials.

Materials and methods

Study site characteristics and sampling of groundwater

Groundwater was obtained from karstic limestone aquifers at the Hainich Critical Zone Exploratory (CZE) in Thuringia (Germany) within the framework of the Collaborative Research Center (CRC) AquaDiva [34]. Further details about location and geological setting are reported elsewhere [33, 34]. Sampling wells H14, H32, H42, H43, H52 and H53 at varying depths provide access to oxygen-deficient (less than 15 µM DO) groundwater from an upper mudstone-dominated low-flow aquifer (HTU) and wells H13, H31, H41 and H51 (mean values 154 to 672 µM DO) provide access to oxic groundwater from a lower fast-flow limestone-dominated karstified aquifer (HTL) [32, 34] (Figure 1A, Figure 1B). The main recharge zone for groundwater is located at the hilltop of the transect near wells H13 and H14. Groundwater collected from regular monthly sampling between 2018 to 2021 was obtained using submersible motor pumps (Grundfos, Denmark). Hydrochemical parameters (pH, T, TOC, DOC) were determined as described in [32, 33]. NH_4^+ (sum of NH_3 and NH_4^+), NO_2^- and NO_3^- concentrations were measured with colorimetric methods [35, 36] from groundwater filtered through 0.2 µm-pore size sterile polyvinylidine fluoride filters. . Concentrations of free NH_3 were calculated based on pH and NH_4^+ concentrations (pKa NH_4^+/NH_3 = 9.73 at 10°C). Groundwater for molecular analysis was collected in sterile 10 L fluorinated polyethylene containers and filtered through 0.2 µm polycarbonate filters (Nuclepore, Whatman). Additional groundwater samples for metaproteomic analysis were collected from wells H14, H41, H43 and H52 in January 2019 and filtered through 0.2 µm pore size filters (Merck Millipore, Germany). All filters were immediately frozen on dry ice and stored at -80 °C until extraction.

^{15}N-isotope labelling and rate measurements

Groundwater was collected in sterile 0.5 L glass bottles and filled from the bottom using sterile tubing. Bottles were overfilled with three volumes exchanges before they were capped with silicone septa without headspace to minimize alterations of *in situ* oxygen concentrations. At each sampling campaign, groundwater was collected in triplicate bottles for each well along

with one control and kept at 4°C until further processing within 2 hours. Prior to the ^{15}N

labelling, 10 ml of each sampling bottle was removed for analysis of inorganic nitrogen

concentrations and pH, and replaced with N_2. Bottles containing anoxic groundwater were

amended with 1 % sterile oxygen (4 ml 100% oxygen) to stimulate nitrification under oxygen-

deficient conditions. Groundwater of control bottles was sterile filtered through 0.2 μm filters

(Supor, Pall Corporation, USA). Sterile filtered ^{15}N-labeled ammonium sulfate solution

(($^{15}NH_4)_2SO_4$) (98 at%, Cambridge Isotope Laboratories, Tewksbury) as the substrate source

for nitrifiers was added to all 0.5 L bottles to a final concentration of 50 μM, at a natural

background of 0.5 to 55 μM NH_4^+. Nitrification rate measurements were conducted according

to Overholt *et al.* [21] and detailed descriptions are provided in Supplementary Methods.

Incubation experiments in the presence of nitrification inhibitors

The ammonia and nitrite oxidation activity of CMX, and the nitrite oxidation activity of NOB

were selectively inhibited by 1 mM potassium chlorate ($KClO_3$) [7], and 100 μM allylthiourea

(ATU) was used to inhibit the ammonia oxidation activity of CMX and AOB [7, 37]. Nitrification

rate measurements with ^{15}N tracer were carried out over five days as described above.

Groundwater from well H41 was collected in February 2020 and amended with 50 μM NH_4^+

as ($^{15}NH_4)_2SO_4$ and one of the nitrification inhibitors along with untreated controls.

Groundwater for mesocosm experiments was collected in March and May 2021 and was filled

in sterile 2 L glass bottles without headspace to assess potential growth of ammonia oxidizers

with NH_4^+ in the presence of nitrification inhibitors at the same concentrations as before.

Groundwater from oxic well H41 (10 μM NH_4^+ *in situ*) was adjusted to 10 and 50 μM NH_4^+, and

groundwater from hypoxic well H53 (36 μM NH_4^+ *in situ*) was used without additional

supplement. All samples were incubated for 15 to 25 days at 15°C without agitation in the

dark. Subsamples for nitrogen compound analysis were taken at the onset and every two to

three days until the end of the incubation to monitor changes of the nitrogen chemistry

(Supplementary Methods). Incubation of well H41 with an initially 10 μM NH_4^+ was adjusted

again to 15 µM NH_4^+ after 15 days. After incubation, the groundwater was filtered on 0.2 µm

polycarbonate filters and stored at -80°C for later extraction of nucleic acids.

Nucleic acid extraction, cDNA-synthesis and quantitative PCR

Genomic DNA and total RNA of microbial biomass from the 0.2 µm filters were extracted using

the DNeasy PowerSoil kit (Qiagen, Germany) and the RNeasy PowerWater kit (Qiagen,

Germany), respectively, according to the manufacturer's specifications. RNA was further

processed with the TURBO DNA-free kit (Thermo Fischer Scientific, Germany) to remove

remnant DNA in the samples before cDNA synthesis [38]. Reverse transcription was

performed using the GoScript Reverse Transcription System (Promega, USA). Gene and

transcript abundances of *amoA* from CMX *Nitrospira*, AOB, AOA and of *nxrB* from *Nitrospira*-

like NOB were determined using a CFX96 qPCR cycler (Bio-Rad, USA). Primer pairs for

quantification were comamoAF/ comamoAR for CMX *amoA* [11], amoA-1F/ amoA-2R for AOB

amoA [39], Arch-amoAF/ Arch-amoAR for AOA *amoA* [40] and nxrB169f/ nxrB638r for

Nitrospira-like *nxrB* [41]. More details are provided in the Supplementary Methods. For testing

of potential correlations between CMX *amoA* and Nitrospira *nxrB* genes, we assumed on

average one copy of *amoA* and 1.5 copies of *nxrB* per cell. Calculation of the correction factor

1.5 was based on the analysis of putative canonical *Nitrospira*-affiliated MAGs from the

Hainich groundwater metagenome, which harbored 1 to 2 *nxrB* gene copies per genome [21]

Amplicon sequencing of *amoA* and 16S rRNA genes and sequence analysis

Amplicon libraries of CMX, AOB and AOA *amoA* genes were created using the NEBNext Ultra

DNA Library Prep Kit for Illumina (New England Biolabs, MA) according to the manufacturer's

guideline. The same primers as for qPCR were used for CMX and AOB *amoA*, while primers

Arch-amoA-104F/ Arch-amoA-616Rmod [42, 43] were chosen for sequencing of AOA *amoA*

genes due to shorter fragment length (Supplementary Methods). Sequencing was conducted

on an Illumina MiSeq using v3 chemistry. After trimming primer sequences, raw reads were

first quality filtered with DADA2 1.22.0 [44] and further processed with Mothur v.1.46.1 [45]

(Supplemental Methods). OTUs were assigned at 95% sequence identity for CMX, 95% for

AOB [46] and 96% for AOA [42] (see Dataset S3).

Amplicon libraries of the bacterial 16S rRNA gene were constructed as described elsewhere

[47] to assess the relative abundance of nitrite oxidizers present in the groundwater, including

non-*Nitrospira* NOB. Raw sequence reads were processed using DADA2 1.22.0. Taxonomy

was assessed using the SILVA reference data base release 138.1 [48]. Sequence analysis

was performed using the R package phyloseq [49].

Inferring metabolic potential from metagenome assembled genomes of CMX *Nitrospira* and other nitrifiers

To assess metabolic capabilities of the two CMX clades, genomic features of metagenome

assembled genomes (MAGs) were investigated. Groundwater sampling for metagenomic

sequencing and generation of MAGs were described in detail in [21, 50]. Here, Hainich

groundwater MAGs, which were taxonomically classified as *Nitrospira* on genus level and 243

Nitrospira reference genomes downloaded from NCBI (July 2021) were aligned and screened

for both *amoA* and *nxrB* genes with HMMER in Anvi'o v7.1 [21] using custom AmoA and NxrB

databases containing sequences from the FunGene repository [51].

After sorting good quality MAGs (> 90 % completeness, except *N. inopinata* with 42.25 %)

and dereplication, 35 CMX *Nitrospira* genomes were kept for final analysis. The affiliation of

those genomes to CMX clade A and clade B was achieved by alignment of protein sequences

of bacterial single copy core genes detected with default 'Bacteria_71' hmm profile using

MUSCLE [52] in Anvi'o and subsequent construction of a maximum likelihood phylogenetic

tree using FastTree with 1000 bootstrap iterations. The tree was then processed with iTOL for

visualization [53]. The genomic repertoire of representative groundwater nitrifier MAGs was

inferred by KEGG orthology (KO) previously generated using kofamscan [21] (Supplementary

Methods). Relative transcriptional activity of each investigated MAG was extracted from

Overholt *et al.* [21], who determined the transcriptomic coverages for each open reading frame

from each MAG from previously generated metatranscriptomic libraries of the groundwater

[30].

Metaproteomic analysis

Proteins were extracted from filters using a phenol-chloroform based protocol as previously

described [54]. Proteins were subjected to SDS polyacrylamide gel electrophoresis,

subsequently in-gel tryptic cleavage and LC-MS/MS analysis was performed as previously

described [55] (Supplemental Methods). Translated amino acid sequences from all genes of

the MAGs obtained from the metagenomic datasets [21, 50] were used as a reference

database for protein identification. Taxonomic and functional annotations of identified peptides

were transferred from the metagenomic dataset [21, 50]. Mass spectrometry proteomics data

have been deposited into the ProteomeXchange Consortium via the PRIDE [30] partner

repository.

Statistical analysis

All statistical analyses were conducted using R 4.12 (R Core Team, 2020) and packages FSA

[57], ComplexHeatmap [58] and vegan [59]. Analysis details are explained in the

Supplementary Methods. To analyze communities of CMX, AOB and AOA together, the

absolute *amoA* read abundances of OTUs > 2% of all reads for each ammonia oxidizer

community were determined as followed: The percentage of each OTU was calculated from

the relative abundance obtained by *amoA* amplicon sequencing and was multiplied by the

total *amoA* counts from corresponding samples measured with qPCR.

Results

Nitrification activity and distribution of nitrifiers across the different groundwater wells

Nitrification rates in groundwater varied over two orders of magnitude ranging from 0.54 – 136 nmol N L^{-1} d^{-1} across the wells (Figure 1C). We observed the highest rates at oxic well H41 with 93 ± 24 nmol N L^{-1} d^{-1} and high activity at hypoxic well H52 with 58 ± 4.0 nmol N L^{-1} d^{-1}. Rates were positively correlated with NH_4^+ concentrations in the groundwater (r_s = 0.67, p < 0.05), which were on average below 40 µM and significantly higher in the hypoxic wells (Figure 1B). NO_3^- concentrations ranged from 1.7 to 38 µM in hypoxic and from 80 to 384 µM in oxic groundwater and showed no correlation with the rates across the transect.

CMX *amoA* genes accounted for 16 ± 7.2 to 75 ± 12.3 % of the total groundwater *amoA* genes detected (mean values per well, see Dataset S2). Proportions of CMX *amoA* and AOB *amoA* genes increased substantially to maxima of 95% and 34% at wells with higher NH_4^+ concentrations (H41, H51, H52, H53), and gene abundances were positively correlated to NH_4^+ (r_s = 0.48, p < 0.01; r_s = 0.44, p < 0.01), respectively (Figure 1E, Figure 2D). In contrast, AOA dominated the ammonia-oxidizing community by up to 90% in wells with lower NH_4^+ concentrations located closer to the main recharge area (H13 and H14). Oxic well H41 and anoxic well H53 harbored the highest absolute *amoA* gene and transcript abundances of CMX and AOB (Figure 1D, Figure S 1A). Nitrification rates were strongly correlated with AOB *amoA* gene abundances (r_s = 0.87, p < 0.001) (Figure 2D), and AOB also had the highest *amoA* transcript/ gene ratios across all wells (Figure S 1B).

Across all wells, abundances of CMX *amoA* genes and *Nitrospira nxrB* genes encoding nitrite oxidoreductase were positively correlated to each other (r_s = 0.72, p < 0.001) (Figure S 1D). Assuming one *amoA* and 1.5 *nxrB* copies per genome (see Dataset S4), CMX accounted for 18 to 44 % and 48 to 70 % of the *Nitrospira* population at oxic well H41 and hypoxic well H52, respectively (Figure S 1C), which exhibited the highest observed nitrification rates (Figure 1C).

Furthermore, CMX and *Nitrosomonadaceae* had the highest relative abundance of peptides

at wells H41 and H52 (Figure S 2). Out of the *Nitrosomonadaceae* affiliated peptides, 88.3%

were assigned to *Nitrosomonas*.

The ammonia oxidizer community composition showed formation of clusters depending on

local groundwater chemistry and the location in the transect (Figure 2A). The CMX community

was dominated by three OTUs affiliated to clade A and six OTUs affiliated to clade B, of which

clade A OTUs were closely related to *Ca.* Nitrospira nitrificans (Figure 2C). These clade A

OTUs increased in abundance at wells H41, H51, H52 and H53 and accounted for up to 50%

of the CMX community (Figure 2B). CMX clade A affiliated OTUs were also positively

correlated to the nitrification rates (r_s = 0.72 − 0.88, p < 0.05), while only two low abundant

clade B affiliated OTUs correlated with the rates (Figure 2D). The AOB community was mainly

dominated by AOB OTU1 affiliated to *Nitrosomonas ureae*, which was the only AOB phylotype

showing a strong correlation with nitrification rates (r_s = 0.88, p < 0.01). The most dominant

AOA OTU1 related to *Nitrosoarchaeum koreense* accounted for an average of 70% of all AOA

across the groundwater sites (Figure S 3). Clustering based on the correlation with

hydrochemical parameters suggested similar ecological preferences of (i) AOA and some

clade B affiliated CMX OTUs, and (ii) *Nitrosomonas ureae* affiliated AOB OTU1 and CMX

clade A affiliated OTUs (Figure 2D). Among potential nitrite oxidizers, *Nitrospira* sp. were

consistently present in all the samples, while 16S rRNA gene sequences affiliated with

Nitrospinaceae, *Cand*. Nitrotoga, and *Nitrobacter* occurred only occasionally, and relative

abundances of these groups were at least one order of magnitude lower than those of

Nitrospira (Figure S 4).

Metabolic capabilities of CMX *Nitrospira* clades and AOB

We analyzed nine CMX affiliated metagenome assembled genomes (MAGs) from the Hainich

groundwater [21], which were phylogenomically affiliated to two CMX clade A MAGs and

seven clade B MAGs (Figure 3A). Both clade A MAGs (H51-bin291-1 and H41-bin216-1) were

closely related to *Ca.* Nitrospira nitrificans on the genome level and showed 99% amino acid

sequence identity to the AmoA of CMX OTU5 (Figure 2C). AmoA sequences of Clade B

representative MAGs showed closest relation to AmoA of CMX OTU2, OTU3, OTU8 and

OTU9.

Genes encoding essential components for ammonia oxidation, nitrite oxidation and urea

degradation were shared in MAGs representing both CMX clades (Figure 3B) and most of

these genes were highly transcribed in the groundwater. Metaproteomic analysis revealed the

presence of peptides related to AmoA, AmoB, AmoC, Hao and NxrA of *Nitrospira*,

demonstrating that nitrification-related proteins of CMX *Nitrospira* populations were formed *in*

situ. We found different genomic repertoires between CMX clade A and clade B affiliated

Nitrospira MAGs containing genes for sulfur and alternative energy metabolism such as

oxidation of formate and hydrogen, nitrogen transporter types and cellular regulation of

reactive oxygen species. CMX clade B MAGs exclusively harbored genes involved in

conversion of bicarbonate to carbon dioxide, oxidation of thiosulfate via the *sox* pathway,

dissimilatory sulfate reduction, as well as genes encoding two superoxide dismutases (SOD1/

SOD2) and a formate transporter (*focA*). Transporters for cellular import of urea as well as for

NO_2^-/NO_3^- were present in representative genomes from both CMX clade A and clade B. We

found that clade B MAGs encoded Amt-type NH_4^+ transporters, while clade A MAGs employed

Rh-type NH_4^+ transporters. MAGs from both clades harbored genes for ATP citrate lyase (*acl*),

which were transcriptionally active, indicating an active role in carbon fixation.

Compared to the CMX and AOA affiliated MAGs, transcription of *amo* genes was more than

100 times higher for AOB affiliated MAG H51-bin202-1 (Figure S 4). The AmoA sequence of

this MAG had 93% sequence identity to the AmoA of the dominant, *N. ureae* affiliated AOB-

OTU1 obtained from amplicon sequencing in this study (Figure S 5), suggesting that both

approaches probably identified the same population of AOB. This MAG employed Rh-type

NH_4^+ transporters as did representatives of CMX clade A, while AOA affiliated MAGs had Amt-

type NH_4^+ transporters as representatives of CMX clade B (Figure S 4).

Contributions of ammonia-oxidizing groups to ammonia oxidation activity

Short-term incubations in the presence of the nitrite oxidation inhibitor chlorate did not affect the ammonia oxidation rates (74.5 ± 9.4 nmol N L^{-1} d^{-1}) compared to the rates in controls without inhibitor (75.9 ± 3.7 nmol N L^{-1} d^{-1}) (Figure 4A). Chlorate is supposed to inhibit nitrite oxidation of NOB, and both ammonia oxidation and nitrite oxidation by CMX, as nitrite oxidoreductase catalyzes the formation of toxic chlorite from chlorate [7]. In the presence of the ammonia oxidation inhibitor allylthiourea, ammonia oxidation rates were not significantly different from zero. Allylthiourea is supposed to block ammonia oxidation, as it inhibits the ammonia monooxygenases of AOB as well as of CMX but affects those of AOA much less at the 100 µM concentrations used in this study [37, 60].

Long-term incubations (15-28 days) were performed to further investigate the potential growth and shifts in community composition of CMX, AOB, AOA and *Nitrospira*-affiliated NOB under the inhibitor treatments, using oxic groundwater from well H41 (condition 1: oxic/ 10 µM NH_4^+; condition 2: oxic/ 50 µM NH_4^+) and hypoxic groundwater from H53 (condition 3: hypoxic/ without supplement). Ammonia oxidation was always completely inhibited in the presence of allylthiourea, regardless of groundwater source or initial NH_4^+ concentrations (Figure S 6). NH_4^+ was only oxidized in incubations without inhibitor and in the chlorate treatment, while nitrite oxidation was only measured in incubations without inhibitor. Condition 3 without inhibitor showed complete consumption of NH_4^+ after 13 days, while it took 19 and 25 days under condition 1 and 2, respectively.

In incubations without inhibitor, CMX increased about two-fold (6.5×10^5 and 2.5×10^6 *amoA* L^{-1} after 25 and 20 days of incubation) compared to initial groundwater (2.8×10^5 and 1.44×10^6 *amoA* L^{-1}) under condition 1 and 2, respectively, while we did not detect growth of CMX under condition 3 (Figure 4B). CMX *amoA* gene abundance also increased in the allylthiourea treatment under condition 1 but decreased in the chlorate treatments. *Nitrospira*-affiliated NOB targeted using *nxrB* genes had similar growth patterns across condition and treatment types as CMX, and CMX accounted for an estimated 14 to 63 % of the *Nitrospira*-affiliated NOB

population. AOB *amoA* abundance increased by one to two orders of magnitude in all

untreated and chlorate treatments of the different conditions, while they decreased in the

presence of allylthiourea. AOA *amoA* gene abundance showed only minor variations during

all incubation settings. CMX OTU1 affiliated to CMX clade A (Figure 4C) and AOB-OTU1

related to *Nitrosomonas ureae* (Figure 4D) increased in abundance after incubations without

inhibitor. These two OTUs were already dominant in the ammonia oxidizer communities of the

original groundwater.

Discussion

We present evidence that despite numerical predominance of CMX bacteria, they contribute only a small fraction to ammonia oxidation in the studied oligotrophic groundwater. Instead, ammonia oxidation in this system appeared to be largely driven by AOB. Ranging from 0.54 up to 136 nmol N L^{-1} d^{-1}, the measured nitrification rates (ammonia oxidation + nitrite oxidation) matched those from oligotrophic lakes and open marine environments including oxygen minimum zones [61–64], but were lower than in rivers, and estuaries [65, 66] (Table S1). The observed nitrification rates did not reflect the spatial heterogeneity of groundwater NO_3^- concentrations, suggesting that surface-derived inputs mask the effects of *in situ* NO_3 formation on groundwater NO_3^- loads in these carbonate-rock aquifers.

The oligotrophic conditions in the groundwater appeared to select for ammonia oxidizers with the highest reported NH_3 affinities of their respective group. As the three groups differ in their NH_3 affinity, with 0.049 to 0.083 µM for *Ca.* Nitrospira inopinata [15] (corresponding to 11 to 19 µM NH_4^+ at pH 7.3), 0.3 to 4 µM for the *Nitrosomonas oligotropha* lineage [15, 20] (corresponding to 55 to 829 µM NH_4^+ at pH 7.3), and < 0.01 µM for AOA closely related to *Nitrosoarchaeum koreensis* [15] (corresponding to < 2.3 µM NH_4^+ at pH 7.3), fluctuating resource availability at overall limiting conditions may have enabled their coexistence. Previous metagenomics analysis work at this study site identified CMX as key nitrifiers and demonstrated that measured groundwater carbon fixation rates could fully be accounted for by carbon fixation associated to nitrification, assuming nitrification and carbon fixation kinetics of CMX *Nitrospira* [21]. Similarly, in our study, CMX accounted for up to 80% of the ammonia oxidizer community at the groundwater well showing the highest nitrification activity, supporting our original assumption that CMX were the most competitive ammonia-oxidizing group. Our inhibitor-based approach strongly suggested that ammonia oxidation appeared to be largely driven by AOB, even though CMX outnumbered AOB by a factor of 10 in the samples used for rate measurements.

Treatment with 1 mM chlorate in nitrification assays was expected to inhibit the nitrite oxidation

activity of canonical nitrite oxidizers as well as both steps of nitrification in CMX, since

reduction of chlorate to toxic chlorite from the activity of nitrite oxidoreductase would eventually

also inhibit ammonia oxidation in the same cell [7]. Ammonia oxidation was not affected in the

chlorate treatment, but was completely inhibited in the treatment with allylthiourea, which

specifically targets the ammonia monooxygenase and acts as a reversible inhibitor preventing

the oxidation of NH_4^+ [60]. At a concentration of below 160 µM, it selectively inhibits the activity

of AOB and CMX but not of AOA [37]. Consequently, this inhibitor approach strongly

suggested that AOB were primarily responsible for the highest observed ammonia oxidation

activity at well H41. A strong bias of our results by potential residual activity of CMX in the

presence of chlorate appears unlikely. Although *Nitrospira* is capable of detoxifying hazardous

chlorite by chlorite dismutases [67], the chlorate concentrations used here obviously resulted

in growth inhibition and cell death, which is in line with previous observations in wetland

sediment mesocosms [28]. Similarly, it appears unlikely that our results were caused by

excessive growth of AOB, since nitrification rates were linear across these five-days

incubations. A negligible contribution of AOA to ammonia oxidation was indicated by no

detectable activity in the presence of allylthiourea.

We cannot completely rule out that activity by AOB was enhanced in the chlorate treatment

due to absence of competition with CMX. This is one of the pitfalls associated with inhibition

experiments and could have led to an overestimation of the contribution of AOB in our case.

In fact, increased activity and growth of AOB in soil mesocosms where AOA were inhibited

was recently demonstrated by Zhao *et al*. [68]. Our transcriptional and metaproteomics-based

findings indicate a very active role of AOB in ammonia oxidation, supporting a substantial

contribution of this group to ammonia oxidation rates. Ammonium concentrations at well H41,

which had the highest nitrification activity, were around 10 µM NH_4^+ *in situ* and 50 µM were

used for rate measurements, corresponding to 0.04 or 0.22 µM free NH_3, respectively. The

NH_3 affinities and per cell ammonia oxidation rates differ between AOB and CMX [15]. It

remains unclear to what extend competition for NH_4^+ would restrict their ammonia oxidation

rates under the given ammonia-limited conditions. These conditions presumably affect AOB

more than CMX, and if this would result in increased activity of AOB when CMX were

suppressed due to inhibition. According to Vilardi *et al.*, CMX and AOB can fill independent

niches in the nitrifying community, which would reduce potential ammonia oxidation

competition between both groups [69]. More detailed knowledge of the exact ammonia

oxidation kinetics of the groundwater AOB and CMX representatives is needed to further

disentangle the mechanisms of their coexistence.

Despite the presumably small contribution of CMX to ammonia oxidation rates, we obtained

strong evidence of their active role in overall nitrification. CMX expressed genes and formed

peptides involved in both ammonia and nitrite oxidation *in situ*, which was confirmed by growth

of CMX in the presence of NH_4^+ in long-term incubations. However, it remains unclear if the

major energy source to sustain cell metabolism originated from complete ammonia oxidation

or rather from nitrite oxidation. In fact, a recent subsurface study detected transcriptional

activity in association with nitrite oxidation but not ammonia oxidation for a CMX-related MAG

[70], while CMX *Nitrospira* in water from rapid sand filters did not prefer to oxidize external

nitrite alone [7]. As we found NO_2^- transporters in representative groundwater MAGs from both

CMX clades, the cells may also be capable of taking up the NO_2^- formed by AOB, supporting

their primary role as nitrite oxidizers along with *Nitrosomonas ureae* and *Nitrosospira* sp.

affiliated AOB as key ammonia oxidizers.

AOB with faster ammonia oxidation rates than CMX [3, 15] did not outcompete CMX for NH_4^+

but simply processed a larger fraction of the available NH_4^+, thereby driving the ammonia

oxidation rates and providing the resulting NO_2^- to both CMX and canonical NOB. Overall, the

genus *Nitrospira* clearly dominated the NOB community, as NOB affiliated with

Nitrospinaceae, *Cand*. Nitrotoga, and *Nitrobacter* occurred only occasionally and were one or

two orders of magnitude less abundant than *Nitrospira*. The question remains unclear, why

CMX which accounted for more than 50% of the *Nitrospira* population were apparently favored

as nitrite oxidizers over canonical nitrite-oxidizing *Nitrospira* in the groundwater. Physiological

studies of *Ca.* Nitrospira inopinata demonstrated a much lower affinity to external NO_2^-

compared to canonical *Nitrospira* [15], a trait which is, however, not shared among all CMX

Nitrospira [71]. A recent microcosm-based study of drinking water biofilter media suggested

that CMX outcompete canonical NOB in natural nitrifier communities from nitrogen-limited

environments over time [69]. However, co-occurrence of CMX and canonical *Nitrospira*

appears to be common across a broad range of terrestrial and aquatic habitats [27], and their

respective roles in nitrification might be highly variable.

In contrast to our findings, CMX *Nitrospira* were identified as the driver of the two steps of

nitrification in groundwater-fed rapid gravity sand filters (RGSF) [7, 72]. Moreover, they made

a similar contribution as AOB to ammonia oxidation in wetland soils [28]. At low NH_4^+

concentrations of 71 µM and 1.1 µmol g^{-1}, CMX also dominated the ammonia oxidizer

communities in the RGSF [7] and in coastal wetland sediments [28] , respectively. However,

our results demonstrate that high abundance of CMX in an environment does not necessarily

imply that they also control the complete nitrification process. The competitive advantage of

CMX has been associated with a biofilm-associated lifestyle such as in RGSF. However,

division of labor in nitrification may remain the preferred strategy in the planktonic communities

of groundwater environments at low NH_4^+ concentrations.

Distribution patterns of ammonia oxidizers across the groundwater wells suggested different

niche preferences for AOB and CMX clade A compared to AOA and some clade B affiliated

CMX. Both AOA and CMX clade B dominated in groundwater closer to the major recharge

area with lower NH_4^+ availability. AOA can be the most abundant ammonia oxidizers in soils

[73] and several studies reported CMX clade B in soil habitats [9, 74]. Consequently, transfer

from soils to shallow groundwater could result in the observed high fractions of AOA and CMX

clade B in the ammonia oxidizer communities of the respective wells [47]. In turn, deeper and

more distant groundwater favored both AOB and clade A affiliated CMX under slightly

increased availability of NH_4^+ and lower DO concentrations.

The observed ecophysiological niche differentiation of CMX clade A and clade B across

different groundwater conditions was not reflected by differences in central nitrogen

metabolism of representative MAGs [21, 30] and metaproteomic data from the groundwater.

Genes encoding enzymes of ammonia and nitrite oxidation were highly transcribed and

respective peptides represented both CMX clades. However, differences in the encoded NH_4^+

transporters could be linked to different ecological preferences. Rh-type NH_4^+ transporters in

CMX clade A convey a better adaptation to higher NH_4^+ concentrations than Amt-type

transporters. As Rh-type transporters have higher uptake capacity but lower NH_3 affinity than

Amt-type transporters [25] and are also encoded in AOB [75], CMX clade A might have a

competitive advantage against clade B in the groundwater when more NH_4^+ is available. Along

with NH_4^+ uptake efficiency, also oxygen tolerance appeared to drive the distribution patterns

of the two clades, as CMX clade B but not clade A encode two types of superoxide dismutases

(SOD1, SOD2) [25, 26]. Finally, given the broader metabolic versatility of clade B including

the potential for formate and thiosulfate utilization, CMX clade B representatives might be less

strongly tied to energy generation by nitrification compared to the clade A representatives.

In conclusion, our findings point to a strong contribution of AOB to ammonia oxidation activity

in oligotrophic groundwater, despite numerical predominance of CMX. Given the limitations of

differential inhibitor assays, the exact contribution of CMX to ammonia oxidation *in situ*

remains unclear. Conditions of low NH_4^+ availability and varying oxygen concentrations

appeared to generally support coexistence of CMX, AOB and AOA with representatives

closely related to those with the highest NH_4^+ affinities in their respective groups. Our results

suggest that high substrate affinity, high growth yields, and a potentially major role in nitrite

oxidation enable CMX *Nitrospira* to maintain consistently high populations in the groundwater.

We further propose niche differentiation of CMX clade A and B with a more important role in

nitrification for clade A and a higher metabolic versatility, including the use of alternative

electron donors, for clade B.

References

1.   Kuypers MMM, Marchant HK, Kartal B. The microbial nitrogen-cycling network. *Nat Rev*
Microbiol . 2018. Nature Publishing Group. , **16**: 263–276

2.   Koch H, van Kessel MAHJ, Lücker S. Complete nitrification: insights into the
ecophysiology of comammox Nitrospira. *Appl Microbiol Biotechnol* 2018; **103**: 177–189.

3.   Lehtovirta-Morley LE. Ammonia oxidation: ecology, physiology, biochemistry and why
they must all come together. *FEMS Microbiol Lett* 2018; **365**.

4.   Daims H, Lebedeva E V., Pjevac P, Han P, Herbold C, Albertsen M, et al. Complete
nitrification by Nitrospira bacteria. *Nature* 2015; **528**: 504–509.

5.   Van Kessel MAHJ, Speth DR, Albertsen M, Nielsen PH, Op Den Camp HJM, Kartal B,
et al. Complete nitrification by a single microorganism. *Nature* 2015; **528**: 555–559.

6.   Fowler SJ, Palomo A, Dechesne A, Mines PD, Smets BF. Comammox *Nitrospira* are
abundant ammonia oxidizers in diverse groundwater-fed rapid sand filter communities.
Environ Microbiol 2018; **20**: 1002–1015.

7.   Gülay A, Fowler SJ, Tatari K, Thamdrup B, Albrechtsen H-J, Al-Soud WA, et al. DNA-
and RNA-SIP Reveal Nitrospira spp. as Key Drivers of Nitrification in Groundwater-Fed
Biofilters. *MBio* 2019; **10**.

8.   Spasov E, Tsuji JM, Hug LA, Doxey AC, Sauder LA, Parker WJ, et al. High functional
diversity among Nitrospira populations that dominate rotating biological contactor
microbial communities in a municipal wastewater treatment plant. *ISME J* 2020; **14**:
1857–1872.

9.   Li C, Hu HW, Chen QL, Chen D, He JZ. Niche differentiation of clade A comammox
Nitrospira and canonical ammonia oxidizers in selected forest soils. *Soil Biol Biochem*
2020; **149**: 107925.

10.   Osburn ED, Barrett JE. Abundance and functional importance of complete ammonia-
oxidizing bacteria (comammox) versus canonical nitrifiers in temperate forest soils. *Soil*
Biol Biochem 2020; **145**: 2015–2018.

11.   Zhao Z, Huang G, He S, Zhou N, Wang M, Dang C, et al. Abundance and community
composition of comammox bacteria in different ecosystems by a universal primer set.
Sci Total Environ 2019; **691**: 146–155.

12.   Shi Y, Jiang Y, Wang S, Wang X, Zhu G. Biogeographic distribution of comammox
bacteria in diverse terrestrial habitats. *Sci Total Environ* 2020; **717**: 137257.

13.   Liu S, Wang H, Chen L, Wang J, Zheng M, Liu S, et al. Comammox Nitrospira within
the Yangtze River continuum: community, biogeography, and ecological drivers. *ISME*
J 2020; **14**: 2488–2504.

14.   Yu C, Hou L, Zheng Y, Liu M, Yin G, Gao J, et al. Evidence for complete nitrification in
enrichment culture of tidal sediments and diversity analysis of clade a comammox
Nitrospira in natural environments. *Appl Microbiol Biotechnol* 2018; **102**: 9363–9377.

15.   Kits DK, Sedlacek CJ, Lebedeva E V., Han P, Bulaev A, Pjevac P, et al. Kinetic analysis
of a complete nitrifier reveals an oligotrophic lifestyle. *Nature* 2017; **549**: 269–272.

16.   Roots P, Wang Y, Rosenthal AF, Griffin JS, Sabba F, Petrovich M, et al. Comammox
Nitrospira are the dominant ammonia oxidizers in a mainstream low dissolved oxygen
nitrification reactor. *Water Res* 2019; **157**: 396–405.

17.   Martens-Habbena W, Berube PM, Urakawa H, De La Torre JR, Stahl DA. Ammonia
oxidation kinetics determine niche separation of nitrifying Archaea and Bacteria. *Nature*
2009; **461**: 976–979.

18.   Prosser JI, Nicol GW. Archaeal and bacterial ammonia-oxidisers in soil: the quest for
niche specialisation and differentiation. *Trends Microbiol* . 2012. Elsevier Current
Trends. , **20**: 523–531

19.   Herber J, Klotz F, Frommeyer B, Weis S, Straile D, Kolar A, et al. A single
Thaumarchaeon drives nitrification in deep oligotrophic Lake Constance. *Environ*
Microbiol 2020; **22**: 212–228.

20.   Sedlacek CJ, McGowan B, Suwa Y, Sayavedra-Soto L, Laanbroek HJ, Stein LY, et al.
A Physiological and Genomic Comparison of Nitrosomonas Cluster 6a and 7 Ammonia-
Oxidizing Bacteria. *Microb Ecol* 2019; **78**: 985–994.

21.   Overholt WA, Trumbore S, Xu X, Bornemann TL V., Probst AJ, Krüger M, et al. Carbon
fixation rates in groundwater similar to those in oligotrophic marine systems. *Nat Geosci*
2022; **15**: 561–567.

22.   Opitz S, Küsel K, Spott O, Totsche KU, Herrmann M. Oxygen availability and distance
to surface environments determine community composition and abundance of
ammonia-oxidizing prokaroytes in two superimposed pristine limestone aquifers in the
Hainich region, Germany. *FEMS Microbiol Ecol* 2014; **90**: 39–53.

23.   Van Der Wielen PWJJ, Voost S, Van Der Kooij D. Ammonia-oxidizing bacteria and
archaea in groundwater treatment and drinking water distribution systems. *Appl Environ*
Microbiol 2009; **75**: 4687–4695.

24.   Palomo A, Dechesne A, Cordero OX, Smets BF. Evolutionary Ecology of Natural
Comammox Nitrospira Populations. *mSystems* 2022; **7**.

25.   Palomo A, Pedersen AG, Fowler SJ, Dechesne A, Sicheritz-Pontén T, Smets BF.
Comparative genomics sheds light on niche differentiation and the evolutionary history
of comammox Nitrospira. *ISME J* 2018; **12**: 1779–1793.

26.   Poghosyan L, Koch H, Lavy A, Frank J, van Kessel MAHJ, Jetten MSM, et al.
Metagenomic recovery of two distinct comammox Nitrospira from the terrestrial
subsurface. *Environ Microbiol* 2019; **21**: 3627–3637.

27.   Palomo A, Dechesne A, Pedersen AG, Smets BF. Genomic profiling of Nitrospira
species reveals ecological success of comammox Nitrospira. *Microbiome* 2022; **10**.

28.   Sun D, Tang X, Li J, Liu M, Hou L, Yin G, et al. Chlorate as a comammox Nitrospira
specific inhibitor reveals nitrification and N2O production activity in coastal wetland. *Soil*
Biol Biochem 2022; **173**: 108782.

29.   Wang DQ, Zhou CH, Nie M, Gu JD, Quan ZX. Abundance and niche specificity of
different types of complete ammonia oxidizers (comammox) in salt marshes covered
by different plants. *Sci Total Environ* 2021; **768**: 144993.

30.   Wegner CE, Gaspar M, Geesink P, Herrmann M, Marz M, Küsel K. Biogeochemical
regimes in shallow aquifers reflect the metabolic coupling of the elements nitrogen,
sulfur, and carbon. *Appl Environ Microbiol* 2019; **85**: e02346-18.

31.   Herrmann M, Rusznyák A, Akob DM, Schulze I, Opitz S, Totsche KU, et al. Large
Fractions of CO 2 -Fixing Microorganisms in Pristine Limestone Aquifers Appear To Be
Involved in the Oxidation of Reduced Sulfur and Nitrogen Compounds. *Appl Environ*
Microbiol 2015; **81**: 2384–2394.

32.   Lehmann R, Totsche KU. Multi-directional flow dynamics shape groundwater quality in
sloping bedrock strata. *J Hydrol* 2020; **580**: 124291.

33.   Kohlhepp B, Lehmann R, Seeber P, Küsel K, Trumbore SE, Totsche KU. Aquifer
configuration and geostructural links control the groundwater quality in thin-bedded
carbonate-siliciclastic alternations of the Hainich CZE, central Germany. *Hydrol Earth*
Syst Sci 2017; **21**: 6091–6116.

34.   Küsel K, Totsche KU, Trumbore SE, Lehmann R, Steinhäuser C, Herrmann M. How
deep can surface signals be traced in the Critical Zone? Merging biodiversity with
biogeochemistry research in a central German Muschelkalk landscape. *Front Earth Sci*
2016; **4**: 32.

35.   Scheiner D. A modified version of the sodium salicylate method for analysis of
wastewater nitrates. *Water Res* 1974; **8**: 835–840.

36.   Seelmeyer G. Deutsche Einheitsverfahren zur Wasseruntersuchung. Herausgegeben
im Auftrage der Fachgruppe Wasserchemie in der Gesellschaft Deutscher Chemiker E.
V. Bearbeitet von L. W. Haase, H. Stooff, G. Gad, W. Wesly u.a. Verlag Chemie,
Weinheim/Bergstraße 1954, 180. *Mater Corros* 1954; **5**: 357–357.

37.   Shen T, Stieglmeier M, Dai J, Urich T, Schleper C. Responses of the terrestrial
ammonia-oxidizing archaeon Ca. Nitrososphaera viennensis and the ammonia-
oxidizing bacterium Nitrosospira multiformis to nitrification inhibitors. *FEMS Microbiol*
Lett 2013; **344**: 121–129.

38.   Herrmann M, Hädrich A, Küsel K. Predominance of thaumarchaeal ammonia oxidizer
abundance and transcriptional activity in an acidic fen. *Environ Microbiol* 2012; **14**:
3013–3025.

39.   Rotthauwe JH, Witzel KP, Liesack W. The ammonia monooxygenase structural gene
amoA as a functional marker: molecular fine-scale analysis of natural ammonia-
oxidizing populations. *Appl Environ Microbiol* 1997; **63**: 4704–12.

40.   Francis CA, Roberts KJ, Beman JM, Santoro AE, Oakley BB. Ubiquity and diversity of
ammonia-oxidizing archaea in water columns and sediments of the ocean. *Proc Natl*
Acad Sci 2005; **102**: 14683–14688.

41.   Pester M, Maixner F, Berry D, Rattei T, Koch H, Lücker S, et al. NxrB encoding the beta
subunit of nitrite oxidoreductase as functional and phylogenetic marker for nitrite-
oxidizing Nitrospira. *Environ Microbiol* 2014; **16**: 3055–3071.

42.   Alves RJE, Minh BQ, Urich T, Von Haeseler A, Schleper C. Unifying the global
phylogeny and environmental distribution of ammonia-oxidising archaea based on
amoA genes. *Nat Commun* 2018; **9**: 1517.

43.   Tourna M, Freitag TE, Nicol GW, Prosser JI. Growth, activity and temperature
responses of ammonia-oxidizing archaea and bacteria in soil microcosms. *Environ*
Microbiol 2008; **10**: 1357–1364.

44.   Callahan BJ, McMurdie PJ, Rosen MJ, Han AW, Johnson AJA, Holmes SP. DADA2:
High-resolution sample inference from Illumina amplicon data. *Nat Methods* 2016; **13**:
581–583.

45.   Schloss PD, Westcott SL, Ryabin T, Hall JR, Hartmann M, Hollister EB, et al.
Introducing mothur: open-source, platform-independent, community-supported
software for describing and comparing microbial communities. *Appl Environ Microbiol*
2009; **75**: 7537–7541.

46.   Purkhold U, Wagner M, Timmermann G, Pommerening-Röser A, Koops HP. 16S rRNA

and *amoA*-based phylogeny of 12 novel betaproteobacterial ammonia-oxidizing
isolates: Extension of the dataset and proposal of a new lineage within the
nitrosomonads. *Int J Syst Evol Microbiol* 2003; **53**: 1485–1494.

47. Krüger M, Potthast K, Michalzik B, Tischer A, Küsel K, Deckner FFK, et al. Drought and
rewetting events enhance nitrate leaching and seepage-mediated translocation of
microbes from beech forest soils. *Soil Biol Biochem* 2021; **154**: 108153.

48. Quast C, Pruesse E, Yilmaz P, Gerken J, Schweer T, Yarza P, et al. The SILVA
ribosomal RNA gene database project: improved data processing and web-based tools.
Nucleic Acids Res 2013; **41**: 590–596.

49. McMurdie PJ, Holmes S. Phyloseq: an R package for reproducible interactive analysis
and graphics of microbiome census data. *PLoS One* 2013; **8**: e61217.

50. Chaudhari NM, Overholt WA, Figueroa-Gonzalez PA, Taubert M, Bornemann TL V.,
Probst AJ, et al. The economical lifestyle of CPR bacteria in groundwater allows little
preference for environmental drivers. *Environ Microbiome* 2021; **16**.

51. Fish JA, Chai B, Wang Q, Sun Y, Brown CT, Tiedje JM, et al. FunGene: The functional
gene pipeline and repository. *Front Microbiol* 2013; **4**: 291.

52. Edgar RC. MUSCLE: Multiple sequence alignment with high accuracy and high
throughput. *Nucleic Acids Res* 2004; **32**: 1792–1797.

53. Letunic I, Bork P. Interactive tree of life (iTOL) v5: An online tool for phylogenetic tree
display and annotation. *Nucleic Acids Res* 2021; **49**: W293–W296.

54. Taubert M, Overholt WA, Heinze BM, Matanfack GA, Houhou R, Jehmlich N, et al.
Bolstering fitness via CO2 fixation and organic carbon uptake: mixotrophs in modern
groundwater. *ISME J* 2021; **16**: 1153–1162.

55. Taubert M, Stöckel S, Geesink P, Girnus S, Jehmlich N, von Bergen M, et al. Tracking
active groundwater microbes with D2O labelling to understand their ecosystem
function. *Environ Microbiol* 2018; **20**: 369–384.

56. Team RC. A Language and Environment for Statistical Computing. *R Found Stat*
Comput . 2020. R Foundation for Statistical Computing, Vienna, Austria. , **2**:
https://www.R--project.org

57. Ogle D, Wheeler P, Dinno A. FSA: Fisheries Stock Analysis. R package version 0.8.31.
2020.

58. Gu Z, Eils R, Schlesner M. Complex heatmaps reveal patterns and correlations in
multidimensional genomic data. *Bioinformatics* 2016; **32**: 2847–2849.

59. Oksanen J, Kindt R, Legendre P, O'Hara B, Simpson GL, Solymos PM, et al. The vegan
package. *Community Ecol Packag* 2008; 190.

60. Bédard C, Knowles R. Physiology, biochemistry, and specific inhibitors of CH4, NH4+,
and CO oxidation by methanotrophs and nitrifiers. *Microbiol Rev* 1989; **53**: 68–84.

61. Small GE, Bullerjahn GS, Sterner RW, Beall BFN, Brovold S, Finlay JC, et al. Rates
and controls of nitrification in a large oligotrophic lake. *Limnol Oceanogr* 2013; **58**: 276–
286.

62. Newell SE, Babbin AR, Jayakumar A, Ward BB. Ammonia oxidation rates and
nitrification in the Arabian Sea. *Global Biogeochem Cycles* 2011; **25**.

63. Beman JM, Chow CE, King AL, Feng Y, Fuhrman JA, Andersson A, et al. Global
declines in oceanic nitrification rates as a consequence of ocean acidification. *Proc Natl*

Acad Sci U S A 2011; **108**: 208–213.

64.   Bristow LA, Dalsgaard T, Tiano L, Mills DB, Bertagnolli AD, Wright JJ, et al. Ammonium
and nitrite oxidation at nanomolar oxygen concentrations in oxygen minimum zone
waters. *Proc Natl Acad Sci* 2016; **113**: 10601–10606.

65.   Gribsholt B, Boschker HTS, Struyf E, Andersson M, Tramper A, De Brabandere L, et
al. Nitrogen processing in a tidal freshwater marsh: A whole-ecosystem 15N labeling
study. *Limnol Oceanogr* 2005; **50**: 1945–1959.

66.   Miranda J, Balachandran KK, Ramesh R, Wafar M. Nitrification in Kochi backwaters.
Estuar Coast Shelf Sci 2008; **78**: 291–300.

67.   Kostan J, Sjöblom B, Maixner F, Mlynek G, Furtmüller PG, Obinger C, et al. Structural
and functional characterisation of the chlorite dismutase from the nitrite-oxidizing
bacterium ' Candidatus Nitrospira defluvii': Identification of a catalytically important
amino acid residue. *J Struct Biol* 2010; **172**: 331–342.

68.   Zhao J, Bello MO, Meng Y, Prosser JI, Gubry-Rangin C. Selective inhibition of ammonia
oxidising archaea by simvastatin stimulates growth of ammonia oxidising bacteria. *Soil
Biol Biochem* 2020; **141**: 107673.

69.   Vilardi KJ, Cotto I, Sevillano M, Dai Z, Anderson CL, Pinto A. Comammox Nitrospira
bacteria outnumber canonical nitrifiers irrespective of electron donor mode and
availability in biofiltration systems. *FEMS Microbiol Ecol* 2022; **98**.

70.   Mosley OE, Gios E, Close M, Weaver L, Daughney C, Handley KM. Nitrogen cycling
and microbial cooperation in the terrestrial subsurface. *ISME J* 2022; **16**: 2561–2573.

71.   Sakoula D, Koch H, Frank J, Jetten MS, van Kessel MA, Lücker S. Enrichment and
physiological characterization of a novel comammox Nitrospira indicates ammonium
inhibition of complete nitrification. *ISME J* 2020; **15**: 1010–1024.

72.   Tatari K, Gülay A, Thamdrup B, Albrechtsen HJ, Smets BF. Challenges in using
allylthiourea and chlorate as specific nitrification inhibitors. *Chemosphere* 2017; **182**:
301–305.

73.   Cardarelli EL, Bargar JR, Francis CA. Diverse Thaumarchaeota Dominate Subsurface
Ammonia-oxidizing Communities in Semi-arid Floodplains in the Western United
States. *Microb Ecol* 2020; **80**: 778–792.

74.   Zhu G, Wang X, Wang S, Yu L, Armanbek G, Yu J, et al. Towards a more labor-saving
way in microbial ammonium oxidation: A review on complete ammonia oxidization
(comammox). *Sci Total Environ* . 2022. Elsevier. , **829**: 154590

75.   Weidinger K, Neuhäuser B, Gilch S, Ludewig U, Meyer O, Schmidt I. Functional and
physiological evidence for a Rhesus-type ammonia transporter in Nitrosomonas
europaea. *FEMS Microbiol Lett* 2007; **273**: 260–267.

76.   Perez-Riverol Y, Csordas A, Bai J, Bernal-Llinares M, Hewapathirana S, Kundu DJ, et
al. The PRIDE database and related tools and resources in 2019: Improving support
for quantification data. *Nucleic Acids Res* 2019; **47**: D442–D450.

Data availability

Sequence data generated in this study was submitted to the European Nucleotide Sequence Archive with project number PRJEB57321 for *amoA* genes under accessions ERS13665998 to ERS13665900, and with project number PRJEB63087 for bacterial 16S rRNA genes under accessions ERS15585018 to ERS15584991.

The mass spectrometry proteomics data have been deposited to the ProteomeXchange Consortium via the PRIDE [76] partner repository with the dataset identifier PXD039573.

Datasets containing the calculation of the groundwater nitrification rates, gene abundances, hydrochemical measurements and *amoA* OTU sequences will be released upon publication of the manuscript.

Acknowledgements

This study was part of the Collaborative Research Centre 1076 AquaDiva (CRC AquaDiva) of the Friedrich Schiller University Jena, funded by the Deutsche Forschungsgemeinschaft under project number 218627073. Climate chambers to conduct experiments under controlled temperature conditions and the infrastructure for high-throughput sequencing were financially supported by the Thüringer Ministerium für Wirtschaft, Wissenschaft und Digitale Gesellschaft (TMWWDG; project B 715-09075 and project 2016 FGI 0024 "BIODIV"). Martin Taubert gratefully acknowledges funding by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) under Germany´s Excellence Strategy – EXC 2051 – Project-ID 390713860.

We thank Heiko Minkmar, Falko Gutmann, Jens Wurlitzer, Stefan Krümmling, Swantantar Kumar, Patricia Geesink, Lena Carstens and Linda Gorniak for support of field and laboratory work and Gundula Rudolph for ion chromatography measurements. We acknowledge Kai Uwe Totsche and Robert Lehmann for field site management and provision of physicochemical data from groundwater.

Author contributions

701   MH and MK designed the study. MK conducted all molecular work, incubation experiments

702   and data analysis. BT and LB performed IRMS measurements and analysis. NC and WO

703   performed metagenomic data analysis. MT, NJ and MvB conducted metaproteomic data

704   analysis. MK and MH wrote the manuscript with contributions of all authors.

706   Declaration of competing interest

The authors declare that there was no financial or personal conflict of interest in the

preparation of this study.

Figures

Figure 1

Figure 1: **A)** Schematic cross-section of the Hainich groundwater monitoring transect showing two
superimposed aquifer assemblages accessible by 10 wells with various depths. **B)** Overview of the
concentrations of dissolved oxygen, ammonium, and nitrate from oxic (blue) and hypoxic/ anoxic
(green) groundwater wells (n = 24 per well). Single dots represent one measurement, the crossbar
indicates the mean of all values, and the error bars show the variation from the mean. Outliers are
beyond the error bars. Lowercase letter code indicates significant differences based on Dunn's multiple
comparison test ($p < 0.05$). **C)** Nitrification rates assessed by ^{15}N based labeling of groundwater from
five days incubation close to in situ conditions (n ≥ 3 per well; n.d. = not detected; n.a. = not available).
D) Gene abundances of *amoA* per L and **E)** fractions of *amoA* genes in groundwater from different
groups of ammonia-oxidizing prokaryotes in 10 groundwater wells (n ≥ 3).

Figure 2

Figure 2: **A)** Redundancy analysis (RDA) demonstrates association between distribution of ammonia-
oxidizing community (relative *amoA* read abundances of CMX, AOB and AOA OTUs which account for
< 2% sequence reads of each community were normalized to *amoA* gene abundances to generate
absolute *amoA* read abundances) and significant hydrochemical parameters fitting (*p* < 0.05) with
nitrifier community distribution across eight groundwater wells (n ≥ 3). Dotted ellipses show the
confidence of clusters assuming a multivariant normal distribution. **B)** Community composition of CMX
Nitrospira showing dominant OTUs (< 2% of sequence reads) affiliated to clade A and clade B across
eight groundwater wells (n ≥ 3 per well). Each bar displays the mean absolute *amoA* read abundance
of OTUs. **C)** Maximum likelihood phylogenetic tree of deduced AmoA amino acid sequences of CMX
Nitrospira depicting evolutionary relationship between reference genomes, and sequences of OTUs
and MAGs from Hainich groundwater. The tree was constructed using the JTT substitution model with
gamma distribution and 1000 bootstrap iterations. Bootstrap support values greater than 75% are
denoted with black dots. An AmoA sequence from *Nitrosomonas oligotropha* served as outgroup
indicated by the arrow. **D)** Pairwise correlations between total *amoA* gene abundances of CMX, AOB
and AOA, absolute *amoA* read abundances of OTUs and groundwater hydrochemical parameters (TIC
= total inorganic carbon, TOC = total organic carbon, DOC = dissolved organic carbon, ORP = redox
potential, EC = electrical conductivity) using Spearman's rank correlation. Hierarchical clustering is
based on Euclidean distance. Color code of correlation coefficient r$_s$ displays strength of association
using following significance levels: * *p* < 0.05, ** *p* < 0.01, *** *p* < 0.001.

Figure 3

749   0.5

Figure 3: **A)** Maximum likelihood tree of bacterial core protein sequences of CMX *Nitrospira* from nine
MAGs originating from Hainich groundwater depicting the evolutionary relationship to reference
genomes. The tree was constructed using the JTT substitution model with gamma distribution and 1000
bootstrap iterations. Bootstrap support values greater than 75% are shown as black dots. AmoA
sequences from *Nitrosomonas oligotropha* and *Nitrosomonas europaea* served as an outgroup
indicated by the arrow. **B)** Metabolic capacities of nine groundwater CMX *Nitrospira* MAGs inferring
genomic potential for nitrogen and sulfur energy metabolism, and corresponding transporter types as
well as protective systems against radical oxygen species, and genes involved in carbon fixation (rTCA
= reductive tricarboxylic acid cycle). Sizes of circles display the transcript coverage of each gene
mapped from groundwater metatranscriptome. If more than one gene was present, maximum
transcriptional coverage is shown. Filled circles indicate peptides from metaproteomic data matching
for displayed genes. rTCA = reverse tricarboxylic acid cycle.

Figure 4

Figure 4: Groundwater incubations from wells H41 and H53 supplemented with different NH_4^+ concentrations and nitrification inhibitors across 5 days (short-term) and 25 to 35 days (long-term). **A)** Nitrification rates from short-term incubation of oxic groundwater supplemented with 50 µM NH_4^+ and selective inhibition with chlorate and allylthiourea (n.d. = not detected). Dots depict single rates of each replicate. Crossbar shows the average of three replicates and error bars show the variation between rates. **B)** Fold-change of *amoA* genes from ammonia oxidizers and *nxrB* genes from *Nitrospira*-like nitrite oxidizers after long-term incubation of groundwater. Single points represent one replicate and crossbar shows the average. Error bars display variance between conditions (n = 3). Condition 1: oxic groundwater supplemented with 10 µM NH_4^+. Condition 2: oxic groundwater supplemented with 50 µM NH_4^+. Condition 3: hypoxic groundwater without additional NH_4^+. **C)** Composition of CMX *Nitrospira*

775   clade A and clade B affiliated OTUs of groundwater and after long-term incubation (n = 3). Bars show
776   the mean relative *amoA* read abundance of OTUs representing > 2%. **D)** Composition of bacterial
777   ammonia-oxidizing community and taxonomic affiliation of OTUs.