DANGER ANALYSIS: RISK-AVERSE ON/OFF-TARGET ASSESSMENT FOR CRISPR EDITING WITHOUT A REFERENCE GENOME

Kazuki Nakamae[1,2,*] and Hidemasa Bono[1,3,*]

[1] Genome Editing Innovation Center, Hiroshima University, Hiroshima 739-0046, Japan
[2] PtBio Inc., Hiroshima 739-0046, Japan
[3] Graduate School of Integrated Sciences for Life, Hiroshima University, Hiroshima 739-0046, Japan

* Correspondence should be addressed to H.B. bonohu@hiroshima-u.ac.jp. Correspondence may also be addressed to K.N. kazuki-nakamae@hiroshima-u.ac.jp.

Kazuki Nakamae, Ph.D.
Genome Editing Innovation Center, Hiroshima University,
3-10-23 Kagamiyama, Higashi-Hiroshima, Hiroshima 739-0046, Japan
E-mail: kazuki-nakamae@hiroshima-u.ac.jp
Tel: +81-82-424-4013
Fax: +81-82-424- 3990

Hidemasa Bono, Ph.D.
Graduate School of Integrated Sciences for Life, Hiroshima University,
3-10-23 Kagamiyama, Higashi-Hiroshima city, Hiroshima 739-0046, Japan
E-mail: bonohu@hiroshima-u.ac.jp
Tel.: +81-82-424-4013

ABSTRACT

The CRISPR-Cas9 system has successfully achieved site-specific gene editing in organisms

ranging from humans to bacteria. The technology efficiently generates mutants, allowing for

phenotypic analysis of the on-target gene. However, some conventional studies did not

investigate whether deleterious off-target effects partially affect the phenotype. Herein, we

present a novel phenotypic assessment of CRISPR-mediated gene editing: Deleterious and

ANticipatable Guides Evaluated by RNA-sequencing (DANGER) analysis. Using RNA-seq data,

this bioinformatics pipeline can elucidate genomic on/off-target sites on mRNA-transcribed

regions related to expression changes and then quantify phenotypic risk at the gene

ontology (GO) term level. We demonstrated the risk-averse on/off-target assessment in

RNA-seq data from gene-edited samples of human cells and zebrafish brains. Our DANGER

analysis successfully detected off-target sites, and it quantitatively evaluated the potential

contribution of deleterious off-targets to the transcriptome phenotypes of the edited

mutants. Notably, DANGER analysis harnessed *de novo* transcriptome assembly to perform

risk-averse on/off-target assessments without a reference genome. Thus, our resources

would help assess genome editing in non-model organisms, individual human genomes, and

atypical genomes from diseases and viruses. In conclusion, DANGER analysis facilitates the

safer design of genome editing in all organisms with a transcriptome.

19
20
INTRODUCTION

The CRISPR-Cas9 system was initially adapted as a bacterial immune system[1,2]. Over the

past decade, this system has been developed as a programmable nuclease that enables site-

specific modification of the genomes of various organisms, including humans)[3–5]), insects)[6,7]),

microalgae([8,9]), and bacteria([10]). Engineered CRISPR-Cas9 undertakes genomic modification

using two components: RNA-guided Cas9 nuclease and single-guide RNA (sgRNA)([11,12]). The

Cas9-sgRNA complex generates indels near the target site (on-target site), where the 19–20

bases of the 5′ ends of sgRNA (protospacer) and the protospacer adjacent motif (PAM) of

Cas9 protein bind([11–14]). Recently, many CRISPR-Cas9 applications, such as Cas9 nickase

(Cas9n)([12]), dead Cas9 (dCas9)([15]), base editors([16,17]), and prime editors([18]), have been

developed. Furthermore, CRISPR-Cas9-mediated genome editing was found to be efficient,

with the editing efficiency exceeding 50% over time[19]. Thus, CRISPR technology has

dramatically facilitated a reverse genetics approach involving phenotypic analysis using

CRISPR-Cas9-based mutants of a user-targeted gene[20–23].

However, genome editing using CRISPR technology presents two challenges that have not

been addressed in previous studies. First, phenotypic effects caused by unexpected CRISPR

dynamics are not quantitively monitored. CRISPR-Cas9 is well known for unexpected

sequence editing (off-target site) with mismatches when compared to protospacers and

PAM[5]. Off-target gene editing results in incorrectly edited mRNA, unexpected phenotypes,

and decreased expression of unrelated genes. Some reports predicted and detected off-

target editing using genomic PCR and DNA sequencing analysis[14,24–26], but most studies

have not assessed the phenotypic effect of the detected off-targets. Second, CRISPR

technology generally depends on basic genomic data, including the reference genome.

CRISPR technology has potential applications in organisms with incompletely characterized

genomes. However, the design of site-specific sgRNAs requires the factual genomic

sequence of materials to be treated with CRISPR technology. This hindrance also emerges in

the human genome, particularly in the genomes of patients and cancer genomes. These

genomes are assumed to be completely distinct from the reference genome[27,28]. The off-

target is always "unexpected." Thus, we need a method to observe factual genomic

sequences and reduce potential off-target effects.

We devised a method to overcome the two challenges above: phenotypic risk and

dependence on a reference genome. Phenotypic risk can be assessed by phenotype analysis

using gene ontology (GO) annotation[29,30]. GO has been widely used for several

decades[20,23,31–34]. Recently, many RNA sequencing (RNA-seq) data and mapped genes have

been annotated with GO terms to characterize the transcriptome phenotype under a specific

condition of the organism. This process is known as enrichment analysis[35]. We expected

that we could quantitatively assess the phenotypic risk of off-target genes if each off-target

gene with decreased expression was annotated with GO terms. Moreover, *de novo*

transcriptome assembly technology can address the dependency problem of reference

genomes. The *de novo* transcriptome assembly can generate transcriptome sequences

without the reference genome using RNA-seq data[36–40]. We identified factual genomic

sequences in mRNA-transcribed regions using *de novo* transcriptome assembly from gene-

edited organisms and cells.

In this study, we combined *de novo* transcriptome assembly and GO annotation analysis in

CRISPR editing to establish a DNA on/off-target assessment, including phenotype risk

analysis without a reference genome. We named it **D**eleterious and **AN**ticipatable **G**uides

Evaluated by **R**NA-sequencing (**DANGER**) analysis (Figure 1). This bioinformatics pipeline can

elucidate genomic on/off-target sites based on de novo transcriptome assembly using RNA-

seq data. Then, it identifies the *deleterious off-targets*, defined as off-targets on the mRNA-

transcribed regions that represent the downregulation of expression in edited samples

compared to wild-type (WT) ones. Furthermore, our pipeline can quantify phenotypic risk at

the GO term level by calculating a newly defined indicator of phenotypic risk by the

deleterious off-targets, named the D-index.

MATERIAL AND METHODS

Implementation of DANGER Analysis

Our pipeline of DANGER analysis is composed of several processes: "Quality control &

Adapter trimming," "rRNA Removal," "de novo transcriptome assembly," " Removal of

redundancy," "Detection of on-target and potential off-target sites," "Expression

quantification," "Search for deleterious off-target sites," "Identification of ORFs and Genes,"

"GO analysis," and "Validation for phenotypic risk" (Figure 2A).

DANGER analysis examines paired-end RNA-seq data derived from wild-type (WT) and

edited samples using the processes depicted in Figure 1. The pipeline generates a *de novo*

transcriptome assembly, an expression profile of transcripts belonging to on/off-target sites,

and an estimation of the phenotypic risk for off-target sites. The script has been uploaded to

our GitHub repository (https://github.com/KazukiNakamae/DANGER_analysis). The analyses

were performed on Docker with Ubuntu v. 22.04.1, LTS, and 235 GB of memory. The scripts

for this processing pipeline were released as a Docker image

(https://hub.docker.com/r/kazukinakamae/dangeranalysis), enabling operation on various

operating systems beyond Linux using this Docker image. Each process is explained in detail

below.

Quality Control & Adapter Trimming. Quality control and adapter trimming were performed

using Cutadapt v. 1.18([41]), which also removed low-quality reads. The adapter sequences

used were "AGATCGGAAGAG."

Ribosomal RNA (rRNA) Removal. The residual rRNA reads were filtered using bbduk v. 38.18

(https://sourceforge.net/projects/bbmap/). Each sample was filtered twice. In the first and

second filters, we used SSU and LSU rRNA datasets from SILVA v. 119.1 (https://www.arb-

silva.de). The dataset was downloaded from the CRISPRroots

(https://rth.dk/resources/crispr/crisprroots/).

De novo Transcriptome Assembly. The *de novo* transcriptome assembly was performed using

Trinity v. 2.12.0([38]). The merged read files were composed of RNA-seq data derived from WT

samples. Transcriptome completeness was assessed using BUSCO v. 5.2.2_cv1([42]). The BUSCO

evaluated the competence of assembly using estimation of similarity to gene database

(BUSCO genes) and classified hit sequence into "complete" (including "single-copy" and

"duplicated"), "fragmented," and "missing." The databases used for conserved mammalian

BUSCO genes were "mammalia_odb10" and "actinopterygii_odb10" for human and zebrafish

assemblies, respectively.

Removal of redundancy. The expression of RNA-seq data derived from WT samples was

quantified in advance to remove transcripts with low expression levels. Quantification was

performed with align_and_estimate_abundance.pl of Trinity v. 2.12.0. The removal of

transcripts was performed with filter_low_expr_transcripts.pl of Trinity v. 2.12.0.

Detection of on-target and potential off-target sites. Detection of on/off-target sites in *de*

novo transcriptome assembly was performed with Crisflash v.1.2.0([43]). Our off-target

detection focused on up to 8 or up to 11nt mismatches and NGG or NRR PAM because the

previous off-target reports indicated that off-target sites with ≥5 nt mismatches and

NGG/NAG/NGA/NAA PAM exist[24,44], although not with a high frequency. Specifically, we

searched for potential off-target sites by executing the following command:

"crisflash -g Result_denovo_transcriptome_ {*de novo* transcriptome assembly generated from

WT RNA-seq data} -s {sequence of protospacer and PAM} -o {output file of Crisflash} -m {the

maximum number of mismaches users consider} -p {PAM} -t {the number of thread} -C."

The output file of Crisflash has on-/off-target locations, off-target sequences, and mismatch

numbers included in the tab-delimited format (Cas-OFFinder format). The on-target

locations were extracted using perfect matching with the expected genomic sequence of

Cas9-sgRNA binding. The potential off-target sites were classified by the mismatch number

in each text file.

Expression quantification. Using filtered *de novo* transcriptome assembly, we quantified the

expression of RNA-seq data derived from each sample. Quantification was performed using

align_and_estimate_abundance.pl of Trinity v. 2.12.0. The transcripts per million (TPM)

dataset was constructed from "RSEM.isoforms.results" of each output directory and was

saved as a single CSV file.

The ratio of TPM values on a transcript was calculated with the following formula:

22
$$(TPM\ ratio) = \frac{(Average\ of\ TPM\ of\ Edited\ samples)}{(Average\ of\ TPM\ of\ WT\ samples)} \qquad (1)$$

A transcript with a TPM ratio of less than the user-defined value (t) was determined as

downregulated TPM (dTPM). Furthermore, in place of TPM, it is compatible with profiling

based on Differentially Expressed Genes (DEG). The DEG analysis was performed to compare

different analysis methods. The read count data were extracted from the

"RSEM.isoforms.results" of each output directory in the section "Implementation of DANGER

analysis: Expression quantification." The raw count data were normalized by the Tag Count

Comparison (TCC) R package[45] with the parameter "norm.method="tmm,"

test.method="edger," iteration=30, FDR=0.1, floorPDEG=0.05" to detect DEG between RNA-

seq data derived from WT and Edited samples. The MA plot was constructed using an in-

house R script. If a DEG transcript had a negative log-ratio of normalized counts (M-value)

and its p-value fell below the user-defined value (α), it was determined to be downregulated

DE (dDE). It was saved as a single CSV file.

Search for deleterious off-target sites. Our pipeline defined an off-target site where the

transcript was annotated with dTPM or dDE as a deleterious off-target site, which could be

also paraphrased as "actual off-target site." We counted the number of deleterious off-target

sites using an in-house Python script, which required the off-target site profile and the TPM

ratio or DEG described above.

Identification of ORFs and Genes. Our pipeline identifies open reading frames (ORFs) in the

filtered de novo transcript sequences and predict the corresponding amino acid sequences.

If these predicted sequences exhibit significant homology with protein sequences in a

database, we assign them as genes. The open reading frames (ORFs) and genes of the

filtered *de novo* transcript were estimated with TransDecoder v. 5.5.0 and ggsearch v. 36.3.8g

using a protein database. The process was based on the Systematic Analysis for

Quantification of Everything (SAQE) pipeline (https://github.com/bonohu/SAQE)[33]. In

particular, "11TransDecoder.sh", "12GetRefProts.sh", "15ggsearch.sh", "15parseggsearch.sh",

and "15mkannotbl.pl" were used with the supplemental script "00_prepare_faa_4Fanflow. sh"

in the GitHub repository (https://github.com/RyoNozu/Sequence_editor). The referred

protein databases were the Ensembl databases of all translations resulting from Ensembl

genes in humans and zebrafish. The DANGER analysis database can be manually customized

with the organism from which the analyzed RNA-seq data was extracted.

GO enrichment analysis. GO annotations of gene ontologies were performed using an in-

house R script using the org.Hs.eg.db and org.Dr.eg.db R packages for humans and zebrafish,

respectively. Enrichment analysis followed by GO annotations was performed using the

topGO R package against genes whose off-target sites were determined to be deleterious

off-target sites. Enrichment analyses were performed per off-target mismatch number.

Finally, the enrichment tables were merged into a single table, named the DANGER table,

with the mismatch number annotated.

Validation for phenotypic risk. We defined the following value (D-index) to evaluate the

phenotypic risk posed by deleterious off-target effects per GO term.

$$(D - index) = \sum_{m=0}^{m_{max}} N(m) \times exp(4 - m) \quad (2)$$

where m indicates the number of mismatches. The N(m) represents the total number of

genes , which have m bases mismatches, included a specific GO term. The m_{max} is the

maximum number of mismatches that a user considers. The D-index considers both the

phenotypic risk and the frequency of off-target effects. Based on previous reports, we used

the exponential function to express the frequency of off-targets because the frequency of

off-targets tends to decrease exponentially as the mismatch number of off-target sites

increases([24,46]). Based on a GUIDE-seq study[24], most reported off-targets possess mismatches

of four or fewer nucleotides (Supplementary Table S1). Therefore, the exponent value is

represented as a decreasing function by subtracting the number of mismatches from four,

and by minimizing the impact of mismatches with five or more nucleotides, we have enabled

the probabilistic risk assessment at an appropriate level. Calculations were performed using

an in-house Python script.

Validation of D-index

We established a validation methodology of statistical significance for the D-index,

determined as described above, for each set of GO terms using a permutation test. The

permutation test involved random shuffling of the expression profile and off-target profile based on a given seed value, and the D-index was computed based on these randomly shuffled profile data. We will refer to this D-index as a pseudo-D-index. After repeating this process of creating pseudo-D-indexes 100 times, we made a distribution of pseudo-D-indices for each set of GO terms. A D-index outside the $(1 − L) \times 100$ % confidence interval of this distribution and higher than the mean value was defined as a "significant D-index." Additionally, we implemented a script to measure the false positive rate of the permutation test. In false-positive detection, ten additional shuffled data sets were generated using seed values different from the ones used to create the shuffled expression and mismatch profiles, and the newly calculated pseudo-D indices from these data were calculated. If they met the criteria for a significant D-index in the above distribution, the pseudo-D-indices were defined as "significant pseudo-D-index" and counted. The ratio of the significant pseudo-D-indices to the total number of new pseudo-D-indices was defined as the false positive rate. Multiplying this false-positive rate by the total number of original D-indices allows us to estimate the number of expected false D-indices, and subtracting this from the total number of original D-indices enables us to count the number of expected true D-indices.

The analyses were performed on Docker with Ubuntu v. 22.04.1, LTS, and 235 GB of memory. The scripts for this processing pipeline were released as a Docker image (https://hub.docker.com/r/kazukinakamae/dangertest), enabling operation on various operating systems beyond Linux by using this Docker image.

Datasets

Two RNA-seq datasets were analyzed to evaluate our pipeline. First, we collected paired-end RNA-seq datasets, which had a total of >100 M reads for all WT samples and an average length of >100 nt, to ensure good quality, which indicated a percentage of complete benchmarking universal single-copy orthologs (BUSCO) genes of >70%, of transcriptome completeness after *de novo* transcriptome assembly. The datasets were downloaded from the Sequence Read Archive (SRA) (Table 1).

GRIN2B. The dataset was extracted from human iPSC-derived cortical neurons with or without indels generated by paired Cas9 nickase (Cas9n)-single-guide RNA (sgRNA) (GRIN2B-FW and GRIN2B-REV sgRNAs) on the GRIN2B locus[23]. Previous studies have established clones with indels resulting in loss-of-function (LOF) and reduced dosage (RD). However, we focused only on LOF samples that had been edited and analyzed using our pipeline. Gorodkin and Seemann have previously reported that off-target sites affect the expression profile of LOF samples using reference-based RNA-seq analysis. Our study used the GRIN2B dataset to benchmark *de novo* transcriptome assembly based and reference-based RNA-seq analyses[44]. Moreover, we profiled the on/off-target assessment of GRIN2B-REV sgRNA.

Park7. The dataset was extracted from the zebrafish brain with or without biallelic indels generated by a single Cas9-sgRNA at the park7 locus (which encodes DJ-1)[20]. The analyzed F2 mutants were generated from a cross between two heterozygous F1 mutants. We used the dataset as an *in vivo* example of the DANGER analysis pipeline in a simple CRISPR-Cas9-mediated knock-out experiment.

Statistical analysis

Plots were made using Microsoft Office and housemade Python scripts. The Exact Fisher's test was performed for the p-value was calculated accordingly using *fisher_exact()* of the *scipy* package in Python. The 2-tailed Welch's t-test was performed for the p-value was calculated accordingly using *ttest_ind(equal_var=False)* of the *scipy* package in Python. We used G power software for the statistical power (1-ß) calculation. The Venn diagrams were generated using web software ("Calculate and draw custom Venn diagrams": https://bioinformatics.psb.ugent.be/webtools/Venn/).

RESULTS & DISCUSSION

Assessment of CRISPR-Cas9 off-targets using DANGER Analysis for RNA-seq Data from *in vitro* Differentiated Human iPSC

We investigated whether our DANGER analysis pipelines could detect deleterious off-target

sites without information on the reference genome. First, we applied DANGER analysis to the

GRIN2B dataset, which was extracted from human iPSC-derived cortical neurons with or

without in-frame deletions at the GRIN2B locus[23]. The obtained *de novo* transcriptome

assembly contained five isoforms in which on-target sequences of sgRNA (GRIN2B-REV)

(Figure 2B) were located. The assembly comprised 342,910 contigs and exhibited BUSCO

transcriptome completeness of 79.1% (Figure 2C). Previous studies on *de novo* transcriptome

assembly using Trinity reported 64.7%, 77.1%, 80%, and 87% complete BUSCO genes in

higher animals such as *Homo sapiens*[36], *Castor fiber* L.[37], *Mirounga angustirostris*[39], and

Dromiciops gliroides[40], respectively. We successfully obtained a *de novo* transcriptome with

standard quality. Consequently, the pipeline performed an exhaustive search using Crisflash,

yielding 33,878 potential off-target sites with up to 8 bases mismatches (MM) and NGG PAM,

using transcriptome assembly.

Next, we quantified the transcriptome-wide expression of each of the four RNA-seq samples

from the WT and Edited GRIN2B loci. Our pipeline examines whether the potential off-target

site, which is output of Crisflash, is in the transcript and whether it also reduces the

expression value. The pipeline considers those potential off-targets that have confirmed

down-expression as deleterious off-targets, in other words, actual off-targets. The reduction

of expression may occur because nonsense-mediated mRNA decay (NMD)[47] destroys

incomplete transcript sequences resulting from off-targeting, or alignment software fails to

map the incomplete transcript sequences to the untreated transcript sequence[48]. The

DANGER analysis screens transcripts with lower expression levels in edited samples

compared to WT samples. In general, there are several criteria for estimating expression

using RNA-seq. We implemented two criteria, "downregulated different expression (dDE)"

and "dTPM," for the detection of downregulated transcripts (Figure 3A; two callouts). Our

dDE criterion uses DEG analysis with TCC normalization (see Materials and Methods). In the

case of DEs in a transcript with a negative M-value and a p-value that was less than the

threshold value (α) in the MA plot (Figure 3A; right callout), we defined the transcript as

"dDE." On the other hand, the "TPM" criterion uses the normalized value, named TPM, as first

defined by [49] and calculates the ratio of TPM between WT and the edited samples. When

the ratio of a transcript is less than the threshold value (t), we defined the transcript as

"dTPM" (Figure 3A; left callout). We confirmed the number of transcripts with dDE (α = 0.001)

or dTPM (t = 0.4) annotation that contained off-target sites (Figure 3A; Venn diagram). There

were 730 transcripts with dDE off-targets. A total of 12,747 transcripts with dTPM off-targets

were detected, which was approximately 17-fold more than those with dDE off-targets. Our

DANGER analysis aims to serve as a screening tool that emphasizes maximizing the

estimation of potential risks by capturing as many sites suspected of phenomena as

knockout or knockdown of off-target genes. From this perspective, the dTPM approach can

estimate the deleterious off-target effects to the greatest extent, more so than the dDE

approach. Furthermore, we focused on the off-targets detected in common by dTPM and

dDE. We observed their mismatch count (Supplementary Figure S1A). As a result, we

confirmed that the off-targets seen in common were distributed at an even ratio for each

mismatch count compared to the off-targets detected only by dTPM (Supplementary Figure

S1B). Given that the number of mismatches affects the likelihood of off-target occurrences,

there is no correlation between the common off-targets and their occurrence rate. While

there are differences in the number and types of transcripts detected by dTPM and dDE, no

evidence suggests that these differences result from a flaw in either approach. Therefore, our

pipeline has adopted both methods.

The Gorodkin and Seemann group previously established the on/off-target assessment

pipeline (CRISPRroots)[44]. They then used STAR[50] to perform expression analysis on the

same RNA-seq data of GRIN2B using reference-based mapping. They suggested that two

off-target sites (ALK: gcAgaCTGGtTGGAAGCaCCNGG, GBA2: cccTTCcGGccGGAAGCGCCNGG)

were binding with the Cas9-GRIN2B-REV sgRNA (AGATTCTGGGTGGAAGCGCCNGG)-DNA

seed and were linked to downregulated expression (namely, "RISK: CRITICAL"). The definition

of off-targets in their study was similar to our idea of deleterious off-targets. Thus, we

compared our list of deleterious off-targets in dTPM and dDE with the two off-targets

detected using CRISPRroots. As a result, in the DANGER analysis, the off-target of

CRISPRroots in the ALK locus was detected by both dTPM (t = 0.4) and dDE (α = 0.001)

criteria considering up to eight base mismatches with NGG PAM, whereas the off-target of

CRISPRroots in GBA2 was not detected by any criteria (Figure 3B). The absence of GBA2 was

understandable because the off-target site on the GBA2 locus was in the promoter region

rather than the GBA2 transcript. Additionally, more stringent expression analyses were

conducted using dTPM (t = 0.2) and dDE (α = 0.0001). As a result, although 125 off-target

genes were detected, neither dTPM nor dDE could identify ALK as an off-target gene

(Supplementary Figure S2). This result suggested the possibility of an increase in false

negatives when the threshold for detection of expression decreases resulting from off-target

effects is overly strict. It is therefore considered appropriate to set t = 0.4 in dTPM and α =

0.001 in dDE. Our *de novo* transcriptome approach focuses solely on the potential off-target

genes in the transcribed region of the genome. Additionally, unlike traditional reference-

based RNA-seq analysis, this approach can provide novel insights. For example, the off-

target search of DANGER analysis detected a transcript with an off-target site downstream

GALR2 locus (Figure 3C). The genome database annotation was not the off-target site of the

transcribed region (XM_047436984.1). However, the *de novo* transcriptome assembly

included the site in the transcript (TRINITY_DN86617_c0_g1_i1). This result indicates that our

DANGER analysis pipeline can detect *bona fide* transcripts, which have never been annotated

in the reference genome database because of cell-specific transcription, personal genomic

variants, and inadequate genomic locus study. The detection range of our DANGER analysis

is limited to the transcribed region. However, our pipeline using dTPM and dDE detected

13,237 and 813 off-target sites with zero to eight mismatches in identified and unidentified

transcripts, respectively. 2,236 and 407 gene-annotated off-target sites with four to eight

mismatches, respectively. In contrast, genome-wide and reference-based RNA-seq analysis

(CRISPRroots) features only two off-target sites in genes with six mismatches. Thus, we

demonstrated that the DANGER analysis could generate a comprehensive and factual record

of off-target sites (Figure 3D).

Finally, our pipeline evaluated the phenotypic risk of deleterious off-targets. Various studies

have used Gene Ontology (GO) analysis to assess phenotypic effects[20,23,31–34]. However, this

study also considered off-target frequency because deleterious off-targets with many

mismatches were expected to occur[5,46]. Thus, our pipeline counted off-target genes

associated with specific GO terms per mismatch number (Figure 4A) and then combined the

results with the mismatch effect by calculating the D-index per GO term (Figure 4B; see

Materials and methods). The D-index is calculated by multiplying the number of genes

containing a GO term for each mismatch number by a decreasing exponential function with

the mismatch number as the exponent. This approach allows us to consider the number of

genes hit by the GO terms and the number of mismatches. Moreover, it can suppress the

influence of the number of genes hit when the mismatch number is large. The formula can

prioritize evaluating the number of genes hit when the mismatch number is small. With these

two characteristics, the D-index represents a unique off-target metric that considers the

impact on phenotype. The sum of the D-index value (total D-index) in detected GO terms

was 6,228 (N = 9,896) (Figure 4C, Supplementary Table S2) in the GRIN2B dataset. The

DANGER analysis was used to evaluate the phenotypic risk at the GO level using RNA-seq of

the human GRIN2B dataset data without any reference genomes.

Evaluation of D-index and optimization of DANGER analysis

Using the D-index, we quantified the phenotypic impact from off-targets at the GO term

level. However, a different threshold must be set for each GO term when evaluating the

statistical significance of the D-index values because GO is a loosely hierarchical annotation

concept, and 'parent' terms appear as annotations of various genes even if there is less

relationship with off-targets. To address this issue, we implemented a permutation test

system to estimate the significance threshold for each GO term (Figure 5). In this system,

after randomly shuffling the expression profile and off-target profile, we repeat the process

of calculating a meaningless D-index (named pseudo-D-index) by applying the D-index

formula 100 times. We then create a null distribution from the 100 pseudo-D-index values

for each GO term and define a Significant D-index as an originally obtained D-index value

that exceeds the $(1 - L) \times 100 \%$ confidence interval of the null distribution. The methodology

and the threshold allow us to extract only the D-indices of the GO terms suspected of having

an off-target-associated impact on the phenotype.

Moreover, we devised a scheme to assess the validity of the permutation test system

(Supplementary Figure S3). In the validation scheme, followed by generating several different

pseudo-D indices, the number of the newly generated pseudo-D indices exceeding the

threshold of the previous null distribution is counted. We defined the ratio of the count to

the total number of D-indices as the false detection rate in the permutation test. We used

the evaluation scheme to verify how the false detection rate changes under various

conditions in DANGER Analysis. No significant influence was observed from the threshold of

the expression analysis (t, α) or the conditions of off-target search (Supplementary Figure

S4A-C). We confirmed L=1E-15 confidence interval threshold reduced the false detection

rate by more than half in the case of L=5E-1. Here, we named the condition of dTPM with

high false positives (up to 11-MM NRR PAM, t=0.4, L=5E-1) as 'Approximate dTPM,' the

condition of dTPM with low false positives (up to 8-MM NGG PAM, t=0.4, L=1E-15) as

'Optimized dTPM,' and the condition of dDE with low false positives (up to 8-MM NGG PAM,

α = 0.001, L=1E-15) as 'Optimized dDE.' When comparing the three conditions, the

optimized dDE showed the lowest false detection rate (Figure 6A). Next, we calculated the

total number of D-indices, significant D-indices, and true significant D-indices (Expected True

D-indices) estimated from the False Detection Rate for these three conditions. The total

number of D-indices was more than twice as high for dTPM as for dDE, while the number of

Expected True D-indices was less than half for dTPM compared to dDE (Figure 6B). The result

indicates that the dTPM criterion is a 'sharply-narrowing-down' approach used for initial

screening, while dDE is a 'meticulously trimming' approach used for a rigorous selection

process. Generally, the dDE approach is recommended for use in human RNA-seq data

because the criterion is expected to present fewer false detections in significant D-indices.

However, in model organisms and non-model organisms with less comprehensive GO

annotations than humans, the dDE approach may not yield a sufficient D-index list for the

evaluation. Thus, the dTPM approach, which can obtain more D-indices, is expected to be

more effective in RNA-seq data from less characterized organisms than humans. We

evaluated the consistency of D-indices and Significant D-indices detected by both optimized

dTPM and optimized dDE. The concordance of each D-index and Significant D-index was

approximately 52% and 1.3%, respectively (Figure 6C), which can be attributed to the fact

that optimized dTPM considers more than five times the gene-annotated off-target sites

compared to optimized dDE (Figure 3D). However, the significant D-indices commonly

detected by optimized dTPM and optimized dDE corresponded to the top 16 significant D-

indices in dTPM (Figure 3C-D). The result suggested that the value of the D-index not only

served as an indicator of the phenotypic impact from off-targets but could also be an

indicator of the strength of its consistency. Thus, it is recommended to conduct follow-up

analyses focusing mainly on the top-ranking D-indices in optimized dTPM.

Assessment of CRISPR-Cas9 on/off-target using DANGER analysis for RNA-seq data from *in vivo* tissue of zebrafish

We performed DANGER analysis using the RNA-seq data from human cells edited by one of the Cas9n-sgRNAs for benchmarking with the previous method (CRISPRroots) and optimized parameters for the DANGER analysis in the previous sections. Next, we investigated whether DANGER analysis could be used to analyze the RNA-seq data from the non-human tissue that had been *in vivo* edited with a single Cas9 nuclease-sgRNA, a more common experimental design for genome editing. Thus, we downloaded and analyzed RNA-seq data from the park7 dataset derived from the zebrafish brain, with and without indels at the park7 locus[20]. Our DANGER analysis successfully built a *de novo* transcriptome assembly with 90.9% complete BUSCO genes and detected on-target sequences in the two transcripts (Figure 7A). Moreover, DANGER analysis revealed a significant downregulation of the transcript with Cas9-sgRNA on-target sequence in the expression quantification (Figure 7B). The original report on the park7 dataset reported downregulated park7 mRNA in the park7 mutant using RNA-seq analysis[20]. This result implied that *de novo* transcriptome assembly and the following expression quantification of our DANGER analysis could generate reliable data consistent with the outcome of standard RNA-seq analysis using a reference genome. Consequently, 19,314 potential off-target sites in all transcripts were detected by optimized dTPM, which were then defined as deleterious off-targets considering the expression profile. There were 4,668 and 70 deleterious off-target sites on all and gene-annotated transcripts, respectively (Figure 7C). The park7 result had no deleterious off-target effects with ≤ 4 mismatches, which meant that frequent off-targeting was not expected in mRNA-transcribed regions. The detected rate of gene-annotated transcripts to all transcripts was more than ten times less than that of optimized dTPM using human GRIN2B (Figure 3D and Figure 7C). The fewer annotations resulted from the poor gene database of zebrafish in comparison with that of humans. Next, our pipeline estimated the D-index per GO term to quantify phenotypic risk. The total D-index was 51 (N=636) (Figure 7D and Supplementary Table S5), which was less than that of human GRIN2B due to fewer annotations and off-target genes. Finally, we validated the D-indices using the permutation test as the same procedure in the last section. Only five significant D-indices were detected (Figure 7E) because the analysis considered only 70 deleterious off-target sites in gene-annotated transcripts. As discussed in

1 the previous section, the park7 analysis has empirically demonstrated that the initially

2 considered number of genes and the total number of D-indices can be small in organisms

3 other than humans. The screening approach of optimized dTPM allows for the acquisition of

4 significant D-indices in poorly annotated data sets.

Comparison of phenotypic risks in the GRIN2B and Park7 datasets

In this study, we evaluated the phenotypic risks associated with off-target transcripts using

the D-index. The number of significant D-indices of the GRIN2B result was approximately 32-

fold larger than that of the park7 result in the optimized dTPM. Furthermore, the DANGER

analysis found off-target genes with four mismatches in the GRIN2B dataset, which is

common in genome-wide off-target studies such as GUIDE-seq and Digenome-seq[24,26] and

numerous deleterious off-target sites on transcript sequences annotated with genes. Thus,

Cas9n-GRIN2B-REV sgRNA may have side effects on the phenotype of differentiated human

iPSC. A follow-up study is required to assess the edited GRIN2B LOF clones using WGS or

alternative genome-wide methods such as GUIDE-seq[24]. Researchers can identify some

clarified points in the future using the result table of the significant D-index (Supplementary

Table S3-4). For example, the GO term "nucleoside monophosphate metabolic process" (GO

ID: GO:0009123) recorded a top significant D-index for the GRIN2B result of optimized dTPM

(Supplementary Table S3). The GO terms of "cell differentiation" (GO ID: GO:0030154), "cell

population proliferation" (GO ID: GO:0008283), and "cell cycle" (GO ID: GO:0007049) were

considered as the phenotype of GRIN2B knock-out in the previous study[23]. However, these

GO terms were listed in the significant D-indices of optimized dDE (Supplementary Table S4,

Table 2). The previous study showed that the gene expression changes of these GO terms

resulted from on-target editing of the GRIN2B locus[23]. However, the results of the DANGER

analysis suggested that off-target editing of additional genes belonging to GO:0030154,

GO:0008283, and GO:0007049 partially contributed to expression changes. Follow-up studies

should include off-target gene analysis of the associated off-targets with the GO terms. On

the other hand, The GO terms of "central nervous system development " (GO ID:

GO:0007417), "brain development" (GO ID: GO:0007420), "cell division" (GO ID: GO:0051301),

and "chromosome segregation" (GO ID: GO:0007059) were not listed in the significant D-

indices of optimized dTPM and optimized dDE, which suggested the GO terms were obvious phenotypes in the GRIN2B knock-out cells. Therefore, DANGER analysis would help reach a reasonable conclusion in genome editing studies.

Limitations

In this section, we discuss the major limitations of the proposed pipeline. First, our method depends on the quality of *de novo* transcriptome assembly using Trinity. Pair-end RNA-seq data with sufficient length and read number must be used to guarantee high-quality assembly (see Material and Methods). If we fail to build an exemplary assembly, producing reliable data for the following analyses, such as on/off-target analysis and expression profiles, becomes difficult. Second, the annotation analysis step in our pipeline may fail to annotate transcripts adequately due to limited information from databases on genes, transcripts, proteins, and gene function. When researchers apply our pipeline to RNA-seq data from organisms with limited genomic knowledge and evidence, we recommend using a database of a model organism with a strong genomic relationship with the organism being analyzed. DANGER analysis can analyze genome-edited samples without a reference genome, but studies of a related model organism with a well-annotated genome are still required. As a third limitation, DANGER analysis cannot strictly distinguish the effects of modifications to on-target genes on other genes from the impacts of off-target gene modifications. Of course, an on-target gene is excluded from the DANGER analysis. Still, it is difficult to distinguish the influence by considering off-target genes whose expression is controlled by the on-target genes. Such an evaluation can only be elucidated through protein interaction and genome analysis conducted using more specialized knowledge by researchers in each field. Therefore, it is appropriate to complete the follow-up studies mentioned in the previous section using a comprehensive analysis. Traditional studies have discussed results based on RNA-seq analysis under the premise that they are solely derived from the effects of on-target gene modifications. However, our DANGER analysis contradicts this assumption, sounding an alarm about the necessity for more specialized investigations and providing the off-target gene information needed for such follow-up analyses.

Possibility of DANGER Analysis as a Simplified Screening Tool, Contributing to More Rigorous Reference-based Phenotypic Risk Assessment

In this study, we developed DANGER analysis as an initial screening tool for maximizing the evaluation of risks to phenotypes. Meanwhile, it is conceivable that we will also need a more rigorous evaluation system for assessing risks to phenotypes. In constructing such an evaluation system, it is believed that a system utilizing information other than RNA-seq data, such as reanalysis of sample genomes by resequencing the genomic DNA rather than *de novo* transcriptome assembly, would be appropriate. Although our DANGER analysis is a tool that only takes RNA-seq data as input to ensure convenience, there is room to apply the partial algorithm (association between off-target genes and GO terms, phenotype risk calculations using the D-index) into such a rigorous reference-based evaluation system for phenotype risks. We believe there is a high possibility that DANGER analysis could become a foundational presence in this new field of phenotype risk assessment.

DANGER analysis Provides a New Perception of the Conventional Genome Editing Process in Medicine, Agriculture, and Biological Research

We demonstrated our DANGER analysis pipeline, as it allows for (i) the detection of potential DNA on/off-target sites in the mRNA-transcribed region on the genome using RNA-seq data, (ii) evaluation of phenotypic effects by deleterious off-target sites based on the evidence provided by gene expression changes, and (iii) quantification of the phenotypic risk at the GO term level, without a reference genome. Thus, DANGER analysis can be performed on various organisms, personal human genomes, and atypical genomes created by diseases and viruses[28]. We believe that the perception resulting from our DANGER analysis has not been observed in the conventional scheme for genome editing. We illustrate a new scheme using DANGER analysis in organisms (I) with and (II) without a reference genome (Figure 8). For example, (I) model organisms such as humans have a reference genome whose information has been generally used for potential on/off-target searches and post-analysis. However, the reference genome is a representative genome and not the personal genome of the patient or cell lines. Reference-based genome editing does not consider unique single nucleotide polymorphisms (SNPs) or spontaneous genomic rearrangements. DANGER analysis can

supply a personal transcriptome-based on/off-target profile to ensure the phenotypic risk of

unexpected off-target mutations. The new workflow of genome editing would be helpful for

ex vivo gene therapy and cancer research because the genome of a cancer cell is generally

characterized by widespread somatic genomic rearrangements([28]). (II) An organism whose

genome has never been comprehensively sequenced and well characterized is not

considered a reasonable subject for genome editing, as site-specific genome editing is

guaranteed without a reference genome. Some groups, however, have used incomplete

genomic information to construct mutants of non-model organisms([51,52]) that have never

been well-characterized in genomics. Such a conventional scheme is hit-or-miss due to the

risk of erroneous knock-out phenotypes in combination with off-targeting of other genes.

DANGER analysis can provide transcriptome phenotype-aware on-/off-target profiles as well

as sequence information of the expressed genes. This information can be used for the safer

design of genome editing, which enables the optimized design of CRISPR editing by

repeatedly looping back through genome-editing experiments without a reference genome.

Furthermore, the DANGER analysis devised in this study employs a simple algorithm based

on mismatch count for identifying off-target candidates. The type of algorithm has also been

utilized in off-target investigations for other programmable nucleases such as CRISPR-

Cas12a, TALEN, and ZFN. DANGER analysis is open-source and freely adjustable. Thus, the

algorithm of this pipeline could be repurposed for the analysis of various genome editing

systems beyond the CRISPR-Cas9 system. We believe that the DANGER analysis pipeline will

expand the scope of genomic studies and industrial applications using genome editing.

DATA AVAILABILITY

The datasets were derived from the following public domain resources:
https://www.ncbi.nlm.nih.gov/geo. The analyses were performed using DANGER analysis
version 1.0. The Script for the DANGER analysis pipeline is available at
https://github.com/KazukiNakamae/DANGER_analysis. In addition, the software provides a
tutorial on reproducing the results presented in this article on the Readme page. The Docker
image of DANGER_analysis is also available at
https://hub.docker.com/repository/docker/kazukinakamae/dangeranalysis/general.

AUTHOR CONTRIBUTIONS

Kazuki Nakamae: Conceptualization, Software, Formal analysis, Methodology, Validation,

Visualization, Writing of the original draft. Hidemasa Bono: Supervision, Writing – Review,

and Editing.

ACKNOWLEDGEMENTS

We would like to thank all the laboratory members at Hiroshima University, PtBio Inc., and

RIKEN for their valuable comments. We deeply appreciate Dr. Ryo Nozu for providing us with

a supplemental script (https://github.com/RyoNozu/Sequence_editor). We would like to

thank Editage (www.editage.com) for English language editing.

FUNDING

This work was supported by the Center of Innovation for Bio-Digital Transformation (BioDX),

the open innovation platform for industry-academia co-creation (COI-NEXT), the Japan

Science and Technology Agency (JST), COI-NEXT [grant number JPMJPF2010 to H.B.], and

JSPS KAKENHI (grant number 21K17855 to K.N.).

CONFLICT OF INTEREST

K.N. was employed by PtBio Inc. H.B. was a consultant for PtBio Inc. H.B. has a financial

interest in PtBio Inc.

REFERENCES

1. Wiedenheft, B., Sternberg, S.H., and Doudna, J.A. (2012). RNA-guided genetic silencing
systems in bacteria and archaea. Nature *482*, 331–338. 10.1038/nature10886.

2. Terns, M.P., and Terns, R.M. (2011). CRISPR-Based Adaptive Immune Systems. Curr
Opin Microbiol *14*, 321–327. 10.1016/j.mib.2011.03.005.

3. Jinek, M., East, A., Cheng, A., Lin, S., Ma, E., and Doudna, J. (2013). RNA-programmed
genome editing in human cells. eLife *2*, e00471. 10.7554/eLife.00471.

4.   Gillmore, J.D., Gane, E., Taubel, J., Kao, J., Fontana, M., Maitland, M.L., Seitzer, J.,
O'Connell, D., Walsh, K.R., Wood, K., et al. (2021). CRISPR-Cas9 In Vivo Gene
Editing for Transthyretin Amyloidosis. New England Journal of Medicine *385*, 493–502.
10.1056/NEJMoa2107454.

5.   Hsu, P.D., Scott, D.A., Weinstein, J.A., Ran, F.A., Konermann, S., Agarwala, V., Li, Y.,
Fine, E.J., Wu, X., Shalem, O., et al. (2013). DNA targeting specificity of RNA-guided
Cas9 nucleases. Nat Biotechnol *31*, 827–832. 10.1038/nbt.2647.

6.   Bassett, A.R., Tibbit, C., Ponting, C.P., and Liu, J.-L. (2013). Highly Efficient Targeted
Mutagenesis of Drosophila with the CRISPR/Cas9 System. Cell Rep *4*, 220–228.
10.1016/j.celrep.2013.06.020.

7.   Shirai, Y., Piulachs, M.-D., Belles, X., and Daimon, T. (2022). DIPA-CRISPR is a simple
and accessible method for insect gene editing. Cell Reports Methods *2*, 100215.
10.1016/j.crmeth.2022.100215.

8.   Nymark, M., Sharma, A.K., Sparstad, T., Bones, A.M., and Winge, P. (2016). A
CRISPR/Cas9 system adapted for gene editing in marine algae. Sci Rep *6*, 24951.
10.1038/srep24951.

9.   Yoshimitsu, Y., Abe, J., and Harayama, S. (2018). Cas9-guide RNA ribonucleoprotein-
induced genome editing in the industrial green alga Coccomyxa sp. strain KJ.
Biotechnology for Biofuels *11*, 326. 10.1186/s13068-018-1327-1.

10. Jiang, W., Bikard, D., Cox, D., Zhang, F., and Marraffini, L.A. (2013). CRISPR-assisted
editing of bacterial genomes. Nat Biotechnol *31*, 233–239. 10.1038/nbt.2508.

11.Jinek, M., Chylinski, K., Fonfara, I., Hauer, M., Doudna, J.A., and Charpentier, E. (2012).
A Programmable Dual-RNA–Guided DNA Endonuclease in Adaptive Bacterial
Immunity. Science *337*, 816–821. 10.1126/science.1225829.

12. Cong, L., Ran, F.A., Cox, D., Lin, S., Barretto, R., Habib, N., Hsu, P.D., Wu, X., Jiang,
W., Marraffini, L.A., et al. (2013). Multiplex Genome Engineering Using CRISPR/Cas
Systems. Science *339*, 819–823. 10.1126/science.1231143.

13. Mojica, F.J.M., Díez-Villaseñor, C., García-Martínez, J., and Almendros, C. (2009).
Short motif sequences determine the targets of the prokaryotic CRISPR defence system.
Microbiology *155*, 733–740. 10.1099/mic.0.023960-0.

14. Fu, Y., Sander, J.D., Reyon, D., Cascio, V.M., and Joung, J.K. (2014). Improving
CRISPR-Cas nuclease specificity using truncated guide RNAs. Nat Biotechnol *32*, 279–
284. 10.1038/nbt.2808.

15. Qi, L.S., Larson, M.H., Gilbert, L.A., Doudna, J.A., Weissman, J.S., Arkin, A.P., and
Lim, W.A. (2013). Repurposing CRISPR as an RNA-Guided Platform for Sequence-
Specific Control of Gene Expression. Cell *152*, 1173–1183. 10.1016/j.cell.2013.02.022.

16. Komor, A.C., Kim, Y.B., Packer, M.S., Zuris, J.A., and Liu, D.R. (2016). Programmable
editing of a target base in genomic DNA without double-stranded DNA cleavage. Nature
533, 420–424. 10.1038/nature17946.

17. Nishida, K., Arazoe, T., Yachie, N., Banno, S., Kakimoto, M., Tabata, M., Mochizuki, M., Miyabe, A., Araki, M., Hara, K.Y., et al. (2016). Targeted nucleotide editing using hybrid prokaryotic and vertebrate adaptive immune systems. Science *353*, aaf8729. 10.1126/science.aaf8729.

18. Anzalone, A.V., Randolph, P.B., Davis, J.R., Sousa, A.A., Koblan, L.W., Levy, J.M., Chen, P.J., Wilson, C., Newby, G.A., Raguram, A., et al. (2019). Search-and-replace genome editing without double-strand breaks or donor DNA. Nature *576*, 149–157. 10.1038/s41586-019-1711-4.

19. Gagnon, J.A., Valen, E., Thyme, S.B., Huang, P., Ahkmetova, L., Pauli, A., Montague, T.G., Zimmerman, S., Richter, C., and Schier, A.F. (2014). Efficient Mutagenesis by Cas9 Protein-Mediated Oligonucleotide Insertion and Large-Scale Assessment of Single-Guide RNAs. PLOS ONE *9*, e98186. 10.1371/journal.pone.0098186.

20. Hughes, G.L., Lones, M.A., Bedder, M., Currie, P.D., Smith, S.L., and Pownall, M.E. (2020). Machine learning discriminates a movement disorder in a zebrafish model of Parkinson's disease. Dis Model Mech *13*, dmm045815. 10.1242/dmm.045815.

21. Liu, D., Awazu, A., Sakuma, T., Yamamoto, T., and Sakamoto, N. (2019). Establishment of knockout adult sea urchins by using a CRISPR-Cas9 system. Development, Growth & Differentiation *61*, 378–388. 10.1111/dgd.12624.

22. Koike-Yusa, H., Li, Y., Tan, E.-P., Velasco-Herrera, M.D.C., and Yusa, K. (2014). Genome-wide recessive genetic screening in mammalian cells with a lentiviral CRISPR-guide RNA library. Nat Biotechnol *32*, 267–273. 10.1038/nbt.2800.

23. Bell, S., Maussion, G., Jefri, M., Peng, H., Theroux, J.-F., Silveira, H., Soubannier, V., Wu, H., Hu, P., Galat, E., et al. (2018). Disruption of GRIN2B Impairs Differentiation in Human Neurons. Stem Cell Reports *11*, 183–196. 10.1016/j.stemcr.2018.05.018.

24. Tsai, S.Q., Zheng, Z., Nguyen, N.T., Liebers, M., Topkar, V.V., Thapar, V., Wyvekens, N., Khayter, C., Iafrate, A.J., Le, L.P., et al. (2015). GUIDE-seq enables genome-wide profiling of off-target cleavage by CRISPR-Cas nucleases. Nat Biotechnol *33*, 187–197. 10.1038/nbt.3117.

25. Luo, X., He, Y., Zhang, C., He, X., Yan, L., Li, M., Hu, T., Hu, Y., Jiang, J., Meng, X., et al. (2019). Trio deep-sequencing does not reveal unexpected off-target and on-target mutations in Cas9-edited rhesus monkeys. Nat Commun *10*, 5525. 10.1038/s41467-019-13481-y.

26. Kim, D., Bae, S., Park, J., Kim, E., Kim, S., Yu, H.R., Hwang, J., Kim, J.-I., and Kim, J.-S. (2015). Digenome-seq: genome-wide profiling of CRISPR-Cas9 off-target effects in human cells. Nat Methods *12*, 237–243. 10.1038/nmeth.3284.

27. Levy, S., Sutton, G., Ng, P.C., Feuk, L., Halpern, A.L., Walenz, B.P., Axelrod, N., Huang, J., Kirkness, E.F., Denisov, G., et al. (2007). The Diploid Genome Sequence of an Individual Human. PLOS Biology *5*, e254. 10.1371/journal.pbio.0050254.

28. Vogelstein, B., Papadopoulos, N., Velculescu, V.E., Zhou, S., Diaz, L.A., and Kinzler, K.W. (2013). Cancer Genome Landscapes. Science *339*, 1546–1558. 10.1126/science.1235122.

29. Ashburner, M., Ball, C.A., Blake, J.A., Botstein, D., Butler, H., Cherry, J.M., Davis, A.P., Dolinski, K., Dwight, S.S., Eppig, J.T., et al. (2000). Gene Ontology: tool for the unification of biology. Nat Genet *25*, 25–29. 10.1038/75556.

30. The Gene Ontology resource: enriching a GOld mine (2020). Nucleic Acids Res *49*, D325–D334. 10.1093/nar/gkaa1113.

31. Tamura, K., and Bono, H. (2022). Meta-Analysis of RNA Sequencing Data of Arabidopsis and Rice under Hypoxia. Life *12*, 1079. 10.3390/life12071079.

32. Toga, K., Yokoi, K., and Bono, H. (2022). Meta-Analysis of Transcriptomes in Insects Showing Density-Dependent Polyphenism. Insects *13*, 864. 10.3390/insects13100864.

33. Bono, H. (2021). Meta-Analysis of Oxidative Transcriptomes in Insects. Antioxidants *10*, 345. 10.3390/antiox10030345.

34. Suzuki, T., Ono, Y., and Bono, H. (2021). Comparison of Oxidative and Hypoxic Stress Responsive Genes from Meta-Analysis of Public Transcriptomes. Biomedicines *9*, 1830. 10.3390/biomedicines9121830.

35. Subramanian, A., Tamayo, P., Mootha, V.K., Mukherjee, S., Ebert, B.L., Gillette, M.A., Paulovich, A., Pomeroy, S.L., Golub, T.R., Lander, E.S., et al. (2005). Gene set enrichment analysis: A knowledge-based approach for interpreting genome-wide expression profiles. Proceedings of the National Academy of Sciences *102*, 15545–15550. 10.1073/pnas.0506580102.

36. Hölzer, M., and Marz, M. (2019). De novo transcriptome assembly: A comprehensive cross-species comparison of short-read RNA-Seq assemblers. Gigascience *8*, giz039. 10.1093/gigascience/giz039.

37. Lipka, A., Paukszto, L., Majewska, M., Jastrzebski, J.P., Panasiewicz, G., and Szafranska, B. (2019). De novo characterization of placental transcriptome in the Eurasian beaver (Castor fiber L.). Funct Integr Genomics *19*, 421–435. 10.1007/s10142-019-00663-6.

38. Grabherr, M.G., Haas, B.J., Yassour, M., Levin, J.Z., Thompson, D.A., Amit, I., Adiconis, X., Fan, L., Raychowdhury, R., Zeng, Q., et al. (2011). Full-length transcriptome assembly from RNA-Seq data without a reference genome. Nat Biotechnol *29*, 644–652. 10.1038/nbt.1883.

39. Khudyakov, J.I., Champagne, C.D., Meneghetti, L.M., and Crocker, D.E. (2017). Blubber transcriptome response to acute stress axis activation involves transient changes in adipogenesis and lipolysis in a fasting-adapted marine mammal. Sci Rep *7*, 42110. 10.1038/srep42110.

40. Nespolo, R.F., Gaitan-Espitia, J.D., Quintero-Galvis, J.F., Fernandez, F.V., Silva, A.X., Molina, C., Storey, K.B., and Bozinovic, F. (2018). A functional transcriptomic analysis in the relict marsupial Dromiciops gliroides reveals adaptive regulation of protective functions during hibernation. Molecular Ecology *27*, 4489–4500. 10.1111/mec.14876.

41. Martin, M. (2011). Cutadapt removes adapter sequences from high-throughput sequencing reads. EMBnet.journal *17*, 10–12. 10.14806/ej.17.1.200.

42. Manni, M., Berkeley, M.R., Seppey, M., Simão, F.A., and Zdobnov, E.M. (2021).
BUSCO Update: Novel and Streamlined Workflows along with Broader and Deeper
Phylogenetic Coverage for Scoring of Eukaryotic, Prokaryotic, and Viral Genomes. Mol
Biol Evol *38*, 4647–4654. 10.1093/molbev/msab199.

43. Jacquin, A.L.S., Odom, D.T., and Lukk, M. (2019). Crisflash: open-source software to
generate CRISPR guide RNAs against genomes annotated with individual variation.
Bioinformatics *35*, 3146–3147. 10.1093/bioinformatics/btz019.

44. Corsi, G.I., Gadekar, V.P., Gorodkin, J., and Seemann, S.E. (2021). CRISPRroots: on-
and off-target assessment of RNA-seq data in CRISPR–Cas9 edited cells. Nucleic Acids
Res *50*, e20. 10.1093/nar/gkab1131.

45. Sun, J., Nishiyama, T., Shimizu, K., and Kadota, K. (2013). TCC: an R package for
comparing tag count data with robust normalization strategies. BMC Bioinformatics *14*,
219. 10.1186/1471-2105-14-219.

46. Fu, R., He, W., Dou, J., Villarreal, O.D., Bedford, E., Wang, H., Hou, C., Zhang, L.,
Wang, Y., Ma, D., et al. (2022). Systematic decomposition of sequence determinants
governing CRISPR/Cas9 specificity. Nat Commun *13*, 474. 10.1038/s41467-022-28028-
x.

47. Kurosaki, T., Popp, M.W., and Maquat, L.E. (2019). Quality and quantity control of gene
expression by nonsense-mediated mRNA decay. Nat Rev Mol Cell Biol *20*, 406–420.
10.1038/s41580-019-0126-2.

48. Srivastava, A., Malik, L., Sarkar, H., Zakeri, M., Almodaresi, F., Soneson, C., Love, M.I.,
Kingsford, C., and Patro, R. (2020). Alignment and mapping methodology influence
transcript abundance estimation. Genome Biol *21*, 239. 10.1186/s13059-020-02151-8.

49. Wagner, G.P., Kin, K., and Lynch, V.J. (2012). Measurement of mRNA abundance using
RNA-seq data: RPKM measure is inconsistent among samples. Theory Biosci. *131*, 281–
285. 10.1007/s12064-012-0162-3.

50. Dobin, A., Davis, C.A., Schlesinger, F., Drenkow, J., Zaleski, C., Jha, S., Batut, P.,
Chaisson, M., and Gingeras, T.R. (2013). STAR: ultrafast universal RNA-seq aligner.
Bioinformatics *29*, 15–21. 10.1093/bioinformatics/bts635.

51. Chang, K.S., Kim, J., Park, H., Hong, S.-J., Lee, C.-G., and Jin, E. (2020). Enhanced lipid
productivity in AGP knockout marine microalga Tetraselmis sp. using a DNA-free
CRISPR-Cas9 RNP method. Bioresource Technology *303*, 122932.
10.1016/j.biortech.2020.122932.

52. Sun, Q., Lin, L., Liu, D., Wu, D., Fang, Y., Wu, J., and Wang, Y. (2018). CRISPR/Cas9-
Mediated Multiplex Genome Editing of the BnWRKY11 and BnWRKY70 Genes in
Brassica napus L. International Journal of Molecular Sciences *19*, 2716.
10.3390/ijms19092716.

TABLE AND FIGURES LEGENDS

Tables

Table 1. RNA-seq datasets evaluated for testing the DANGER analysis pipeline.

Table 2. D-index summary of GO terms related to the enrichment analysis of previous

GRIN2B research.

Figures

Figure 1. Scheme of CRISPR-Cas9 targeting, deleterious off-target editing, and DANGER

analysis.

Figure 2. Overview of DANGER analysis and on-target region constructed by *de novo*

transcriptome assembly. A. Bioinformatic workflow of DANGER analysis. Our analysis requires

RNA-seq data derived from WT and Edited (each n≥3). DANGER analysis has two steps in the

workflow: (1) *de novo* transcriptome assembly (light green background color) and (2)

annotation analysis (light yellow background color). The *de novo* transcriptome assembly

step is processed with Trinity and preprocessing tools such as cutadapt and bbduk.sh.

Crisflash performs the search of on/off-target sequences. The RSEM quantifies gene

expression in Edited RNA-seq samples in comparison to the WT *de novo* transcriptome (dot

allow). The step of annotation analysis was involved processing with TransDedoder, ggsearch,

org.XX.eg.db (e.g., org.Hs.eg.db in the transcriptome related to humans), and topGO. We

implemented specific modules, colored in pink, for considering the phenotypic effect of

deleterious off-targets. B. Comparison between the hg38 reference genome and transcript

sequence constructed by *de novo* assembly of RNA-seq samples derived from WT iPSC-

derived cortical neurons on the GRIN2B on-target region. The on-target region of the hg38

reference genome is illustrated with annotations of the *GRIN2B* CDS, the protospacer, and

the NGG PAM sequence of SpCas9. The detected GRIN2B isoforms (1–5) are lined up in

green in the black box. The Cas9-sgRNA binding sites are highlighted in blue. C. Genome

completeness of *de novo* transcriptome assembly RNA-seq data derived from WT iPSC-

derived cortical neurons was assessed using conserved mammal BUSCO genes

(mammalia_odb10). The result was shortly 79.1% of "complete," 20.7% of "single-copy," 58.4%

of "duplicated," 3.2% of "fragmented," and 17.7% of "missing" (n = 9226).

Figure 3. The benchmark for expression analysis methods compared with reference-based

RNA-seq analysis using RNA-seq data derived from WT and GRIN2B edited iPSC-derived

cortical neurons. A. Comparison of different expression analyses. A Venn diagram comparing

the *de novo* transcripts (duplicate counts on a predicted ORF basis), which had potential off-

target sites with up to 8 nt mismatches, was detected by the dTPM and dDE approaches.

dTPM indicates that the expression is decreased based on the ratio of TPM counts between

WT and Edited samples (left callout). dDE means the expression is reduced based on DEG

analysis between WT and Edited samples (right callout). B. Comparison of *de novo*

transcriptome assembly-based and reference-based analysis on the deleterious off-target

detection. A Venn diagram comparing the off-target genes identified from *de novo*

transcriptome analysis (dTPM (t = 0.4) and dDE (α = 0.001) approaches) and reference-based

RNA-seq analysis (CRISPRroots, "RISK: CRITICAL"). C. Genomic sequence map of off-target

located outside of GALR2 mRNA. The sequence is a part of the hg38 reference genome with

annotations of GALR2 mRNA (XM_047436984.1) and the *de novo* transcript

(TRINITY_DN86617_c0_g1_i1) and an off-target site with three mismatches compared to the

on-target sequence. D. Summary of deleterious off-target sites detected by *de novo*

transcriptome analysis (dTPM) and reference-based RNA-seq analysis (CRISPRroots, "RISK:

CRITICAL"). D. The counts of off-target sites are annotated with genes and classified by

mismatch number related to the on-target sequence. The brackets indicate the number of

transcripts, including those with and without identified gene annotations. The number of

genes and transcripts with ≤4 nt mismatches is colored red.

Figure 4. The result of risk assessment in DANGER analysis using RNA-seq data derived from

WT and GRIN2B edited iPSC-derived cortical neurons. A. An example of the annotation table

for DANGER analysis. The table includes GO ID, GO term, number of mismatches (n), and the

counts of n-MM off-target genes belonging to a specific GO term. B. The formula for

phenotypic off-target risk (D-index). An example of the calculation is shown on a yellow

background. C. Distribution of the D-index of each GO term (orange). The sum of all D-

indexes and the number of D-indices (N) were labeled on the top right.

Figure 5. A scheme for permutation testing to evaluate the validity of the D-index. The thin

black arrow indicates the manipulation of rearranging values from the original expression

and off-target profile to the permutation data. The black cross represents the computation

for applying the D-index formula to the above expression profile and the below off-target

profile data. The workflow is shown as the orange allows.

Figure 6. Evaluation of permutation test for DANGER analysis and comparison between

dTPM and dDE. A. Comparison of false detection rates among approximate dTPM (up to 11-

MM NRR PAM, t=0.4, L=5E-1), optimized dTPM (up to 8-MM NGG PAM, t=0.4, L=1E-15),

and optimized dDE (up to 8-MM NGG PAM, α = 0.001, L=1E-15) in GO categories. BP, CC,

and MF indicate GO categories of Biological Process, Cellular Component, and Molecular

Function, respectively. Error bars represent SEM; asterisk indicates the statistical significance

of two-sided Welch's t-test; cross indicates statistical power (1-ß) >0.8. Mean ± s.d. of n = 10

permutation data set. B. Comparison of amount of GO terms of all D-index, significant D-

index, and expected true D-index among approximate dTPM (up to 11-MM NRR PAM, t=0.4,

L=5E-1), optimized dTPM (up to 8-MM NGG PAM, t=0.4, L=1E-15), and optimized dDE (up

to 8-MM NGG PAM, α = 0.001, L=1E-15), respectively. C. Comparison of D-index and

significant D-index between optimized dTPM and optimized dDE. A Venn diagram

comparing the counts of D-index and significant D-index between optimized dTPM and

optimized dDE. D. The list of the top 16 significant D-indices in the optimized dTPM. The D-

index values are indicated by bar graphs adjacent to the GO terms.

Figure 7. DANGER analysis result using RNA-seq data derived from WT and park7 (dj1)

Edited brains of *Danio rerio*. A. Comparison between the GRCz11 reference genome and

transcript sequence constructed by *de novo* assembly of RNA-seq samples derived from WT

brain on park7 on-target region. The on-target region of the GRCz11 reference genome is

illustrated with annotations of the park7 CDS, the protospacer, and the NGG PAM sequence

of SpCas9. The detected park7 isoforms (1-2) are lined up in green in the black box. The

Cas9-sgRNA binding sites are highlighted in blue. B. Comparison of TPM values of park7. The

TPM was measured from WT and Edited RNA-seq samples (Each n=3); data were expressed

as the means±SEM. *** p-value < 0.001 of two-sided Welch's t-test. C. The gene counts are

classified by mismatch number related to the on-target sequence. The brackets indicate the

number of transcripts, including those with and without identified gene annotations. The

number of genes and transcripts with ≤4 nt mismatches is colored red. D. Distribution of the

D-index of each GO term associated with Biological Process (orange). The sum of all D-

indices and the number of D-indices (N) is labeled on the top right. E. The list of all

significant D-indices in the optimized dTPM. The D-index values are indicated by bar graphs

adjacent to the GO terms. The colors of the bar indicate GO categories belonging to the GO

terms.

Figure 8. Our proposal for the usage of DANGER-analysis in organisms with and without a

reference genome. The workflow is shown as black arrows. The dotted black arrows indicate

the front of the arrow and refer to the arrow base information. The image of the book is

from TogoTV (© 2016 DBCLS TogoTV, CC-BY-4.0,

https://creativecommons.org/licenses/by/4.0/).

1   **Table1.**

Dataset	Layout	Length of raw reads [nt]	Total raw reads (Spots)	The average length of raw reads [nt]	Run	WT/Edited	Ref
GRIN2B	paired-end	125	44,621,372	125	SRR7187518	WT	[23]
		125	39,290,710		SRR7187519	WT	
		125	42,356,322		SRR7187520	WT	
		125	37,725,951		SRR7187521	WT	
		125	40,423,942	125	SRR7187524	Edited	
		125	43,003,607		SRR7187525	Edited	
		125	35,134,095		SRR7187526	Edited	
		125	39,725,694		SRR7187527	Edited	
park7 (dj1)	paired-end	151	55,801,577	151	SRR9886606	WT	[20]
		151	58,680,708		SRR9886607	WT	
		151	52,584,965		SRR9886608	WT	
		151	57,060,988	151	SRR9886609	Edited	
		151	48,882,852		SRR9886610	Edited	
		151	53,633,192		SRR9886611	Edited	

2
3

1 **Table2.**

GO term	Optimized dTPM in GRIN2B			Optimized dDE in GRIN2B		
	Significant D-index	Rank	%Rank	Significant D-index	Rank	%Rank
cell differentiation	-	-	-	4.507514299	54	4.9046
cell population proliferation (cell proliferation)	-	-	-	1.204410358	245	22.2525
cell cycle	-	-	-	1.140013876	264	23.9782
central nervous system development	-	-	-	-	-	-
brain development	-	-	-	-	-	-
cell division	-	-	-	-	-	-
chromosome segregation	-	-	-	-	-	-

1 **Figure 1.**

Figure 2.

Figure 3.

(): number of transcripts *: multiple gene annotations from one transcript

Mismatch number	0	1	2	3	4	5	6	7	8
DANGER analysis (dTPM t = 0.4, up tp 8-MM, NGG PAM)	0	0	0	0	2 (6)	16 (67)	99 (418)	431 (2,310)	1,688 (10,436)
DANGER analysis (dDEG α = 0.001, up to 8-MM, NGG PAM)	0	0	0	0	2* (1)	1 (5)	25 (35)	79 (152)	300 (620)
CRISPRroots (RISK: CRITICAL)	0	0	0	0	0	0	2	0	0

1 **Figure 4.**

1 **Figure 5.**

1   **Figure 6.**

1 **Figure 7.**

1   **Figure 8.**