Synaptic deregulation of cholinergic projection neurons causes olfactory dysfunction across 5 fly Parkinsonism models

Ulrike Pech[1,2,†], Jasper Janssens[1,3,4,†], Nils Schoovaerts[1,2], Sabine Kuenen[1,2], Samira Makhzami[1,3,4], Gert Hulselmans[1,3,4], Suresh Poovathingal[1,3,4], Adekunle T. Bademosi[1,2], Jef Swerts[1,2], Sven Vilain[1,2], Stein Aerts[1,3,4] and Patrik Verstreken[1,2,*]

[1] VIB-KU Leuven Center for Brain & Disease Research

[2] KU Leuven, Department of Neurosciences, Leuven Brain Institute

[3] KU Leuven, Department of Human Genetics, Leuven Brain Institute

[4] VIB Center for AI & Computational Biology (VIB.AI)

3000 Leuven, Belgium

[*] corresponding author: patrik.verstreken@kuleuven.be

[†] These authors contributed equally to this work

Abstract

The classical diagnosis of Parkinsonism is based on motor symptoms that are the consequence of nigrostriatal pathway dysfunction and reduced dopaminergic output. However, a decade prior to the emergence of motor issues, patients frequently experience non-motor symptoms, such as a reduced sense of smell (hyposmia). The cellular and molecular bases for these early defects remain enigmatic. To explore this, we developed a new collection of five fruit fly models representing various forms of familial Parkinsonism and conducted single-cell RNA sequencing on young entire brains of these models. Interestingly, cholinergic projection neurons are the most vulnerable cells and genes associated with presynaptic function are the most deregulated. Additional single nucleus sequencing of three specific brain regions of Parkinson's disease patients recapitulates these finding. Indeed, these disturbances lead to early synaptic dysfunction, notably affecting cholinergic olfactory projection neurons crucial for olfactory function in flies. Our research uncovers that one of the earliest indicators of disease in 5 different models of familial Parkinsonism is synaptic dysfunction in higher-order cholinergic projection neurons, and it suggests common synaptic molecular pathways contribute to the development of hyposmia.

Keywords: Parkinson's disease (PD), Single-cell RNA sequencing, Cholinergic neurons, Olfaction

Introduction

Parkinson's disease (PD) is a complex multifactorial neurodegenerative disease that affects millions. Dopaminergic neuron (DAN) death in the *Substantia nigra* causes motor defects in patients. However, 97 % of patients suffer from non-motoric dysfunction decades earlier, including constipation, REM sleep disorder and hyposmia (Ansari & Johnson, 1975; Chase & Markopoulou, 2020; Doty et al., 1988; Parkinson, 2002; Schapira et al., 2017). These non-motor defects in PD are often overlooked because they occur before disease diagnosis and thus, they do not belong to the normal clinical work-up. Furthermore, non-motoric defects are largely non-responsive to dopamine replacement therapy suggesting other cell types are involved (Kalia & Lang, 2015).

Hyposmia, the reduced ability to discern odors, is one of the earliest defects associated with familial and sporadic Parkinsonism (Chase & Markopoulou, 2020; Doty, 2012; Doty et al., 1988; Haehner et al., 2009). These patients show difficulties to classify odors: they often perceive banana odor as motor oil or beer odor as fruit in standardized odor identification tests (UPSIT) (Double et al., 2003). These observations suggest that hyposmia in Parkinsonism is caused by a defect in higher-order perception, consistent with the observations that first-order olfactory receptor cells of the nasal epithelium are not majorly affected in patients (Mueller et al., 2005; Witt et al., 2009). The synaptic site of second-order olfactory neurons, the olfactory bulb, is modulated by basal forebrain cholinergic neurons and local GABAergic and dopaminergic interneurons (Harvey & Heinbockel, 2018). While these DAN do not degenerate in Parkinsonism, it has been suggested there is local impairment of cholinergic transmission in the olfactory bulb of Parkinsonism patients and vertebrate models (Huisman et al., 2004; Mundiñano et al., 2013; S. Zhang et al., 2015), but the molecular defects are elusive.

There are only few models to study hyposmia in Parkinsonism and most are based on the use of toxins or expression of mutant alpha-Synuclein (Taguchi et al., 2020). While this may recapitulate aspects of the disease, they bias the disease mechanisms to known phenomena associated with the specific model. Modelling approaches to Parkinsonism are further hindered due to the large number of mutated genes. These mutations affect various biological functions and therefore it is currently hard to unveil the specific and possibly common pathways and cell types that are at the basis of the early defects in the disease. So far, mutations in more than 25 genes have been identified that cause familial forms of Parkinsonism (Brooker et al., 2024). These genes cause defects in a diverse range of processes including mitochondrial function,

endocytosis, lysosomal function, autophagy and protein translation (Kalia & Lang, 2015). Nonetheless, these mutations do result in overlapping motor and non-motor defects, including hyposmia. This raises the question of whether defective mechanisms across the genetic space of Parkinsonism converge on common biological processes and affect similar cell types beyond DAN. In this study, we generated and utilized five well-controlled *Drosophila* Parkinsonism models to investigate cellular and molecular dysfunction preceding motor defects that occur later in the life of these animals.

Results

A new collection of five familial PD knock-in models.

The characterization of cells and pathways affected in PD is hindered because patient populations are heterogeneous and because mutations in numerous genes have been linked to familial forms of the disease (Brooker et al., 2024). To overcome these problems, we created a new collection of *Drosophila* PD models that was extensively backcrossed into one isogenic background (>10 generations). We selected homologues of 5 PD genes that represent different affected pathways and clinical manifestations in this disease (Kalia & Lang, 2015) and replaced the *Drosophila lrrk, rab39, auxilin (aux), synaptojanin (synj)* and *Pink1* genes by wild-type and pathogenic mutant human or *Drosophila* cDNA sequences (CDS). We confirm that the wild-type and pathogenic mutant human CDS (*LRRK2, RAB39B* and *PINK1*) or *Drosophila* CDS (*DNAJC6 (Aux)* and *SYNJ1 (Synj)*) are expressed at endogenous levels (Fig. 1a, Supplemental Fig. 1a, Supplemental File 1). To determine if our new models recapitulate known PD-relevant phenotypes we tested them in an array of assays that were previously used to analyze other fly PD models. Wild-type PD-gene knock-ins are similar to controls, indicating the cDNA knock-ins (human or fly) recapitulate normal gene function. Conversely, the PD knock-in models show defects akin to those seen in previously described mutants of these genes. For example, we find reduced lifespan under starvation, disturbances in mitochondrial membrane potential and electrophysiological defects in electroretinogram (ERG) recordings (Hindle et al., 2013; Hoffmann & Harshman, 1999; Morais et al., 2009; Mortiboys et al., 2010; Ng et al., 2012; Vanhauwaert et al., 2017) (Fig. 1b, Supplemental Fig. 1b-d). Importantly, we also tested the flies for their performance in a startle-induced negative geotaxis assay (SING). This is a motor behavior that critically depends on specific central brain dopaminergic neurons (**p**osterior-**a**nterior-**m**edial dopaminergic neurons (DAN): "PAM"). Previous work showed that PAM-

dependent SING was unaffected in young flies that express alpha-Synuclein, but that SING declines as these flies age (Riemensperger et al., 2013). Similarly, none of our models shows a SING defect at a young age (5 d, young) (Fig.1b). However, as flies age (25 d, old), the mutant knock-ins display impaired performance in this DAN-dependent locomotion assay, while the performance of controls is not affected (Fig. 1b). This suggests that the mutants we used in this study suffer from a progressive locomotion defect that has been linked to DAN dysfunction.

Cholinergic olfactory projection neurons are impaired across 5 PD fly models.

To define cell types affected by pathogenic PD mutations early in life, we used an unbiased approach and conducted single cell RNA sequencing of entire brains of young (5 d) animals (Supplemental File 2). We pooled 108k newly sequenced cells from these 5 mutants and controls with the 118k wild-type cells from our original atlas leading to 186 cell clusters, of which 81 were annotated to known cell types (Davie et al., 2018; Janssens et al., 2022) (Fig. 2a). PD mutant cells equally mix with the cells of the original fly brain cell atlas and the frequency of recovered cell types is similar between control and mutant brains (including DAN; Fig. 2b-f, Supplemental Fig. 2a, b). We further do not find any cell type that shows higher or lower expression of the knocked-in CDS (Supplemental Fig. 2c). Thus, cellular identity and cellular composition are preserved in young PD fly models.

For every cluster, we then compared gene expression between cells from mutants and controls using two independent algorithms (Wilcoxon and DESEQ2, (Wang et al., 2019)), and find both methods correlate well (average Spearman correlation of signed p-value > 0.8 to > 0.95; Supplemental Fig. 2d). However, the number of differentially expressed genes (DEG) is largely dependent on the number of cells present in a cluster. To correct for this bias, we fitted a negative binomial model that correlated the number of DEG and cell cluster size in each mutant (Supplemental Fig. 2e, Supplemental File 3). Using the residuals of this model we find eight common predominantly affected cell types across the 5 mutants (Supplemental File 4). We also find cell types uniquely affected in some mutants, e.g., glial subtypes that were affected in $PINK1^{P399L}$. The commonly affected cell types include cholinergic olfactory projection neurons (OPN), Kenyon cells, cholinergic neurons of the visual system (T1 neurons), and other yet-to-be-identified cell types (Fig. 2b-f, black cells). Hence, while our mutants are modelling different genetic origins of disease, we find some cell types are commonly affected. OPN are here of particular interest because they are higher-order projection neurons that control innate olfactory

processing and odor classification (Masse et al., 2009). Furthermore, hyposmia is one of the earliest defects associated with familial and sporadic Parkinsonism (Chase & Markopoulou, 2020; Doty, 2012) with patients having difficulties to classify odors in standardized UPSIT tests (Double et al., 2003; Mueller et al., 2005; Witt et al., 2009).

Synaptic genes and pathways are deregulated in cholinergic brain regions of fly PD models and human PD patients.

We next compared the identity of the DEG found in PD fly model cholinergic projection neurons to the identity of the (homologous) genes differentially expressed in patient brain samples rich in cholinergic neurons. We analyzed *nucleus basalis of Meynert* (NBM)*, nucleus accumbens* (Nac) and *putamen* (Put) from five brains of PD patients (all have idiopathic *LRRK2* risk mutations) and five brains from unaffected age-matched individuals (without PD-relevant genetic variants). Our analysis identified 37 distinct cell types in these brain regions (Supplemental Fig. 3a), and most of the cell types have a similar frequency when comparing PD vs non-PD condition. An exception is a microglia subtype (MG #20, Supplemental Fig. 3b), which is 900% increased in frequency in PD patient brains compared to non-PD samples. We next analyzed neuronal subtypes, including cholinergic neurons (Supplemental Fig. 3c, d) and identified the DEG in these cells. Many of the DEG are genes encoding synaptic proteins that are also listed in the synapse-specific portal SynGo (Koopmans et al., 2019) (Fig. 3a, a'). We then asked how they compare to the DEG found in affected cholinergic projection neurons of our five fly models (Fig. 3a"). We therefore first listed the deregulated genes in affected cell types for each fly model. There are many DEG unique to each PD mutant, but there is also overlap in DEG between the models. To define this overlap across fly PD mutants, we ranked genes according to their signed differential expression for each of the 5 PD models (up or downregulated) and then summed their rank across the mutants retaining the genes in the top 5% and the bottom 5% as the common highly up- or downregulated genes (Supplemental File 5). Additionally, when we compare this list of deregulated fly genes to the DEG-orthologues in PD patients, there is remarkable overlap. For example, in the NBM*,* an area associated with PD (Arendt et al., 1983), 20% of the DEG that have an orthologous gene in the fly are also found among the most deregulated genes across PD fly models.

Next, we analyzed the list of human-fly overlapping DEG using Gorilla gene ontology enrichment (Eden et al., 2009). Markedly, this shows that the processes most strikingly enriched

in both fly models and human NBM are those related to synaptic function (Fig. 3b). Indeed, more than half of the DEG found in human and flies are present in the synapse-specific portal SynGo (Koopmans et al., 2019) (Fig. 3a"). We found enrichment for presynaptic function, including synaptic exocytosis, axon guidance, synapse organization, ion transport and protein homeostasis (Fig. 3b, c). Additionally, we also found enrichment of genes involved in RNA regulation and mitochondrial function that are also important for the functioning of synaptic terminals (Gorenberg & Chandra, 2017; Morais et al., 2009; Snead & Eliezer, 2019; Uytterhoeven et al., 2011). In summary, cholinergic neurons of young familial fly PD models and human PD patients both show preferential transcriptional deregulation of genes that regulate synaptic function (Fig. 3c).

Young PD models suffer from synaptic impairment of cholinergic OPN.

The transcriptional deregulation of genes encoding synaptic proteins predicts synaptic alterations in the affected cell types. We therefore expressed the Ca^{2+}-sensor GCaMP3 in the OPN, one of the majorly affected cell types in our PD flies, and monitored synaptic terminal function through a small window in the head capsule of living flies upon stimulation (Fig 4a, b). Neuronal stimulation (10 mM Nicotine) results in a robust synaptic Ca^{2+} signal at OPN synapses in controls (Fig. 4c, d (black points), quantified Fig. 4g (gray bar)) and in animals with the wild-type PD gene knocked-in (e.g. hLRRK2 in Fig. 4c, d (dark red points), quantified in Fig. 4g (left bar)). In contrast, the five PD-mutant knock-ins show significantly weaker Ca^{2+}-responses (e.g. hLRRK2^{G2019S} in Fig. 4c, d (pink points), and all 5 mutants quantified in Fig. 4g ("OPN>wt CDS -")). Next, we tested if this defect is cell-autonomous and created PD knock-in mutants expressing wild-type CDS *selectively* in their OPN (Fig. 4g ("OPN>wt CDS +")). These animals show synaptic Ca^{2+} signals similar to controls, indicating the PD genes are required in OPN of these young animals to maintain robust synaptic function. Note that while *hLRRK2* knock-in flies are very similar to wild-type controls (Fig. 4g (left bar)), the *hLRRK2^{G2029S}* knock-in flies are not rescued by OPN-specific expression of *hLRRK2* (Fig. 4g (right bar)). This is in agreement with the *G2019S* mutation being dominant (West et al., 2005; Zimprich et al., 2004).

Previous work indicated that functional defects of OPN often cause morphological changes at the level of the synaptic area in the calyx (Kremer et al., 2010). We thus also assessed the synaptic area of OPN in the calyx (based on the GFP fluorescence area of the GCaMP3 sensor). Interestingly, the PD mutants have a significantly larger area compared to controls, despite very

similar GFP expression levels (Fig. 4e, f, h). Again, this defect is cell-autonomous, because it is rescued by OPN-selective expression of the wild-type CDS, except -again- for $LRRK2^{G2019S}$ agreeing with it being a dominant mutant (Fig. 4h). Thus, OPN of young PD mutants display cell-autonomous presynaptic defects: even though the synaptic area is increased, synaptic Ca^{2+} signals are diminished.

Hyposmia is prevalent in young PD models.

While old (25 d) PD mutants show PAM-dependent locomotion dysfunction in the SING assay (Fig. 1b (SING old)), young (5 d) PD mutants do not show SING defects (Fig. 1b (SING young)). These data indicate PAM neurons are functional in young PD mutants. Yet, these young mutants suffer from OPN-synaptic dysfunction (Fig. 4). We thus wondered if our young PD models have olfactory performance defects. We subjected flies to a choice between two odors (test 1: motor oil and banana *or* test 2: beer and wine). Control and wild-type knock-in flies show a robust odor preference in these two odor tests (Fig. 5a, b). Conversely, the pathogenic mutant CDS knock-in flies behave indecisive (Fig. 5a, b). Hence, young PD mutants show olfactory behavior defects.

We then asked if this defect is cell-autonomous and expressed the relevant wild-type CDS *selectively* in the OPN of the pathogenic knock-in mutant flies. This manipulation restores olfactory performance back to control levels, except in $LRRK2^{G2019S}$ (Fig. 5c, d), again agreeing with it being a dominant mutant. Note that $lrrk^{KO}$ flies also show an olfactory performance defect and that expressing $hLRRK2$ cDNA specifically in the OPN of $lrrk^{KO}$ rescues this defect (Fig. 5c, d). This indicates hLRRK2 is a functional protein that, in this context, can compensate for the function of the fly orthologue. Finally, we demonstrate that the rescue of the olfactory performance in PD mutants is specific for OPN, because expressing wild-type CDS in another affected cell type, "T1-interneurons of the visual system", of knock-in PD animals (we chose to test this in $Rab39B^{G193R}$ mutants), does not improve olfactory performance (Fig. 5c, d (5th row)). Hence, a cell autonomous defect in cholinergic OPN causes olfactory defects across PD mutants.

Discussion

Parkinsonism is currently diagnosed based on motor defects that only occur following significant DAN degeneration. However, patients suffer from non-motoric problems decades earlier, including sleep problems and hyposmia (the inability to properly discern odors) (Yamakado &

Takahashi, 2024). Most of these non-motoric problems are not responsive to dopamine replacement therapy, suggesting other transmitter systems are involved. Using unbiased methodology, we find cholinergic system failure as one of the earliest defects across 5 different Parkinsonism fly models and we show that synaptic failure of specific cholinergic projection neurons (OPN) causes non-motor problems, including hyposmia. Also in patients, cholinergic defects in several brain regions have been observed, including the olfactory bulb and the occipital cortex (Kuhl et al., 1996; Shimada et al., 2009). These defects correlate with several non-motor symptoms, including impaired olfactory and visual processing (Bohnen et al., 2010). We now place cholinergic failure firmly ahead of dopaminergic system failure in flies, and we suggest that dysfunction of specific cholinergic projection neurons may be a prodromal signal of disease. Our work opens the possibility to explore the early detection of specific cholinergic interneuron dysfunction, e.g. using PET tracing, as an entry point to move diagnosis and intervention ahead of dopaminergic degeneration.

The connection between early cholinergic neuron dysfunction and subsequent failure of the DAN system in disease progression remains uncertain. However, dopaminergic neurons have a high density of nicotinic acetylcholine receptors, and evidence suggests that stimulating these receptors enhances DAN survival (Brimblecombe et al., 2018). This is supported by epidemiological data indicating that nicotine exposure from chronic smoking protects against PD (Janssens et al., 2022; Wang et al., 2019), and various studies, including our own unpublished data, showing that direct nicotine supplementation mitigates DAN dysfunction in mouse and fly PD models (Chambers et al., 2013; Fares et al., 2023; Olsen et al., 2023). These findings imply that factors affecting the function of cholinergic neurons might, by absence of insufficient innervation, lead to DAN problems and degeneration, warranting further exploration of the underlying molecular mechanisms.

Our findings reveal a common trigger relevant for both familial and idiopathic cases. Observations of Lewy body pathology in postmortem samples suggest the disease may nucleate at specific sites and then propagate. This includes spreading of the disease between the gut and the brain as well as early aggregation of alpha-Synuclein, a synaptic protein, in the olfactory bulb (Braak, Tredici, et al., 2003). The presence of protein aggregates in these areas is consistent with protein homeostasis and synaptic defects that ultimately contribute to the aggregation of alpha-Synuclein (Chu et al., 2009; Friedman et al., 2012; Sarkar et al., 2007; Webb et al., 2003). In idiopathic disease, there is some evidence this is because the gut and olfactory bulb synapses are

more easily exposed to environmental toxins that cause protein misfolding and aggregation (Braak, Rüb, et al., 2003). Fly olfactory projection neurons, like human basal forebrain cholinergic neurons and striatal cholinergic interneurons, are also characterized by large neuronal arbors, numerous synaptic contacts and high activity (C. J. Wilson et al., 1990; R. I. Wilson, 2013; Wu et al., 2014). Such neurons are thought to be particularly vulnerable to synaptic problems (Ashrafi et al., 2014; Vijayan & Verstreken, 2017) and our work now indicates that the defects induced by different familial PD mutations converge onto synapses. This can be through different upstream mechanisms (i.e. the causal genes affect different cellular processes), but what they have in common is that this ultimately induces synaptic failure. This is particularly relevant in such elaborate cholinergic projection neurons where synapses are often at a long distance from their cell body and synapses need to operate relatively independently. As such there can be different mechanisms at play and our single-cell sequencing shows the deregulation of several synaptic pathways in the PD knock-in mutants. While several genes mutated in familial forms of PD have proposed synaptic functions (endocytosis, synaptic autophagy, etc.), others may be affecting the synapse indirectly (e.g. mitochondria, transport, etc.). Future work will now need to elucidate which synaptic processes are affected and how this has an impact on cholinergic projection neuron function and cholinergic transmission.

Abbreviations

DAN   Dopaminergic neurons

DEG   Differential expressed genes

ERG   Electroretinograms

KC   Kenyon cells

Nac   Nucleus accumbens

NBM   Nucleus basalis meynert

OPN   Olfactory projection neurons

PD   Parkinson's disease

Put   Putamen

SING   Startle induced negative geotaxis

Declarations

Ethical approval: Usage of post-mortem brain samples was ethically approved (EC reference NH019 2019-02-01).

Competing interests: Authors declare that they have no competing interests.

Funding: Research support was provided by ERC CoGs (PV and SA) and ERC AdG (PV), the Research Foundation Flanders (FWO), a Methusalem grant of the Flemish government, IMI2, Opening the Future and VIB. UP was supported by a fellowship from DFG; AB was supported by an EMBO long term fellowship and SV was supported by a fellowship from FWO. PV is an alumnus of the FENS Kavli Network of Excellence.

Acknowledgments
We thank Ann Geens, Eren Can Eksi, Vinoy Vijayan, Liesbeth Deaulmerie, Willem van den Bergh, Marianna Decet and Sara Aibar, Bart De Strooper, Joris de Wit, Roman Prashberger, the members of the Verstreken and Aerts labs and the VIB BioImaging core for help and comments. We thank Atefeh Pooryasin and the Bloomington Drosophila Stock Center (NIH P40OD018537) for fly lines. Human tissue was kindly provided by the Parkinson's UK Brain Bank at UCL, funded by Parkinson's UK, a charity registered in England and Wales (258197) and in Scotland (SC037554).

Author contributions:
Conceptualization: UP, SV, PV
Methodology: UP, JJ, NS, SK, SM, GH, SP, JS, SV
Investigation: UP, JJ, AB

Visualization: UP, JJ, PV

Funding acquisition: PV, SA, UP, SV, AB

Project administration: PV

Supervision: SA, PV

Writing – original draft: UP, PV

Writing – review & editing: all

Figure legends

Fig. 1. A collection of familial PD knock-in models, *related to Supplemental figure 1.*

(**a**) Scheme of the knock-in strategy where the first common exon in all *Drosophila* transcripts (left) was replaced by an attP-flanked mini-*white* gene using CRISPR/Cas9 mediated homologous recombination, creating a null mutant, and then replaced by wild-type or pathogenic mutant human or fly CDS using PhiC31 mediated integration (right). The chromosomal positions of the genes are indicated. All knock-ins are in the same isogenic genetic background (cantonized w^{1118}). (**b**) Phenotypic analysis of PD mutants (1) Median survival under starvation. n>50 animals per genotype. (2) Mitochondrial membrane potential measured by ratiometric TMRE fluorescence at neuromuscular junction (NMJ) boutons in 3^{rd} instar larvae. n≥4 animals per genotype and 10 boutons from ≥3 NMJs per animal. (3) Depolarisation amplitude quantified from electroretinograms (ERGs), recorded after 7 days of light exposure. n≥20 animals per genotype. (4) Quantification of startle-induced-negative geotaxis (SING) at 5±1 d after eclosion (young) and 25±1 d after eclosion (old). 95% of *pink1* mutants died < 25 d and were tested at 15 d. Values are normalized to control (see methods). Variance of control measurements are in grey. Bars are mean ± s.e.m; *, p<0.05 ANOVA/Dunnett's test.

Fig. 2. Single-cell RNA sequencing reveals that olfactory projection neurons are impaired across young PD fly models, *related to Supplemental figure 2.*

(**a**) tSNE of 226k cells characterized by single-cell RNA sequencing of entire young (5 day old) fly brains (controls and PD knock-in mutants). Colors indicate the cell types; 186 of which were identified (See methods). Key cell types are encircled, including dopaminergic neurons (DAN, blue), Mushroom body Kenyon cells (orange) and olfactory projection neurons (OPN, green). (**b-f**) tSNE of the cells from 5 selective PD knock-in mutants (5 day old). Black cells are those with

a significant transcriptomic change. Key cell types labeled in (a) are indicated. Note that OPN are consistently affected across mutants, while DAN are not at this early stage.

Fig. 3. Common DEGs in cholinergic neuron-rich brain regions of human PD patients and PD fly models, *related to Supplemental figure 3*.

(**a-a"**) Schematic of the sunburst plot indicating GO terms for each sector (**a**) and the mapping of the DEGs in *nucleus basalis of Meynert (NMB), nucleus accumbens* and *putamen* brain samples idiopathic PD patients (with *LRRK2* risk mutations) and controls (**a'**) and mapping of the DEGs found commonly in fly and human samples (**a"**). (**b**) Gene Ontology analysis of DEG in cholinergic neurons of young PD fly models and NBM neurons of PD patients. Redundant terms were removed. Color: adjusted p-value. (**c**) Schematic of the DEGs found commonly in fly PD models (blue) and human PD samples (black) manually sorted according to their previously described synaptic functions.

Fig. 4. Synaptic defects in OPN of PD mutants.

(**a**) Schematic of a head-fixed awake fly for live Ca^{2+}-imaging through a window in the head capsule. (**b**) Confocal image of GCaMP3-fluorescence expressed in OPN using GH146-Gal4, indicating the locations of the cell bodies, antennal lobe and the calyx (Scale bar: 50 µm). (**c-d**) Confocal image of the stimulus-induced change of fluorescence (peak amplitude) in the synaptic region of the calyx of a control and a *hLRRK2^{G2019S}* knock-in animal (c) and quantification of fluorescence change ($\pm$ SEM) over time (d); Arrowhead: time of stimulus application (10 mM Nicotine, See methods). (**e, f**) Images of GFP fluorescence marking the synaptic area of OPN in the calyx of control (e) and *hLRRK2^{G2019S}* knock-ins (f); Scale bar is 20 µm. (**g**) Quantification of GCaMP3 peak amplitude at OPN synapses in the calyx following stimulation (10 mM Nicotine) in controls, wild-type CDS knock-ins, and in the PD knock-in mutants where the wild-type CDS is not (-) or is (+) expressed in OPN (*OPN>wt CDS*). (**h**) Quantification of the GFP fluorescence area of OPN synapses in the calyx (based on GCaMP3 signal) in controls, wild-type CDS knock-ins, and in the PD knock-in mutants where the wild-type CDS is not (-) or is (+) expressed in OPN (*OPN>wt CDS*). For (g, h): n≥5 animals per genotype. *, p<0.05 in ANOVA/Dunnett.

Fig. 5. OPN dysfunction causes hyposmia in PD mutants.

(**a, b**) Olfactory performance of PD knock-in flies and controls when given the choice between a blend of motor oil and banana odors (a) or a blend of beer and wine odors (b). (**c, d**) Olfactory performance of PD knock-in flies and $LRRK^{KO}$ when the relevant wildtype CDS is not (-) or is (+) expressed selectively in OPN (or T1 cholinergic interneurons, T1>hR39B, as a negative control). For (a-d): Bars are mean ± s.e.m. n≥5 assays, ≥200 flies per genotype. *, p<0.05 in ANOVA/Dunnett.

Methods

Resource availability

Lead contact

For further information and requests for resources and reagents contact Patrik Verstreken (patrik.verstreken@kuleuven.be).

Material availability

Plasmids and fly lines generated in this study are available on request.

Data and code availability

Codes and data sets generated in this study are stored online https://www.ncbi.nlm.nih.gov/ geo/query/acc.cgi?acc=GSE235332.

Experimental model details

The "control" we refer to is a semi-isogenized w^{1118} backcrossed to CantonS for 10 generations (cantonized *w*). We sequenced the entire genome of the resulting fly line and defined there are no deleterious mutations in homologues of known causative Parkinson's disease genes or in homologues of human genes located close to GWAS loci. All our knock-ins (see below), UAS (see below) and Gal4 lines were backcrossed for >10 generations to this control. All generated fly lines were verified by multiple PCRs and PCR-product sequencing.

Flies were reared on standard yeast food medium at 25° C and kept on 12/12 h light/dark cycles. Flies were kept in mixed populations of analogous density and were flipped onto fresh food every three days. Brains for RNA experiments, including single cell sequencing, were dissected between 11 and 12 h Zeitgeber time, behavioral and physiology experiments were performed between 18 and 20 h Zeitgeber time. Males and females were tested together, quantified

separately, and results pooled if there was no statistical difference. Male hPINK1 pathogenic mutant knock-in flies are sterile, requiring balancing of the stock and consequently they produce only male hemizygotes to be quantified in experiments.

Generation of *Drosophila* PD models

A complete list of primers, oligos, gBlocks and fly strains used will be provided. To clone inserts into plasmids NEBuilder® HiFi DNA Assembly (NEB) was used.

Unique gRNAs (Supplemental File 1) for *LRRK2*, *PINK1*, *Rab39B, Aux* and *Synj* were identified by http://crispr.mit.edu/ and introduced into pCFD4: U6:1-gRNA U6:3-gRNA (Port et al., 2014) by following an established protocol: http://www.crisprflydesign.org/wp-content/uploads/2014/06/

Cloning-with-pCFD4.pdf.

Homology arms of 1 kB surrounding the first common exon of the different transcripts from the gene of interest were cloned into pWhite-STAR (Choi et al., 2009). For *Aux*, homology arms surrounded the entire *Aux* gene. HiFi DNA Assembly (NEB) was performed with four fragments: mini-white IMCE-cassette, left homology arm, right homology arm and the plasmid backbone. The homology arms were either generated by gBlocks (IDT), which were modified in order to make them compatible with the requirements of IDT, or amplified by Q5 PCR from genomic fly DNA of the target genotype (CantonS *w*). The mini-white IMCE-cassette and vector backbone were generated by restriction digest on pWhite-STAR (Choi et al., 2009) with AvrII and XhoI. In order to avoid cutting of the donor plasmid, the PAM sequence of the target was mutated, if located in the homology arms. Mostly NGG to NAA modifications were used, however different modifications were used when necessary (presence of PAM in coding region). The donor plasmids pDoC-miniwhite-gene (Donor cassette) contains an Integrase mediated exchange (IMCE) cassette surrounded by two homology arms that allows for HDR (homology directed repair). HDR occurs when two double strand breaks are introduced in the DNA surrounding the first (or second) common exon of all transcripts. This HDR replaced the first common exon by the IMCE-cassette. Site of HDR was directed by the choice of the homology arms in a way such that the two attP sites for IMCE were situated inside evolutionary non-conserved regions.

Rescue plasmids were used to exchange the mini-white IMCE-cassette with the coding DNA sequence (CDS) of target genes. All rescue plasmids contain: 1) a stretch of genomic sequence

that will be referred to as 'utr': this sequence was deleted together with the removed exon of the target gene and now is being placed back to repair the locus; 2) CDS of the target gene and 3) a poly(A)-tail. These 3 elements are surrounded by two attB sites. The rescue plasmids are named pReC-*gene* or pReC2-*gene* and are target gene-specific. The backbone: pReC was generated by linearizing pUC19 with SapI and EcoRI followed by insertion of two gBlocks: attB-MCS-L and MCS-attB-R. We also created pReC2, an optimized version that allows for easier cloning of the utr and CDS as it already contains the SV40 poly(A), by linearizing pUC19 with SapI and EcoRI followed by insertion of two gBlocks: attB-2xMCS-L and 2xMCS-attB-R.

LRRK2: pReC was linearized with SapI. gBlock LRRK2:utr-P2A-SapI-SV40, containing the utr a P2A site (since we knocked-out exon 2), SapI restriction site (for subsequent cloning of the CDS) and an SV40 poly(A)) was introduced. The resulting plasmid was linearized with SapI. Next the CDS of *LRRK2* was amplified from 2XMyc-LRRK2-WT with F_hLRRK2_cDNA and R_hLRRK2_cDNA and introduced into the linearized plasmid. The resulting plasmid is called pReC-hLRRK2. pReC-hLRRK2G2019S and pReC-hLRRK2Y1699C were created by using Q5® Site-Directed Mutagenesis (NEB) with primers F_mut_LRRK2_G2019S, R_mut_LRRK2_G2019S and Fmut_LRRK2_Y1699C, R_mut_LRRK2_Y1699C respectively on pReC-hLRRK2.

PINK1: pReC was linearized with SapI. gBlock: PINK1:utr-SapI containing the utr and SapI restriction site (for subsequent cloning of the CDS) is cloned into pReC. The resulting plasmid was linearized with AflIII. Next, the gBlock PINK1:SV40poly(A) containing the human Sv40 poly(A) was introduced into the linearized plasmid. This plasmid is linearized with SapI. Next the CDS of human *PINK1* was amplified from pLenti6-DEST PINK1-V5 WT with F_hPink1_cDNA and R_hPINK1_cDNA and introduced into the linearized plasmid. The resulting plasmid is called pReC-hPINK1. pReC-hPINK1R145Q and pReC-hPINK1L347P were created by using Q5® Site-Directed Mutagenesis (NEB) with primers F_mut_PINK1_L347P, R_mut_PINK1_L347P and F_mut_PINK1_P399L, R_mut_PINK1_P399L respectively on pReC-hPINK1. While we created wild-type CDS knock-ins for rescue experiments of all Parkinson's disease models, *Pink1* we used the expression of human *PINK1* with UAS-hPINK1 and nsyb-Gal4. This is because, unlike the two pathogenic variants, we were unsuccessful in targeting the wild-type human *PINK1* CDS to the fly *pink1* locus.

Synj: pReC was linearized with SapI. gBlock Synj_utr-SapI containing the utr and SapI restriction site (for subsequent cloning of the CDS) is cloned into pReC. The resulting plasmid was linearized with AflIII. gBlock Synj:SV40poly(A) containing the human Sv40 poly(A) was introduced into the linearized plasmid. This plasmid is linearized with SapI. Next the CDS of *Drosophila synj* and *synj^R258Q* were amplified from respectively UAS-Synj+ and UAS-SynjRQ (Vanhauwaert et al., 2017) with FW Synj and RC Synj and are introduced into the linearized plasmid by utilizing HiFi assembly. The resulting plasmids are called pReC-dSynj and pReC-dSynjR258Q.

Rab39B: pReC2 was linearized with XhoI and XbaI. A gblock: Rab39_utr was inserted into the pReC2. Next, this plasmid was linearized with SapI and a gBlock containing the human *RAB39B* CDS was cloned. The resulting plasmid is called pReC2-hRab39. pReC2-hRab39T168K and pReC2-hRab39G192R were created by using Q5® Site-Directed Mutagenesis (NEB) with primers F_mut_hRab39_T168K, R_mut_hRab39_T168K and F_mut_hRab39_G192R, R_mut_hRab39_G192R respectively on pReC2-hRab39.

Aux: The entire *Drosophila auxilin* gene region was cloned from BAC CH322-22D05 into pRec using BbsI, with an HA-tag at the N-terminal of the gene. AuxR927G was produced by Q5® Site-Directed Mutagenesis (NEB). See also (Jacquemyn et al., 2023).

UAS-aux: pUAST.attB was linearized with XhoI and XbaI; gBlock: Aux was cloned into the this plasmid. See also (Jacquemyn et al., 2023).

Constructs were injected in house or by BestGene (BestGene Inc, CA, US).

Quantitative RT-PCR

For each sample, 20 heads of 5 day (5 d) old males were collected, RNA was extracted and transcribed using Maxwell RSC instrument and Maxwell RSC RNA kit (Promega) and quantified by subsequent qPCR using a LightCycler 480, 480 SYBR Green master mix (Roche).

Survival assay

Flies were collected into populations of 10 males and 10 females the day after eclosion. Five days after eclosion, these populations were transferred into vials containing 1 % agarose for wet starvation. Dead flies were scored and removed every 8 h, the remaining flies were flipped on fresh agarose every 24 h. The assay was repeated with at least two different batches of a given genotype. Median survival and survival curves were determined using GraphPad Prism.

Mitochondrial membrane potential (TMRE) assay

Measurements of mitochondrial membrane potential were performed with the potentiometric dye tetramethyl rhodamine ethyl ester (TMRE). Male third-instar larvae were dissected in HL3 (in mM: 110 NaCl, 5 KCl, 10 MgCl2 • 6H2O, 10 NaHCO3, 30 sucrose, 5 trehalose, and 10 HEPES, pH 7.2). Larval fillets were incubated for 15 min in the presence of 50 nM TMRE (Abcam). Subsequently, the TMRE solution was discarded and fillets were rinsed three times with normal HL3 without TMRE. Mitochondrial labeling of TMRE was imaged on a Nikon spinning disk confocal microscope with a 40x water dipping objective 0.8 NA, excitation wavelength of 561 nm and an emission filter 595/50 nm. We imaged at 0.5 μm Z-steps entire NMJ branches. Larvae were stained a mutant and control larva were processed and imaged side-by-side. TMRE labeling intensity was quantified in individual synaptic boutons using Fiji (Schindelin et al., 2012). Because of variability between sessions, every mutant was normalized to the controls stained and measured in the same session.

Electroretinograms (ERG)

ERGs were recorded as previously described (Slabbaert et al., 2016). Flies were immobilized on glass microscope slides, by use of liquid Pritt glue. For recordings, glass electrodes (borosilicate, 1.5 mm outer diameter) filled with 3 M NaCl were placed in the thorax as a reference and on the fly eye for recordings. Responses to repetitive 1 s light stimuli were recorded using Axosope 10.7 and analyzed using Clampfit 10.7 software (Molecular Devices) and Igor Pro 6.37. To assess the ERG response after exposure to constant light, 3 day old flies were placed under continuous illumination (1300 lux) for 7 days at 25 °C prior to ERG data acquisition as described.

Startle-induced negative geotaxis

Geotaxis locomotion was quantified in mixed populations of ~20 flies (Benzer, 1967) with modifications described by (Inagaki et al., 2010). Flies were flipped into the apparatus without anesthesia and allowed to adjust for some minutes to experimental environment (24° C, 50 % humidity). They were tapped down into the first lower tube and allowed to climb into the upper tube for 30 s, at which time the flies that reached the upper tube were moved to tube 2. The procedure was repeated five times and the SING scored: (#flies in tube 1 + (#flies in tube 2 * 2) + (#flies in tube 3 * 3) + (#flies in tube 4 *4) + (#flies in tube 5 * 5)) / (#total flies *5).

Modeling to define the number of cells to be sequenced

We randomly sampled between 0 and 50,000 cells, in steps of 100, from the annotated brain cell dataset generated in (Davie et al., 2018). For each sample we calculated the number of cell types that were retrieved for different thresholds of detection. This process was repeated 50 times for convergence and Michaelis-Menten kinetics were fit through the data points.

Brain dissection for single cell sequencing

30 fly brains (15 females and 15 males, except *Pink1* mutants, where only males were used) were dissected on ice and collected in 100 µl ice cold DPBS solution, centrifuged at 800 x *g* for 5 min, and placed into 50 µL of dispase (3 mg/mL, Sigma-Aldrich_D4818-2mg) and 75 µl collagenase I (100 mg/mL, Invitrogen_17100-017). Brains were dissociated in a Thermoshaker (Grant Bio PCMT) for 2 h at 25 °C and 500 rpm. The enzymatic reaction was reinforced by pipetting every 15 min. Subsequently, cells were washed with 1000 µl ice cold DPBS solution and resuspended in 400 µl DPBS 0.04 % BSA. Cell suspensions were passed through a 10 µM pluriStrainer (ImTec Diagnostics_435001050), cell viability and concentration were assessed by the LUNA-FL™ Dual Fluorescence Cell Counter, and cells were immediately used for the 10x run.

10x Genomics high-throughput sequencing

Before sequencing, the fragment size of every library was analyzed on a Bioanalyzer high sensitivity chip. The libraries were diluted to 2 nM and quantified by qPCR using primers against p5-p7 sequence. 10x libraries were sequenced twice: shallow on NextSeq500 (Illumina) to determine library quality and more in depth using NovaSeq6000 (Illumina). The targeted saturation for each sample was around 60%, and additional sequencing runs were performed if

needed. Each sequencing run used the following sequencing parameters: 28 bp read 1 – 8 bp index 1 (i7) – 91 bp read 2.

10x Data Preprocessing

The 10x fly brain samples were each processed (alignment, barcode assignment and UMI counting) with Cell Ranger (version 3.0.2) count pipeline. The Cell Ranger reference index was built upon the 3[rd] 2017 FlyBase release (*D. melanogaster* r6.16) (Gramates et al., 2017). To count the human mutated genes, sequences from the construct were added as artificial chromosomes to the index. For every sample all sequencing runs were aggregated using CellRanger and the *–recovered-cells* parameter was specified as 5000.

Data filtering and clustering

CellRanger generated a gene expression matrix, which was used as input for VSN pipelines (Zenodo, v0.27.0) (Flerin et al., 2021) for the first filtering and data clean up. These pipelines are written in Nextflow DSL2 making them highly reproducible, and the config file used for the fly samples can be found in Supplemental File 2. The VSN pipeline removed predicted doublets using scrublet (Wolock et al., 2019), followed by a cleaning of ambient RNA reads using DecontX (S. Yang et al., 2020), similar to the procedures used in the Fly Cell Atlas (Li et al., 2022).

Afterwards the data was loaded into scanpy (1.8.2, (Jenett et al., 2012)). Cells were filtered using min_genes=200, max_genes=7000, min_counts=500 and max_counts=30000, and finally with percentage of mitochondrial counts below 15. The data were log-normalized with a scale factor of 10^4 and scaled (maximum values = 10), regressing out the number of UMIs and the percentage of mitochondrial genes as latent variables using a linear regression model. Highly variable genes were selected using sc.pp.highly_variable_genes with default parameters.

We used sc.tl.pca to calculate 250 principal components on the highly variable genes, from which the first 100 components were selected for follow-up analysis based on an elbow plot. Since the data contains many samples (n=65) covering a variety of different ages and genotypes, we noticed batch effects and low mixing with default parameters. Therefore, we used Harmony [ref] to correct the principal components. For this we used sce.pp.harmony_integrate with max_iter_harmony=20, using the genotype + age combination as batch variable. The batch effect-corrected components were then used as input to calculate neighbors using

sc.pp.neighbors with n_neighbors=30, followed by Leiden clustering for resolutions between 2 and 20. Subsequently, tSNE and UMAP visualizations were calculated.

Seurat

Dimensionality reduction and clustering were performed using the Seurat package in R (Satija et al., 2015). The processed and filtered matrices of all samples were combined together with an annotated dataset of 57k cells of the adult brain containing two wild-type genotypes (Davie et al., 2018), leading to a total number of ~170k cells. In the first steps of the Seurat pipeline, each genotype was processed separately. The data were log-normalized with a scale factor of 104 and scaled, regressing out the number of UMIs and the percentage of mitochondrial genes as latent variables using a negative binomial model. For further downstream analysis, the 2000 most variable genes were selected using FindVariableGenes and method='vst' with default parameters. Then 3000 integration features were determined across all genotypes with SelectIntegrationFeatures and used in the rest of the analysis. At this point we looped over a number of components to test the dimensionality of the dataset, ranging between 30 and 100 components. These components were used as input for the integration functions, PCA, tSNE and clustering. Integration was performed by linking cells between datasets (so-called anchors, function FindIntegrationAnchors), followed by integrating the datasets one by one (IntegrateData). The scaled and integrated matrix was used as input for PCA to reduce the dimensions. The principal components were then used to calculate the non-linear dimensionality reduction tSNE (RunTSNE), and to perform clustering with Louvain Shared-Nearest Neighbor (SNN) graph algorithm (FindNeighbors, FindClusters). Different resolutions ranging between 1 and 8 were tested. In general, results are only slightly affected by the number of components used (Seurat tutorial), so we decided to compare the tSNE and clusters qualitatively, settling on 70 as the final number.

Label transfer

In the previous step different resolutions were used in clustering, leading to an increasing number of clusters detected. To start annotating these and to find a final clustering resolution, we compared the detected clusters with the annotations from the co-clustered published dataset (Davie et al., 2018; Jenett et al., 2012) and annotations from the Fly Cell Atlas (Li et al., 2022). In addition, we also used the neural net classifier (Davie et al., 2018) using default parameters as

described. Finally, we compared the clusters with sorted cell types from (Davis et al., 2020) using lasso regression as described below. These data were used to annotate clusters in resolution 20, after which clusters with the same annotation were merged. For unknown cells, we chose to use Leiden resolution 5 as basis annotation to not split clusters too much and keep enough cells to perform meaningful differential expression analysis.

In total we found 188 clusters, of which 83 were annotated (median size = 1,011 cells, total = 140,778 cells) and 105 unknown (median size = 810 cells, total = 145,010 cells).

Lasso regression

To expand the published atlas, we analyzed the bulk transcriptomes of multiple FAC-sorted neuronal cell types (Davie et al., 2018). Regression models have been previously used to compare bulk transcriptome profiles to aggregated cluster profiles, including non-negative least squares (Davie et al., 2018). Here we use lasso regression to predict single cell cluster transcriptome profiles using the bulk profiles as variables. Therefore, each single cell cluster is decomposed as a weighted sum of bulk profiles. The regularization of the lasso model, allows for a higher signal to noise ratio and clearer assignments. Clear matches between a cluster and a bulk profile are detected as high coefficients (>0.1) in the sum. The regression model was fit using gene signatures derived from Seurat as features. Before fitting, all datasets were scaled to counts per million CPM using the common genes, independent of previously applied normalization techniques. The calculations were performed using the SciPy (Virtanen et al., 2020) and Scikit-learn packages in Python.

Human brain samples

Usage of post-mortem brain samples was ethically approved (EC reference NH019 2019-02-01). All tissue samples were obtained from the Parkinson's UK Brain Bank at the Imperial College London. The Parkinson's disease cases used in this study all carried a mutation in the *LRRK2* gene (P1542S, M1646T or R1325Q). One case carried a mutation in *PARKIN* in addition and one case carried a mutation in *EIF4G* in addition. Control samples are from cases without pre- or post mortem Parkinson's disease-associated pathology and diagnosis.

Human nuclei isolation

Nuclei of human samples were prepared as described in (Hodge et al., 2019; Slyper et al., 2020).

Calling SNPs

In the human data, 10 different genotypes (5 PD cases, 5 controls) were mixed across 3 regions. To perform demultiplexing, the SNPs are required per sample. Since no genotype information was available, we performed bulk RNA-seq on all samples and called SNPs on the transcriptome. For this we used BCFFtools (v1.11, (Li et al., 2022)). We performed a pileup using the common loci of the 1000 Genomes project (1000 Genomes Project Consortium et al., 2015; Gramates et al., 2017) on the Hg38 iGenomes genome (-d 8000, -Ou), calling SNPs with the bcftools call function (-mv, -Ob). Using the reheader function we then updated the bcf file to make it compatible with demuxlet. We detected 146,848 SNPs, ranging between 25k and 51k per genotype (see table). Next, the bcf file was sorted in the same order as the bam files and bcftools +fill-tags was used to add frequencies.

Genotype	SNPs
C23	25,222
C45	41,379
C72	37,607
C73	40,651
C87	48,896
PD106	38,281
PD188	62,987
PD194	46,071
PD348	44,195
PD423	51,902

Demuxlet

Bam files were filtered using SAMtools (1.11, (Li et al., 2022)) to only contain reads overlapping with the vcf file. Then popscle dsc-pileup was used to create a pileup file for input in popscle demuxlet (Kang et al., 2018). All cells labeled as doublets were removed (~4-7 %), and cells assigned to singlets were kept (25-43 %). Additionally, a large proportion of cells (~49-69 %) were labelled as ambiguous. These cells were assigned to their most likely singlet identity

and kept in the remaining analysis. In a next step, we only kept possible genotype-experiment combinations, ending up with 97,489 cells.

Data filtering and clustering

CellRanger generated a gene expression matrix of 151,368 cells. After removing doublets and wrongly assigned cells, 97,489 cells remained. The data was loaded into scanpy (Wolf et al., 2018) and further filtered using min_genes=500, max_genes=7000, min_counts=1000 and max_counts=30000, and finally with percentage of mitochondrial counts below 5. We lowered the allowed percentage of mitochondrial counts since the human data was obtained from snRNA-seq. The data were log-normalized with a scale factor of 10^4 and scaled (maximum values = 10), regressing out the number of UMIs and the percentage of mitochondrial genes as latent variables using a linear regression model. Highly variable genes were selected using sc.pp.highly_variable_genes with default parameters.

We used sc.tl.pca to calculate 200 principal components on the highly variable genes, from which the first 75 components were selected for follow-up analysis based on an elbow plot. We again used Harmony [ref] to correct the principal components for batch effects. The batch effect-corrected components were then used as input to calculate neighbors using sc.pp.neighbors with n_neighbors=30, followed by Leiden clustering for resolutions between 1 and 10. Subsequently, tSNE and UMAP visualizations were calculated.

Using marker genes for main neuronal and glial populations we chose to combine Leiden resolution 1 for glia and Leiden resolution 3 for neurons.

Gene set enrichment was performed using the AUCell function from the pyscenic package (0.10.4) in Python. The genes detected in affected cell types of *Drosophila* models were converted to human homologues using the FlyBase database, selecting homologues with a DIOPT score>=5. Significant enrichment was determined using a two-tailed t-test.

Differential expression

For every cluster, cells from each mutant genotype were compared against cells from the wild-type control. In the fly, OPN subtypes were merged before calculating differentially expressed

genes (DEG). To control for biases in methods, we used Wilcoxon and DESEQ2 (Love et al., 2014). The Wilcoxon test was performed with the scanpy wrapper sc.tl.rank_genes_groups. For DESEQ2 we used rpy2 to call the R package DESEQ2 in Python. DESEQ2 was originally designed to be used on bulk RNA-seq data. Therefore, we grouped cells into pseudobulks based on their experimental run by summing their gene expression profiles. Only pseudobulks consisting of at least 10 cells were kept. We did not select any additional variables to regress.

To detect any method biases, we calculated the spearman correlation of the signed p-value (log-foldchange multiplied with log(p-val)), showing that both methods differ in the exact number of DEG, but agree quite well in their ranking of up- and downregulated genes. We found larger differences in the number of DEG called in the human data (0-10 in DESEQ2, 200-1800 in Wilcoxon) but the rankings remained highly correlated (r~0.8). The Wilcoxon test is more sensitive to genes expressed in all cells in the cluster, while DESEQ2 can be oversensitive to genes expressed in only a minor fraction of cells.

Additional runs were made in which the number of cells per cluster was down sampled to correct against cluster-size based biases.

Modeling of DEG

Since the number of DEG has been shown to be highly correlated with the number of cells in single-cell RNA-seq, we used a modeling approach to find outlier clusters with more than expected DEG. Since the data is over dispersed count data, we fit a negative binomial model to the data, with number of differential genes (nDEG) as dependent variable and number of cells in the mutant and number of cells in the control as independent variables:

$$E[y_i|\alpha, \mu_i] = \mu_i$$
$$VAR[y_i|\alpha, \mu_i] = \mu_i + \alpha\mu_i^2$$
$$\log(\mu_i) = Intercept + B_1 * \log(\min(X_{1i}, X_{2i}))$$

With y_i being the number of differential genes detected in cluster i, X_{1i} the log of the number of cells of cluster i in the mutant and X_{2i} the number of cells of cluster i in the wild type.

Models were fit using the statsmodels package in Python and goodness of fit was measured using the Log-Likelihood (Supplemental File 3). The number of UMIs per cluster was not included as independent variable, since it has been implicated with cell size and thus could remove biological effects. Residuals were calculated as the difference between the measured number of differential genes and the predicted number, wherein a positive residual means that the cluster has more differential genes than expected.

$$r_i = \widehat{\mu}_i - y_i$$

Olfactory behavior assay

Olfactory behavior was quantified in mixed populations of ~50 5±1 day old flies. Flies were flipped into the experimental setup without anesthesia and allowed to adjust for some minutes to the experimental environment (24 °C, 50 % humidity). They were placed into a custom-built T-Maze (modified from (Tully & Quinn, 1985)), where they were allowed to freely walk between two tubes for one minute. Throughout the test, an aquarium pump delivers two separated, constant airstreams (1 l/min) which are directed through an odor vial before entering the tube. Odor vials contained: 2 ml motor oil, 2 g fresh banana, 2 ml undiluted beer (Jupiler), or 2 ml wine (LesTruffiers, 2015) diluted 1:1000. Afterwards the olfactory preference index was scored: (#flies in tube with odor A - #flies in tube with odor B)/#total flies. Tests were pseudo-randomized by switching the position of odor tubes every trial. Control flies were tested each experimental session.

In vivo Ca^{2+}-imaging and synaptic area

5±1 day old (unless noted otherwise) male flies were placed into a custom-build chamber and the brains were imaged at 5 Hz through a window in the head capsule using a wide field fluorescence microscope (Nikon) equipped with a Hamamatsu camera and a 20×/NA=0.9 water-immersion objective. 100 mM nicotine was injected into the Ringer's solution covering the brain (final concentration ~10 mM). Images were analyzed using Fiji (Schindelin et al., 2012). Fluorescence intensity was quantified in a region of interest (ROI, d=10 μm) that was placed in the center of the synaptic area (calyx for OPNs, mushroom body lobes for dopaminergic PAM neurons) and normalized by subtracting background fluorescence. Basal fluorescence F_0 in the synaptic region was determined by averaging five frames before stimulus onset and the change

of fluorescence over time was calculated by $F-F_0/F_0$. Flies of related genotypes and controls were always measured within the same sessions, in alternating order.

The area of the calyx was determined by measuring the (2D) surface of the entire visible fluorescent area.

Quantification and statistical analysis

GraphPad Prism was used for visualization and statistical analysis of results from all but the sequencing experiments. Males and females were quantified separately, and results pooled later because they did not show significant differences. Data sets were checked for normal distribution and then analyzed with ANOVA and subsequent multiple pairwise comparisons using Dunnett's test when comparing to a general control, or Bonferroni-correction of p-values when comparing different control-test pairs.

Key resources table

REAGENT RESOURCE	SOURCE	IDENTIFIER
Bacterial and Virus Strains		
Oneshot top10 cells	ThermoFisher Scientific	C404010
Chemicals, Peptides, and Recombinant Proteins		
Ringer's solution	(Estes et al., 1996)	
TMRE	Abcam, Cambridge, UK	ab113852
Dispase	Sigma Aldrich	D4818
Collagenase	Invitrogen, CA, US	17100-017
Trypsin-EDTA	Invitrogen, CA, US	25300054
Nicotine	Sigma Aldrich	N3876
Critical Commercial Assays		
NEBuilder® HiFi DNA Assembly	New England Biolabs, Inc (NEB)	NEB #E2621
Maxwell RSC instrument and kit	Promega	AS8500, AS1340
Chromium Single Cell 3' Library & Gel Bead Kit v2	10x Genomics	PN-120237
Chromium Single Cell A	10x Genomics	PN-1000009

Chip Kit		
Chromium i7 Multiplex Kit	10x Genomics	PN-120262
Chromium Next GEM Single Cell 3' Kit v3	10x Genomics	PN-1000075
Chromium Next GEM Chip B Single Cell Kit	10x Genomics	PN-1000073
Single Index Kit T Set A	10x Genomics	PN-1000213
Experimental Models: Fly Strains		
GH146-Gal4 (OPN-Gal4)	(Stocker et al., 1997)	BDSC_30026; FlyBase: FBti0016783
GMR31F10-Gal4 (T1-Gal4)	(Jenett et al., 2012)	BDSC_49685; FlyBase:
GMR58E02-Gal4 (PAM-Gal4)	(Jenett et al., 2012)	BDSC_41347;FlyBase: FBti0137105
UAS-GCaMP3	(Tian et al., 2009)	BDSC_32236; BDSC_32116
UAS-hLRRK2	(Venderova et al., 2009)	
UAS-hPINK1	(Yang et al., 2006)	
UAS-YFP-Rab39	(Zhang et al., 2007)	BDSC_9825; FlyBase: FBti0100878
UAS-Synj	(Vanhauwaert et al., 2017)	
CG5010$^{k.o.}$ (*chchd2$^{k.o.}$*)	this study	
LRRK$^{k.o.}$	this study	
hLRRK2	this study	
hLRRK2^{G2019S}	this study	
hLRRK2^{Y1699C}	this study	
PINK1$^{k.o.}$	this study	
hPINK1^{P399L}	this study	
hPINK1^{L347P}	this study	
Rab39$^{k.o.}$	this study	
hRab39B	this study	
hRab39B^{G192R}	this study	
hRab39B^{T168K}	this study	
Aux$^{k.o.}$	this study	
hDNAJC6	this study	
Aux^{wt}	this study	

Aux^{R927G}	this study	
Synj$^{K.O.}$	this study	
hSynj1	this study	
Synj^{WT}	this study	
Synj^{R258Q}	this study	
Plasmids		
pCFD4: U6:1-gRNA U6:3-gRNA	(Port et al., 2014)	RRID:Addgene_49411
pUC19	(Norrander et al., 1983)	RRID:Addgene_50005
pUAST.attB	(Bischof et al., 2007)	GenBank:EF362409.1
pFL44S{w+}-attB	(Khuong et al., 2013)	N/A
pWhite-STAR	(Choi et al., 2009)	N/A
2XMyc-LRRK2-WT	(Greggio et al., 2008)	RRID:Addgene_25361
pLenti6-DEST PINK1-V5 WT	(Beilina et al., 2005)	RRID:Addgene_13320
Software and Algorithms		
Fiji	(Schindelin et al., 2012)	https://imagej.net/Fiji
Cell Ranger (version 3.0.2)	10xGenomics	https://support.10xgenomics.com/single-cell-gene-expression/software/downloads/latest
Seurat	(Satija et al., 2015)	
GOrilla	(Eden et al., 2009)	http://cbl-gorilla.cs.technion.ac.il/
GraphPad Prism	GraphPad software, CA, US	https://www.graphpad.com/scientific-software/prism/
Other and deposited data		
list of oligonucleotides		Supplemental File 1
list of software used		Supplemental File 6
SCope Resource Website	(Davie et al., 2018) and this study	http://scope.aertslab.org
full fly single-cell-RNA-seq data	this study	https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE228843
full human single-cell-RNA-seq data	this study	https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE235332

References

1000 Genomes Project Consortium, Auton, A., Brooks, L. D., Durbin, R. M., Garrison, E. P., Kang, H. M., Korbel, J. O., Marchini, J. L., McCarthy, S., McVean, G. A., & Abecasis, G. R. (2015). A global reference for human genetic variation. *Nature*, *526*(7571), 68–74. https://doi.org/10.1038/nature15393

Ansari, K. A., & Johnson, A. (1975). Olfactory function in patients with Parkinson's disease. *Journal of Chronic Diseases*, *28*(9), 493–497. https://doi.org/10.1016/0021-9681(75)90058-2

Arendt, T., Bigl, V., Arendt, A., & Tennstedt, A. (1983). Loss of neurons in the nucleus basalis of Meynert in Alzheimer's disease, paralysis agitans and Korsakoff's disease. *Acta Neuropathologica*, *61*(2), 101–108. https://doi.org/10.1007/bf00697388

Ashrafi, G., Schlehe, J. S., LaVoie, M. J., & Schwarz, T. L. (2014). Mitophagy of damaged mitochondria occurs locally in distal neuronal axons and requires PINK1 and Parkin. *The Journal of Cell Biology*, *206*(5), 655–670. PubMed. https://doi.org/10.1083/jcb.201401070

Beilina, A., Van Der Brug, M., Ahmad, R., Kesavapany, S., Miller, D. W., Petsko, G. A., & Cookson, M. R. (2005). Mutations in PTEN-induced putative kinase 1 associated with recessive parkinsonism have differential effects on protein stability. *Proceedings of the National Academy of Sciences of the United States of America*, *102*(16), 5703–5708. https://doi.org/10.1073/pnas.0500617102

Benzer, S. (1967). BEHAVIORAL MUTANTS OF Drosophila ISOLATED BY COUNTERCURRENT DISTRIBUTION. *Proceedings of the National Academy of Sciences of the United States of America*, *58*(3), 1112–1119. PubMed. https://doi.org/10.1073/pnas.58.3.1112

Bischof, J., Maeda, R. K., Hediger, M., Karch, F., & Basler, K. (2007). An optimized transgenesis system for Drosophila using germ-line-specific phiC31 integrases. *Proceedings of the National Academy of Sciences of the United States of America*, *104*(9), 3312–3317. https://doi.org/10.1073/pnas.0611511104

Bohnen, N. I., Müller, M. L. T. M., Kotagal, V., Koeppe, R. A., Kilbourn, M. A., Albin, R. L., & Frey, K. A. (2010). Olfactory dysfunction, central cholinergic integrity and cognitive impairment in Parkinson's disease. *Brain : A Journal of Neurology*, *133*(Pt 6), 1747–1754. PubMed. https://doi.org/10.1093/brain/awq079

Braak, H., Rüb, U., Gai, W. P., & Del Tredici, K. (2003). Idiopathic Parkinson's disease: Possible routes by which vulnerable neuronal types may be subject to neuroinvasion by an unknown pathogen. *Journal of Neural Transmission*, *110*(5), 517–536. https://doi.org/10.1007/s00702-002-0808-2

Braak, H., Tredici, K. D., Rüb, U., de Vos, R. A. I., Jansen Steur, E. N. H., & Braak, E. (2003). Staging of brain pathology related to sporadic Parkinson's disease. *Neurobiology of Aging*, *24*(2), 197–211. https://doi.org/10.1016/s0197-4580(02)00065-9

Brimblecombe, K. R., Threlfell, S., Dautan, D., Kosillo, P., Mena-Segovia, J., & Cragg, S. J. (2018). Targeted Activation of Cholinergic Interneurons Accounts for the Modulation of Dopamine by Striatal Nicotinic Receptors. *eNeuro*, *5*(5), ENEURO.0397-17.2018. https://doi.org/10.1523/ENEURO.0397-17.2018

Brooker, S. M., Naylor, G. E., & Krainc, D. (2024). Cell biology of Parkinson's disease: Mechanisms of synaptic, lysosomal, and mitochondrial dysfunction. *Current Opinion in Neurobiology*, *85*, 102841. https://doi.org/10.1016/j.conb.2024.102841

Chambers, R. P., Call, G. B., Meyer, D., Smith, J., Techau, J. A., Pearman, K., & Buhlman, L. M. (2013). Nicotine increases lifespan and rescues olfactory and motor deficits in a Drosophila model of Parkinson's disease. *Behavioural Brain Research*, *253*, 95–102. https://doi.org/10.1016/j.bbr.2013.07.020

Chase, B. A., & Markopoulou, K. (2020). Olfactory Dysfunction in Familial and Sporadic Parkinson's Disease. *Frontiers in Neurology*, *11*, 447–447. PubMed. https://doi.org/10.3389/fneur.2020.00447

Choi, C. M., Vilain, S., Langen, M., Van Kelst, S., De Geest, N., Yan, J., Verstreken, P., & Hassan, B. A. (2009). Conditional Mutagenesis in Drosophila. *Science*, *324*(5923), Article 5923. https://doi.org/10.1126/science.1168275

Chu, Y., Dodiya, H., Aebischer, P., Olanow, C. W., & Kordower, J. H. (2009). Alterations in lysosomal and proteasomal markers in Parkinson's disease: Relationship to alpha-synuclein inclusions. *Neurobiology of Disease*, *35*(3), 385–398. https://doi.org/10.1016/j.nbd.2009.05.023

Davie, K., Janssens, J., Koldere, D., De Waegeneer, M., Pech, U., Kreft, Ł., Aibar, S., Makhzami, S., Christiaens, V., Bravo González-Blas, C., Poovathingal, S., Hulselmans, G., Spanier, K. I., Moerman, T., Vanspauwen, B., Geurs, S., Voet, T., Lammertyn, J., Thienpont, B., … Aerts, S. (2018). A Single-Cell Transcriptome Atlas of the Aging Drosophila Brain. *Cell*, *174*(4), 982-998.e20. PubMed. https://doi.org/10.1016/j.cell.2018.05.057

Davis, F. P., Nern, A., Picard, S., Reiser, M. B., Rubin, G. M., Eddy, S. R., & Henry, G. L. (2020). A genetic, genomic, and computational resource for exploring neural circuit function. *eLife*, *9*, e50901. PubMed. https://doi.org/10.7554/eLife.50901

Doty, R. L. (2012). Olfactory dysfunction in Parkinson disease. *Nature Reviews Neurology*, *8*(6), 329–339. https://doi.org/10.1038/nrneurol.2012.80

Doty, R. L., Deems, D. A., & Stellar, S. (1988). Olfactory dysfunction in parkinsonism. *Neurology*, *38*(8), 1237–1237. https://doi.org/10.1212/wnl.38.8.1237

Double, K. L., Rowe, D. B., Hayes, M., Chan, D. K. Y., Blackie, J., Corbett, A., Joffe, R., Fung, V. S., Morris, J., & Halliday, G. M. (2003). Identifying the Pattern of Olfactory Deficits in Parkinson Disease Using the Brief Smell Identification Test. *Archives of Neurology*, *60*(4), 545. https://doi.org/10.1001/archneur.60.4.545

Eden, E., Navon, R., Steinfeld, I., Lipson, D., & Yakhini, Z. (2009). GOrilla: A tool for discovery and visualization of enriched GO terms in ranked gene lists. *BMC Bioinformatics*, *10*, 48–48. PubMed. https://doi.org/10.1186/1471-2105-10-48

Estes, P. S., Roos, J., van der Bliek, A., Kelly, R. B., Krishnan, K. S., & Ramaswami, M. (1996). Traffic of dynamin within individual Drosophila synaptic boutons relative to compartment-specific markers. *The Journal of Neuroscience: The Official Journal of the Society for Neuroscience*, *16*(17), 5443–5456. https://doi.org/10.1523/JNEUROSCI.16-17-05443.1996

Fares, M. B., Alijevic, O., Johne, S., Overk, C., Hashimoto, M., Kondylis, A., Adame, A., Dulize, R., Peric, D., Nury, C., Battey, J., Guedj, E., Sierro, N., Mc Hugh, D., Rockenstein, E., Kim, C., Rissman, R. A., Hoeng, J., Peitsch, M. C., … Mathis, C. (2023). Nicotine-mediated effects in neuronal and mouse models of synucleinopathy. *Frontiers in Neuroscience*, *17*, 1239009. https://doi.org/10.3389/fnins.2023.1239009

Flerin, C., Davie, K., Hulselmans, G., & De Waegeneer, M. (n.d.). *Flerin C, Chris, Davie K, Hulselmans G, De Waegeneer M. vib-singlecell-nf/vsn-pipelines: V0.27.0 (v0.27.0). Zenodo. Https://doi.org/10.5281/zenodo.5751297.* [Computer software].

Friedman, L. G., Lachenmayer, M. L., Wang, J., He, L., Poulose, S. M., Komatsu, M., Holstein, G. R., & Yue, Z. (2012). Disrupted autophagy leads to dopaminergic axon and dendrite degeneration and promotes presynaptic accumulation of α-synuclein and LRRK2 in the brain. *The Journal of Neuroscience : The Official Journal of the Society for Neuroscience*, *32*(22), 7585–7593. PubMed. https://doi.org/10.1523/JNEUROSCI.5809-11.2012

Gorenberg, E. L., & Chandra, S. S. (2017). The Role of Co-chaperones in Synaptic Proteostasis and Neurodegenerative Disease. *Frontiers in Neuroscience*, *11*, 248–248. PubMed. https://doi.org/10.3389/fnins.2017.00248

Gramates, L. S., Marygold, S. J., Santos, G. D., Urbano, J.-M., Antonazzo, G., Matthews, B. B., Rey, A. J., Tabone, C. J., Crosby, M. A., Emmert, D. B., Falls, K., Goodman, J. L., Hu, Y., Ponting, L., Schroeder, A. J., Strelets, V. B., Thurmond, J., Zhou, P., & the FlyBase Consortium. (2017). FlyBase at 25: Looking to the future. *Nucleic Acids Research*, *45*(D1), D663–D671. PubMed. https://doi.org/10.1093/nar/gkw1016

Greggio, E., Zambrano, I., Kaganovich, A., Beilina, A., Taymans, J.-M., Daniëls, V., Lewis, P., Jain, S., Ding, J., Syed, A., Thomas, K. J., Baekelandt, V., & Cookson, M. R. (2008). The Parkinson disease-associated leucine-rich repeat kinase 2 (LRRK2) is a dimer that undergoes intramolecular autophosphorylation. *The Journal of Biological Chemistry*, *283*(24), 16906–16914. https://doi.org/10.1074/jbc.M708718200

Haehner, A., Boesveldt, S., Berendse, H. W., Mackay-Sim, A., Fleischmann, J., Silburn, P. A., Johnston, A. N., Mellick, G. D., Herting, B., Reichmann, H., & Hummel, T. (2009). Prevalence of smell loss in Parkinson's disease – A multicenter study. *Parkinsonism & Related Disorders*, *15*(7), 490–494. https://doi.org/10.1016/j.parkreldis.2008.12.005

Harvey, J. D., & Heinbockel, T. (2018). Neuromodulation of Synaptic Transmission in the Main Olfactory Bulb. *International Journal of Environmental Research and Public Health*, *15*(10), 2194. PubMed. https://doi.org/10.3390/ijerph15102194

Hindle, S., Afsari, F., Stark, M., Middleton, C. A., Evans, G. J. O., Sweeney, S. T., & Elliott, C. J. H. (2013). Dopaminergic expression of the Parkinsonian gene LRRK2-G2019S leads to non-autonomous visual neurodegeneration, accelerated by increased neural demands for energy. *Human Molecular Genetics*, *22*(11), 2129–2140. PubMed. https://doi.org/10.1093/hmg/ddt061

Hodge, R. D., Bakken, T. E., Miller, J. A., Smith, K. A., Barkan, E. R., Graybuck, L. T., Close, J. L., Long, B., Johansen, N., Penn, O., Yao, Z., Eggermont, J., Höllt, T., Levi, B. P., Shehata, S. I., Aevermann, B., Beller, A., Bertagnolli, D., Brouner, K., … Lein, E. S. (2019). Conserved cell types with divergent features in human versus mouse cortex. *Nature*, *573*(7772), 61–68. https://doi.org/10.1038/s41586-019-1506-7

Hoffmann, A. A., & Harshman, L. G. (1999). Desiccation and starvation resistance in Drosophila: Patterns of variation at the species, population and intrapopulation levels. *Heredity*, *83*(6), 637–643. https://doi.org/10.1038/sj.hdy.6886490

Huisman, E., Uylings, H. B. M., & Hoogland, P. V. (2004). A 100% increase of dopaminergic cells in the olfactory bulb may explain hyposmia in Parkinson's disease. *Movement Disorders*, *19*(6), 687–692. https://doi.org/10.1002/mds.10713

Inagaki, H. K., Kamikouchi, A., & Ito, K. (2010). Methods for quantifying simple gravity sensing in Drosophila melanogaster. *Nature Protocols*, *5*(1), 20–25. https://doi.org/10.1038/nprot.2009.196

Jacquemyn, J., Kuenen, S., Swerts, J., Pavie, B., Vijayan, V., Kilic, A., Chabot, D., Wang, Y.-C., Schoovaerts, N., Corthout, N., & Verstreken, P. (2023). Parkinsonism mutations in DNAJC6 cause lipid defects and neurodegeneration that are rescued by Synj1. *Npj Parkinson's Disease*, *9*(1), Article 1. https://doi.org/10.1038/s41531-023-00459-3

Janssens, J., Aibar, S., Taskiran, I. I., Ismail, J. N., Gomez, A. E., Aughey, G., Spanier, K. I., De Rop, F. V., González-Blas, C. B., Dionne, M., Grimes, K., Quan, X. J., Papasokrati, D., Hulselmans, G., Makhzami, S., De Waegeneer, M., Christiaens, V., Southall, T., & Aerts, S. (2022). Decoding gene regulation in the fly brain. *Nature*, *601*(7894), 630–636. https://doi.org/10.1038/s41586-021-04262-z

Jenett, A., Rubin, G. M., Ngo, T.-T. B., Shepherd, D., Murphy, C., Dionne, H., Pfeiffer, B. D., Cavallaro, A., Hall, D., Jeter, J., Iyer, N., Fetter, D., Hausenfluck, J. H., Peng, H., Trautman, E. T., Svirskas, R. R., Myers, E. W., Iwinski, Z. R., Aso, Y., … Zugates, C. T. (2012). A GAL4-driver line resource for Drosophila neurobiology. *Cell Reports*, *2*(4), 991–1001. PubMed. https://doi.org/10.1016/j.celrep.2012.09.011

Kalia, L. V., & Lang, A. E. (2015). Parkinson's disease. *The Lancet*, *386*(9996), Article 9996. https://doi.org/10.1016/S0140-6736(14)61393-3

Kang, H. M., Subramaniam, M., Targ, S., Nguyen, M., Maliskova, L., McCarthy, E., Wan, E., Wong, S., Byrnes, L., Lanata, C. M., Gate, R. E., Mostafavi, S., Marson, A., Zaitlen, N., Criswell, L. A., & Ye, C. J. (2018). Multiplexed droplet single-cell RNA-sequencing using natural genetic variation. *Nature Biotechnology*, *36*(1), 89–94. PubMed. https://doi.org/10.1038/nbt.4042

Khuong, T. M., Habets, R. L. P., Kuenen, S., Witkowska, A., Kasprowicz, J., Swerts, J., Jahn, R., van den Bogaart, G., & Verstreken, P. (2013). Synaptic PI(3,4,5)P3 is required for Syntaxin1A clustering and neurotransmitter release. *Neuron*, *77*(6), 1097–1108. https://doi.org/10.1016/j.neuron.2013.01.025

Koopmans, F., van Nierop, P., Andres-Alonso, M., Byrnes, A., Cijsouw, T., Coba, M. P., Cornelisse, L. N., Farrell, R. J., Goldschmidt, H. L., Howrigan, D. P., Hussain, N. K., Imig, C., de Jong, A. P. H., Jung, H., Kohansalnodehi, M., Kramarz, B., Lipstein, N., Lovering, R. C., MacGillavry, H., … Verhage, M. (2019). SynGO: An Evidence-Based, Expert-Curated Knowledge Base for the Synapse. *Neuron*, *103*(2), 217-234.e4. https://doi.org/10.1016/j.neuron.2019.05.002

Kremer, M. C., Christiansen, F., Leiss, F., Paehler, M., Knapek, S., Andlauer, T. F. M., Förstner, F., Kloppenburg, P., Sigrist, S. J., & Tavosanis, G. (2010). Structural long-term changes at mushroom body input synapses. *Current Biology: CB*, *20*(21), 1938–1944. https://doi.org/10.1016/j.cub.2010.09.060

Kuhl, D. E., Minoshima, S., Fessler, J. A., Ficaro, E. P., Wieland, D. M., Koeppe, R. A., Frey, K. A., & Foster, N. L. (1996). In vivo mapping of cholinergic terminals in normal aging, Alzheimer's disease, and Parkinson's disease. *Annals of Neurology*, *40*(3), 399–410. https://doi.org/10.1002/ana.410400309

Li, H., Janssens, J., De Waegeneer, M., Kolluru, S. S., Davie, K., Gardeux, V., Saelens, W., David, F. P. A., Brbić, M., Spanier, K., Leskovec, J., McLaughlin, C. N., Xie, Q., Jones, R. C., Brueckner, K., Shim, J., Tattikota, S. G., Schnorrer, F., Rust, K., … Zinzen, R. P. (2022). Fly Cell Atlas: A single-nucleus transcriptomic atlas of the adult fruit fly. *Science (New York, N.Y.)*, *375*(6584), eabk2432. https://doi.org/10.1126/science.abk2432

Love, M. I., Huber, W., & Anders, S. (2014). Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. *Genome Biology*, *15*(12), 550–550. PubMed. https://doi.org/10.1186/s13059-014-0550-8

Masse, N. Y., Turner, G. C., & Jefferis, G. S. X. E. (2009). Olfactory Information Processing in Drosophila. *Current Biology*, *19*(16), R700–R713. https://doi.org/10.1016/j.cub.2009.06.026

Morais, V. A., Verstreken, P., Roethig, A., Smet, J., Snellinx, A., Vanbrabant, M., Haddad, D., Frezza, C., Mandemakers, W., Vogt-Weisenhorn, D., Van Coster, R., Wurst, W., Scorrano, L., & De Strooper, B. (2009). Parkinson's disease mutations in PINK1 result in decreased Complex I activity and deficient synaptic function. *EMBO Molecular Medicine*, *1*(2), Article 2. https://doi.org/10.1002/emmm.200900006

Mortiboys, H., Johansen, K. K., Aasly, J. O., & Bandmann, O. (2010). Mitochondrial impairment in patients with Parkinson disease with the G2019S mutation in LRRK2. *Neurology*, *75*(22), 2017–2020. https://doi.org/10.1212/wnl.0b013e3181ff9685

Mueller, A., Abolmaali, N. D., Hakimi, A. R., Gloeckler, T., Herting, B., Reichmann, H., & Hummel, T. (2005). Olfactory bulb volumes in patients with idiopathic Parkinson's disease □ a pilot study. *Journal of Neural Transmission*, *112*(10), 1363–1370. https://doi.org/10.1007/s00702-005-0280-x

Mundiñano, I.-C., Hernandez, M., DiCaudo, C., Ordoñez, C., Marcilla, I., Tuñon, M.-T., & Luquin, M.-R. (2013). Reduced cholinergic olfactory centrifugal inputs in patients with neurodegenerative disorders and MPTP-treated monkeys. *Acta Neuropathologica*, *126*(3), 411–425. https://doi.org/10.1007/s00401-013-1144-3

Ng, C.-H., Guan, M. S. H., Koh, C., Ouyang, X., Yu, F., Tan, E.-K., O'Neill, S. P., Zhang, X., Chung, J., & Lim, K.-L. (2012). AMP kinase activation mitigates dopaminergic dysfunction and mitochondrial abnormalities in Drosophila models of Parkinson's disease. *The Journal of Neuroscience□: The Official Journal of the Society for Neuroscience*, *32*(41), 14311–14317. PubMed. https://doi.org/10.1523/JNEUROSCI.0499-12.2012

Norrander, J., Kempe, T., & Messing, J. (1983). Construction of improved M13 vectors using oligodeoxynucleotide-directed mutagenesis. *Gene*, *26*(1), 101–106. https://doi.org/10.1016/0378-1119(83)90040-9

Olsen, A. L., Clemens, S. G., & Feany, M. B. (2023). Nicotine-Mediated Rescue of α-Synuclein Toxicity Requires Synaptic Vesicle Glycoprotein 2 in Drosophila. *Movement Disorders: Official Journal of the Movement Disorder Society*, *38*(2), 244–255. https://doi.org/10.1002/mds.29283

Parkinson, J. (2002). An Essay on the Shaking Palsy. *Journal of Neuropsychiatry*, *14*(2), 223–236. https://doi.org/10.1176/appi.neuropsych.14.2.223

Port, F., Chen, H.-M., Lee, T., & Bullock, S. L. (2014). Optimized CRISPR/Cas tools for efficient germline and somatic genome engineering in Drosophila. *Proceedings of the National Academy of Sciences of the United States of America*, *111*(29), Article 29. https://doi.org/10.1073/pnas.1405500111

Riemensperger, T., Issa, A.-R., Pech, U., Coulom, H., Nguyễn, M.-V., Cassar, M., Jacquet, M., Fiala, A., & Birman, S. (2013). A Single Dopamine Pathway Underlies Progressive Locomotor Deficits in a Drosophila Model of Parkinson Disease. *Cell Reports*, *5*(4), 952–960. https://doi.org/10.1016/j.celrep.2013.10.032

Sarkar, S., Davies, J. E., Huang, Z., Tunnacliffe, A., & Rubinsztein, D. C. (2007). Trehalose, a Novel mTOR-independent Autophagy Enhancer, Accelerates the Clearance of Mutant Huntingtin and α-Synuclein. *Journal of Biological Chemistry*, *282*(8), 5641–5652. https://doi.org/10.1074/jbc.m609532200

Satija, R., Farrell, J. A., Gennert, D., Schier, A. F., & Regev, A. (2015). Spatial reconstruction of single-cell gene expression data. *Nature Biotechnology*, *33*(5), 495–502. PubMed. https://doi.org/10.1038/nbt.3192

Schapira, A. H. V., Chaudhuri, K. R., & Jenner, P. (2017). Erratum: Non-motor features of Parkinson disease. *Nature Reviews Neuroscience*, *18*(8), 509–509. https://doi.org/10.1038/nrn.2017.91

Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch, T., Preibisch, S., Rueden, C., Saalfeld, S., Schmid, B., Tinevez, J.-Y., White, D. J., Hartenstein, V., Eliceiri, K., Tomancak, P., & Cardona, A. (2012). Fiji: An open-source platform for biological-image analysis. *Nature Methods*, *9*(7), 676–682. PubMed. https://doi.org/10.1038/nmeth.2019

Shimada, H., Hirano, S., Shinotoh, H., Aotsuka, A., Sato, K., Tanaka, N., Ota, T., Asahina, M., Fukushi, K., Kuwabara, S., Hattori, T., Suhara, T., & Irie, T. (2009). Mapping of brain acetylcholinesterase alterations in Lewy body disease by PET. *Neurology*, *73*(4), 273–278. https://doi.org/10.1212/wnl.0b013e3181ab2b58

Slabbaert, J. R., Kuenen, S., Swerts, J., Maes, I., Uytterhoeven, V., Kasprowicz, J., Fernandes, A. C., Blust, R., & Verstreken, P. (2016). Shawn, the Drosophila Homolog of SLC25A39/40, Is a Mitochondrial Carrier That Promotes Neuronal Survival. *The Journal of Neuroscience: The Official Journal of the Society for Neuroscience*, *36*(6), 1914–1929. https://doi.org/10.1523/JNEUROSCI.3432-15.2016

Slyper, M., Porter, C. B. M., Ashenberg, O., Waldman, J., Drokhlyansky, E., Wakiro, I., Smillie, C., Smith-Rosario, G., Wu, J., Dionne, D., Vigneau, S., Jané-Valbuena, J., Tickle, T. L., Napolitano, S., Su, M.-J., Patel, A. G., Karlstrom, A., Gritsch, S., Nomura, M., … Regev, A. (2020). A single-cell and single-nucleus RNA-Seq toolbox for fresh and frozen human tumors. *Nature Medicine*, *26*(5), 792–802. https://doi.org/10.1038/s41591-020-0844-1

Snead, D., & Eliezer, D. (2019). Intrinsically disordered proteins in synaptic vesicle trafficking and release. *The Journal of Biological Chemistry*, *294*(10), 3325–3342. PubMed. https://doi.org/10.1074/jbc.REV118.006493

Stocker, R. F., Heimbeck, G., Gendre, N., & de Belle, J. S. (1997). Neuroblast ablation in Drosophila P[GAL4] lines reveals origins of olfactory interneurons. *Journal of Neurobiology*, *32*(5), 443–456. https://doi.org/10.1002/(sici)1097-4695(199705)32:5<443::aid-neu1>3.0.co;2-5

Taguchi, T., Ikuno, M., Yamakado, H., & Takahashi, R. (2020). Animal Model for Prodromal Parkinson's Disease. *International Journal of Molecular Sciences*, *21*(6), 1961. PubMed. https://doi.org/10.3390/ijms21061961

Tian, L., Hires, S. A., Mao, T., Huber, D., Chiappe, M. E., Chalasani, S. H., Petreanu, L., Akerboom, J., McKinney, S. A., Schreiter, E. R., Bargmann, C. I., Jayaraman, V., Svoboda, K., & Looger, L. L. (2009). Imaging neural activity in worms, flies and mice with improved GCaMP calcium indicators. *Nature Methods*, *6*(12), 875–881. https://doi.org/10.1038/nmeth.1398

Tully, T., & Quinn, W. G. (1985). Classical conditioning and retention in normal and mutantDrosophila melanogaster. *Journal of Comparative Physiology A*, *157*(2), 263–277. https://doi.org/10.1007/bf01350033

Uytterhoeven, V., Kuenen, S., Kasprowicz, J., Miskiewicz, K., & Verstreken, P. (2011). Loss of Skywalker Reveals Synaptic Endosomes as Sorting Stations for Synaptic Vesicle Proteins. *Cell*, *145*(1), 117–132. https://doi.org/10.1016/j.cell.2011.02.039

Vanhauwaert, R., Kuenen, S., Masius, R., Bademosi, A., Manetsberger, J., Schoovaerts, N., Bounti, L., Gontcharenko, S., Swerts, J., Vilain, S., Picillo, M., Barone, P., Munshi, S. T., de Vrij, F. M., Kushner, S. A., Gounko, N. V., Mandemakers, W., Bonifati, V., Meunier, F. A., … Verstreken, P. (2017). The SAC1 domain in synaptojanin is required for autophagosome maturation at presynaptic terminals. *The EMBO Journal*, *36*(10), 1392–1411. PubMed. https://doi.org/10.15252/embj.201695773

Venderova, K., Kabbach, G., Abdel-Messih, E., Zhang, Y., Parks, R. J., Imai, Y., Gehrke, S., Ngsee, J., Lavoie, M. J., Slack, R. S., Rao, Y., Zhang, Z., Lu, B., Haque, M. E., & Park, D. S. (2009). Leucine-Rich Repeat Kinase 2 interacts with Parkin, DJ-1 and PINK-1 in a Drosophila melanogaster model of Parkinson's disease. *Human Molecular Genetics*, *18*(22), 4390–4404. https://doi.org/10.1093/hmg/ddp394

Vijayan, V., & Verstreken, P. (2017). Autophagy in the presynaptic compartment in health and disease. *Journal of Cell Biology*, *216*(7), Article 7. https://doi.org/10.1083/jcb.201611113

Virtanen, P., Gommers, R., Oliphant, T. E., Haberland, M., Reddy, T., Cournapeau, D., Burovski, E., Peterson, P., Weckesser, W., Bright, J., van der Walt, S. J., Brett, M., Wilson, J., Millman, K. J., Mayorov, N., Nelson, A. R. J., Jones, E., Kern, R., Larson, E., … SciPy 1.0 Contributors. (2020). SciPy 1.0: Fundamental algorithms for scientific computing in Python. *Nature Methods*, *17*(3), 261–272. https://doi.org/10.1038/s41592-019-0686-2

Wang, T., Li, B., Nelson, C. E., & Nabavi, S. (2019). Comparative analysis of differential gene expression analysis tools for single-cell RNA sequencing data. *BMC Bioinformatics*, *20*(1), 40–40. PubMed. https://doi.org/10.1186/s12859-019-2599-6

Webb, J. L., Ravikumar, B., Atkins, J., Skepper, J. N., & Rubinsztein, D. C. (2003). α-Synuclein Is Degraded by Both Autophagy and the Proteasome. *Journal of Biological Chemistry*, *278*(27), 25009–25013. https://doi.org/10.1074/jbc.m300227200

West, A. B., Moore, D. J., Biskup, S., Bugayenko, A., Smith, W. W., Ross, C. A., Dawson, V. L., & Dawson, T. M. (2005). Parkinson's disease-associated mutations in leucine-rich repeat kinase 2 augment kinase activity. *Proceedings of the National Academy of Sciences of the United States of America*, *102*(46), 16842–16847. PubMed. https://doi.org/10.1073/pnas.0507360102

Wilson, C. J., Chang, H. T., & Kitai, S. T. (1990). Firing patterns and synaptic potentials of identified giant aspiny interneurons in the rat neostriatum. *The Journal of Neuroscience : The Official Journal of the Society for Neuroscience*, *10*(2), 508–519. PubMed. https://doi.org/10.1523/JNEUROSCI.10-02-00508.1990

Wilson, R. I. (2013). Early olfactory processing in Drosophila: Mechanisms and principles. *Annual Review of Neuroscience*, *36*, 217–241. PubMed. https://doi.org/10.1146/annurev-neuro-062111-150533

Witt, M., Bormann, K., Gudziol, V., Pehlke, K., Barth, K., Minovi, A., Hähner, A., Reichmann, H., & Hummel, T. (2009). Biopsies of olfactory epithelium in patients with Parkinson's disease. *Movement Disorders*, *24*(6), 906–914. https://doi.org/10.1002/mds.22464

Wolf, F. A., Angerer, P., & Theis, F. J. (2018). SCANPY: Large-scale single-cell gene expression data analysis. *Genome Biology*, *19*(1), 15. https://doi.org/10.1186/s13059-017-1382-0

Wolock, S. L., Lopez, R., & Klein, A. M. (2019). Scrublet: Computational Identification of Cell Doublets in Single-Cell Transcriptomic Data. *Cell Systems*, *8*(4), 281-291.e9. PubMed. https://doi.org/10.1016/j.cels.2018.11.005

Wu, H., Williams, J., & Nathans, J. (2014). Complete morphologies of basal forebrain cholinergic neurons in the mouse. *eLife*, *3*, e02444–e02444. PubMed. https://doi.org/10.7554/eLife.02444

Yamakado, H., & Takahashi, R. (2024). Experimental Animal Models of Prodromal Parkinson's Disease. *Journal of Parkinson's Disease*. https://doi.org/10.3233/JPD-230393

Yang, S., Corbett, S. E., Koga, Y., Wang, Z., Johnson, W. E., Yajima, M., & Campbell, J. D. (2020). Decontamination of ambient RNA in single-cell RNA-seq with DecontX. *Genome Biology*, *21*(1), 57–57. PubMed. https://doi.org/10.1186/s13059-020-1950-6

Yang, Y., Gehrke, S., Imai, Y., Huang, Z., Ouyang, Y., Wang, J.-W., Yang, L., Beal, M. F., Vogel, H., & Lu, B. (2006). Mitochondrial pathology and muscle and dopaminergic neuron degeneration caused by inactivation of Drosophila Pink1 is rescued by Parkin. *Proceedings of the National Academy of Sciences of the United States of America*, *103*(28), 10793–10798. https://doi.org/10.1073/pnas.0602493103

Zhang, J., Schulze, K. L., Hiesinger, P. R., Suyama, K., Wang, S., Fish, M., Acar, M., Hoskins, R. A., Bellen, H. J., & Scott, M. P. (2007). Thirty-one flavors of Drosophila rab proteins. *Genetics*, *176*(2), 1307–1322. https://doi.org/10.1534/genetics.106.066761

Zhang, S., Xiao, Q., & Le, W. (2015). Olfactory dysfunction and neurotransmitter disturbance in olfactory bulb of transgenic mice expressing human A53T mutant α-synuclein. *PloS One*, *10*(3), e0119928–e0119928. PubMed. https://doi.org/10.1371/journal.pone.0119928

Zimprich, A., Biskup, S., Leitner, P., Lichtner, P., Farrer, M., Lincoln, S., Kachergus, J., Hulihan, M., Uitti, R. J., Calne, D. B., Stoessl, A. J., Pfeiffer, R. F., Patenge, N., Carbajal, I. C., Vieregge, P., Asmus, F., Müller-Myhsok, B., Dickson, D. W., Meitinger, T., … Gasser, T. (2004). Mutations in LRRK2 cause autosomal-dominant parkinsonism with pleomorphic pathology. *Neuron*, *44*(4), 601–607. https://doi.org/10.1016/j.neuron.2004.11.005

a

Chiasm glia

Cortex glia

T2a

Tyraminergic

Ensheathing glia

Peptidergic

Serotonergic

Mi15

CB DAN

Central Brain Dopaminergic neurons (incl. PAM)

Lawf1

Lawf2

Tm9

Tm4

Astrocyte-like

LC17p

LC4

LLPC1

LC12p

PB2

LPC1

DCN

Tm20

L5

AstA/NPF

LPLC1

LPLC2

LC6

Tm2

L1

Mi1

L3 L2

TmY14p

Dm11

Clk-1

Clk-2

Dm8

PB1

LC10a

LC10b

L4

Dm9

Pm4

Photoreceptors

TmY4p

Poxn

α/β-KC

Perineurial glia

Plasmatocytes

LPi-34

Dm2

T4/T5

γ-KC

Kenyon cells

PB4/5

Dm3

TmY3

Mi4

Dm12

dFB_A

Dm1

dFB_B

Tm5c

Mi9

Tm3

OPN

Olfactory Projection neurons

T3

Dm10

TmY8p

T2

C3

Tm29

Pm1

AstA/Nplp1

Lai

Pm2

TmY5a

C2

Tm5ab

α'/β'-KC

Kenyon cells

Tm1

T1

● Black: Cells with a significant number of DEGs

b hLRRK2^{G2019S}

c hPINK1^{P399L}

d hRab39B^{G192R}

e Aux^{R927G}

f Synj^{R258Q}

a

transport

metabolism

synapse organisation

synaptic signaling

process in the postsynapse

process in the presynapse

a' humanDEGs

-log10 Q-value
too few genes
ns
2
6
≥7

a" overlap DEGs human fly

-log10 Q-value
too few genes
ns
2
10
≥14

b

30 -log10 p-value
0

	hLRRK2^G2019S	hPINK1^P399L	hRab39B^G192R	Aux^R927G	Synj^R258Q	Human NBM
Synapse organization						
Signal transduction						
Trans-synaptic signaling						
Axon guidance						
SNARE binding						
MAPK cascade						
Hippo signaling						
Wnt signaling pathway						
Autophagy						
Protein folding						

c

Cytoskeleton

TUBA1A	*alpha-tubX*
TUBA2A,2B,4B	*beta-tubX*
DMD	*Dys*
ROCK1,2	*rok*
MAP1A	*futch*
MAPT	*tau*
PAK5	*mbt*

Proteostasis

HSP90AB1	*hsp83*
HSPA5	*hsc70-4*
RB1CC1	*Atg17*

Active zone

TSPOAP1	*rbp*
RIMS3	*rim*
CADPS	*cadps*

Channels

KCNN2,3	*sk*
CACNA1A,B,	*Ca alpha1d*
KCNB1 C,D,E	*shab*
KCNC1	*shaw*
KCNH1	*eag*
SCN1A,2A, 8A	*para*

Synaptic Trafficking

SH3GL2	*EndoA*
CPLX1	*cpx*
SNAP25,23	*SNAP25,23*
STXBP5	*tomo*
SYT1,2	*syt1*
SYT4	*syt4*
SYT7	*syt7*
PPP3CA	*canA14F,*
RCAN1,2	*sra* *pp2B14D*
AP1G2	*AP1-gamma*
EFR3B	*stmA*
HOOK1,2	*hook*
RAB11A	*rab11*

Postsynaptic organisation

IQSEC1,2	*siz*
DLG1	*Dlg*
ACHE	*alpha-Est9*
UNC79	*unc79*
CHRM2,3	*mAchR-A*
GRM1,3,5,8	*mmt*
GRIA3,4	*GluRIA,B*
TTYH3	*tty*

Synaptic Adhesion

NLGN2, 4Y	*Nlg2,3*
ROBO	*robo*
NRXN3	*Nrx1*
EPHA3,4,5,	*Eph*
PTPRD 6,7	*Lar*
DSCAM	*Dscam1,2,4*
NCK2	*dock*

human gene fly homologue

a

b
Calyx
Antennal lobe
OPN cell bodies
OPN>GCaMP

c
Control
hLRRK2^{G2019S}
ΔF (%)
250
0

d
ΔF/F$_0$ (%)
Control
hLRRK2
hLRRK2^{G2019S}
time (s)
0   90

e
Control

f
hLRRK2^{G2019S}

g
ΔF/F$_0$ (%)
400
0
nd

h
Calyx area (%)
200
0
nd

Control
hLRRK2
hLRRK2^{G2019S}
hPINK1
hPINK1^{P399L}
hRab39B
hRab39B^{G192R}
Aux
Aux^{R927G}
Synj
Synj^{R258Q}

OPN>*wt CDS*   − +   − +   − +   − +   − +

a

Control

hLRRK2
hLRRK2^{G2019S}

hPINK1
hPINK1^{P399L}

hRab39B
hRab39B^{G192R}

Aux
Aux^{R927G}

Synj
Synj^{R258Q}

-0.4   0   0.4

b

-0.4   0   0.4

Olfactory performance in PD fly mutants

c

hLRRK2^{G2019S} −/+

LRRK^{KO} −/+

hPINK1^{P399L} −/+

hRab39B^{G192R} −/+

hRab39B$^{G192R}_{T1>hR39B}$ −/+

Aux^{R927G} −/+

Synj^{R258Q} −/+

-0.4   0   0.4
Preference index

d

-0.4   0   0.4
Preference index

OPN-specific rescue of olfactory performance defects