PHR1 and PHL1 mediate rapid high-light responses and acclimation to high photosynthetic activity

Lukas Ackermann[1], Agnes Fekete[2], Laura Schröder[3], Thomas Nägele[3], Alina J. Hieber[1], Monika Müller[1], Martin J. Mueller[2], Maria Klecker[1*]

[1] Plant Physiology, University of Bayreuth, 95447 Bayreuth, Germany

[2]Julius-von-Sachs-Institute of Biosciences, Biocenter, Pharmaceutical Biology, University of Würzburg, 97082 Würzburg, Germany

[3]Faculty of Biology, Plant Evolutionary Cell Biology, LMU München, 82152 Planegg, Germany

[*]Address correspondence to: Maria Klecker, Department of Plant Physiology, University of Bayreuth, Universitätsstrasse 30, D-95447 Bayreuth, Germany. e-mail: maria.klecker@uni-bayreuth.de

ORCID IDs

Maria Klecker	0000-0001-5508-2363
Lukas Ackermann	0009-0009-8604-6029
Agnes Fekete	0000-0003-3058-7570
Martin J. Mueller	0000-0001-8122-5460
Thomas Nägele	0000-0002-5896-238X
Alina J. Hieber	0009-0009-0407-8968

ABSTRACT

•   Rapid acclimation towards high light is vital for the prevention of phototoxic stress to

plant tissues. Photosynthesis transiently consumes inorganic phosphate (P_i). We therefore asked

whether changes in intracellular P_i pools can act as a signal for reprogramming gene expression

upon increased light intensity.

•   The function of triose phosphate utilization for intracellular P_i homeostasis was

deciphered by investigation of mutants defective in both photoassimilate partitioning and low-

P_i signaling. Next, we determined subcellular P_i levels and transcript accumulation upon short-

term high light. Physiological consequences were examined by analyses of ATP, carbohydrates,

and an untargeted lipid profiling approach.

•   The capacity for triose phosphate utilization clearly affected systemic P_i signaling.

High-light treatment caused a rapid depletion specifically of chloroplast P_i levels paralleled by

induction of transcripts dependent on PHOSPHATE STARVATION RESPONSE 1 (PHR1)

and PHR1-LIKE 1. Among the high-light induced targets of PHR1, *SRG3/GDPD1* is involved

in phospholipid catabolism. Lipid profiling revealed differences between WT and *srg3* mutant

plants upon high light, including changes in linolenic acid and photoprotective zeaxanthin.

•   We conclude that photosynthetic activity regulates the low-P_i response machinery in the

nucleus to implement high-light acclimation. This facilitates the liberation of cellular P as well

as the maintenance of membrane lipid integrity.

Key words

46   PHR1, phosphate starvation, light acclimation, photosynthesis, glycerophosphodiester

47 phosphodiesterase, membrane remodeling, non-photochemical quenching (NPQ)

Introduction

Light energy is provided to photoautotrophic organisms at variable intensities that can

fall below or exceed the requirements of the photosynthetic machinery at unpredictable times

during the day period. This necessitates an enzymatic machinery capable of coping with rapid

changes in metabolic flux, as well as the containment of overexcitation-induced hazards, such

as lipid peroxidation (Havaux & Niyogi, 1999). Both challenges are met not only by post-

transcriptional regulation of enzyme activity (reviewed in König et al. (2012), Matiolli et al.

(2022)), but also by pronounced changes in mRNA abundance that occur within 30 minutes

upon an increase in irradiance (Vogel *et al.*, 2014; Suzuki *et al.*, 2015; Huang *et al.*, 2019). The

latter mechanism depends on fast signal transmission between the chloroplast and the nuclear

transcription machinery. Such communication is mediated by 'primary' retrograde signals

(Dietz, 2015), including redox cues and reactive oxygen species which are produced during

photosynthetic reactions (Bechtold *et al.*, 2008; Dietz *et al.*, 2016). Additionally, increased

electron transport activity and light-induced Calvin-Benson cycle activation promote the

accumulation of primary products of carbon fixation (Dietz & Heber, 1984, 1986; Borghi *et al.*,

2019) which may themselves act as signaling molecules (Häusler *et al.*, 2014; Moore *et al.*,

2014). Hence, induction of high-light specific gene expression partially depends on the triose

phosphate/phosphate translocator (TPT) that mediates the exchange of triose phosphates and 3-

phosphoglycerate (3-PGA) with inorganic phosphate (P_i) at the chloroplast envelope (Schneider

et al., 2002; Vogel *et al.*, 2014; Weise *et al.*, 2019). A central function of the TPT in light

acclimation and retrograde signaling was corroborated by the severe high-light dependent

phenotypes of double mutants defective in *TPT* and *ADG1* (for *ADP GLUCOSE*

PYROPHOSPHORYLASE 1), encoding a small subunit of the ADP-glucose pyrophosphorylase

complex (AGPase) which is required for the biosynthesis of transitory starch (Schmitz *et al.*,

2012). Phenotypes of *adg1-1 tpt* double mutants include growth retardation, low diurnal

changes in carbohydrate levels, and high chlorophyll fluorescence. These observations point

towards an important role of photosynthate partitioning for retrograde signaling, however the

identity of the primary sensors as well as signal transduction mechanisms are still unclear.

Phosphorus makes up about 2-30 permille of the plant dry mass (Kumar *et al.*, 2019),

as it is incorporated into a large subset of biomolecules. Systemic signaling of P_i availability

and P_i distribution within the plant body mainly depend on a conserved gene family encoding

PHOSPHATE STARVATION RESPONSE 1 (PHR1) and at least 14 PHR1 LIKE (PHL)

transcription factors of the GARP coiled-coiled type (Rubio *et al.*, 2001; Zhou *et al.*, 2008;

Bustos *et al.*, 2010; Thibaud *et al.*, 2010; Sun *et al.*, 2016; Safi *et al.*, 2017). Simultaneous loss

of *PHR1* and *PHL1* function affects the expression of roughly 70 % of the P_i starvation

responsive genes in the shoots of Arabidopsis (*Arabidopsis thaliana)* (Bustos *et al.*, 2010). A

decrease in cellular P_i concentration impairs production of the inositol pyrophosphate species

InsP8 in the cytosol, relieving PHR1/PHLs of suppression by SPX (for SYG1/Pho81/XPR1)

domain-containing proteins, thus linking transcription factor activity to P_i availability (Puga *et*

al., 2014; Dong *et al.*, 2019; Zhu *et al.*, 2019; Ried *et al.*, 2021). The transcriptional response

to P_i scarcity leads to scavenging of external and internal P_i sources through phosphatase

induction (Morcuende *et al.*, 2007) and altered lipid metabolism (Misson *et al.*, 2005; Pant *et*

al., 2015; Yang *et al.*, 2025), anthocyanin biosynthesis (Nilsson *et al.*, 2012; Liu *et al.*, 2022;

Li *et al.*, 2023), and protection from photodamage (Nilsson *et al.*, 2012), to name a selection.

The largest pool of cellular P reserves resides in the vacuole, and it has been shown in rice

(*Oryza sativa*), that vacuolar P_i efflux into the cytosol is mediated by two proteins of the

glycerol 3-phosphate transporter family which are transcriptionally induced by PHR proteins

(Xu *et al.*, 2019). Three homologs have been identified in Arabidopsis and are termed VPE1-3

(Ramaiah *et al.*, 2011; Xu *et al.*, 2019), but their contribution to P_i homeostasis has not been

addressed so far in dicotyledonous plants. PHR1/PHLs are characterized as transcriptional

activators (Nilsson *et al.*, 2007; Bustos *et al.*, 2010). Accordingly, direct targets of PHR1

(Bustos *et al.*, 2010; Castrillo *et al.*, 2017) are strongly upregulated upon P_i starvation and

include genes such as *PHOSPHATE STARVATION-INDUCED GENE 2* (*PS2*; AT1G73010)

(Hanchi *et al.*, 2018), *MONOGALACTOSYLDIACYLGLYCEROL SYNTHASE 3* (*MGD3*;

AT2G11810) (Kobayashi *et al.*, 2004), and *GLYCEROPHOSPHODIESTER*

PHOSPHODIESTERASE 1/SENESCENCE-RELATED GENE 3 (*GDPD1/SRG3*; AT3G02040)

(Cheng *et al.*, 2011). The latter encodes a plastid-localized enzyme with broad-spectrum

glycerophosphodiester phosphodiesterase activity releasing primary alcohols and glycerol

3-phosphate from deacylated phospholipids (Cheng *et al.*, 2011). SRG3 is supposed to mainly

act in the catalysis of phosphatidylcholine to provide glycerol-3-phosphate for DGDG

biogenesis, for glycolytic utilization, or liberation of P_i by subsequent acid phosphatase activity

(Cheng *et al.*, 2011). A function in phospholipid homeostasis was recently corroborated by

Wojciechowska et al. (2024) who proposed that the transcriptional repressor PROTODERMAL

FACTOR2 directly links phospholipid availability to the repression of both *SRG3* and *SPX1*.

The *SRG3* gene was shown to be important for seedling fitness and maintenance of cellular P_i

levels under P_i starvation conditions (Cheng *et al.*, 2011), but how exactly SRG3 contributes to

nutrient homeostasis is not understood.

P_i is unique among the macronutrients in that it is the only root-supplied element which

is directly required for photosynthesis (Heldt & Rapley, 1970; Dietz & Heber, 1984). It has

been shown experimentally that under high-CO_2 and high-light conditions, photosynthesis can

indeed become limited by P_i supply to the chloroplast, both under P_i depleted and nutrient-rich

conditions (Walker & Osmond, 1986; Dietz & Foyer, 1986; Sivak & Walker, 1986). However,

whether rapid increase in photosynthetic activity directly affects P_i homeostasis of the cell, is

still unclear. In this study, we show that increments in light intensity rapidly deplete chloroplast

P$_i$ reserves *in planta*. Strikingly, although other cellular P$_i$ pools are not affected by short-term

high light, this is accompanied by a nuclear transcriptional response characteristic of P$_i$

depletion which depends on *PHR1* and *PHL1*. Transcript analysis of the *adg1 tpt-2* mutant links

PHR(-like) activation to triose phosphate allocation. Processes regulated by PHR1/PHL1 upon

high light contribute to photosynthetic acclimation since *phr1 phl1* mutants are sensitive to

high-light growth conditions, and showed altered carbohydrate and anthocyanin accumulation

compared to the wildtype (WT) when the light intensity was increased. Moreover, analyses of

the starch biosynthesis-deficient *adg1 phr1 phl1* triple mutant revealed an important function

of P$_i$ signaling when triose phosphate utilization is constrained. Induction of genes involved in

lipid metabolism such as *SRG3* by PHR1/PHL1 furthermore contributes to fatty acid

homeostasis and membrane photoprotection, as revealed by lipid profiling of the *srg3* mutant

upon high light. Together, we propose that PHR1/PHL1 mediate cellular P$_i$ release and rapid

photosynthetic acclimation upon increases in light intensity, probably activated by direct

sensing of chloroplast P$_i$ levels upon triose phosphate overproduction.

Results

Triose phosphate utilization affects P_i signaling

Photoassimilate allocation and utilization in the chloroplast and cytosol are supposed to limit the rate of P_i recycling from photosynthetic products (McClain & Sharkey, 2019). In line with this assumption, P_i deprivation has been shown to stimulate starch accumulation in a *PHR1*-dependent manner, which involves AGPase activity (Nilsson *et al.*, 2007). To address whether partitioning of phosphorylated assimilates contributes to P_i homeostasis of the cell, we examined the phenotypes of mutants defective in starch biosynthesis (*adg1-1*) and chloroplastic triose phosphate export (*tpt-2*) during P_i starvation in the presence of 14.61 mM (0.5 %) sucrose. Starch accumulation in response to P_i depletion was abolished in both the *phr1-3 phl1* double mutant and the *adg1-1* mutant (Figure 1A), confirming that PHR(-like) signaling as well as the small AGPase subunit ADG1 are required for increased starch production under P_i starvation. Concomitantly, P_i-depleted *adg1-1* mutants accumulated high levels of monosaccharides (glucose and fructose, Figure 1A). Loss of TPT function caused significantly higher levels of starch and glucose under P_i deprivation as compared to the WT, which resumed a trend that was also observed under control conditions and likely reflects increased day-time starch turnover in the *tpt-2* mutant (Schneider *et al.*, 2002). Interestingly, over-accumulation of carbohydrates by *adg1-1* and *tpt-2* mutants under P_i deprivation was ceased in the double mutant *adg1 tpt-2*, indicating that retrograde signaling of carbohydrate status is involved in this process. Likewise, *adg1 tpt-2* double mutants failed to accumulate anthocyanins upon P_i deprivation (Supporting Information Fig. S1A, B).

In order to test if nuclear gene regulation by PHR1/PHL1 is important for the adjustment of P homeostasis when P_i recycling from photosynthetic intermediates is impaired, we generated the triple mutant *adg1-1 phr1-3 phl1* (hereafter referred to as *adg1 phr1 phl1*). This mutant exhibited a pronounced growth defect which was the result of reduced leaf size and number (Figure 1B, C). Next, *adg1 phr1 phl1* plants flowered later than the parental lines (Supporting Information Fig. S1C). These findings indicate that starch production functions synergistically with PHR1/PHL1 in maintaining cellular P_i and/or carbohydrate homeostasis.

We noted that growth retardation of *adg1 phr1 phl1* resembled the phenotype of the *adg1 tpt-2* mutant (Figure 1B). To test the hypothesis that *adg1 tpt-2* might likewise be affected in P_i homeostasis, we measured P_i contents of aboveground tissues and determined P_i starvation gene expression in vegetative plants grown on soil. Surprisingly, we found that *adg1 tpt-2* double mutants over-accumulated P_i compared to the *adg1-1* single mutant (Figure 1D).

Counter-intuitively, quantitative RT-PCR (qRT-PCR) analysis revealed that transcripts of P_i

starvation marker genes including *SRG3*, *PS2*, *VPE1* (AT3G47420), *SPX1*, and *MGD3*

(Supporting Information Fig. S2) were all constitutively up-regulated in *adg1 tpt-2* (Figure 1E).

Additionally, transcript abundance of the circadian regulator *BASIC LEUCINE ZIPPER 63*

(*bZIP63*) (Baena-González *et al.*, 2007; Frank *et al.*, 2018), which is repressed by P_i depletion

in the WT (Supporting Information Fig. S2), was also reduced in rosettes of *adg1 tpt-2* as well

as *adg1-1* when grown on soil (Figure 1E). Interestingly, *tpt-2* on the contrary showed slightly

higher expression levels of *bZIP63* than WT plants. Thus, P_i signaling pathways are

substantially affected when triose phosphate utilization and allocation in the chloroplast are

compromised.

It was reported that the *adg1 tpt-2* mutant is sensitive towards higher light intensities,

which could be overcome by exogenously supplied sucrose (Schmitz *et al.*, 2012; Heinrichs *et*

al., 2012). Given the phenotypic similarities between *adg1 tpt-2* and *adg1 phr1 phl1,* we asked

if the growth defect of *adg1 phr1 phl1* was also dependent on light intensity. In fact, we

observed high-light dependent growth restriction of *adg1 phr1 phl1* seedlings that was

comparable to *adg1 tpt-2*, both under low and high P_i conditions (Figure 1F; Supporting

Information Fig. S3). As for *adg1 tpt-2*, the difference in growth performance between low-

and high-light conditions was alleviated by providing sucrose to *adg1 phr1 phl1* mutants

(Figure 1F), a phenomenon that was also observed for root growth of the mutant seedlings

(Supporting Information Fig. S3). Remarkably, high-light dependent growth restriction was

also seen for both the *adg1-1* and *phr1-3 phl1* mutants, although to a lower extend than in the

cases of *adg1 tpt-2* and *adg1 phr1 phl1* (Figure 1F). Again, the difference in growth

performance between low and high light was attenuated by exogenous sucrose, indicating that

carbohydrate homeostasis is disturbed in *adg1-1*, *phr1-3 phl1*, and the triple mutant when

exposed to high light.

We concluded from these experiments that triose phosphate allocation has pronounced

effects on the nuclear P_i starvation response. Vice versa, P_i signaling is required for acclimation

to triose phosphate oversupply in high light or when starch production is genetically disabled.

High light elicits P_i starvation gene expression in the WT

The results of our reverse genetic analyses indicated that P_i homeostasis is affected by

photoassimilate availability and utilization. This raised the question whether P_i signaling was

also modulated in WT plants under non-starved conditions by naturally occurring changes in

the photosynthetic rate such as upon a shift in light intensity. In order to address this, we

interrogated published transcriptomic data collected from seedling shoots under P_i starvation

(Bustos *et al.*, 2010) and high-light treatments under nutrient-rich conditions (Huang *et al.*,

2019). We found that out of 379 transcripts upregulated ($\log_2$ fold change >2) by 30 min of high

light, 75 were also upregulated by P_i starvation (Figure 2A). For longer high-light treatments of

24 hours, 131 out of 564 transcripts were also P_i-responsive (Figure 2A). In both cases, the

overlap was found to be significant at $P < 0.0001$ in hypergeometric testing using Fisher's exact

test. In order to estimate whether the observed overlap between high-light and low-P_i induced

transcripts might be due to generally higher nutrient requirements under high-light conditions,

we next included a set of nitrogen-responsive transcripts in the comparison (Krapp *et al.*, 2011).

We detected a significant overlap of 33 transcripts that are differentially expressed in the shoots

under low nitrogen conditions with transcripts upregulated by a high-light treatment of 24 hours

duration (Figure 2A). In contrast, no significant enrichment of nitrogen-responsive transcripts

among high-light induced transcripts was observed for 30-min high-light treated plants (10 out

of 379). Thus, short-term high-light treatment specifically triggered low-P_i responses, while

long-term high-light stress seems to elevate general nutrient demand of the plant.

We next asked whether the induction of P_i-starvation responsive genes by short-term

high light required the activity of PHR1(-like) transcription factors and thus was truly caused

by low-P_i signaling pathways. To this end, we grew WT and *phr1-1 phl1* mutant plants under

nutrient-rich conditions on soil in an 8/16 hours (light/dark) photoregime. After 5 weeks, the

photon flux density was increased to $450\pm30\ \mu mol\ m^{-2}\ s^{-1}$ (high light) at 1 hour after

commencement of the photoperiod, while the control group was kept under growth light

conditions of $70\pm5\ \mu mol\ m^{-2}\ s^{-1}$. We conducted qRT-PCR analysis of samples harvested after

20, 45, and 125 min of treatment and detected pronounced upregulation of *SRG3*, *SPX1*, and

VPE1 transcripts in WT plants already after 20 min of exposure to high light compared to

control plants, as well as a slight increase in *MGD3* and *PS2* transcript abundance (Figure 2B).

In all five cases, the upregulation of gene expression was attenuated within 125 min in high

light, confirming that response to P_i shortage constitutes an early consequence of increased

irradiance. Crucially, P_i-starvation marker transcripts were not responsive to light conditions in

phr1-1 phl1 mutant plants (Figure 2B), suggesting that PHR1/PHL1 are required for high-light

mediated regulation of these genes. Instead, we observed a tendency towards constitutively

higher transcript levels of most tested genes in the *phr1-1 phl1* mutants under control

conditions, particularly within the later timepoints of the analysis. This is likely caused by

activation of the remaining PHL genes in this mutant exhibiting chronically low P_i levels.

PHR1 expression itself was reported to depend on light (Liu *et al.*, 2017). Importantly
however, increasing the light intensity to 450±30 µmol m^{-2} s^{-1} did not affect *PHR1* transcript
levels under our experimental conditions (Supporting Information Fig. S4), consistent with the
accumulation of *PHR1* transcript being only responsive to changes in the very low fluence range
(Liu *et al.*, 2017).

GLUCOSE-6-PHOSPHATE/PHOSPHATE TRANSLOCATOR 2 (*GPT2*) is a light
responsive gene (Athanasiou *et al.*, 2010) which was also described to be induced by P$_i$
starvation (Morcuende *et al.*, 2007). Furthermore, high-light induced expression of *GPT2* was
previously described to depend on *TPT* function (Kunz *et al.*, 2010; Weise *et al.*, 2019), and we
noted that the promotor sequence of *GPT2* contains a PHR1 binding motif (GTATATTC) close
to the transcriptional start site. However, in our high-light experiment, *GPT2* was upregulated
in both WT and *phr1-1 phl1* after 45 minutes, and transcript levels continued to rise upon stress
during the course of the analysis (Figure 2B). This showed that PHR1/PHL1 are not required
for high-light mediated induction of *GPT2* expression. Moreover, we were not able to detect an
effect of the *gpt2-1* allele (Niewiadomski *et al.*, 2005) on carbohydrate accumulation under P$_i$
depletion (Supporting Information Fig. S5), arguing against a major role of this gene for sugar
homeostasis under P$_i$ starvation. Thus, *GPT2* induction under high light is independent of
PHR1(-like) signaling, and *phr1 phl1* mutants are not generally impaired in the transcriptional
response to high light. Nevertheless, high-light treatment at early timepoints specifically
triggers low-P$_i$ signaling resulting in the upregulation of transcripts such as *SRG3*, *SPX1*, and
VPE1.

Short-term high light affects chloroplast P$_i$ levels and sucrose accumulation

Upregulation of P$_i$ starvation marker genes upon short-term high light suggested the
existence of a light-induced signal resulting in the activation of PHR1(-like) transcription
factors. Since PHR1(-like) activity is primarily regulated by the P$_i$ status of the cell, we asked
whether cellular P$_i$ reserves were affected by high light, thus relieving PHR(-like) transcription
factors of SPX-mediated inhibition. To address this, we performed non-aqueous fractionation
of plant rosettes subjected to high light for 20 min or kept under control conditions, and
determined subcellular concentrations of free P$_i$. Under control conditions, WT and *phr1-1 phl1*
mutant plants clearly differed in vacuolar P$_i$ reserves (14 % of WT level in *phr1-1 phl1*),
chloroplast P$_i$ content (50 %), as well as the cytosolic and nuclear P$_i$ fraction (63 %) (Figure
3A). Strikingly, high-light treatment caused a decline in chloroplast P$_i$ levels by 54 % in the

WT (Figure 3A), decreasing to 0.4 µmol P_i g^{-1} dry weight which corresponds to a concentration

of about 0.12 mM P_i (Santarius & Heber, 1965). Consequently, P_i levels of high-light treated

chloroplasts in the WT approximated the P_i levels measured in chloroplasts of *phr1-1 phl1*

under control conditions. In contrast, no light-dependent effect was seen for P_i levels in the

double mutant. Importantly, P_i concentrations of the cytosolic/nuclear and vacuolar fractions

were not significantly altered by high light in either WT or *phr1-1 phl1* mutant plants (Figure

3A). Hence, elevated light rapidly induced P_i depletion in the WT, however the effect remained

restricted to the chloroplast, and therefore does not qualify as a direct cause of PHR1/PHL1

activation upon high light.

PHR1(-like) regulation by inositol pyrophosphates relates to the metabolism of these signaling

compounds in the cytosol (Wild *et al.*, 2016) which depends on both P_i and ATP levels (Zhu *et*

al., 2019; Riemer *et al.*, 2021). Given that cytosolic P_i concentrations were unchanged by 20

min high-light treatments, we asked if light increase affected cellular ATP levels. While

phr1-1 phl1 double mutants exhibited a tendency towards lover ATP levels that was more

pronounced upon high-light treatment, light increase did not significantly affect ATP or ADP

levels in the WT at 20 or 90 min of treatment (Figure 3B). Hence, activation of PHR1(-like)

transcription by short-term high light appears to be independent of cytosolic P_i and ATP levels.

It has been reported that P_i starvation responses are modulated by sugar availability (Franco-

Zorrilla *et al.*, 2005; Müller *et al.*, 2005, 2007; Karthikeyan *et al.*, 2007). Since sugars are also

involved in the signaling of photosynthetic status (Häusler *et al.*, 2014; Li & Sheen, 2016;

Zirngibl *et al.*, 2023), we tested if short-term high-light treatment was sufficient to induce

changes in sugar levels in leaves. At the timepoint tested, i.e. 80 min after onset of the

photoperiod, contents of sucrose under control conditions were found to be lower in leaves of

phr1-1 phl1 as compared to the WT (Figure 3C). 20 min high-light treatment resulted in a

further increase in sucrose (Figure 3C), but not glucose or fructose (Supporting Information

Fig. S6) levels in the WT. A similar fold-change of sucrose contents was observed in

phr1-1 phl1 mutant leaves upon high light, although the increase in absolute levels was not

significant (Figure 3C). Thus, sucrose levels respond to high-light treatment on a very short

timescale, suggesting that they might act as early signaling compounds, and potentially might

also affect light-dependent P_i starvation responses.

Analysis of transcriptomic data revealed upregulation of both P_i and nitrogen starvation

responses upon medium-term (24 h) high light (Figure 2A). To address if this was attributable

to enhanced nutrient requirements caused by high light, we determined total P_i contents of

rosettes subjected to 24 and 48 h of light increase. In fact, shoot P_i levels of WT plants were

slightly reduced within 48 h of high light as compared to control light conditions (Figure 3D).

This correlated with significantly enhanced plant biomass measured upon 48 h of high light

compared to control treatments (Figure 3D), suggesting that light increase stimulated growth

which was not immediately satisfied by additional nutrient acquisition. In contrast to WT plants,

P_i contents in leaves of *phr1-1 phl1* mutant plants remained at low levels independently of the

light treatment (Figure 3D).

Together, our results indicate that induction of P_i starvation responses upon light increase

occurs in at least two phases: Short-term (20-45 min) responses are triggered by a signal

independent of cytosolic/nuclear P_i concentrations. This short-term signal communicates

chloroplast P_i levels to the nucleus, and/or potentially involves cellular sucrose levels. In the

medium term however, a shift to higher light intensities causes cellular P_i depletion through an

imbalance of growth and nutrient uptake.

Carbohydrate accumulation is affected in *phr1 phl1* mutants

We asked if the induction of P_i starvation gene expression was relevant for the

acclimation of photosynthetic processes to increased radiation. To this end, we first performed

pulse-amplitude modulated (PAM) chlorophyll fluorometry on rosette leaves subjected to

increased light. The efficiency of photosystem II photochemistry (ΦPSII) dropped to the same

extent in both WT and *phr1-1 phl1* mutants upon exposure to increased illumination for

125 min (Figure 4A). After 125 min of treatment, also the maximum efficiency of PSII (F_v/F_m)

had decreased significantly in the stressed plants of both genotypes (Figure 4B). As for ΦPSII,

no difference was observed for F_v/F_m between WT and *phr1-1 phl1*, indicating that thylakoid

reactions were not affected in *phr1-1 phl1* mutants.

Next, we determined the changes in carbohydrate contents that occurred within 90 min

upon an increase in photon flux density from 70 ± 5 μmol m^{-2} s^{-1} to 450 ± 30 μmol m^{-2} s^{-1} in

rosette leaves of 5-weeks-old WT and *phr1-1 phl1* mutant plants. As shown in Figure 4C, starch

contents under control conditions were higher in *phr1-1 phl1* mutants than in the WT. This may

be related to the fact that *phr1-1 phl1* mutants manifest lower P_i levels in the chloroplasts

(Figure 3A), a condition that allosterically activates AGPase (Figueroa *et al.*, 2022). High-light

treatment caused an increase in starch contents in both WT and mutant (Figure 4C). However,

the fold change of starch content upon high light was lower in *phr1-1 phl1* than in WT plants.

This was accompanied by slightly lower levels of monosaccharides (glucose and fructose), but

not sucrose, in the double mutant upon 90 min of light increase (Figure 4E). Notably, differences in sucrose levels that were seen between WT and *phr1-1 phl1* under control growth at 80 min after the onset of the photoperiod (Figure 3C), were not detected at 150 min into the photoperiod (Figure 4E), indicating that sucrose levels are disturbed in *phr1-1 phl1* during the early morning.

In conclusion, carbon fixation and/or allocation is slightly compromised in *phr1-1 phl1* mutants upon high light, while WT-like sucrose levels are maintained during the later course of the day, independent of light conditions.

PHR1/PHL1 are involved in high-light-mediated anthocyanin accumulation

Since both P_i starvation and high-light stress trigger anthocyanin production, we next analyzed the role of *PHR1/PHL1* signaling in anthocyanin accumulation during high-light acclimation. As shown in Figure 4D, anthocyanin contents of *phr1 phl1* leaves after 3 days of high-light exposure were significantly lower than in WT leaves or in the single mutants *phr1-3* and *phl1*, while loss of *SPX1/2* function did not have any statistically significant effect. PHR1 has been implicated in flavonoid biosynthetic gene induction under low P_i (He *et al.*, 2021; Liu *et al.*, 2022). We did not detect significant differences in the induction of *LDOX*, *DFR*, *PAP1*, or *MYB111* transcripts between WT and *phr1-1 phl1* mutants at 24 or 48 h of light increase (Supporting Information Fig. S7). Interestingly however, *PAP2* expression was markedly upregulated in *phr1-1 phl1* compared to the WT at 24 h of high light (Supporting Information Fig. S7), indicating that the regulation of anthocyanin accumulation is disturbed at a different level than biosynthetic gene expression in *phr1-1 phl1*.

It has been reported that high-light induced anthocyanin biosynthesis depends on sucrose signaling (Zirngibl *et al.*, 2023). Given that sucrose levels were similar between WT and *phr1-1 phl1* at 90 min of high light (Figure 4E), we reasoned that sugar-mediated anthocyanin production might be impaired at the signaling level in *phr1-1 phl1*. To address this, we exposed 7-days old seedlings grown on nutrient-rich media to sucrose- or sorbitol-containing media with different P_i supplies for 67 hours. As depicted in Figure 4F, 3 days of mere P_i depletion were not sufficient to trigger measurable anthocyanin production in either genotype. When sorbitol was added to the media, WT and *phr1-1 phl1* seedlings accumulated anthocyanins to the same level (Figure 4F), indicating that WT and *phr1-1 phl1* are indistinguishable in terms of anthocyanin production upon osmotic stress. Crucially, when sorbitol containing media were compared to media containing sucrose at same-osmotic effect

concentrations, *phr1-1 phl1* mutants failed to produce anthocyanins above the level of the

osmotic effect when the growth medium lacked P_i (Figure 4F). Thus, *phr1-1 phl1* mutants are

specifically defective in sucrose-mediated anthocyanin production already at early stages of P_i

depletion.

Taken together, *PHR1/PHL1* functions are required for adequate high-light mediated

anthocyanin accumulation, which might be related to a role in sucrose signaling upon cellular

P_i restriction.

SRG3 expression is required for lipid homeostasis upon light increase

Among the *PHR1/PHL1*-dependent transcripts analyzed under short-term high light,

SRG3 showed particularly strong upregulation (Figure 2B), and it has been suggested that the

SRG3 promoter is directly activated by PHR1 (Cheng *et al.*, 2011). To test this hypothesis, we

performed promoter-transactivation assays in mesophyll protoplasts. PHR1 was able to induce

the expression of a luciferase reporter fused to the promoters of *SRG3*, *MGD3*, *GPT2*, and *SPX1*

(lacking the naturally occurring *Nco*I restriction site, *proSPX1^{GC}*) (Figure 5A). Notably

however, the responsiveness of *proGPT2* to *PHR1* expression was considerably lower than

observed with the constructs harboring *proSPX1^{GC}, proSRG3* or *proMGD3* upstream of the

reporter CDS. This finding was consistent with the observation that *GPT2* was induced

independently of *PHR1/PHL1* under high light (Figure 2B), and suggests that *SRG3*, like

MGD3 and *SPX1*, is a veritable target gene of PHR1.

We next asked whether PHR1-mediated induction of *SRG3* was relevant for lipid

metabolism upon light increase, given the proposed role of SRG3 in phospholipid degradation

pathways. To answer the question, we compared the lipid profile of a four-week-old WT and

srg3 loss-of-function mutant (Cheng et al., 2011) grown in soil, which had been exposed to

high light for two or seven hours. The two-hour exposure was an early timepoint of acclimation,

close to the onset of gene expression, while the seven-hour exposure corresponded to the end

of the light period to allow for a stronger metabolic response. Total lipid extracts of shock-

frozen leaf material were analyzed by ultra-performance liquid chromatography coupled to a

quadrupole time-of-flight mass spectrometer equipped with electrospray ionization source. The

first principal component of the PCA analysis (Figure 5B) on the aligned 2478 lipid features

(Supporting Information Table S1) accounted for 50 % of the total variance, suggesting

differences in the lipidome of *srg3* compared to the WT. Therefore, we profiled the main

structural lipids monogalactosyl diacylglycerols (MGDG), digalactosyl diacylglycerols

(DGDG), sulfoquinovosyl diacylglycerols (SQDG), phosphatidylcholines (PC),
phosphatidylglycerols (PG) and phosphatidylinositols (PI) in WT and *srg3* after 2 h and 7 h of
light stress. Their total levels were comparable (Supporting Information Fig. S8A), which is
consistent with the results obtained for *srg3* mutants under P_i starvation conditions (Cheng *et*
al., 2011). Notably, the levels of MGDG(34:1), DGDG(34:1) and DGDG(34:2) were slightly
higher in both WT and *srg3* after 7 h of high light (Supporting Information Fig. S9), whereas
the levels of the phospholipids determined were comparable (Supporting Information Fig. S10).
In addition, profiling of sphingolipids revealed non-significant higher levels of ceramides in
srg3 compared to WT in all four investigated conditions, but hydroxy ceramides and glucosyl
ceramides were WT-like in the mutant (Supporting Information Fig. S8B).

We also performed lipid profiling of acyl-MGDGs and Arabidopsides in the extracts,
as galactolipids can be acylated or oxygenated (Ibrahim *et al.*, 2011). We could only identify
acyl-MGDG-54:8 and Arabidopside A (1-OPDA, 2-dnOPDA MGDG), and their levels were
significantly higher in *srg3* after 7 h of high light stress compared to the other groups (Figure
5C). This result encouraged us to perform ANOVA on the annotated 2478 lipid features
(Supporting Information Table S2) in order to identify more *srg3*-dependent high-light
responsive lipids. We first selected lipid features that were significantly different between WT
plants treated with 7 h of high light and untreated plants. ANOVA revealed 81 lipid features
(fold change $\geq$ 2, $P \leq 0.05$) whose intensities were different after exposure to high light for 7 h
(Supporting Information Table S2). Of the 81 lipid features that respond to high light in the
WT, 18 were significantly different in *srg3* compared to WT after 7 h of high light (Supporting
Information Table S3). One of them was identified as linolenic acid, whose levels were higher
in *srg3* compared to WT after 7 h but not 2 h of high light (Figure 5C), like Arabidopside A
and MGDG-54:8. Interestingly, a lipid feature was constitutively higher in *srg3* and
accumulated upon high-light treatment (Figure 5C), which we could tentatively assign to a
glycosylated SQDG species (SQDG-34:1+$C_5H_8O_4$; Supporting Information Table S3). The
identification of this unknown lipid species will need to be verified in the future using tandem
mass spectrometry and nuclear magnetic resonance spectrophotometry after purification.

One of the *srg3*-dependent metabolite features showing highest light-responsiveness
was identified as zeaxanthin (Figure 5C; Supporting Information Table S3), a photoprotective
carotenoid and component of the xanthophyll cycle (Demmig-Adams & Adams, 1996; Havaux
& Niyogi, 1999). In *srg3*, zeaxanthin accumulation following 7 h of high light was reduced by
one third as compared to the WT, indicative of a reduced capacity for light energy dissipation
in chloroplast membranes of *srg3*. To further investigate this, we performed PAM chlorophyll

fluorescence measurements and determined the light response of the non-photochemical
quenching (NPQ) parameter in WT, *srg3*, and *phr1-1 phl1* mutant plants. Indeed, the extend of
NPQ was slightly but significantly reduced in both the *srg3* and the *phr1-1 phl1* mutants
compared to the WT (Figure 5F). Taken together, *SRG3* expression is required for maintaining
fatty acid homeostasis and the establishment of light protection upon high-light stress.

SRG3 contributes to triacylglycerol accumulation upon light increase

The observation that linolenic acid accumulated in *srg3* leaves upon high light (Figure
5C) raised the question as to whether containment of free fatty acids was impaired by *srg3*
mutation. One strategy to reduce the levels of cytotoxic fatty acids in plant cells is their
incorporation into triacylglycerols (TAG) under stress, including P_i starvation (Pant *et al.*, 2015;
Yang & Benning, 2018). We therefore quantified total TAG levels and found a pronounced
accumulation after 7 h of light treatment in the wildtype (Figure 5D). In *srg3* mutants, high-
light induced TAG accumulation was significantly lower. The differences between WT and
mutant were due to reduced accumulation of TAG species with long-chain fatty acids exhibiting
higher degree of desaturation (54:6, 54:7, 54:8, 54:9) in *srg3* (Supporting Information Fig. S12),
which would be accordant with linolenic acid esterification. TAG species with polyunsaturated
54-carbon acyl chains also showed the highest changes in the WT upon high light (Figure 5E),
a pattern that is reminiscent of heat-stress induced TAG accumulation (Mueller *et al.*, 2015).
To control if high light caused a heat response in the treated plants, we analyzed transcript levels
of *HEAT SHOCK PROTEIN 70* (*HSP70*) and *HSP18.2*, previously used to discriminate light
and heat responses (Huang *et al.*, 2019). High light under our experimental conditions caused
accumulation of *HSP70* and, to a lesser extent, *HSP18.2* transcripts which was mostly confined
to the 45 min timepoint of treatment (Supporting Information Fig. S13). While *HSP70* induction
upon high light was also seen by Huang *et al.* (2019), *HSP18.2* transcript accumulation is
indicative of a mild heat response under the conditions used in this study. Thus, we cannot fully
exclude that TAG accumulation observed upon high-light treatments was triggered by higher
temperature at the leaf surface. Nevertheless, the late appearance of TAG enrichment after 7
hours of high-light treatment, as opposed to the moderate and transient induction of *HSP18.2*
at 45 min of treatment, suggests that the observed changes might indeed be triggered by high-
light treatment independently of temperature effects. Our analysis of TAG accumulation in *srg3*
mutant plants furthermore suggest that *SRG3* gene function is required for normal occurrence
of stress-induced TAG enrichment.

Discussion

P_i is intimately linked to photosynthetic metabolism, both as a substrate for
photophosphorylation, and as a potent kinetic regulator of carbon assimilation (Furbank &
Lilley, 1980; Stitt *et al.*, 1983; Preiss, 1984; Hendriks *et al.*, 2003; Marcus *et al.*, 2005).
Additionally, P_i starvation responses affect the photosynthetic machinery of plants (Giersch &
Robinson, 1987; Fredeen *et al.*, 1990; Liu *et al.*, 2024) which is partially achieved at the
transcriptional level (Wu *et al.*, 2003; Morcuende *et al.*, 2007; Bustos *et al.*, 2010; Nilsson *et*
al., 2012). Nevertheless, it has remained elusive whether changes in the metabolic P_i demand
of the chloroplast relay to the nuclear P_i starvation response. The observations presented herein
establish that the P_i response machinery involving the transcription factors PHR1 and PHL1 is
activated by an increase in photosynthetic activity, and we provide evidence that *PHR1/PHL1*
contribute to the rapid metabolic adjustment during changes in light intensity (Figure 6).

Increase in light intensity creates a transient low-P_i signal

Transcriptional responses to changes in light intensity can be observed on the timescale
of minutes (Vogel *et al.*, 2014; Huang *et al.*, 2019), but the signaling cascades behind are
incompletely elucidated. We noted that the transcriptomic changes upon a shift to high light
overlap with those observed during growth under P_i-limited conditions (Figure 2A).
Interestingly, a previous study addressing intraspecies variation of the transcriptional response
to increased irradiance recognized the induction of P_i-starvation responsive genes upon high
light in two out of three tested accessions of Arabidopsis, and PHR1-target gene induction
coincided with photosynthetic efficiency of the ecotypes (van Rooijen *et al.*, 2018). Among the
high-light and low-P_i induced transcripts, we quantified the expression of *SRG3*, *SPX1*, and
VPE1 during the early phase of high-light acclimation and determined that these transcripts
transiently accumulate already at 20 min of exposure to increased photon flux density (Figure
2B). Crucially, gene induction was absent in *phr1-1 phl1* double mutants, qualifying this short-
term high-light response as dependent on nuclear low-P_i signaling. Hence, plant cells are
capable of exploiting the transcriptional potential of the low-P_i response module in order to
react to environmental stimuli, even under nutrient-replenished conditions. Notably, high-light
mediated induction was not seen to the same extend for all PHR1 targets examined, as
expression of *MGD3* and *PS2* were only marginal (Figure 2B). This suggests that short-term
high light affects only a subset of the transcriptional P_i starvation response, potentially
reflecting differential chromatin accessibility of the genes (Barragán-Rosillo *et al.*, 2021).

How are PHR1 and PHL1 activated by an increase in light intensity? It has been established that photosynthetic activity, through the production of phosphorylated assimilates, reduces the P_i pool available for photophosphorylation (Sharkey, 1985; Walker & Sivak, 1986; Robinson & Giersch, 1987; McClain & Sharkey, 2019), but the effect on other cellular compartments than the chloroplast has remained obscure. Our analysis of subcellular P_i contents revealed a reduction of chloroplast P_i by more than one half upon an increase in light intensity of 20 min duration (Figure 3A). Meanwhile, no decline in the cytosolic/nuclear or the vacuolar fractions was observed, proving high capacity of the cell to control subcellular P_i distribution. This finding however argues against InsP8-dependent PHR1(-like) activation under these conditions. Given the large difference in vacuolar P_i levels compared to the cytoplasm, we cannot fully rule out that a difference in the P_i levels between treated and control cytosolic preparations was masked by minor vacuolar contaminations. It is also possible that a drop in cytosolic P_i levels sufficient to trigger a peak in PHR1(-like) activity occurs transiently prior to the sampling timepoint and therefore precedes transcript accumulation seen at 20 min of treatment. However, our finding is consistent with the assumption that P_i exchange between chloroplast and cytosol (Santarius & Heber, 1965), and even more so between cytosol and vacuole (Woodrow *et al.*, 1984) are rather slow processes. Additionally, the rapid increase in sucrose levels that can be observed upon high light (Figure 3C) indicates that the capacity for P_i liberation via sucrose biosynthesis might be sufficient to balance cytosolic P_i levels against depletion through triose phosphate import.

Intriguingly, our observations would be coherent with the possibility that P_i levels of the chloroplast are directly sensed to promote PHR1(-like) activation upon high light. Evidence for a role of chloroplast P_i levels in the acclimation of the photosynthetic machinery was recently provided by Raju et al. (2024) who investigated stromal P_i levels of mutants defective in P_i transporters of the chloroplast envelope using a FRET-based P_i sensor. Here, P_i levels in the stroma of *tpt*, *pht2;1*, and double mutants correlated with ΦPSII and NPQ parameters, while proton motif force and proton efflux efficiency of the ATP synthase, as determined by electrochromic shift measurements, were not affected after 2 min of a dark-light transition. The latter finding is corroborated by our result that ATP levels of WT plants are unaffected upon light increase, although ATP synthesis in the chloroplast would be expected to be kinetically constrained at the P_i concentrations measured in our system (Sharkey, 1985; McClain & Sharkey, 2019). Chloroplast P_i acting as an original retrograde signal would serve as an explanation for high constitutive PHR1(-like) activation in the *adg1 tpt-2* mutant (Figure 1E). Here, metabolically available P_i levels are likely very low given that the mutant is defective in

major pathways of triose phosphate utilization. Hence, bulk release of P_i in *adg1 tpt-2* might be
restricted to glycolysis and vacuolar phosphatase activity acting on phosphorylated organic
compounds such as phosphoenolpyruvate (Ohnishi *et al.*, 2018). This would cause unequal
intracellular pool sizes of free P_i, characterized by high vacuolar P_i stores (Figure 1D), and
particularly low P_i levels in the chloroplast. It will be insightful to examine the subcellular
distribution of P_i in the *adg1 tpt-2* mutant during future studies in order to elucidate the function
of chloroplast P_i in retrograde signaling, and in the pleiotropic phenotypes of the *adg1 tpt-2*
mutant.

PHR1/PHL1 act in the network of sugar signaling

Blocking starch production in the *phr1-3 phl1* double mutant by introduction of the
adg1-1 allele substantially restricted growth (Figure 1B, C, F). In line with this, *phr1-1 phl1*
mutants over-accumulated starch at the adult stage (Figure 4C), indicating that starch
biosynthesis aids metabolism when cellular P_i levels are chronically depleted as it is observed
in *phr1 phl1* mutants (Figure 3A, D). In fact, it is estimated that the bulk of P sequestered in
photoassimilates is liberated via starch and sucrose biosynthesis (McClain & Sharkey, 2019).
In line with the assumption that chloroplast P_i reserves are exploited through photosynthetic P_i
consumption, growth restriction of *phr1-3 phl1*, *adg1-1*, and particularly *adg1 phr1 phl1*
seedlings was more pronounced under high-light growth conditions (Figure 1F). However
counter-intuitively, this was alleviated by an additional carbohydrate source supplied via the
growth medium (Figure 1F), suggesting that metabolic alterations in these mutants extend
beyond an insufficient capacity for P_i recycling for photoassimilation. It has been concluded
from the studies on *adg1 tpt-2* mutants that sugars act as retrograde signaling molecules
communicating photosynthetic status to the nuclear genome (Heinrichs *et al.*, 2012; Häusler *et
al.*, 2014; Schmitz *et al.*, 2014). Like in *adg1 tpt-2* (Schmitz *et al.*, 2014), acclimation to high-
light growth conditions might be impaired in *adg1 phr1 phl1*, and to a lesser extend in *phr1 phl1*
mutants, resulting from a depletion of carbohydrates that occurs in the long-term of light
exposure. In fact, already under the moderate growth light conditions, we noted time-of-day
dependent deficits in sucrose levels in adult *phr1-1 phl1* plants (Figure 3C, 4E), similar to
adg1 tpt-2 mutants that lack diurnal peaks in sucrose concentrations (Schmitz *et al.*, 2012).
Moreover, *phr1 phl1* mutants exhibit a phenotype of early leaf senescence (Bustos *et al.*, 2010;
Wang *et al.*, 2023) that is reminiscent of the symptoms caused by carbohydrate depletion.
Interestingly, Schmitz et al. (2014) also noted that *adg1-1* mutation was associated with

transcriptomic changes affecting lipid metabolism as we report here for the *PHR1/PHL1*-
mediated high-light response. Finally, *phr1 phl1* mutants were specifically insensitive towards
sucrose in terms of anthocyanin production under early P_i-depleted growth conditions (Figure
4F). We therefore propose a function of PHR1/PHL1 in sugar signaling, thus contributing to
the adjustment of the C/P balance.

PHR1/PHL1 promote acclimation to higher light intensities

Our analysis of the *PHR1/PHL1* dependent transcriptional response to high light
focused on *SRG3*, *SPX1*, and *VPE1*. *VPE1* induction, putatively encoding a vacuolar efflux
transporter (Xu *et al.*, 2019), immediately suggests itself as a strategy for intracellular P_i
redistribution in favor of the cytosol by exploitation of the vacuolar P_i stores. This would serve
to provide additional P_i to metabolic pathways, which seems adequate since light increase
represents a condition that accelerates growth rate and implies higher nutrient demand in the
mid-term (Figure 3D). In contrast, *SPX1* as a highly responsive P_i starvation marker gene (Wang
et al., 2023) functions in negative feed-back of P_i starvation responses (Puga *et al.*, 2014). Such
fine-tuning of PHR1(-like) activation under light increase seems appropriate given the
fluctuating nature of this environmental factor.

The pronounced PHR1/PHL1 dependent induction of *SRG3* was further investigated
within this study. The mechanistic contribution of SRG3 to P_i homeostasis is indistinct, and it
has been reported in other plant species that GDPD isoforms play a role in phospholipid
homeostasis (Dang *et al.*, 2025), but also in hormone signaling and the modulation of root
morphology upon P_i starvation (Mai *et al.*, 2021; Hu *et al.*, 2024; Verma *et al.*, 2025). We
demonstrated here that the *SRG3* promotor is activated by over-expression of *PHR1* in
mesophyll protoplasts (Figure 5A), indicating that it constitutes a true target of the transcription
factor. To elucidate a potential physiological function of *SRG3* expression for lipid metabolism
in the light acclimation response of shoots, we set out to identify lipid features that were altered
by *srg3* mutation upon high-light treatment using an untargeted approach. Comparing WT and
srg3 rosette leaves, we detected differences in the lipid fingerprints particularly after longer
treatment duration (Figure 5B). Among the molecules that over-accumulated in *srg3* mutants
in a high-light dependent manner, we identified linolenic acid, the galactolipid derivatives acyl-
MGDG (54:8), and Arabidopside A, as well as a molecule that was constitutively upregulated
in *srg3* and putatively represents glucosylated SQDG (Figure 5C). While the latter lipid species
has to our knowledge not been reported before, accumulation of the remaining species draws

the picture that *srg3* mutants experience stress during acclimation to higher light intensities.
Both acyl-MGDG and Arabidopside accumulation have been reported to be associated with
stress conditions (Buseman *et al.*, 2006; Vu *et al.*, 2014; Nilsson *et al.*, 2016), and it was
suggested that the pool of acyl-MGDG increases upon stress to sequester oxidized fatty acids
(Vu *et al.*, 2014). Moreover, Arabidopside A has been ascribed a role in the regulation of
chlorophyll degradation during senescence processes (Weber, 2002; Hisamatsu *et al.*, 2006).
Thus, the lipid fingerprint of *srg3* upon light increase implies that the gene might be required
to maintain the integrity of photosynthetic membrane components upon changes in light
availability. The accumulation of linolenic acid in leaves of light-stressed *srg3* plants
furthermore indicates that the processing of fatty acids is disturbed in the mutant. After 7 h of
high light, we observed an increase in galactolipid species with low degree of desaturation in
WT and mutant plants (Supporting Information Fig. S9), an aspect of short-term high-light
acclimation that has been reported before (Burgos *et al.*, 2011). Such membrane remodeling
involves lipid degradation and can be assumed to generate free fatty acids with higher degree
of desaturation, including linolenic acid. Hence, SRG3 might fulfill a function in the
containment of such products derived from membrane remodeling. The product of SRG3
activity, glycerol-3-phosphate (Cheng *et al.*, 2011), is a substrate for membrane lipid
biosynthesis, as well as for TAG production (Bates, 2016), both of which also require fatty acid
precursors. To test whether any of these pathways is constrained in *srg3*, giving rise to higher
linolenic acid levels, we quantified the pools of predominant membrane lipids and TAGs. While
the levels of major lipid classes were unaffected by *srg3* mutation (Supporting Information Fig.
S8), we detected TAG accumulation in the WT after 7 h of light increase which was slightly
reduced in *srg3* mutants (Figure 5C). Consistent with a potential involvement of SRG3 in
plastoglobule formation, TAG profiles revealed that the species with strongest accumulation
under our high-light conditions also contained linolenic acid (Figure 5E, Supporting
Information Fig. S12). Notably, linolenic acid is also a component of the acyl-MGDG (54:8)
species which was enriched in *srg3* upon high light, confirming high availability of this fatty
acid in *srg3* under stress. Elevated levels of free fatty acids can be cytotoxic probably due to
their amphipathic and detergent-like properties leading to the inhibition of photosynthesis and
damage to PSII (Kunz *et al.*, 2009). Thus, our results suggest that SRG3 contributes to the
detoxification of free linolenic acid that is released during high light-induced membrane
remodeling by incorporation of excess linolenic acid into TAGs. Notably, linolenic acid
accumulation was previously also observed in Arabidopsis plants subjected to salinity stress,
and this was connected to the control of ion homeostasis via the regulation of H^+-ATPase

activity (Han *et al.*, 2017; Abdelrahman *et al.*, 2021). Hence, instead of a direct involvement of

SRG3 in the processing of fatty acids, linolenic acid accumulation in *srg3* might also be a

secondary effect related to perturbed membrane remodeling.

Interestingly, a compound that accumulated upon light increase in the WT, but was

reduced by *srg3* mutation, was identified as zeaxanthin (Figure 5C), a molecule that is involved

in the protection against excess excitation energy as part of the xanthophyll cycle (Demmig-

Adams & Adams, 1996; Havaux & Niyogi, 1999; Sacharz *et al.*, 2017). In line with a function

of both *SRG3* and *PHR1/PHL1* for proper operation of the xanthophyll cycle, assessment of the

NPQ component of chlorophyll a fluorescence revealed deficits in both mutant genotypes

(Figure 5F). It is well-known that the membrane lipid composition is important for the

xanthophyll cycle, and both lipid class and fatty acid composition determine faithful operation

of all components (Goss & Latowski, 2020). Hence, it is tempting to hypothesize that *SRG3*

expression, induced by the nuclear low-P_i response machinery, might be required for the

adjustment of the membrane environment to ensure proper function of the enzymes involved in

the xanthophyll cycle. Strikingly, this might be directly related to the changes in chloroplast P_i

levels seen under high light (Figure 3A), given the recent observations on NPQ levels of P_i-

transport mutants made by Raju et al. (2024) which were mentioned above. We therefore

propose a signaling pathway originating from photosynthetic activity that engages PHR1 and

PHL1 to induce *SRG3* expression for the acclimation of the photosynthetic membranes towards

high light.

Conclusions

High-light stress requires immediate responses to avert damage to cellular components.

As a new mode of signaling excess photosynthetic activity, we describe a function of the low-

P_i response machinery involving PHR1 and PHL1 transcription factors under nutrient replete

growth conditions. Our results indicate that chloroplast P_i sequestration into phosphorylated

photoassimilates, and/or the accumulation of these photoassimilates, create a retrograde signal

resulting in the activation of PHR1 and PHL1 in the cytosol and nucleus. This serves to mobilize

cellular P_i, and to support acclimation responses including anthocyanin production, and the

adjustment of lipid metabolism to ensure photoprotection and fatty acid homeostasis.

Materials and Methods

Plant material

All Arabidopsis plants used in this study are in the background of the Columbia-0 accession. The *phr1-3* (AT4G28610; SALK_067629C) (Nilsson *et al.*, 2007) and *phl1* (AT5G29000; SAIL_731_B09) (Klecker *et al.*, 2014) lines were described previously and the derived homozygous double mutant *phr1-3 phl1* was a kind gift of Angelika Mustroph (University of Bayreuth). The double mutant *phr1-1 phl1* (Bustos *et al.*, 2010) was backcrossed to *phl1* in order to remove a transgene containing the *NPTII* cassette which was previously introduced for isolation of the *phr1-1* allele (Rubio *et al.*, 2001). All experiments were performed with the homozygous double mutant *phr1-1 phl1* lacking the *NPTII* cassette. The double mutant *spx1 spx2* was described earlier (Puga *et al.*, 2014). Knockout lines *adg1-1* (At5g48300) (Lin *et al.*, 1988), *tpt-2* (At5g46110; SALK_073707.54.25.x) and *adg1 tpt-2* (Schmitz *et al.*, 2012) were kindly provided by Rainer E. Häusler (University of Cologne). *gpt2-1* (AT1G61800; GK-454H06) and *srg3/gdpd1-1* (AT3G02040; SALK_087106) were obtained from The Nottingham Arabidopsis Stock Centre (NASC) and were described previously (Niewiadomski *et al.*, 2005; Cheng *et al.*, 2011). Primer sequences used for genotyping are listed in Table S4. Homozygous insertion mutants were verified using the following primer combinations: *PHR1* WT: PHR1_F/PHR1_R; *phr1-3* T-DNA: LBb1/PHR1_R; *PHL1* WT: PHL1_F/PHL1_R; *phl1* T-DNA: PHL1_F/LB3; *TPT* WT: TPT_F/TPT_R; *tpt-2* T-DNA: tpt-2_F/LBb1; *GPT2* WT: GPT2_F/GPT2_R; *gpt2-1* T-DNA: GK-LB/GPT2_R; *SRG3* WT: qSRG3_F/SRG3_R; *srg3* T-DNA: qSRG3_F/LBb1. To generate the *adg1 phr1 phl1* triple mutant, *adg1-1* was crossed to *phr1-3 phl1*. Homozygous *adg1-1* mutants were identified by screening for starch-free phenotypes by iodine staining at the end of the photoperiod. Homozygosity of the *phr1-1* allele was verified by Sanger sequencing (Macrogen Europe, Amsterdam) of the PCR product of PHR1_F/PHR1_R using primer PHR1_R.

Cultivation conditions

Seeds were stratified for 64 hours at 4°C in the dark. For vegetative growth, plants were germinated and grown in an 8 h/16 h (light/dark) cycle and 23°C/22°C day/night temperature on soil consisting of seeding compost, universal soil, and vermiculite mixed in a 3:3:1 ratio. For high-light experiments, the plants were additionally fertilized once directly after pricking at the age of 10 days using a complete mineral mixture (Wuxal Super, Aglukon Spezialduenger,

Duesseldorf, Germany) according to the manufacturer's instructions. For seedling experiments,

seeds were surface sterilized by chlorine gas exposure and transferred to solidified Arabidopsis

growth media as described in the figure legends. Seedlings were grown under 16 h/8 h light/dark

cycles. If not indicated otherwise, growth light was set to 90 ± 10 µmol m^{-2} sec^{-1} (for seedling

growth and for experiments shown in Figure 1B-E), or 70 ± 5 µmol m^{-2} s^{-1} (before high-light

treatments at 450 ± 30 µmol m^{-2} s^{-1} of rosette plants), and monitored using an illuminance meter.

Growth and treatments of rosette plants for high-light experiments were conducted in a Percival

SE-41L plant growth incubator equipped with dimmable fluorescent lamps under temperature

control.

P$_i$-limitation experiments

For P$_i$-limitation experiments, seeds were germinated on medium containing MS salts

at full strength together with 0.8 % (w/v) phytoagar (Duchefa Biochemie, Haarlem, The

Netherlands) and 0.5 % (w/v) sucrose. After 5-10 days of growth (see figure legends), seedlings

were transferred to P$_i$ media (0.5 % (w/v) sucrose; 20 mM 2-(N-Morpholino)-ethane sulphonic

acid; 2.5 mM KNO$_3$; 1 mM MgSO$_4$; 1 mM Ca(NO$_3$)$_2$; 2.5 mM KH$_2$PO$_4$; 25 µM Fe-EDTA;

35 µM H$_3$BO$_3$; 7 µM MnCl$_2$; 0.25 µM CuSO$_4$; 0.5 µM ZNSO$_4$; 0.1 µM NaMoO$_4$; 5 µM NaCl;

0.005 µM CoCl$_2$; pH 6) solidified by 0.8 % (w/v) agar (adopted from Härtel et al. (2000)). For

P$_i$ depleting conditions, KH$_2$PO$_4$ was omitted, and Fe-EDTA was reduced to 10 µM in order to

minimize low-P$_i$ induced iron toxicity (Ward *et al.*, 2008). Seedling shoots were harvested after

7-8 days of growth on P$_i$ differing media (see figure legends). For the experiment shown in

Figures 1F and Supporting Information Fig. S3, P$_i$ medium was used containing either 0.25 mM

KH$_2$PO$_4$/10 µM Fe-EDTA, 2.5 mM KH$_2$PO$_4$/25 µM Fe-EDTA, or 2.5 mM KH$_2$PO$_4$/25 µM Fe-

EDTA +50 mM sucrose. For the experiment shown in Figure 4F, seedlings were germinated on

MS salts at ½ strength without added sucrose and grown for 7 days before transfer to P$_i$-

differing conditions.

Vector construction

Standard molecular cloning procedures were applied. Primer sequences used for cloning

are listed in table S4. The firefly *LUCIFERASE* reporter plasmid *pBT10-proMGD3::LUC$_{Firefly}$*

and the *pHBTL-p35S::3HA-GFP* effector plasmid (Klecker *et al.*, 2014), as well as the *pBT10-*

p35S::LUC$_{Renilla}$ normalization plasmid (Bäumler *et al.*, 2019) have been described elsewhere.

For details on cloning *proSPX1^{GC}, proSRG3, proMGD3, proGPT2,* and the *PHR1* effector

construct, refer to the supplemental methods.

Anthocyanin determination

Anthocyanin contents were determined with few changes as described in

(Vandenbussche *et al.*, 2007). In brief, 40-70 mg of seedling shoots or rosette leaf material were

frozen in liquid nitrogen and ground in a bead mill (MM400, Retsch, Haan, Germany). The

frozen powder was resuspended in 300 µl of methanol containing 1 % HCl. After vortex-mixing

for 15 sec, cell debris was pelleted by centrifugation for 15 min at >14,000 g, 4°C. The

supernatant was mixed with 200 µl water and 500 µl chloroform by vortexing for 15 sec. After

15 min centrifugation at 17,900 g, 4°C, the upper phase was diluted in methanol to measure the

absorbance at 535 nm and 630 nm using a spectrophotometer (Specord 200 Plus, Analytik Jena,

Jena, Germany).

Measurements of starch, soluble sugars, and adenylates

747 Carbohydrates were determined by the Warburg optical test and adenylates were

748 assessed based on luciferase activity. Both procedures were described previously (Mustroph *et*

749 *al.*, 2006), for details see supplemental methods.

Determination of P$_i$ contents

752 Free P$_i$ was determined in leaf tissue based on the method described by (Ames, 1966)

753 with some modifications partially based on (Sakuraba *et al.*, 2018). For the experiment shown

754 in Figure 3D, leaf material was lyophilized prior to extraction, and 50 µl extraction buffer per

755 mg plant material was used. For details see supplemental methods.

Non-aqueous fractionation of leaf material

For the analysis of subcellular P$_i$ distribution, 12 plant rosettes (total of >1 g) were

harvested into vials precooled by liquid nitrogen and dry ice, with an averaged harvesting time

of 3 seconds (max. 7 seconds) per rosette. The plant material was ground to a fine powder and

lyophilized. The freeze-dried powder was suspended in a mixture of tetrachlorethylene and n-

heptane and fractionated across a non-aqueous density gradient as described before (Hernandez

et al., 2023). Following fractionation, marker enzyme activities for plastids (alkaline

pyrophosphatase), cytosol (UDP-glucose pyrophosphorylase), and vacuole (acidic

phosphatase) were determined photometrically. Phosphate amounts were determined as

described above for each fraction and correlated with the marker enzyme activities using the

using the NAFalyzer app (https://github.com/cellbiomaths/NAFalyzer).

RNA extraction and cDNA synthesis

Plant material was harvested under the respective experimental conditions and instantly

frozen in liquid nitrogen. The tissue was ground in a bead mill, and RNA was extracted using

Bioline TRIsure (Meridian Bioscience, Cincinnati, United States) according to the

manufacturer's instructions. The quality of the RNA was assessed by screening A_{260} and A_{280}.

For qRT-PCR analysis, 1 µg of RNA was treated with DNAse I (Thermo Fisher Scientific,

Waltham, United States) according to product instructions before application in reverse

transcription using oligo(dT)$_{15}$ and RevertAid Reverse Transcriptase (Thermo Fisher

Scientific). Complementary DNA was diluted by a factor of 50 for use as template in qPCR

analysis. qPCR was performed in technical triplicates using SsoAdvanced Universal SYBR

Green Supermix (Bio-Rad, Hercules, United States) on a CFX Connect Real-time System (Bio-

Rad). Transcript levels were normalized to the levels of *PP2A* transcript and calculated as

$1000 \cdot 2^{-\Delta CT}$. Primer sequences are listed in table S4.

Protoplast isolation, transfection and transactivation assay

Leaf mesophyll protoplasts were isolated using the "Tape-Arabidopsis Sandwich"

method (Wu *et al.*, 2009) with minor modifications. For details, see supplemental methods.

Chlorophyll fluorescence measurements

For the determination of the quantum yield of photosystem II photochemistry

$(F_m'-F')/F_m'$ (ΦPSII), as well as the maximum efficiency of photosystem II (F_v'/F_m'), and the

NPQ parameter (or SV_N; calculated by the Junior-PAM software using the Stern-Volmer

equation for fluorescence quenching, Gilmore and Yamamoto (1991)), pulse-amplitude-

modulated fluorescence measurements (PAM) of chlorophyll fluorescence were conducted

using a Junior-PAM (Walz, Effeltrich, Germany). Here, to calculate chlorophyll fluorescence

parameters using the saturating pulse method (Schreiber *et al.*, 1986), one large rosette leaf was

held between two magnets and subjected to 10 seconds of actinic light irradiance set to control

light intensity (70±5 µmol m^{-2} sec^{-1}), followed by a saturation light pulse.

Lipid analysis

Lipid analysis was performed according to (Mueller *et al.*, 2015). Progenesis QI

(version 2.1, Waters) was used for data pre-processing and PCA was performed in the

MetaboAnalyst platform (version 6.0, www.metaboanalyst.ca).

Statistical analysis

Statistical analyses were performed with Microsoft Office Excel using the Real

Statistics Resource Pack software (www.real-statistics.com). Two-factor ANOVA and

hypergeometric testing were performed using RStudio (version 2024.09.1+394) available at

https://www.r-project.org/.

Funding information

This research was funded by the University of Bayreuth and by the University of

Würzburg.

Acknowledgements

We thank Stephan Clemens and Angelika Mustroph for helpful discussions and

comments on the manuscript, as well as for providing infrastructure and equipment. We thank

Susanne Berger for help with choosing experimental conditions used for lipid analyses.

Author contributions

M.K., conceptualization; A.F., T.N., and M.K., methodology; L.A., A.F., T.N., and

M.K., formal analysis; L.A., A.F., T.N., A.H., M.M., and M.K., investigation; M.K., writing–

original draft; all authors, writing–review and editing; A.F. and M.K., visualization; M.J.M and

M.K., supervision; M.K., project administration.

Competing interests

None declared.

Data availability

The source data of lipid analyses are available in the Supporting information of this article.

Supplemental Data

Supporting Information Fig. S1 Phenotypes of *adg1-1 phr1-3 phl1* (TM).

Supporting Information Fig. S2 Expression of selected P_i-starvation responsive genes depends on *PHR1/PHL1* under P_i depletion.

Supporting Information Fig. S3 Root growth of *adg1 phr1 phl1* and *adg1 tpt-2* responds to exogenously applied sucrose.

Supporting Information Fig. S4 *PHR1* transcript levels are not affected by short-time high-light exposure.

Supporting Information Fig. S5 Changes in sugar and starch contents under P_i depletion.

Supporting Information Fig. S6 Monosaccharide levels after 20 min of high light.

Supporting Information Fig. S7 Anthocyanin biosynthetic and regulatory gene expression upon high light in WT and *phr1-1 phl1*.

Supporting Information Fig. S8 Total levels of lipid classes in WT (Col-0) and *srg3* mutants upon shift to high light (HL).

Supporting Information Fig. S9 Contents of glycosylglycerol lipid species in WT (Col-0) and *srg3* mutants upon shift to high light (HL) relative to fresh weight (FW).

Supporting Information Fig. S10 Contents of glycerophospholipid species in WT (Col-0) and *srg3* mutants upon shift to high light (HL) relative to fresh weight (FW).

Supporting Information Fig. S11 Levels of zeaxanthin and putative glycosylated SQDG determined in opposite ESI ion mode compared to Figure 5C.

Supporting Information Fig. S12 Levels of triacylglycerol (TAG) species in WT (Col-0) and *srg3* mutants upon shift to high light (HL).

Supporting Information Fig. S13 Levels of high-temperature responsive transcripts *HSP70*

and *HSP18.2* in WT (Col-0) and *phr1-1 phl1* mutant plants upon high light treatment.

Supporting Information Table S1 Annotated lipid features in WT (Col-0) and *srg3* mutants

following a shift to high light.

Supporting Information Table S2 Lipid features whose levels were significantly altered upon

shifting to the WT (Col-0).

Supporting Information Table S3 Heat-responsive lipid features whose levels differed

significantly between WT (Col-0) and *srg3* mutants.

Supporting Information Table S4 List of primer sequences used in this study.

Supplemental Methods and References

REFERENCES

Abdelrahman M, Nishiyama R, Tran CD, Kusano M, Nakabayashi R, Okazaki Y, Matsuda F, Chávez Montes RA, Mostofa MG, Li W, *et al.* 2021. Defective cytokinin signaling reprograms lipid and flavonoid gene-to-metabolite networks to mitigate high salinity in Arabidopsis. *Proceedings of the National Academy of Sciences of the United States of America* **118**.

Ames BN. **1966**. [10] Assay of inorganic phosphate, total phosphate and phosphatases. In: Methods in Enzymology. 115–118.

Athanasiou K, Dyson BC, Webster RE, Johnson GN. **2010**. Dynamic acclimation of photosynthesis increases plant fitness in changing environments. *Plant physiology* **152**: 366–73.

Baena-González E, Rolland F, Thevelein JM, Sheen J. **2007**. A central integrator of transcription networks in plant stress and energy signalling. *Nature* **448**: 938–42.

Barragán-Rosillo AC, Peralta-Alvarez CA, Ojeda-Rivera JO, Arzate-Mejía RG, Recillas-Targa F, Herrera-Estrella L. **2021**. Genome accessibility dynamics in response to phosphate limitation is controlled by the PHR1 family of transcription factors in Arabidopsis. *Proceedings of the National Academy of Sciences of the United States of America* **118**.

Bates PD. **2016**. Understanding the control of acyl flux through the lipid metabolic network of plant oil biosynthesis. *Biochimica et Biophysica Acta (BBA) - Molecular and Cell Biology of Lipids* **1861**: 1214–1225.

Bäumler J, Riber W, Klecker M, Müller L, Dissmeyer N, Weig AR, Mustroph A. **2019**. AtERF#111/ABR1 is a transcriptional activator involved in the wounding response. *The Plant journal : for cell and molecular biology* **100**: 969–990.

Bechtold U, Richard O, Zamboni A, Gapper C, Geisler M, Pogson B, Karpinski S, Mullineaux PM. **2008**. Impact of chloroplastic- and extracellular-sourced ROS on high light-responsive gene expression in Arabidopsis. *Journal of experimental botany* **59**: 121–33.

Borghi GL, Moraes TA, Günther M, Feil R, Mengin V, Lunn JE, Stitt M, Arrivault S. **2019**. Relationship between irradiance and levels of Calvin-Benson cycle and other intermediates in the model eudicot Arabidopsis and the model monocot rice. *Journal of Experimental Botany* **70**: 5809–5825.

Burgos A, Szymanski J, Seiwert B, Degenkolbe T, Hannah MA, Giavalisco P, Willmitzer

L. **2011**. Analysis of short-term changes in the Arabidopsis thaliana glycerolipidome in
response to temperature and light. *Plant Journal* **66**: 656–668.

Buseman CM, Tamura P, Sparks AA, Baughman EJ, Maatta S, Zhao J, Roth MR, Esch
SW, Shah J, Williams TD, *et al.* **2006**. Wounding stimulates the accumulation of glycerolipids
containing oxophytodienoic acid and dinor-oxophytodienoic acid in arabidopsis leaves. *Plant*
Physiology **142**: 28–39.

Bustos R, Castrillo G, Linhares F, Puga MI, Rubio V, Pérez-Pérez J, Solano R, Leyva A,
Paz-Ares J. **2010**. A central regulatory system largely controls transcriptional activation and
repression responses to phosphate starvation in Arabidopsis. *PLoS genetics* **6**: e1001102.

Castrillo G, Teixeira PJPL, Paredes SH, Law TF, de Lorenzo L, Feltcher ME, Finkel OM,
Breakfield NW, Mieczkowski P, Jones CD, *et al.* **2017**. Root microbiota drive direct
integration of phosphate stress and immunity. *Nature* **543**: 513–518.

Cheng Y, Zhou W, El Sheery NI, Peters C, Li M, Wang X, Huang J. **2011**. Characterization
of the Arabidopsis glycerophosphodiester phosphodiesterase (GDPD) family reveals a role of
the plastid-localized AtGDPD1 in maintaining cellular phosphate homeostasis under phosphate
starvation. *Plant Journal* **66**: 781–795.

Dang DTT, Nguyen LT, Tran A-T, Tran TTT, Tran T-A, Vo KTX, Jeon J-S, Van Vu T,
Kim J-Y, Nguyen NP, *et al.* **2025**. A novel glycerophosphodiester phosphodiesterase 13 is
involved in the phosphate starvation-induced phospholipid degradation in rice. *Plant*
physiology and biochemistry : PPB **228**: 110230.

Demmig-Adams B, Adams WW. **1996**. The role of xanthophyll cycle carotenoids in the
protection of photosynthesis. *Trends in Plant Science* **1**: 21–26.

Dietz K-J. **2015**. Efficient high light acclimation involves rapid processes at multiple
mechanistic levels. *Journal of experimental botany* **66**: 2401–14.

Dietz KJ, Foyer C. **1986**. The relationship between phosphate status and photosynthesis in
leaves : Reversibility of the effects of phosphate deficiency on photosynthesis. *Planta* **167**:
376–81.

Dietz K, Heber U. **1984**. Rate-limiting factors in leaf photosynthesis. I. Carbon fluxes in the
calvin cycle. *Biochimica et Biophysica Acta (BBA) - Bioenergetics* **767**: 432–443.

Dietz K-J, Heber U. **1986**. Light and CO2 limitation of photosynthesis and states of the
reactions regenerating ribulose 1,5-bisphosphate or reducing 3-phosphoglycerate. *Biochimica*

et Biophysica Acta (BBA) - Bioenergetics **848**: 392–401.

Dietz K-J, Turkan I, Krieger-Liszkay A. **2016**. Redox- and Reactive Oxygen Species-
Dependent Signaling into and out of the Photosynthesizing Chloroplast. *Plant physiology* **171**:
1541–50.

Dong J, Ma G, Sui L, Wei M, Satheesh V, Zhang R, Ge S, Li J, Zhang T-E, Wittwer C, *et*
al. **2019**. Inositol Pyrophosphate InsP8 Acts as an Intracellular Phosphate Signal in
Arabidopsis. *Molecular plant* **12**: 1463–1473.

Figueroa CM, Asencion Diez MD, Ballicora MA, Iglesias AA. **2022**. Structure, function, and
evolution of plant ADP-glucose pyrophosphorylase. *Plant molecular biology* **108**: 307–323.

Franco-Zorrilla JM, Martín AC, Leyva A, Paz-Ares J. **2005**. Interaction between
phosphate-starvation, sugar, and cytokinin signaling in Arabidopsis and the roles of cytokinin
receptors CRE1/AHK4 and AHK3. *Plant Physiology* **138**: 847–857.

Frank A, Matiolli CC, Viana AJC, Hearn TJ, Kusakina J, Belbin FE, Wells Newman D,
Yochikawa A, Cano-Ramirez DL, Chembath A, *et al.* **2018**. Circadian Entrainment in
Arabidopsis by the Sugar-Responsive Transcription Factor bZIP63. *Current biology : CB* **28**:
2597-2606.e6.

Fredeen AL, Raab TK, Rao IM, Terry N. **1990**. Effects of phosphorus nutrition on
photosynthesis in Glycine max (L.) Merr. *Planta* **181**: 399–405.

Furbank RT, Lilley RM. **1980**. Effects of inorganic phosphate on the photosynthetic carbon
reduction cycle in extracts from the stroma of pea chloroplasts. *Biochimica et biophysica acta*
592: 65–75.

Giersch C, Robinson SP. **1987**. Regulation of photosynthetic carbon metabolism during
phosphate limitation of photosynthesis in isolated spinach chloroplasts. *Photosynthesis*
Research **14**: 211–227.

Gilmore AM, Yamamoto HY. **1991**. Zeaxanthin Formation and Energy-Dependent
Fluorescence Quenching in Pea Chloroplasts under Artificially Mediated Linear and Cyclic
Electron Transport. *Plant physiology* **96**: 635–43.

Goss R, Latowski D. **2020**. Lipid Dependence of Xanthophyll Cycling in Higher Plants and
Algae. *Frontiers in plant science* **11**: 455.

Han X, Yang Y, Wu Y, Liu X, Lei X, Guo Y. **2017**. A bioassay-guided fractionation system
to identify endogenous small molecules that activate plasma membrane H + -ATPase activity

in Arabidopsis. *Journal of Experimental Botany* **68**: 2951–2962.

Hanchi M, Thibaud M-C, Légeret B, Kuwata K, Pochon N, Beisson F, Cao A, Cuyas L,
David P, Doerner P, *et al*. 2018. The Phosphate Fast-Responsive Genes PECP1 and PPsPase1
Affect Phosphocholine and Phosphoethanolamine Content. *Plant physiology* **176**: 2943–2962.

Härtel H, Dormann P, Benning C. 2000. DGD1-independent biosynthesis of extraplastidic
galactolipids after phosphate deprivation in Arabidopsis. *Proceedings of the National Academy*
of Sciences of the United States of America **97**: 10649–54.

Häusler RE, Heinrichs L, Schmitz J, Flügge UI. 2014. How sugars might coordinate
chloroplast and nuclear gene expression during acclimation to high light intensities. *Molecular*
Plant **7**: 1121–1137.

Havaux M, Niyogi KK. 1999. The violaxanthin cycle protects plants from photooxidative
damage by more than one mechanism. *Proceedings of the National Academy of Sciences of the*
United States of America **96**: 8762–8767.

He Y, Zhang X, Li L, Sun Z, Li J, Chen X, Hong G. 2021. SPX4 interacts with both PHR1
and PAP1 to regulate critical steps in phosphorus-status-dependent anthocyanin biosynthesis.
The New phytologist **230**: 205–217.

Heinrichs L, Schmitz J, Flügge UI, Häusler RE. 2012. The mysterious rescue of adg1-1/tpt-
2 - An Arabidopsis thaliana double mutant impaired in acclimation to high light - By
exogenously supplied sugars. *Frontiers in Plant Science* **3**.

Heldt HW, Rapley L. 1970. Specific transport of inorganic phosphate, 3-phosphoglycerate
and dihydroxyacetonephosphate, and of dicarboxylates across the inner membrane of spinach
chloroplasts. *FEBS letters* **10**: 143–148.

Hendriks JHM, Kolbe A, Gibon Y, Stitt M, Geigenberger P. 2003. ADP-glucose
pyrophosphorylase is activated by posttranslational redox-modification in response to light and
to sugars in leaves of Arabidopsis and other plant species. *Plant physiology* **133**: 838–49.

Hernandez JS, Dziubek D, Schröder L, Seydel C, Kitashova A, Brodsky V, Nägele T. 2023.
Natural variation of temperature acclimation of Arabidopsis thaliana. *Physiologia plantarum*
175: e14106.

Hisamatsu Y, Goto N, Hasegawa K, Shigemori H. 2006. Senescence-promoting effect of
arabidopside A. *Zeitschrift fur Naturforschung - Section C Journal of Biosciences* **61**: 363–366.

Hu D, Cui R, Wang K, Yang Y, Wang R, Zhu H, He M, Fan Y, Wang L, Wang L, *et al.*

2024. The Myb73-GDPD2-GA2ox1 transcriptional regulatory module confers phosphate
deficiency tolerance in soybean. *Plant Cell* **36**: 2176–2200.

Huang J, Zhao X, Chory J. **2019**. The Arabidopsis Transcriptome Responds Specifically and
Dynamically to High Light Stress. *Cell Reports* **29**: 4186-4199.e3.

Ibrahim A, Schütz AL, Galano JM, Herrfurth C, Feussner K, Durand T, Brodhun F,
Feussner I. **2011**. The alphabet of galactolipids in Arabidopsis thaliana. *Frontiers in Plant*
Science **2**: 1–24.

Karthikeyan AS, Varadarajan DK, Jain A, Held MA, Carpita NC, Raghothama KG.
2007. Phosphate starvation responses are mediated by sugar signaling in Arabidopsis. *Planta*
225: 907–918.

Klecker M, Gasch P, Peisker H, Dörmann P, Schlicke H, Grimm B, Mustroph A. **2014**. A
Shoot-Specific Hypoxic Response of Arabidopsis Sheds Light on the Role of the Phosphate-
Responsive Transcription Factor PHOSPHATE STARVATION RESPONSE1. *Plant*
physiology **165**: 774–790.

Kobayashi K, Awai K, Takamiya K, Ohta H. **2004**. Arabidopsis type B
monogalactosyldiacylglycerol synthase genes are expressed during pollen tube growth and
induced by phosphate starvation. *Plant physiology* **134**: 640–8.

König J, Muthuramalingam M, Dietz K-J. **2012**. Mechanisms and dynamics in the
thiol/disulfide redox regulatory network: transmitters, sensors and targets. *Current opinion in*
plant biology **15**: 261–8.

Krapp A, Berthomé R, Orsel M, Mercey-Boutet S, Yu A, Castaings L, Elftieh S, Major H,
Renou JP, Daniel-Vedele F. **2011**. Arabidopsis roots and shoots show distinct temporal
adaptation patterns toward nitrogen starvation. *Plant Physiology* **157**: 1255–1282.

Kumar V, Sharma A, Bhardwaj R, Thukral AK. **2019**. Elemental Composition of Plants
and Multivariate Analysis. *National Academy Science Letters* **42**: 45–50.

Kunz HH, Häusler RE, Fettke J, Herbst K, Niewiadomski P, Gierth M, Bell K, Steup M,
Flügge U-I, Schneider A. **2010**. The role of plastidial glucose-6-phosphate/phosphate
translocators in vegetative tissues of Arabidopsis thaliana mutants impaired in starch
biosynthesis. *Plant biology (Stuttgart, Germany)* **12 Suppl 1**: 115–28.

Kunz H-H, Scharnewski M, Feussner K, Feussner I, Flügge U-I, Fulda M, Gierth M. **2009**.
The ABC transporter PXA1 and peroxisomal beta-oxidation are vital for metabolism in mature

leaves of Arabidopsis during extended darkness. *The Plant cell* **21**: 2733–49.

Li H, He K, Zhang Z, Hu Y. **2023**. Molecular mechanism of phosphorous signaling inducing
anthocyanin accumulation in Arabidopsis. *Plant Physiology and Biochemistry* **196**: 121–129.

Li L, Sheen J. **2016**. Dynamic and diverse sugar signaling. *Current Opinion in Plant Biology*
33: 116–125.

Lin TP, Caspar T, Somerville C, Preiss J. **1988**. Isolation and Characterization of a Starchless
Mutant of Arabidopsis thaliana (L.) Heynh Lacking ADPglucose Pyrophosphorylase Activity.
Plant physiology **86**: 1131–5.

Liu Z, Wu X, Wang E, Liu Y, Wang Y, Zheng Q, Han Y, Chen Z, Zhang Y. **2022**. PHR1
positively regulates phosphate starvation-induced anthocyanin accumulation through direct
upregulation of genes F3'H and LDOX in Arabidopsis. *Planta* **256**: 42.

Liu Y, Xie Y, Wang H, Ma X, Yao W, Wang H. **2017**. Light and Ethylene Coordinately
Regulate the Phosphate Starvation Response through Transcriptional Regulation of
PHOSPHATE STARVATION RESPONSE1. *The Plant cell* **29**: 2269–2284.

Liu S, Xu Z, Essemine J, Liu Y, Liu C, Zhang F, Iqbal Z, Qu M. **2024**. GWAS unravels
acid phosphatase ACP2 as a photosynthesis regulator under phosphate starvation conditions
through modulating serine metabolism in rice. *Plant communications* **5**: 100885.

Mai NTP, Mai CD, Nguyen H Van, Le KQ, Duong LV, Tran TA, To HTM. **2021**.
Discovery of new genetic determinants of morphological plasticity in rice roots and shoots
under phosphate starvation using GWAS. *Journal of plant physiology* **257**: 153340.

Marcus Y, Altman-Gueta H, Finkler A, Gurevitz M. **2005**. Mutagenesis at two distinct
phosphate-binding sites unravels their differential roles in regulation of Rubisco activation and
catalysis. *Journal of Bacteriology* **187**: 4222–4228.

Matiolli CC, Soares RC, Alves HLS, Abreu IA. **2022**. Turning the Knobs: The Impact of
Post-translational Modifications on Carbon Metabolism. *Frontiers in plant science* **12**: 781508.

McClain AM, Sharkey TD. **2019**. Triose phosphate utilization and beyond: From
photosynthesis to end product synthesis. *Journal of Experimental Botany* **70**: 1755–1766.

Misson J, Raghothama KG, Jain A, Jouhet J, Block MA, Bligny R, Ortet P, Creff A,
Somerville S, Rolland N, *et al.* 2005. A genome-wide transcriptional analysis using
Arabidopsis thaliana Affymetrix gene chips determined plant responses to phosphate
deprivation. *Proceedings of the National Academy of Sciences of the United States of America*

102: 11934–11939.

Moore M, Vogel MO, Dietz KJ. **2014**. The acclimation response to high light is initiated
within seconds as indicated by upregulation of AP2/ERF transcription factor network in
Arabidopsis thaliana. *Plant Signaling and Behavior* **9**: 1–4.

Morcuende R, Bari R, Gibon Y, Zheng W, Pant BD, Bläsing O, Usadel B, Czechowski T,
Udvardi MK, Stitt M, *et al.* **2007**. Genome-wide reprogramming of metabolism and regulatory
networks of Arabidopsis in response to phosphorus. *Plant, Cell and Environment* **30**: 85–112.

Mueller SP, Krause DM, Mueller MJ, Fekete A. **2015**. Accumulation of extra-chloroplastic
triacylglycerols in Arabidopsis seedlings during heat acclimation. *Journal of Experimental*
Botany **66**: 4517–4526.

Müller R, Morant M, Jarmer H, Nilsson L, Nielsen TH. **2007**. Genome-wide analysis of the
arabidopsis leaf transcriptome reveals interaction of phosphate and sugar metabolism. *Plant*
Physiology **143**: 156–171.

Müller R, Nilsson L, Nielsen LK, Nielsen TH. **2005**. Interaction between phosphate starvation
signalling and hexokinase-independent sugar sensing in Arabidopsis leaves. *Physiologia*
Plantarum **124**: 81–90.

Mustroph A, Boamfa EI, Laarhoven LJJ, Harren FJM, Albrecht G, Grimm B. **2006**.
Organ-specific analysis of the anaerobic primary metabolism in rice and wheat seedlings. I:
Dark ethanol production is dominated by the shoots. *Planta* **225**: 103–14.

Niewiadomski P, Knappe S, Geimer S, Fischer K, Schulz B, Unte US, Rosso MG, Ache P,
Flügge UI, Schneider A. **2005**. The Arabidopsis plastidic glucose 6-phosphate/phosphate
translocator GPT1 is essential for pollen maturation and embryo sac development. *Plant Cell*
17: 760–775.

Nilsson AK, Fahlberg P, Johansson ON, Hamberg M, Andersson MX, Ellerström M.
2016. The activity of HYDROPEROXIDE LYASE 1 regulates accumulation of galactolipids
containing 12-oxo-phytodienoic acid in Arabidopsis. *Journal of Experimental Botany* **67**:
5133–5144.

Nilsson L, Lundmark M, Jensen PE, Nielsen TH. **2012**. The Arabidopsis transcription factor
PHR1 is essential for adaptation to high light and retaining functional photosynthesis during
phosphate starvation. *Physiologia plantarum* **144**: 35–47.

Nilsson L, Müller R, Nielsen TH. **2007**. Increased expression of the MYB-related transcription

factor, PHR1, leads to enhanced phosphate uptake in Arabidopsis thaliana. *Plant, cell & environment* **30**: 1499–512.

Ohnishi M, Anegawa A, Sugiyama Y, Harada K, Oikawa A, Nakayama Y, Matsuda F, Nakamura Y, Sasaki R, Shichijo C, et al. 2018. Molecular Components of Arabidopsis Intact Vacuoles Clarified with Metabolomic and Proteomic Analyses. *Plant & cell physiology* **59**: 1353–1362.

Pant BD, Burgos A, Pant P, Cuadros-Inostroza A, Willmitzer L, Scheible WR. **2015**. The transcription factor PHR1 regulates lipid remodeling and triacylglycerol accumulation in Arabidopsis thaliana during phosphorus starvation. *Journal of Experimental Botany* **66**: 1907–1918.

Preiss J. **1984**. Starch, sucrose biosynthesis and partition of carbon in plants are regulated by orthophosphate and triose-phosphates. *Trends in Biochemical Sciences* **9**: 24–27.

Puga MI, Mateos I, Charukesi R, Wang Z, Franco-Zorrilla JM, de Lorenzo L, Irigoyen ML, Masiero S, Bustos R, Rodríguez J, et al. 2014. SPX1 is a phosphate-dependent inhibitor of Phosphate Starvation Response 1 in Arabidopsis. *Proceedings of the National Academy of Sciences of the United States of America* **111**: 14947–52.

Raju AS, Kramer DM, Versaw WK. **2024**. Genetically manipulated chloroplast stromal phosphate levels alter photosynthetic efficiency. *Plant Physiology* **196**: 385–396.

Ramaiah M, Jain A, Baldwin JC, Karthikeyan AS, Raghothama KG. **2011**. Characterization of the phosphate starvation-induced glycerol-3-phosphate permease gene family in Arabidopsis. *Plant physiology* **157**: 279–91.

Ried MK, Wild R, Zhu J, Pipercevic J, Sturm K, Broger L, Harmel RK, Abriata LA, Hothorn LA, Fiedler D, et al. 2021. Inositol pyrophosphates promote the interaction of SPX domains with the coiled-coil motif of PHR transcription factors to regulate plant phosphate homeostasis. *Nature communications* **12**: 384.

Riemer E, Qiu D, Laha D, Harmel RK, Gaugler P, Gaugler V, Frei M, Hajirezaei M-R, Laha NP, Krusenbaum L, et al. 2021. ITPK1 is an InsP6/ADP phosphotransferase that controls phosphate signaling in Arabidopsis. *Molecular plant* **14**: 1864–1880.

Robinson S, Giersch C. **1987**. Inorganic Phosphate Concentration in the Stroma of Isolated Chloroplasts and Its Influence on Photosynthesis. *Australian Journal of Plant Physiology* **14**: 451.

van Rooijen R, Harbinson J, Aarts MGM. **2018**. Photosynthetic response to increased
irradiance correlates to variation in transcriptional response of lipid-remodeling and heat-shock
genes. *Plant direct* **2**: e00069.

Rubio V, Linhares F, Solano R, Martín AC, Iglesias J, Leyva A, Paz-Ares J. **2001**. A
conserved MYB transcription factor involved in phosphate starvation signaling both in vascular
plants and in unicellular algae. *Genes & development* **15**: 2122–33.

Sacharz J, Giovagnetti V, Ungerer P, Mastroianni G, Ruban A V. 2017. The xanthophyll
cycle affects reversible interactions between PsbS and light-harvesting complex II to control
non-photochemical quenching. *Nature plants* **3**: 16225.

Safi A, Medici A, Szponarski W, Ruffel S, Lacombe B, Krouk G. **2017**. The world according
to GARP transcription factors. *Current Opinion in Plant Biology* **39**: 159–167.

Sakuraba Y, Kanno S, Mabuchi A, Monda K, Iba K, Yanagisawa S. **2018**. A phytochrome-
B-mediated regulatory mechanism of phosphorus acquisition. *Nature plants* **4**: 1089–1101.

Santarius KA, Heber U. **1965**. Changes in the intracellular levels of ATP, ADP, AMP and P1
and regulatory function of the adenylate system in leaf cells during photosynthesis. *Biochimica*
et biophysica acta **102**: 39–54.

Schmitz J, Heinrichs L, Scossa F, Fernie AR, Oelze M-L, Dietz K-J, Rothbart M, Grimm
B, Flügge U-I, Häusler RE. **2014**. The essential role of sugar metabolism in the acclimation
response of Arabidopsis thaliana to high light intensities. *Journal of experimental botany* **65**:
1619–36.

Schmitz J, Schöttler MA, Krueger S, Geimer S, Schneider A, Kleine T, Leister D, Bell K,
Flügge U-I, Häusler RE. **2012**. Defects in leaf carbohydrate metabolism compromise
acclimation to high light and lead to a high chlorophyll fluorescence phenotype in Arabidopsis
thaliana. *BMC plant biology* **12**: 8.

Schneider A, Häusler RE, Kolukisaoglu U, Kunze R, van der Graaff E, Schwacke R,
Catoni E, Desimone M, Flügge U-I. **2002**. An Arabidopsis thaliana knock-out mutant of the
chloroplast triose phosphate/phosphate translocator is severely compromised only when starch
synthesis, but not starch mobilisation is abolished. *The Plant journal : for cell and molecular*
biology **32**: 685–99.

Schreiber U, Schliwa U, Bilger W. **1986**. Continuous recording of photochemical and non-
photochemical chlorophyll fluorescence quenching with a new type of modulation fluorometer.
Photosynthesis Research **10**: 51–62.

Sharkey TD. **1985**. O(2)-insensitive photosynthesis in c(3) plants : its occurrence and a
possible explanation. *Plant physiology* **78**: 71–5.

Sivak MN, Walker DA. **1986**. Photosynthesis in vivo can be limited by phosphate supply. *New*
Phytologist **102**: 499–512.

Stitt M, Huber S, Kerr P. **1987**. Control of Photosynthetic Sucrose Formation. In:
Photosynthesis. Elsevier, 327–409.

Stitt M, Wirtz W, Heldt HW. **1983**. Regulation of Sucrose Synthesis by Cytoplasmic
Fructosebisphosphatase and Sucrose Phosphate Synthase during Photosynthesis in Varying
Light and Carbon Dioxide. *Plant physiology* **72**: 767–74.

Sun L, Song L, Zhang Y, Zheng Z, Liu D. **2016**. Arabidopsis PHL2 and PHR1 Act
Redundantly as the Key Components of the Central Regulatory System Controlling
Transcriptional Responses to Phosphate Starvation. *Plant physiology* **170**: 499–514.

Suzuki N, Devireddy AR, Inupakutika MA, Baxter A, Miller G, Song L, Shulaev E, Azad
RK, Shulaev V, Mittler R. **2015**. Ultra-fast alterations in mRNA levels uncover multiple
players in light stress acclimation in plants. *The Plant journal : for cell and molecular biology*
84: 760–72.

Thibaud M-C, Arrighi J-F, Bayle V, Chiarenza S, Creff A, Bustos R, Paz-Ares J, Poirier
Y, Nussaume L. **2010**. Dissection of local and systemic transcriptional responses to phosphate
starvation in Arabidopsis. *The Plant journal : for cell and molecular biology* **64**: 775–89.

Vandenbussche F, Habricot Y, Condiff AS, Maldiney R, Van Der Straeten D, Ahmad M.
2007. HY5 is a point of convergence between cryptochrome and cytokinin signalling pathways
in Arabidopsis thaliana. *Plant Journal* **49**: 428–441.

Verma L, Pandey M, Bhatia C, Mehra P, Singh B, Giri J. **2025**. Phosphate deficiency
inducible OsGDPD5 affects root growth by regulating sugar-auxin crosstalk. *The Plant*
journal : for cell and molecular biology **121**: e17249.

Vogel MO, Moore M, König K, Pecher P, Alsharafa K, Lee J, Dietz KJ. **2014**. Fast
retrograde signaling in response to high light involves metabolite export, MITOGEN-
ACTIVATED PROTEIN KINASE6, and AP2/ERF transcription factors in Arabidopsis. *Plant*
Cell **26**: 1151–1165.

Vu HS, Roth MR, Tamura P, Samarakoon T, Shiva S, Honey S, Lowe K, Schmelz EA,
Williams TD, Welti R. **2014**. Head-group acylation of monogalactosyldiacylglycerol is a

common stress response, and the acyl-galactose acyl composition varies with the plant species and applied stress. *Physiologia Plantarum* **150**: 517–528.

Walker DA, Osmond CB. **1986**. Measurement of photosynthesis in vivo with a leaf disc electrode: correlations between light dependence of steady-state photosynthetic O 2 evolution and chlorophyll a fluorescence transients. *Proceedings of the Royal Society of London. Series B. Biological Sciences* **227**: 267–280.

Walker DA, Sivak MN. **1986**. Photosynthesis and phosphate: a cellular affair? *Trends in Biochemical Sciences* **11**: 176–179.

Wang Z, Zheng Z, Zhu Y, Kong S, Liu D. **2023**. PHOSPHATE RESPONSE 1 family members act distinctly to regulate transcriptional responses to phosphate starvation. *Plant Physiology* **191**: 1324–1343.

Ward JT, Lahner B, Yakubova E, Salt DE, Raghothama KG. **2008**. The effect of iron on the primary root elongation of Arabidopsis during phosphate deficiency. *Plant physiology* **147**: 1181–91.

Weber H. **2002**. Fatty acid-derived signals in plants. *Trends in plant science* **7**: 217–24.

Weise SE, Liu T, Childs KL, Preiser AL, Katulski HM, Perrin-Porzondek C, Sharkey TD. **2019**. Transcriptional Regulation of the Glucose-6-Phosphate/Phosphate Translocator 2 Is Related to Carbon Exchange Across the Chloroplast Envelope. *Frontiers in plant science* **10**: 827.

Wild R, Gerasimaite R, Jung JY, Truffault V, Pavlovic I, Schmidt A, Saiardi A, Jacob Jessen H, Poirier Y, Hothorn M, *et al.* **2016**. Control of eukaryotic phosphate homeostasis by inositol polyphosphate sensor domains. *Science* **352**: 986–990.

Wojciechowska I, Mukherjee T, Knox-Brown P, Hu X, Khosla A, Subedi B, Ahmad B, Mathews GL, Panagakis AA, Thompson KA, *et al.* **2024**. Arabidopsis PROTODERMAL FACTOR2 binds lysophosphatidylcholines and transcriptionally regulates phospholipid metabolism. *New Phytologist*.

Woodrow IE, Raymond Ellis J, Jellings A, Foyer CH. **1984**. Compartmentation and fluxes of inorganic phosphate in photosynthetic cells. *Planta* **161**: 525–30.

Wu P, Ma L, Hou X, Wang M, Wu Y, Liu F, Deng XW. **2003**. Phosphate starvation triggers distinct alterations of genome expression in arabidopsis roots and leaves. *Plant Physiology* **132**: 1260–1271.

Wu FH, Shen SC, Lee LY, Lee SH, Chan MT, Lin CS. **2009**. Tape-arabidopsis sandwich -

A simpler arabidopsis protoplast isolation method. *Plant Methods* **5**: 1–10.

Xu L, Zhao H, Wan R, Liu Y, Xu Z, Tian W, Ruan W, Wang F, Deng M, Wang J, *et al.*

2019. Identification of vacuolar phosphate efflux transporters in land plants. *Nature Plants* **5**:

84–94.

Yang Y, Benning C. **2018**. Functions of triacylglycerols during plant development and stress.

Current Opinion in Biotechnology **49**: 191–198.

Yang B, Peng Y, Zhang G, Liu R, Hao S, Ren Y, Lu S, Wang X, Guo L. **2025**. Regulation

of glucosylceramide synthase and sphingolipid remodeling in the plant response to phosphate

deficiency. *The Plant cell* **37**.

Zhou J, Jiao F, Wu Z, Li Y, Wang X, He X, Zhong W, Wu P. **2008**. OsPHR2 is involved in

phosphate-starvation signaling and excessive phosphate accumulation in shoots of plants. *Plant*

physiology **146**: 1673–86.

Zhu J, Lau K, Puschmann R, Harmel RK, Zhang Y, Pries V, Gaugler P, Broger L, Dutta

AK, Jessen HJ, *et al.* **2019**. Two bifunctional inositol pyrophosphate kinases/phosphatases

control plant phosphate homeostasis. *eLife* **8**: 1–25.

Zirngibl M-E, Araguirang GE, Kitashova A, Jahnke K, Rolka T, Kühn C, Nägele T,

Richter AS. **2023**. Triose phosphate export from chloroplasts and cellular sugar content

regulate anthocyanin biosynthesis during high light acclimation. *Plant communications* **4**:

100423.

1223

Figures

Figure 1 Mutants with defects in photoassimilate partitioning show altered P_i responses. **A,**
Contents of glucose, fructose, sucrose, and starch in shoots upon P_i depletion relative to fresh
weight (**FW**). Seedlings of WT (Col-0), *adg1-1*, *tpt-2*, *phr1-3 phl1*, *adg1 tpt-2* and
adg1 phr1-3 phl1 genotype were grown for 10 days on rich medium containing 0.5 % sucrose
before transfer to media with 0.5 % sucrose and either 2.5 mM (**Control**) or 0 mM (**P_i**
depletion) KH_2PO_4 added. Growth was continued for 7 days. Pools of seedling shoots were
harvested 10.25 h after onset of the 16-h photoperiod. Means ± standard deviations; $n = 3$
independent experiments. Letter code indicates statistical differences at $P < 0.05$. One-way
ANOVA with Tukey HSD follow-up test and Bonferroni alpha correction for contrasts. **B, C,**
Phenotype of the *adg1 phr1 phl1* triple mutant. Plants were grown at 23°C and a light intensity
of 100±10 µmol m^{-2} s^{-1}. **B,** Rosette habitus of plants grown under short-day (8 h light period)
and long-day (16 h light period) conditions. Pictures were taken of representative plants after
27 (short day) and 25 (long day) days of growth. Bar, 2.5 cm. **C,** Leaves of 42 day-old plants
grown under an 8-h light regime were aligned according to age (starting with the cotyledons on
the left hand side). Background was manually removed for better visualization. **D, E,**
Constitutive P_i-starvation transcription in rosette leaves of *adg1 tpt-2*. Plants were grown on
soil at a photon flux density of 90±10 µmol m^{-2} s^{-1} and sampled 135 min after onset of the 8-h
photoperiod after 5 weeks of growth. **D,** P_i contents of rosette leaves. Means ± standard
deviations are depicted. $n = 3$ independent experiments. Kruskal-Wallis test with DUNN's
follow-up test and DUNN/Sidak alpha correction for contrasts; $P < 0.01$. **E,** qRT-PCR analysis
of P_i starvation responsive genes in rosette leaves of WT, *adg1-1*, *tpt-2*, and *adg1 tpt-2*.
Transcript levels are $1000 \cdot 2^{-\Delta CT}$ relative to *PP2A*. Bars show means ± standard deviations. $n = $
3 independent experiments. One-way ANOVA with Tukey HSD follow-up test and Bonferroni
alpha correction for contrasts; $P < 0.01$. **F**, High-light sensitivity of *adg1-1*, *phr1-3 phl1* and
derived mutants. Seedlings were grown for two weeks in a 16/8 h (light/dark) cycle under low
light (60±5 µmol m^{-2} s^{-1}) or high light (350±30 µmol m^{-2} s^{-1}) on agar-solidified media
containing 0.25 mM P_i, 2.5 mM P_i, or 2.5 mM P_i +50 mM sucrose. Shoot FW were calculated
as percent of mean shoot FW of WT seedlings grown under the respective conditions. Bars
represent means ± standard deviations; $n = 3$ independent experiments comprising means of
each 4-8 seedling shoots. Statistical analyses were performed using Student's *t* test with 2-tailed
distribution, unpaired with unequal variance. Significant differences between data are indicated
as ***$P < 0.001$, **$P < 0.01$, *$P < 0.05$, n.s., not significant.

1258

1259

Figure 2 High light triggers P_i-depletion responsive gene expression. **A**, Overlap of differential

gene expression in shoots upon high light, P_i starvation, and nitrogen starvation. Venn diagrams

show numbers of transcripts upregulated by high light or P_i starvation, and of genes

differentially expressed under N starvation. Transcript data are taken from Huang et al. (2019)

(high light), Bustos et al. (2010) (P_i depletion), and Krapp et al. (2011) (N depletion). **B**, qRT-

PCR analyses of selected transcripts in rosette leaves from WT (Col-0) and *phr1-1 phl1*

mutants. Plants were grown on soil at 23°C for 5 weeks in an 8-h light regime at

70±5 µmol m^{-2} s^{-1} before light intensity was set to 450±30 µmol m^{-2} s^{-1} (High light, yellow bars)

starting 1 h after onset of the photoperiod on day 36. The control group was kept at 70±5 µmol

m^{-2} s^{-1} (blue bars). Each 2-3 fully developed leaves (leaves no. 10-13) were harvested per sample

at the indicated treatment times. Transcript levels were calculated relative to *PP2A* as

$1000 \cdot 2^{-\Delta CT}$. Bars represent means ± standard deviations, *n* = 3 independent experiments.

Statistical analyses were performed using Student's *t* test with 2-tailed distribution, unpaired

with unequal variance. Significant differences are indicated as ***P <0.001, **P <0.01, *P

<0.05, n.s., not significant.

1275

1276

Figure 3 High light affects cellular P_i pools and sucrose levels. **A-C,** Plants were grown and treated as described in Fig. 2B. **A,** Subcellular P_i concentrations in WT and *phr1-1 phl1* mutant plants upon high light relative to dry weight (**DW**). Rosettes were harvested after 20 min of treatment. P_i contents were determined following non-aqueous fractionation of cellular compartments. *n* = 4 (*phr1-1 phl1*) or 5 (Col-0) independent experiments. **B,** Changes in ATP/ADP levels after shift to high light relative to fresh weight (**FW**). *n* = 6 plants from 3 (90 min) or 4 (20 min) independent experiments. **C,** Sucrose levels upon 20 min of high light; *n* = 8 plants from 4 independent experiments; absolute values and fold changes (**FC**, high light/control) are shown. Significant difference in FC was tested using Student's *t* test with 2-tailed distribution, unpaired with unequal variance. **D,** High light affects total shoot P_i and

growth rates on the medium-term. WT and *phr1-1 phl1* mutants were grown for 4 weeks as

described for Fig. 2B before light intensity was shifted to 450 ± 30 µmol m^{-2} s^{-1} (high light) at

4 h after onset of the photoperiod. Control plants were kept under growth light conditions (70 ± 5

µmol m^{-2} s^{-1}). Material was harvested after 24 and 48 h. Total P$_i$ contents in rosettes relative to

DW and average rosette DWs are depicted; $n = 10$ plants from 5 individual experiments. All

bars show means ± standard deviations. Except for FC shown in **C**, all statistical analysis were

performed using 2-factor ANOVA with Tukey HSD post-hoc test; $P < 0.05$; n.s., not significant.

1294

1295

Figure 4 *phr1-1 phl1* mutants show altered carbohydrate responses. Unless specified, plants were grown and treated as described for Fig. 2B. Data points and bars depict means ± standard deviations. **A, B**, PAM measurements were performed to determine the effect of high light (**HL**) on chlorophyll a fluorescence parameters. $n = 3$ independent experiments (representing means of each 3 leaves measured). **A,** Quantum yield of photosystem II photochemistry (**ΦPSII**) in light-adapted leaves after 20, 45, and 125 min of treatment. Significance was tested using One-

way ANOVA with Tukey HSD follow-up test and Bonferroni alpha correction for contrasts;

n.s., not significant. **B,** Plants subjected to high-light stress were dark-adapted for 20 min after

125 min of stress duration to determine the maximum efficiency of photosystem II (F_v/F_m). 2-

factor ANOVA with Tukey HSD post-hoc test; $P < 0.05$. **C, E,** Changes in carbohydrate levels

in WT (Col-0) and *phr1-1 phl1* mutants after 90 min of high-light treatment. 2-factor ANOVA

with Tukey HSD post-hoc test; $P < 0.05$. Fold changes (**FC**) of sugar/starch contents were

calculated for each genotype as HL/mean of control. Statistical analyses of FC were performed

with Student's *t* test with two-tailed distribution, unpaired with unequal variance. Significant

differences are indicated as ** $P < 0.01$, n.s. not significant. **C,** Starch levels relative to fresh

weight (**FW**); *n* = 9 plants from 3 independent experiments. **D,** Anthocyanin (cyanidin 3-

glucoside) levels after 3 days of high-light exposure. Plants were grown for 32 days until onset

of the treatment. Leaf material was harvested after additional 70 h. Pigment contents were

calculated relative to FW. *n* = 3 independent experiments; statistical analyses were conducted

for each treatment group using One-way ANOVA with Tukey HSD follow-up test and

Bonferroni alpha correction for contrasts, $P < 0.01$. **E,** Levels of glucose, fructose, and sucrose

after 90 min of high-light treatment. *n* = 11-12 plants from 4 independent experiments. **F,**

phr1-1 phl1 mutants are defective in sugar-induced anthocyanin production under P_i-limiting

conditions. Seedlings were grown on nutrient-rich media (no sucrose or sorbitol added) for 7

days under a 16-h light regime before transfer to media with either 0 or 2.5 mM KH_2PO_4 added

and containing either no additives (control), 90 mM sucrose, or 160.7 mM sorbitol (same

osmotic strength). Seedling shoots were harvested for anthocyanin determination after 67 h of

growth on differing media. *n* = 6 pools of seedling shoots from 3 independent experiments; 2-

factor ANOVA with Tukey post-hoc test; $P < 0.05$.

1325

1326

1327

Figure 5 Effect of *srg3* mutation on the lipid composition upon light increase. **A,** Promoter

transactivation assay in mesophyll protoplasts showing that the promoters of *SPX1*, *MGD3*, and

SRG3 are activated by PHR1. Promoter sequences were fused to the firefly *LUC* reporter gene

(*proSRG3::LUC_Firefly*; *proSPX1^GC^::LUC_Firefly; proMGD3::LUC_Firefly*, *proGPT2::LUC_Firefly*) and

co-transfected with either *PHR1* or *GFP*. Luminescence units were normalized to the

luminescence of *Renilla LUC* which was additionally co-expressed from the Cauliflower

mosaic virus *35S* promoter, and calculated relative to the luminescence level of GFP-expressing

samples. **RLU**, relative luminescence level. Results for each promoter construct are from

independently triplicated experiments. Means ± standard deviations; $n = 6$ from 3 independent

experiments. One-way ANOVA with Tukey HSD follow-up test and Bonferroni alpha

correction for contrasts; ***$P < 0.001$, **$P < 0.01$. **B-E**, Plants of WT (Col-0) and *srg3*

genotype were grown and treated as described for Fig. 2B, but leaves were harvested at 2 and

7 h of treatment. Lipid species were analyzed using LC-MS. **B**, Principal component analysis

of samples from 2 h and 7 h high-light (**HL**) treated or control plants of WT and *srg3* genotype.

C, Levels of *srg3*-dependent HL-responsive lipid features normalized to leaf fresh weight

(**FW**). $n = 4$ independent experiments, except for acyl-MGDG: $n = 3$-4. **D, E,** $n = 3$-4. **D,**

Contents of triacylglycerols (**TAG**) in WT and *srg3* leaves after high-light treatments. **E**,

Changes in TAG pools upon 7 h of light increase (7 h HL vs. 7 h control) in the WT. Fold

changes (Col-0_HL_7h/Col-0_C_7h) are indicated by color code and were calculated for TAG

species sorted by the total number of carbon atoms of the acyl chains (**C**) and the total number

of double bonds (**DB**). **nd**, not detected. **F**, Light response of the NPQ parameter measured by

PAM chlorophyll fluorometry. Plants of WT (Col-0), *srg3*, and *phr1-1 phl1* genotype were

grown for 40 days in an 8-h light regime at 70 ± 5 µmol m^{-2} s^{-1} and dark-adapted for ≥30 min

before measurement. NPQ was plotted against the fluence rate of photosynthetically active

radiation (**PAR**). $n = 3$ independent experiments representing means of 2 plants. **C-F**, 2-factor

ANOVA with Tukey HSD post-hoc test; $P < 0.05$.

1354

1355

Figure 6 Model of P_i signaling under light increase. **Left**, Under normal growth light conditions, production of triose phosphates in the Calvin Benson cycle and their utilization in downstream processes such as starch and sucrose biosynthesis are balanced, and the rate of P_i recycling matches its consumption by the chloroplast metabolism (Stitt *et al.*, 1987; McClain & Sharkey, 2019). **Right**, Upon shift to higher fluence rates, phosphorylated photosynthetic intermediates accumulate (Borghi *et al.*, 2019), resulting in a decrease of chloroplast P_i concentrations. Chloroplast P_i levels or changes in the pools of photosynthetic products generate a signal that leads to PHR1(-like) activation. This causes P_i-starvation responsive gene expression including the induction of *SRG3* and *VPE1*, leading to P_i release from intracellular pools, and changes in lipid metabolism enabling the containment of free fatty acids and xanthophyll cycle operation, as well as anthocyanin accumulation in the medium-term. **TP**, triose phosphate; **F6P**, Fructose 6-phosphate; **G1P**, Glucose 1-phosphate.

1368