PHR1 and PHL1 mediate rapid high-light responses and acclimation to triose phosphate oversupply

Lukas Ackermann[1], Monika Müller[1], Alina Johanna Hieber[1], Maria Klecker[1*]

[1] Plant Physiology, University of Bayreuth, 95440 Bayreuth, Germany

[*]Address correspondence to: Maria Klecker, Department of Plant Physiology, University of Bayreuth, Universitätsstrasse 30, D-95447 Bayreuth, Germany. e-mail: maria.klecker@uni-bayreuth.de

Short title: PHR1 and PHL1 signal photosynthetic stress

ORCID IDs

Maria Klecker   0000-0001-5508-2363

Lukas Ackermann   0009-0009-8604-6029

ABSTRACT

Fluctuations in light intensity require immediate metabolic adjustment which includes reprogramming of both plastidial and nuclear gene expression, but the signaling pathways behind such responses are not fully understood. Here we report the identification of an early high-light responsive pathway in *Arabidopsis thaliana* that depends on PHOSPHATE STARVATION RESPONSE 1 (PHR1) and PHR1-LIKE 1 (PHL1) transcription factors involved in low phosphate (P_i) signaling. High-light treatment rapidly induced the accumulation of PHR1-responsive transcripts in wildtype plants grown under nutrient-sufficient conditions, but not in *phr1 phl1* double knockout plants. Differences in starch accumulation and ATP levels were detected between wildtype and *phr1 phl1* mutants subjected to high light, suggesting a link between P_i signaling, carbohydrate partitioning, and energy status during stress. In line with a function of PHR1/PHL1 upon triose phosphate accumulation, we observed that blocking starch biosynthesis in the *phr1 phl1* double mutant, by introducing the *agd1-1* allele, causes a severe growth defect. Phenotypes of the *adg1 phr1 phl1* triple mutant such as high-light sensitivity and growth restriction in the absence of exogenously supplied sucrose resemble the previously described double mutant *adg1 tpt-2*, lacking a functional copy of the TRIOSE PHOSPHATE/PHOSPHATE TRANSLOCATOR (TPT), and we show that P_i responses are disturbed in *adg1 tpt-2*. We propose that P_i sequestration by photosynthesis and import of P_i into the chloroplast transiently depletes cytosolic P_i reserves upon sudden increases in light intensity. The low-P_i sensing machinery in the nucleus consequently implements early high-light transcriptional responses, qualifying P_i as a new operational retrograde signal.

Introduction

Light as a source of energy with highly fluctuating availability needs to be efficiently captured by the photoautotrophic metabolism. This necessitates an enzymatic machinery capable of coping with rapid changes in metabolic flux, as well as the containment of overexcitation-induced hazards. Both challenges are met not only by post-transcriptional regulation of enzyme activity (reviewed in König et al. (2012), Matiolli et al. (2022)), but also by pronounced changes in mRNA abundance that occur within 30 minutes upon an increase in irradiance (Vogel et al., 2014; Suzuki et al., 2015; Huang et al., 2019). The latter mechanism depends on fast signal transmission between the chloroplast and the nuclear transcription machinery. Such communication is mediated by 'primary' retrograde signals (Dietz, 2015),

including redox cues and reactive oxygen species like H_2O_2 and singlet oxygen which are
produced during photosynthetic reactions (Bechtold et al., 2008; Dietz et al., 2016).
Additionally, increased electron transport activity and light-induced Calvin cycle activation
promote the accumulation of primary products of carbon fixation (Dietz and Heber, 1984; Dietz
and Heber, 1986) which may themselves act as signaling molecules (Häusler et al., 2014; Moore
et al., 2014). Hence, induction of high-light specific gene expression partially depends on the
triose phosphate/phosphate translocator (TPT) that mediates the exchange of triose phosphates
or 3-phosphoglycerate (3-PGA) with inorganic phosphate (P_i) at the chloroplast envelope
(Schneider et al., 2002; Vogel et al., 2014; Weise et al., 2019). A central function of the TPT in
light acclimation and retrograde signaling was corroborated by the severe high-light dependent
phenotypes of double mutants defective in the *TPT* and *ADG1* (for *ADP GLUCOSE*
PYROPHOSPHORYLASE 1), encoding a small subunit of the ADP-glucose pyrophosphorylase
complex (AGPase) which is required for the biosynthesis of transitory starch (Schmitz et al.,
2012). Phenotypes of *adg1-1 tpt* double mutants include growth retardation, low diurnal
changes in carbohydrate levels, and high chlorophyll fluorescence. These observations point
towards an important role of photosynthate partitioning for retrograde signaling, however the
identity of the primary sensors as well as signal transduction mechanisms are still unclear.

Phosphorus makes up about 2-30 permille of the plant dry mass (Kumar et al., 2019), as it is
incorporated into a large subset of biomolecules. Systemic signaling of P_i availability and P_i
distribution within the plant body mainly depend on a conserved gene family encoding
PHOSPHATE STARVATION RESPONSE 1 (PHR1) and PHR1 LIKE (PHL) transcription
factors of the MYB coiled-coiled type (Rubio et al., 2001; Zhou et al., 2008; Bustos et al., 2010;
Thibaud et al., 2010; Sun et al., 2016). Simultaneous loss of *PHR1* and *PHL1* function affects
the expression of at least 68 percent of the P_i starvation responsive genes in the shoots of
Arabidopsis (*Arabidopsis thaliana)* (Bustos et al., 2010). A decrease in cellular P_i concentration
impairs production of the inositol pyrophosphate species InsP8, relieving PHR1/PHLs of
suppression by SPX (for SYG1/Pho81/XPR1) domain-containing proteins, thus linking
transcription factor activity to P_i availability (Puga et al., 2014; Dong et al., 2019; Zhu et al.,
2019; Ried et al., 2021). The transcriptional response to P_i scarcity leads to scavenging of
external and internal P_i sources through phosphatase induction (Morcuende et al., 2007) and
altered lipid metabolism (Misson et al., 2005; Pant et al., 2015b), anthocyanin biosynthesis
(Nilsson et al., 2012; Liu et al., 2022; Li et al., 2023), and protection from photodamage
(Nilsson et al., 2012), to name a selection. PHR1/PHLs are characterized as transcriptional
activators (Nilsson et al., 2007; Bustos et al., 2010). Accordingly, direct targets of PHR1

(Bustos et al., 2010; Castrillo et al., 2017) are strongly upregulated upon P_i starvation and

include genes such as *PHOSPHATE STARVATION-INDUCED GENE 2* (*PS2*; AT1G73010)

(Hanchi et al., 2018), *MONOGALACTOSYLDIACYLGLYCEROL SYNTHASE 3* (*MGD3*;

AT2G11810)　(Kobayashi　et　al.,　2004),　and　*GLYCEROPHOSPHODIESTER*

PHOSPHODIESTERASE 1/SENESCENCE-RELATED GENE 3 (*SRG3*; AT3G02040) (Cheng

et al., 2011). Nevertheless, a subset of transcripts is specifically repressed under P_i-depleted

conditions which can be accounted for by the induction of microRNAs (reviewed in Paz-Ares

et al. (2022)), the induction of transcriptional repressors, as well as the modulation of genome

accessibility (Barragán-Rosillo et al., 2021) through PHR1/PHLs activity.

P_i is unique among the macronutrients in that it is the only root-supplied element which is

directly consumed by photosynthesis (Heldt and Rapley, 1970; Dietz and Heber, 1984). It has

been shown experimentally that under high-CO_2 and high-light conditions, photosynthesis can

indeed become limited by P_i supply to the chloroplast, both under P_i depleted and nutrient-rich

conditions (Dietz and Foyer, 1986; Sivak and Walker, 1986). However, whether rapid increase

in photosynthetic activity affects P_i homeostasis of the cell, is still unclear. In this study, we

show that increments in light intensity trigger an early nuclear transcriptional response

characteristic of P_i depletion which depends on *PHR1* and *PHL1*. Processes regulated by

PHR1/PHL1 upon increased illumination contribute to photosynthetic acclimation since

phr1 phl1 mutants showed metabolic alterations compared to the wildtype when subjected to

high-light stress. Moreover, analysis of the starch biosynthesis-deficient *adg1 phr1 phl1* triple

mutant revealed an important function of P_i signaling when triose phosphate utilization is

impaired. Aspects of the growth defects observed for *adg1 phr1 phl1* phenocopy the *adg1 tpt-2*

mutant, suggesting that triose phosphate partitioning strongly affects P_i homeostasis. Together,

we propose that PHR1/PHL1 mediate rapid photosynthetic acclimation upon sudden increases

in light intensity by using P_i as a previously unknown operational retrograde signal.

Results

Acclimation to P_i starvation is disturbed in *adg1 tpt-2*

Photoassimilate allocation and utilization in the chloroplast and cytosol mainly determine the rate of P_i recycling from photosynthetic products. To address how the partitioning of phosphorylated assimilates affects P_i homeostasis, we examined the phenotypes of mutants defective in starch biosynthesis (*adg1-1*) and chloroplastic triose phosphate export (*tpt-2*) during P_i starvation in the presence of 14.61 mM (0.5 %) sucrose, and used anthocyanin production as a read-out for the induction of the P_i starvation response. In accordance with previous reports (Nilsson et al., 2007), wildtype seedlings accumulated anthocyanins when transferred to P_i-deficient conditions (Figure 1A, B). This was not observed for the *phr1-3 phl1* mutant. While *tpt-2* mutant seedlings were indistinguishable from the wildtype in terms of anthocyanin content, pigmentation was significantly enhanced upon P_i starvation in *adg1-1*. Surprisingly, this effect was fully reversed in the double mutant *adg1 tpt-2* which even accumulated much lower amounts of anthocyanins than the wildtype (Figure 1B). Nevertheless, root responses to P_i depletion such as root hair production were normal in *adg1 tpt-2* (Figure 1A). Hence, increased anthocyanin production of *adg1-1* upon P_i starvation almost fully relied on photoassimilate transport over the chloroplast envelope. To test whether increased anthocyanin accumulation in *adg1-1* also depends on the signaling pathway executed by PHR1/PHLs, we created a triple mutant by crossing *adg1-1* and *phr1-3 phl1*. The anthocyanin content did not increase in the *adg1 phr1 phl1* triple mutant in response to P_i starvation (Figure 1B), confirming a crucial function of PHR1/PHL1 signaling in this process even in the absence of ADG1 activity.

P_i deprivation is also known to induce starch accumulation in a *PHR1*-dependent manner (Figure 1C and Supplemental Figure 1), which likely involves AGPase activity (Nilsson et al., 2007). Furthermore, also soluble sugar contents rise in wildtype plants upon P_i limitation (Pant et al., 2015a). We reasoned that contrasting pigmentation behavior of the *adg1-1* and *adg1 tpt-2* seedlings under P_i-depleted conditions might be caused by differential sugar accumulation (Zirngibl et al., 2023). First, we analyzed starch contents of seedling shoots under P_i-sufficient and -deficient conditions. While shoots of *tpt-2* showed slightly higher levels of starch than wildtype, P_i depletion did not result in considerable starch production in *adg1-1* or any of the derived mutant lines (Figure 1C), confirming that the small AGPase subunit ADG1 is required for increased starch accumulation under P_i starvation. In agreement with impaired starch biosynthesis, hexose levels were increased in *adg1-1* shoots under P_i depletion compared to the

wildtype, and we observed a tendency towards higher sucrose levels in *adg1-1* (Figure 1C).

None of these effects were seen in the *adg1 phr1 phl1* triple mutant, suggesting that sugar

metabolism in *adg1-1* was under control of PHR1/PHL1 during P_i depletion (Figure 1C).

Surprisingly, sugar levels were not correlated to anthocyanin production in either *tpt-2* or *adg1*

tpt-2: While P_i-depleted shoots of *tpt-2* contained higher glucose levels than wildtype, the

contents of all three soluble sugar species analyzed were similar between shoots of *adg1 tpt-2*

and the wildtype when grown on sucrose-supplemented media (Fig. 1C). Thus, anthocyanin

production in response to P_i starvation was partially uncoupled from cellular sugar contents.

Taken together, PHR1/PHL1 activity is required, but not sufficient for anthocyanin production

under P_i starvation, which furthermore depends on adequate metabolite partitioning between

chloroplasts and the cytosol. Furthermore, the phenotypes of the *adg1 tpt-2* double mutant

indicate that P_i starvation responses are disturbed in this mutant.

P_i starvation gene expression is triggered by high light

The low-P_i phenotypes of *adg1-1* and *adg1 tpt-2* mutant lines indicated that P_i starvation

responses are affected by photoassimilate distribution and utilization. This raised the question

whether photoassimilate turnover also impacts on P_i homeostasis under non-starved conditions.

More specifically, P_i signaling might be modulated by naturally occurring changes in triose

phosphate metabolism, for example upon an increase in light intensity. To address this, we

tested low-P_i marker gene expression upon sudden increments in photon flux density. Using

quantitative RT-PCR (qRT-PCR), we first confirmed that in our experimental setup transcript

levels of *SRG3*, *PS2*, *SPX1*, and *MGD3* were all strongly induced by P_i starvation, which mainly

depended on *PHR1/PHL1* (Supplemental Figure 2A). Additionally, we analyzed transcript

levels of a putative vacuolar P_i efflux transporter, *VPE1* (AT3G47420) (Ramaiah et al., 2011;

Xu et al., 2019), and confirmed upregulation of this gene in the wildtype subjected to P_i

starvation which was again strongly attenuated in *phr1-1 phl1* mutants (Supplemental Figure

2A). Furthermore, for the promoter sequences of *SPX1* (lacking the naturally occurring *Nco*I

restriction site, *proSPX1^{GC}*), *MGD3*, and *SRG3*, we observed that PHR1 was able to induce the

expression of *promoter::LUC* fusion constructs when transiently expressed in mesophyll

protoplasts (Supplemental Figure 2B), confirming that *SPX1*, *MGD3*, and *SRG3* are targets of

PHR1.

To examine responses to enhanced illumination, we grew wildtype and *phr1-1 phl1*

mutant plants under nutrient-rich conditions on soil in an 8/16 hours (light/dark) photoregime.

After 5 weeks, photon flux density was increased to 320±30 µmol m^{-2} s^{-1} (high light) while the

control group was kept under growth light conditions of 80±5 µmol m^{-2} s^{-1}. Using qRT-PCR,

we detected a pronounced upregulation of *SRG3*, *SPX1*, and *VPE1* transcripts in wildtype plants

already after 20 min of exposure to high light compared to control plants, as well as a slight

increase in *MGD3* and *PS2* transcript abundance (Figure 2). In all five cases, the upregulation

of gene expression was attenuated within 125 min in high light (Figure 2), indicating that

response to P$_i$ shortage constitutes an early consequence of increased irradiance. Notably,

induction of P$_i$ starvation marker transcripts was not observed in *phr1-1 phl1* mutant plants

(Figure 2). *PHR1* expression itself was reported to depend on light (Liu et al., 2017).

Importantly however, increasing the light intensity to 320±30 µmol m^{-2} s^{-1} did not affect *PHR1*

transcript levels under our experimental conditions (Supplemental Figure 3), consistent with

the accumulation of *PHR1* transcript being only responsive to changes in the very low fluence

range (Liu et al., 2017).

GLUCOSE-6-PHOSPHATE/PHOSPHATE TRANSLOCATOR 2 (*GPT2*) is a light

responsive gene (Athanasiou et al., 2010) which was also described to be induced by P$_i$

starvation (Morcuende et al., 2007). Furthermore, high-light induced expression of *GPT2* was

previously described to depend on *TPT* function (Kunz et al., 2010; Weise et al., 2019), and we

noted that the promotor sequence of *GPT2* contains a PHR1 binding motif (GTATATTC) close

to the transcriptional start site. Therefore, we tested if the *GPT2* promotor could be activated

by PHR1 in a transient expression system using the *LUC* reporter gene. In fact, *PHR1*

expression induced LUC reporter activity when protoplasts were co-transfected with the

proGPT2::LUC_Firefly construct (Supplemental Figure 2B). However, the responsiveness of

proGPT2 to *PHR1* expression was considerably lower than observed with the constructs

harboring *proSPX1^{GC}, proSRG3* or *proMGD3* upstream of the reporter CDS. Consistent with

weak PHR1-mediated induction of *proGPT2* in protoplast assays, and in contrast to the other

tested P$_i$-starvation responsive genes, *GPT2* was upregulated by high light in both wildtype and

phr1-1 phl1 after 45 minutes, and transcript levels continued to rise upon stress during the

course of the analysis (Figure 2). This indicated that PHR1/PHL1 are not involved in high-light

mediated induction of *GPT2* expression. Moreover, we were not able to detect an effect of the

gpt2-1 allele (Niewiadomski et al., 2005) on carbohydrate accumulation under P$_i$ depletion

(Supplemental Figure 1), arguing against a major role of this gene for sugar homeostasis under

P$_i$ starvation. Thus, metabolite transport over the chloroplast membrane likely creates another

signal which is responsible for *GPT2* induction under high light. While this pathway is not

affected by *phr1-1 phl1* mutation, the double mutant fails to induce other P_i-depletion
responsive genes which are upregulated in the wildtype very early after a shift in light intensity.

Metabolic responses to increased light are impaired in *phr1 phl1* mutants

Given that transcription of P_i-starvation responsive genes was rapidly induced by high
light, we asked if this response was of physiological significance during acclimation to
increased radiation. To investigate this, we determined the changes in adenylates, sugar, and
starch contents that occurred within 90 min upon an increase in photon flux density from
$80\pm5\ \mu\mathrm{mol\ m^{-2}\ s^{-1}}$ to $320\pm30\ \mu\mathrm{mol\ m^{-2}\ s^{-1}}$ in rosette leaves of 5-weeks-old wildtype and
phr1-1 phl1 mutant plants. As shown in Figure 3A, increasing the light intensity caused a slight
drop in ATP levels in leaves of *phr1-1 phl1* mutants, while no significant changes were seen in
wildtype plants. Notably, the ratio of ATP/ADP was not affected by either the genotype or the
high-light treatment (Figure 3A). In contrast, fructose, glucose, and sucrose levels increased in
leaves of both wildtype and *phr1-1 phl1* mutants subjected to high light as compared to control
light conditions (Figure 3B). Interestingly, total levels of glucose and fructose were slightly
lower in *phr1-1 phl1* leaves subjected to high light compared to wildtype leaves (Figure 3B).
However, the relative increments in sugar levels upon stress did not differ significantly between
wildtype and *phr1-1 phl1* mutants (Figure 3C).

Unlike sugar levels, starch content under control conditions was higher in *phr1-1 phl1*
mutants than in wildtype plants (Figure 3D). This may be related to the fact that *phr1-1 phl1*
mutants manifest lower P_i levels in the shoots (Wang et al., 2018a) which allosterically activates
AGPase (Figueroa et al., 2022). Exposure to higher light intensities further increased starch
content in both wildtype and *phr1-1 phl1* (Figure 3D). However, while the starch content of
wildtype leaves increased by 1.98-fold relative to control conditions, *phr1-1 phl1* mutant plants
exhibited a significantly lower increase by only 1.62-fold (Figure 3D). Thus, sequestration of
photoassimilates in transitory starch under high light conditions was affected by loss of
PHR1/PHL1 function.

Since both P_i starvation and high-light stress trigger anthocyanin production, we next
analyzed the ability of *phr1 phl1* mutants to accumulate anthocyanins during high-light
acclimation. As shown in Figure 3E, anthocyanin contents of *phr1 phl1* leaves after 3 days of
high-light exposure were significantly lower than in wildtype leaves or in the single mutants
phr1-3 and *phl1*, while loss of *SPX1/2* function did not have any statistically significant effect.
To exclude that *phr1 phl1* mutant plants were generally impaired in anthocyanin biosynthesis,

we subjected seedlings to chemically induced oxidative stress. Here, anthocyanin contents were
not significantly different between wildtype, *phr1-3 phl1*, and *spx1 spx2* seedlings after 5 days
of growth on paraquat-containing media (Supplemental Figure 4), indicating that anthocyanin
biosynthesis was functional during reactive oxygen stress in *phr1-3 phl1* mutants.

Next, we asked whether metabolic abnormalities in the *phr1 phl1* mutants were related
to differences in photosynthetic performance at the level of the thylakoid reactions. Therefore,
we performed pulse-amplitude modulated (PAM) chlorophyll fluorometry on rosette leaves
subjected to increased light. The efficiency of photosystem II photochemistry (ΦPSII) dropped
to the same extent in both wildtype and *phr1-1 phl1* mutants upon exposure to increased
illumination for 125 min (Figure 4A). After 125 min of treatment, also the maximum efficiency
of PSII (F_v/F_m) had decreased significantly in the stressed plants of both genotypes (Figure 4B).
As for ΦPSII, no difference was observed for F_v/F_m between wildtype and *phr1-1 phl1*. Thus,
PHR1/PHL1 function likely affected adenylate levels, starch metabolism and anthocyanin
pigmentation of leaves under high-light stress in a manner independent of the thylakoid
reactions.

255   *PHR1/PHL1* function is important when triose phosphate utilization is limited

Our observations indicated that P_i-starvation signaling might be important during a shift
to higher light intensities. We hypothesized that this might be related to transient imbalances of
triose phosphate production and P_i recycling that occur upon changes in photosynthetic activity.
In order to address this, we tried to genetically mimic the otherwise transient state of insufficient
P_i recycling from photosynthetic products. In the *adg1-1* mutant, disruption of starch
biosynthesis leads to continuous limitation in triose phosphate utilization. Hence, we examined
the phenotypes of the *adg1 phr1 phl1* triple mutant to assess the importance of PHR1/PHL1
signaling during triose phosphate overaccumulation.

Strikingly, when grown under conditions with a short photoperiod (8 hours),
adg1 phr1 phl1 mutant plants exhibited a strong growth defect that was apparent in a reduction
in both leaf size and leaf number compared to *adg1-1* single and *phr1-3 phl1* double mutants
(Figure 5A, B). Under long-day conditions (16 hours photoperiod), the reduction in rosette size
was less pronounced, but *adg1 phr1 phl1* mutant plants exhibited premature leaf senescence
with symptoms already visible at the age of 25 days (Figure 5A, arrow). Accelerated leaf
senescence is also seen in *phr1 phl1* double mutants particularly under P_i deprived conditions
(Bustos et al., 2010) (Figure 1A), but initiates later at the age of approximately 35 days when

grown under standard conditions on soil (Figure 5B) (Wang et al., 2018a). Furthermore,

transition to flowering was much delayed in *adg1 phr1 phl1* compared to the parental lines

(Figure 5C). Notably, delayed onset of the regenerative phase is a phenotype already inherent

to both the *adg1-1* (Matsoukas et al., 2013) and *phr1 phl1* (Supplemental Figure 5) mutants.

We noted that the photoperiod-dependent reduction in rosette size seen for

adg1 phr1 phl1 was reminiscent of the phenotype of the *adg1 tpt-2* mutant (Figure 5A). Thus,

we tested the ability of *adg1 phr1 phl1* to adapt to higher light intensities, since high-light

sensitivity was also reported for the *adg1 tpt-2* mutant (Schmitz et al., 2012). In fact, seedling

shoot growth of *adg1 phr1 phl1* and *adg1 tpt-2* was reduced to a similar extent when grown

under high-light conditions (Figure 5D and Supplemental Figure 6). Moreover, growth of

adg1 phr1 phl1 mutants under high light was promoted by exogenously supplied sucrose

(Figure 5D), as it was described for *adg1 tpt-2* (Heinrichs et al., 2012; Schmitz et al., 2012).

When seedlings were grown on vertically positioned petri dishes, it became evident that sucrose

strongly stimulated root growth of both *adg1 phr1 phl1* and *adg1 tpt-2* particularly under high-

light conditions (Supplemental Figure 7). Interestingly, when analyzing vertically grown

seedlings, we observed a tendency for high-light sensitivity of shoot growth also for the *adg1-*

1 single and *phr1-3 phl1* double mutant lines (Supplemental Figure 6A). Likewise, this effect

could be alleviated by exogenously supplied sucrose. Notably, high-light sensitivity of *phr1-3*

phl1 was more pronounced under high-P_i (2.5 mM) than low-P_i (0.25 mM) growth conditions

(Supplemental Figure 6A), supporting the hypothesis that PHR1/PHL1 play a role for high-

light adaptation also under P_i sufficiency. It was concluded from these experiments that lack of

PHR1/PHL1 compromises plant growth under conditions of high triose phosphate

accumulation such as high light or when starch biosynthesis is genetically blocked.

Carbohydrate signaling is affected during P_i deprivation

Seedling growth of *phr1-3 phl1* (Supplemental Figure 6A) and *adg1 phr1 phl1* (Figure

5D and Supplemental Figure 7) was supported by sucrose supplementation particularly under

high light, and high-light induced starch accumulation was reduced in *phr1-1 phl1* mutants

(Figure 3D). Both observations suggest that *phr1 phl1* mutation leads to alterations in

carbohydrate metabolism. To address the question of how carbohydrate signaling might be

modulated under conditions when PHR1 activity is high, we re-analyzed published

transcriptome data obtained from seedlings subjected to P_i depletion (Bustos et al., 2010), and

from rosette plants exposed to a low CO_2 environment (Bläsing et al., 2005). Among the

transcripts which were most strongly upregulated in shoots of *phr1-1 phl1* under P$_i$ depletion
relative to the wildtype, the majority was repressed by P$_i$ depletion in the wildtype
(Supplemental Figure 8). This supports a function of PHR1/PHL1 in the downregulation of
certain transcripts under P$_i$ depletion, as previously suggested (Bustos et al., 2010).
Interestingly, we noted that many of these transcripts were also responsive to carbon starvation
(Bläsing et al., 2005) (Supplemental Figure 8).

We chose four of the transcripts which were upregulated in *phr1-1 phl1* compared to
the wildtype under P$_i$ starvation for further analysis: The circadian regulators *BASIC LEUCINE*
ZIPPER 63 (*bZIP63*) (Baena-González et al., 2007; Frank et al., 2018) and *REVEILLE 1*
(*RVE1*) (Rawat et al., 2009), as well as *BETA-XYLOSIDASE1* (*BXL1*), and
SENESCENCE1/DARK INDUCIBLE1 (*SEN1/DIN1*), a putative target gene of bZIP63
(Matiolli et al., 2011). In qRT-PCR analysis of seedlings grown on sucrose-containing media
with or without P$_i$, both *bZIP63* and *BXL1* were down-regulated in shoots of wildtype seedlings
exposed to P$_i$ depletion, whereas no significant changes were detected for *phr1-1 phl1* mutants
(Figure 5E), potentially reflecting sugar accumulation in the wildtype (Supplemental Figure 1).
In contrast, transcript levels of *SEN1* and *RVE1* were highly increased in *phr1-1 phl1* mutants
under P$_i$ deprivation, but no significant changes were seen in the wildtype background (Figure
5F). Thus, expression of carbohydrate responsive genes, including circadian-clock associated
genes, differs between wildtype and *phr1-1 phl1* mutants under P$_i$ depletion.

325 Transcription in the *adg1 tpt-2* mutant exhibits a constitutive P$_i$-starvation signature

The phenotypic similarities of the *adg1 phr1 phl1* and *adg1 tpt-2* mutants suggested that
triose phosphate partitioning and P$_i$ signaling share some physiological consequences,
supporting that triose phosphate utilization affects cellular P$_i$ homeostasis. To assess if
PHR1/PHLs activity was altered in *adg1 tpt-2*, we analyzed P$_i$-starvation marker gene
expression in rosette leaves of plants grown for 5 weeks under an 8-hours light regime and a
moderate light intensity of $90\pm10\,\mu$mol m^{-2} s^{-1}. Surprisingly, *SRG3*, *PS2*, *SPX1*, *VPE1* and
MGD3 were all significantly upregulated in shoots of soil-grown *adg1 tpt-2* plants compared
to wildtype or the single *adg1-1* and *tpt-2* mutants (Figure 6A), indicating high activity of PHR1
in shoots of *adg1 tpt-2*. Moreover, *bZIP63* expression was significantly reduced in both *adg1-1*
and *adg1 tpt-2*, whereas we observed a slight increase in *bZIP63* transcript levels in rosette
leaves of *tpt-2* mutants (Figure 6A). To test if altered expression of P$_i$-starvation associated
genes in *adg1 tpt-2* was linked to reduced P$_i$ levels in the shoots of the mutant, we next

determined P_i contents in 5-weeks-old rosette leaves grown on soil under moderate light.
Interestingly, the P_i content was not reduced in shoots of *adg1 tpt-2* compared to the wildtype,
but P_i levels were even found to be higher than in *adg1-1* single mutants (Figure 6B). Thus, P_i
loading of the shoots was not impaired in *adg1 tpt-2*, and P_i-starvation responsive genes were
instead constitutively upregulated in *adg1 tpt-2* rosette leaves, indicating high PHR1 activity
even under nutrient-sufficient physiological conditions in this mutant.

Discussion

The observations presented herein establish that the P_i response machinery involving
the transcription factors PHR1 and PHL1 is capable of sensing early high-light stress, and we
provide evidence that *PHR1/PHL1* contribute to the rapid adjustment of carbohydrate
metabolism during changes in light intensity. P_i consumption by photosynthesis is thus
hypothesized to generate a chloroplast-to-nucleus retrograde signal. In fact, P_i qualifies as a
subcellular signal by a number of criteria: 1) $[P_i]$ changes immediately and gradually in response
to photoassimilation (Stitt et al., 1980; Dietz and Heber, 1984; Robinson and Giersch, 1987);
2) multiple transport proteins at the organellar membranes (Fabiańska et al., 2019) ensure rapid
transmission of the $[P_i]$ signal; 3) signal propagation is based on diffusion and does not rely on
acceptor and donor molecules; 4) P_i exhibits low toxicity within a wide concentration range; 5)
a highly responsive signaling machinery is present in the cytosol and nucleus.

Increase in light intensity creates a transient low-P_i signal

Under conditions where carboxylation exceeds subsequent metabolic capacities of the
chloroplast, such as high light (Laisk et al., 1991), low temperature (Stitt and Grosse, 1988),
and low oxygen (Klecker et al., 2014), large pools of 3-phosphoglycerate may build up which
can enter the cytosol via the TPT (Schneider et al., 2002). By the nature of the transport
mechanism, this depletes the cytosol of P_i, and it is assumed that P_i import from the vacuole is
too slow to balance against very short-term changes in P_i demand (Woodrow et al., 1984). In
accordance with these considerations, we found that transcription exhibits a P_i-starvation
signature in Arabidopsis rosette leaves subjected to light increase already after 20 min of stress
(Figure 2). This response depended on *PHR1/PHL1* function, indicating that light stress
activated the cytosolic and nuclear P_i sensing machinery (Figure 7). PHR1 target genes which
showed enhanced expression in our experimental setup included *SRG3*, *SPX1*, and *VPE1*, and

all three transcripts also showed early enrichment upon high-light treatment within a time-resolved transcriptome analysis previously described by Huang et al. (2019). Both *SRG3* and *VPE1* expression might be suitable to rapidly increase P_i availability in the cytosol via glycerophosphodiester turnover (Cheng et al., 2011) and vacuolar P_i efflux (Xu et al., 2019), respectively. In contrast, *SPX1* upregulation potentially contributes to the rapid tempering of the transcriptional response which was quenched already after 2 hours of treatment (Figure 2). Next to negative feed-back by SPX1, adjustment of triose phosphate turnover by post-translational activation of sucrose (Stitt et al., 1983; Preiss, 1984) and starch (Preiss, 1988; Hendriks et al., 2003) biosynthesis would be expected to attenuate any condition of light-induced P_i depletion within relatively short time. Therefore, long-term P_i-starvation responses such as lipid remodeling are rather unlikely to occur upon light intensity increases as long as higher P_i demand can be sufficiently covered by uptake via the roots or release of P_i storage within the mesophyll cell. Consistent with this assumption, induction of *MGD3* transcript was moderate in our experimental setup (Figure 2). Interestingly however, van Rooijen et al. (2018) observed upregulation of PHR1 target genes with involvement in lipid remodeling after 1 hour of irradiance increase. Thus, it will be exciting to learn from future experiments whether P_i signaling also contributes to the changes that can be seen in the lipid composition of leaves under different light conditions (Yu et al., 2021).

P_i sensing impacts on carbohydrate and energy metabolism upon light increase

The induction of genes related to P_i mobilization upon high light might be considered a transcriptional feedback response initiated in order to sustain substrate supply for metabolic reactions. Classification of P_i as a retrograde signal requires, beyond that, P_i signaling to affect further physiological processes by calibrating nuclear gene expression according to the status of the plastids. Crucially, this study shows that mutants defective in *PHR1* and *PHL1* which fail to mount a transcriptional response related to P_i starvation, develop metabolic abnormalities in the leaves within only 1.5 hours of high-light treatment (Figure 3A-D), and exhibit lower levels of anthocyanin pigmentation after 3 days of treatment (Figure 3E). ATP levels measured after 1.5 hours of high-light exposure were significantly lower in leaves of *phr1-1 phl1* mutant plants compared to wildtype plants (Figure 3A), whereas ΦPSII values were indistinguishable (Figure 4A). Given that ΦPSII can be interpreted as a measure of the linear electron transport rate (Maxwell and Johnson, 2000), this indicates that ATP synthesis at the thylakoids was likely not affected by loss of *PHR1/PHL1* function. This is consistent with the assumption that P_i

sequestration upon photosynthetic activity does not compromise stromal P_i levels and thus ATP

synthesis as long as phosphorylated metabolites are exchanged with the cytosol via P_i antiport

systems (Robinson and Giersch, 1987). In contrast, it would be conceivable that the P_i

concentration outside of the chloroplast might drop below levels required for efficient

mitochondrial ATP synthesis in the *phr1-1 phl1* mutant. In fact, a decline in ATP levels can

also be observed in plants under P_i-deficient growth conditions (Zhu et al., 2019; Riemer et al.,

2021), but the reason for this is currently unknown.

While starch levels under standard light conditions were slightly higher in *phr1-1 phl1*

rosettes compared to wildtype, the high-light induced relative increase in starch content was

considerably higher in wildtype leaves compared to the mutant (Figure 3D). This delay in the

activation of starch biosynthesis is remarkable, given that post-translational activation of

AGPase by the 3-PGA/P_i ratio is probably high in *phr1-1 phl1* leaves which manifest

chronically low P_i contents (Wang et al., 2018a). Similar to what can be seen under P_i

deprivation (Supplemental Figure 1) (Nilsson et al., 2007), the difference in starch

accumulation observed between wildtype and *phr1-1 phl1* mutants upon transfer to high light

was not reflected by changes in soluble sugar contents (Figure 3B, C). Together with our finding

that photosynthetic capacity is wildtype-like in *phr1-1 phl1* mutants subjected to high-light

stress (Figure 4), this suggests that starch production is rather not limited by photosynthetic

precursor production in *phr1-1 phl1*. Hence, it is likely that PHR1/PHL1 directly regulate

photoassimilate partitioning towards starch biosynthesis. Expression of *APL3* (AT4G39210)

encoding a large subunit of AGPase, or *GGBS1* (AT1G32900), coding for a granule-bound

starch synthase have been described to be induced by P_i starvation (Morcuende et al., 2007;

Bustos et al., 2010; Wang et al., 2018b), and might contribute to both P_i-starvation and high-

light induced starch accumulation.

P_i signaling is crucial under conditions of triose-phosphate oversupply

Prevention of starch production during P_i restriction leads to higher sugar and

anthocyanin levels in the shoots (Figure 1). The observation is consistent with sugars being

formed upon recycling of phosphorylated metabolites (Morcuende et al., 2007), and starch

serving as a carbon sink in P_i-deprived plants (Hammond and White, 2008; MacNeill et al.,

2017). Neither sugar accumulation, nor transitory starch or anthocyanin production take place

in *phr1 phl1* mutant seedlings subjected to P_i depletion (Figure 1 and Supplemental Figure 1).

Nevertheless, prohibition of starch biosynthesis by introduction of the *adg1-1* allele into this

mutant background produced a pronounced growth defect already visible at the seedling stage

(Figure 5A, B, D and Supplementary Figures 6, 7). Growth of the *adg1 phr1 phl1* triple mutant

was further impaired by higher light intensities (Figure 5D and Supplemental Figure 6), a

condition that reinforces triose phosphate generation. These observations lead to the conclusion

that *PHR1/PHL1* function is crucial in the *adg1-1* knockout situation, demonstrating the

necessity for low-P_i signaling when photoassimilate consumption is constrained. Next to

growth retardation, the *adg1 phr1 phl1* mutant showed a delay in the onset of flowering (Figure

5C), and benefited from exogenous sucrose supply in terms of both shoot and root growth

(Figure 5D and Supplemental Figures 6, 7). Both late flowering (Wahl et al., 2013) and

inhibition of primary root growth (Smith and Stitt, 2007) indicate that *adg1 phr1 phl1* plants

suffer from carbohydrate starvation in the absence of external sucrose supply.

Interestingly, transcriptional changes observed on wildtype and *phr1-1 phl1* mutant

plants under P_i deprivation also indicate that PHR1/PHL1 maintain sugar homeostasis when P_i

availability is low (Figure 5E, F and Supplemental Figure 8). In agreement, strong increments

in the levels of amino acids and oligosaccharides such as raffinose and kestose were reported

for wildtype but not *phr1* mutants under P_i depletion (Pant et al., 2015a). Notably,

oligosaccharide production facilitated by PHR1/PHL1 might serve as an alternative

carbohydrate sink in the *adg1-1* mutant, and could support carbon supply of the starchless

mutant during the night. This would be consistent with the observation that *adg1 phr1 phl1* was

more strongly constrained by diel cycles with short light phases (Figure 5A). Similarly,

oligosaccharide and amino acid accumulation triggered by the low-P_i signaling module could

support carbohydrate homeostasis during high-light stress by maintaining P_i/C(/N) balance, and

might even be suitable to prepare for potential osmotic stress as a likely consequence of high

light in a natural environment. In line with this, a function of PHR1/PHL1 in proline

accumulation and drought resistance has been reported (Aleksza et al., 2017; Scheible et al.,

2023). It is difficult to dissect direct effects of P_i signaling from secondary consequences of P_i-

deprived metabolism. Thus, metabolic differences of *adg1 phr1 phl1* mutants grown under

short and long photoperiods might reveal new insights into how carbon fluxes are modulated

by P_i signaling.

P_i signaling is affected in *adg1 tpt-2*

Strikingly, growth defects and high-light sensitivity of *adg1 phr1 phl1* are paralleled by

the phenotypes of the *adg1 tpt-2* mutant (Figure 5A, D and Supplemental Figures 6, 7),

indicating that photoassimilate allocation and P_i signaling to some extent overlap in terms of

their physiological consequences. Even under P_i sufficient growth conditions, we found that

adg1 tpt-2 exhibits a signature of P_i-starvation responsive gene expression (Figure 6), raising

the question how insufficient export of photosynthetic products could generate a situation of

high PHR1 activity. In the *adg1 tpt-2* mutant, a pronounced fraction of cytosolic/chloroplastic

P_i can be assumed as metabolically unavailable, given that starch biosynthesis is prevented by

adg1-1 mutation, while sucrose production during the course of the day remains at much lower

levels compared to *adg1-1* (Schmitz et al., 2012) (Figure 7). Hence, bulk release of P_i in *adg1*

tpt-2 might be restricted to glycolysis and vacuolar phosphatase activity acting on

phosphorylated intermediates such as phosphoenolpyruvate (Ohnishi et al., 2018). This would

cause unequal intracellular pool sizes of free P_i, facilitating PHR1 activity in the

cytosol/nucleus, while vacuolar P_i stores are high (Figure 6).

In fact, imbalances of intracellular P_i pools would affect both *PHR1*-dependent and

independent (Osorio et al., 2019; Nam et al., 2021) P_i-responsive pathways. Such inadequate

signaling could even contribute to the pleiotropic defects observed for *adg1 tpt-2* mutants

(Schmitz et al., 2012), for example by altering lipid composition, or by the regulation of

photosynthesis-associated genes (Morcuende et al., 2007; Bustos et al., 2010; Barragán-Rosillo

et al., 2021; Nam et al., 2021). Interestingly, we also found that transcript levels of the circadian

regulator bZIP63 were downregulated both by P_i depletion (Figure 5E), and in the *adg1 tpt-2*

mutant (Figure 6). Given that *adg1 tpt-2* is characterized by overall low carbohydrate levels

(Schmitz et al., 2012), *bZIP63* expression appears to be uncoupled from sugar levels in this

mutant situation. Our observations open up the possibility that P_i signaling could be involved

in this regulation which will be an exciting topic to address in the future.

Conclusions

High-light stress requires immediate responses to avert damage to cellular components.

As a potential mode of sensing excess photosynthetic activity, we describe a function of the P_i-

sensing machinery involving PHR1 and PHL1 transcription factors under nutrient replete

growth conditions. We propose that P_i sequestration into phosphorylated photoassimilates and

metabolite exchange across the chloroplast membrane causes transient P_i restriction in the

cytosol and nucleus. This activates PHR1/PHL1 proteins to mobilize cellular P_i and to support

acclimation responses such as starch biosynthesis. The finding that P_i signaling is disturbed in

the *adg1 tpt-2* mutant emphasizes that PHR1/PHL activity constitutes a so-far underestimated
contributor to retrograde signaling of photosynthetic status.

Materials and Methods

Plant material

All Arabidopsis plants used in this study are in the background of the Columbia-0
accession. The *phr1-3* (AT4G28610; SALK_067629C) (Nilsson et al., 2007) and *phl1*
(AT5G29000; SAIL_731_B09) (Klecker et al., 2014) lines were described previously and the
derived homozygous double mutant *phr1-3 phl1* was a kind gift of Angelika Mustroph
(University of Bayreuth). The double mutant *phr1-1 phl1* (Bustos et al., 2010) was backcrossed
to *phl1* in order to remove a transgene containing the *NPTII* cassette which was previously
introduced for isolation of the *phr1-1* allele (Rubio et al., 2001). All experiments were
performed with the homozygous double mutant *phr1-1 phl1* lacking the *NPTII* cassette. The
double mutant *spx1 spx2* was described earlier (Puga et al., 2014). Knockout lines *adg1-1*
(At5g48300) (Lin et al., 1988), *tpt-2* (At5g46110; SALK_073707.54.25.x) and *adg1 tpt-2*
(Schmitz et al., 2012) were kindly provided by Rainer E. Häusler (University of Cologne).
gpt2-1 (AT1G61800; GK-454H06) was obtained from The Nottingham Arabidopsis Stock
Centre (NASC) and was described previously (Niewiadomski et al., 2005). Primer sequences
used for genotyping are listed in Table S1. Homozygous insertion mutants were verified using
the following primer combinations: *PHR1* WT: PHR1_F/PHR1_R; *phr1-3* T-DNA:
LBb1/PHR1_R; *PHL1* WT: PHL1_F/PHL1_R; *phl1* T-DNA: PHL1_F/LB3; *TPT* WT:
TPT_F/TPT_R; *tpt-2* T-DNA: *tpt-2*_F/LBb1; *GPT2* WT: GPT2_F/GPT2_R; *gpt2-1* T-DNA:
GK-LB/GPT2_R. To generate the *adg1 phr1 phl1* triple mutant, *adg1-1* was crossed to *phr1-3*
phl1. Homozygous *adg1-1* mutants were identified by screening for starch-free phenotypes by
iodine staining at the end of the photoperiod. Homozygosity of the *phr1-1* allele was verified
by Sanger sequencing (Macrogen Europe, Amsterdam) of the PCR product of
PHR1_F/PHR1_R using primer PHR1_R.

Cultivation conditions

530   Seeds were stratified for 64 hours at 4°C in the dark. For vegetative growth, plants were
531 germinated and grown in an 8 h/16 h (light/dark) cycle on soil consisting of seeding compost,
532 universal soil, and vermiculite mixed in a 3:3:1 ratio. For high-light experiments, the plants

were additionally fertilized once directly after pricking at the age of 10 days using a complete mineral mixture (Wuxal Super, Aglukon Spezialduenger, Duesseldorf, Germany) according to the manufacturer's instructions. For seedling experiments, seeds were surface sterilized by chlorine gas exposure and transferred to solidified Arabidopsis growth media as described in the figure legends. Seedlings were grown under 16 h/8 h light/dark cycles. For paraquat treatments, seedlings were germinated on agar-solidified media containing full-strength Murashige and Skoog (MS) salts and grown for 10 days before transfer to MS media with or without 1 µM paraquat dichloride hydrate (#36541, Riedel-de Haën, Hannover, Germany). If not indicated otherwise, growth light was set to 90 ± 10 µmol m^{-2} sec^{-1} and monitored using an illuminance meter.

P$_i$-starvation experiments

For P$_i$-starvation experiments, seeds were germinated on medium containing MS salts at full strength together with 0.8 % (w/v) phytoagar (Duchefa Biochemie, Haarlem, The Netherlands) and 0.5 % (w/v) sucrose. After 5-10 days of growth (see figure legends), seedlings were transferred to P$_i$ media (0.5 % (w/v) sucrose; 20 mM 2-(N-Morpholino)-ethane sulphonic acid; 2.5 mM KNO$_3$; 1 mM MgSO$_4$; 1 mM Ca(NO$_3$)$_2$; 2.5 mM KH$_2$PO$_4$; 25 µM Fe-EDTA; 35 µM H$_3$BO$_3$; 7 µM MnCl$_2$; 0.25 µM CuSO$_4$; 0.5 µM ZNSO$_4$; 0.1 µM NaMoO$_4$; 5 µM NaCl; 0.005 µM CoCl$_2$; pH 6) solidified by 0.8 % (w/v) agar (adopted from Härtel et al. (2000)). For P$_i$ depleting conditions, KH$_2$PO$_4$ was omitted, and Fe-EDTA was reduced to 10 µM in order to minimize low-P$_i$ induced iron toxicity (Ward et al., 2008). Seedling shoots were harvested after 7-8 days of growth on P$_i$ differing media (see figure legends). For horizontally grown seedlings, MS salts were used at ½ concentrations and 0.6 % (w/v) agar were applied. For the experiment shown in Figures S6 and S7, P$_i$ medium was used containing either 0.25 mM KH$_2$PO$_4$/10 µM Fe-EDTA, 2.5 mM KH$_2$PO$_4$/25 µM Fe-EDTA, or 2.5 mM KH$_2$PO$_4$/25 µM Fe-EDTA +50 mM sucrose.

Vector construction

Standard molecular cloning procedures were applied. Primer sequences used for cloning are listed in table S1. The firefly *LUCIFERASE* reporter plasmid *pBT10-proMGD3::LUC$_{Firefly}$* and the *pHBTL-p35S::3HA-GFP* effector plasmid (Klecker et al., 2014), as well as the *pBT10-p35S::LUC$_{Renilla}$* normalization plasmid (Bäumler et al., 2019) have been described elsewhere.

565   For details on cloning *proSPX1^{GC}, proSRG3*, *proMGD3*, *proGPT2,* and the *PHR1* effector

566   construct, refer to the supplemental methods.

Anthocyanin determination

Anthocyanin contents were determined with few changes as described in

(Vandenbussche et al., 2007). In brief, 40-70 mg of seedling shoots or rosette leaf material were

frozen in liquid nitrogen and ground in a bead mill (MM400, Retsch, Haan, Germany). The

frozen powder was resuspended in 300 µl of methanol containing 1 % HCl. After vortex-mixing

for 15 sec, cell debris was pelleted by centrifugation for 15 min at >14,000 g, 4°C. The

supernatant was mixed with 200 µl water and 500 µl chloroform by vortexing for 15 sec. After

15 min centrifugation at 17,900 g, 4°C, the upper phase was diluted in methanol to measure the

absorbance at 535 nm and 630 nm using a spectrophotometer (Specord 200 Plus, Analytik Jena,

Jena, Germany).

Measurements of starch, soluble sugars, and adenylates

580   Carbohydrates were determined by the Warburg optical test and adenylates were

581 assessed based on luciferase activity. Both procedures were described previously (Mustroph et

582 al., 2006), for details see supplemental methods.

Determination of P_i contents

585   Free P_i was determined in leaf tissue based on the method described by (Ames, 1966)

586 with some modifications partially based on (Sakuraba et al., 2018). For detail see supplemental

587 methods.

RNA extraction and cDNA synthesis

Plant material was harvested under the respective experimental conditions and instantly

frozen in liquid nitrogen. The tissue was ground in a bead mill, and RNA was extracted using

Bioline TRIsure (Meridian Bioscience, Cincinnati, United States) according to the

manufacturer's instructions. The quality of the RNA was assessed by screening A_{260} and A_{280}.

For qRT-PCR analysis, 1 µg of RNA was treated with DNAse I (Thermo Fisher Scientific,

Waltham, United States) according to product instructions before application in reverse

transcription using oligo(dT)$_{15}$ and RevertAid Reverse Transcriptase (Thermo Fisher Scientific). Complementary DNA was diluted by a factor of 50 for use as template in qPCR analysis. qPCR was performed in technical triplicates using SsoAdvanced Universal SYBR Green Supermix (Bio-Rad, Hercules, United States) on a CFX Connect Real-time System (Bio-Rad). Transcript levels were normalized to the levels of *PP2A* transcript and calculated as $1000 \cdot 2^{-\Delta CT}$. Primer sequences are listed in table S1.

Protoplast isolation, transfection and transactivation assay

Leaf mesophyll protoplasts were isolated using the "Tape-Arabidopsis Sandwich" method (Wu et al., 2009) with minor modifications. For details, see supplemental methods.

Chlorophyll fluorescence measurements

For the determination of the quantum yield of photosystem II photochemistry $(F_m'-F')/F_m'$ (ΦPSII), as well as the maximum efficiency of photosystem II (F_v'/F_m'), pulse-amplitude-modulated fluorescence measurements (PAM) of chlorophyll fluorescence were conducted using a Junior-PAM (Walz, Effeltrich, Germany). Here, to calculate chlorophyll fluorescence parameters using the saturating pulse method (Schreiber et al., 1986), one large rosette leaf was held between two magnets and subjected to 10 seconds of actinic light irradiance set to control light intensity (80 ± 5 µmol m^{-2} sec^{-1}), followed by a saturation light pulse.

Statistical analysis

Statistical analyses were performed with Microsoft Office Excel using the Real Statistics Resource Pack software (www.real-statistics.com).

Funding information

This research was funded by the University of Bayreuth.

Acknowledgements

We thank Stephan Clemens and Angelika Mustroph for helpful discussions and comments on the manuscript, as well as for providing infrastructure and equipment. We thank Kristina Munzert for help with experiments.

Author contributions

M. K. designed the research, analyzed data, and wrote the paper; L. A., M. M., A. H., and M. K. performed research.

Supplemental Data

Supplemental Figure 1 Changes in sugar and starch contents under P_i depletion.

Supplemental Figure 2 Expression of selected P_i-starvation responsive genes is activated by PHR1/PHL1.

Supplemental Figure 3 *PHR1* transcript levels are not affected by short-time high-light exposure.

Supplemental Figure 4 Anthocyanin production of P_i signaling mutants under ROS stress.

Supplemental Figure 5 Delayed onset of flowering in *phr1-1 phl1* mutants.

Supplemental Figure 6 Quantification of shoot fresh weights of seedlings grown under low and high light.

Supplemental Figure 7 Root growth of *adg1 phr1 phl1* and *adg1 tpt-2* responds to exogenously supplied sucrose.

Supplemental Figure 8 Carbohydrate responsive gene expression is affected under P_i starvation.

Supplemental Table 1 List of primer sequences used in this study.

Supplemental Methods

REFERENCES

Aleksza D, Horváth G V., Sándor G, Szabados L (2017) Proline accumulation is regulated by transcription factors associated with phosphate starvation. Plant Physiol **175**: 555–567

Ames BN (1966) [10] Assay of inorganic phosphate, total phosphate and phosphatases. Methods Enzymol. pp 115–118

Athanasiou K, Dyson BC, Webster RE, Johnson GN (2010) Dynamic acclimation of photosynthesis increases plant fitness in changing environments. Plant Physiol **152**: 366–73

Baena-González E, Rolland F, Thevelein JM, Sheen J (2007) A central integrator of transcription networks in plant stress and energy signalling. Nature **448**: 938–42

Barragán-Rosillo AC, Peralta-Alvarez CA, Ojeda-Rivera JO, Arzate-Mejía RG, Recillas-Targa F, Herrera-Estrella L (2021) Genome accessibility dynamics in response to phosphate limitation is controlled by the PHR1 family of transcription factors in Arabidopsis. Proc Natl Acad Sci U S A **118**: e2107558118

Bäumler J, Riber W, Klecker M, Müller L, Dissmeyer N, Weig AR, Mustroph A (2019) AtERF#111/ABR1 is a transcriptional activator involved in the wounding response. Plant J **100**: 969–990

Bechtold U, Richard O, Zamboni A, Gapper C, Geisler M, Pogson B, Karpinski S, Mullineaux PM (2008) Impact of chloroplastic- and extracellular-sourced ROS on high light-responsive gene expression in Arabidopsis. J Exp Bot **59**: 121–33

Bläsing OE, Gibon Y, Günther M, Höhne M, Morcuende R, Osuna D, Thimm O, Usadel B, Scheible W-R, Stitt M (2005) Sugars and circadian regulation make major contributions to the global regulation of diurnal gene expression in Arabidopsis. Plant Cell **17**: 3257–81

Bustos R, Castrillo G, Linhares F, Puga MI, Rubio V, Pérez-Pérez J, Solano R, Leyva A, Paz-Ares J (2010) A central regulatory system largely controls transcriptional activation and repression responses to phosphate starvation in Arabidopsis. PLoS Genet **6**: e1001102

Castrillo G, Teixeira PJPL, Paredes SH, Law TF, de Lorenzo L, Feltcher ME, Finkel OM, Breakfield NW, Mieczkowski P, Jones CD, et al (2017) Root microbiota drive direct integration of phosphate stress and immunity. Nature **543**: 513–518

Cheng Y, Zhou W, El Sheery NI, Peters C, Li M, Wang X, Huang J (2011) Characterization

680 of the Arabidopsis glycerophosphodiester phosphodiesterase (GDPD) family reveals a role

681 of the plastid-localized AtGDPD1 in maintaining cellular phosphate homeostasis under

682 phosphate starvation. Plant J **66**: 781–95

683 **Dietz K-J** (2015) Efficient high light acclimation involves rapid processes at multiple

684 mechanistic levels. J Exp Bot **66**: 2401–14

685 **Dietz K-J, Heber U** (1986) Light and CO2 limitation of photosynthesis and states of the

686 reactions regenerating ribulose 1,5-bisphosphate or reducing 3-phosphoglycerate.

687 Biochim Biophys Acta - Bioenerg **848**: 392–401

688 **Dietz K-J, Turkan I, Krieger-Liszkay A** (2016) Redox- and Reactive Oxygen Species-

689 Dependent Signaling into and out of the Photosynthesizing Chloroplast. Plant Physiol **171**:

690 1541–50

691 **Dietz K, Heber U** (1984) Rate-limiting factors in leaf photosynthesis. I. Carbon fluxes in the

692 calvin cycle. Biochim Biophys Acta - Bioenerg **767**: 432–443

693 **Dietz KJ, Foyer C** (1986) The relationship between phosphate status and photosynthesis in

694 leaves : Reversibility of the effects of phosphate deficiency on photosynthesis. Planta **167**:

695 376–81

696 **Dong J, Ma G, Sui L, Wei M, Satheesh V, Zhang R, Ge S, Li J, Zhang T-E, Wittwer C, et**

697 **al** (2019) Inositol Pyrophosphate InsP8 Acts as an Intracellular Phosphate Signal in

698 Arabidopsis. Mol Plant **12**: 1463–1473

699 **Fabiańska I, Bucher M, Häusler RE** (2019) Intracellular phosphate homeostasis - A short

700 way from metabolism to signaling. Plant Sci **286**: 57–67

701 **Figueroa CM, Asencion Diez MD, Ballicora MA, Iglesias AA** (2022) Structure, function,

702 and evolution of plant ADP-glucose pyrophosphorylase. Plant Mol Biol **108**: 307–323

703 **Frank A, Matiolli CC, Viana AJC, Hearn TJ, Kusakina J, Belbin FE, Wells Newman D,**

704 **Yochikawa A, Cano-Ramirez DL, Chembath A, et al** (2018) Circadian Entrainment in

705 Arabidopsis by the Sugar-Responsive Transcription Factor bZIP63. Curr Biol **28**: 2597-

706 2606.e6

707 **Hammond JP, White PJ** (2008) Sucrose transport in the phloem: integrating root responses to

708 phosphorus starvation. J Exp Bot **59**: 93–109

709 **Hanchi M, Thibaud M-C, Légeret B, Kuwata K, Pochon N, Beisson F, Cao A, Cuyas L,**

710 **David P, Doerner P, et al** (2018) The Phosphate Fast-Responsive Genes PECP1 and

PPsPase1 Affect Phosphocholine and Phosphoethanolamine Content. Plant Physiol **176**: 2943–2962

Härtel H, Dormann P, Benning C (2000) DGD1-independent biosynthesis of extraplastidic galactolipids after phosphate deprivation in Arabidopsis. Proc Natl Acad Sci U S A **97**: 10649–54

Häusler RE, Heinrichs L, Schmitz J, Flügge UI (2014) How sugars might coordinate chloroplast and nuclear gene expression during acclimation to high light intensities. Mol Plant **7**: 1121–1137

Heinrichs L, Schmitz J, Flügge U-I, Häusler RE (2012) The Mysterious Rescue of adg1-1/tpt-2 - an Arabidopsis thaliana Double Mutant Impaired in Acclimation to High Light - by Exogenously Supplied Sugars. Front Plant Sci **3**: 265

Heldt HW, Rapley L (1970) Specific transport of inorganic phosphate, 3-phosphoglycerate and dihydroxyacetonephosphate, and of dicarboxylates across the inner membrane of spinach chloroplasts. FEBS Lett **10**: 143–148

Hendriks JHM, Kolbe A, Gibon Y, Stitt M, Geigenberger P (2003) ADP-glucose pyrophosphorylase is activated by posttranslational redox-modification in response to light and to sugars in leaves of Arabidopsis and other plant species. Plant Physiol **133**: 838–49

Hilgers EJA, Schöttler MA, Mettler-Altmann T, Krueger S, Dörmann P, Eicks M, Flügge UI, Häusler RE (2018) The combined loss of triose phosphate and xylulose 5-phosphate/phosphate translocators leads to severe growth retardation and impaired photosynthesis in arabidopsis thaliana tpt/xpt double mutants. Front Plant Sci **9**: 1–24

Huang J, Zhao X, Chory J (2019) The Arabidopsis Transcriptome Responds Specifically and Dynamically to High Light Stress. Cell Rep **29**: 4186-4199.e3

Klecker M, Gasch P, Peisker H, Dörmann P, Schlicke H, Grimm B, Mustroph A (2014) A Shoot-Specific Hypoxic Response of Arabidopsis Sheds Light on the Role of the Phosphate-Responsive Transcription Factor PHOSPHATE STARVATION RESPONSE1. Plant Physiol **165**: 774–790

Kobayashi K, Awai K, Takamiya K, Ohta H (2004) Arabidopsis type B monogalactosyldiacylglycerol synthase genes are expressed during pollen tube growth and induced by phosphate starvation. Plant Physiol **134**: 640–8

König J, Muthuramalingam M, Dietz K-J (2012) Mechanisms and dynamics in the

thiol/disulfide redox regulatory network: transmitters, sensors and targets. Curr Opin Plant

Biol **15**: 261–8

Kumar V, Sharma A, Bhardwaj R, Thukral AK (2019) Elemental Composition of Plants

and Multivariate Analysis. Natl Acad Sci Lett **42**: 45–50

Kunz HH, Häusler RE, Fettke J, Herbst K, Niewiadomski P, Gierth M, Bell K, Steup M,

Flügge U-I, Schneider A (2010) The role of plastidial glucose-6-phosphate/phosphate

translocators in vegetative tissues of Arabidopsis thaliana mutants impaired in starch

biosynthesis. Plant Biol (Stuttg) **12 Suppl 1**: 115–28

Laisk A, Siebke K, Gerst U, Eichelmann H, Oja V, Heber U (1991) Oscillations in

photosynthesis are initiated and supported by imbalances in the supply of ATP and

NADPH to the Calvin cycle. Planta **185**: 554–62

Lei M, Liu Y, Zhang B, Zhao Y, Wang X, Zhou Y, Raghothama KG, Liu D (2011) Genetic

and genomic evidence that sucrose is a global regulator of plant responses to phosphate

starvation in Arabidopsis. Plant Physiol **156**: 1116–30

Li H, He K, Zhang Z, Hu Y (2023) Molecular mechanism of phosphorous signaling inducing

anthocyanin accumulation in Arabidopsis. Plant Physiol Biochem **196**: 121–129

Lin TP, Caspar T, Somerville C, Preiss J (1988) Isolation and Characterization of a

Starchless Mutant of Arabidopsis thaliana (L.) Heynh Lacking ADPglucose

Pyrophosphorylase Activity. Plant Physiol **86**: 1131–5

Liu Y, Xie Y, Wang H, Ma X, Yao W, Wang H (2017) Light and ethylene coordinately

regulate the phosphate starvation response through transcriptional regulation of

PHOSPHATE STARVATION RESPONSE1. Plant Cell **29**: 2269–2284

Liu Z, Wu X, Wang E, Liu Y, Wang Y, Zheng Q, Han Y, Chen Z, Zhang Y (2022) PHR1

positively regulates phosphate starvation-induced anthocyanin accumulation through

direct upregulation of genes F3'H and LDOX in Arabidopsis. Planta **256**: 42

MacNeill GJ, Mehrpouyan S, Minow MAA, Patterson JA, Tetlow IJ, Emes MJ (2017)

Starch as a source, starch as a sink: the bifunctional role of starch in carbon allocation. J

Exp Bot **68**: 4433–4453

Matiolli CC, Soares RC, Alves HLS, Abreu IA (2022) Turning the Knobs: The Impact of

Post-translational Modifications on Carbon Metabolism. Front Plant Sci **12**: 781508

Matiolli CC, Tomaz JP, Duarte GT, Prado FM, Del Bem LEV, Silveira AB, Gauer L,

773 **Corrêa LGG, Drumond RD, Viana AJC, et al** (2011) The Arabidopsis bZIP gene
774   AtbZIP63 is a sensitive integrator of transient abscisic acid and glucose signals. Plant
775   Physiol **157**: 692–705

776 **Matsoukas IG, Massiah AJ, Thomas B** (2013) Starch metabolism and antiflorigenic signals
777   modulate the juvenile-to-adult phase transition in Arabidopsis. Plant Cell Environ **36**:
778   1802–11

779 **Maxwell K, Johnson GN** (2000) Chlorophyll fluorescence--a practical guide. J Exp Bot **51**:
780   659–68

781 **Misson J, Raghothama KG, Jain A, Jouhet J, Block MA, Bligny R, Ortet P, Creff A,**
782   **Somerville S, Rolland N, et al** (2005) A genome-wide transcriptional analysis using
783   Arabidopsis thaliana Affymetrix gene chips determined plant responses to phosphate
784   deprivation. Proc Natl Acad Sci U S A **102**: 11934–9

785 **Moore M, Vogel MO, Dietz KJ** (2014) The acclimation response to high light is initiated
786   within seconds as indicated by upregulation of AP2/ERF transcription factor network in
787   Arabidopsis thaliana. Plant Signal Behav **9**: 1–4

788 **Morcuende R, Bari R, Gibon Y, Zheng W, Pant BD, Bläsing O, Usadel B, Czechowski T,**
789   **Udvardi MK, Stitt M, et al** (2007) Genome-wide reprogramming of metabolism and
790   regulatory networks of Arabidopsis in response to phosphorus. Plant Cell Environ **30**: 85–
791   112

792 **Mustroph A, Boamfa EI, Laarhoven LJJ, Harren FJM, Albrecht G, Grimm B** (2006)
793   Organ-specific analysis of the anaerobic primary metabolism in rice and wheat seedlings.
794   I: Dark ethanol production is dominated by the shoots. Planta **225**: 103–14

795 **Nam H-I, Shahzad Z, Dorone Y, Clowez S, Zhao K, Bouain N, Lay-Pruitt KS, Cho H,**
796   **Rhee SY, Rouached H** (2021) Interdependent iron and phosphorus availability controls
797   photosynthesis through retrograde signaling. Nat Commun **12**: 7211

798 **Niewiadomski P, Knappe S, Geimer S, Fischer K, Schulz B, Unte US, Rosso MG, Ache P,**
799   **Flügge UI, Schneider A** (2005) The Arabidopsis plastidic glucose 6-phosphate/phosphate
800   translocator GPT1 is essential for pollen maturation and embryo sac development. Plant
801   Cell **17**: 760–775

802 **Nilsson L, Lundmark M, Jensen PE, Nielsen TH** (2012) The Arabidopsis transcription factor
803   PHR1 is essential for adaptation to high light and retaining functional photosynthesis
804   during phosphate starvation. Physiol Plant **144**: 35–47

Nilsson L, Müller R, Nielsen TH (2007) Increased expression of the MYB-related
transcription factor, PHR1, leads to enhanced phosphate uptake in Arabidopsis thaliana.
Plant Cell Environ **30**: 1499–512

Ohnishi M, Anegawa A, Sugiyama Y, Harada K, Oikawa A, Nakayama Y, Matsuda F,
Nakamura Y, Sasaki R, Shichijo C, et al (2018) Molecular Components of Arabidopsis
Intact Vacuoles Clarified with Metabolomic and Proteomic Analyses. Plant Cell Physiol
59: 1353–1362

Osorio MB, Ng S, Berkowitz O, De Clercq I, Mao C, Shou H, Whelan J, Jost R (2019)
SPX4 Acts on PHR1-Dependent and -Independent Regulation of Shoot Phosphorus Status
in Arabidopsis. Plant Physiol **181**: 332–352

Pant B-D, Pant P, Erban A, Huhman D, Kopka J, Scheible W-R (2015a) Identification of
primary and secondary metabolites with phosphorus status-dependent abundance in
Arabidopsis, and of the transcription factor PHR1 as a major regulator of metabolic
changes during phosphorus limitation. Plant Cell Environ **38**: 172–87

Pant BD, Burgos A, Pant P, Cuadros-Inostroza A, Willmitzer L, Scheible WR (2015b) The
transcription factor PHR1 regulates lipid remodeling and triacylglycerol accumulation in
Arabidopsis thaliana during phosphorus starvation. J Exp Bot **66**: 1907–1918

Paz-Ares J, Puga MI, Rojas-Triana M, Martinez-Hevia I, Diaz S, Poza-Carrión C,
Miñambres M, Leyva A (2022) Plant adaptation to low phosphorus availability: Core
signaling, crosstalks, and applied implications. Mol Plant **15**: 104–124

Preiss J (1984) Starch, sucrose biosynthesis and partition of carbon in plants are regulated by
orthophosphate and triose-phosphates. Trends Biochem Sci **9**: 24–27

Preiss J (1988) Biosynthesis of starch and its regulation. *In* J Preiss, ed, Biochem. Plants, 14th
ed. Academic Press, San Diego, pp 181–254

Puga MI, Mateos I, Charukesi R, Wang Z, Franco-Zorrilla JM, de Lorenzo L, Irigoyen
ML, Masiero S, Bustos R, Rodríguez J, et al (2014) SPX1 is a phosphate-dependent
inhibitor of Phosphate Starvation Response 1 in Arabidopsis. Proc Natl Acad Sci U S A
111: 14947–52

Ramaiah M, Jain A, Baldwin JC, Karthikeyan AS, Raghothama KG (2011)
Characterization of the phosphate starvation-induced glycerol-3-phosphate permease gene
family in Arabidopsis. Plant Physiol **157**: 279–91

Rawat R, Schwartz J, Jones MA, Sairanen I, Cheng Y, Andersson CR, Zhao Y, Ljung K,
Harmer SL (2009) REVEILLE1, a Myb-like transcription factor, integrates the circadian
clock and auxin pathways. Proc Natl Acad Sci U S A **106**: 16883–8

Ried MK, Wild R, Zhu J, Pipercevic J, Sturm K, Broger L, Harmel RK, Abriata LA,
Hothorn LA, Fiedler D, et al (2021) Inositol pyrophosphates promote the interaction of
SPX domains with the coiled-coil motif of PHR transcription factors to regulate plant
phosphate homeostasis. Nat Commun **12**: 384

Riemer E, Qiu D, Laha D, Harmel RK, Gaugler P, Gaugler V, Frei M, Hajirezaei M-R,
Laha NP, Krusenbaum L, et al (2021) ITPK1 is an InsP6/ADP phosphotransferase that
controls phosphate signaling in Arabidopsis. Mol Plant **14**: 1864–1880

Robinson S, Giersch C (1987) Inorganic Phosphate Concentration in the Stroma of Isolated
Chloroplasts and Its Influence on Photosynthesis. Aust J Plant Physiol **14**: 451

van Rooijen R, Harbinson J, Aarts MGM (2018) Photosynthetic response to increased
irradiance correlates to variation in transcriptional response of lipid-remodeling and heat-
shock genes. Plant direct **2**: e00069

Rubio V, Linhares F, Solano R, Martín AC, Iglesias J, Leyva A, Paz-Ares J (2001) A
conserved MYB transcription factor involved in phosphate starvation signaling both in
vascular plants and in unicellular algae. Genes Dev **15**: 2122–33

Sakuraba Y, Kanno S, Mabuchi A, Monda K, Iba K, Yanagisawa S (2018) A phytochrome-
B-mediated regulatory mechanism of phosphorus acquisition. Nat plants **4**: 1089–1101

Scheible W-R, Pant P, Pant BD, Krom N, Allen RD, Mysore KS (2023) Elucidating the
unknown transcriptional responses and PHR1 mediated biotic and abiotic stress tolerance
during phosphorus-limitation. J Exp Bot **11**: erad009

Schmitz J, Schöttler MA, Krueger S, Geimer S, Schneider A, Kleine T, Leister D, Bell K,
Flügge U-I, Häusler RE (2012) Defects in leaf carbohydrate metabolism compromise
acclimation to high light and lead to a high chlorophyll fluorescence phenotype in
Arabidopsis thaliana. BMC Plant Biol **12**: 8

Schneider A, Häusler RE, Kolukisaoglu Ü, Kunze R, Van der Graaff E, Schwacke R,
Catoni E, Desimone M, Flügge UI (2002) An Arabidopsis thaliana knock-out mutant of
the chloroplast triose phosphate/phosphate translocator is severely compromised only
when starch synthesis, but not starch mobilisation is abolished. Plant J **32**: 685–699

Schreiber U, Schliwa U, Bilger W (1986) Continuous recording of photochemical and non-
photochemical chlorophyll fluorescence quenching with a new type of modulation
fluorometer. Photosynth Res **10**: 51–62

Sivak MN, Walker DA (1986) Photosynthesis in vivo can be limited by phosphate supply.
New Phytol **102**: 499–512

Smith AM, Stitt M (2007) Coordination of carbon supply and plant growth. Plant Cell Environ
30: 1126–49

Stitt M, Grosse H (1988) Interactions between Sucrose Synthesis and CO2 Fixation IV.
Temperature-dependent adjustment of the relation between sucrose synthesis and CO2
fixation. J Plant Physiol **133**: 392–400

Stitt M, Wirtz W, Heldt HW (1980) Metabolite levels during induction in the chloroplast and
extrachloroplast compartments of spinach protoplasts. Biochim Biophys Acta **593**: 85–
102

Stitt M, Wirtz W, Heldt HW (1983) Regulation of Sucrose Synthesis by Cytoplasmic
Fructosebisphosphatase and Sucrose Phosphate Synthase during Photosynthesis in
Varying Light and Carbon Dioxide. Plant Physiol **72**: 767–74

Sun L, Song L, Zhang Y, Zheng Z, Liu D (2016) Arabidopsis PHL2 and PHR1 Act
Redundantly as the Key Components of the Central Regulatory System Controlling
Transcriptional Responses to Phosphate Starvation. Plant Physiol **170**: 499–514

Suzuki N, Devireddy AR, Inupakutika MA, Baxter A, Miller G, Song L, Shulaev E, Azad
RK, Shulaev V, Mittler R (2015) Ultra-fast alterations in mRNA levels uncover multiple
players in light stress acclimation in plants. Plant J **84**: 760–72

Thibaud M-C, Arrighi J-F, Bayle V, Chiarenza S, Creff A, Bustos R, Paz-Ares J, Poirier
Y, Nussaume L (2010) Dissection of local and systemic transcriptional responses to
phosphate starvation in Arabidopsis. Plant J **64**: 775–89

Vandenbussche F, Habricot Y, Condiff AS, Maldiney R, Van Der Straeten D, Ahmad M
(2007) HY5 is a point of convergence between cryptochrome and cytokinin signalling
pathways in Arabidopsis thaliana. Plant J **49**: 428–441

Vogel MO, Moore M, König K, Pecher P, Alsharafa K, Lee J, Dietz KJ (2014) Fast
retrograde signaling in response to high light involves metabolite export, MITOGEN-
ACTIVATED PROTEIN KINASE6, and AP2/ERF transcription factors in Arabidopsis.

Plant Cell **26**: 1151–1165

Wahl V, Ponnu J, Schlereth A, Arrivault S, Langenecker T, Franke A, Feil R, Lunn JE,
Stitt M, Schmid M (2013) Regulation of flowering by trehalose-6-phosphate signaling in
Arabidopsis thaliana. Science **339**: 704–7

Wang Z, Zheng Z, Song L, Liu D (2018a) Functional Characterization of Arabidopsis PHL4
in Plant Response to Phosphate Starvation. Front Plant Sci **9**: 1432

Wang ZQ, Zhou X, Dong L, Guo J, Chen Y, Zhang Y, Wu L, Xu M (2018b) iTRAQ-based
analysis of the Arabidopsis proteome reveals insights into the potential mechanisms of
anthocyanin accumulation regulation in response to phosphate deficiency. J Proteomics
184: 39–53

Ward JT, Lahner B, Yakubova E, Salt DE, Raghothama KG (2008) The effect of iron on
the primary root elongation of Arabidopsis during phosphate deficiency. Plant Physiol
147: 1181–91

Weise SE, Liu T, Childs KL, Preiser AL, Katulski HM, Perrin-Porzondek C, Sharkey TD
(2019) Transcriptional Regulation of the Glucose-6-Phosphate/Phosphate Translocator 2
Is Related to Carbon Exchange Across the Chloroplast Envelope. Front Plant Sci **10**: 827

Woodrow IE, Raymond Ellis J, Jellings A, Foyer CH (1984) Compartmentation and fluxes
of inorganic phosphate in photosynthetic cells. Planta **161**: 525–30

Wu FH, Shen SC, Lee LY, Lee SH, Chan MT, Lin CS (2009) Tape-arabidopsis sandwich -
A simpler arabidopsis protoplast isolation method. Plant Methods **5**: 1–10

Xu L, Zhao H, Wan R, Liu Y, Xu Z, Tian W, Ruan W, Wang F, Deng M, Wang J, et al
(2019) Identification of vacuolar phosphate efflux transporters in land plants. Nat plants
5: 84–94

Yu L, Fan J, Zhou C, Xu C (2021) Chloroplast lipid biosynthesis is fine-tuned to thylakoid
membrane remodeling during light acclimation. Plant Physiol **185**: 94–107

Zhou J, Jiao F, Wu Z, Li Y, Wang X, He X, Zhong W, Wu P (2008) OsPHR2 is involved in
phosphate-starvation signaling and excessive phosphate accumulation in shoots of plants.
Plant Physiol **146**: 1673–86

Zhu J, Lau K, Puschmann R, Harmel RK, Zhang Y, Pries V, Gaugler P, Broger L, Dutta
AK, Jessen HJ, et al (2019) Two bifunctional inositol pyrophosphate
kinases/phosphatases control plant phosphate homeostasis. Elife **8**: 1–25

Zirngibl M-E, Araguirang GE, Kitashova A, Jahnke K, Rolka T, Kühn C, Nägele T,

Richter AS (2023) Triose phosphate export from chloroplasts and cellular sugar content

regulate anthocyanin biosynthesis during high light acclimation. Plant Commun **4**: 100423

Figure 1 P_i starvation responses are affected by photoassimilate partitioning. **A-C,** Seedlings of wildtype (Col-0), *adg1-1*, *tpt-2*, *phr1-3 phl1*, *adg1 tpt-2* and *adg1 phr1-3 phl1* (**TM**) genotype were grown for 7 (A), 5 (B), or 10 (C) days on rich medium including 0.5 % sucrose before transfer to media with 0.5 % sucrose and either 2.5 mM (**Control**) or 0 mM (**P_i depletion**) KH_2PO_4 added. Growth was continued for 8 (A), or 7 (B, C) days. **A,** Seedling phenotypes under P_i deficient growth conditions. Bar, 2 cm. **B,** Anthocyanin contents of seedling shoots. Pigment levels are depicted as units of absorbance (A535 - A630) relative to shoot fresh weights. Bars represent means ± standard deviations; *n* = 5-6 pools of seedling shoots from 3 independent experiments; **C,** Contents of glucose, fructose, sucrose and starch. Pools of seedling shoots were harvested 10.25 hours after onset of the 16-hours photoperiod.

945   Means ± standard deviations; $n = 3$ independent experiments. **B, C,** Statistical analyses were

946   performed with One-way ANOVA with Tukey HSD follow-up test and Bonferroni alpha

947   correction for contrasts; $P < 0.05$.

Figure 2 PHR1-target genes are rapidly induced by high light. qRT-PCR analyses of indicated
transcripts in rosette leaves from wildtype (Col-0) and *phr1-1 phl1* mutants. Plants were grown
on soil for 5 weeks in an 8-hours light regime at 80 ± 5 µmol m^{-2} s^{-1} before light intensity was

set to 320 ± 30 µmol m^{-2} s^{-1} (High light, yellow bars) starting 1 hour after onset of the

photoperiod on day 36. The control group was kept at 80 ± 5 µmol m^{-2} s^{-1} (blue bars). Per sample,

2-3 fully developed leaves (leaves no. 10-13) were harvested at the indicated treatment times.

Transcript levels were calculated relative to *PP2A* as $1000\cdot2^{-\Delta CT}$. Bars represent means

± standard deviations, *n* = 3 independent experiments. Statistical analyses were performed

using Student's *t* test with 2-tailed distribution, unpaired with unequal variance. Significant

differences are indicated as ***P <0.001, **P <0.01, *P <0.05, ns, not significant.

Figure 3 High-light induced metabolic changes in *phr1-1 phl1*. **A-D,** Levels of the indicated metabolites in leaves of wildtype (Col-0, light grey bars) and *phr1-1 phl1* mutants (dark grey bars) are depicted. Plants were grown and treated as described in Figure 2. Fully expanded leaves were sampled after 90 min of exposure to a photon flux density of either 80±5 µmol m^{-2} s^{-1} (**Control**) or 320±30 µmol m^{-2} s^{-1} (**HL**). Bars show means ± standard deviations; statistical analyses were performed with Student's *t* test with two-tailed distribution, unpaired with unequal variance. Significant differences are indicated as *** P <0.001, ** P <0.01, * P <0.05, ns not significant. **A,** $n = 6$ plants from 3 independent experiments; significant differences indicated by letter code: P <0.05. **B, C,** $n = 11-12$ plants from 4 independent experiments. **C, D,** Fold changes of sugar/starch contents were calculated for each genotype as HL/mean of control. **Gluc** glucose; **Fruc** fructose; **Suc** sucrose. **D,** $n = 9$ plants from 3 independent experiments. **E,** Anthocyanin levels determined in leaves of the indicated genotypes after 3 days of high-light exposure. Plants were grown for 32 days under low light conditions (70±10 µmol m^{-2} s^{-1}) in an 8-hours light regime. Subsequently, light intensity was increased for the treatment group (High light, yellow bars) to 450±30 µmol m^{-2} s^{-1}, while the

977   control group (blue bars) was kept under growth light conditions. Leaf material was harvested

978   after additional 70 hours. Pigment contents are expressed as absorbance units (A535 - A630)

979   relative to shoot fresh weight. $n = 3$ independent experiments; statistical analyses were

980   conducted using One-way ANOVA with Tukey HSD follow-up test and Bonferroni alpha

981   correction for contrasts, P <0.01; ns, not significant.

Figure 4 Effect of high light on chlorophyll fluorescence in the *phr1-1 phl1* mutant. Plants of wildtype (Col-0) and *phr1-1 phl1* genotype were grown as described in Figure 2. After 5 weeks of growth, the light intensity was increased to 320 ± 30 µmol m^{-2} s^{-1} for high-light (**HL**) treatments, whereas the control group was kept under growth light conditions (80 ± 5 µmol m^{-2} s^{-1}). $n = 3$ independent experiments (representing means of each 3 leaves measured). Significance was tested using One-way ANOVA with Tukey HSD follow-up test and Bonferroni alpha correction for contrasts; ns, not significant. **A,** PAM measurements were performed to determine the quantum yield of photosystem II photochemistry (ΦPSII) on light-adapted leaves after 20, 45, and 125 min of treatment. **B,** Plants subjected to high-light stress were dark-adapted for 20 min after 125 min of stress duration to determine the maximum efficiency of photosystem II (F_v/F_m). Bars show means $\pm$ standard deviations; $P < 0.01$.

Figure 5 Carbohydrate metabolism is affected by *phr1 phl1* mutation. **A-D,** Phenotypes of the

adg1 phr1 phl1 triple mutant (**TM**). **A-C,** Plants were grown at 23°C and a light intensity of

$100\pm10\,\mu mol\,m^{-2}\,s^{-1}$. **A,** Rosette habitus of plants grown under short-day (8 hours light period)

and long-day (16 h light period) conditions. Pictures were taken of representative plants after

27 (short day) and 25 (long day) days of growth. The arrow indicates a senescent leaf. Bar,

2.5 cm. **B,** Leaves of 42 day-old plants grown under an 8-hours light regime were aligned

according to age (starting with the cotyledons on the left hand side). **C,** Inflorescences of

wildtype (Col-0), *adg1-1*, *phr1-3 phl1*, and derived genotypes showing delay of the transition

to flowering in the TM. Pictures were taken of representative plants after 42 days of growth

under a 16-hours light regiment. **B, C,** Backgrounds were manually removed for better

visualization. **D**, High-light sensitivity of TM and growth promotion by exogenous sucrose.

Seedlings were grown in long-day conditions under low light (50 ± 5 µmol m^{-2} s^{-1}) or high light

(320 ± 30 µmol m^{-2} s^{-1}) on rich media with or without sucrose added. Media-containing square

petri dishes were positioned horizontally. Pictures were taken after 14 days of growth. **E, F,**

Carbohydrate responsive gene expression is affected under P_i starvation. qRT-PCR analysis of

carbohydrate-responsive transcripts which are downregulated in shoots of the wildtype (E), or

upregulated in shoots of *phr1-1 phl1* mutant seedlings (F) grown for 7 days on rich medium

containing 0.5 % sucrose, and for additional 8 days on media with 0.5 % sucrose and either

2.5 mM (control, light grey bars), or 0 mM (P_i depletion, blue bars) KH_2PO_4 added. Transcript

levels were calculated relative to the level of *PP2A* transcript as $1000\cdot2^{-\Delta CT}$. Bars represent

means $\pm$ standard deviations; $n = 3$ independent experiments. One-way ANOVA with Tukey

HSD follow-up test and Bonferroni alpha correction for contrasts; ** P <0.01, * P <0.05, ns,

no significant difference.

1019

Figure 6 Constitutive P_i starvation transcription in rosette leaves of *adg1 tpt-2*. **A, B,** Plants were grown on soil at a photon flux density of 90 ± 10 µmol m^{-2} s^{-1} and sampled 135 min after onset of the 8-hours photoperiod after 5 weeks of growth. **A,** qRT-PCR analysis of the transcript levels of P_i starvation marker genes in rosette leaves of wildtype (Col-0), *adg1-1*, *tpt-2*, and *adg1 tpt-2*. Transcript levels are $1000 \cdot 2^{-\Delta CT}$ relative to *PP2A*. Bars show means ± standard deviations. $n = 3$ independent experiments. One-way ANOVA with Tukey HSD follow-up test and Bonferroni alpha correction for contrasts; $P < 0.01$. **B,** P_i contents of rosette leaves. Means ± standard deviations are depicted. $n = 3$ independent experiments. Kruskal-Wallis test with DUNN's follow-up test and DUNN/Sidak alpha correction for contrasts; $P < 0.01$.

1029

1030

Figure 7 Model for PHR1 activation during imbalances in triose phosphate production and
consumption. In wildtype mesophyll cells subjected to an increase in light intensity (left panel),
triose phosphate (TP) production by the Calvin-Benson-Bassham cycle transiently exceeds the
rates of TP utilization by starch and sucrose biosynthesis causing sequestration of free P_i.
Increased export of TPs to the cytosol via the TPT leads to a decrease in P_i concentration in the
cytosol which results in PHR1 activation. High PHR1 activity facilitates the mobilization of P_i
from internal sources including the stimulation of starch accumulation, and, together with a
signal derived from TPs or 3-phosphoglycerate (3-PGA) accumulation, PHR1 triggers
anthocyanin production. In the *adg1-1* knockout (middle panel), TPs accumulate in the cytosol
due to the lack of starch biosynthesis. High TP/3-PGA levels in the cytosol promote
anthocyanin production and sucrose biosynthesis. High sucrose levels enhance P_i-starvation
responses (Lei et al., 2011), while TP turnover confined to the cytosol keeps PHR1 activation
at a moderate level. In *adg1 tpt-2* (right hand side panel), TP export relies on the xylulose
5-phosphate/P_i translocator (XPT) (Hilgers et al., 2018), keeping sucrose biosynthesis at low
levels (Schmitz et al., 2012). TPs are partially consumed by glycolysis and imported back into
the chloroplast by the phosphoenolpyruvate (PEP)/P_i translocator (PPT). Another portion of
TPs and PEP is transferred to the vacuole where P_i is released by phosphatase activity.
However, overall release of free P_i in the cytosol is low thus keeping PHR1 activity at high
levels.