Title: Intragenic DNA inversions expand bacterial coding capacity

Authors: Rachael B. Chanin[1]†, Patrick T. West[1]†, Ryan M. Park[1], Jakob Wirbel[1], Gabriella Z.

M. Green[1], Arjun M. Miklos[1], Matthew O. Gill[2], Angela S. Hickey[2], Erin F. Brooks[1], Ami S.

Bhatt[1,2]*

Affiliations:

† These authors contributed equally to this work, listed alphabetically by last name

[1]Department of Medicine (Hematology, Blood and Marrow Transplantation); Stanford, USA.

[2]Department of Genetics, Stanford University; Stanford, USA.

*Corresponding author. Email: asbhatt@stanford.edu

Abstract: Bacterial populations that originate from a single bacterium are not strictly clonal.

Often, they contain subgroups with distinct phenotypes. Bacteria can generate heterogeneity

through phase variation: a preprogrammed, reversible mechanism that alters gene expression

levels across a population. One well studied type of phase variation involves enzyme-mediated

inversion of specific intergenic regions of genomic DNA. Frequently, these DNA inversions flip

the orientation of promoters, turning ON or OFF adjacent coding regions within otherwise

isogenic populations. Through this mechanism, inversion can affect fitness, survival, or group

dynamics. Here, we develop and apply bioinformatic approaches to discover thousands of

previously undescribed phase-variable regions in prokaryotes using long-read datasets. We

identify 'intragenic invertons', a surprising new class of invertible elements found entirely within

genes, in bacteria and archaea. To date, inversions within single genes have not been described.

Intragenic invertons allow a gene to encode two or more versions of a protein by flipping a DNA

sequence within the coding region, thereby increasing coding capacity without increasing

genome size. We experimentally characterize specific intragenic invertons in the gut commensal

Bacteroides thetaiotaomicron, presenting a 'roadmap' for investigating this new gene-

diversifying phenomenon.

One-Sentence Summary: Intragenic DNA inversions, identified using long-read sequencing

datasets, are found in many phyla across the prokaryotic tree of life.

Introduction

Adaptation is a cornerstone of survival for any species. In the complex gut microenvironment, bacteria experience many stressors including nutritional and niche competition, oxidative and nitrosative stress, and antibiotics. To overcome these challenges, bacteria may activate specific response programs which alter transcriptional or translational profiles promoting survival during these conditions. Additionally, bacterial daughter cells may acquire mutations, such as single nucleotide variations or small insertions or deletions, within genes. These gene alterations can then promote survival in the right circumstances. For example, mutations in drug targets, efflux pumps, or their regulators can provide increased resistance to antibiotics [1,2]. While many of these gene-varying mutations in bacteria are semi-reproducible, meaning that nucleotide alterations will occur in the same region of a genome under a similar environmental stressor, most are not reversible and may be costly when the stimulus is removed [3].

Beyond mutations and small insertions and deletions, there are only a few known mechanisms for introducing gene variation in bacteria. These mechanisms include: alternative translational start sites or terminators, which enable the encoding of two or more different gene products from a single mRNA [4,5]; slipped-strand mispairing, which introduces replicative or translational changes that can alter bacterial gene sequence length [6]; and diversity generating retroelements [7], which can diversify a gene during reverse transcription and recombine in a novel gene variant. Outside of these rare gene-varying events, the typical prokaryotic 'one gene, one gene product' rule generally holds and is in stark contrast to nearly all Eukaryotes, in which a large proportion of transcribed genes can undergo alternative splicing to generate multiple protein isoforms from one gene.

One fairly prevalent mechanism of reversible adaptation in bacteria is phase variation. This is a preprogrammed and reversible mechanism that generates phenotypic diversity in a clonal population [8]. Phase variation can promote cooperativity by sharing resources between subgroups in a metabolically efficient way or through bet hedging by diversifying a population to protect from complete elimination in future selective events. One type of preprogrammed variation occurs through DNA inversion. Site-specific recombinases recognize a pair of inverted repeats in genomic DNA and invert the intervening DNA sequence [9]. In the first described

example, DNA inversion of a promoter sequence resulted in the switching of expression from

one flagellar antigen (H1) to another (H2) in *Salmonella enterica* serovar Typhimurium. The

change in antigen expression determined whether the bacterium was bound by antiserum, and

thus was termed a 'phase-determining event'[10–12]. This DNA inversion, and the many others that

have since been discovered, play critical adaptive roles in both commensal and pathogenic

bacteria.

For decades, these invertible loci were identified individually. Then, computational

approaches enabled higher-throughput discovery of these 'invertons' across the genomes of a

small subset of specific bacterial species [13,14]. In 2019, Jiang *et al.* developed an elegant method

that facilitated broad scale identification of 4,686 intergenic invertons (i.e., invertons between

genes) through a search of 54,875 bacterial reference genomes [15], utilizing short-read mapping as

evidence; however, short-reads cannot span entire invertons, which can range in lengths of up to

multiple kbps [15]. The long length of many of these invertons causes short-read based inverton

detection methods to be lower in sensitivity, as methods to detect invertons rely on reads that

span one boundary of a given inverton. This means that only a small proportion of reads provide

usable evidence for inverton detection. Similarly to Jiang *et al.*, in early 2023, Milman *et al.* used

a computational model to predict over 11,000 potential invertons that partially overlap with

genes (partial intergenic) in >35,000 bacterial species. Partial intergenic invertons are sometimes

referred to as shufflon systems; they function by flipping out homologous domains of enzymes,

which can change their specificity [16,17]. Once Milman *et al.* predicted the candidate invertons,

they then manually inspected publicly available long-read datasets, which led to the validation of

22 of the >11,000 predicted invertons [18]. Taken together, these two studies demonstrate that

computational approaches can be a powerful method to identify invertible elements within many

phyla. Furthermore, the ubiquity of both intergenic and shufflon-type invertons in bacterial

genomes highlights their likely importance in affecting bacterial gene regulation and phenotypes.

While previous work has demonstrated the presence of intergenic and shufflon-type

invertons, there are no reports to our knowledge of invertons that occur entirely within genes.

Such invertons would represent a novel mechanism of preprogrammed gene variation in bacteria.

Furthermore, with the advent of long-read sequencing technologies, and improvements in their

accuracy, developing a long-read inverton finding workflow would be expected to improve

sensitivity in inverton discovery and detection. Building off these concepts, here we find that the

same mechanisms that underlie intergenic and partially intergenic invertons can occur entirely within a gene. These intragenic invertons expand bacterial coding capacity by either recoding protein sequences within the inverted region or introducing premature stop codons. In both cases, intragenic invertons result in a single gene being able to produce two or more different protein products. We develop PhaVa, a long-read based tool to identify intragenic, intergenic, and partial intergenic invertons. By applying PhaVa to long read sequencing data for ~30,000 bacterial isolates from ~4,000 unique species, we find that intragenic invertons occur in many phyla across the prokaryotic tree of life. In particular, we focus on *Bacteroides thetaiotaomicron*, a model enteric commensal, and validate 10 intragenic invertons experimentally with particular focus on the inverton contained within the thiamine biosynthesis protein *thiC*. Finally, we make both the PhaVa software package and all of the identified invertons (intragenic, intergenic, and partial intergenic) publicly available.

Results

Most knowledge regarding bacterial genes and their regulation is based on bacteria that are studied in laboratory conditions. Because of this, invertons that provide a fitness advantage *in vivo* but may not be advantageous to fitness *in vitro* have likely been overlooked [19–21]. We therefore hypothesized that there are currently unknown gut-relevant invertons. To test our hypothesis, we endeavored to identify invertons in metagenomic sequencing data from longitudinally collected human stool samples from 149 adult and 21 pediatric patients undergoing hematopoietic cell transplantation [22,23] (Fig. 1A). These samples were selected given the varying and complex environments enteric bacteria would encounter over time, with many different stressors present such as chemotherapy, antibiotic treatment, variation in food intake, and inflammation. We hypothesized that these factors might induce inverton flipping.

In our efforts to comprehensively annotate invertons from this metagenomic data set, in which there are many different organisms represented in each sample, we first decided to examine invertons in organisms within the taxon Bacteroidetes. Bacteroidetes species are prevalent and typically highly abundant in the human gut, and many organisms within this taxon have known intergenic invertons [15]. To orthogonally confirm sequencing-based observations in subsequent microbiological and genetic experiments, we focused our analysis on *B.*

thetaiotaomicron, a genetically tractable species suitable for downstream experimental

manipulation. To identify invertons in *B. thetaiotaomicron*, we used PhaseFinder [15], a short-read,

reference-based inverton detection pipeline with *B. thetaiotaomicron* VPI-5482 (BTh) as the

reference genome, and with relaxed filters to increase sensitivity (see methods, Fig. 1A). As an

internal control to assess whether PhaseFinder could sensitively detect BTh invertons in our

metagenomic samples, we examined BTh's capsular polysaccharide (CPS) genes, a known set of

invertible loci. BTh has 8 loci that encode different CPS, 5 of which are controlled by invertible

promoters [24–26]. CPS are important mediators of phage susceptibility [27] and can modulate the host

immune system [28–30]. Using PhaseFinder on the patient sample datasets, we found read evidence

of all 5 CPS invertons in both the reference and inverted (flipped) orientations (fig. S1),

demonstrating that PhaseFinder is able to detect known invertons in these metagenomic samples

and that these samples have enough *Bacteroides* sequencing depth to identify invertons. Of note,

in the reference BTh genome, the invertible promoter for each of these 5 loci is in the 'OFF'

state by virtue of it being oriented in the opposite direction of the CPS genes. Similarly, *in vitro*

transcriptional analyses support the finding that the majority of invertible CPS loci are in the

OFF orientation [31], suggesting that in laboratory conditions, these loci are not transcriptionally

active. Finding read evidence of inversion for all invertible CPS loci suggests that the *in vivo*

patient datasets are an ideal environment to detect invertible events that are rare in laboratory-

grown bacteria but may be prevalent in bacteria living in more 'natural' ecological settings.

In addition to known intergenic invertons such as those in the CPS loci, we also found

read evidence of intragenic invertons in BTh across 132 short-read metagenomic samples (Fig.

1B). We use the term 'intragenic inverton' to describe invertible regions found entirely within

single genes. To date, the only description of invertible DNA sequences entirely within a gene

are in isolated cases of very short (7 bp) flips within mitochondrial DNA in certain pathogenic

states [32]. These 7 bp mitochondrial DNA flips are postulated to be the consequence of an

enzyme-independent event, and thus are different from what we predict here to be an invertase-

mediated, preprogrammed inversion. In the intragenic invertons that we observed, there were

two predicted consequences. In some cases, the intragenic inverton resulted in a portion of the

protein being "re-coded" (Fig. 1C). For example, we observed a 57 bp inversion in BT0375, the

invertase that is believed to flip the adjacent CPS1 invertible promoter. This intragenic inversion

changes the amino acid sequence of the 'flipped' region, and might alter the binding specificity

of the invertase, possibly changing the invertible repeats (IRs) that it targets for flipping or its

binding affinity for its cognate IRs. In other cases, the intragenic inverton resulted in the

introduction of a 'premature' stop codon, affecting the prediction of protein coding open reading

frames (ORFs). Often, inversion resulted in two predicted ORFs (called with Prodigal [33]). For

example, the inverton in the hybrid two-component system BT3786 occurs between two

predicted protein folding domains, and thus might untether the "sensing" and "response"

elements (Fig. 1D) of this signaling protein. However, we also observed intragenic inversions

that resulted in zero, one, three, or more ORFs. Taken together, we describe the discovery of

intragenic invertons and identify two types of invertons - those that are 'recoding' and those that

cause a 'premature stop'.

To validate these predicted intragenic invertons, we analyzed the DNA sequences in

these gene regions *in vitro*. We extracted DNA from wild-type BTh grown in either rich or

defined media and designed PCR primer sets that enabled us to amplify either the reference or

the inverted version (fig. S2). We tested 59 of the 63 predicted intragenic invertons and

confirmed that 10 of them had DNA molecules in both the reference and inverted orientation in

our laboratory-grown population of BTh (Fig. 1B, Table 1, Data S1). The 49 unconfirmed

intragenic invertons may be due to the absence of cues in the growth conditions required to flip

the locus to the inverted orientation and/or false positives from the metagenomic read-based

evidence.

As genomic structural variation often involves highly repetitive or low complexity

regions, short-reads are often not long enough to resolve these sequences [34], and thus short-read

based approaches would be predicted to have limited sensitivity. We, therefore, developed a

long-read based inverton predictor, PhaVa. PhaVa maps long-reads against both a forward

(identical to reference) and reverse orientation version of potential invertons (Fig. 2A). PhaVa's

accuracy is high because it requires long-reads that span the entire length of a given inverton in

order to make a 'call' about its orientation. To ensure accurate performance of PhaVa, we

optimized read mapping parameters by simulating long-read datasets from ten bacterial genomes

at various sequencing depths (Fig. 2B-C). The reads were generated from a reference genome,

and thus no invertons are expected and any detected would be false positives. In general, the

false positive rate was very low (mean false positive count per simulated sample of <0.1 in 9/10

species), with the exception of reads simulated from the *Bordetella pertussis* genome (Fig. 2D).

Further investigation revealed the false positives detected in *B. pertussis* were due to a single

putative inverton with inverted repeats longer than 750 bps, of which only a smaller portion of

the total length were detected by 'einverted', the computational tool used to detected inverted

repeats (fig. S3). In summary, our long-read based inverton predictor, PhaVa, demonstrates high

accuracy in resolving complex genomic structural variations, with only rare instances of false

positives observed.

To find invertons across prokaryotic genomes, we ran PhaVa on ~30,000 prokaryotic

isolate long-read datasets deposited on SRA. We limited our analysis to readsets belonging to

Bacteria or Archaea and with 50 Mbp or more of total sequencing, which resulted in our final

analysis containing results from ~4,000 unique species (fig. S4). The vast majority of these

datasets represented bacteria, with only 42 archaeal long-read sequencing datasets. In total, we

identified 4622 unique invertons, 3,468 of which are intergenic. Of note, compared to Jiang *et al.*

[15], we find invertons at a higher rate per sequencing dataset (0.15 vs 0.07) and per individual

isolate (1.15 v 0.09) highlighting the increased sensitivity of long-reads for detecting this type of

structural variation. Like Jiang *et al.* [15], we found that Bacteroidetes have a relatively large

number of intergenic invertons (673, Fig. 3A) and intergenic invertons per genome (2.26, Fig.

3B). Fusobacteria, Gammaproteobacteria, and Verrucomicrobia also have high numbers of

intergenic invertons per genome (Fig. 3B), with Verrucomicrobia having the highest number per

genome overall at 5.55 intergenic invertons per genome. In our dataset, Verrucomicrobia is

composed of only *Akkermansia* strains. As increases in *Akkermansia* abundance correlate with

protection against metabolic disease [35,36], there is interest in its use as a probiotic. However,

Akkermansia strains exhibit broad phenotypic diversity and differential gut colonization ability

[37], which may be attributable, in part, to the orientation of these varied intergenic invertons. In

addition to the intergenic invertons, we also identified 733 partial intergenic invertons (Fig. 3A).

Many of these partial intergenic invertons may form shufflon systems, and thus, as expected,

these invertons are significantly longer than intergenic or intragenic invertons (Fig. 3C, p=7.1e-

293 and p=7.6e-67 with a t-test, respectively). This finding of 733 partial intergenic invertons

adds to the 22 long-read-validated intergenic invertons recently reported by Milman *et al*. Thus,

our analysis of ~30,000 prokaryotic isolate long-read datasets from diverse species uncovered

both known and novel intergenic and partial intergenic invertons, shedding light on the

remarkable structural variability within prokaryotic genomes and emphasizing the heightened

sensitivity of long-read sequencing in this context.

Beyond intergenic and partial intergenic invertons, we also found evidence of intragenic

invertons across multiple phyla, including the major gut microbiome-related phyla,

Proteobacteria, Firmicutes and Bacteroidetes (Fig. 3A-B). We found the largest number of

intragenic invertons, 118, in Gammaproteobacteria, including from organisms such as

Escherichia coli and *Salmonella*; this is largely due to the abundance of samples for these

organisms in SRA and our dataset (~4,000 *E. coli* samples (fig. S4)), given that

Gammaproteobacteria have a relatively small number of intragenic invertons detected per

genome (0.09, Fig. 3B). Few long-read datasets for Archaea were available with 36 and 6 for

Euryarchaeota and Crenarchaeota, respectively. Despite this, 12 putative archaeal invertons were

identified; ten intergenic, one partial intergenic, and an intragenic inverton that introduces an

early stop in a adenylosuccinate synthase gene in *Salarchaeum sp. JOR-1* (42 total archaeal

genomes searched, Fig. 3A-B, Data S2). Chromosomal invertons have only been minimally

investigated in Archaea. However, our study and a recent computational analysis of phase

variable Type 1 restriction modification systems by Atack *et al.* [17] suggest that inverton-mediated

phase variation may be an important, yet understudied, regulatory mechanism in this domain.

The mean number of intragenic invertons per genome varied greatly between different phyla

(Fig. 3B) with Tenericutes, Betaproteobacteria, and Actinobacteria having a relatively high

number of intragenic invertons detected per genome, at 0.19, 0.32, and 0.21, respectively. The

distribution of inversion proportions of individual intragenic invertons was different from that of

intergenic invertons (Fig. 3D); intergenic inversions typically appeared to be in either an "ON" or

"OFF" state in a given sample - suggesting that all of the organisms within that population

shared the same biological 'state' of that inverton. By contrast, intragenic invertons more

commonly had inversion proportions somewhere between 0 and 1 (Fig. 3D), indicating presence

of both the forward and reverse orientations within a given 'clonal' sample. Invertons with a

100% or near 100% proportion in the 'reverse' orientation may also represent those that can no

longer be flipped, either due to mutations in the IR or loss of the invertase that flips the inverton.

Having cataloged these intragenic invertons, we next investigated whether specific gene

types or functions were enriched for the presence of intragenic invertons by doing a clade-

resolved enrichment analysis. We calculated gene set enrichments (using Pfam clan definitions

as gene sets) per genome, species, and genus, combining the genes from all genomes in a

specific clade (Fig. 3E, S5). We found six Pfam clans enriched across several genera with the

strongest and most consistent enrichments for the Pfam clans CL0123 (Helix-turn-Helix) and

CL0219 (RNase-H-like) (fig. S5). This indicates that intragenic invertons occur more frequently

than would be expected by chance in genes that have DNA binding or DNA/RNA modifying

activity.

As noted previously, we postulate that inverton orientation likely relates to the

environment of a bacterium, and that invertons are more likely to be in the non-reference

orientation in organisms that are living in their 'natural' ecological settings. Therefore, we also

ran PhaVa on 210 *de novo* assembled long-read metagenomes from the human gut [38,39], mapping

sequencing reads back to their respective metagenomic assemblies. This enabled us to detect

invertons that may be absent in isolated bacteria grown in laboratory cultures, but present *in vivo*.

Doing so, we identified over 3,500 putative invertons, largely from contigs assigned to the phyla

Bacteroidetes and Firmicutes (fig. S6A). In keeping with our model that invertons are more

likely to be 'active' *in vivo* than *in vitro*, significantly more invertons were identified per species

in the metagenomic samples than in the isolate sequencing samples (fig. S6B). We hypothesize

this is because bacteria grown as isolates in laboratory settings do not experience the wide range

of diverse environmental conditions that they do in their natural, polymicrobial habitats. Our

analysis of the metagenomic data with PhaVa suggests that bioinformatic analysis of isolate

genomes grown in laboratory conditions likely underestimates the number and range of invertons

that exist in microbes. Therefore, the invertons called from the isolate datasets can be thought of

as a 'minimal set', as isolate conditions may not be the ideal setting to uncover phase variable

regions relative to metagenomic samples or co-cultures.

Both short-read and long-read based analyses of metagenomic datasets revealed that

intragenic invertons exist. However, the biological consequences of inversion of these invertons

to the non-reference orientation is not known. Thus, to evaluate the phenotypic consequences of

a particularly prevalent inverton, we focused on an intragenic inverton that introduces a

premature-stop codon in the BTh BT0650 gene (thiamine biosynthesis protein ThiC) (Fig. 4).

Thiamine is an essential cofactor in many cellular biochemical processes and is essential for

nearly all organisms. Some organisms, such as humans and certain gut microbes, are fully reliant

on dietary, host, or other microbial sources for vitamins or their building blocks; others, like

many gut microbes, including BTh, have the capacity to biosynthesize thiamine, albeit at a large

energetic cost [40,41]. Thus, thiamine availability has been hypothesized to strongly influence

microbial community composition [42]. We chose to characterize the intragenic inverton in *thiC* as

this gene has a defined function in thiamine biosynthesis [43,44]. Specifically, the *thiC* gene

product, which encodes the enzyme 2-methyl-4-amino-5-hydroxymethylpyrimidine phosphate

(HMP-P) kinase, catalyzes the conversion of aminoimidazole ribotide (AIR) to 4-amino-5-

hydroxymethyl-2-methylpyrimidine (HMP) and forms a key wing in thiamine biosynthesis. In

addition to having a defined role, we detected intragenic inversion in both DNA and RNA in our

laboratory grown BTh strain (Fig. 4B).We predicted that the non-reference orientation of the

inverton introduces a premature stop codon in the *thiC* mRNA, which would result in a truncated

protein containing only the N-terminal "thiC associated domain" of the protein (Fig. 4A). The

exact function of this domain of the protein is not known, but it is required for enzyme function.

As noted previously, BTh can grow in the absence of exogenous thiamine as it can synthesize

thiamine *de novo,* however, strains that lack ThiC lose this ability. We hypothesized that

inversion of the invertible locus in *thiC* would interfere with thiamine biosynthesis, and would

phenocopy the ThiC null mutant.

To test the biological consequences of inversion, we generated 'locked' versions of the

thiC inverton that prevent inversion from occurring within the gene. Traditionally, locking

elements in a specific orientation is accomplished by mutating the nucleotides in the inverted

repeat regions required for inversion or by deleting the inverted sequences entirely.

Unfortunately, for intragenic invertons, deletion of these sequences or complete mutations would

alter the corresponding amino acid sequences and confound interpretation. We therefore

exploited the wobble position of the codon to maximize mismatches between the inverted

repeats. By mutating these residues, we introduced mismatches in 6 out of 11 positions of the

inverted repeat (fig. S7A). Using this method, we created a locked forward (reference

orientation) and a locked reverse (flipped intragenic inverton) *thiC* strain. We also generated a

thiC clean deletion strain.

Next, we grew wild-type BTh, locked forward, locked reverse, and the *thiC* knockout

strain in various concentrations of thiamine (Fig. 4C). The locked forward strain phenocopied the

wild-type strain, as it was able to grow to the same optical density regardless of whether

thiamine was added to the media. By contrast, the locked reverse strain mirrored the *thiC* knock-

out strain and was only able to grow to wild-type levels when 0.1 μM or greater thiamine was

added to the media. This finding confirms our expectation that the reverse version of the

intragenic inverton interferes with ThiC function.

312   Having found that the locked reverse strain of the *thiC* intragenic inverton phenocopies

the null mutant, we wondered whether there may be physiological circumstances that favor this

mutant over the wild-type or locked forward strain. A classical approach to assess the relative

fitness of two bacterial strains is to perform a competitive growth experiment. Thus, to test

whether the inverted form of the *thiC* inverton provides a fitness advantage in different

conditions, we competed the locked forward strain against the locked reverse strain in an equal

proportion in varying concentrations of thiamine. Each strain was chromosomally marked with a

different antibiotic resistance cassette. Then we determined the competitive index (CI), which is

the ratio of recovered locked forward bacteria to recovered locked reverse bacteria (Fig. 4D). To

account for any fitness advantages conferred by the antibiotic resistance cassettes, we repeated

the competition with the cassette swapped between the two strains. While results varied slightly

between these two complementary versions of the experiment, they were generally concordant.

Specifically, we found that as thiamine concentration increases in the media the advantage

conferred by the locked forward version of *thiC* was first diminished and then abolished at

concentrations above 0.01 μM. In one version, the locked reverse significantly outcompeted the

locked forward strain at 1 and 10 μM, whereas in the other the version the reverse significantly

outcompeted the locked forward at 0.01, 0.1, and 10 μM (fig. S7 B-C). Notably, at

physiologically relevant thiamine concentrations, in the human intestine 0.02-2uM [45], the locked

reverse strain was more fit than the locked forward strain. This finding complements previous

work showing that auxotrophs have a fitness advantage in conditions containing a low level of

exogenous metabolites when competing against prototrophic strains [46]. The reversible nature of

invertons would allow a subgroup to switch between phenotypes, whereas a simple loss of

function mutation would not. Taken together, we find evidence of a physiologically relevant

condition in which an intragenic inversion within the *thiC* gene may provide an energetic or

other form of competitive advantage, and thus might be adaptive.

Discussion

Bacterial genomes densely encode functional genetic programs as well as multiple layers

of bioregulation. These layers of programming can be accessed by varying transcription,

translation, or through genomic restructuring. One mechanism of genome restructuring and

resultant 'genomic plasticity' is through enzyme-mediated DNA inversions. Such inversions can

regulate transcription of specific genetic loci through the flipping of promoter orientation

[10,15,47,48]. Furthermore, DNA inversions can also regulate shufflon systems that recombine

modular domains of bacterial protein-encoding genes to alter enzyme specificity [18,49,50]. To date,

entirely within-gene DNA inversions have not been described in prokaryotes. While not present

in every genome, the identification of enzyme-mediated intragenic inversions is important as it

represents another mechanism of genetic variation, and a way in which a single genetic locus can

encode multiple genes.

Here, we used short-read metagenomic data and a database of publicly available isolate

long-read sequencing to identify intragenic invertons in prokaryotes. In addition to using an

existing short-read inverton calling program, we developed a long-read inverton finding pipeline,

PhaVa, to more sensitively enumerate invertons. We identified intragenic invertons across the

prokaryotic tree of life, in both bacteria and the small number of archaea that we evaluated.

Using BTh as a model organism, we experimentally validated 10 intragenic invertons identified

from our short-read metagenomic analysis. We further assessed the consequence of inversion by

characterizing the phenotypic effects of an intragenic inverton found in the BTh thiamine

biosynthesis gene *thiC*. Thiamine is an essential cofactor for many central metabolic processes

and is bio-energetically costly to produce. Many microbes encode salvage, transport, and

biosynthetic pathways [51]. In BTh, thiamine acquisition and biosynthesis is highly regulated at

both the transcriptional and translational level [43,52]. Here, we find that thiamine biosynthesis also

appears to be regulated at the genomic structural level. The *thiC* intragenic inverton induces a

premature stop codon and we found that the truncated 'reverse' isoform has impaired growth in

thiamine-limited conditions. However, we also found that at physiological concentrations of

thiamine found in the human intestinal lumen, organisms encoding a locked 'reverse' isoform of

thiC have a competitive growth advantage over the locked 'forward' isoform. This supports the

presence of a novel mechanism of thiamine biosynthesis regulation and suggests a possible

ecological explanation for the existence of a 'toggle-able' switch of isoforms.

While the advantages of each identified intragenic inverton will differ depending on the

coding region affected, there are three general biological consequences of phase variation. One

reason an organism may have preprogrammed heterogeneity is to enable division of labor. By

generating subgroups within a population, members of the community may produce public goods

at a potential cost to their own fitness but for the benefit of the group as a whole [53–56]. While

altruism in bacterial interspecies relationships is often unstable, as cheaters will take advantage

of the public goods and outcompete, intraspecies altruism could bypass this as the losers in this

scenario can be repopulated by the winners [57]. The second type of heterogeneity producing

behavior is via a bet hedging strategy [58]. Diverse subgroups are generated allowing for survival

in future selection events. One classic example of this is the CPS switching that occurs in many

gut bacteria. As different CPS have varying susceptibility to phage predation, a diversified

population allows the species to persist in the presence of phage [27]. Third, bacteria of a given

species and strain may exist in various biogeographic 'niches', where neighboring bacteria, host

cells, nutrient access, and stressors might vary - thus, within a given ecological system such as

the intestinal lumen, different subcommunities of bacteria may benefit from employing different

bioregulatory programs. Future mechanistic work is needed to determine the advantages and

community structure implications of each intragenic inverton.

Although we validated the presence of the intragenic invertons in BTth, we have not

identified which invertases are responsible for each of the validated intragenic invertons that we

described. We suspect that an underlying "molecular grammar" exists and that certain invertases

recognize and flip specific sequences; specificity of invertases for a given sequence likely lie in

the inverted repeats, but might also lie within the inverted regions. In terms of how the

expression of these invertases is controlled and regulated in bacteria, phenotypic diversity is

often generated via two different mechanisms; random [57] or coordinated specialization [59]. It is

possible that invertases function at a basal level and therefore there is a baseline, low level of

inversion that occurs in a small proportion of the population. Alternatively, invertases may be

expressed in response to specific cues or signals. As BTh encodes 56 invertases, future work is

needed to identify which invertase flips these invertons and under which conditions.

Understanding how these elements are regulated and the consequences of inversion could

advance the field of synthetic biology and create new therapeutic targets.

Intragenic invertons that cause recoding mutations present an exciting opportunity to rethink gene variation. In BTh, we molecularly confirmed 2 recoding intragenic invertons (Table 1). Of note, these recoding mutations may help regulate the outer membrane of a bacterial cell potentially altering interactions with the host or other microbes. The first is BT0375, the putative CPS1 invertase. Cross regulation of CPS loci has been well described [60,61]. Changes to the invertase structure could change its binding specificity and alter which regions it flips or its kinetics. Future studies are needed to elucidate how this inversion may add another layer of regulation to which CPS loci are expressed and when they are expressed. The second is *nagA* which encodes an enzyme that catalyzes the deacetylation of N-acetylglucosamine-6-phosphate (GlcNAc-6-P) to glucosamine-6-phosphate (GlcN-6-P). NagA is important for cell wall recycling and can supply GlcN-6-P for glycolysis [62]. As NagA can be allosterically regulated [63], intragenic inversion could alter the binding of its allosteric regulator, altering this process; this might result in changing the metabolism of the cell or the outer membrane structure. Bacterial outer membranes play crucial roles in microbial interactions, niche establishment, and immune modulation. Intragenic invertons may add another layer of regulation and future studies are needed to study their effects.

While we find fairly extensive evidence of intragenic invertons using sequencing based approaches and explore some of them in detail, this work has limitations. First, our analysis of invertons across the prokaryotic tree of life was performed on previously sequenced isolates. While growth conditions for most of these samples are not readily available, we presume that most of these isolates were grown in rich laboratory media; these nutrient- and micronutrient-replete conditions may not recapitulate physiological conditions in which invertases are active or reverse orientations are favorable. However, there are currently limited long-read datasets available from physiological conditions for a wide range of prokaryotic organisms. We therefore may have only identified a minimal set of invertons in this study and we estimate the full "invertome" likely includes a larger number of elements. Second, if the invertons we identified are not representative of the true capacity for inversion, our gene set enrichment analysis may also not identify the types of genes hit most frequently in physiological conditions. Third, both PhaVa and PhaseFinder are reliant upon a reference genome or *de novo* assembly for read mapping and selection of a particular sequence for read mapping can affect inverton discovery. Detection of invertons is thus restricted to the genomic sequence common between the input

sequenced strain and the reference. Finally, PhaVa uses relatively strict mapping parameters and

if the selected reference is distantly related to the sequenced strain, read mapping quality will

decrease and reduce the discovery rate. However, using a *de novo* assembly instead may result in

missing 'fully inverted' invertons relative to reference strains, which may be of interest.

Despite these limitations, intragenic invertons are an exciting new mechanism for genetic

variation and adaptation in bacteria. In this manuscript, we present a 'roadmap' for more in depth

investigation of a specific invertible intragenic locus. Our initial analysis of long-read isolate

data provides a minimal set of invertons, including intragenic, intergenic, and partial intergenic

invertons. We expect future niche-specific investigation of inverton-containing organisms to

identify additional invertons. Additionally, we anticipate that future studies of intragenic

inverton will uncover new layers of bioregulation in prokaryotes, and more thoroughly

demonstrate the many hidden genetic programs that exist within highly plastic bacterial

genomes. More thorough characterization of invertons and other reversible and preprogrammed

types of genomic variation will likely substantially impact several fields of research ranging

from synthetic biology, to microbe-microbe interactions, to microbial physiology, and beyond.

References

1.   Hooper, D. C. & Jacoby, G. A. Mechanisms of drug resistance: quinolone resistance. *Ann.*

N. Y. Acad. Sci. **1354**, 12–31 (2015).

2.   Woodford, N. & Ellington, M. J. The emergence of antibiotic resistance by mutation. *Clin.*

Microbiol. Infect. **13**, 5–18 (2007).

3.   Björkman, J. & Andersson, D. I. The cost of antibiotic resistance from a bacterial

perspective. *Drug Resist. Updat.* **3**, 237–245 (2000).

4.   Meydan, S., Vázquez-Laslop, N. & Mankin, A. S. Genes within Genes in Bacterial

Genomes. *Microbiol Spectr* **6**, (2018).

5.   Zhong, A. *et al.* Toxic antiphage defense proteins inhibited by intragenic antitoxin proteins.

Proc. Natl. Acad. Sci. U. S. A. **120**, e2307382120 (2023).

6.   Moxon, R., Bayliss, C. & Hood, D. Bacterial contingency loci: the role of simple sequence

DNA repeats in bacterial adaptation. *Annu. Rev. Genet.* **40**, 307–333 (2006).

7. Medhekar, B. & Miller, J. F. Diversity-generating retroelements. *Curr. Opin. Microbiol.* **10**, 388–395 (2007).

8. van der Woude, M. W. & Bäumler, A. J. Phase and antigenic variation in bacteria. *Clin. Microbiol. Rev.* **17**, 581–611, table of contents (2004).

9. Trzilova, D. & Tamayo, R. Site-Specific Recombination - How Simple DNA Inversions Produce Complex Phenotypic Heterogeneity in Bacterial Populations. *Trends Genet.* **37**, 59–72 (2021).

10. Zieg, J., Silverman, M., Hilmen, M. & Simon, M. Recombinational switch for gene expression. *Science* **196**, 170–172 (1977).

11. Andrewes, F. W. Studies in group-agglutination I. The salmonella group and its antigenic structure. *J. Pathol. Bacteriol.* **25**, 505–521 (1922).

12. Stocker, B. A. Measurements of rate of mutation of flagellar antigenic phase in Salmonella typhi-murium. *J. Hyg.* **47**, 398–413 (1949).

13. Goldberg, A., Fridman, O., Ronin, I. & Balaban, N. Q. Systematic identification and quantification of phase variation in commensal and pathogenic Escherichia coli. *Genome Med.* **6**, 112 (2014).

14. Sekulovic, O. *et al.* Genome-wide detection of conservative site-specific recombination in bacteria. *PLoS Genet.* **14**, e1007332 (2018).

15. Jiang, X. *et al.* Invertible promoters mediate bacterial phase variation, antibiotic resistance, and host adaptation in the gut. *Science* **363**, 181–187 (2019).

16. Komano, T. Shufflons: multiple inversion systems and integrons. *Annu. Rev. Genet.* **33**, 171–191 (1999).

17. Atack, J. M., Guo, C., Yang, L., Zhou, Y. & Jennings, M. P. DNA sequence repeats identify numerous Type I restriction-modification systems that are potential epigenetic regulators controlling phase-variable regulons; phasevarions. *FASEB J.* **34**, 1038–1051 (2020).

18. Milman, O., Yelin, I. & Kishony, R. Systematic identification of gene-altering programmed inversions across the bacterial domain. *Nucleic Acids Res.* **51**, 553–573 (2023).

19. Chatzidaki-Livanis, M., Coyne, M. J., Roche-Hakansson, H. & Comstock, L. E. Expression of a uniquely regulated extracellular polysaccharide confers a large-capsule phenotype to Bacteroides fragilis. *J. Bacteriol.* **190**, 1020–1026 (2008).

20. Taketani, M., Donia, M. S., Jacobson, A. N., Lambris, J. D. & Fischbach, M. A. A Phase-
Variable Surface Layer from the Gut Symbiont Bacteroides thetaiotaomicron. *MBio* **6**,
e01339–15 (2015).

21. Troy, E. B., Carey, V. J., Kasper, D. L. & Comstock, L. E. Orientations of the Bacteroides
fragilis capsular polysaccharide biosynthesis locus promoters during symbiosis and
infection. *J. Bacteriol.* **192**, 5832–5836 (2010).

22. Severyn, C. J. *et al.* Microbiota dynamics in a randomized trial of gut decontamination
during allogeneic hematopoietic cell transplantation. *JCI Insight* **7**, (2022).

23. Siranosian, B. A. *et al.* Rare transmission of commensal and pathogenic bacteria in the gut
microbiome of hospitalized adults. *Nat. Commun.* **13**, 586 (2022).

24. Martens, E. C., Chiang, H. C. & Gordon, J. I. Mucosal glycan foraging enhances fitness and
transmission of a saccharolytic human gut bacterial symbiont. *Cell Host Microbe* **4**, 447–
457 (2008).

25. Martens, E. C., Roth, R., Heuser, J. E. & Gordon, J. I. Coordinate regulation of glycan
degradation and polysaccharide capsule biosynthesis by a prominent human gut symbiont.
J. Biol. Chem. **284**, 18445–18457 (2009).

26. Krinos, C. M. *et al.* Extensive surface diversity of a commensal microorganism by multiple
DNA inversions. *Nature* **414**, 555–558 (2001).

27. Porter, N. T. *et al.* Phase-variable capsular polysaccharides and lipoproteins modify
bacteriophage susceptibility in Bacteroides thetaiotaomicron. *Nat Microbiol* **5**, 1170–1181
(2020).

28. Round, J. L. *et al.* The Toll-like receptor 2 pathway establishes colonization by a
commensal of the human microbiota. *Science* **332**, 974–977 (2011).

29. Neff, C. P. *et al.* Diverse Intestinal Bacteria Contain Putative Zwitterionic Capsular
Polysaccharides with Anti-inflammatory Properties. *Cell Host Microbe* **20**, 535–547 (2016).

30. Mazmanian, S. K., Liu, C. H., Tzianabos, A. O. & Kasper, D. L. An immunomodulatory
molecule of symbiotic bacteria directs maturation of the host immune system. *Cell* **122**,
107–118 (2005).

31. Porter, N. T., Canales, P., Peterson, D. A. & Martens, E. C. A Subset of Polysaccharide
Capsules in the Human Symbiont Bacteroides thetaiotaomicron Promote Increased
Competitive Fitness in the Mouse Gut. *Cell Host Microbe* **22**, 494–506.e8 (2017).

32. Musumeci, O. *et al.* Intragenic inversion of mtDNA: a new type of pathogenic mutation in a
patient with mitochondrial myopathy. *Am. J. Hum. Genet.* **66**, 1900–1904 (2000).

33. Hyatt, D. *et al.* Prodigal: prokaryotic gene recognition and translation initiation site
identification. *BMC Bioinformatics* **11**, 119 (2010).

34. West, P. T., Chanin, R. B. & Bhatt, A. S. From genome structure to function: insights into
structural variation in microbiology. *Curr. Opin. Microbiol.* **69**, 102192 (2022).

35. Dao, M. C. *et al.* Akkermansia muciniphila and improved metabolic health during a dietary
intervention in obesity: relationship with gut microbiome richness and ecology. *Gut* **65**,
426–436 (2016).

36. Greninger, A. L. *et al.* Clinical metagenomic identification of Balamuthia mandrillaris
encephalitis and assembly of the draft genome: the continuing case for reference genome
sequencing. *Genome Med.* **7**, 113 (2015).

37. Becken, B. *et al.* Genotypic and Phenotypic Diversity among Human Isolates of
Akkermansia muciniphila. *MBio* **12**, (2021).

38. Chen, L. *et al.* Short- and long-read metagenomics expand individualized structural
variations in gut microbiomes. *Nat. Commun.* **13**, 3175 (2022).

39. Maghini, D. *et al.* Achieving quantitative and accurate measurement of the human gut
microbiome. *bioRxiv* 2022.09.28.509972 (2022) doi:10.1101/2022.09.28.509972.

40. Rodionov, D. A. *et al.* Micronutrient Requirements and Sharing Capabilities of the Human
Gut Microbiome. *Front. Microbiol.* **10**, 1316 (2019).

41. Sharma, V. *et al.* B-Vitamin Sharing Promotes Stability of Gut Microbial Communities.
Front. Microbiol. **10**, 1485 (2019).

42. Yatsunenko, T. *et al.* Human gut microbiome viewed across age and geography. *Nature*
486, 222–227 (2012).

43. Costliow, Z. A. & Degnan, P. H. Thiamine Acquisition Strategies Impact Metabolism and
Competition in the Gut Microbe Bacteroides thetaiotaomicron. *mSystems* **2**, (2017).

44. Martinez-Gomez, N. C. & Downs, D. M. ThiC is an [Fe-S] cluster protein that requires
AdoMet to generate the 4-amino-5-hydroxymethyl-2-methylpyrimidine moiety in thiamin
synthesis. *Biochemistry* **47**, 9054–9056 (2008).

45. Said, H. M. Intestinal absorption of water-soluble vitamins in health and disease. *Biochem.*
J **437**, 357–372 (2011).

46. D'Souza, G. *et al.* Less is more: selective advantages can explain the prevalent loss of
biosynthetic genes in bacteria. *Evolution* **68**, 2559–2570 (2014).

47. Abraham, J. M., Freitag, C. S., Clements, J. R. & Eisenstein, B. I. An invertible element of
DNA controls phase variation of type 1 fimbriae of Escherichia coli. *Proc. Natl. Acad. Sci.*
U. S. A. **82**, 5724–5727 (1985).

48. Weinacht, K. G. *et al.* Tyrosine site-specific recombinases mediate DNA inversions
affecting the expression of outer surface proteins of Bacteroides fragilis. *Mol. Microbiol.*
53, 1319–1330 (2004).

49. Nakayama-Imaohji, H. *et al.* Identification of the site-specific DNA invertase responsible
for the phase variation of SusC/SusD family outer membrane proteins in Bacteroides
fragilis. *J. Bacteriol.* **191**, 6003–6011 (2009).

50. Li, J.-W., Li, J., Wang, J., Li, C. & Zhang, J.-R. Molecular Mechanisms of hsdS Inversions
in the cod Locus of Streptococcus pneumoniae. *J. Bacteriol.* **201**, (2019).

51. Jurgenson, C. T., Ealick, S. E. & Begley, T. P. Biosynthesis of Thiamin Pyrophosphate.
EcoSal Plus **3**, (2009).

52. Rodionov, D. A., Vitreschak, A. G., Mironov, A. A. & Gelfand, M. S. Comparative
genomics of thiamin biosynthesis in procaryotes. New genes and regulatory mechanisms. *J.*
Biol. Chem. **277**, 48949–48959 (2002).

53. Veening, J.-W. *et al.* Transient heterogeneity in extracellular protease production by
Bacillus subtilis. *Mol. Syst. Biol.* **4**, 184 (2008).

54. Colizzi, E. S., van Dijk, B., Merks, R. M. H., Rozen, D. E. & Vroomans, R. M. A.
Evolution of genome fragility enables microbial division of labor. *Mol. Syst. Biol.* **19**,
e11353 (2023).

55. Chai, Y., Chu, F., Kolter, R. & Losick, R. Bistability and biofilm formation in Bacillus
subtilis. *Mol. Microbiol.* **67**, 254–263 (2008).

56. Marlow, V. L. *et al.* The prevalence and origin of exoprotease-producing cells in the
Bacillus subtilis biofilm. *Microbiology* **160**, 56–66 (2014).

57. Ackermann, M. *et al.* Self-destructive cooperation mediated by phenotypic noise. *Nature*
454, 987–990 (2008).

58. Grimbergen, A. J., Siebring, J., Solopova, A. & Kuipers, O. P. Microbial bet-hedging: the
power of being different. *Curr. Opin. Microbiol.* **25**, 67–72 (2015).

59.   Kalamara, M., Spacapan, M., Mandic-Mulec, I. & Stanley-Wall, N. R. Social behaviours by
Bacillus subtilis: quorum sensing, kin discrimination and beyond. *Mol. Microbiol.* **110**,
863–878 (2018).

60.   Chatzidaki-Livanis, M., Weinacht, K. G. & Comstock, L. E. Trans locus inhibitors limit
concomitant polysaccharide synthesis in the human gut symbiont Bacteroides fragilis. *Proc.*
Natl. Acad. Sci. U. S. A. **107**, 11976–11980 (2010).

61.   Chatzidaki-Livanis, M., Coyne, M. J. & Comstock, L. E. A family of transcriptional
antitermination factors necessary for synthesis of the capsular polysaccharides of
Bacteroides fragilis. *J. Bacteriol.* **191**, 7288–7295 (2009).

62.   Park, J. T. & Uehara, T. How bacteria consume their own exoskeletons (turnover and
recycling of cell wall peptidoglycan). *Microbiol. Mol. Biol. Rev.* **72**, 211–27, table of
contents (2008).

63.   White, R. J. & Pasternak, C. A. The purification and properties of N-acetylglucosamine 6-
phosphate deacetylase from Escherichia coli. *Biochem. J* **105**, 121–125 (1967).

64.   Bacic, M. K. & Smith, C. J. Laboratory maintenance and cultivation of bacteroides species.
Curr. Protoc. Microbiol. **Chapter 13**, Unit 13C.1 (2008).

65.   Zhu, W. *et al.* Xenosiderophore Utilization Promotes Bacteroides thetaiotaomicron
Resilience during Colitis. *Cell Host Microbe* **27**, 376–388.e8 (2020).

66.   Rice, P., Longden, I. & Bleasby, A. EMBOSS: the European Molecular Biology Open
Software Suite. *Trends Genet.* **16**, 276–277 (2000).

67.   Li, H. Minimap2: pairwise alignment for nucleotide sequences. *Bioinformatics* **34**, 3094–
3100 (2018).

68.   Yang, C., Chu, J., Warren, R. L. & Birol, I. NanoSim: nanopore sequence read simulator
based on statistical characterization. *Gigascience* **6**, 1–6 (2017).

69.   De Coster, W., D'Hert, S., Schultz, D. T., Cruts, M. & Van Broeckhoven, C. NanoPack:
visualizing and processing long-read sequencing data. *Bioinformatics* **34**, 2666–2669
(2018).

70.   Jumper, J. *et al.* Highly accurate protein structure prediction with AlphaFold. *Nature* **596**,
583–589 (2021).

71.   Aramaki, T. *et al.* KofamKOALA: KEGG Ortholog assignment based on profile HMM and
adaptive score threshold. *Bioinformatics* **36**, 2251–2252 (2020).

72.   Eddy, S. R. Accelerated Profile HMM Searches. *PLoS Comput. Biol.* **7**, e1002195 (2011).

73.   Paysan-Lafosse, T. *et al.* InterPro in 2022. *Nucleic Acids Res.* **51**, D418–D427 (2023).

74.   Benjamini, Y. & Hochberg, Y. Controlling the false discovery rate: A practical and

powerful approach to multiple testing. *J. R. Stat. Soc.* **57**, 289–300 (1995).

75.   Prjibelski, A., Antipov, D., Meleshko, D., Lapidus, A. & Korobeynikov, A. Using SPAdes

De Novo Assembler. *Curr. Protoc. Bioinformatics* **70**, e102 (2020).

76.   Lin, Y. *et al.* Assembly of long error-prone reads using de Bruijn graphs. *Proc. Natl. Acad.*

Sci. U. S. A. **113**, E8396–E8405 (2016).

77.   Wood, D. E., Lu, J. & Langmead, B. Improved metagenomic analysis with Kraken 2.

Genome Biol. **20**, 257 (2019).

78.   Li, W. & Godzik, A. Cd-hit: a fast program for clustering and comparing large sets of

protein or nucleotide sequences. *Bioinformatics* **22**, 1658–1659 (2006).

79.   Koropatkin, N. M., Martens, E. C., Gordon, J. I. & Smith, T. J. Starch catabolism by a

prominent human gut symbiont is directed by the recognition of amylose helices. *Structure*

16, 1105–1115 (2008).

80.   Simon, R., Priefer, U. & Pühler, A. A Broad Host Range Mobilization System for In Vivo

Genetic Engineering: Transposon Mutagenesis in Gram Negative Bacteria. *Biotechnology*

1, 784–791 (1983).

81.   Pal, D., Venkova-Canova, T., Srivastava, P. & Chattoraj, D. K. Multipartite regulation of

rctB, the replication initiator gene of Vibrio cholerae chromosome II. *J. Bacteriol.* **187**,

7167–7175 (2005).

Fig. 1. Short-read metagenomic datasets reveal intragenic invertons in *Bacteroides thetaiotaomicron* (BTh). (**A**) An overview of the analysis pipeline for identifying putative invertons in short-read datasets. (**B**) A heatmap of the inversion proportion of intragenic invertons in BTh. Samples with no intragenic invertons were removed. Rows labeled with a gene name represent intragenic invertons with PCR and Sanger sequencing evidence of inversion. (**C-D**) Genome diagrams for confirmed intragenic invertons in BTh. Gray bars indicate putative invertons without sequencing support. Red bars indicate invertons with sequencing evidence. (**C**) Left - Genome diagram of the region surrounding the BT0375 recoding intragenic inverton, and a domain diagram of the BT0375 gene with the location of the inverton IRs indicated. Right - AlphaFold overlay of the BT0375 forward (blue) pLDDT 89.91 and reverse (green) pLDDT 85.24. The region that is recoded is circled in red. (**D**) Genome diagram of the region surrounding the BT3786 premature stop codon intragenic inverton, and a domain diagram of the gene. The consequence of the inversion and resulting two predicted ORFs are indicated in the domain diagram.

Gene	Annotation	Consequence
BT0375	integrase	recoding
BT0579	putative transcription regulator	premature stop codon
BT0629	Mn2+ and Fe2+ transport protein	premature stop codon
BT0650	thiamine biosynthesis protein ThiC	premature stop codon
BT0675	N-acetylglucosamine-6-phosphate deacetylase NagA	recoding
BT1600	BexA, membrane protein	premature stop codon
BT1620	SusD homolog	premature stop codon
BT3385	putative helicase	premature stop codon
BT3786	two-component system sensor histidine kinase/response regulator, hybrid (one-component system)	premature stop codon
BT3938	ATP-dependent DNA helicase RecQ	premature stop codon

657
658 **Table 1. Confirmed intragenic invertons BTh.** Intragenic invertons confirmed *in vitro* in BTh
659 are listed. Invertons from short-read datasets were called with PhaseFinder on metagenomic
660 samples (see Fig.1). The predicted consequence of inversion is also listed.

661
Fig. 2. Developing and optimizing PhaVa, a long-read based, accurate inverton caller. (A)
Schematic of PhaVa's workflow. Putative invertons are identified, and long-reads are mapped to
both a forward (highlighted by the black dashed lines) and reverse orientation (highlighted by the
gray dashed lines) version of the inverton and surrounding genomic sequence, similar to
PhaseFinder. Reads that do not map across the entire inverton and into the flanking sequence on
either side, or have poor mapping characteristics are removed. See methods for details. **(B-C)**
Optimizing cutoffs for the minimum number of reverse reads as both a raw number and
percentage of all reads, to reduce false positive inverton calls with simulated reads. Cell color
and number represent **(B)** the false positive rate per simulated readset and **(C)** the total number of
unique false positives across all simulated datasets. **(D)** False positives in simulated data plotted
per species. All measurements were made with a minimum of three reverse reads cutoff and
varying the percentage of minimum reverse reads cutoff. Dashed line indicates the minimum
reverse reads percent cutoff used for isolate and metagenomic datasets.

Fig. 3. PhaVa analysis of isolate long-read sequencing data reveals intragenic inversions are
prevalent across the bacterial tree of life. (**A**) The total number of invertons found within
various bacterial phyla from 29,989 publicly available long-read isolate sequencing datasets.
Green bars refer to intergenic invertons. Orange bars refer to intragenic invertons. Blue bars refer
to partial intergenic invertons. Asterisks denote phyla within Archaea. Inset corresponds to the
portion of the bar graph outlined in dotted lines. (**B**) The mean number of invertons found per
genome within a phylum, of genomes that had at least one inverton. Asterisks denote phyla
within Archaea. (**C**) The distribution of lengths of identified invertons, grouped by inverton type
(intergenic, partial intergenic - denoted 'partial', and intragenic). Median value is indicated by
gray dots. Partial length distribution was found to be significantly different from intergenic
(p=0.0) and intragenic (p=4.5e-146) with a t-test (**D**) The distribution of inversion rates of
identified invertons, defined as the percentage of reads mapped in the reverse orientation.
Median value is indicated by gray dots. (**E**) Pfam clan enrichment across several genera. Dot size
and fill color is proportional to the mean log-odds ratio, an effect size measure for the
enrichment, and the length of the line indicates the fraction of included genera in which an
enrichment score for the specific clan could be calculated.
693

Fig. 4. Consequences of inversion in thiamine biosynthesis protein (*thiC*). (A) Schematic showing the location of the *thiC* intragenic inverton (red bar). Inverton flipping results in a premature stop codon located between two protein-folding domains in ThiC. Black arrows indicate the binding location of primers used to determine the orientation of inverton. **(B)** PCR confirmation of the *thiC* intragenic inverton in both genomic DNA and reverse transcribed RNA (cDNA). PCR products of the expected size were extracted and confirmed with Sanger sequencing. **(C)** BTh strains were grown in defined media with the indicated concentrations of thiamine. The maximum optical density of each strain reached was recorded. Each point

represents the average of six replicates conducted across two separate experiments. Mean and
standard deviation are shown. Locked forward (blue line), locked reverse (gray line), *thiC*
knockout (purple line), and wild-type (black line) are presented. (**D**) Locked strains were
competed against each other in thiamine-containing media. The competitive growth experiment
was performed in two different ways with the antibiotic resistance marker cassettes flipped
between the two versions. Black bars indicate the locked forward strain marked with
erythromycin resistance and locked reverse strain marked with tetracycline resistance. White
bars indicate the locked forward strain marked with tetracycline resistance and locked reverse
strain marked with erythromycin resistance. The competitive index was determined. Geometric
mean and geometric standard deviation are shown for 8-12 replicates across 4-6 independent
experiments.
714

Materials and Methods

Strains and media

The bacterial strains used in this study are listed in Table S1. *E. coli* strains were routinely grown in LB Miller media (Fisher) at 37 °C. When necessary, carbenicillin was added at 100 μg/mL. BTh was grown anaerobically (90% Nitrogen, 5% carbon dioxide, 5% hydrogen) in an anaerobic chamber (Sheldon Manufacturing) in hemin (5 μg/mL) and L-cysteine (1 mg/mL) supplemented Brain Heart Infusion (Sigma) media (BHIS) or Varel-Bryant broth (VB)[64]. When necessary, the antibiotics tetracycline (2.5 μg/mL), erythromycin (25 μg/mL), or gentamicin (200 μg/mL) were added to the media. Thiamine HCL (Sigma) was added at the specified concentrations. All media used to grow BTh was preincubated in the anaerobic chamber overnight.

Construction of *thiC* clean deletion and locked strains

The *thiC* clean deletion and locked strains were generated via allelic exchange as previously described [65]. For Δ*thiC*, 600 - 700 base pair flanking regions of the coding region were amplified using Q5 high fidelity polymerase (New England Biolabs). Recombinant DNA used in this study is listed in Table S2. For locked strains, plasmid overhangs, flanking regions, locked repeats and intervening forward or inverted sequences were synthesized (Twist Biosciences) (Data S8). Regions were assembled into pExchange-tdk using the HiFi DNA Assembly Kit (New England Biolab). Plasmid inserts were verified using Sanger sequencing (ElimBio). Sequence confirmed plasmids were propagated in *E. coli* DH5α λ*pir*. *E. coli* S17-1 λ*pir* was used as a donor strain for conjugation into BTh Δ*tdk*. Exconjugants with chromosomal integration of plasmids were recovered on BHIS plates containing gentamicin and erythromycin. Second crossover events were selected by using BHIS FudR (200 μg/mL 5-fluoro-2-deoxy-uridine). Deletion and locked versions were confirmed by PCR.

To generate differentially resistant *thiC* locked strains, the suicide vectors pNBU2_tet and pNBU2_erm were used. *E. coli* S17-1 λ*pir* harboring the plasmids were used as donor strains for

conjugation. Single crossover events were selected by plating on gentamicin plates containing
either erythromycin or tetracycline respectively.

<u>Validating inversion in DNA</u>

Intragenic inverton confirmation primers were designed using NCBI Primer Blast under default
settings with the addition of adding in a GC clamp. PCR product size was targeted to be between
300 and 600 base pairs. The common and reverse primer were oriented on the same strand of the
reference genome and the forward primer was located on the complementary strand. The
common primer was located in between the two inverted repeats (fig. S2, primers listed in Data
S6). Four of the predicted invertons were not experimentally tested, as target-specific primers
could not be generated within the above constraints.

DNA was isolated from wild-type BTh cultures grown for 18 hours in either BHIS or VB media.
DNA was isolated using a chemical and enzymatic lysis. Glass beads (0.1 mm) were added to
bacterial pellets along with 700 µl of extraction buffer (50 mM Tris pH 7.5, 1 mM EDTA, 100
mM NaCl, 1% (w/v) SDS) and 25 µl of Proteinase K (10 mg/mL). Pellets were vortexed for 20
seconds and incubated at 55ºC for 60 min. 700 µl of phenol:chloroform:isoamyl alcohol (25:24:1
by volume) was added to the mixture prior to incubating at room temperature for 5 minutes.
Phases were separated by centrifuge at 10,000 rpm for 5 minutes. The aqueous upper layer was
collected and transferred to a new tube. 5 µl of RNAse A (10 mg/mL) was added and incubated
at 37ºC for 15 minutes. An equal volume of phenol:chloroform:isoamyl alcohol was added,
mixed, and incubated at room temperature for 5 minutes. Phases were separated as above and the
aqueous phase was added to a new tube containing an equal volume of chloroform: isoamyl
alcohol (24:1 by volume). Tubes were mixed and incubated at room temperature for 5 minutes
prior to phase separation via centrifugation. The aqueous phase was added to a new tube along
with 45 µl of 3M sodium acetate and 1 mL cold 100% ethanol. DNA was precipitated overnight
at -20ºC . Pellets were washed twice with 1mL of cold 70% ethanol. Dried pellets were
resuspended in water.

PCR reactions were performed using Q5 high fidelity polymerase (68 °C annealing temp, 10
second annealing time, and 30 second extension time). PCR reactions were run on an 1.2%
agarose gel. If multiple bands were visible, bands of the expected size were gel purified using the

Qiagen Gel Extraction kit. If a single band of the expected size was observed, the PCR reaction

was purified using the Monarch PCR Cleanup Kit (New England Biolabs). DNA was sent for

Sanger sequencing. Sequencing was compared to the *in silico* prediction.

<u>Validating *thiC* inversion in RNA</u>

RNA was isolated from wild-type BTh cultures grown for 18 hours in BHIS media. 5mL cultures

were quenched using 500 μL phenol/ethanol solution (90% [vol/vol] ethanol and 10% [vol/vol]

saturated phenol pH 4-5). Pellets were spun down and stored at -80 °C until extraction. Pellets

were lysed in 250 μL PBS and 10 μL of lysozyme (10 mg/mL) at 37 °C for 30 minutes. 30 μl

20% SDS was added prior to an additional 30 minute incubation. 1.5 mL Trizol was added to the

mixture and incubated at room temperature for 10 minutes. Chloroform (0.5 mL) was added to

each sample and inverted vigorously for 15 seconds. The aqueous phase was taken from

centrifuged samples and an equal volume of 100% ethanol was added. RNA was purified using

the Zymo RNA clean kit. DNA was removed using Ambion Turbo DNAse. cDNA was made

using Taqman Reverse Transcription reagents (Invitrogen) according to the manual. A no reverse

transcriptase control was performed to ensure that all DNA was removed. PCR was performed to

determine orientation of inverton as above. Correctly sized bands were sent for Sanger

sequencing.

<u>BTh growth in thiamine concentrations</u>

BTh wild-type, *thiC* locked forward, *thiC* locked reverse, and *thiC* knockout strains were grown

overnight in BHIS media. Aliquots of each were then washed twice in preincubated PBS

containing cysteine (1 mg/mL). Strains were inoculated at an OD600 of 0.05 in VB media

containing the indicated concentration of Thiamine in a 96-well flat bottom plate. Readings were

taken in a Stratus plate reader (Cerillo) every ten minutes. Non-inoculated VB media from each

time point was used as a blank. The maximum OD600 value achieved per well was determined.

<u>Competitive growth assay</u>

Marked BT0650 locked strains were grown overnight in BHIS with appropriate antibiotics.

Strains were washed twice with preincubated PBS containing cysteine (1 mg/mL). A glass

dilution tube containing 3 mL of VB with indicated concentrations of thiamine was inoculated

with 1 x 10^3 CFU/mL of each strain. After 40 hrs of growth at 37 ℃ in the anaerobic chamber,

CFU/mL was determined by plating on selective agar. A competitive index was calculated by

dividing the recovered CFU/mL of the locked forward by the CFU/mL of the locked reverse

strain corrected by the inoculum.

<u>Identifying invertons in BTh with PhaseFinder</u>

Two short-read datasets were used for identifying invertons in BTh, 416 samples from 149 adult

HMT patients ([23], BioProject PRJNA707487) and 142 samples from 21 pediatric HMT patients

([22], BioProject PRJNA787952). Each individual short-read dataset was analyzed with

PhaseFinder [15] with the VPI-5482 reference genome and default parameters to identify putative

invertons in BTh. Invertons were included in further analysis if they had at least 5 reads mapping

to the reverse orientation of the inverton, and had reads mapping to the reverse orientation in at

least three different samples. Inverton-gene overlaps and partial overlaps were found using a

custom script, now incorporated in PhaVa in the 'Create' step, and the gene annotations from the

VPI-5482 genbank file (.gbff).

<u>The PhaVa algorithm</u>

Inverted repeats are identified with einverted, part of the EMBOSS suite [66]. For each putative

inverton, two sequences are then created: one where the sequence between identified inverted

repeat pairs is inverted (reverse) and one where it is not (forward), along with flanking sequence

on either side, similar to PhaseFinder. Long-reads are mapped against the created sequence with

minimap2 [67] and must pass several filters to be included as evidence of inversion. 1) reads must

have a MAPQ score of $\geq$ 2 to eliminate multimapping reads. 2) Reads must span the entire length

of the inverton and at least 30 bps into the flanking sequence on either side. 3) The mismatch rate

along the length of a read must be below a maximum mismatch rate. The mismatch rate is

considered separately over the length of an inverton and over flanking sequence, to avoid reads

that map well to only one region or the other. An adjustable mismatch rate is used instead of a

flat mismatch cutoff to account for both the variable length of long-reads and the high

sequencing error rate of current long-read sequencing technologies relative to short-read

sequencing. After mapping, reads mapped to the inverted and non-inverted sequences are tallied

and optional post-mapping filters are applied. 1) A minimum number of total reads mapped to

the 'reverse' sequence and 2) a minimum proportion of total mapped reads mapped to the

'reverse' sequence.

<u>Simulating long-read datasets for optimizing PhaVa</u>

For benchmarking, ten bacterial species were selected, in part based on the relevance in the

human microbiome. For each species, a reference genome and reference long-read dataset were

obtained from NCBI (File S3). Long-reads were mapped against their respective reference

genome with minimap2 and the mappings were used as input for the 'characterization stage' of

NanoSim [68] in genome mode. The resulting NanoSim models were used to generate simulated

long-read datasets in the 'simulation stage' in genome mode. Reads were generated from the

unmodified reference genome, and so no evidence of inversion for any inverton is expected, and

any invertons identified by PhaVa would be false positives. For each species, five coverage

levels and six replicates of each coverage level were generated (Data S7) totaling in 300

simulated long-read datasets. *Streptomyces eurocidicus* and *Enterococcus faecalis* simulated

readsets were generated at relatively deeper coverages due to poor read mapping characteristics

from the selected reference long-read datasets resulting in a smaller proportion of reads passing

PhaVa read mapping filters. Simulated readsets were then analyzed with PhaVa and used to

estimate false positive rates and optimize post-mapping filters.

<u>Identifying putative invertons from public long-read sequencing data with PhaVa</u>

Candidate isolate long-read sequencing datasets were identified on NCBI with the following

search criteria: "(Bacteria[Organism] OR Archaea[Organism]) AND ("pacbio smrt"[Platform]

OR "oxford nanopore"[Platform]) AND genomic[Source]". Datasets were further filtered by

removing datasets with the "amplicon" flag, and removing datasets with less than 50 Mbp of

sequencing in total (Data S5). Individual read datasets were downloaded with fastq-dump, a part

of the sratoolkit (https://trace.ncbi.nlm.nih.gov/Traces/sra/sra.cgi?view=software). Nanostat [69]

was run on each remaining readset to measure dataset characteristics. For each unique taxid

represented in the resulting readsets, a reference genome and paired genbank file (.gbff) were

selected by identifying a genome with the highest level of completion for that species, and the

least number of contigs. In the case of reference genomes with equal quality based on these

parameters, the first identified was selected. Long-read datasets were then analyzed with PhaVa

with default parameters. Gene overlaps and partial gene overlaps were identified by comparing

coordinates of genbank file annotations with inverton coordinates, a function available for use in

PhaVa (fig. S8).

<u>AlphaFold prediction</u>

Structural predictions of the amino acid sequences for the forward and reverse orientations of the

intragenic inverton within BT0375 were generated using AlphaFold [70] v2.2.0. The required

databases were downloaded on March 3rd, 2022 and the max template date was set to 2020-05-

14. The top ranked structures were then visualized and aligned in PyMOL.

<u>Gene set enrichment analysis</u>

In order to assess which functional groups were enriched for genes harboring intragenic

invertons, we performed a clade-resolved gene set enrichment analysis. We first annotated genes

with KEGG KOs using the kofamKOALA tool [71] and with Pfam domains by running HMMER3

[72] with the Pfam domain database. KEGG pathways and modules were filtered for those that

were present in bacterial genomes and Pfam clan definitions were downloaded from the Interpro

website [73]. We then calculated enrichments per genome and additionally per species and per

genus, for those combining the genes from all genomes in a specific clade. At each level, we

filtered out groups with fewer than 5 intragenic invertons (fig. S5), resulting in 10 genomes, 12

species, and 19 genera being included for downstream analysis. Alternatively, we also considered

genes with both intragenic or partial intergenic invertons, resulting in 47 genomes, 52 species, and 54

genera being tested. In each group, we tested for each pathway if the genes annotated with this

pathway were enriched for those carrying invertons by using a one-sided Fisher test. Pathways,

the genes of which did not harbor any invertons in a specific group, were skipped for the

enrichment analysis for a given group. Multiple testing correction was performed with the

Benjamini-Hochberg procedure [74].

<u>Identifying putative invertons from long-read metagenomes.</u>

200 hybrid short-read and long-read human stool metagenomic datasets were accessed from

BioProject PRJNA820119 [38]. Each hybrid dataset was assembled using SPAdes [75] with the '-

meta' flag and long-reads provided with the '--nanopore' option. An additional ten nanopore

888 long-read human stool microbiome metagenomic datasets from BioProject PRJNA940499 [39]

889 were assembled with Flye [76]. Assembled contigs will be deposited at

890 https://doi.org/10.5281/zenodo.7662825 after publication. Gene annotations for assemblies were

891 obtained with Prodigal [33] using the '-meta' flag. Contig taxonomic classifications were obtained

892 with Kraken2 [77]. Each long-read dataset was then analyzed with PhaVa with default parameters,

893 using its respective de novo assembly as its reference assembly. Resulting inverton calls were

894 dereplicated by clustering the inverton with 1000 bp flanking sequence upstream and

895 downstream at 99% average nucleotide identity with CD-HIT [78].

Supplemental Figures

899
900

Fig. S1. Inversion proportion of CPS loci invertons in BTh. Inversion proportions of CPS loci
invertons in HCT metagenomic samples measured with PhaseFinder. Samples with no inversions
in the five CPS invertons were removed.

905
906
907
908

Fig. S2. Inverton confirmation PCR primer design. A Forward and Reverse primer bind to
regions of the genome upstream and downstream of the inverton on opposite strands. The
Common primer binds the DNA inside of the inverton, between the inverted repeats. When the
DNA is in the forward orientation, the Common and Forward primer will generate a PCR
product. When the inverton flips, the Common and Reverse primer will generate a PCR product.

Fig. S3. Very long (>750bp), near perfect, inverted repeats can lead to false positives. (A)
Alignment of inverton NZ_CP025371.1:2124719-2124870-2125316-2125467, with its invertible
sequence inverted, against the *B. pertussis* genome leads to perfect alignment of flanking and IR
regions as expected. 'Reference genome' refers to the *B. pertussis* reference genome sequence.
'Inverton reversed' refers to the putative inverton sequence and flanking sequence, with the
invertible sequence inverted. Red dashed lines indicate boundaries of the invertible sequence,
black dashed lines indicate boundaries of the inverted repeats as detected by einverted, and
purple dashed lines indicate the true boundary of inverted repeats. (B) Alignment of the reverse
complement of the entire inverton NZ_CP025371.1:2124719-2124870-2125316-2125467 with
its invertible sequence inverted and flanking sequence, against the *B. pertussis* genome leads to
near perfect alignment (6 mismatches) spanning far into the flanking sequence to the true
boundary of the inverted repeats, allowing for reads to map regardless of inverton orientation.
(C) Example with toy nucleotide sequences. Red nucleotides indicate mismatches.

Fig. S4. Overview of SRA long-read isolate sequencing samples analyzed with PhaVa. (A) The number of unique species represented in the dataset, grouped by phylum. (**B**) The raw number of sequencing samples, grouped by phylum. (**C**) Histogram of sequencing samples per species. Species with particularly large numbers of samples are labeled. (**D**) A histogram of sequencing depths for all long-read isolate sequencing samples.

Fig. S5. Intragenic invertons are rare across genomes yet consistently enriched in some Pfam clans. (A) Histograms showing the number of clades (genomes, species, or genera) at various numbers of invertons indicate that invertons are rare, as only one to three invertons can be detected in the majority of clades. Only clades with at least five invertons (red line; number of clades is indicated in the top-right corner of each subplot) were included for the subsequent enrichment analysis. **(B)** KEGG pathways and Pfam clans were tested for enrichment of intragenic (or partial intergenic) invertons in included clades, using a one-sided Fisher's exact test per clade (see Methods). Enrichment was only calculated for sets with at least five invertons associated with genes in the set. Histograms show the number of sets with enrichment score at the number of included clades, showing that most enrichments could be calculated for single

clades only. For example, all KEGG pathways associated with enough intragenic invertons for
an enrichment analysis on genome-level were specific for each genome. Sets with enrichment
scores across at least five clades (red line) are labeled with their corresponding identifiers. **(C)**
Heatmap showing the log-odds ratio (effect size for the enrichment of intragenic invertons)
across included clades for the six Pfam clans that have enrichment scores on genus-level (see
panel B). Stars indicate significance of the enrichment as calculated by Fisher's exact test and
corrected for multiple hypothesis testing using the Benjamini-Hochberg procedure.
962

Fig. S6. PhaVa analysis of 210 long-read metagenomes from human stool. (**A**) Counts of invertons identified with PhaVa in 210 stool samples, grouped by phylum and the type of inverton. (**B**) Comparisons of the number of invertons (per genome) found in metagenomic datasets vs. SRA isolate sequencing samples. Total refers to all invertons identified, regardless of taxonomic classification. The distribution of inverton counts per species were found to be significantly different between metagenomes and isolate samples in both the Total and Firmicutes comparisons (p=3.35e-05 and p=0.005 respectively) with a Kolmogorov–Smirnov test. Other individual phyla were not compared due to small species counts with invertons in metagenomic samples.

Fig. S7. Locked *thiC* intragenic inverton construction and growth competition. (**A**)
Generation of locked intragenic invertons. The forward and locked forward *thiC* IR nucleotide
sequences are shown. When possible, the wobble position of each codon corresponding to the IR
was mutated to increase mismatches between the two palindromic sequences while maintaining
the amino acid sequence. Nucleotides that were mutated are highlighted in gray. (**B-C**) Locked
thiC strains were competed against each other in thiamine-containing media in a 1:1 ratio. After
40 hours, the abundance of each strain was enumerated using selective agar. Black bars indicate
the locked forward strain and white bars indicate the locked reverse strain. Recovered
abundances shown here correspond with the competitive index shown in Fig. 4D. In (**B**) the
locked forward strain is marked with an erythromycin resistant cassette and the locked reverse
strain is marked with a tetracycline resistant cassette. In (**C**) the locked forward strain is marked
with a tetracycline resistant cassette and the locked reverse strain is marked with an
erythromycin resistant cassette. Geometric mean and geometric standard deviation are shown for
replicates conducted across 4-6 independent experiments. For each thiamine concentration a ratio
paired t test was performed on the locked forward and locked reverse abundances. ***, $p <$
0.001; **, $p < 0.01$; *, $p < 0.05$.

Figure S8: Inputs and outputs of a variation_wf PhaVa run. Output tables of particular interest are labeled and shown below the diagram with example output.

Strain name	Source	Identifier
Bacteroides thetaiotaomicron VPI-5482 Δ*tdk*	[79]	WT
Bacteroides thetaiotaomicron VPI-5482 Δ*tdk* ΔBT0650	this study	RC131
Bacteroides thetaiotaomicron VPI-5482 Δ*tdk* BT0650 locked RV	this study	RC149
Bacteroides thetaiotaomicron VPI-5482 Δ*tdk* BT0650 locked FW	this study	RC134
Bacteroides thetaiotaomicron VPI-5482 Δ*tdk* BT0650 locked FW *NBU2::NBU2_tet*	this study	RC165
Bacteroides thetaiotaomicron VPI-5482 Δ*tdk* BT0650 locked FW *NBU2::NBU2_erm*	this study	RC 166
Bacteroides thetaiotaomicron VPI-5482 Δ*tdk* BT0650 locked RV *NBU2::NBU2_erm*	this study	RC164
Bacteroides thetaiotaomicron VPI-5482 Δ*tdk* BT0650 locked RV *NBU2::NBU2_tet*	this study	RC163
E. coli S17-1 λ*pir*; *zxx::*RP4 2-(Tetr::Mu) (Kanr::Tn7) λ*pir*	[80]	S17-1 λpir
E. coli DH5α *λpir; F- endA1 hsdR17 (r-m+) supE44 thi-1 recA1 gyrA relA1 Δ(lacZYA-argF)U189 φ80lacZΔM15 λpir*	[81]	DH5α λpir

996
997 **Table S1. Strains used in this study**
998

Recombinant DNA	Identifier	Source
pKNOCK-*bla-ermGb*::*tdk*	pExchange	[79]
pExchange BT0650 KO	pRBC20	this study
pExchange BT0650 locked FW	pRBC21	this study
pExchange BT0650 locked RV	pRBC22	this study
pNBU2_tet	tetR	[24]
pNBU2_erm	ermR	[24]

Table S2. Recombinant DNA used in this study

Acknowledgments: We thank Nora Enright, Danica Schmidtke, Aravind Natarajan, Jack Diaz Shanahan, Dylan Maghini, Mai Dvorak, Alvin Han, Meena Chakraborty, and Bhatt Lab members for helpful conversations and scientific advice regarding this project. We also thank Wenhan Zhu for plasmids (pNBU2_tet and pNBU2_erm) and Sebastian Winter for the strain (DH5α) that were used in the context of this project. We also thank Daniel Haft and Francoise Thibaud-Nissen at NCBI for helpful discussion about accessing SRA long-read datasets.

Funding:

National Institutes of Health R01 AI148623 (ASB)

National Institutes of Health R01 AI143757 (ASB)

Stand Up 2 Cancer Foundation

National Institutes of Health T32 training Grant HG000044 (RBC)

National Institutes of Health T32 training Grant HL120824 (PTW)

Author Contributions:

Conceptualization: RBC, PTW, ASB

Methodology: RBC, PTW, JW

Investigation: RBC, PTW, JW, RMP, GZMG, ASH, MOG, EFB, AMM

Visualization: RBC, PTW, JW, RMP

Funding acquisition: ASB

Project administration: RBC, PTW, ASB

Supervision: RBC, PTW, ASB

Writing – original draft: RBC, PTW, ASB

Writing – review & editing: RBC, PTW, JW, RMP, GZMG, ASH, MOG, EFB, AMM, ASB

Competing Interests: Authors declare that they have no competing interests.

Data and materials availability: PhaVa is available at (https://github.com/patrickwest/PhaVa). Short-read adult HCT stool sequencing data was previously published and is available at (NCBI BioProject ID PRJNA707487). Short-read pediatric HCT stool sequencing data was previously published and is available at (NCBI BioProject ID PRJNA787952). Long-read metagenomic sequencing data was previously published and is available at BioProject PRJNA820119 and BioProject PRJNA940499. Assembled metagenomic contigs will be made available after

1033 publication at https://doi.org/10.5281/zenodo.7662825. A list of accession numbers for long-read

1034 isolate sequencing data is available in supplementary file Data S5.

Supplementary Information is available for this paper.

Extended Data Tables

Data S1. *B. theta* **intragenic invertons identified from short-read metagenomic sequencing samples.**

Data S2. Archaeal invertons identified from long-read isolate sequencing samples.

Data S3. Invertons identified from long-read isolate sequencing samples.

Data S4. Dereplicated invertons identified from long-read metagenomic sequencing samples.

Data S5. List of accession numbers and associated metadata for long-read isolate sequencing samples.

Data S6. Primers used in this study.

Data S7. Simulated read datasets.

Data S8. Additional sequences.

Correspondence and requests for materials should be addressed to Ami Bhatt (asbhatt@stanford.edu).