A high-resolution view of RNA endonuclease cleavage in *Bacillus subtilis*

James C. Taggart[1,4], Julia Dierksheide[1], Hannah LeBlanc[1], Jean-Benoît Lalanne[1,2,5], Sylvain Durand[3],

Frédérique Braun[3], Ciarán Condon[3], and Gene-Wei Li[1,#]

[1]Department of Biology, Massachusetts Institute of Technology, Cambridge, MA 02142, USA

[2]Department of Physics, Massachusetts Institute of Technology, Cambridge, MA 02142, USA

[3]Expression Génétique Microbienne (EGM), CNRS, Université Paris Cité, Institut de Biologie Physico-

Chimique, 13 rue Pierre et Marie Curie, 75005 Paris, France

[4]Current address: Department of Systems Biology, Harvard Medical School, Boston, MA, USA

[5]Current address: Department of Genome Sciences, University of Washington, Seattle, WA, USA.

[#]corresponding author (gwli@mit.edu)

ABSTRACT

RNA endonucleases are the rate-limiting initiator of decay for many bacterial mRNAs. However, the

positions of cleavage and their sequence determinants remain elusive even for the well-studied *Bacillus*

subtilis. Here we present two complementary approaches – transcriptome-wide mapping of

endoribonucleolytic activity and deep mutational scanning of RNA cleavage sites – that reveal distinct

rules governing the specificity among *B. subtilis* endoribonucleases. Detection of RNA terminal

nucleotides in both 5′- and 3′-exonuclease-deficient cells revealed >10^3 putative endonucleolytic cleavage

sites with single-nucleotide resolution. We found a surprisingly weak consensus for RNase Y targets, a

contrastingly strong primary sequence motif for EndoA targets, and long-range intramolecular secondary

structures for RNase III targets. Deep mutational analysis of RNase Y cleavage sites showed that the

specificity is governed by many disjointed sequence features, each with mild contributions. Our results

highlight the delocalized nature of mRNA stability determinants and provide a strategy for elucidating

endoribonuclease specificity *in vivo*.

INTRODUCTION

Predicting protein abundance from an arbitrary gene sequence remains an aspirational goal for quantitative biology. Decades of research has explored the encoding of rates of transcription and translation initiation (Taggart et al., 2021), resulting in predictive models for how RNA polymerase and the ribosome recognize promoters and ribosome binding sites (Brewster et al., 2012; Salis, 2011; Salis et al., 2009; Urtecho et al., 2019). Comparatively poorly characterized, however, is post-transcriptional control of RNA abundance by ribonucleases, factors which exert widespread quantitative control on bacterial gene expression levels (Chen et al., 2015; Khemici et al., 2015; Moffitt et al., 2016). Understanding the encoding of ribonuclease cleavage sites in bacterial mRNA sequences is therefore a critical question in quantitative biology.

In the Gram-positive bacterium *Bacillus subtilis*, cleavage by an endoribonuclease is thought to initiate the decay of a large fraction of cellular mRNAs (Durand et al., 2012a; Lehnik-Habrink et al., 2011a, 2012; Shahbabian et al., 2009). The cleaved RNA fragments typically have a 5′-phosphate and a 3′-hydroxyl group, which are subsequently degraded by a 5′-3′ exoribonuclease (RNase J1) and by a combination of 3′-5′ exoribonucleases (PNPase, RNase R, RNase PH, and YhaM) (Trinquier et al., 2020), respectively. The endoribonuclease RNase Y appears to drive the turnover of the majority of mRNAs in *B. subtilis*, and its depletion results in the accumulation of hundreds of transcripts (Durand et al., 2012a; Lehnik-Habrink et al., 2011a; Shahbabian et al., 2009). For several well-studied mRNAs, RNase Y cleavage is specific to certain positions and can drive operon mRNA maturation (Braun et al., 2017; Commichau et al., 2009; DeLoughery et al., 2018; Noone et al., 2014). Unfortunately, though RNase Y has been identified as a key player shaping the transcriptomes of Gram-positive bacteria, our understanding of which mRNA sequences *B. subtilis* RNase Y cleaves remains relatively limited. Studies have suggested *B. subtilis* RNase Y recognizes single stranded RNA, is stimulated by downstream secondary structure, and has a preference for adenosine and uridine-rich sequence (Braun et al., 2017; Shahbabian et al., 2009). Orthologs of RNase Y in *Staphylococcus aureus* and *Streptococcus pyogenes*, organisms for which RNase Y is not essential and appears to play a more limited role compared to *B.*

subtilis, additionally demonstrate a preference for cleavage downstream of a guanosine (Khemici et al.,

2015) (Benda et al., 2021). However, these limited sequence preferences cannot explain the highly

specific locations of cleavage that have been observed for the *B. subtilis* RNase Y (DeLoughery et al.,

2018).

In addition to RNase Y, several other endoribonucleases contribute to mRNA decay. RNase III

has been suggested to cleave and initiate decay for dozens of mRNAs in *B. subtilis* (DiChiara et al.,

2016). Although the *E. coli* RNase III strictly cleaves double-stranded RNAs, the *B. subtilis* RNase III has

been implicated to deviate from this rule in a way that remains unknown (DiChiara et al., 2016). Other

minor endoribonucleases, such as the RNase toxin EndoA (a homolog of *E. coli* MazF), may also initiate

mRNA decay and produce different 5′ and 3′ moieties (Cerullo et al., 2022; Ingle et al., 2022; Leroy et al.,

2017). Their positions of cleavage and subsequent decay processes remain poorly described (Pellegrini et

al., 2005).

A number of studies have mapped the target repertoire of major bacterial endoribonucleases

(Altuvia et al., 2018; Broglia et al., 2020; Chao et al., 2017; DiChiara et al., 2016; Gordon et al., 2017;

Khemici et al., 2015). These measurements detect the terminal nucleotides of RNA decay intermediates

and analyze their abundance upon endoribonuclease inactivation. The results are confounded by several

factors. First, because cleaved decay intermediates are typically unstable, the detected end signals are

often close to background levels and may be biased by their own stability. Second, the sequential actions

of several endoribonucleases and/or exoribonucleases on the same mRNA may result in RNA ends that

are shifted from the initial position that is cleaved by the endoribonuclease under investigation. As a

result, low-throughput approaches are often required to precisely determine exact positions of cleavage

for each target (Braun et al., 2017; Bruscella et al., 2011; DiChiara et al., 2016; Yao and Bechhofer,

2010).

In this work, we first present a global approach to map bacterial endoribonuclease cleavage sites

by combining high-resolution RNA end-mapping with genetic perturbation of all known exonucleases

participating in *B. subtilis* mRNA decay. By stabilizing mRNA decay intermediates in exonuclease

deficient *B. subtilis* cells, this "stabilized end-sequencing" approach revealed thousands of putative

positions of endoribonuclease cleavage and a previously unknown nuclease activity targeting RNA 5′

ends. Furthermore, using a panel of endoribonuclease knockouts, we dramatically expanded the known

target repertoire for RNase Y, RNase III, and EndoA. Our results captured a revised EndoA cleavage

motif, elucidated long-range RNA folds that dictate RNase III cleavage, and a surprisingly weak

consensus among RNase Y cleavage sites, despite their high specificity. To further investigate the

sequence determinants for RNase Y cleavage, we developed a deep mutational scanning approach to

interrogate individual cleavage sites. Interestingly, the residues that affect cleavage activity are scattered

across a region over 40 nucleotides long. These results suggest that the highly specific RNase Y cleavage

does not follow the conventional model of protein-nucleic acid interactions mediated by compact

sequence or structural motifs. Instead, the "cleavage code" is specified by a collection of weak

interactions embedded in an extended region.

RESULTS

Stabilized end-sequencing reveals positions of endoribonucleolytic activity

In the absence of additional nucleolytic activity, one can map the positions of cleavage by an

endoribonuclease by simply observing the termini of its cleavage products. *In vivo*, however,

exonucleases typically rapidly act on endonuclease cleavage products, rendering the original RNA

endonuclease cleavage sites difficult to detect. If all exoribonucleolytic activity targeting either 3′ or 5′

ends is ablated, endoribonuclease cleavage at a particular site should result in stably accumulating

products whose ends correspond to the position of cleavage (Figure 1A). Strains combining knockouts of

all known exonucleases in either the 5′ or 3′ direction are viable in *B. subtilis*, meaning such a

measurement is theoretically possible in this species (Figaro et al., 2013; Oussenko et al., 2005). If RNA

ends are separately mapped in both a knockout of RNase J1, the only known 5′-3′ exonuclease, and a

knockout of all four known 3′-5′ exonucleases (a knockout of *yhaM, rph, pnpA,* and *rnr*, hereafter referred

to as a 4-exo knockout), exact positions of endoribonucleolytic activity can be captured as pairs of

adjacent 3′ and 5′ ends that appear across the two strains.

Based on this principle, we implemented a "stabilized end-sequencing" approach to create a

transcriptome-wide map of endoribonuclease activity by capturing the 3′ hydroxyl and 5′ monophosphate

groups that are generated by endonucleases such as RNases Y and III (the primary endonucleases

targeting mRNAs in *B. subtilis*) (Figure 1A; (Bechhofer and Deutscher, 2019; Leroy et al., 2017)). To

quantitatively capture 3′ hydroxyl groups, we utilized a previously established method for RNA 3′

ligation and deep sequencing, hereafter referred to as 3′ end sequencing (Figure 1A, left; (Dar et al., 2016;

Herzel et al., 2022; Mandell et al., 2021)). To complement 3′ end mapping, we specifically captured 5′

monophosphates through T4 RNA Ligase 1 treatment, allowing quantification of cleavage-derived 5′ ends

by deep sequencing, hereafter referred to as 5′ end sequencing (Figure 1A, right). By performing 3′ end

sequencing on a 4-exo knockout (strain CCB396) and 5′ end sequencing on an RNase J1 knockout (strain

CCB434), we captured 3′/5′ end pairs derived from endoribonuclease activity transcriptome-wide.

Combining these exoribonuclease perturbations with endoribonuclease knockouts, we can attribute these

cleavage signatures to specific enzymes. End-enriched RNA sequencing (Rend-seq) can additionally be

performed to quantify expression of putative endoribonuclease targets and provide an orthogonal RNA

end-mapping approach (Lalanne et al., 2018).

This stabilized end-sequencing strategy robustly captures positions of previously documented

endoribonucleolytic cleavage sites. Among 13 previously characterized sites cleaved by either RNase Y,

RNase III, or Rae1 that were individually detected (Braun et al., 2017; Bruscella et al., 2011; DiChiara et

al., 2016; Durand et al., 2015; Leroy et al., 2017; Noone et al., 2014; Yao and Bechhofer, 2010), we

identified at least one clear 3′/5′ pair at or near the annotated cleavage position (Figure 1B, Figure S1). 5′

end sequencing of RNase Y+J1 or III+J1 double mutant strains (CCB760 and BG879) confirmed that we

can detect cleavage by both endoribonucleases and accurately assign the enzymes responsible for

cleavage (Figure 1B).

Importantly, the paired end-mapping strategy provides clear advantages over measurements of a

single end type. Because we capture both RNA ends generated by endoribonuclease activity, multiple

adjacent positions of cleavage can be resolved, as appears to be the case for the first characterized

substrate of Rae1, *S1025* (Figure 1B, bottom). Without these two paired channels of information, it is

difficult to determine whether signal at adjacent positions is derived from multiple positions of

endoribonucleolytic cleavage or residual exoribonuclease activity by unknown enzymes, the latter of

which may be occurring after endoribonucleolytic cleavage of the RoxS RNA (Figure 1B, middle).

Precisely resolving cleavage positions becomes particularly relevant in cases such as the

endoribonucleolytic cleavages observed in *hbs*, where many nearby positions are known to be subject to

RNase Y cleavage (Figure S1, (Braun et al., 2017)).

Using these data, we first identified putative positions of cleavage within transcripts whose

abundance is known to be regulated by RNA endonuclease activity. Recent work has shown that the RNA

decay machinery in *B. subtilis* is subject to extensive autoregulation and cross-regulation, suggesting that

cleavage sites should be detectable within the RNAs encoding decay-associated proteins (DeLoughery et

al., 2018; DiChiara et al., 2016; Korobeinikova et al., 2023). This expectation is borne out in our

stabilized end-sequencing, where multiple putative positions of endoribonucleolytic cleavage are detected

within the transcripts encoding RNases Y, III, and the Y-complex, a complex of three small proteins

(YlbF, YmcA, and YaaT) that modulates RNase Y activity (Deloughery et al., 2016) (Figure 2, Figure

S2). Further, a number of additional transcripts, including that encoding sporulation master regulator

SinR, have been experimentally characterized as stabilized upon deletion of RNase Y (Deloughery et al.,

2016; Lehnik-Habrink et al., 2011a). Paired peaks are clearly visible within each of these transcripts (Fig

2, Fig S2). Taken together, our ability to robustly identify known and previously unknown positions of

cleavage suggests that stabilized end-sequencing provide a high-resolution, genome-wide view of RNA

endonuclease activity in *B. subtilis*.

Systematic identification of positions of endoribonuclease cleavage

To systematically identify putative positions of endonuclease cleavage, we implemented a computational

approach to call abutting pairs of peaks in 3′ and 5′ end sequencing across our 4-exo and RNase J1

knockout data, respectively (Figure S3A). To do so, we assessed the signal at each position relative to the

average in a local window, calling positions exceeding a signal-to-background threshold as peaks.

Positions for which a peak is called in both 5′ and 3′ end sequencing with a 1-nucleotide offset (a 5′ peak

downstream of a 3′ peak) are called as peak pairs and represent putative sites of endoribonucleolytic

activity. This approach reveals 1798 putative positions of cleavage in mRNAs. It should be noted that we

are unable to identify cleavage events proximal to the ends of transcription units, because short RNA

decay fragments (<35 nt) are excluded during 5′ end sequencing library preparation (see Methods).

To assign putative cleavage positions to the activity of specific endoribonucleases, we identified

the positions that have diminished end-signals when an endoribonuclease is knocked out (Fig S3B). To

account for changes in expression levels in strains with and without the endoribonuclease, we first

normalized end-sequencing signals to local RNA abundance measured by Rend-seq in the same strain.

We then quantified the sensitivity of the expression-normalized signal to endoribonuclease knockout, by

computing the ratio of 5′-end signals (in the absence of RNase J1) between strains with and without the

endoribonuclease (Materials and Methods). This "sensitivity" score will be elevated for positions cleaved

by the corresponding endoribonuclease, as was previously shown by DiChiara et al. (DiChiara et al.,

2016).

***B. subtilis* RNase III cleaves within long-range intramolecular structures**

We first characterized the cleavage profile of RNase III, motivated by previous reports of unexpected

cleavage patterns in *B. subtilis* (DiChiara et al., 2016). RNase III plays well-characterized roles in the

processing of noncoding RNAs (DiChiara et al., 2016; Herskovitz and Bechhofer, 2000; Oguro et al.,

1998) and the regulation of mRNAs (Durand et al., 2012b; Jahn and Brantl, 2013; Meißner et al., 2016).

Several potential mRNA targets of this enzyme have previously been proposed, but only a small number

of cleavage sites in these mRNAs have been identified (DiChiara et al., 2016; Gordon et al., 2017).

Intriguingly, such cleavage sites often did not appear as pairs at a defined 2-nt spacing within RNA

duplexes, contrasting with the hallmark of RNase III activity in *E. coli* (Altuvia et al., 2018). We reasoned

that a more systematic, high-resolution mapping of RNase III cleavage sites in *B. subtilis* will help resolve

this issue.

Our stabilized end-sequencing data revealed 68 RNase III cleavage sites located on 42 mRNAs

(Table S3). These sites constitute a small fraction of the putative endonuclease cleavage sites we

identified (5%) in the background strain (phage-cured (Gilet et al., 2015)), consistent with the notion that

RNase III targets a small subset of mRNAs (Fig. 3B). Despite the sparseness of these sites, they are often

present in the same mRNAs as other RNase III sites (45 of 68) (Figure 3A, Figure S1, and Table S3).

These results suggest that the *B. subtilis* RNase III acts in a manner similar to the *E. coli* RNase III in

cutting a pair of positions in RNA duplexes.

Indeed, of the 44 of these co-occurring cleavage sites that occur within 1 kb of each other, a large

majority (36 of 44) are predicted to reside in RNA secondary structures. The extent of secondary

structures is substantial, with up-to 505 nt between cleavage sites (Figure 3D). Despite the large contour

length, the cleavage positions are almost always (32 of 36) separated by 2 base pairs on the opposite sides

of RNA duplexes, which range in length from 10 to 36 basepairs (Figure 3C E, and F). 4 of the 36 co-

occurring sites are cleaved at 2 adjacent positions, which may still result from the 2-bp rule on individual

RNA molecules but are difficult to resolve in our ensemble measurement. It is possible that the minority

of sites that do not form a clear intra-molecular duplex (32 of 68) participate in inter-molecular pairing,

but the presence of such interactions remains to be determined.

We next sought to determine what sequence features within these specific secondary structures

might be recognized by RNase III. To increase the number of RNase III sites used in this analysis, we

included putative cleavage positions with a lower threshold on 5′ end sequencing depth (see Methods for

details, Tables S5,6 for a list of putative cleavage positions with this relaxed threshold, and Figures S3C,E

for endonuclease sensitivity distributions), restricting our analysis only to pairs of such putative cleavage

positions that were predicted to pair with 2-bp spacing. Using this approach, we identified an additional

10 pairs of RNase III cleavage sites for a total of 26 pairs. The genomic distance between these sites is

similar to those initially identified (Figure S3D).

From these pairs of RNase III cleavage sites, we calculated the position-wise sequence content

and along the predicted RNA duplex, as well as the fraction of sites for which each position is paired,

using the two cleavage positions as our reference points (Figure 3G). Through this, we generated a *B.*

subtilis RNase III sequence-structure motif as has previously been reported for *E. coli* (Altuvia et al.,

2018, Figure 3H). Like the *E. coli* homolog, the positions surrounding *B. subtilis* RNase III cleavage sites

appear to be consistently paired, with the probability of pairing increasing towards the cleavage positions

(reaching >90% at the -2/+2 positions) (Figure 3G). Similarly, a number of sequence features appear to be

recognized similarly by these enzymes, including position-specific preferences for A-U and G-C base-

pairing within the proximal box (PB) and an increased occurrence of U-G wobble pairing within the distal

box (DB) (Altuvia et al., 2018; Pertzev and Nicholson, 2006). At each position from -6 to 6, at least one

of the maximum-frequency base pairs is conserved between these species (Figure 3H, green text). In

contrast to *E. coli*, however, the base pairs between cleavage positions do not demonstrate a clear

preference for C-G/G-C pairing in our data (Altuvia et al., 2018) (Figure 3H, red box).

Together, these findings suggest that the *B. subtilis* RNase III displays similar substrate

preferences to its evolutionarily distant *E. coli* homolog (Altuvia et al., 2018; Pertzev and Nicholson,

2006). Furthermore, the presence of extensive long-range RNA secondary structure for RNase III targets

suggests that mRNA base pairing across hundreds of nucleotides is common in this organism and can

directly influence gene expression regulation (Braun et al., 2017; Ruiz de los Mozos et al., 2013).

Induction of EndoA activity in exoribonuclease knockouts reveals additional 5′ end trimming

activity in *B. subtilis*

To our surprise, manual inspection of our Rend-seq data in all strains with an RNase J1 or 4-exo knockout

showed prominent peaks across the transcriptome within the motif U/ACAUA (/ denotes the cleavage

position, Figure 4A, B). This sequence resembles an extended version of the canonical U/ACAU cleavage

motif of the endonuclease EndoA, an RNA interferase of the MazF family associated with the EndoA-

EndoAI type II toxin-antitoxin system (Park et al., 2011; Pellegrini et al., 2005). The presence of adjacent

3′ and 5′ ends in Rend-seq data separating the first U/A dinucleotide within this motif strongly suggests

endoribonuclease activity (Figure 4A, B), possibly through activation of this toxin in the exoribonuclease

deficient strains used in this study. This U/ACAUA-specific cleavage is not observed in Rend-seq data

from wild-type *B. subtilis* (Figure S4A). Consistent with EndoA-mediated cleavage, which leaves a 5′

hydroxyl group (Bechhofer and Deutscher, 2019; Pellegrini et al., 2005), we did not observe

corresponding signals in our 5′ end sequencing data, which is specific to 5′ monophosphate groups,

shown for a representative site in the *pgk* gene (Figure 4E). Because Rend-seq relies on cDNA

circularization rather than RNA ligation to capture RNA 5′ ends, it is agnostic to the 5′ moiety generated

following cleavage (Lalanne et al., 2018). To further confirm that these cleavage sites are driven by

EndoA, which is encoded by the *ndoA* gene, we repeated these measurements upon deletion of *ndoA* in an

RNase J1 deficient strain and found complete ablation of the cleavage signature in Rend-seq (Figure 4C,

D). Furthermore, consistent with recent work indicating that EndoA recognizes a 6-mer motif and not a

previously described 5-mer motif (Guler et al., 2024), we found that only U/ACAU followed by an

adenosine leads to cleavage (Figure 4A, B).

Though our monophosphate-specific 5′ end sequencing approach did not detect the hydroxylated

5′ ends generated by EndoA, it unexpectedly captured peaks 2 nucleotides downstream of the canonical

EndoA cleavage positions (Figure 4E). These 5′ monophosphorylated ends only appear when EndoA is

present (Figure 4D, E), suggesting that two nucleotides are removed following cleavage by EndoA. To

test if this trimming activity is general against all 5′ hydroxyl groups, not simply EndoA sites, we looked

at the 5′ OH generated by the self-cleaving *glmS* ribozyme, whose downstream cleavage product is

subject to subsequent degradation by RNase J1 (Collins et al., 2007; Winkler et al., 2004). In both a

knockout of RNase J1 and a double knockout of RNase J1 and EndoA, 5′ end sequencing shows a clear 5′

monophosphorylated peak 2 nucleotides downstream of the site of self-cleavage in *glmS*, suggesting that

this novel trimming activity is independent of EndoA (Figure 4F). The trimming activity is also

independent of RNases Y, III, J1, the Y-complex constituent YlbF, or the RNA pyrophosphohydrolase

RppH, as downstream peaks are readily detectable in strains lacking these enzymes (Figure 4G). Finally,

this activity cannot be explained by RNase J2, as 5′-mapped Rend-seq measurements of a double

knockout of RNase J1 and J2 also show a clear peak two nucleotides downstream of the canonical *glmS*

cleavage position (Figure 4H).

We observed multiple rounds of 2-nt trimming at some, but not all, EndoA cleavage sites,

suggesting sequence-specific trimming activities (Figure S4B,C). To determine the sequence specificity,

we examined the downstream nucleotide content following the U/ACAUA EndoA motif (Figure 4I),

focusing on our *rny* knockout and RNase J1 depletion strain. This strain was chosen for its particularly

clear EndoA activity, which is likely driven by further stabilization of EndoA cleavage products

following inactivation of an additional downstream decay pathway, though EndoA activity may also be

increased. For sites with an EndoA motif followed by a U or C, monophosphorylated 5′ ends are found at

both 2 and 4 nucleotides downstream of the initial cleavage, indicating discrete trimming (Figure 4I). For

U/ACAUA motifs followed by an A (particularly by AA), additional 5′ ends are also generated five or

more nucleotides downstream of the initial cleavage. By contrast, motifs followed by a G only produce 5′

ends at 2 nucleotides downstream. These 5′ ends are not driven by changes to the EndoA cleavage

position, as the 3′ end density in the 4-exo knockout remains constant regardless of the downstream

sequence context (Figure 4J). These data are consistent with successive rounds of nucleotide removal

from the RNA 5′ end following endoribonucleolytic cleavage, with a preference for trimming sequences

rich in adenosine and against those rich in guanosine. We are unable to rule out that differential stability

following EndoA cleavage and 5′ end trimming may also influence the apparent sequence preferences we

observe, however, and further work is required to reveal the identity and mode of action of this enzyme.

Cleavage 2 or 4 nucleotides downstream of the canonical EndoA cleavage site coincides with the

U/A and C/A dinucleotides that are recognized by the common laboratory contaminant RNase A. Several

lines of evidence argue against such a contaminant generating the trimming behavior we characterize in

this work. First, RNase A cleavage generates RNA termini with 5′ hydroxyl or 3′ phosphate groups,

similar to EndoA (Cuchillo et al., 2011). These ends cannot be captured by our RNA 5′ and 3′ end

sequencing approaches (Figure 4E). Second, the 5′ trimming activity observed at *glmS* occurs within a

C/G dinucleotide, suggesting that this activity is not restricted to U/A and C/A dinucleotides. Finally, our

observation of 5′ end trimming is highly consistent across many replicate measurements of strains lacking

RNase J1, as is an observed lack of trimming when this enzyme is present. It is thus highly unlikely that

this trimming is driven by *in vitro* RNase activity downstream of cell lysis and RNA extraction.

Determinants of cleavage by RNase Y

Though RNase Y is thought to be the primary endonuclease driving RNA decay in *B. subtilis*, only a

small number of cleavage positions have been mapped in this bacterium, and what specifies RNase Y

cleavage is poorly understood (Trinquier et al., 2020). To define the substrate features driving RNase Y

cleavage, we sought to identify a set of putative cleavage sites that are dependent on the presence of this

enzyme. For the 1330 putative endonuclease cleavage positions identified in our W168 background, we

calculated their RNase Y sensitivity using RNase J1 deficient cells with or without an RNase Y knockout.

This metric was bimodally distributed, and any cleavage signature with a sensitivity value over 10.56 was

called as RNase Y dependent based on a double-Gaussian fit of these data (Figure 5A). Using this

approach, we identified 596 cleavage sites across 387 genes that can confidently assigned to RNase Y

sites (Table S4).

This set of putative RNase Y sites sheds light onto the substrate features that guide RNase Y

activity. Frequency and information content logos constructed from the RNA sequences surrounding

putative RNase Y sites suggest a sequence preference for cleavage downstream of a purine, with a

particular preference for guanosine, and upstream of an adenosine (Figure 5B, top, middle). Similarly,

considering k-mers of length ≤4, there is most significant enrichment for the dinucleotide sequence GA

spanning the cleavage site, with an additional preference for uridine in the -2 and -3 positions (Figure 5B,

bottom). This observed specificity is unlikely to be a result of biases in our methodology; identification of

these cleavage sites requires concordance between 3′ and 5′ end sequencing datasets that are generated

through distinct enzymatic reactions and adapter sequences. This sequence preference is additionally

consistent with the sequence context in several previously mapped cleavage sites (Braun et al., 2017;

Shahbabian et al., 2009), as well as the measured sequence preferences of RNase Y in *S. aureus* and *S.*

pyogenes, despite the more limited role this enzyme plays in those organisms (Broglia et al., 2020;

Khemici et al., 2015). However, the GA dinucleotide motif does not appear to be absolutely required, as

63% of the putative cleavage sites do not have this motif.

In a broader window around cleavage sites, there is a clear depletion for GC-containing sequence

(Figure 5C) and a lower propensity to form secondary structures (Figure 5D). Computationally predicted

folding of 40-nt RNAs surrounding positions of cleavage shows weaker structures for RNase Y sites

compared to randomly sampled positions at uncleaved GA dinucleotides (Figure 5D). Interestingly,

sequences centered just downstream of this lowly-structured region appear to fold more strongly than

background, suggesting an enrichment for secondary structure proximal to the position of cleavage

(Figure 5D). The downstream signal is recapitulated in the GC content within this window (Figure 5C),

and is consistent with the influence of secondary structure on previously detected RNase Y cleavage

events (Marincola and Wolz, 2017; Shahbabian et al., 2009). Taken together, these data indicate that *B.*

subtilis RNase Y targets share a weak motif comprising a GA dinucleotide surrounded by AU-rich

sequences that have reduced secondary structure.

The complete Y-complex is not a strict requirement for RNase Y activity

Recent work has shown that the Y-complex, comprising proteins YlbF, YaaT, and YmcA, can act as a

specificity factor for RNase Y for the maturation of nearly two-dozen operon mRNAs (DeLoughery et al.,

2018). To understand if this complex plays a role in specifying the observed RNase Y cleavage events,

we deleted the *ylbF* gene in both an RNase J1 knockout (CCB434) and 4-exo deficient background

(CCB396). We then repeated our Rend-seq and end-sequencing measurements in these backgrounds,

calculating the sensitivity of each putative cleavage site to YlbF (Figure 5E). Compared to RNase Y, few

sites appeared dependent on the presence of YlbF, suggesting that the Y-complex plays a limited role in

influencing RNA turnover outside of its known role in mRNA processing. Consistent with this

observation, knocking out *ylbF* has a limited effect on the *B. subtilis* transcriptome (Figure S4D), as has

been previously observed (DeLoughery et al., 2018).

Surprisingly, canonical maturation events, such as those within the *cggR-gapA*, *glnRA*, and *atp*

operons are still observed in the absence of YlbF, albeit strongly attenuated (Figure 5E-H). Previously,

the YlbF-dependence of these sites were determined using Rend-seq for cells that maintain the entire

repertoire of exonucleases, which is less sensitive at detecting cleavage events. Here, by ablating

exonucleases in cells lacking YlbF, we observed weak but detectable signals of cleavage products using

Rend-seq, as well as unambiguous peaks in both 5′ and 3′ end sequencing (Figure 5F-K). These results

underscore the sensitivity of our approach and suggest a revision of the role of the Y-complex. That is,

instead of being a strictly necessary specificity factor for RNase Y cleavage, the Y-complex plays an

enhancing role at these operon maturation events.

Deep mutational scanning of RNase Y substrates

Our census of RNase Y cleavage events showed a degenerate motif, which is insufficient to explain its

highly precise site selection for previously established mRNA targets. To identify additional sequence or

structural features required, we developed a massively parallel reporter assay (MPRA) that allows high-

throughput measurement of cleavage efficiency among a library of 10^4-10^5 mutated cleavage sites (Figure

6, Table S7). The reporter consists of a >100-nt region surrounding a cleavage site inserted into a stable

native mRNA encoding the gene *aprE*, truncated such that it lacks an active site (Figure 6A). These

reporter constructs are integrated at a neutral genomic locus (*ganA*) and contain a randomized 15-

nucleotide barcode associated with each mutant variant, allowing relative quantification of each variant's

expression by RNA and DNA sequencing.

To simultaneously quantify cleavage activity at each mutant cleavage site, a pooled library of *B.*

subtilis cells expressing our MPRA construct were grown and harvested in exponential phase for genomic

DNA (gDNA) and RNA extraction (Figure 6B). Purified RNA is then subjected to reverse transcription

and a size selection that separates cleaved from full-length RNA, allowing specific detection of each

isoform. A reduction in cleavage should result in greater accumulation of the full-length RNA, with

concomitant decrease in cleaved RNA. The barcode abundance in the uncleaved RNA fraction for each

variant, normalized by its frequency in the pool as measured by gDNA sequencing, thus provides a

readout of cleavage efficiency for that mutated sequence (Figure 6B).

RNase Y cleavage is specified by distributed substrate features

We initially focused our MPRA measurements on a cleavage site found within the *cggR* gene, which in its

native context drives differential expression within the glycolytic operon by producing a stable

downstream product following cleavage (Ludwig et al., 2001; Meinken et al., 2003) (Figure 5F, G). The

cleaved downstream product, which includes the *gapA* gene, is thought to be stabilized by an RNA

hairpin structure at the cleaved end (henceforth referred to as the downstream hairpin) (Meinken et al.,

2003), but whether proximal structure also plays a major role in specifying the cleavage site by *B. subtilis*

RNase Y remains unknown. To capture surrounding sequences that may contribute to this cleavage, we

inserted 148 bp of this sequence into our reporter, approximately centering the cleavage position within

this window (Figure 6C). The site of cleavage, which in its native context resides in the *cggR* open

reading frame, was similarly translated in the same reading frame of the reporter. Also adhering to the

native context, the site of cleavage was followed by a stop codon and a short open reading frame derived

from the start of the downstream *gapA* gene (Figure 6C). Using this approach, we characterized the effect

of mutations from positions -30 to +36 relative to the cleavage position. After sequencing of the gDNA

and RNA, 91,580 uniquely barcoded variants were detected across all experiments. These data cover

virtually all possible single-nucleotide changes (Figure 6D) and 32% and 44% of all double-nucleotide

changes across experiments detecting full-length and cleaved RNAs, respectively (Figure S5A, B). For

each mutant, we average the effects over different barcode variants (median 44 barcodes for single-

nucleotide changes) to reduce noise and barcode-driven variations (Figure S5C, D).

Mutations across a wide region surrounding the *cggR* cleavage site showed modest but detectable

impact on steady-state levels of the full-length mRNA (Figure 6D). Between -30 and +1 nucleotide

relative to the cleavage site, only a small number of mutations appear to reduce cleavage efficiency. The

strongest effects are seen at the cytosine at position -11 (denoted as C[-11]), and the effect sizes are at

most twofold. Interestingly, mutations of G at position -1 showed no statistically significant effect, even

though it is the strongest consensus among RNase Y cleavage sites across the transcriptome. The fact that

the G[-1] is not strictly required is consistent with our finding that 37% of the RNase Y cleavage sites

detected do not have a guanine at this position (Figure 5B). Therefore, mutations upstream of the *cggR*

cleavage site lend further support to our transcriptome-wide results that showed a weak consensus.

Downstream of the cleavage site, mutations in the hairpin region also produced limited effects on

full-length RNA levels. Single-nucleotide substitutions in this region did not have significant effects

(Figure 6D). Combinations of substitutions which further destabilize the hairpin also did not affect the

apparent cleavage efficiency (Figure 6H). Surprisingly, the only mutations in this structured region that

reduce cleavage are specific substitutions in a bulge of the downstream hairpin, including an A[+6]G

substitution (fourfold) and a C[+24]U substitution (twofold) (Figure 6D). These two single-substitution

mutants close the bulge and complete the hairpin, and other bulge-closing double substitutions at these

two locations also showed reduced cleavage. Together, these results suggest that the downstream hairpin

is not essential for cleavage, and that a stronger hairpin can in fact negatively impact cleavage.

Immediately downstream of the hairpin, an AU-rich region appears to play a role in cleavage.

Mutations to AUAAAUAU located at position +28 to +35 lead to increases in full-length RNA level by

up to 4-fold. In the folded RNA, this distal AU-rich region is brought to be adjacent to the cleavage site

by the intervening hairpin (Figure 6F). Although it is plausible that such spatial proximity enables the

distal region to recruit RNase Y or facilitate its activity, the fact that mutations in the hairpin region have

no detectable effects suggests that other mechanisms may be at play.

To further elucidate the role of the downstream hairpin, we tested the hypothesis that this

secondary structure mainly serves as a stabilizer for the cleaved product that encodes *gapA,* by preventing

exoribonucleolytic decay (Meinken et al., 2003). This role is distinct from the model for the

Staphylococcus aureus cggR-gapA operon, where the downstream hairpin has been suggested to specify

RNase Y cleavage sites (Marincola and Wolz, 2017; Le Scornet et al., 2024). To distinguish these roles,

we conducted deep mutational scanning on the accumulation of processed mRNA, measuring 349,266

mutant variants in total, and compared it to the results on full-length mRNA levels. Unlike full-length

mRNA, the level of the downstream processed mRNA should depend on both cleavage efficiency and its

own half-life post-cleavage.

Indeed, most mutations observed to increase full-length mRNA level resulted in reduced

accumulation of the cleavage product, but not vice versa (Figure 6E). The effects of C[-11] mutations,

bulge-closing mutation, and distal AU-rich region are recapitulated when we detected the downstream

cleavage product. However, in contrast to the full-length RNA, mutations in the downstream hairpin

region were among the most deleterious to cleavage product accumulation. Consistent with the secondary

structure playing a role, double-substitution variants that complement these mutations (Figure 6F) have

dramatically reduced effect sizes, and so are stem mutations that maintain base pairing (through wobble

interactions, Figure 6E, red arrows). Considering all variants, including those with two or more

substitutions, we found a clear trend that a stronger downstream hairpin is predictive of the cleaved RNA

level (Figure 6G), in stark contrast with a lack of correlation for full-length RNA (Figure 6H). Finally, we

repeated our measurements of cleaved RNA abundance in the context of an RNase J1 depletion, and

found that, with the exception of closing the bulge, substitutions within the downstream stem no longer

have a measurable impact when exoribonucleolytic activity against the processed RNA is attenuated

(Figure S6A). Together, these results suggest that the downstream hairpin plays a much more substantial

role dictating cleavage product abundance than the efficiency of RNase Y cleavage.

To assess whether the observed role of proximal secondary structure is specific to the *cggR*

operon, we repeated the described measurements and analysis for a second cleavage site, located within

the *glnR* gene of the *glnRA* operon (Figure S7A). As before, this cleavage occurs after the first gene in an

operon and a stem-loop structure is predicted to form at the 5′ end of a stabilized downstream cleavage

product (Figure S7B). Measuring accumulation of the cleaved RNA, we captured 54,431 variants with

mutations occurring from positions -5 to +31 relative to the position of cleavage. For nucleotides inside

the stem, both single mutations (Figure S7C) and higher-order substitutions that disrupt this RNA folding

(Figure S7D) attenuated the accumulation of the cleavage product (up to approximately 4-fold),

recapitulating the pattern seen for *cggR*. When we quantified the accumulation of full-length RNA

(Figure S7E) or cleaved RNA upon RNase J1 depletion (Figure S7F), we found that disruption of the

stem did not appear to stabilize the full-length RNA. Together, these results suggest that the behavior seen

for *cggR* may be shared with other similarly structured mRNA processing sites.

Finally, we placed the *cggR* cleavage site into a second reporter construct encoding the

exogenous gene *tetM* to assess whether the features driving cleavage product accumulation are context-

dependent. Although most key effects are observed in both reporter constructs, the *tetM* reporter showed

deleterious substitutions in C[-19]G and, to a lesser degree, A[-6]G (Figure S6B). The nucleotide C[-19]

is situated in an otherwise purine-rich region, AGA **C** GAAGGAG, inside the *cggR* open reading frame.

We initially hypothesized that the C[-19]G substitution may affect cleavage by slowing down elongating

ribosomes through its Shine-Dalgarno-like sequence. However, the deleterious effects were still observed

when we introduced an upstream stop codon, suggesting that these substitutions do not act through

translation but other context-specific means (Figure S6C). These results reinforce the notion that the

determinants of RNase Y cleavage are not localized to a narrow region on the mRNA.

DISCUSSION

In this work, we present the first genome-scale map of *B. subtilis* endoribonuclease cleavages at single-

nucleotide resolution. Using stabilized end-sequencing, we identified over a thousand putative cleavage

events, allowing us to learn substrate preferences of the central endonucleases involved in *B. subtilis*

mRNA decay. This work revealed the importance of long-range RNA base pairing for RNase III cleavage

and confirmed that the toxin EndoA recognizes a hexameric rather than pentameric motif *in vivo*. We

additionally uncovered evidence for and potential specificity of a new ribonucleolytic activity previously

unknown in *B. subtilis*, motivating further work to understand RNA 5′ end processing and decay. Finally,

we identified hundreds of new positions of cleavage by the core endonuclease driving mRNA decay,

RNase Y. Combining this global cleavage mapping with a new massively parallel reporter assay, we

demonstrate that cleavage by this enzyme is specified by features distributed over many positions across

regions tens of nucleotides in length. This stands in stark contrast to enzymes like EndoA, which

recognizes a strict consensus cleavage sequence, and RNase III, which has a strongly stereotyped

cleavage behavior within secondary structures.

The broad suite of features that contribute to RNase Y processing at the *cggR* site helps explain

the highly degenerate motif that we observed through global cleavage site mapping. Our data support a

model in which RNase Y cleavage is localized not just by the sequence proximal to the cleaved bond, but

rather by combined contributions of features distributed as many as tens of nucleotides away from the

cleavage site. The ability for features distal in primary sequence to strongly influence cleavage may be

mediated by RNA folding or by higher-order RNA-protein and protein-protein interactions between these

features and components of the bacterial RNA degradosome. As such, the particular positions recognized

to specify cleavage at each site may be variable and highly context-specific. Interestingly, a common

feature we identified between many RNase Y sites, the presence of a -1G, only have a modest effect on

the accumulation of processed *cggR* RNA in our reporter system, if any. This discrepancy may suggest a

similar model to what is observed in in *S. aureus,* where an upstream guanosine fine-tunes the position of

cleavage but is not required for RNase Y activity (Le Scornet et al., 2024).

The absence of a clear consensus motif specifying RNase Y activity echoes previous findings for

RNase E in *E. coli* (Chao et al., 2017; Lin-Chao et al., 1994; McDowall et al., 1994). These observations

suggest that the code for RNA decay fundamentally differs from transcription and translation, whose rates

are influenced by well-defined interactions between the gene expression machinery and localized

sequence features within either promoters or ribosome binding sites (Taggart et al., 2021). Instead, decay-

initiating RNases such as RNase Y appear to be strongly influenced by features distal to the position of

cleavage, possibly facilitated by RNA secondary structure and RNA-protein interactions. mRNA stability

may also be strongly influenced by features extrinsic to primary RNA sequence. Decay can occur

concurrently with and may be influenced by transcription and translation (Condon, 2003; Yarchuk et al.,

1992), and the nonuniform distribution of decay machinery in bacterial cells also means that protein

binding and RNA localization may have a strong influence on RNA stability (Hamouche et al., 2020;

Lehnik-Habrink et al., 2011b; Moffitt et al., 2016). Robust prediction of endoribonuclease cleavage from

primary sequence will thus likely require integration of more modes of measurement and interrogation of

more cleavage sites than our current end-mapping approach and MPRA allow.

Although the presented global map of endonuclease activity expands our list of known

endoribonuclease cleavage positions by orders of magnitude, we do not expect it to be comprehensive.

First, due to size-selection steps in the end-sequencing protocol, we are unable to detect cleavage sites

within 15 nucleotides of a 5′ end or 40 nucleotides of a 3′ end of a transcription unit. Additionally, while

all known exoribonuclease activity in one direction is ablated in each of our experiments, secondary

endoribonuclease cleavage followed by activity of the unperturbed exoribonucleases acting in the

direction opposite may destabilize cleavage products, rendering them undetectable. Finally, activity by

unknown exoribonucleases may cause broadening or shifting of 3′ or 5′ ends, separating the peak pairs

and obscuring potential cleavage positions (as seen in Figure 4). The absence of signal in these data

therefore is not necessarily an indication that turnover of a particular RNA is independent of

endoribonucleolytic cleavage.

Unexpectedly, our analysis of EndoA cleavage sites, as well as the *glmS* ribozyme, revealed a

previously uncharacterized 5′ end trimming activity in *B. subtilis.* We have confirmed that this activity is

not dependent on the primary enzymes targeting mRNAs for decay, including RNases Y, III, J1, J2,

RppH, or EndoA. Whether this nuclease acts in a 5′-3′ exonucleolytic manner or through

endoribonucleolytic cleavage near 5′ ends remains to be determined. Further work is additionally

necessary to understand whether this enzyme acts to facilitate RNase J1 cleavage at substrates with a 5′

hydroxyl moiety, as such behavior would shape our understanding of how systems like the *glmS*

ribozyme regulate gene expression. Determining what gene encodes this activity is thus of great interest

moving forward.

Our finding that the full Y-complex is not strictly required for processing of *B. subtilis* operons

suggests a refinement of our understanding of this complex in RNA maturation and decay. Although our

data support that loss of a component of the Y-complex strongly attenuates cleavage at the operons it

canonically regulates, persistence of weak cleavage signatures suggests that the presence of YlbF is not a

strict requirement for RNase Y activity at these sites. This residual cleavage may represent dispensability

of the Y-complex as a whole or dispensability of YlbF itself for Y-complex activity. Indeed, recent work

by Dubnau *et al.* has suggested that, of the Y-complex members, only YaaT (RicT) stably associates with

RNase Y to form the functional complex required for operon maturation (Dubnau et al., 2023). It remains

to be tested whether deletion of YaaT eliminates the low-level cleavage activity we detect in the absence

of YlbF.

Mutational scanning of the *cggR* mRNA processing site revealed that downstream secondary

structure primarily influences *cggR* operon stoichiometry through stabilization of the product of cleavage

rather than driving cleavage itself. This result, while consistent with early studies of this operon in *B.*

subtilis (Meinken et al., 2003), may appear to contradict studies in *S. aureus* that suggest presence of a

downstream hairpin is a key driver of RNase Y cleavage (Marincola and Wolz, 2017; Le Scornet et al.,

2024). It is worth noting, however, that *S. aureus* RNase Y targets a more limited set of RNAs than its *B.*

subtilis homolog and regulation of cleavage may differ between the two enzymes (Khemici et al., 2015;

Marincola et al., 2012). Additionally, the single base substitutions investigated here are a milder

perturbation than used in other studies of the *cggR* operon (Le Scornet et al., 2024), and when secondary

structure is strongly ablated in our experiments, a mild stabilization of the full-length construct is

observed (Figure 6H). It is thus possible that although the effect of secondary structure on cleavage

product stability is more pronounced, downstream RNA structure is playing a dual role in this system.

The approach described in this work has the potential to be generalized to study RNA decay in

other bacteria for which exonuclease activity can be ablated or transiently depleted, representing a new

approach to discover factors influencing RNA decay. Additionally, application of this approach to

conditions beyond exponential growth in rich medium will provide an insight into regulation of

endonuclease activity and RNA decay more broadly. The high-resolution picture of RNA decay initiation

provided by this approach, particularly when integrated with future measurements of RNA half-life, will

inform future quantitative predictions of RNA stability and gene expression levels from primary sequence

in Gram-positive bacteria.

MATERIALS AND METHODS

Strains and strain construction

For all Rend-seq, 5′ end sequencing, and 3′ end sequencing experiments, excluding the Rend-seq

measurement of an *rnjB* deletion with RNase J1 depletion, derivatives of W168 were used (Table S1). All

transformations for strain construction were performed using standard protocols relying on natural *B.*

subtilis competence and flanking homology mediated recombination using the media and protocol

described in (Parker et al., 2020). BJT070 was created through SPP1 phage transduction of the *ylbF::erm*

allele from BJT034 into CCB396 (Yasbin and Young, 1974). To create BJT074, gDNA from BJT034 was

first transformed into SSB1002 to generate a *ylbF* knockout in a W168 background, BJT069. From this,

SPP1 phage transduction from CCB434 was used to construct an *rnjA ylbF* double knockout, BJT074.

CCB390 was constructed by first transferring an inducible RNase J1 allele (first described in (Britton et

al., 2007)) into W168 (strain SSB1002), followed by deletion of *rnjA* through transformation with gDNA

from CCB434. To construct BJT231, genomic DNA from GLB230 was transformed into CCB390,

supplementing with 2% xylose in the liquid and solid media throughout the transformation protocol. To

construct BJT129, CCB390 was transformed with genomic DNA from an *rppH* knockout, strain

BKE30630 (Koo et al., 2017) .

A supercompetent *B. subtilis* strain BJT200, based on (Zhang and Zhang, 2011), was used for

experiments that do not require depletion or induction of RNase J1. This strain was constructed from

W168 *B. subtilis* strain SSB1002 through insertion of an anhydrotetracycline (aTc) inducible *comK* allele

at the *ganA* locus by transformation of ScaI-linearized plasmid pJT117. For experiments in which an

inducible RNase J1 was required, strain BJT201 was utilized. Similar to BJT200, this strain contains an

inducible *comK* gene. BJT201 is a derivative of strain CCB390, a W168 *B. subtilis* strain with xylose

inducible RNase J1, constructed through transformation with pJT117. These strains were used for all

MPRA experiments except the *tetM-cggR* data, which were collected in a *trpC* 168 strain. Plasmid

pJT117 and MPRA plasmids pJT028 and pJT124 were constructed by isothermal assembly.

Cell growth and harvesting (Rend-seq, 5′-, and 3′-end sequencing)

All Rend-seq, 5′-, and 3′-end sequencing experiments were performed using cells grown in 2xYT,

supplemented with 2% xylose or glucose for RNase J1 induction or depletion as necessary. Except when

we sought to deplete RNase J1, strains with an inducible RNase J1 allele were consistently grown in the

presence of 2% xylose. For all sequencing experiments, overnight cultures in 2xYT were back-diluted to

an OD_{600} of 0.0002 in 20 mL prewarmed 2xYT and grown to an OD_{600} of 0.2, at which point they were

harvested. For any strain with an RNase J1 knockout, cultures were vortexed prior to and following

inoculation to minimize clumping. To achieve RNase J1 depletion, 10 times the required volume of

overnight culture was collected, washed twice with 1 mL pre-warmed 2xYT, and resuspended in 1 mL

2xYT. 100 µL of this resuspension was used to inoculate the final culture. Samples were harvested for

sequencing by adding 7 mL of culture to 7 mL of ice cold methanol, pelleting, decanting, and flash

freezing with liquid nitrogen. Cell pellets were stored at -80°C prior to RNA extraction.

End-enriched RNA sequencing and 3′ end sequencing

Prior to RNA extraction, with the exception of our first Rend-seq experiment on the exonuclease-

deficient strains (profiling wild-type 168, 4-exo, and Δ*rnjA* strains with and without *ylbF* knockout), cell

pellets were washed with 10-15 mL ice cold 10 mM Tris pH 7.0 to remove contaminating nucleic acids

that precipitate from 2xYT in methanol. For Rend-seq, RNA was extracted from cell pellets using either

an RNeasy Mini Kit (Qiagen) with on-column DNase treatment, an RNeasy Plus Mini Kit (Qiagen) or

through RNAsnap extraction (Stead et al., 2012) followed by Turbo DNase treatment. All 3′ end

sequencing samples were purified through RNAsnap. In the RNAsnap protocol, cells were resuspended in

500 µL extraction solution (95% Formamide, 1% BME, 0.025% SDS, 18 mM EDTA) and transferred to a

2 mL tube containing 200 µL chilled 0.1 mm diameter zirconia beads. These tubes were vortexed for 10

minutes using a horizontal vortex adapter and subsequently incubated at 95 °C for 7 minutes. Samples

were spun at 15,000 g, after which 400 µL was removed and diluted 1:5 with 1.6 mL DEPC-treated

water. This mixture was precipitated with isopropanol and resuspended in 10 mM Tris, pH 7.0. At least

30 µg of RNA was then treated for 30 minutes at 37°C with 10 µL Turbo DNase in a 100 µL reaction.

Following RNA extraction and gDNA depletion, 20 µg of purified RNA was treated

with MICROBExpress Bacterial mRNA Enrichment Kit (ThermoFisher) per the manufacturer's

instructions and precipitated with isopropanol.

Rend-seq libraries were prepared as described previously (Lalanne et al., 2018). In short, rRNA

depleted RNA was fragmented for 25 seconds (ThermoFisher #AM8740) and fragments of sizes 15-45

were size selected on a 15% TBE-Urea polyacrylamide gel (ThermoFisher). RNA was dephosphorylated

with T4 PNK (NEB) and 3 pmol of this dephosphorylated RNA was ligated to an adenylated adaptor

using truncated T4 RNA ligase 2 K277Q (Jonathan Weissman). Ligated RNA was purified by denaturing

polyacrylamide size selection and reverse transcribed using SuperScript III (ThermoFisher) with a custom

primer oCJ485. This cDNA was then circularized using CircLigase (Lucigen) and barcodes and full

Illumina adapters were added through PCR using Phusion (NEB).

3′ end sequencing libraries were prepared in a manner similar to Rend-seq, with deviation in the

order of reactions (Herzel et al., 2022). Prior to fragmentation, 1.8 µg of rRNA depleted RNA was ligated

to an adenylated adapter as described for Rend-seq. This ligation product was then fragmented and RNA

species of sizes 33-63 nt were purified via denaturing polyacrylamide gel extraction. From this point

onwards, the library preparation is identical to Rend-seq. Note that this protocol lacks a

dephosphorylation step prior to 3′ end ligation, such that RNA ends with a 3′ phosphate will not be

captured by this protocol. Both Rend-seq and 5′ end sequencing libraries were sequenced with 50 or 75 bp

single end reads on either a HiSeq 2000 or NextSeq500.

Multiplexed 5′ end sequencing

For all 5′ end sequencing samples, pellets were washed, and RNA was extracted using the RNAsnap

protocol and Turbo DNase treated as described for the Rend-seq and 3′ end sequencing experiments.

rRNA was depleted from 10 μg of purified RNA with the MICROBExpress Bacterial mRNA Enrichment

Kit, per manufacturer's instructions, precipitated, and resuspended in 10 mM Tris pH 7.0. 600 ng rRNA

depleted RNA was then ligated to 260 pmol of the RNA adapter oJT900 (Table S2) for 18 hours at 22 °C

using T4 RNA Ligase 1 (based on (Subtelny et al., 2014)).

Following ligation, RNA was fragmented for 2 minutes (ThermoFisher) and fragments of either

size 41-71 (corresponding to 15-45 plus a 26-nucleotide adapter) or 60-90 were isolated through

denaturing polyacrylamide gel extraction. The larger size window was chosen to increase effective

sequencing depth by removing dimerized fragments of oJT900, which were observed in high abundance

in sequencing libraries prepared with a 41-71 nt size selection. RNA fragments were dephosphorylated,

polyadenylated, and reverse transcribed following the BaM-seq multiplexed RNA sequencing protocol

(Johnson et al., 2023). In short, in a single tube RNA fragments are sequentially dephosphorylated with

T4 PNK (NEB), polyadenylated with *E. coli* poly(A) polymerase (NEB) and reverse transcribed with

SuperScript III (ThermoFisher) and a barcoded reverse transcription primer. The cDNA from each sample

generated by this reaction can then be pooled and purified by denaturing polyacrylamide gel

electrophoresis. To add Illumina adapters, pooled cDNA was amplified between 5-10 cycles with primers

oDP128 and oDP161 (Johnson et al., 2023) using Q5 DNA polymerase and purified by polyacrylamide

gel electrophoresis. 5′ end sequencing libraries were sequenced with 75 bp single end reads on a

NextSeq500.

Rend-seq, 5′-, and 3′-end sequencing data processing

Adapter or poly(A) sequence was trimmed from end sequencing and Rend-seq libraries using CutAdapt

(Martin, 2011) with parameters "-a 'CTGTAGGCACCATCAAT' -m 15" or "-a A{15} -m 15 -u 8 --

rename='{id} UMI={cut_prefix}'". 5′ end sequencing reads were additionally collapsed using the 8 bp

unique molecular identifier (UMI) at the start of each read using custom Python scripts, requiring

identical UMI sequences and read sequences within a hamming distance of 1 to be considered duplicates.

Trimmed reads were then mapped to the *S. cerevisiae* genome (SacCer3) using bowtie v1.1.2 with

parameters "-v 1 -k 1 –best –un" with retained unmapped reads subsequently mapped to the *B. subtilis*

genome (NC_000964.3) with the same parameters (Langmead et al., 2009). Mapping against yeast

removes contaminating 2xYT-derived sequences and allows estimation of the degree of contamination in

a particular sample. Using custom Python scripts, we converted the bowtie output to wig files, attributing

signal from reads with a mismatch at their first position to the adjacent nucleotide to address mismatches

due to non-templated addition during reverse transcription.

Identification of putative endonuclease sites

To identify putative sites of endoribonucleolytic cleavage, we identified positions for which an adjacent 5′

and 3′ end could be detected in our end sequencing approaches. To do so, we first normalized each dataset

to the total number of millions of CDS-mapped reads. At each position, we identified peaks in read

density by calculating a ratio of 5′ end signal at that position to the 90% winsorized average in a window

50 nucleotides upstream (for 3′ ends) or downstream (for 5′ ends) of that position, excluding a 2-

nucleotide gap on either side of the peak. For any adjacent positions (allowing a gap of 1 position) with a

ratio over background greater than 7.5, these positions were grouped for the purposes of downstream 5′

end signal quantification. At each position with >10 reads per million CDS-mapped reads (RPM) in 5′

end sequencing, we detected paired 3′ ends by calculating an equivalent ratio centered around the position

one nucleotide upstream in the 3′ end sequencing of the 4-exo deletion. Additionally, in the 50-nucleotide

window downstream of each position, we calculated the local density of mapped 3′ ends of the RNase J1

deletion Rend-seq data to establish that we were considering positions only in genes expressed in this

strain. Putative cleavage sites were called as positions for which the average local Rend-seq depth

exceeded 0.05 RPM and both the 3′ and 5′ end sequencing signals had a ratio of 7.5 over background.

Because the sequencing methods used in this work are unable to capture short RNA fragments

(<15 nt for Rend-seq and 3′ end sequencing and <15-34 nt for 5′ end sequencing, depending on the exact

size selection criteria used), we did not consider positions within 40 nucleotides of an RNA 3′ end. To

identify these ends, we called peaks in 3′-mapped RNase J1 Rend-seq data using a z-score based peak

calling method. At each position, a z-score is calculated as the signal at that position minus the average

signal in a window of 50 nucleotides on either side of the peak (excluding a gap of 2 nucleotides on either

side of the peak), divided by the standard deviation within this window. Any position for which the

calculated score exceeds 8 is flagged as a peak (i.e. an RNA 3′ end) in our Rend-seq data. Putative

cleavage positions located up to 40 nucleotides upstream of one of these 3′ end positions are discarded.

To count the number of putative cleavage positions across all our datasets, positions that were called in

either the W168 (CCB434) or phage-cured W168 (BG880) background were considered. If two adjacent

positions were called across these datasets, they were counted only a single position.

To attribute putative cleavage positions to specific endoribonucleases, we calculated the ratio of

5′ end sequencing signal at each position in an endonuclease knockout with RNase J1 deletion (or

depletion) to the average Rend-seq signal of the same strain in a corresponding window 50 nucleotides on

downstream of that position, excluding a 2-nt gap. To account for differences in sequencing depth

between samples, this 5′ end sequencing and Rend-seq signal at each position was normalized to the total

CDS-mapped reads in each sample. Additionally, signal from positions which were previously flagged as

grouped in the 5′ end sequencing were summed, with the cleavage assigned upstream of the 5′-most

position in the group. This ratio of 5′ end sequencing to Rend-seq signal is calculated for the

corresponding RNase J1 deficient strain that expresses the considered endonuclease (e.g. a phage-cured

RNase J1 deletion when considering RNase III). The ratio in the endoribonuclease expressing strain is

then divided by the ratio in the endoribonuclease knockout strain to produce a "sensitivity" score. This

approach is similar to that used to map RNase III sites in previous work (DiChiara et al., 2016).

Putative cleavage positions with a sensitivity score that exceeds 10.56 in a particular

endoribonuclease knockout are called as cleavage sites for that enzyme. This threshold was empirically

determined by fitting of the distribution of log-transformed sensitivity values for the RNase Y knockout

to the sum of two Gaussian distributions and finding the point at which 95% of this fit is contributed by

the underlying Gaussian with the larger mean (see Figure 5A). For this endonuclease assignment analysis,

any position for which the local Rend-seq density in the endonuclease knockout strain (as calculated

above) falls below 0.05 RPM is excluded to avoid spuriously calling positions in genes that are

downregulated in this condition. When calculating the fraction of putative cleavage sites that are

dependent on a particular protein, only those cleavage sites that were identified in an RNase J1 knockout

of the same background were considered (e.g. only sites appearing in the phage-cured RNase J1 knockout

strain BG880 were used when calculating the fraction of sites sensitive to RNase III knockout, which was

generated in the same phage-cured W168 background, see Table S1).

Determination of RNase III site duplex length, genomic distance, and structural distance

To identify pairs of RNase III cleavage positions that might represent intramolecular pairing, we first

identified all RNase III-dependent cleavage positions within 1 kb of one another. The distance between

sites in the primary sequence is the genomic distance as reported in Figure 3D. We then extracted the

intervening RNA sequence plus a 50 nucleotide buffer on either side for each pair of positions and

calculated its minimum free energy (MFE) structure using RNAfold from ViennaRNA v2.4.14 with

default parameters (Lorenz et al., 2011). We then used this MFE structure and a custom Python script to

calculate the length of the overhang generated (if any) by RNase III cleavage at this pair of sites. This

length is reported as structural distance in Figure 3E. The reported duplex length for each structure was

manually annotated, allowing for a maximum gap of two unpaired bases between base pairs.

RNase III sequence-structure motif analysis

For analyses of sequence content within RNA secondary structures recognized by RNase III, we

considered an expanded set of putative cleavage sites, using a threshold of >2 reads per million CDS-

mapping reads in 5′ end sequencing (rather than the previous >10 RPM). Results of putative cleavage

position identification and endonuclease sensitivity calculation using this threshold are summarized in

Tables S5,6. From this set of putative cleavage sites, identified pairs of sites within 1 kb of one another

and calculated a MFE secondary structure as described above. As before, we then used custom Python

scripts to quantify the overhang length generated by the two RNase III cleavages and retained any sites

which produced a 2-bp 3′ overhang.

Using the predicted MFE structures for these pairs of sites, we identified the base pair at each

position within the structure relative to the two positions of cleavage. The position-wise frequency of each

type of base-pairing interaction (A-U, U-A, G-C, C-G, U-G, or G-U) and fraction of sites for which that

position was paired was then determined. In cases where an asymmetric bulge was observed, those bases

were included as additional unpaired positions, rather than skipping over them in our indexing of base-

pairs. To compare our data to those previously-published for *E. coli* (Altuvia et al., 2018), we calculated

the difference between the frequency of each pairing type in our data and the equivalent observed at that

position by Altuvia et al.

Local secondary structure prediction for RNase Y sites

To determine folding propensity of sequences proximal to endoribonuclease cleavage sites, the *B. subtilis*

genome was tiled with 40-mers and folded with RNAfold from the ViennaRNA v2.4.14 package with

default parameters (Lorenz et al., 2011). The MFE values generated from this folding were assigned to

the center position within each k-mer and used to generate wig files containing positional folding

propensity. These values were extracted in 150 nucleotide windows centered around putative

endoribonuclease cleavage positions and a position-wise mean was calculated. If a cleavage site was

identified within 75 nucleotides of a 3′ or 5′ end present in wild-type W168 Rend-seq data (identified in

strain SSB1002 by the previously described z-score based peak calling method), that cleavage site was

not included in this analysis. To generate a background model that controls for sequence context in the -

1/+1 positions, this analysis was repeated 100 times sampling an equivalent number of random positions

within coding regions that are centered on GA dinucleotides (the most enriched dinucleotide flanking

putative RNase Y sites), excluding sites called as cleavage positions. From these predicted position-wise

average folding curves, the 25th and 75th percentile at each position across all curves was used to generate

the shaded region on Figure 5D.

Information content and k-mer logo generation

To generate sequence logos, the local sequence in a 30-nucleotide window surrounding all putative

cleavage sites for a given enzyme were identified and aggregated into a FASTA file. A background set of

30-mers evenly sampled every 100 nucleotides from *B. subtilis* coding regions (with sampling beginning

after the start codon) was generated as a background set. kpLogo v1.1 (Wu and Bartel, 2017) was then

used to generate information content and k-mer logos with parameters "-bgfile %s -startPos 16 -pc 0.01",

with %s pointing to our background set.

763
MPRA plasmid library construction

Plasmid libraries were constructed through restriction cloning of randomized mRNA processing site

sequences into suitable reporter plasmids (pJT028 or pJT124 for *tetM* or *aprE* experiments). Inserts for

plasmid libraries were constructed through a two-step protocol using mutated oligonucleotides generated

by IDT as a template. Oligonucleotides were designed with a 9% error rate (3% per non-native

nucleotide) over the specified positions (see Table S7 for enumeration of experiments and their variable

regions). In the first step, 10 fmol of oligo was PCR amplified for 12 cycles using Q5 DNA polymerase

(NEB). To produce a *cggR*-derived insert with mutations downstream of the cleavage site (experiments

2a, 2b, 4a, 4b, and 6), oJT837 was amplified with oJT851/852. To prepare the *glnR* insert (experiments

8a, 8b, 9), oJT839 was amplified with oJT857/858. For the *cggR* insert with upstream mutations

(experiments 1a, 1b, 3, 5, 7), instead of PCR amplification, oJT592 and oJT593 were mixed and extended

through one cycle of denaturing, annealing, and extension with Q5. These products were diluted 1:10 into

a second reaction in which they were amplified and barcoded through 4 cycles of PCR with Q5 using

oJT548/594 and oJT855/859 for all *cggR* and *glnR* inserts respectively. Amplified, barcoded inserts were

then cleaned up with a DNA Clean & Concentrator 5 column (Zymo Research).

To construct the plasmid pool, the amplified insert and 4 μg plasmid backbone were digested

with SalI-HF and NheI-HF for 60 minutes at 37°C and purified with a DNA Clean & Concentrator 5

column. 400 ng cut plasmid and insert were mixed at an approximately 1:10 ratio and ligated through a 5-

15 minute treatment with QuickLigase (NEB) at 25°C. This reaction was then cleaned up with a DNA

Clean & Concentrator 5 column and transformed into electrocompetent *E. coli* (NEB 10-beta or NEB 5-

alpha cells). Transformants were plated on LB with 100 μg/mL carbenicillin in 245 mm square bioassay

dishes (Corning) and grown at 37°C overnight. Cells were then scraped off the plates and pelleted for

ZymoPure II Maxiprep plasmid extraction (Zymo Research).

787
Transformation and integration of MPRA library into B. subtilis genome

To introduce our plasmid libraries into *B. subtilis*, we utilized an inducible competence system mediated

by an aTc-inducible *comK* allele introduced into BJT200 and BJT201, based on (Zhang and Zhang,

2011). A single colony was picked into LB (with 2% xylose for BJT201) and allowed to grow to an OD_{600}

of 1.0 at 37°C with vigorous shaking. At this point, 10 ng/mL aTc was added to the culture to induce

expression of ComK. Cells were shaken at 37°C for an additional 2 hours to allow induction of

competence, at which point 100 μL per mL culture of plasmid pool linearized with either ScaI-HF or AfeI

(NEB) was added to the culture. This restriction digest was prepared per the manufacturers protocol with

1 μg plasmid per 50 μL reaction. The cells were cultured with DNA for 1.5 hours, at which point the

cultures were pelleted at room temperature, resuspended in residual media, and plated on 245 mm

bioassay dishes containing 100 μg/mL spectinomycin or 5 μg/mL chloramphenicol. The *cggR-tetM*

library was transformed using standard protocols relying on natural competence. After an overnight

growth at 37°C, cells were scraped into LB with the appropriate antibiotic and approximately 500 million

cells were back-diluted into 200 mL LB with antibiotic. This culture was grown for approximately 4-8

hours, and several 1 mL aliquots of culture were mixed with equal volume 40% glycerol and frozen at -

80°C to prepare stocks for future experiments.

Cell growth and harvesting (MPRA)

For all MPRA experiments, cells were grown in LB medium with an inducer (20 ng/mL

anhydrotetracycline or 2% xylose) when noted. For all experiments in an inducible RNase J1 background,

2% xylose was consistently used to maintain RNase J1 expression, which was substituted for 2% glucose

when RNase J1 depletion was required. To collect samples, a single cryovial of a *B. subtilis* library was

thawed, washed twice with 10 mL LB to remove glycerol, and grown overnight in 200 mL of LB at 37°C

in a 2.8 L flask with vigorous shaking. These cultures were then back-diluted to an OD_{600} of 0.0003 in

200 mL LB with library inducer and grown at 37°C until an OD_{600} of 0.3, at which point samples were

harvested. Two cell pellets were harvested per sample, one for gDNA extraction and one for RNA

extraction. For RNA, 7 mL culture was added to 7 mL ice cold methanol, pelleted, decanted, and flash-

frozen in liquid nitrogen. For gDNA, 14 mL of culture was pelleted, decanted, and frozen in liquid

nitrogen. Cell pellets were stored at -80°C until nucleic acid extraction.

MPRA RNA and DNA sequencing

Three sequencing libraries were prepared for each sample: one to quantify the barcodes in cellular RNA,

one to quantify barcodes in the gDNA, and one to map barcodes to variant sequence. To prepare the RNA

barcode quantification library, RNA was first extracted from frozen cell pellets with gDNA depletion

using an RNeasy Plus Mini Kit (Qiagen). For experiments 1a, 1b, 2a, 2b, and 9 (see Table S7), 40 µg of

total RNA was reverse transcribed using primer oJT442 and SuperScript III. This cDNA was then run on

a 6% TBE-Urea gel, and the following size selection ranges were gel purified: 300-500 nt for full-length

cggR and *glnR* constructs (experiments 1a, 2a, 9) and 100-250 nt for cleaved *cggR* constructs

(experiments 1b and 2b). For the remaining experiments, cleaved RNA was isolated by size selection of

RNAs ranging from 200-400 nt for *tetM-cggR* sites, 150-300 nt for *aprE-cggR* sites, and 150-350 nt for

aprE-glnR sites on a 6% TBE-Urea gel (Novex). The size selected product was reverse transcribed using

primer oJT442 and SuperScript III (Thermo Fisher) and size selected on 6% TBE-Urea gel, extracting

approximately 120 to 200 nucleotides for *cggR* libraries and 100 to 200 for *glnR*. For all experiments,

size-selected cDNA was amplified using Phusion DNA polymerase (NEB), adding a barcode and

Illumina adapters with a Truseq indexing primer and an isoform-specific primer (see Table S2 for primer

sequences). These isoform-specific primers were oJT441/oJT1290 for cleaved/full-length products from

cggR constructs or oJT890/oJT1291 for cleaved/full-length products from *glnR* constructs. The resulting

libraries were sequenced with 50 or 75 bp single end reads on an Illumina NextSeq500.

To prepare both libraries for gDNA barcode quantification and barcode to variant sequence

mapping, a single gDNA sample was prepared from a frozen cell pellet using the Wizard Genomic DNA

Purification Kit (Promega). Both gDNA-derived libraries were generated using a two-step Phusion PCR

protocol. An initial amplicon was generated through 2 cycles of PCR using from 4 µg of purified nucleic

acid per 50 µL PCR reaction (reaction scaled up as needed to minimize UMI collapse in more complex

libraries) with primers oJT444/445 for *cggR* upstream mutations, oJT1125/445 for *cggR* downstream

mutations, and oJT891/445 for *glnR* downstream mutations. The untranslated *cggR* library required a

longer amplicon to capture the upstream stop codon and was instead amplified with oJT942 and oJT445.

This first PCR reaction was cleaned up with a double-sided Select-a-Size DNA Clean & Concentrator

column (Zymo) and a DNA Clean & Concentrator 5 column (Zymo). A quarter of this reaction was then

used for the second round of PCR, in which two different reactions were used to prepare barcode

quantification and barcode to variant sequence mapping libraries. For barcode quantification, the second-

round amplification was performed using the same primers as used for quantifying cleaved RNA

abundance (oJT441 or oJT890 with our Truseq indexing primer) and size selected on an 8% TBE gel.

These libraries were sequenced with 50 or 75 bp single end reads on an Illumina NextSeq500. Libraries

mapping barcodes to variant sequence were amplified using primers oJT446/447, size selected on an 8%

TBE gel, and sequenced with either 75 bp or 150 bp paired end reads on an Illumina MiSeq.

Barcode to variant sequence mapping and barcode quantification

Barcode quantification and variant sequence identification was handled using custom scripts written in

Python. To determine the variant sequence associated with each barcode, the paired end reads from the

gDNA-derived libraries were directly parsed as FASTQ files, with one read capturing the barcode and the

other the variable region of the construct. Variants with at least one mutation in the constant region

proceeding the barcode were discarded. To avoid rare mutations or sequencing errors confounding our

variant sequence mapping, variant sequence assignment required at least three reads per barcode. If,

among these reads, the second-most frequent sequence was at least 25% as frequent as the top sequence,

the barcode was discarded entirely.

Barcodes were counted by directly parsing the single-end sequencing reads derived from both

gDNA and RNA in a FASTQ format, with the barcode appearing at a fixed position within the read. As

before, any variant with mutations in the constant sequence proceeding the barcode was discarded.

Additionally, any two barcodes of identical sequence and unique molecular identifier were counted as a

single read. The number of unique reads mapping to each variant barcode was calculated for both the

RNA and gDNA-derived samples, and the ratio of RNA to gDNA-derived reads was calculated to extract

the relative accumulation of processed mRNA for each variant. Barcodes can then be grouped by

associated mutations, thresholded based on read count, and normalized to the median wild-type barcode's

ratio.

873
874

AUTHOR CONTRIBUTIONS:
J.T. and G.-W.L. conceptualized project, designed experiments and analysis, and wrote the manuscript
with input from all co-authors. J.T. collected data and performed analysis. H.L. and J.D. contributed to the
design and implementation of the MPRA experiments. J.-B.L. collected preliminary Rend-seq data for
exonuclease knockouts. C.C., S.D., and F. B. constructed strains, provided feedback on analysis, and
helped in conceptualization of project.

881
ACKNOLWEDGEMENTS:
The authors thank David Bechhofer for sharing the phage-cured RNase III knockout strains, Byoung-Mo
Koo and Carol Gross for sharing the RNase J2 knockout strain, the MIT BioMicro Center for advice and
high-throughput DNA sequencing, Sean McGeary for feedback on early versions of the manuscript, and

Michael Laub and David Bartel for feedback in developing this project. Thanks additionally to all
members of the Li Lab for valuable discussion. This research is supported by a National Science
Foundation Graduate Research Fellowship (to J.C.T.) and the NSF CAREER Award MCB1844668. The
C.C. lab is supported by funds from the CNRS (UMR 8261), Université Paris Cité, and the Agence
Nationale de la Recherche (CoNoCo and LabEx Dynamo).

REFERENCES:

Altuvia, Y., Bar, A., Reiss, N., Karavani, E., Argaman, L., and Margalit, H. (2018). In vivo cleavage rules
and target repertoire of RNase III in Escherichia coli. Nucleic Acids Res. *46*, 10380–10394.
Bechhofer, D.H., and Deutscher, M.P. (2019). Bacterial ribonucleases and their roles in RNA metabolism.
Crit. Rev. Biochem. Mol. Biol. *54*, 242–300.
Benda, M., Woelfel, S., Faßhauer, P., Gunka, K., Klumpp, S., Poehlein, A., Kálalová, D., Šanderová, H.,
Daniel, R., Krásný, L., et al. (2021). Quasi-essentiality of RNase Y in Bacillus subtilis is caused
by its critical role in the control of mRNA homeostasis. Nucleic Acids Res. *49*, 7088–7102.
Braun, F., Durand, S., and Condon, C. (2017). Initiating ribosomes and a 5- /3-UTR interaction control
ribonuclease action to tightly couple B. Subtilis hbs mRNA stability with translation. Nucleic
Acids Res. *45*, 11386–11400.
Brewster, R.C., Jones, D.L., and Phillips, R. (2012). Tuning promoter strength through RNA polymerase
binding site design in Escherichia coli. PLoS Comput. Biol. *8*, e1002811.
Britton, R.A., Wen, T., Schaefer, L., Pellegrini, O., Uicker, W.C., Mathy, N., Tobin, C., Daou, R., Szyk,
J., and Condon, C. (2007). Maturation of the 5′ end of Bacillus subtilis 16S rRNA by the essential
ribonuclease YkqC/RNase J1. Mol. Microbiol. *63*, 127–138.
Broglia, L., Lécrivain, A., Renault, T.T., Ahmed-begrich, R., Rhun, A. Le, and Charpentier, E. (2020). An
RNA-seq based comparative approach reveals the transcriptome-wide interplay between 3'-to-5'
exoRNases and RNase Y. Nat. Commun. *11*.
Bruscella, P., Shahbabian, K., Laalami, S., and Putzer, H. (2011). RNase Y is responsible for uncoupling
the expression of translation factor IF3 from that of the ribosomal proteins L35 and L20 in
Bacillus subtilis. Mol. Microbiol. *81*, 1526–1541.
Cerullo, F., Filbeck, S., Patil, P.R., Hung, H.C., Xu, H., Vornberger, J., Hofer, F.W., Schmitt, J., Kramer,
G., Bukau, B., et al. (2022). Bacterial ribosome collision sensing by a MutS DNA repair ATPase
paralogue. Nature *603*, 509–514.
Chao, Y., Li, L., Girodat, D., Förstner, K.U., Said, N., Corcoran, C., Śmiga, M., Papenfort, K., Reinhardt,
R., Wieden, H.J., et al. (2017). In Vivo Cleavage Map Illuminates the Central Role of RNase E in
Coding and Non-coding RNA Pathways. Mol. Cell *65*, 39–51.
Chen, H., Shiroguchi, K., Ge, H., and Xie, X.S. (2015). Genome-wide study of mRNA degradation and
transcript elongation in E scherichia coli. Mol. Syst. Biol. *11*, 781.
Collins, J.A., Irnov, I., Baker, S., and Winkler, W.C. (2007). Mechanism of mRNA destabilization by the
glmS ribozyme. Genes Dev. *21*, 3356–3368.
Commichau, F.M., Rothe, F.M., Herzberg, C., Wagner, E., Hellwig, D., Lehnik-Habrink, M., Hammer,
E., Völker, U., and Stülke, J. (2009). Novel activities of glycolytic enzymes in Bacillus subtilis:
interactions with essential proteins involved in mRNA processing. Mol. Cell. Proteomics *8*,
1350–1360.
Condon, C. (2003). RNA Processing and Degradation in Bacillus subtilis. Microbiol. Mol. Biol. Rev. *67*,
157–174.
Cuchillo, C.M., Nogués, M.V., and Raines, R.T. (2011). Bovine pancreatic ribonuclease: fifty years of the
first enzymatic reaction mechanism. Biochemistry *50*, 7835–7841.
Dar, D., Shamir, M., Mellin, J.R., Koutero, M., Stern-Ginossar, N., Cossart, P., and Sorek, R. (2016).
Term-seq reveals abundant ribo-regulation of antibiotics resistance in bacteria. Science *352*,
aad9822.
Deloughery, A., Dengler, V., Chai, Y., and Losick, R. (2016). Biofilm formation by Bacillus subtilis
requires an endoribonuclease-containing multisubunit complex that controls mRNA levels for the

937   matrix gene repressor SinR. Mol. Microbiol. *99*, 425–437.

938 DeLoughery, A., Lalanne, J.-B., Losick, R., and Li, G.-W. (2018). Maturation of polycistronic mRNAs
939   by the endoribonuclease RNase Y and its associated Y-complex in *Bacillus subtilis*. Proc. Natl.
940   Acad. Sci. *115*, E5585–E5594.

941 DiChiara, J.M., Liu, B., Figaro, S., Condon, C., and Bechhofer, D.H. (2016). Mapping of internal
942   monophosphate 5′ ends of Bacillus subtilis messenger RNAs and ribosomal RNAs in wild-type
943   and ribonuclease-mutant strains. Nucleic Acids Res. *44*, 3373–3389.

944 Dubnau, E., DeSantis, M., and Dubnau, D. (2023). Formation of a stable RNase Y-RicT (YaaT) complex
945   requires RicA (YmcA) and RicF (YlbF). MBio *14*, e0126923.

946 Durand, S., Gilet, L., Bessières, P., Nicolas, P., and Condon, C. (2012a). Three essential ribonucleases-
947   RNase Y, J1, and III-control the abundance of a majority of bacillus subtilis mRNAs. PLoS
948   Genet. *8*.

949 Durand, S., Gilet, L., and Condon, C. (2012b). The essential function of B. subtilis RNase III is to silence
950   foreign toxin genes. PLoS Genet. *8*, e1003181.

951 Durand, S., Braun, F., Lioliou, E., Romilly, C., Helfer, A.-C., Kuhn, L., Quittot, N., Nicolas, P., Romby,
952   P., and Condon, C. (2015). A Nitric Oxide Regulated Small RNA Controls Expression of Genes
953   Involved in Redox Homeostasis in Bacillus subtilis. PLOS Genet. *11*, e1004957.

954 Figaro, S., Durand, S., Gilet, L., Cayet, N., Sachse, M., and Condon, C. (2013). Bacillus subtilis mutants
955   with knockouts of the genes encoding ribonucleases RNase Y and RNase J1 are viable, with
956   major defects in cell morphology, sporulation, and competence. J. Bacteriol. *195*, 2340–2348.

957 Gan, J., Tropea, J.E., Austin, B.P., Court, D.L., Waugh, D.S., and Ji, X. (2006). Structural insight into the
958   mechanism of double-stranded RNA processing by ribonuclease III. Cell *124*, 355–366.

959 Gilet, L., Dichiara, J.M., Figaro, S., Bechhofer, D.H., and Condon, C. (2015). Small stable RNA
960   maturation and turnover in Bacillus subtilis. Mol. Microbiol. *95*, 270–282.

961 Gordon, G.C., Cameron, J.C., and Pfleger, B.F. (2017). RNA sequencing identifies new RNase III
962   cleavage sites in Escherichia coli and reveals increased regulation of mRNA. MBio *8*.

963 Guler, P., Bendori, S.O., Borenstein, T., Aframian, N., Kessel, A., and Eldar, A. (2024). Arbitrium
964   communication controls phage lysogeny through non-lethal modulation of a host toxin–antitoxin
965   defence system. Nat. Microbiol. *9*, 150–160.

966 Hamouche, L., Billaudeau, C., Rocca, A., Chastanet, A., Ngo, S., Laalami, S., and Putzer, H. (2020).
967   Dynamic membrane localization of RNase Y in bacillus subtilis. MBio *11*.

968 Herskovitz, M.A., and Bechhofer, D.H. (2000). Endoribonuclease RNase III is essential in Bacillus
969   subtilis. Mol. Microbiol. *38*, 1027–1033.

970 Herzel, L., Stanley, J.A., Yao, C.-C., and Li, G.-W. (2022). Ubiquitous mRNA decay fragments in E. coli
971   redefine the functional transcriptome . Nucleic Acids Res. *50*, 5029–5046.

972 Ingle, S., Chhabra, S., Chen, J., Lazarus, M.B., Luo, X., and Bechhofer, D.H. (2022). Discovery and
973   initial characterization of YloC, a novel endoribonuclease in Bacillus subtilis. RNA *28*, 227–238.

974 Jahn, N., and Brantl, S. (2013). One antitoxin-two functions: SR4 controls toxin mRNA decay and
975   translation. Nucleic Acids Res. *41*, 9870–9880.

976 Johnson, G.E., Parker, D.J., Lalanne, J., Parker, M.L., and Li, G. (2023). BaM-seq and TBaM-seq , highly
977   multiplexed and targeted RNA-seq protocols for rapid , low-cost library generation from bacterial
978   samples. *5*, 1–11.

979 Khemici, V., Prados, J., Linder, P., and Redder, P. (2015). Decay-Initiating Endoribonucleolytic Cleavage
980   by RNase Y Is Kept under Tight Control via Sequence Preference and Sub-cellular Localisation.
981   PLoS Genet. *11*, 1–27.

982 Koo, B.M., Kritikos, G., Farelli, J.D., Todor, H., Tong, K., Kimsey, H., Wapinski, I., Galardini, M.,
983   Cabal, A., Peters, J.M., et al. (2017). Construction and Analysis of Two Genome-Scale Deletion
984   Libraries for Bacillus subtilis. Cell Syst. *4*, 291-305.e7.

985 Korobeinikova, A., Laalami, S., Berthy, C., and Putzer, H. (2023). RNase Y Autoregulates Its Synthesis
986   in Bacillus subtilis. Microorganisms *11*, 1374.

987 Lalanne, J.B., Taggart, J.C., Guo, M.S., Herzel, L., Schieler, A., and Li, G.W. (2018). Evolutionary

988    Convergence of Pathway-Specific Enzyme Expression Stoichiometry. Cell *173*, 749-761.e38.

989 Langmead, B., Trapnell, C., Pop, M., and Salzberg, S.L. (2009). Ultrafast and memory-efficient
990    alignment of short DNA sequences to the human genome. Genome Biol. *10*.

991 Lehnik-Habrink, M., Schaffer, M., Mäder, U., Diethmaier, C., Herzberg, C., and Stülke, J. (2011a). RNA
992    processing in Bacillus subtilis: Identification of targets of the essential RNase Y. Mol. Microbiol.
993    *81*, 1459–1473.

994 Lehnik-Habrink, M., Newman, J., Rothe, F.M., Solovyova, A.S., Rodrigues, C., Herzberg, C.,
995    Commichau, F.M., Lewis, R.J., and Stülke, J. (2011b). RNase Y in Bacillus subtilis: A natively
996    disordered protein that is the functional equivalent of RNase E from Escherichia coli. J. Bacteriol.
997    *193*, 5431–5441.

998 Lehnik-Habrink, M., Lewis, R.J., Mäder, U., and Stülke, J. (2012). RNA degradation in Bacillus subtilis:
999    An interplay of essential endo- and exoribonucleasesmmi. Mol. Microbiol. *84*, 1005–1017.

1000 Leroy, M., Piton, J., Gilet, L., Pellegrini, O., Proux, C., Coppée, J., Figaro, S., and Condon, C. (2017).
1001    Rae1/YacP, a new endoribonuclease involved in ribosome-dependent mRNA decay in
1002    Bacillus subtilis . EMBO J. *36*, 1167–1181.

1003 Lin-Chao, S., Wong, T.-T., Mcdowall, K.J., and Cohens, S.N. (1994). Effects of Nucleotide Sequence on
1004    the Specificity of me-dependent and RNase E-mediated Cleavages of RNA I Encoded by the
1005    pBR322 Plasmid. J. Biol. Chem. *269*, 10797–10803.

1006 Lorenz, R., Bernhart, S.H., Zu Siederdissen, C.H., Tafer, H., Flamm, C., Stadler, P.F., and Hofacker, I.L.
1007    (2011). ViennaRNA Package 2.0. Algorithms Mol. Biol. *6*, 1–14.

1008 Ludwig, H., Homuth, G., Schmalisch, M., Dyka, F.M., Hecker, M., and Stülke, J. (2001). Transcription of
1009    glycolytic genes and operons in Bacillus subtilis: Evidence for the presence of multiple levels of
1010    control of the gapA operon. Mol. Microbiol. *41*, 409–422.

1011 Mandell, Z.F., Oshiro, R.T., Yakhnin, A. V, Kashlev, M., Kearns, D.B., and Babitzke, P. (2021). NusG is
1012    an intrinsic transcription termination factor that stimulates motility and coordinates global gene
1013    expression with NusA. Elife *10:e61880*.

1014 Marincola, G., and Wolz, C. (2017). Downstream element determines RNase y cleavage of the saePQRS
1015    operon in Staphylococcus aureus. Nucleic Acids Res. *45*, 5980–5994.

1016 Marincola, G., Schäfer, T., Behler, J., Bernhardt, J., Ohlsen, K., Goerke, C., and Wolz, C. (2012). RNase
1017    Y of Staphylococcus aureus and its role in the activation of virulence genes. Mol. Microbiol. *85*,
1018    817–832.

1019 Martin, M. (2011). Cutadapt removes adapter sequences from high-throughput sequencing reads.
1020    EMBnet.Journal *17*, 10.

1021 McDowall, K.J., Lin-Chaol, S., and Cohen, S.N. (1994). A+U content rather than a particular nucleotide
1022    order determines the specificity of RNase E cleavage. J. Biol. Chem. *269*, 10790–10796.

1023 Meinken, C., Blencke, H.M., Ludwig, H., and Stülke, J. (2003). Expression of the glycolytic gapA operon
1024    in Bacillus subtilis: Differential syntheses of proteins encoded by the operon. Microbiology *149*,
1025    751–761.

1026 Meißner, C., Jahn, N., and Brantl, S. (2016). In Vitro characterization of the type i toxin-antitoxin system
1027    bsrE/SR5 from Bacillus subtilis. J. Biol. Chem. *291*, 560–571.

1028 Moffitt, J.R., Pandey, S., Boettiger, A.N., Wang, S., and Zhuang, X. (2016). Spatial organization shapes
1029    the turnover of a bacterial transcriptome. Elife *5*, 1–22.

1030 Noone, D., Salzberg, L.I., Botella, E., Bäsell, K., Becher, D., Antelmann, H., and Devine, K.M. (2014). A
1031    highly unstable transcript makes CwlO D,L-endopeptidase expression responsive to growth
1032    conditions in bacillus subtilis. J. Bacteriol. *196*, 237–247.

1033 Oguro, A., Kakeshita, H., Nakamura, K., Yamane, K., Wang, W., and Bechhofer, D.H. (1998). Bacillus
1034    subtilis RNase III cleaves both 5'- and 3'-sites of the small cytoplasmic RNA precursor. J. Biol.
1035    Chem. *273*, 19542–19547.

1036 Oussenko, I.A., Abe, T., Ujiie, H., Muto, A., and Bechhofer, D.H. (2005). Participation of 3′-to-5′
1037    Exoribonucleases in the Turnover of Bacillus subtilis mRNA Participation of 3 J -to-5 J
1038    Exoribonucleases in the Turnover of Bacillus subtilis mRNA. J. Bacteriol. *187*, 2758–2767.

Park, J.-H., Yamaguchi, Y., and Inouye, M. (2011). Bacillus subtilis MazF-bs (EndoA) is a UACAU-
specific mRNA interferase. FEBS Lett. *585*, 2526–2532.
Parker, D.J., Lalanne, J.B., Kimura, S., Johnson, G.E., Waldor, M.K., and Li, G.W. (2020). Growth-
Optimized Aminoacyl-tRNA Synthetase Levels Prevent Maximal tRNA Charging. Cell Syst. *11*,
121-130.e6.
Pellegrini, O., Mathy, N., Gogos, A., Shapiro, L., and Condon, C. (2005). The Bacillus subtilis ydcDE
operon encodes an endoribonuclease of the MazF/PemK family and its inhibitor. Mol. Microbiol.
56, 1139–1148.
Pertzev, A. V, and Nicholson, A.W. (2006). Characterization of RNA sequence determinants and
antideterminants of processing reactivity for a minimal substrate of Escherichia coli ribonuclease
III. Nucleic Acids Res. *34*, 3708–3721.
Ruiz de los Mozos, I., Vergara-Irigaray, M., Segura, V., Villanueva, M., Bitarte, N., Saramago, M.,
Domingues, S., Arraiano, C.M., Fechter, P., Romby, P., et al. (2013). Base Pairing Interaction
between 5′- and 3′-UTRs Controls icaR mRNA Translation in Staphylococcus aureus. PLoS
Genet. *9*.
Salis, H.M. (2011). The ribosome binding site calculator. Methods Enzym. *498*, 19–42.
Salis, H.M., Mirsky, E.A., and Voigt, C.A. (2009). Automated design of synthetic ribosome binding sites
to control protein expression. Nat. Biotechnol. *27*, 946–950.
Le Scornet, A., Jousselin, A., Baumas, K., Kostova, G., Durand, S., Poljak, L., Barriot, R., Coutant, E.,
Pigearias, R., Tejero, G., et al. (2024). Critical factors for precise and efficient RNA cleavage by
RNase Y in Staphylococcus aureus. PLOS Genet. *20*, 1–30.
Shahbabian, K., Jamalli, A., Zig, L., and Putzer, H. (2009). RNase Y, a novel endoribonuclease, initiates
riboswitch turnover in Bacillus subtilis. EMBO J. *28*, 3523–3533.
Stead, M.B., Agrawal, A., Bowden, K.E., Nasir, R., Mohanty, B.K., Meagher, R.B., and Kushner, S.R.
(2012). RNAsnap^{TM}: a rapid, quantitative and inexpensive, method for isolating total RNA from
bacteria. Nucleic Acids Res. *40*, e156.
Subtelny, A.O., Eichhorn, S.W., Chen, G.R., Sive, H., and Bartel, D.P. (2014). Poly(A)-tail profiling
reveals an embryonic switch in translational control. Nature *508*, 66–71.
Taggart, J.C., Lalanne, J.B., and Li, G.W. (2021). Quantitative Control for Stoichiometric Protein
Synthesis. Annu. Rev. Microbiol. *75*, 243–267.
Trinquier, A., Durand, S., Braun, F., and Condon, C. (2020). Regulation of RNA processing and
degradation in bacteria. Biochim. Biophys. Acta - Gene Regul. Mech. *1863*, 194505.
Urtecho, G., Tripp, A.D., Insigne, K.D., Kim, H., and Kosuri, S. (2019). Systematic Dissection of
Sequence Elements Controlling σ70 Promoters Using a Genomically Encoded Multiplexed
Reporter Assay in Escherichia coli. Biochemistry *58*, 1539–1551.
Winkler, W.C., Nahvi, A., Roth, A., Collins, J.A., and Breaker, R.R. (2004). Control of gene expression
by a natural metabolite-responsive ribozyme. Nature *428*, 281–286.
Wu, X., and Bartel, D.P. (2017). kpLogo: positional k-mer analysis reveals hidden specificity in
biological sequences. Nucleic Acids Res. *45*, W534–W538.
Yao, S., and Bechhofer, D.H. (2010). Initiation of decay of Bacillus subtilis rpsO mRNA by
endoribonuclease RNase Y. J. Bacteriol. *192*, 3279–3286.
Yarchuk, O., Jacques, N., Guillerez, J., and Dreyfus, M. (1992). Interdependence of translation,
transcription and mRNA degradation in the lacZ gene. J. Mol. Biol. *226*, 581–596.
Yasbin, R.E., and Young, F.E. (1974). Transduction in Bacillus subtilis by Bacteriophage SPP1 . J. Virol.
14, 1343–1348.
Zhang, K., and Nicholson, A.W. (1997). Regulation of ribonuclease III processing by double-helical
sequence antideterminants. Proc. Natl. Acad. Sci. U. S. A. *94*, 13437–13441.
Zhang, X.Z., and Zhang, Y.H.P. (2011). Simple, fast and high-efficiency transformation system for
directed evolution of cellulase in Bacillus subtilis. Microb. Biotechnol. *4*, 98–105.
1088
1089

1090
1091
Figure 1. Workflow and validation of endoribonuclease cleavage mapping approach
(A) Deletion of exoribonucleases results in stable accumulation of RNA decay intermediates. A schematic
mRNA is shown with endoribonuclease cleavage (red scissors) occurring, with remaining
exoribonucleases indicated in gray. RNAs are shown with 5′-P and 3′-OH characteristic of the most
common endoribonucleases in *B. subtilis*. Newly generated RNA ends can be captured through RNA
ligation allowing mapping of cleavage positions by 3′ and 5′ end sequencing. Adapters used to capture
RNA ends indicated in purple for 3′ end sequencing and yellow for 5′ end sequencing, with matching
color scheme in schematic data below.
(B) Validation of endonuclease cleavage site detection using known positions of endoribonucleolytic
cleavage by RNase III, RNase Y, and Rae1 (DiChiara et al., 2016; Durand et al., 2015; Leroy et al.,
2017). Yellow indicates 5′ end sequencing data and purple indicates 3′ end sequencing data. Red dotted
line represents manually annotated cleavage positions. Plotted are reads per million CDS-mapping reads,
normalized to the average 3′-mapped Rend-seq RPM in this window.
1105

1106
1107
Figure 2. Cleavage within transcripts known to be destabilized by endoribonucleolytic cleavage. 5′
and 3′ end sequencing data are shown in yellow and purple, respectively. Plotted are reads per million
CDS-mapping reads. Manually annotated putative cleavage sites (positions of adjacent 3′ and 5′ read
density) are highlighted with arrows. Insets show highlighted sites at single-nucleotide resolution, with
the Y-axis in each direction rescaled to the maximal value within the inset region. Data are aligned to a
schematic representation of the relevant genomic locus. Annotated promoter and transcriptional
terminators, indicated by bent arrows and lollipops, respectively, were identified using Rend-seq in
SSB1002. Sequence context and RNase Y dependence of highlighted sites shown in Figure S2.
Consistent with a recent report (Korobeinikova et al., 2023), multiple 5′ and 3′ end signals were observed
in the 5′ UTR of *rny*.

1118
1119
Figure 3. *B. subtilis* RNase III cleaves within long-range intramolecular secondary structures
(A) Identification of RNase III cleavage positions in the *priA* mRNA. 5′ and 3′ end sequencing data are
shown in yellow and purple, respectively. Identified positions of cleavage are highlighted in red. Plotted
are reads per million CDS-mapping reads, normalized to the average 3′-mapped Rend-seq RPM in this
window.

(B) Results of systematic identification of RNase III sites. Histogram shows distribution of endonuclease
sensitivities for called peak pairs in 3′/5′ end sequencing of exoribonuclease knockouts (see Figure S3 and
Methods), with black dashed line indicating threshold for calling dependence on RNase III. 68 sites
exceeded this defined threshold. For 43 sites we were unable to calculate a sensitivity score due to an
absence of 5′ end sequencing counts in our knockout. These sites are called as RNase III sensitive and are
counted within the ">4" bin of the histogram.
(C) Predicted secondary structure of the sequence surrounding the identified RNase III sites within the
priA mRNA. Positions of cleavage are indicated with red arrows, with numbers corresponding to those in
(A). Structural distance between these cleavage positions, defined as the defined as the length of the 3′
overhang generated by the cleavages, is annotated in red. Genomic distance, defined as the distance
between the cleavage sites in the primary sequence, is annotated in blue.
(D) Genomic distance (as defined in C) between pairs of unambiguous RNase III cleavage positions
identified within 1 kb of one another and predicted to fall on opposite sides of an RNA stem with a 2 bp
spacing between cleavage positions (32 of 68 positions).
(E) Structural distance (as defined in C) generated by RNase III cleavage at both positions within the
pairs of cleavage positions plotted in (D).
(F) Predicted secondary structures of the sequence surrounding the identified RNase III sites within the
mRNAs encoding the *yqeU*, *atpA*, and *asnB* genes. Insets are plotted as in (A).
(G) Consensus sequence-structure motif of RNase III targets in *B. subtilis*. The predicted structures for
each pair of RNase III-sensitive cleavage sites that generate a 2-nt 3′ overhang were aligned to the
positions of cleavage, "1" and "2". The position-wise frequency of each type of base-pairing interaction is
shown. Note that the base pairing frequencies sum to the fraction of sites paired at each position. The
most frequent base pair at each position is shown in the schematic on the left, with lower-case characters
used to designate positions for which the most frequent base pair occurs <20% of the time. When two
base pair frequencies are within 0.02 of one another, both are separated by a slash (e.g., A-U/G-C). The
gray regions labeled "PB", "MB", and "DB" correspond to the proximal, middle, and distal boxes (Gan et
al., 2006; Zhang and Nicholson, 1997).
(H) Comparison of base pairing frequences between *B. subtilis* and *E. coli* (Altuvia et al., 2018). The
consensus *E. coli* sequence-structure motif is shown on the left with top base pairs at each position which
are shared between *B. subtilis* and *E. coli* highlighted in green. Red box highlights the decrease in C-G or
G-C pairing in *B. subtilis* relative to *E. coli*.
1156

1157
Figure 4. Revision of EndoA cleavage specificity and evidence for new ribonuclease activity in *B.*
subtilis
(A, B) 3′ (A) and 5′ (B) mapped Rend-seq signal across all UACAU motifs in the genome, separated by
downstream nucleotide. 5′ end sequencing data derived from an RNase J1 deletion (CCB434) and 3′ end
sequencing derived from a 4-exo knockout (CCB396). Rend-seq data at each site is normalized to first 8
(for 3′-mapped) or last 8 (for 5′-mapped) positions within the window and a position-wise mean and
standard deviation (shaded interval) are calculated with 90% winsorization. Motif instances with fewer

than 1 read per position or fewer than 10 reads within the normalization window are not considered.
Following this filtering, the number of considered sites ranges from 264 to 435 (A) or 292 to 455 (B) per
motif. The black vertical line indicates the position of cleavage by EndoA.
(C, D) 5′-mapped Rend-seq signal in an RNase J1 depletion (CCB390) (C) or and RNase J1 depletion
with deletion of *ndoA* (BT231) (D). Analysis performed identically to (A, B). Following filtering, the
number of considered sites ranges from 191 to 288 (C) or 86 to 117 (D) per motif.
(E) Rend-seq (top) and end sequencing (bottom) data at a representative EndoA cleavage site located in
the gene *pgk* for a 4-exo knockout (CCB396), RNase J1 depletion (CCB390), and RNase J1 depletion
with an *ndoA* knockout (BJT231). 5′-mapped data shown in yellow (5′ end sequencing) or orange (Rend-
seq) and 3′-mapped data shown in purple (3′ end sequencing) or blue (Rend-seq). Red dotted line
indicates position of cleavage. Plotted are reads per million CDS-mapping reads. 5′/3′-end sequencing
data are normalized to the average 3′-mapped Rend-seq signal in this window.
(F) Rend-seq (top) and end sequencing (bottom) data at the position of cleavage by the *glmS* ribozyme. 5′-
mapped data shown in yellow (5′ end sequencing) or orange (Rend-seq) and 3′-mapped data shown in
purple (3′ end sequencing) or blue (Rend-seq). Strains included are identical to panel E. Red dotted line
indicates position of cleavage. Data are plotted as in (E).
(G) 5′-end sequencing data at the position of cleavage by the *glmS* ribozyme for knockouts of additional
RNA decay-associated proteins. All knockouts are coupled to either a depletion or deletion of RNase J1.
Knocked out genes include *rnjA* (RNase J1, strain CCB434), *rny* (RNase Y, strain CCB760), *ylbF* (YlbF,
strain BJT074), *rppH* (RppH, strain BJT129), and *rnc* (RNase III, strain BG879). Data are plotted as in
(E), with the exception of *rppH*, which is normalized to an *rnjA* knockout alone, rather than the same
genetic background, due to a lack of corresponding Rend-seq data.
(H) Rend-seq data at the position of cleavage by the *glmS* ribozyme in a 4-exo knockout (CCB396),
RNase J1 knockout (CCB434), and RNase J1 depletion with a *rnjB* knockout (GLB186). Red dotted line
indicates position of cleavage. 3′-mapped data are shown in blue and 5′-mapped data are shown in orange.
Plotted are reads per million CDS-mapping reads.
(I, J) 5′ end sequencing (I) and Rend-seq 3′-mapped (J) signal across all UACAUA EndoA cleavage
motifs, separated by downstream sequence. 5′ end sequencing data derived from an RNase J1 depletion
with *rny* knockout (CCB760) and 3′ end sequencing derived from a 4-exo knockout (CCB396). The
signal in a 20 nt window around each cleavage site was normalized to its maximal value and a position-
wise average across all normalized windows was calculated with 90% winsorization. UACAUA instances
for which the local density in corresponding Rend-seq data is less than 1 read per position were discarded.
Following this filtering, the number of considered sites ranges from 10 to 54 per row (I) or 10 to 62 per
row (J). The red vertical line indicates the position of cleavage by EndoA.
1199
1200

1201
1202
Figure 5. Sequence and structural features of putative RNase Y substrates
(A) Systematic identification of putative RNase Y cleavage sites. Histogram shows distribution of
endonuclease sensitivities for called peak pairs in 3′/5′ end sequencing of exoribonuclease knockouts (see
Figure S3 and Methods). Solid red line shows results of fitting the sum of two Gaussian distributions to
this underlying log-transformed data, with the dashed lines showing the two constituent distributions
produced by this fitting. The dashed vertical line indicates the lowest endonuclease sensitivity value
(10.56) for which 95% of the Gaussian sum is derived from the right constituent distribution, a value
above which sites are considered dependent on RNase Y. 596 of 1301 considered putative cleavage
signatures exceed this threshold. For 85 sites we were unable to calculate a sensitivity score due to an

absence of 5′ end sequencing counts in the knockout. These sites are called as RNase Y sensitive and are
counted within the ">4" bin of the histogram.
(B) Local sequence context around positions of RNase Y cleavage (N=596). Nucleotide frequency (top),
information content (middle) and k-mer (bottom) logos in a 30 nucleotide window around called RNase Y
sites. In k-mer logo, the most enriched k-mers are shown above the sequence and the most depleted
below. Significant positions (Bonferroni corrected P<0.01, one-sided binomial test), are highlighted in
red.
(C) Local GC content around positions of RNase Y cleavage (N=484). Sites near transcript boundaries
are not considered (see Methods). Dotted gray line indicates average %GC across *B. subtilis* genome.
Vertical red line indicates position of cleavage. Yellow region indicates positions shown in (B).
(D) Predicted folding near RNase Y cleavage positions (N=484). Sites near transcript boundaries are not
considered (see Methods). Blue line shows position-wise average minimal free energy (MFE) predicted
by RNAfold for 40-mers tiling cleavage sites. Gray band indicates per-position interquartile range of
curves resulting from folding randomly sampled GA-dinucleotide centered windows within coding
regions, excluding putative cleavage positions (see Methods). Red vertical line indicates putative position
of cleavage.
(E) Systematic identification of YlbF-dependent endoribonuclease cleavage sites. Histogram shows
distribution of endonuclease sensitivities for called peak pairs in 3′/5′ end sequencing of exoribonuclease
knockouts (see Figure S3 and Methods). The dashed vertical line indicates the endonuclease sensitivity
threshold used to call RNase Y sites. Red arrows indicate endonuclease sensitivity values of canonical
processing sites within the *cggR*, *glnR*, and *atpI* transcripts. For 9 sites we were unable to calculate a
sensitivity score due to an absence of 5′ end sequencing counts in the knockout. These sites are called as
YlbF sensitive and are counted within the ">4" bin of the histogram.
(F-K) End mapping data showing evidence for inefficient but detectable endoribonucleolytic cleavage at
canonical RNase Y processing sites within the transcripts encoding *glnR* (F, G), *cggR* (H, I), and *atpI* (J,
K) in the absence of YlbF. (F, H, and J) show 5′/3′-end sequencing and (G, I, and K) show Rend-seq. 5′-
mapped data shown in yellow (5′ end sequencing) or orange (Rend-seq) and 3′-mapped data shown in
purple (3′ end sequencing) or blue (Rend-seq). Red dotted line indicates positions of endoribonucleolytic
cleavage. Plotted are reads per million CDS-mapping reads. 5′/3′-end sequencing data are normalized to
the average 3′-mapped Rend-seq signal in this window.
1242

1243
1244
Figure 6. Mutational scanning of the *cggR-gapA* operon cleavage site
(A) Schematic design of MPRA construct. Cleavage region to mutate is indicated in dark gray with
yellow used to indicate mutated region. Variant barcode indicated in green. Translated regions are
indicated with a thick border. Promoters and transcriptional terminators are indicated by bent arrows and
lollipops, respectively.
(B) Workflow for measurement of mRNA processing. Protocol begins from *B. subtilis* culture containing
a pool of genetically encoded cleavage site variants, and splits into gDNA and RNA barcode

quantification protocols. The relative barcoded RNA abundance is calculated as the ratio of RNA-derived
to gDNA-derived reads.
(C) Schematic of the $cggR_{148}$ construct inserted into the $aprE$ MPRA transcript. $cggR$-derived sequence is
colored gray, with the mutated positions highlighted in yellow. Translated regions are indicated with a
thick border, and the variant barcode is indicated in green.
(D) Impact of all single-nucleotide mutations on the accumulation of barcoded full-length RNA. Boxplots
show variation between barcodes of identical variant sequence. Number of barcodes captured for each
mutation is indicated above plot. Whiskers indicate 5^{th} and 95^{th} percentile. Gray shaded region indicates
interquartile range for variants of wild-type sequence. Red line indicates position of cleavage. Blue
arrows highlight mutations which close the bulge in the $cggR$ downstream hairpin. Four variants (no more
than one per mutation) have a value of zero are thus not visualized.
(E) Impact of all single-nucleotide mutations on the accumulation of barcoded cleavage product. Boxplots
show variation between barcodes of identical variant sequence. Number of barcodes captured for each
mutation (N) is indicated by the bar height (top). Whiskers indicate 5^{th} and 95^{th} percentile. Gray shaded
region indicates interquartile range for variants of wild-type sequence. Red line indicates position of
cleavage and brown bars indicate regions predicted to pair in the formation of a downstream stem-loop
structure (as illustrated in F). Red arrows indicate mutations which are predicted to generate G•U wobble
pairing, and the blue arrow highlights a mutation which corrects the bulge in the downstream stem-loop
structure (B). Four variants (no more than one per mutation) have a value of zero are thus not visualized.
(F) Predicted stem-loop structure at 5′ end of processed RNA. Inset shows impact of double mutants on
accumulation of processed RNA. Double mutant averages calculated over any variants with the two
indicated mutations. Any variants which contain additional mutations that on their own result in greater
than 10% change in processed RNA accumulation are discarded.
(G) Relationship between predicted strength of downstream secondary structure and accumulation of
barcoded cleavage product. The experiment each data point is derived from is denoted with a triangle
(experiment 2b, Table S3), circle (4a), or square (6). The vertical dashed line indicates the ΔG of the
unmutated sequence. Five variants fall outside of the bounds of this plot.
(H) Relationship between predicted strength of downstream secondary structure and accumulation of
barcoded full-length RNA. All data derived from experiment 2a (Table S7). The vertical dashed line
indicates the ΔG of the unmutated sequence. Five variants fall outside of the bounds of this plot.
1282

Figure 7. Determinants of endoribonuclease cleavage identified in this work

(A) Summary of the identified cleavage determinants of the three endoribonucleases profiled in this work. RNase III appears to cleave within double-stranded RNA sequences, including many duplexes formed through long range (>100 nt) intramolecular interactions, consistently leaving a 2 nucleotide 3′ overhang (left, "III"). EndoA appears to cleave specifically at a UACAUA primary sequence motif (right, "EndoA"). RNase Y cleavage appears to be targeted towards AU-rich regions of low secondary structure, appearing to prefer cleavage within a G/A dinucleotide (center, "Y"). Interrogation of the cleavage sites within *cggR* and *glnR* suggest that a -1G is dispensable for RNase Y cleavage, and for a given site the sequence and structural elements that drive cleavage (orange) may be distributed over many tens of nucleotides. Further, for some mRNA processing sites such as those in *cggR* or *gapA*, a downstream stem-loop structure (blue) appears to primarily drive stabilization of the cleaved isoform.

Figure S1. Validation of approach through detection of known positions of endonuclease activity. 5′ and 3′ end sequencing data at known positions of cleavage by RNases Y and III. Datasets considered are a 4-exo knockout (CCB396), *rnjA* knockout (CCB434), and either deletion or depletion of RNase J1 with a knockout of RNase Y (CCB760) or III (BG879). 5′-mapped data shown in yellow and 3′-mapped data shown in purple. Plotted are reads per million CDS-mapping reads, normalized to the average 3′-mapped Rend-seq RPM in this window. A manually annotated position of cleavage based on sequencing data is shown with a red dotted line, and the published position is marked in gray of cleavage if this position disagrees with our annotation. In transcripts with many nearby peak pairs such as *hbs*, additional cleavage positions may be visible beyond that which is highlighted.

1308
Figure S2. **Sequence context and RNase Y/III dependence of cleavage positions within mRNAs**
known to be destabilized by endoribonucleases. Cleavage positions match order of those shown in
Figure 2. Datasets considered are a 4-exo knockout (CCB396), *rnjA* knockout (CCB434), rnjA rnc double
knockout in an SPβ and Skin-cured background (BG879), and depletion of RNase J1 with a knockout of
rny (CCB760). 5′ end sequencing data shown in yellow and 3′ end sequencing data shown in purple.
Plotted are reads per million CDS-mapping reads, normalized to the average 3′-mapped Rend-seq RPM in
this window. A manually annotated position of cleavage based on sequencing data is shown with a red
dotted line. Data from the *rnc*- or *rny*-deficient strain are not shown for sites within the transcripts
encoding these genes.

1318
Figure S3. Schematic representation of strategy for calling putative cleavage positions and
assigning them to the activity of endoribonucleases
(A) Strategy for calling paired 3′ and 5′ peaks in end sequencing data, with sites exceeding a peak-to-
background ratio of 7.5 in the 3′/5′ at adjacent positions called as putative cleavage positions.
(B) Strategy for attributing sites to specific endonucleases. Peak heights in 5′ end sequencing are
normalized to local density in a corresponding 5′-mapped Rend-seq dataset. A ratio of normalized peak
heights between exoribonuclease strains with and without an endoribonuclease knockout (termed
"endonuclease sensitivity") is calculated as a metric for dependence on this endoribonuclease. Note that
prior to illustrated calculation, 5′ end sequencing data (h_{WT_5end} and $_{hKO\_5end}$) are normalized by the sum of
5′ end sequencing reads mapped to putative cleavage positions, and the Rend-seq background (bg_{Rend}) is
normalized to the total CDS-mapped reads to account for differences in sequencing depth between
samples.

(C) Results of systematic identification of RNase III sites with a relaxed 5′ end sequencing depth
threshold (as used in Figure 3G, H). Histogram shows distribution of endonuclease sensitivities for called
peak pairs in 3′/5′ end sequencing of exoribonuclease knockouts (see Figure S3A and Methods), with
black dashed line indicating threshold for calling dependence on RNase III. 162 sites exceeded this
defined threshold. For 122 sites we were unable to calculate a sensitivity score due to an absence of 5′ end
sequencing counts in our knockout. These sites are called as RNase III sensitive and are counted within
the ">4" bin of the histogram.
(D) Genomic distance between pairs of unambiguous RNase III cleavage positions identified within 1 kb
of one another and predicted to fall on opposite sides of an RNA stem, using both restrictive and relaxed
5′ end sequencing depth thresholds.
(E) Results of systematic identification of RNase Y sites with a relaxed 5′ end sequencing depth
threshold. Histogram shows distribution of RNase Y sensitivities, plotted as described in Figure 5A. 1351
sites exceeded this defined threshold. For 408 sites we were unable to calculate a sensitivity score due to
an absence of 5′ end sequencing counts in our knockout. These sites are called as RNase Y sensitive and
are counted within the ">4" bin of the histogram.
1346

1347
Figure S4. EndoA and YlbF have a limited influence on wild-type *B. subtilis* mRNA decay
(A) UACAUA-specific EndoA cleavage is not detectable in wild-type *B. subtilis.* 3′ (left) and 5′ (right)
mapped Rend-seq signal across all UACAU motifs in the genome, separated by downstream nucleotide.
Data are derived from wild-type W168. Rend-seq data at each site is normalized to first 8 (for 3′-mapped)
or last 8 (for 5′-mapped) positions within the window and a position-wise mean and standard deviation
are calculated with 90% winsorization. Motif instances with fewer than 1 read per position or fewer than
10 reads within the normalization window are not considered. Following this filtering, the number of
considered sites ranges from 193 to 313 (for 3′-mapped) and 203 to 311 (for 5′-mapped) per motif. The
dashed vertical line indicates the position of cleavage by EndoA.
(B, C) Representative EndoA sites showing presence (B) or absence (C) of subsequent trimming after the
first two nucleotides. Yellow indicates 5′ end sequencing data and purple indicates 3′ end sequencing
data. Plotted are reads per million CDS-mapping reads, normalized to the average 3′-mapped Rend-seq
RPM in this window. Red dotted line represents manually annotated EndoA cleavage positions. Note that
cleavage product accumulation is dependent on EndoA (bottom two panels).
(D) mRNA abundances for all annotated *B. subtilis* genes in a wild-type and *ylbF* W168 background.
Lowly expressed genes (<128 total reads) are excluded. RPKM calculated as reads per thousand bases of
the gene per million total reads that map to *B. subtilis* coding regions. Inset shows differential expression
distribution (Δ*ylbF*/WT RPKM) for genes with genes with (red) or without (black) a YlbF-dependent
putative cleavage site.
1367

1368
1369
Figure S5. Variability and sequence coverage within our MPRA datasets
(A) Cumulative distribution of the number of unique barcodes associated with each possible sequence
containing a single substitution relative to the wild-type *cggR* or *glnR* operon sequence. Undetected
sequences are included with a pseudocount of 0.1 prior to $\log_2$ transformation.
(B) Cumulative distribution of the number of unique barcodes associated with each possible sequence
containing two substitutions relative to the wild-type *cggR* or *glnR* operon sequence. Undetected
sequences are included with a pseudocount of 0.1 prior to $\log_2$ transformation.
(C) Variation in MRPA readout as a function of genomic DNA sequencing depth. The normalized MPRA
RNA abundance for each variant with a wild-type sequence was calculated as illustrated in Figure 6A.
These values were grouped by the number of unique reads mapping to that variant in the genomic DNA
barcode sequencing with a bin width of 50 reads. The coefficient of variation within each group is shown.
The gray vertical dashed line indicates the 50 read cutoff used for experiments 4b, 8a, 8b, and 9, red
vertical dashed line indicates the 100 read cutoff used in analysis of all remaining experiments. See Table
S7 for more information.
(D) The number of variants within each bin for each dataset show in (A). Dashed lines show thresholds as
described in (A).
1386

1387
Figure S6. Features driving RNase Y cleavage within *cggR* are consistent across multiple MPRA contexts

(A) Impact of all single-nucleotide mutations on the accumulation of barcoded cleaved *aprE-cggR*148 RNA in the context of RNase J1 depletion. Boxplots show variation between barcodes of identical variant sequence. Number of barcodes captured for each mutation is indicated above plot. Whiskers indicate 5th and 95th percentile. Gray shaded region indicates interquartile range for variants of wild-type sequence. Red line indicates position of cleavage. Ten variants have a value of zero (no more than one per mutation) and are thus not visualized.

(B) Impact of all single-nucleotide mutations on the accumulation of barcoded cleavage product when the *cggR*148 construct is appended to a new scaffold RNA, *tetM*. Plotted as in (A). Two variants (no more than one per mutation) have a value of zero are thus not visualized.

(C) Impact of all single-nucleotide mutations in the 30 nucleotides upstream of the cggR148 cleavage site on the accumulation of barcoded cleavage product when a stop codon is inserted upstream of the inserted sequence. *tetM* scaffold was used in this experiment, as in (C). Plotted as in (A).

1402
Figure S7. Determinants of *glnR-glnA* mRNA processing by RNase Y.
(A) Schematic of the *glnR*₁₇₈ construct inserted into the *aprE* MPRA transcript. *glnR*-derived sequence is
colored dark gray, with the mutated positions highlighted in yellow. Translated regions are indicated with
a thick border, and the variant barcode is indicated in green.

(B) Predicted stem-loop structure at 5′ end of processed RNA. Brown bar corresponds to regions
highlighted in panels C, E, and F.
(C) Impact of all single-nucleotide mutations on the accumulation of barcoded cleaved RNA. Boxplots
show variation between barcodes of identical variant sequence. Number of barcodes captured for each
mutation is indicated above plot. Whiskers indicate 5th and 95th percentile. Gray shaded region indicates
interquartile range for variants of wild-type sequence. Red line indicates position of cleavage. Brown bar
corresponds to positions predicted to form the stem of the structure showed in (B). Nine variants (no more
than two per mutation) have a value of zero are thus not visualized.
(D) Relationship between predicted strength of downstream secondary structure and accumulation of
barcoded cleavage product. All data are derived from experiment 8a (Table S7). The vertical dashed line
indicates the ΔG of the unmutated sequence. 17 variants fall outside of the bounds of this plot.
(E) Impact of all single-nucleotide mutations on the accumulation of barcoded full-length RNA, plotted as
described in (C). 30 variants (no more than two per mutation) have a value of zero are thus not visualized.
(F) Impact of all single-nucleotide mutations on the accumulation of barcoded cleaved RNA in the
context of RNase J1 depletion, plotted as described in (C). Three variants (no more than one per
mutation) have a value of zero are thus not visualized.
1423

1424 **Table S1. List of strains used in this study**
1425
1426 **Table S2. List of oligonucleotide sequences used in this study**
1427
1428 **Table S3. List of identified RNase III cleavage sites**
1429
1430 **Table S4. List of identified RNase Y cleavage sites**
1431
1432 **Table S5. List of putative RNase III cleavage sites with relaxed 5' end sequencing depth threshold**
1433
1434 **Table S6. List of putative RNase Y cleavage sites with relaxed 5' end sequencing depth threshold**
1435
1436 **Table S7. Summary of MPRA experiments conducted in this study**