Linking B-factor and temperature-induced conformational transition

Fernando de Sá Ribeiro[1] and Luís Maurício T. R. Lima[1,2*]

[1] Pharmaceutical Biotechnology Laboratory (pbiotech), Faculty of Pharmacy, Federal University of Rio de Janeiro, Rio de Janeiro, RJ, 21941-902, Brazil.
[2] National Institute for Metrology, Quality and Technology (INMETRO), INMETRO, Duque de Caxias, RJ, 25250-020, Brazil.

Running title: Thermal-dependence of B-factor

*To whom correspondence should be addressed

Authors Address and Contact
- Luis Mauricio T. R. Lima – Pharmaceutical Biotechnology Laboratory (pbiotech), Faculty of Pharmacy, Federal University of Rio de Janeiro – UFRJ, CCS, Bss24, Ilha do Fundão, 21941-590, Rio de Janeiro, RJ, Brazil. Phone/Fax: (+55-21) 3938-6639 – Mauricio@pharma.ufrj.br. Social Media: @pbiotech

AUTHOR LIST
Fernando de Sá Ribeiro – FernandoRibeiroBiomed@gmail.com – ORCID 0000-0002-8521-6745
Luís Maurício T. R. Lima – Mauricio@pharma.ufrj.br – ORCID 0000-0001-6020-0504

ABSTRACT

The crystallographic B-factor, also called temperature factor or Debye-Waller factor, has long been used as a surrogate for local protein flexibility. However, the use of the B-factor as a probe for protein motion still urge reproducibility and further validation against chemical and physical variables. Here we report the investigation of the thermal dependence of the crystallographic B-factor and its correlation with protein conformational changes. We solved the B-factor reproducibility issue at high resolution over a broad temperature range (100 K to 325 K) by using hydrocarbon grease during data collection. We found that the crystallographic protein conformation varies linearly as a function of temperature. Further, the demonstrated that the thermal dependence of B-factor as a function of temperature were similar for all atoms (Cα, N-amide and side chains, demonstrating the lack of correlation between temperature-dependent conformational change and the B-factor or the B-factor thermal constant. Our results reveal a dynamic correlation of B-factor with global rigid body motion.

Keywords: B-factor; crystallography; temperature; ion mobility spectrometry; lysozyme

Abbreviations: HEWL, hen egg white lysozyme; CD, circular dichroism; ESI, Electrospray ionization; MS, mass spectrometry; IMS, ion mobility spectrometry.
.

Introduction

Protein conformational polymorphism and dynamics are part of the proteostasis network involving synthesis and folding, conformational maintenance and degradation [1]. While high resolution structure of macromolecules provides conformational snapshots, molecular dynamics are also required to infer mechanisms required for the understanding of biological systems.

There is increasing prevalence of computational modeling approaches in the study of macromolecular structure and dynamics [2,3]. However, web-bench techniques are still required [4]. Structure determination of macromolecules can be performed by a large set of state-of-the-art methods, such as in solid phase from crystals (nano to micro; powder, poly or single) by diffraction (electron, x-ray, neutron, XFEL), or in solution by NMR and vitreous by cryo-EM. Time-resolved techniques can also infer dynamics directly such as NMR and ultrafast serial femtosecond crystallography (SFX) techniques [5,6].

Conformational polymorphism and dynamics may also be inferred by solution NMR techniques, or from ensemble structure solved at varying conditions such as chemical, including pH, ligands, cosolutes, and hydration changes [7,8] or physical such as temperature and pressure [9–11].

Single-crystal X-ray diffraction (SCXRD) has been used for over 60 years in solving protein structure [12], and holds an outstanding position in structural biology for being a highly reproducible technique [13–15]. While the high-resolution crystal structure reflects the average coordinate position of the proteins in the crystal, the crystallographic **B-factor** would allow inferring conformational flexibility from this ensemble. However, It has been argued that the B-factor is highly inaccurate, requiring experimental and analytical improvements [16].

The B-factor – often called temperature factor [17–20] - relates to the mean-square atomic displacement (X^2) by the Debye-Waller function

[Eq. 1] $B=8\pi^2 X^2$

where X^2 embodies additive crystal contributions from atomic fluctuation (X^2_A), conformational motions (X^2_C), rigid body vibration (X^2_{RB}) and crystal lattice disorders (X^2_{LT}) [21]. The crystallographic B-factor is also susceptible to variables from crystal (such as hydration, cosolutes, cryoprotection), XRD setup (such as beam, detector), gas stream for temperature control and diffuse scattering [14]. Diffraction shows contributions from both ordered lattice and diffuse scattering [22–25], while diffuse scattering can show contributions from varying sources including rigid-body motion [26] and background scattering [14]. For these reasons, direct correlations between B-factor with protein dynamics and consequent propensity for conformational changes are still controverse [27].

Attempts to solve the uncertainty problem in B-factor backs from decades ago. The broad distribution of B-factor can be empirically scaled down by its average B-factor (B_{avg}) resulting in normalized B-factor (B_{norm}) [17] in the form of

[Eq. 2] $B_{norm} = B_{obs} / B_{avg}$

which differ from the *Z-score* as proposed elsewhere [27,28].

We have recently shown that B-factor reproducibility can be achieved by the empirical scaling [15]. In this procedure, we used 20 crystals of lysozyme collected in 2 different XRD setups, as well as reanalyzed data from human insulin from four dissimilar biotechnology processes collected at four dissimilar XRD setups [13]. The B_{norm} for each amino acid showed low standard deviation and were reproducible. We further analyzed data from RCSB, and found that local variations in B_{norm} was related to space group. The

correlation between B-factor and protein dynamic remained to be solved, requiring further investigation preferably by using variables to promoting change in conformation and B-factor, separating contributions from static (temperature independent) and dynamic (temperature-dependent) disorder [21,29].

The temperature influence over protein conformation has long been investigated by crystallography for ribonuclease from 98 to 325 K [29], high-pressure (200 MPa, 2 kbar) cryocooled thaumatin from 80 to 160 K [30], thaumatin from 25 to 300 K [10], myoglobin from 80 to 240 K [31], and hen egg white lysozyme (HEWL) solved at cryogenic and room temperatures and shown to vary in conformation and B-factor [32,33] resulting in expansion of about 0.9 % per 100 K.

Analysis of tetragonal lysozyme data currently deposited in the RCSB evidences a gap in the temperature dimension resulting in two subgroups of structures (**Fig S1**), one in the cryogenic range (about 100 K) and another close to room temperature (at about 300 K, with highest temperature at 308 K, i.e., 35 $^{\circ}$C). We also observe a large dispersion in cell parameters (**Fig. S1A** and **Fig S1B**) and the lack of direct correlation between them (**Fig. S1D**). There is no apparent correlation between crystallization pH and data collection temperature (**Fig. S1E**), or with the average B-factor for the final structural model (**Fig. S1F**), although a clear correlation is found between the Wilson and average structural B-factor (**Fig. S1G**). From these data it becomes evident the absence of precise information on the effects of data collection temperature on crystallographic and structural parameters, requiring a systematic approach to use broad-range temperature as a physical variable in protein crystallography.

A major limitation in using physico-chemical variables in the study of B-factor is the lack of experimental repeatability and reproducibility in raw (non-normalized) B-factor, mostly due to dehydration. In small-molecule crystallography, non-dehydration agents are routinely used, such as oils and greases. In protein crystallography, non-dehydration techniques involves capillaries [34], polymer tubes (such as CrystalCap®, Hampton Research), uncured glue / epoxy cement [29,35], silicon nitride wafers (Silson®) [36], lipid cubic phase (LCP) [37], oil [10,38,39], agarose/gelatin [40], poly-acrylamide [41] mineral oil-based grease [42] shortening [43] and animal fat [44,45].

In this study we aimed to solve the experimental reproducibility of B-factor and elucidate the temperature dependence of B-factor. We solved the structure of tetragonal lysozyme over a broad temperature range from single crystals immersed into hydrocarbon grease, and discuss the potential correlation between conformational changes and B-factor.

Results

Reproducible crystal diffraction at high resolution from 100 K to 325 K

While glycerol is used as cryoprotection, the present study required further stabilization of the crystal for repeatability of the B-factor. For this purpose, we embed the crystal in hydrocarbon grease. We were able to diffract at 1.5 Å resolution from 100 K to 325 K (52 °C) in 25 K intervals, in triplicate for each temperature. Attempts to diffract at higher temperature resulted in progressive loss in order as indicated by loss of diffraction data (not shown). All dataset were obtained under similar data collection strategy, automatically chosen by the controller program, and the crystal structures were solved using the same workflow, avoiding individual interpretations and interventions.

We observed a positive dependence of the unit cell lengths on data collection temperature, for both a=b (**Fig. S2A**) and c (**Fig. S2B**), and a positive correlation between these dimensions (**Fig. S2C**). The increase in unit cell lengths corresponded to a volume expansion of the cell (**Fig. S3A**), as well as in the increase in volume of the protein in the asymmetric unit (**Fig. S3B**), with no significant changes in the protein accessible surface area (**Fig. S3C**). The volumetric changes in cell unit and protein showed positive correlation (**Fig. S3D**), which might be related to contribution from the protein to the expansion of the cell. A positive correlation between ASA and the unit cell volume (**Fig. S3E**) and with the protein volume (**Fig. S3F**) was also found, despite minor correlation between ASA and temperature (**Fig. S3C**).

Loss of diffraction power due to protein unfolding over 325 K

We measured the thermal denaturation of lysozyme in sodium acetate buffer pH 4.6 by circular dichroism (CD). A progressive decrease in ellipticity was observed by increasing temperature from 15 °C to 90 °C in 5 °C steps (**Fig. S4A**). The thermal denaturation curves for each wavelength demonstrate a decrease in the steepness of the transitions as a function of the wavelength, converging to about 360 K for complete thermal denaturation transition (**Fig. S4B**). We adjusted the thermal denaturation curves for each wavelength with a sigmoidal function, obtaining both the half-transition temperature ($T_{50\%}$) and the slope of the transition (**Fig. S4C**). The $T_{50\%}$ for thermal denaturation of lysozyme was found to be about 310 K in the 212 nm to 225 nm wavelength range. These data indicate that the temperature range of 340 K to 360 K are compatible with denaturation of lysozyme in solution in buffer conditions close to the crystallization milieu.

We further probed the overall conformational distribution of lysozyme by ion mobility spectrometry coupled to mass spectrometry by electrospray ionization (ESI-IMS-MS). We evaluated the overall ion mobility distribution for lysozyme at varying charged state as a function of source temperature, from 50 °C to 250 °C in 50 °C intervals (**Fig. S5**). Attempts for measurement with source temperature below 50 °C resulted in spectra with insufficient signal. We have found a progressive drift and broadening in the mobility as a function of increasing temperature, indicative of changes in the protein conformational landscape. Collectively, these spectroscopic data (CD and IMS) indicate the thermal instability of lysozyme in the proximity of 360 K, which may determine a loss in lattice order and consequently diffraction power of lysozyme crystals at higher temperatures.

Temperature induces local conformational changes in crystal phase

A pairwise global structural alignment of the lysozyme structures demonstrates an average root-mean square deviation (RMSD) up to 0.3 Å for Cα (**Fig. S6**). Superposition of the protein structures showed global folding similarity in lysozyme conformation, with local changes for both backbone (**Fig. 1A**) and side chains (**Fig. 1B**). Overall deviation in side chains (**Fig. 1C**), Cα (**Fig. 1D**) and amide nitrogen

(N-amide; **Fig. 1E**) showed a positive linear correlation with temperature. No overall change in secondary structure content were observed in crystal structures as a function of data collection temperature in the range of 100 K to 325 K (**Fig. 1F**).

A general overview of conformational changes as a function of data collection temperature could be further inferred from the change in $C\alpha$ RMSD distribution over the polypeptide chain at varying temperatures using a reference structure at 100 K (**Fig. 2**). The positional drift induced by temperature was analyzed by first-order linear regression (**Fig. 2B; Fig. S7**), and by non-linear regression with a single exponential (**Fig. 2D; Fig. S8**). The thermal constant for conformational changes **k** was distributed as a function of polypeptide chain for first-order linear regression (**Fig. 2C**) and for non-linear exponential regression (**Fig. 2E**). The correlation coefficient for the regressions were higher at regions with higher conformation-temperature constants, and apparently higher for first-order polynomial function compared to exponential function (**Fig. S9**), indicating better adequacy of the linear function for this analysis. The thermal expansion of the lysozyme was discontinuous along the polypeptide chain (**Fig. 2C**), indicating that it occurs preferably in certain regions.

Temperature affects average B-factor uniformly

We further analyzed the changes in B-factor as a function of temperature. The **average** B-factor from data as estimated from the Wilson plot (**Fig. 3A**), and from the final refined structure side-chains (**Fig. 3B**), $C\alpha$ (**Fig. 3C**) or N-amide (**Fig. 3D**) showed similar exponential positive correlation with temperature, increasing about 2.5 times from 100 K to 325 K. The thermal dependence of average observed B-factors (^{avg}B) was adjusted with a single exponential function according to **[Eq. 3]**

[Eq. 3] $B_{obs} = B_0 * e^{(k*T)}$

where B_{obs} is the refined B-factor, B_0 is the B-factor interpolated to zero Kelvin from regression, **k** is the thermal constant, and T is the data collection temperature (in Kelvin), resulting in close average thermal factors of about $^{avg}k = 0.0045$ K^{-1} and in $^{avg}B_0$ of about 5 $Å^2$

Temperature B-factor is a global constant

A detailed picture of B-factor dependence on data collection temperature could be inferred from the change in B-factor **distribution over the polypeptide chain** at varying temperatures for side-chains (**Fig. 4A; Fig. S10**), $C\alpha$ (**Fig. 5A; Fig. S11**) and N-amide (**Fig. 6A; Fig. S12**). These data revealed similar B-factor pattern of *crescendo* and *decrescendo* along the polypeptide chain for all temperature tested.

We further analyzed the changes in B-factor as a function of temperature for each amino acid residue, for their side-chain (**Fig. 4B**), $C\alpha$ (**Fig. 5B**) and N-amide (**Fig. 6B**), and adjusted these temperature curves by a single exponential non-linear regression according to [Eq. 3], which described adequately the thermal-dependent changes in $^{Wilson}B$ and average ^{SC}B, $^{C\alpha}B$ and $^{N-amide}B$ (**Fig. 8**). The distributions of the calculated B_0 along the polypeptide chains (**Fig. 4C**, **Fig. 5C** and **Fig. 6C**) showed similar patterns for the raw B-factor distributions (**Fig. 4A**, **Fig. 5A** and **Fig. 6A**). In opposite, the estimated thermal factor $^{C\alpha B}k$ was found to be roughly constant at about 0.005 K^{-1} along the polypeptide chain for side-chains (**Fig. 4C**), $C\alpha$ (**Fig. 5C**) and N-amide (**Fig. 6C**), in close agreement with those found for ^{avg}k (**Fig. 3**). These data suggests that the B-factor increases uniformly for the whole system (side

chain, Cα and N-amide), and thus confirming it as a **temperature-dependent factor**. However, the thermal factor ^{B}**k** shows no association with the observed B-factor (**Fig. 4C, Fig. 5C, Fig. 6C**), indicating that the absolute B-factor does not predict its temperature dependence. Moreover, the similarity of the thermal factor ^{B}**k** between all amino acid residues (both side-chains, Cα and N-amide) indicates that this parameter may not correlate with the observed temperature-induced protein conformational changes (**Fig. 2**).

Local temperature dependence of B-factor is null

Since the temperature-dependence of the raw B-factor (**B$_{obs}$**) was found to be similar for all protein atoms (**Fig. 4-6**), as well as for the Wilson B-factor (**Fig. 8**), we hypothesize that a uniform thermal contribution may additively influence one component of the mean-square atomic displacement, resulting in the observed raw B-factor.

Converting [Eq. 3] to a log function results in

[Eq.4] $\ln B_{obs} = \ln B_0 + k*T$

Incorporating the contribution of a thermal component B_{other} to the raw B_{obs} becomes

[Eq. 5] $\ln B_{norm} = \ln B_{obs} - \ln B_{other} + k*T$

which rearranges back to

[Eq. 6] $B_{norm} = (B_{obs} / B_{other}) * e^{(k*T)}$

Since the thermal contribution is indeed uniform among all atoms, we may estimate B_{other} as the average of the B_{obs}, i.e., B_{avg}, resulting in

[Eq. 7] $B_{other} = B_{avg}$

and thus

[Eq. 8] $B_{norm} = (B_{obs} / B_{avg}) * e^{(k*T)}$

We normalized the **B$_{obs}$** into **B$_{norm}$** by scaling it down with the average B-factor (**B$_{avg}$**) according to [**Eq. 2**]. The **B$_{norm}$** distribution profile becomes stable over a broad temperature range (100 K to 325 K), for side-chain (**Fig. 4D**), Cα (**Fig. 5D**) and N-amide (**Fig. 6D**), which were similar for all temperatures. The **B$_{norm}$** was roughly constant as a function of temperature for each amino acid side-chain (**Fig. 4E**), Cα (**Fig. 5E**) and N-amide (**Fig. 6E**). We adjusted each temperature curves for each amino acid residue side-chain (**Fig. 4E**), Cα (**Fig. 5E**) and N-amide (**Fig. 6E**) with single exponential function [**Eq. 3**]. The estimated thermal factor ^{Bnorm}**k** was found to fluctuate around zero, for side-chain (**Fig. 4F**), Cα (**Fig. 5F**) and N-amide (**Fig. 6F**). These data suggest that the scaled **B$_{norm}$** embodies only the static, temperature-independent component of the B-factor.

We further explored the hypothesis for a link between conformational changes and B-factor, as depicted in **Fig. 7**. While ^{avg}B and temperature were found to be tightly correlated (**Fig. 3**; R^2>0.8), only a loose correlation was found between thermal expansion coefficient $^{\delta c\alpha/\delta T}$**k** and **B$_0$** ($R^2$=0.371; (**Fig. 7A** and **Fig. 7C**), or between $^{C\alpha B}$**k** (R^2=0.175; **Fig. 7B** and **Fig. 7D**). While a progressive shift in conformation occurs for side-chain (**Fig. S10**), Cα (**Fig. S11**) and N-amide (**Fig. S12**), no major changes in distribution for **B$_{norm}$** seems to occur as a function of temperature.

In order to test the validity to other systems, we analyzed the RCSB for datasets available from single-crystal diffraction under similar controlled conditions and temperature as variable, as inspected from their respective published articles. We have found data from thaumatin crystal structures solved at temperatures from 25 K to 300K. These data showed increasing B-factor as a function of temperature (**Fig. 8A**), which revealed similar pattern for B$_{norm}$ distribution along the amino acid sequence (**Fig. 8B**). Thaumatin change conformation as a function of temperature (**Fig. 8C**), although the expansion

coefficient $^{\delta C\alpha/\delta T}k$ does not correlate with either the absolute B-factor (**Fig. 8D**) or the thermal factor $^{C\alpha B}k$.

Discussion

Inferring conformational changes from a given protein require testing continuous or discontinuous chemical (pH, ligands, drugs, metals, cofactors, cosolutes, point mutation, hydration, among others) and/or physical (temperature, pressure) variables which might induce dynamic conformational changes and accumulation of polymorphs. In the case of crystal structures, it may even result in dissimilar packing and space groups.

Previous work has shown that tetragonal lysozyme present changes in B-factor as a function of temperature in a non-uniform patter along the protein, with smaller changes at regions involved in symmetry-related intermolecular contacts [32,33]. Although the lysozyme (44 mg/mL) was crystallized in similar conditions reported here, at same space group ($P4_32_12$) from 0.7 M NaCl in sodium acetate pH 4.8, main-chain distances over 20 Å were found between structures solved at 298 and 100 K. However, while data collected at 100 K was obtained from crystals mounted in a glass fiber, the data obtained at 298 K was collected from crystal mounted into a capillary tube, resulting in doubts concerning the variable dictating crystallographic changes. Ribonuclease-A type III (RNAse) has been also studied at 1.5 Å resolution, over a broad range of temperature, from 98 K to 320 K [29,46]. However, each crystal was collected under dissimilar conditions of solvent composition (50 % MPD, for 98 K to 220 K, or 50 % methanol, for 240 K to 320 K), crystal mounting technique (capillary or glass fiber), diffractometer, instrument controllers and data collection procedures [29], limiting interpretation concerning the effective variable.

The reproducibility of the crystallographic model of lysozyme at high resolution (1.6 Å) has been shown to be robust when using tetragonal lysozyme obtained in sodium acetate pH 4.6 and NaCl buy hanging drop vapor diffusion technique and using two dissimilar home-source diffractometers [15]. While the final structural models showed reproducible, the raw B-factor was found non-reproducible despite their similar distribution along the polypeptide chain. The normalization of the B-factor by the average B-factor allowed obtaining reproducible B-factor patterns, suggesting that the B-factor could be influenced by variables from crystal handling to data collection. It thus became evident the need for stabilization of crystal allowing a more stable and reproducible data collection.

In the present work, we used same cryoprotectant (glycerol), mounting device (nylon loop) and stabilization (synthetic hydrocarbon grease, Apiezon®) for all dataset, allowing high repeatability of B-factors at each temperature compared to equivalent data collected in the absence of grease [15]. We have reported here a broad data collection temperature investigation of tetragonal lysozyme crystals allowing the observation of dissociation between lysozyme conformational changes and B-factor. These data were possible to obtain thanks to the strategy of crystal stabilization with a hydrocarbon grease, allowing x-ray diffraction at high resolution in a broad temperature range, from 100 K up to 325 K, without loss in lattice order and diffraction power. In fact, since we obtained high resolution (1.5 Å) through the investigated temperature range, one can assume that the B-factor is not dominated by the well-organized lattice [21,29].

As previously envisioned by Wilson [47], the B-factor is susceptible to temperature, and indeed we have confidently observed this phenomena here once solved the experimental reproducibility of the B-factor by using hydrophobic grease. However, when the B_{obs} was normalized to the B_{norm} by scaling down with the B_{avg} (from all atoms of the polypeptide chain) we observed similar B-factor distribution pattern over the polypeptide chain for the temperature range assayed here (100 K to 325 K), both for Cα (^{Cα}B), ($^{N-amide}B$) and side chain (^{SC}B). These data indicate that the inherent protein B_{norm} is not temperature-dependent, and that the total B-factor (including the unmodeled B_{Wilson}) is most likely dominated by dynamic, rigid body contributions [26,29]. The variations in the inherent protein B-factor along the polypeptide chain is likely to be influenced by the atom environment, by immediate local

interactions, and by adjacent amino acids as previously reported [17]. While local conformational changes take place in lysozyme as a function of temperature, it did not correlate with B-factor parameters (**Fig. 7**) given the distribution profile and correlation coefficient [48], suggesting a dissociation between these two parameters, also observed for the unrelated protein model thaumatin (**Fig. 8**) [10].

We envision that the raw B_{obs} is composed by the mostly thermal-insensitive atomic vibration B_A and an additive contribution of a thermal-sensitive rigid body motion B_{RB} along with other contributions to baseline B-factor rising from sources other including crystal handling and data collection, which could be minimized by using a sealant such as hydrocarbon grease.

The present study with our original data from thermal-induced changes in lysozyme (and an intervalidation with thaumatin) demonstrate that changes in B-factor does not correlate with conformational changes. While the raw B-factor may indeed correlate with temperature, the normalized B-factor does not seem to be susceptible by this variable, lacking local changes that could be related to concomitant conformational transitions in the protein.

Although we have shown the extension of our findings with lysozyme to thaumatin, and even being conducted with a rigorous triplicate for each temperature and a standard data collection and processing workflow, we used only one protein (lysozyme) at one crystallization condition and space group. While independent work is desirable to further test the conclusion drawn here and its wider application. In the meanwhile, is time to dissociate the mean-squared atomic displacement Debye-Waller factor (and the converted B-factor) from protein dynamics unless supported by direct data.

Conclusions

We have found that embedding protein crystals within hydrocarbon grease allowed stable and repeatable x-ray diffraction data collection and structural solution in a broad temperature range. We observed a lack of correlation between local protein B-factor and temperature-induced conformational changes, demonstrating a dissociation of the mean-squared deviation with conformational plasticity. More importantly, since the raw B-factor is uniformly influenced by temperature at same constant, it suggests a global, dynamic correlation with rigid body motion and/or any other scattering and crystal property. Further development should consider separation the B-factor components – atomic and system (including rigid body, scattering, among others) - contributing for the composite raw B-factor.

Materials and Methods

Material

HEWL was obtained from Fluka (Cat #62971; Lot #BCG4805V) and kept at -20°C until use. All other reagents were of analytical grade.

Methods
Protein crystallography

Test crystal: Crystals of HEWL were obtained by vapor diffusion, sitting-drop technique, using Corning 3552 plates with 1 μL HEWL 50 mg/mL in water, and 1 μL precipitant, equilibrated against 80 μL precipitant solution in the well, comprising 1.2 M NaCl, 100 mM sodium acetate pH 4.6 (adjusted either with acetic acid or NaOH), at 22 °C ± 2 °C. This method was used throughout the study in order to avoid the large variability inherent to the vapor-diffusion crystallization step [49]. Diffraction-quality crystals grew within 1 day and were used for diffraction after 2 days. Crystals were handled with 20 μm nylon CryoLoop™ (Hampton Research). Each crystal was soaked for 5 to 10 sec in the same crystallization solution from the well, supplemented with glycerol (for 30% v/v final concentration, with proportional dilution of the well solution), following by immersion into Apiezon® N hydrocarbon grease, and mounted onto the goniometer under N_{2g} stream at desired temperature for data collection. Each dataset corresponds to a single crystal at a given temperature.

Since Apiezon® N has a recommended working range of 4 to 303 K (-269 °C to 30 °C), we also tested the Apiezon® T which has recommended use range of 283 to 393 K (10 °C to 120 °C), although it provided no additional benefits, since lysozyme lost diffraction power in temperatures over 55 °C (not shown).

Diffractometer setup and data acquisition: The crystals were subjected to X-ray diffraction and data collection using CuKα radiation at constant exposure time using a 30 W air-cooled |μS microfocus source (Incoatec) mounted on D8-Venture diffractometer (Bruker; at a CENABIO-UFRJ facility) operating at 50 kV and 1.1 mA, and recorded on a Photon II detector (Bruker). Crystals were under N_{2g} stream at desired temperature with flow of 1.2 L/h, controlled by a CryoStream 800 (Oxford Cryogenics). All datasets were collected using 60 sec exposures with 0.5° oscillation/image and at leas 99% completeness assuming identical Friedel pairs, targeting 1.5 Å resolution. Data were collected, indexed, integrated and scaled using *Proteum3* (Bruker AXS Inc.), followed by analysis with *Truncate* [50].

A nonredundant set of information regarding the tetragonal ($P4_32_12$) hen lysozyme crystal structures determined by X-ray diffraction of single crystals was obtained from The Research Collaboratory for Structural Bioinformatics Protein Data Bank (RCSB PDB, rcsb.org), as of February 15, 2022 (**Supporting Information**). After filtering for entries with missing information, data was plotted according to the desired analysis (**Fig. S1**).

Data processing: Data was further processed and analyzed using *C.C.P.4.* v7.0.071 [51]. The crystal structures were solved by isomorphous replacement with 20 cycles of rigid body search with RefMac v5.8.0238 [52] [53] using PDB entry 3A8Z [54] edited for single occupancy and no water as a search model, allowing only a single sodium ion, resulting in a clear solution for a monomer in the asymmetric unit. The solution was further subjected to 10 cycles of restrained refinement using *Refmac* [52]. Real space refinement was conducted by visual inspection of both the map and the model with *C.O.O.T.* v0.8.9.2 [55], adjusting solely misplaced side chains, followed by water addition using 1.2 σ as threshold. Additional restrained refinement with 10 cycles was performed using *Refmac*. All data processing was performed

using default mode in order to avoid bias. Further data analysis was conducted with *Superpose* v1.05 from *C.C.P.4.* Global pairwise alignment of the HEWL structures was performed using ProSMART [56]. Analysis of B-factor was performed with *Baverage* (CCP4).

A summary of crystal parameters along with data collection and refinement statistics are presented in **Supporting Information**. Structural validation of the model performed with PROCHECK. All structure figures were generated with *PyMOL* v2.0 [57]. Graphics were generated with GraphPad Prism v. 8.0.2 for Windows (GraphPad Software, San Diego, California USA, www.graphpad.com). The atomic coordinates have been deposited with the Protein Data Bank and have been assigned the codes depicted in **Supporting information**, along with the RCSB validation report.

Circular Dichroism

Thermal denaturation of HEWL (20 mg/mL) was performed in 50 mM AcONa pH 4.6, in a circular dichroism spectropolarimeter (JASCO® J-810; Tokyo, Japan; LMProt-ICB-UFMG, Brazil), equipped with Peltier Jasco® temperature control system - PFD-425S (Tokyo, Japan), using a 100 µm pathlength quartz cuvette (Uvonic Instruments, Plainview, NY). All the spectra were recorded with three scans from 260 to 190 nm using a 0.2 nm spectral bandwidth, 100 nm/min scan speed, and 1 sec response time. Similar experiments with the respective blank solutions were also carried out in order to allow for proper background subtraction. The temperature was increased at a rate of 5 °C/min, and allowed equilibration for 60 sec before measurements. Temperature curve was analyzed using a sigmoidal 4 parameters function.

Direct Infusion ESI-Traveling Wave – Ion Mobility Spectrometry – Mass Spectrometry (ESI-TWIM-MS)

The lysozyme was diluted to 50 µM in 0.1 % formic acid. The ESI-TWIM-MS measurements were performed in a Traveling Wave Ion Mobility Mass Spectrometer (TWIM-MS, Synapt G1 HDMS, Waters, UK), with perfusion at 20 µL/min in positive ESI with a capillary voltage of 4.0 kV and N_2g used as mobility gas at 0.4 bar, and source temperature varying between 50 °C and 250 °C in 50 °C intervals. The data acquisition was conducted over the range of *m/z* 500 to 3,000 for 5 min. The mass spectrometer calibration was performed with phosphoric acid 0.1 % (v/v) in acetonitrile:H_2O (1:1). Other typical instrumental settings are as described previously [58]. Data were analyzed using DriftScope [59] (Waters Corporation, UK). All measurements were performed at the *Laboratory of Macromolecules – LAMAC / Life Science Metrology Division* – DIMAV, INMETRO, Rio de Janeiro, Brazil.

Structural analysis from RCSB-PDB

In order to further extend our temperature analysis, we searched the RCSB for crystallographic protein structures solved by x-ray diffraction at varying temperatures (**<400 K**) and resolution ≤ 2.0 Å, deposited between 01.01.2000 and 02.28.2022. Further search parameters included average B-factor and Wilson B-factor estimate ≤ 100 $Å^2$ and refinement R-free ≤ 0.56. This search returned 24,755 entries. These entries were searched for works showing works reporting crystal structures solved with at least **five** equivalent entries (such as same protein, pH, crystallization condition, space group) under data collection temperatures spanning most regularly for at least 200 K.

The RCSB was also searched for hen egg white **lysozyme** crystal structures solved by x-ray diffraction in space group **P4$_3$2$_1$2** up to 02.15.2022. This search returned 516 entries which were analyzed for refinement and structural parameters as reported in **Supporting Information**.

Data analysis

Graphics were generated with GraphPad Prism v. 8.0.2 for Windows (GraphPad Software, San Diego, California USA, www.graphpad.com). Significance was considered when p<0.005 [60].

Acknowledgments

We would like to thanks Prof. Mariana Quezado (UFMG) for valuable discussion along the development of this work. We thank Dr. Sandra Scapin (INMETRO) for helpful assistance during ESI-IMS-MS data collection.

Funding

This study was supported by Fundação Carlos Chagas Filho de Amparo à Pesquisa do Estado do Rio de Janeiro (grants E-26/202.998/2017-BOLSA, E-26/200.833/2021-BOLSA, E-26/010.001434/2019-Tematico and E-26/210.195/2020 to LMTRL), by the Conselho Nacional de Desenvolvimento Científico (grant PQ2/311582/2017-6 and 313179/2020-4, to LMTRL), by the Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (CAPES, Finance Code #001) and the Programa Nacional de Apoio ao Desenvolvimento da Metrologia, Qualidade e Tecnologia (PRONAMETRO, 01/2018, to LMTRL) from the Instituto Nacional de Metrologia, Qualidade e Tecnologia (INMETRO). The funding agencies had no role in the study design, data collection and analysis, or decision to publish or prepare of the manuscript.

Conflict of interest

The authors declare no financial or intellectual conflicts of interest with the contents of this article.

Author Contributions

Fernando de Sá Ribeiro - Data curation; Formal analysis; Investigation; Methodology; Validation; Visualization; Writing - review & editing.
Luís Maurício T. R. Lima - Conceptualization; Data curation; Formal analysis; Funding acquisition; Investigation; Methodology; Project administration; Supervision; Validation; Visualization; Roles/Writing - original draft; Writing - review & editing.

References

1. Hipp, M. S., Kasturi, P. & Hartl, F. U. The proteostasis network and its decline in ageing. *Nat. Rev. Mol. Cell Biol.* **20**, 421–435 (2019).

2. Cramer, P. AlphaFold2 and the future of structural biology. *Nat. Struct. Mol. Biol.* **28**, 704–705 (2021).

3. Thorp, H. H. Proteins, proteins everywhere. *Science* **374**, 1415 (2021).

4. Moore, P. B., Hendrickson, W. A., Henderson, R. & Brunger, A. T. The protein-folding problem: Not yet solved. *Science* (2022) doi:10.1126/science.abn9422.

5. Levantino, M. *et al.* Ultrafast myoglobin structural dynamics observed with an X-ray free-electron laser. *Nat. Commun.* **6**, 6772 (2015).

6. Pandey, S. *et al.* Time-resolved serial femtosecond crystallography at the European XFEL. *Nat. Methods* **17**, 73–78 (2020).

7. Basso, S., Fitch, A. N., Fox, G. C., Margiolaki, I. & Wright, J. P. High-throughput phase-diagram mapping via powder diffraction: a case study of HEWL versus pH. *Acta Crystallogr. D Biol. Crystallogr.* **61**, 1612–1625 (2005).

8. Takayama, Y. & Nakasako, M. A few low-frequency normal modes predominantly contribute to conformational responses of hen egg white lysozyme in the tetragonal crystal to variations of molecular packing controlled by environmental humidity. *Biophys. Chem.* **159**, 237–246 (2011).

9. Urayama, P., Phillips, G. N. & Gruner, S. M. Probing substates in sperm whale myoglobin using high-pressure crystallography. *Struct. Lond. Engl. 1993* **10**, 51–60 (2002).

10. Warkentin, M., Badeau, R., Hopkins, J. B. & Thorne, R. E. Spatial distribution of radiation damage to crystalline proteins at 25-300 K. *Acta Crystallogr. D Biol. Crystallogr.* **68**, 1108–1117 (2012).

11. Keedy, D. A. Journey to the center of the protein: allostery from multitemperature multiconformer X-ray crystallography. *Acta Crystallogr. Sect. Struct. Biol.* **75**, 123–137 (2019).

12. Perspectives: Cracking the Phase Problem - The Nobel Prize in Chemistry 1962. *NobelPrize.org* https://www.nobelprize.org/prizes/chemistry/1962/perspectives/.

13. Fávero-Retto, M. P., Palmieri, L. C., Souza, T. A. C. B., Almeida, F. C. L. & Lima, L. M. T. R. Structural meta-analysis of regular human insulin in pharmaceutical formulations. *Eur. J. Pharm. Biopharm. Off. J. Arbeitsgemeinschaft Pharm. Verfahrenstechnik EV* **85**, 1112–1121 (2013).

14. Su, Z. *et al.* Reproducibility of protein x-ray diffuse scattering and potential utility for modeling atomic displacement parameters. *Struct. Dyn. Melville N* **8**, 044701 (2021).

15. Ramos, N. G., Sarmanho, G. F., de Sá Ribeiro, F., de Souza, V. & Lima, L. M. T. R. The reproducible normality of the crystallographic B-factor. *Anal. Biochem.* 114594 (2022) doi:10.1016/j.ab.2022.114594.

16. Carugo, O. B-factor accuracy in protein crystal structures. *Acta Crystallogr. Sect. Struct. Biol.* **78**, 69–74 (2022).

17. Karplus, P. A. & Schulz, G. E. Prediction of chain flexibility in proteins: A tool for the selection of peptide antigens. *Naturwissenschaften* **72**, 212–213 (1985).

18. Rupp, B. *Biomolecular Crystallography: Principles, Practice, and Application to Structural Biology*. (Garland Science, 2009).

19. Giacovazzo, C., Monaco, H. L. & Artioli, G. *Fundamentals of Crystallography: 15*. (Oxford University Press, USA, 2011).

20. *International Tables for Crystallography: Crystallography of biological macromolecules*. vol. F (International Union of Crystallography, 2012).

21. Ringe, D. & Petsko, G. A. Study of protein dynamics by X-ray diffraction. *Methods Enzymol.* **131**, 389–433 (1986).

22. Doucet, J. & Benoit, J. P. Molecular dynamics studied by analysis of the X-ray diffuse scattering from lysozyme crystals. *Nature* **325**, 643–646 (1987).

23. Pérez, J., Faure, P. & Benoit, J.-P. Molecular Rigid-Body Displacements in a Tetragonal Lysozyme Crystal Confirmed by X-ray Diffuse Scattering. *Acta Crystallogr. D Biol. Crystallogr.* **52**, 722–729 (1996).

24. Van Benschoten, A. H. *et al.* Measuring and modeling diffuse scattering in protein X-ray crystallography. *Proc. Natl. Acad. Sci. U. S. A.* **113**, 4069–4074 (2016).

25. Wall, M. E. Internal protein motions in molecular-dynamics simulations of Bragg and diffuse X-ray scattering. *IUCrJ* **5**, 172–181 (2018).

26. de Klijn, T., Schreurs, A. M. M. & Kroon-Batenburg, L. M. J. Rigid-body motion is the main source of diffuse scattering in protein crystallography. *IUCrJ* **6**, 277–289 (2019).

27. Sun, Z., Liu, Q., Qu, G., Feng, Y. & Reetz, M. T. Utility of B-Factors in Protein Science: Interpreting Rigidity, Flexibility, and Internal Motion and Engineering Thermostability. *Chem. Rev.* **119**, 1626–1665 (2019).

28. Carugo, O. How large B-factors can be in protein crystal structures. *BMC Bioinformatics* **19**, 61 (2018).

29. Tilton, R. F., Dewan, J. C. & Petsko, G. A. Effects of temperature on protein structure and dynamics: x-ray crystallographic studies of the protein ribonuclease-A at nine different temperatures from 98 to 320K. *Biochemistry* **31**, 2469–2481 (1992).

30. Kim, C. U., Tate, M. W. & Gruner, S. M. Protein dynamical transition at 110 K. *Proc. Natl. Acad. Sci.* **108**, 20897–20901 (2011).

31. Daniels, B. V., Schoenborn, B. P. & Korszun, Z. R. Myoglobin Solvent Structure at Different Temperatures. *Neutrons Biol.* 325–331 (1996) doi:10.1007/978-1-4615-5847-7_28.

32. Young, A. C. M., Dewan, J. C., Nave, C. & Tilton, R. F. Comparison of radiation-induced decay and structure refinement from X-ray data collected from lysozyme crystals at low and ambient temperatures. *J. Appl. Crystallogr.* **26**, 309–319 (1993).

33. Young, A. C. M., Tilton, R. F. & Dewan, J. C. Thermal expansion of hen egg-white lysozyme. *J. Mol. Biol.* **235**, 302–317 (1994).

34. Dunlop, K. V., Irvin, R. T. & Hazes, B. Pros and cons of cryocrystallography: should we also collect a room-temperature data set? *Acta Crystallogr. D Biol. Crystallogr.* **61**, 80–87 (2005).

35. Dewan, J. C. & Tilton, R. F. Greatly reduced radiation damage in ribonuclease crystals mounted on glass fibers. *J. Appl. Crystallogr.* **20**, 130–132 (1987).

36. de la Mora, E. *et al.* Radiation damage and dose limits in serial synchrotron crystallography at cryo- and room temperatures. *Proc. Natl. Acad. Sci. U. S. A.* **117**, 4142–4151 (2020).

37. Ihara, K. *et al.* Isoprenoid-chained lipid EROCOC17+4: a new matrix for membrane protein crystallization and a crystal delivery medium in serial femtosecond crystallography. *Sci. Rep.* **10**, 19305 (2020).

38. Hope, H. Cryocrystallography of biological macromolecules: a generally applicable method. *Acta Crystallogr. B* **44 (Pt 1)**, 22–26 (1988).

39. Hope, H. *et al.* Cryocrystallography of ribosomal particles. *Acta Crystallogr. B* **45 (Pt 2)**, 190–199 (1989).

40. Nam, K. H. Stable sample delivery in viscous media via a capillary for serial crystallography. *J. Appl. Crystallogr.* **53**, 45–50 (2020).

41. Park, J. *et al.* Polyacrylamide injection matrix for serial femtosecond crystallography. *Sci. Rep.* **9**, 2525 (2019).

42. Sugahara, M. *et al.* Grease matrix as a versatile carrier of proteins for serial crystallography. *Nat. Methods* **12**, 61–63 (2015).

43. Nam, K. H. Shortening injection matrix for serial crystallography. *Sci. Rep.* **10**, 107 (2020).

44. Nam, K. H. Lard Injection Matrix for Serial Crystallography. *Int. J. Mol. Sci.* **21**, 5977 (2020).

45. Nam, K. H. Beef tallow injection matrix for serial crystallography. *Sci. Rep.* **12**, 694 (2022).

46. Bf, R., Am, S., D, R. & Ga, P. Crystalline ribonuclease A loses function below the dynamical transition at 220 K. *Nature* **357**, (1992).

47. Wilson, A. J. C. Determination of Absolute from Relative X-Ray Intensity Data. *Nature* **150**, 152–152 (1942).

48. Anscombe, F. J. Graphs in Statistical Analysis. *Am. Stat.* **27**, 17–21 (1973).

49. Newman, J., Xu, J. & Willis, M. C. Initial evaluations of the reproducibility of vapor-diffusion crystallization. *Acta Crystallogr. D Biol. Crystallogr.* **63**, 826–832 (2007).

50. Popov, A. N. & Bourenkov, G. P. Choice of data-collection parameters based on statistic modelling. *Acta Crystallogr. Sect. D* **59**, 1145–1153 (2003).

51. Winn, M. D. *et al.* Overview of the CCP4 suite and current developments. *Acta Crystallogr. D Biol. Crystallogr.* **67**, 235–242 (2011).

52. Kovalevskiy, O., Nicholls, R. A., Long, F., Carlon, A. & Murshudov, G. N. Overview of refinement procedures within REFMAC5: utilizing data from different sources. *Acta Crystallogr. Sect. Struct. Biol.* **74**, 215–227 (2018).

53. Vagin, A. & Teplyakov, A. Molecular replacement with MOLREP. *Acta Crystallogr. D Biol. Crystallogr.* **66**, 22–25 (2010).

54. Ueno, T. *et al.* Elucidation of Metal-Ion Accumulation Induced by Hydrogen Bonds on Protein Surfaces by Using Porous Lysozyme Crystals Containing RhIII Ions as the Model Surfaces. *Chem. - Eur. J.* **16**, 2730–2740 (2010).

55. Bond, P. S., Wilson, K. S. & Cowtan, K. D. Predicting protein model correctness in Coot using machine learning. *Acta Crystallogr. Sect. Struct. Biol.* **76**, 713–723 (2020).

56. Nicholls, R. A., Fischer, M., McNicholas, S. & Murshudov, G. N. Conformation-independent structural comparison of macromolecules with ProSMART. *Acta Crystallogr. D Biol. Crystallogr.* **70**, 2487–2499 (2014).

57. The PyMOL Molecular Graphics System.

58. de Araújo, T. S. *et al.* Biophysical characterization of two commercially available preparations of the drug containing Escherichia coli L-Asparaginase 2. *Biophys. Chem.* 106554 (2021) doi:10.1016/j.bpc.2021.106554.

59. Williams, J. P., Lough, J. A., Campuzano, I., Richardson, K. & Sadler, P. J. Use of ion mobility mass spectrometry and a collision cross-section algorithm to study an organometallic ruthenium anticancer complex and its adducts with a DNA oligonucleotide. *Rapid Commun. Mass Spectrom. RCM* **23**, 3563–3569 (2009).

60. Ioannidis, J. P. A. The Proposal to Lower P Value Thresholds to .005. *JAMA* **319**, 1429–1430 (2018).

Figures

Figure 1. Global temperature-dependent conformational changes.
Lysozyme structure models solved from data collected at varying temperatures from 100 K to 325 K were aligned and are showed for A) backbone and B) side chains. Colors represents data collection temperature as indicated. n=3 per temperature.

Global (average values for residues 1-129) changes in conformation were inferred from superposition with a reference structure at 100 Kelvin. Continuous line are first order linear regression and dotted lines are 95 % confidence interval. We observed a small (< 1 Å) but progressive and significant increase in global conformational changes as inferred for distance changes in (C) SC (0.001566 Å.K-1; p≤ 0.0001), (D) Cα (0.001110 Å.K^{-1}; p≤ 0.0001) and (E) amide nitrogen (0.0007971 Å.K^{-1}; p= 0.0037) from reference structure. No significant temperature-dependent changes were inferred for total content in helix (p=0.4260), beta (p=0.3935), coil (p=0.0107) and turn (p=0.0458) as shown in (F). Symbol is average and bar is standard deviation (n=3).

Figure 2. Temperature induced conformational changes in lysozyme .
Structures were aligned with a reference structure at 100 K using *Superpose* (CCP4) and distances between Cα were obtained.
A) Changes in Cα distances along protein sequence at varying temperatures (different colors).
B) Changes in Cα distances as a function of temperature for each amino acid residue (different colors) protein sequence at varying temperatures. Lines are first order linear regression, from which were obtained their thermal-dependence of conformational change (δCα; angular coefficient of linear regression).
C) Distribution of thermal-dependent conformational change (δCα) along protein sequence.
D) Changes in Cα distances as a function of temperature for each amino acid residue (different colors) protein sequence at varying temperatures. Lines are exponential non-linear regression, from which were obtained their exponential thermal constant ($^{\delta C\alpha}k$).
E) Distribution of exponential thermal constant ($^{\delta C\alpha}k$) of conformational change along protein sequence. Symbol is average and bar is standard deviation (n=3).

$$B_{obs} = B_0 * e^{(k.T)}$$

A

Wilson$B$, $Å^2$

Temperature, K

$B_0 = 5.682 \pm 0.502$ $Å^2$
$k = 0.004427 \pm 0.000336$ K^{-1}
$R^2 = 0.865$

B

sc$B_{avg1-129}$, $Å^2$

Temperature, K

$B_0 = 5.390 \pm 0.474$ $Å^2$
$k = 0.004598 \pm 0.000333$ K^{-1}
$R^2 = 0.877$

C

C$\alpha$$B_{avg1-129}$, $Å^2$

Temperature, K

$B_0 = 4.874 \pm 0.498$ $Å^2$
$k = 0.004555 \pm 0.000387$ K^{-1}
$R^2 = 0.837$

D

Namide$B_{avg1-129}$, $Å^2$

Temperature, K

$B_0 = 4.740 \pm 0.512$ $Å^2$
$k = 0.004532 \pm 0.000410$ K^{-1}
$R^2 = 0.819$

Figure 3. Temperature-dependent changes in B-factor (B).

Overall changes in B-factor as a function of temperature for **A)** Wilson plot, **B)** side-chains, **C)** Cα and **D)** amide nitrogen.

Values corresponds to the extrapolated B factor at zero Kelvin (B_0) and the thermal coefficient k, as inferred from single exponential (equation above panels) non-linear regression of their respective panels (solid lines; dotted lines are 95% confidence interval). Symbol is average and bar is standard deviation (n=3).

Figure 4. Temperature-dependent changes in side-chain B-factor

A) B-factor distribution for side chain at varying temperatures (different colors).

B) Changes in raw B-factor as a function of temperature for each amino acid side chain (different colors). Lines are exponential non-linear regression, from which were obtained the raw B_0 and the thermal constant k.

C) Distribution of raw B_0 and raw k along protein sequence.

D) Normalized B-factor (B_{norm}) distribution for side chain at varying temperatures (different colors).

E) Changes in B_{norm} as a function of temperature for each amino acid side chain (different colors). Lines are exponential non-linear regression, from which was obtained the thermal constant ^{norm}k.

C) Distribution of ^{norm}k along protein sequence.

Symbol is average and bar is standard deviation (n=3).

Figure 5. Temperature-dependent changes in Cα B-factor
A) B-factor distribution for Cα at varying temperatures (different colors).
B) Changes in raw B-factor as a function of temperature for each amino acid Cα (different colors). Lines are exponential non-linear regression, from which were obtained the raw B_0 and the thermal constant k.
C) Distribution of raw B_0 and raw k along protein sequence.
D) Normalized B-factor (B_{norm}) distribution for Cα at varying temperatures (different colors).
E) Changes in B_{norm} as a function of temperature for each amino acid Cα (different colors). Lines are exponential non-linear regression, from which was obtained the thermal constant ^{norm}k.
F) Distribution of ^{norm}k along protein sequence.
Symbol is average and bar is standard deviation (n=3).

Figure 6. Temperature-dependent changes in amide nitrogen (N-amide) B-factor
A) B-factor distribution for N-amide α at varying temperatures (different colors).
B) Changes in raw B-factor as a function of temperature for each amino acid Cα (different colors). Lines are exponential non-linear regression, from which were obtained the raw B_0 and the thermal constant k.
C) Distribution of raw B_0 and raw k along protein sequence.
D) Normalized B-factor (B_{norm}) distribution for N-amide at varying temperatures (different colors).
E) Changes in B_{norm} as a function of temperature for each amino acid N-amide (different colors). Lines are exponential non-linear regression, from which was obtained the thermal constant ^{norm}k.
F) Distribution of ^{norm}k along protein sequence.
Symbol is average and bar is standard deviation (n=3).

Figure 7. Correlation between changes in conformation B-factor parameters.
Correlation between B-factor parameters (B_0; panels **A** and **C**) and ($^{C\alpha B}k$) and temperature-induced
conformational changes constant ($^{\delta C\alpha/\delta T}k$; panels **B** and **D**). Data correspond to calculated $^{\delta C\alpha}B_0$ from **Fig.
5C** and $^{\delta C\alpha/\delta T}k$ from **Fig. 2C**. Data are plotted in linear (**A** and **B**) or log scale (**C** and D) for ease of
observation of data span.
Lines correspond to first-order linear regression (continuous) and 95 % confidence interval (dotted).

Figure 8. Temperature-dependence of B-factor and conformation of Thaumatin. X-ray structures solved at 25 K (PDB ID 4ek0), 100 K (PDPB ID 4ekb), 180 K (PDB ID 4eko), 240 K (PDB ID 4el2) and 300 K (PDB ID 4el7), as depicted in the Figure legends.

A) B-factor distribution for Cα at varying temperatures (different colors).
B) Normalized B-factor (B_{norm}) distribution for Cα at varying temperatures (different colors).
C) Changes in Cα distances along protein sequence at varying temperatures (different colors).
D) Correlation between B-factor and backbone thermal expansion coefficient ($^{\delta C\alpha/\ \delta T}k$) at varying temperatures (different colors). Each point corresponds to one amino acid.
E) Dependence of thermal factor ($^{C\alpha B}k$) on the thermal expansion coefficient ($^{\delta C\alpha/\ \delta T}k$).
Data from [10].