Portable X-ray Fluorescence for Measuring Lead in the Hair of

Wild Mammals

Evie M. Jones, [*,†] Andrew J. Bengsen, [‡,§] Aaron J. Specht, [⬜] Amelia J. Koch, [†,⊥] Rodrigo K.

Hamede, [†] Menna E. Jones, [†] and Jordan O. Hampton [#,∇]

[†] School of Natural Sciences, University of Tasmania, Private Bag 55, Hobart, TAS, 7001, Australia

[‡] Vertebrate Pest Research Unit, NSW Department of Primary Industries, 4 Marsden Park Rd, Calala, New

South Wales 2340, Australia

[§] Biosphere Environmental Consultants, Tamworth, NSW 2340, Australia

[⬜] School of Health Sciences, Purdue University, 610 Purdue Mall, West Lafayette, Indiana 47907, United States

[⊥] Forest Practices Authority, 30 Patrick St, Hobart, TAS, 7001, Australia

[#] Faculty of Science, University of Melbourne, Parkville, Victoria 3052, Australia

[∇] Harry Butler Institute, Murdoch University, 90 South Street, Murdoch, Western Australia 6150, Australia

[*] Corresponding author.

Email address: evie.jones@utas.edu.au

Abstract

Lead exposure threatens scavenging wildlife globally. For inexpensive estimation of lead concentration in bones from avian scavengers, portable X-ray fluorescence (XRF) devices have been trialed. However, portable XRF has not been validated for tissue lead measurement in non-human mammalian scavengers. We evaluated portable XRF for hair lead measurement in the endangered Tasmanian devil (*Sarcophilus harrisii*). We first analyzed large (~1.0 g) hair samples from 39 deceased animals from southeastern Tasmania via portable XRF and then inductively-coupled plasma mass spectrometry (ICP-MS) (validation study). We then measured lead concentrations via portable XRF only in 61 small (~0.1 g) hair samples from live devils from a plantation landscape (plantation study). Portable XRF measures of hair lead levels were positively correlated with ICP-MS values ($R^2 = 0.61$). In the validation study, 95% of sampled Tasmanian devils had relatively low lead levels (< 2 mg/kg), but the remaining two showed elevated lead levels (> 15 mg/kg). Mean lead levels in the plantation study and validation study were not significantly different. Our preliminary results suggest that portable XRF can provide a useful measure of differences in lead levels in wildife hair over a coarse scale. We provide recommendations for further research and refinement of this method.

Keywords ecotoxicology • marsupial • non-invasive • Pb • scavenger •Tasmanian devil

Synopsis

Portable XRF can provide inexpensive and non-destructive analysis of environmental contaminants in wildlife. We present the first evaluation of portable XRF for analysis of lead contamination in mammalian scavenger hair.

INTRODUCTION

Lead is one of the best-known toxicants, with deleterious effects on humans, animals and plants. Recent research has identified lead-based ammunition as a major source of anthropogenic environmental lead globally,[1,2] representing a significant health hazard to wildlife[3] and potentially humans[4], leading to calls for a transition to non-toxic alternatives[5,6]. This source of lead is a particular threat to scavenging wildlife[7]. As a result of exposure to lead-based ammunition, numerous species of avian scavengers have been found with elevated tissue lead concentrations[3,8,9]. Impacts of lead exposure on avian scavengers include increased mortality[10], suppressed population growth rates[11], altered spatial behavior[12] and inhibition of enzymes[13], but little is known about the health impacts of lead exposure on wild mammalian scavengers. Lead exposure is a global issue, but relatively little research has been conducted in Australia[6]. The exceptions to this trend have been studies of Australia's top avian predator, the wedge-tailed eagle (*Aquila audax*)[14–16], and one marsupial scavenger, the Tasmanian devil (*Sarcophilus harrisii*)[17,18]. However, many more wildlife species may be at risk in Australia, given that wildlife shooting (culling, hunting and commercial harvesting) is ubiquitous in Australia, and relies almost exclusively on lead-based ammunition[6,19].

While scavenging birds have been relatively well-studied, comparatively few studies have assessed the effects of lead pollution from ammunition on wild mammalian scavengers[20]. The notable exception to this trend has been studies of bears (*Ursus* spp.) in the Northern Hemisphere[21–24]. One of the few studies from the Southern Hemisphere demonstrated harmful blood lead levels in Tasmanian devils in captivity[25]. The Tasmanian devil is a morphologically specialized scavenger and the largest marsupial carnivore in the world[26]. They are endangered, with population declines of over 80% since 1996 caused by a transmissible cancer, devil facial tumor disease[27]. The low population size of the species and

their reliance on scavenging makes the potential impact of lead contamination particularly

concerning.

Due to their endangered status, the Tasmanian devil is the focus of intense conservation

efforts, including proactive attempts to identify threatening processes that may prevent or

delay their recovery, such as habitat modification and vehicle collisions[28–30]. There are

concerns that lead exposure may also threaten the recovery of devils. The devil populations

that are likely to be most affected by lead pollution are those inhabiting modified

anthropogenic landscapes where culling of browsing mammals is common. For instance, in

Tasmanian timber plantations (henceforth 'plantations'), culling of marsupial browsers is

regularly conducted prior to planting in coupes to protect seedlings[31]. The browsers targeted

include brushtail possums (*Trichosurus vulpecula*), Bennett's wallabies (*Notamacropus*

rufogriseus), and rufous-bellied pademelons (*Thylogale billardierii*)[32]. All three species

constitute primary prey for devils[33], and currently culling methods involve the use of lead-

based ammunition[18].

The studies of Hivert et al. (2018)[17] and Hutchinson et al. (2023)[18] measured lead

exposure in wild devils via blood and liver, respectively, and reported relatively low lead

concentrations. However, these studies used samples from across Tasmania. No study has

specifically investigated lead exposure in devils living in a landscape subjected to regular

culling of browsers. To our knowledge, all past studies on lead exposure in scavenging

mammals have used invasive sampling, whether samples were collected from live or dead

animals. Tissue types analyzed have included blood[21,22,24,25,34], teeth[23] and liver[18,35]. However,

recent reviews have highlighted the value of non-invasive sampling of heavy metal exposure

in wild mammals[36].

One non-invasive approach to investigating heavy metal exposure in mammals is through

hair analysis, a technique used by some recent studies on scavengers such as brown bears

(*Ursus arctos*)[37] and non-scavenging species such as fruit bats (*Pteropus* spp.)[38], Malay civets

(*Viverra tangalunga*)[39] and roe deer (*Capreolus capreolus*)[40]. Hair reflects an individual's

heavy metal exposure during the period of hair growth[41]. Collecting hair samples from wild

animals can be done remotely using hair snares[42] or during handling of captured wild

animals[38].

Most studies on lead exposure in mammalian scavengers have used *ex situ* analysis via

inductively-coupled plasma mass spectrometry (ICP–MS) in a laboratory[22–24], which is

expensive and time-consuming. X-ray fluorescence (XRF) is another option for analyzing

lead in tissues such as hair[43], horn[44] and bone[45,46], and has long been used in human

medicine[47]. The XRF device analyzes the elemental content of samples through the

fluorescence produced by X-rays from a low energy x-ray tube (50 kV). XRF methods have

been used to determine lead levels in the hair of humans[43,48,49] but not, to our knowledge,

other mammals.

Portable XRF units have recently become commercially available. As they are handheld

and battery-powered, these units are relatively inexpensive, quick and practical for use in the

field, with most functions taking only three minutes to measure samples[14]. Advantages of

portable XRF over ICP-MS include lower cost and non-destructive analysis of samples[14,50].

One important disadvantage of XRF is that the instrument emits ionizing radiation with

corresponding risks associated with health and safety for users[51]. Portable XRF has been

validated for measurement of lead in bones of several avian species, showing strong

correlations with ICP-MS values, such as wedge-tailed eagles[14], California condors

(*Gymnogyps californianus*)[46], Pacific black ducks (*Anas superciliosa*)[52], and common loons

(*Gavia immer*)[45]. However, portable XRF has not been validated for measurement of lead in

the hair of non-human mammals.

Here, we investigated hair lead concentrations in wild Tasmanian devils in two stages. The

first stage was a validation study (comparing portable XRF to traditional ICP-MS methods)

and the second stage was a study of lead exposure in a forestry plantation landscape using

XRF. Our aims were: 1) to determine if portable XRF measures of hair lead levels in

Tasmanian devils are similar enough to ICP-MS results that they can be used as a proxy, 2) to

determine the usefulness of portable XRF as a tool to rapidly identify lead exposure levels in

mammals through non-destructive analysis of hair samples, and 3) to use portable XRF to

non-invasively identify lead exposure in an endangered mammalian scavenger living in a

landscape subject to browser culling operations.

MATERIALS AND METHODS

Validation study. Hair from Tasmanian devils (n = 39) was collected from archived

collections held in the Tasmanian Museum and Art Gallery (TMAG) and the University of

Tasmania (UTAS). Road-killed devils were collected between April 2017 and February 2022

from across Tasmania, concentrated in the southeast, which is dominated by native forest and

agricultural land (Figure 1). Approximately 1 g of hair was shaved from whole frozen devil

carcasses and transported in ice packs to the laboratory for analysis. Lead concentrations in

each hair sample were analyzed with both portable XRF and ICP-MS. As this study aimed in

part to validate portable XRF for use in the field, we did not wash hair samples prior to

analysis, as this would be impractical in the field.

Figure 1. Map of Tasmania, showing where 33 of the 39 Tasmanian devils (*Sarcophilus harrisii*) in
the validation study were collected; and Surrey Hills, a eucalypt plantation where all the hair samples
for the plantation study were collected. The collection locations of the remaining six devils in the
validation study is unknown.

Forestry plantation landscape study. Relatively small (~0.1 g) hair samples were
collected from 61 live wild devils within Surrey Hills, a forestry plantation landscape in
northwest Tasmania (-41.420S, 145.814E) (Figure 1). Surrey Hills is an intensively managed
forestry landscape dominated by *Eucalyptus nitens* plantations[28]. Culling, using lead-based
ammunition, is regularly undertaken at the site prior to planting in coupes to protect seedlings
from browsing by three marsupial species: Bennett's wallaby, rufous-bellied pademelon, and
brushtail possum. All three species are common prey of devils[33] and there is evidence that
culled browsers are scavenged by devils[32]. Culling occurs year-round, and devils are highly
reliant on scavenging throughout the year[53], therefore differences in sampling times are
unlikely to have had a major impact on hair lead levels in devils. Between February 2021 and
March 2022, we captured devils in custom-made 80 cm by 30 cm PVC pipe traps, shaved a 1
cm^2 patch of fur from their right shoulders, and recorded the age (from tooth wear) and sex of

the animal. These samples were analyzed using XRF only, as they were not large enough for

traditional ICP-MS methods. As above, samples were not washed prior to analysis.

XRF measurements. For XRF analysis of hair samples, we used almost identical methods

to those used in past XRF studies of lead exposure in bird bones[14,46,47,52,54] – see those studies

for more detailed descriptions of the process. The portable XRF device used was a Niton

XL3t GOLDD+ (Thermo☐Fisher Scientific, Omaha, USA), with the same settings, filter

combinations and read times as those used to analyze eagle bones in Hampton et al. (2021)[14].

To analyze the spectra produced, we used standard procedures as described in past

studies[14,55].

ICP-MS measurements. The same hair samples (validation study samples only) were

then sent to a commercial laboratory (Edith Cowan University Analytical Facility, Joondalup,

Australia) to determine their lead concentrations via inductively coupled plasma mass

spectrometry (ICP-MS), as per standard procedure in wildlife toxicology studies. Briefly, hair

samples were first freeze-dried, then underwent acid digestion and were homogenized and

processed as per liver and bone samples in Pay et al. (2021)[16]. Likewise, lead concentrations

from the samples were analyzed with ICP-MS using the same methods as Pay et al. (2021)[16].

We used the same Certified Reference Material (CRM) as Pay et al. (2021)[16] as a positive

control. The CRM was digested twice, and two ICP-MS readings were made on each

digestion, producing an average accuracy of 100.2%. We created duplicate reads by re-

analyzing every fifth or sixth sample, yielding an average relative standard deviation (RSD)

of 2.2%. We also conducted duplicate blind sample digestions on five randomly selected

samples (average RSD = 2.0%). The lead concentrations were measured in mg/kg of dry

weight (equivalent to ppm). Limit of detection (LOD) was 0.0015 mg/kg and limit of

quantification (LOQ) was 0.005 mg/kg. No samples had ICP-MS lead concentrations below

the LOD or LOQ.

Statistical analysis. We identified the uncertainty values for XRF measurements as per

past studies[45]. We calculated the uncertainty (σ) of each measurement using the following

function **(eq 1)**:

$$\sigma = \frac{c \times \sqrt{\frac{BKG + Gross}{t}}}{NET}$$

where c is concentration, *BKG* is the background counts as estimated by our fitting, Gross is

the gross counts, t is measurement time, and *NET* is net lead counts from a Gaussian

function[14]. The uncertainty values for XRF give the confidence intervals associated with a

point estimate of elemental concentration. We retained the negative values for lead in hair

because they represent a point estimate of hair lead concentration incorporating

uncertainty[56,57].

We used linear regression to examine the relationship between ICP-MS-derived and XRF-

derived lead concentrations for samples tested via both methods (n = 39). We square-root

transformed both variables to minimize heteroscedasticity, and estimated the relationship

using the following function **(eq 2):**

$$\sqrt{ICP_i} = \beta_0 + \beta_1(\sqrt{XRF_i}) + \epsilon_i$$

where β_0 is the intercept, β_1 is the slope of the relationship, and ϵ_i is the residual error. We

evaluated predictive competence of the regression model with 10-fold cross-validation,

repeated three times. Modelling was conducted using the caret package[58] in R[59].

For the plantation study, we used the equation from the regression analysis (Equation 3

below) to predict ICP-MS values from XRF measurements of lead levels in devil hair. We

used a t-test to identify any difference in lead levels between devils in the validation and

plantation study, and a two-way analysis of variance (ANOVA) to test for differences in lead

levels among devils of different sexes and ages in the plantation study (juvenile < 1 year,

subadult 1–2 years, adult > 2 years). Additionally, we constructed linear models with the

predictors of sex and age to look for any influence of these parameters on predicted ICP-MS

lead levels in devils, using Akaike's Information Criterion (AIC) values to select the best

fitting model[60]. Lead level values were log transformed for these analyses, which were

undertaken in the R statistical environment[59].

RESULTS

We found a positive relationship between lead concentrations in the hair of Tasmanian devils

estimated using portable XRF and ICP-MS (n = 39) (Figure 2). The regression model showed

a reasonable correlation between portable XRF and ICP-MS values, explaining 61% of the

variability in the data over the range of values observed. Expected ICP-MS estimates were

predicted from portable XRF estimates across the range of data from the plantation study (n =

61) using the following function (**eq 3**):

$$ICP\text{-}MS_i = (\beta_0 + \beta_1(\sqrt{XRF_i}))^2$$

where β_0 = -0.205 (SE = 0.11) and β_1 = 0.907 (SE = 0.072) (R^2 = 0.61, t_{37} = 12.651, p <

0.001; RMSE = 0.45).

As the two highest values exerted a high degree of influence over the form of the

regression, we ran a second model excluding these values. This model showed limited,

though still significant, correlation between XRF and ICP-MS values, explaining 43% of the

variability in the data (R^2 = 0.43, t_{37} = 1.869, p < 0.001; RMSE = 0.28).

Most of the samples (94.87% as measured via either method) showed lead concentrations

of < 3 mg/kg. However, the two points with the highest lead levels (> 15 mg/kg) measured

consistently high with both XRF and ICP-MS (Figure 2). XRF estimates tended to be higher

than ICP-MS: mean concentrations were 2.33 mg/kg when measured via XRF versus 1.65

mg/kg when measured via ICP-MS (Table 1).

Figure 2. Correlation between portable XRF and ICP-MS lead concentration measurements in

Tasmanian devil (*Sarcophilus harrisii*) hair, including all observed values. Shaded polygon represents

the 95% prediction interval.

Table 1. Lead Concentration in Hair Samples from 39 Tasmanian Devils (*Sarcophilus Harrisii*),

Estimated using Inductively-Coupled Mass Spectrometry (ICP-MS) and Portable X-Ray Fluorescence

(XRF)

	Mean	**Min**	**Max**	**SD**
ICP-MS (mg/kg)	1.65	0.12	25.34	4.61
Portable XRF (mg/kg)	2.33	0.08	25.16	4.76
Portable XRF uncertainty (mg/kg)	0.58	0.20	1.97	0.39

Table 2 shows portable XRF hair lead measurements and uncertainties from devils in the

plantation landscape. These data were converted to ICP-MS values using equation 3. Two

negative values (-1.39 and -0.48 mg/kg, σ = 2.02 and 1.04 respectively) were excluded from

analysis as they could not be square-root transformed. There was no detectable difference in

mean lead levels between devils in the plantation (mean 1.05 ± 1.02 SD mg/kg) and

validation (mean 1.65 ± 4.61) studies (df = 40, t = 0.79, p = 0.43) (Figure 3). In the plantation

study, there was no evidence of differences in hair lead levels between sexes (df = 1, F =

0.96, p = 0.33) or among ages (df = 2, F = 0.23, p = 0.79) or an interaction between these two

variables (df = 2, F = 1.32, p = 0.28). The best fitting model predicting lead levels in devil

hair in this landscape was the null model, indicating no influence of age or sex on lead levels

in devils (Table 3). However, these results should be interpreted with caution due to low

sample sizes (Figure 4).

Table 2. Lead Concentrations in Hair Derived from Portable XRF and Predicted ICP-MS Levels from

Regression Analysis for Live Tasmanian Devils (*Sarcophilus Harrisii*), Sampled During a Trapping

Program in a Forestry Plantation Landscape (n = 61)

	Mean	Min	Max	SD
Predicted ICP-MS (mg/kg)	1.05	0.01	4.34	1.02
Portable XRF (mg/kg)	1.69	-1.39	6.37	1.53
Portable XRF uncertainty (mg/kg)	0.85	0.25	2.02	0.45

252   **Table 3.** Model Ranking Testing the Importance of Age and Sex for Predicting Lead Levels in

253   Tasmanian Devil (*Sarcophilus Harrisii*) Hair Samples from a Forestry Plantation Landscape. Df =

254   Degrees of Freedom, AIC = Akaike Information Criterion, ΔAIC = Akaike Information Criterion

255   Difference, AICwt = Akaike Information Criterion Weight

Model	df	AIC	ΔAIC	AICwt
Null	2	221.5	0	0.47
Sex	3	222.8	1.24	0.25
Age	4	223.5	1.96	0.18
Age + sex	5	224.7	3.19	0.10

258
259   **Figure. 3.** Comparison of lead concentrations in the hair of Tasmanian devils (*Sarcophilus harrisii*)

260   from the plantation study and the validation study, estimated using portable X-ray fluorescence.

261   Points show individual measurements, squares show means and error bars show SDs.

Figure 4. Portable X-ray fluorescence estimates of lead concentration in the hair of Tasmanian devils

(*Sarcophilus harrisii*) in a timber plantation landscape. Points show individual measurements, squares

show means and error bars show SDs.

DISCUSSION

We present the first investigation of portable XRF for measurement of elemental

concentrations in hair in a non-human species, and the first report of lead exposure in a

morphologically specialized mammalian scavenger as measured via hair lead levels. Using

regression analysis, we showed reasonable correlation (61%) between XRF and ICPS-MS

values. This suggests that portable XRF may be useful as an initial screening method for

mammalian hair to identify individuals with the largest burden of lead exposure versus a

typical exposure profile, though it is does not seem to be useful for precise predictions of lead

exposure levels.

Lead in hair has been found to correlate with lead levels in other tissues such as blood,

liver and bone of other species[61,62]. However, this is not always the case; for example, hair

lead levels did not correlate with liver lead levels in European hedgehogs (*Erinaceus*

europaeus)[62], or liver and kidney samples in caribou (*Rangifer tarandus caribou*)[63], though

the latter may have been influenced by low lead concentrations. Use of hair and other

keratinized matrices to measure lead exposure in mammals should therefore be treated with

caution, as small variations in hair lead may not reflect differences in lead concentrations in

internal tissues or blood[44]. Given these limitations, XRF may be most useful for

distinguishing individuals with substantially elevated hair lead levels from those individuals

with background lead levels, rather than precise measurement of hair lead concentrations.

Our results indicate portable XRF is likely to be sufficient for this type of 'screening' task.

Additionally, characterizing the relationship between lead levels in hair and internal

tissues/blood of mammalian scavengers such as devils would assist with interpreting lead

exposure from hair lead measurements.

Lead levels in devil hair, in both the validation study and the plantation landscape study,

were comparable to other published research on lead exposure in mammals (both wild and

domestic). The levels we found were lower than those found in the hair of domestic dogs

(*Canis familiaris*) used for hunting in Argentina (mean 2.37 ± 6.21 mg/kg)[64], similar to those

in hair from fruit bats in eastern Australia (medians for three species: 1.26/1.64/2.26 mg/kg)[38]

and roe deer in Italy (mean 1.39 ± 1.63 mg/kg)[40], and higher than those found in hair from

Malay civets sampled in Malaysian Borneo (mean 0.52 ± 1.08 mg/kg)[39]. Similar to several of

these studies[38–40], we found no evidence of variation in lead levels in devil hair among sexes

and age classes.

Without a standard classification of harmful lead levels in hair, even apparently low levels

of lead in devil hair may be cause for concern. Two devils, both in the validation study,

showed atypically high hair lead levels: 15.77 and 25.34 mg/kg, when no other devils in

either sample group had > 4.4 mg/kg (none > 2 mg/kg in the validation study). The source of

these unusually high levels is unknown. The generally low hair lead levels are consistent with

other studies showing low lead levels in the blood[25] and liver[18] of wild devils. Mean lead

levels were similar in devils in the plantation study and the validation study, providing little

support for our hypothesis that lead contamination would be higher in devils living in a

modified landscape subject to regular culling.

Grey wolves (*Canis lupus*), cougars (*Puma concolor*) and American black bears (*Ursus*

americanus) in the greater Yellowstone ecosystem likewise showed relatively low lead levels

in blood and tissue samples, indicating that some mammalian scavengers in some areas may

not be impacted by lead exposure even where hunting using lead-based ammunition occurs[34].

Brown bears did show elevated lead levels in this landscape, but lead levels did not increase

during the hunting season[34]. In contrast, brown bears sampled in northern Europe have shown

elevated lead levels in blood and milk[21], which in one study was linked to hunting activity[22],

and lead levels in the teeth of Canadian black bears increase markedly during hunting

seasons[23]. It is clear that further research is required to identify risk factors and determine

harmful lead exposure in free ranging mammalian species[20].

One limitation to the results of this study is that we did not wash the samples in this study

prior to analysis, as we aimed to validate portable XRF for use in the field, and washing the

fur of live-caught animals would be impractical. Samples may therefore have been

contaminated by external lead[41,65,66]. However, lead levels in devil hair in our study were

generally low despite the possibility of external contamination. Nonetheless, we could not

distinguish between endogenous lead incorporated into the hair from the blood, and

exogenous lead from external contamination, in our XRF measurements. We recommend

further research to corroborate our results by comparing ICP-MS and portable XRF values of

washed hair samples. Another limitation relates to operator safety. The handheld XRF emits a

small amount of radiation, which for standard use would be indistinguishable from

background radiation in terms of dosimetry[67]. For those planning on regular use of these

systems in the field, proper dosimetry should be used to monitor any potential exposure to the

user.

In conclusion, Portable XRF has the potential to be developed into a non-destructive, non-

invasive method of analyzing lead levels in mammal hair. We found sufficient correlation

between lead levels in devil hair derived via portable XRF and ICP-MS to identify

differences in lead concentrations over a coarse scale. Our preliminary data suggest portable

XRF for hair lead analysis has promise as a screening test to distinguish animals with marked

lead exposure from those with relatively low lead levels. While ICP-MS remains the more

accurate and precise method to quantify lead exposure levels in non-human mammalian hair,

portable XRF is a non-invasive alternative in situations where invasive sample collection or

destructive analysis is impractical or nonpreferred. These contexts may include when results

are required immediately, for use in the field, when working with threatened species, or for

archived museum specimens. Notably, any attempts to use portable XRF as a field-based tool

for measuring hair lead concentrations in live animals would need to be cognizant of health

and safety risks and regulations associated with the use of an ionizing radiation source[51].

While radiation exposure to the user can be minimized with training and proper use, it is a

consideration when choosing a method for lead monitoring. Future research could test the

relationship between ICP-MS and portable XRF hair lead measurements in other mammalian

scavenging species across a larger lead concentration gradient to refine this method.

ETHICS AND PERMITS

The live animal trapping and sample collection component of this study was approved by the

University of Tasmania Animal Ethics Committee (AEC), approval number 23211, and

conducted under a permit to take threatened fauna for scientific purposes (#TFA 21284) from

the Tasmanian Department of Natural Resources and Environment. Sample collection from

dead animals was performed under permit #TFA 21284. Samples were shipped interstate for

laboratory analysis under a Wildlife Export Permit (#TFA 21284) from the Tasmanian

Department of Natural Resources and Environment.

ACKNOWLEDGEMENTS

We acknowledge the Palawa people as the Traditional Owners of the lands on which this

research was conducted. For access to samples, we acknowledge the following people: Bill

Brown and Sarah Peck from the Department of Natural Resources and Environment

Tasmania, and David Hocking and Nicole Zehntner from the Tasmanian Museum and Art

Gallery (TMAG). For assistance with dissections, we thank Regi Broeren. For laboratory

analysis, we thank Mark Bannister from Edith Cowan University. For fauna licenses needed

to import samples to Victoria, we thank Sue Hadden and Roberta Campbell from the

Victorian Department of Environment, Land and Water. For facilitating XRF hire, we thank

Shellby Thomas-Brookes, Denise Guldemond and Alex Toy from Portable Analytical

Solutions. We thank Forico Pty Ltd for providing accommodation and access to land they

manage for fieldwork.

This work was supported by the Paddy Pallin Foundation [grant number H0029051], the

McKenzie Fellowship Program of the University of Melbourne [grant number 000376], the

Holsworth Wildlife Research Endowment [Grant No. H0027981], the Dr Eric Guiler

Tasmanian Devil Research Grant, and a matched funding scheme through Forests and Wood

Products Australia Limited, with funding provided by the Forest Practices Authority, Forico

Pty Ltd, SFM, Sustainable Timber Tasmania, Timberlands, Norske Skog, and Private Forests

Tasmania. Dr. Aaron Specht was partially funded by a training grant from the National

Institute of Occupational Safety and Health [grant number K01OH012528].

REFERENCES

(1) Arnemo, J. M.; Fuchs, B.; Sonne, C.; Stokke, S. Hunting with Lead Ammunition: A

One Health Perspective. In *Arctic One Health: Challenges for Northern Animals and*

People; Tryland, M., Ed.; Springer International Publishing: Switzerland, 2022; pp 439–

468. https://doi.org/10.1007/978-3-030-87853-5_21.

(2) Sonne, C.; Lam, S. S.; Kanstrup, N. Commentary: The Environmental Threats from

Lead Ammunition. *Eco-Environment & Health* **2023**, *2* (1), 16–17.

https://doi.org/10.1016/j.eehl.2023.02.001.

(3) Pain, D. J.; Mateo, R.; Green, R. E. Effects of Lead from Ammunition on Birds and

Other Wildlife: A Review and Update. *Ambio* **2019**, *48* (9), 935–953.

https://doi.org/10.1007/s13280-019-01159-0.

(4) Pain, D. J.; Cromie, R. L.; Newth, J.; Brown, M. J.; Crutcher, E.; Hardman, P.; Hurst,

L.; Mateo, R.; Meharg, A. A.; Moran, A. C.; Raab, A.; Taggart, M. A.; Green, R. E.

Potential Hazard to Human Health from Exposure to Fragments of Lead Bullets and

Shot in the Tissues of Game Animals. *PLOS ONE* **2010**, *5* (4), e10315.

https://doi.org/10.1371/journal.pone.0010315.

(5) Lambertucci, S. A.; Donázar, J. A.; Hiraldo, F. Poisoning People and Wildlife with

Lead Ammunition: Time to Stop. *Environmental Science & Technology* **2010**, *44* (20),

7759–7760. https://doi.org/10.1021/es102667c.

(6)   Hampton, J. O.; Laidlaw, M.; Buenz, E.; Arnemo, J. M. Heads in the Sand: Public

Health and Ecological Risks of Lead-Based Bullets for Wildlife Shooting in Australia.

Wildlife Research **2018**, *45* (4), 287–306. https://doi.org/10.1071/WR17180.

(7)   Plaza, P. I.; Lambertucci, S. A. What Do We Know about Lead Contamination in Wild

Vultures and Condors? A Review of Decades of Research. *Science of the Total*

Environment **2019**, *654*, 409–417. https://doi.org/10.1016/j.scitotenv.2018.11.099.

(8)   Finkelstein, M. E.; George, D.; Scherbinski, S.; Gwiazda, R.; Johnson, M.; Burnett, J.;

Brandt, J.; Lawrey, S.; Pessier, A. P.; Clark, M.; Wynne, J.; Grantham, J.; Smith, D. R.

Feather Lead Concentrations and 207Pb/206Pb Ratios Reveal Lead Exposure History of

California Condors (*Gymnogyps californianus*). *Environmental Science & Technology*

2010, *44* (7), 2639–2647. https://doi.org/10.1021/es903176w.

(9)   Herring, G.; Eagles-Smith, C. A.; Goodell, J.; Buck, J. A.; Willacker, J. J. Small-

Mammal Shooting as a Conduit for Lead Exposure in Avian Scavengers. *Environmental*

Science & Technology **2021**, *55* (18), 12272–12280.

https://doi.org/10.1021/acs.est.1c01041.

(10) Green, R. E.; Pain, D. J.; Krone, O. The Impact of Lead Poisoning from Ammunition

Sources on Raptor Populations in Europe. *Science of The Total Environment* **2022**, *823*,

154017. https://doi.org/10.1016/j.scitotenv.2022.154017.

(11) Slabe, V. A.; Anderson, J. T.; Millsap, B. A.; Cooper, J. L.; Harmata, A. R.; Restani,

M.; Crandall, R. H.; Bodenstein, B.; Bloom, P. H.; Booms, T.; Buchweitz, J.; Culver,

R.; Dickerson, K.; Domenech, R.; Dominguez-Villegas, E.; Driscoll, D.; Smith, B. W.;

Lockhart, M. J.; McRuer, D.; Miller, T. A.; Ortiz, P. A.; Rogers, K.; Schwarz, M.;

Turley, N.; Woodbridge, B.; Finkelstein, M. E.; Triana, C. A.; DeSorbo, C. R.; Katzner,

T. E. Demographic Implications of Lead Poisoning for Eagles across North America.

Science **2022**, *375* (6582), 779–782. https://doi.org/10.1126/science.abj3068.

(12) Ecke, F.; Singh, N. J.; Arnemo, J. M.; Bignert, A.; Helander, B.; Berglund, Å. M. M.;

Borg, H.; Bröjer, C.; Holm, K.; Lanzone, M.; Miller, T.; Nordström, Å.; Räikkönen, J.;

Rodushkin, I.; Ågren, E.; Hörnfeldt, B. Sublethal Lead Exposure Alters Movement

Behavior in Free-Ranging Golden Eagles. *Environmental Science & Technology* **2017**,

51 (10), 5729–5736. https://doi.org/10.1021/acs.est.6b06024.

(13) Finkelstein, M. E.; Doak, D. F.; George, D.; Burnett, J.; Brandt, J.; Church, M.;

Grantham, J.; Smith, D. R. Lead Poisoning and the Deceptive Recovery of the Critically

Endangered California Condor. *Proceedings of the National Academy of Sciences* **2012**,

109 (28), 11449–11454. https://doi.org/10.1073/pnas.1203141109.

(14) Hampton, J. O.; Specht, A. J.; Pay, J. M.; Pokras, M. A.; Bengsen, A. J. Portable X-Ray

Fluorescence for Bone Lead Measurements of Australian Eagles. *Science of The Total*

Environment **2021**, *789*, 147998. https://doi.org/10.1016/j.scitotenv.2021.147998.

(15) Lohr, M. T.; Hampton, J. O.; Cherriman, S.; Busetti, F.; Lohr, C. Completing a

Worldwide Picture: Preliminary Evidence of Lead Exposure in a Scavenging Bird from

Mainland Australia. *Science of the Total Environment* **2020**, *715*, 135913.

https://doi.org/10.1016/j.scitotenv.2019.135913.

(16) Pay, J. M.; Katzner, T. E.; Hawkins, C. E.; Kock, A. J.; Wiersma, J. M.; Brown, W. E.;

Mooney, N. J.; Cameron, E. Z. High Frequency of Lead Exposure in the Population of

an Endangered Australian Top Predator, the Tasmanian Wedge-Tailed Eagle (*Aquila*

Audax Fleayi). Environmental Toxicology and Chemistry **2021**, *40* (1), 219–230.

https://doi.org/10.1002/etc.4914.

(17) Hivert, L.; Clarke, J.; Peck, S.; Lawrence, C.; Brown, W.; Huxtable, S.; Schaap, D.;

Pemberton, D.; Grueber, C. High Blood Lead Concentrations in Captive Tasmanian

Devils (*Sarcophilus harrisii*): A Threat to the Conservation of the Species? *Australian*

Veterinary Journal **2018**, *96* (11), 442–449. https://doi.org/10.1111/avj.12753.

(18) Hutchinson, D. J.; Jones, E. M.; Pay, J. M.; Clarke, J. R.; Lohr, M. T.; Hampton, J. O. Evidence of Minimal Lead Exposure in a Scavenging Marsupial Species. *Australian Veterinary Journal* **2023**.

(19) Hampton, J. O.; Pay, J. M.; Katzner, T. E.; Arnemo, J. M.; Pokras, M. A.; Buenz, E.; Kanstrup, N.; Thomas, V. G.; Uhart, M.; Lambertucci, S. A.; Krone, O.; Singh, N. J.; Naidoo, V.; Ishizuka, M.; Saito, K.; Helander, B.; Green, R. E. Managing Macropods without Poisoning Ecosystems. *Ecological Management & Restoration* **2022**, *23* (2), 153–157. https://doi.org/10.1111/emr.12555.

(20) Chiverton, L.; Cromie, R.; Kock, R. European Mammal Exposure to Lead from Ammunition and Fishing Weight Sources. *Heliyon* **2022**, *8* (8), e10014. https://doi.org/10.1016/j.heliyon.2022.e10014.

(21) Boesen, A. H.; Thiel, A.; Fuchs, B.; Evans, A. L.; Bertelsen, M. F.; Rodushkin, I.; Arnemo, J. M. Assessment of the LeadCare® Plus for Use on Scandinavian Brown Bears (*Ursus arctos*). *Frontiers in Veterinary Science* **2019**, *6*.

(22) Brown, L.; Fuchs, B.; Arnemo, J. M.; Kindberg, J.; Rodushkin, I.; Zedrosser, A.; Pelletier, F. Lead Exposure in Brown Bears Is Linked to Environmental Levels and the Distribution of Moose Kills. *Science of the Total Environment* **2023**, 162099. https://doi.org/10.1016/j.scitotenv.2023.162099.

(23) Brown, L.; Rosabal, M.; Dussault, C.; Arnemo, J. M.; Fuchs, B.; Zedrosser, A.; Pelletier, F. Lead Exposure in American Black Bears Increases with Age and Big Game Harvest Density. *Environmental Pollution* **2022**, *315*, 120427. https://doi.org/10.1016/j.envpol.2022.120427.

(24) Fuchs, B.; Thiel, A.; Zedrosser, A.; Brown, L.; Hydeskov, H. B.; Rodushkin, I.; Evans, A. L.; Boesen, A. H.; Græsli, A. R.; Kindberg, J.; Arnemo, J. M. High Concentrations of Lead (Pb) in Blood and Milk of Free-Ranging Brown Bears (*Ursus arctos*) in

Scandinavia. *Environmental Pollution* **2021**, *287*, 117595.

https://doi.org/10.1016/j.envpol.2021.117595.

(25) Hivert, L. G.; Clarke, J. R.; Peck, S. J.; Lawrence, C.; Brown, W. E.; Huxtable, S. J.;

Schaap, D.; Pemberton, D.; Grueber, C. E. High Blood Lead Concentrations in Captive

Tasmanian Devils (*Sarcophilus harrisii*): A Threat to the Conservation of the Species?

Australian Veterinary Journal **2018**, *96* (11), 442–449.

https://doi.org/10.1111/avj.12753.

(26) Jones, M.; Archer, M.; Dickman, C. *Predators with Pouches: The Biology of*

Carnivorous Marsupials; CSIRO Publishing, 2003.

(27) Cunningham, C. X.; Comte, S.; McCallum, H.; Hamilton, D. G.; Hamede, R.; Storfer,

A.; Hollings, T.; Ruiz☐Aravena, M.; Kerlin, D. H.; Brook, B. W.; Hocking, G.; Jones,

M. E. Quantifying 25 Years of Disease-Caused Declines in Tasmanian Devil

Populations: Host Density Drives Spatial Pathogen Spread. *Ecology Letters* **2021**, *24*

(5), 958–969. https://doi.org/10.1111/ele.13703.

(28) Jones, E. M.; Koch, A. J.; Mann, D. F.; Hamede, R. K.; Jones, M. E. What Drives the

Abundance of Marsupial Carnivores in Production Forest Landscapes? *Forest Ecology*

and Management **2023**, *529*, 120745. https://doi.org/10.1016/j.foreco.2022.120745.

(29) Jones, M. E.; Jarman, P. J.; Lees, C. M.; Hesterman, H.; Hamede, R. K.; Mooney, N. J.;

Mann, D.; Pukk, C. E.; Bergfeld, J.; McCallum, H. Conservation Management of

Tasmanian Devils in the Context of an Emerging, Extinction-Threatening Disease:

Devil Facial Tumor Disease. *EcoHealth: Conservation Medicine: Human Health:*

Ecosystem Sustainability **2007**, *4* (3), 326. https://doi.org/10.1007/s10393-007-0120-6.

(30) Jones, M. E. Road Upgrade, Road Mortality and Remedial Measures: Impacts on a

Population of Eastern Quolls and Tasmanian Devils. *Wildl. Res.* **2000**, *27* (3), 289–296.

https://doi.org/10.1071/wr98069.

(31) Forest Practices Authority. *State of the Forests Tasmania 2017*; Forest Practices

Authority: Hobart, Tasmania, 2017.

(32) Le Mar, K.; McArthur, C. Relocating Radio-Collared Targeted Marsupials after a 1080-

Poisoning Operation. *Tasforests* **2000**, *12*, 155–160.

(33) Andersen, G. E.; Johnson, C. N.; Barmuta, L. A.; Jones, M. E. Dietary Partitioning of

Australia's Two Marsupial Hypercarnivores, the Tasmanian Devil and the Spotted-

Tailed Quoll, across Their Shared Distributional Range. *PLOS ONE* **2017**, *12* (11),

e0188529. https://doi.org/10.1371/journal.pone.0188529.

(34) Rogers, T. A.; Bedrosian, B.; Graham, J.; Foresman, K. R. Lead Exposure in Large

Carnivores in the Greater Yellowstone Ecosystem. *Journal of Wildlife Management*

2012, *76* (3), 575–582. https://doi.org/10.1002/jwmg.277.

(35) Kelly, J.; Carstensen, M.; Hildebrand, E. C.; Long, L. Assessing Lead Exposure in Free-

Ranging Gray Wolves (*Canis lupus)* in Minnesota, USA. *Journal of Wildlife Diseases*

2021, *57* (4), 917–921. https://doi.org/10.7589/JWD-D-20-00185.

(36) Jota Baptista, C.; Seixas, F.; Gonzalo-Orden, J. M.; Oliveira, P. A. Biomonitoring

Metals and Metalloids in Wild Mammals: Invasive versus Non-Invasive Sampling.

Environmental Science and Pollution Research **2022**, *29* (13), 18398–18407.

https://doi.org/10.1007/s11356-022-18658-5.

(37) Noël, M.; Spence, J.; Harris, K. A.; Robbins, C. T.; Fortin, J. K.; Ross, P. S.;

Christensen, J. R. Grizzly Bear Hair Reveals Toxic Exposure to Mercury through

Salmon Consumption. *Environmental Science & Technology* **2014**, *48* (13), 7560–7567.

https://doi.org/10.1021/es500631g.

(38) Sánchez, C. A.; Penrose, M. T.; Kessler, M. K.; Becker, D. J.; McKeown, A.;

Hannappel, M.; Boyd, V.; Camus, M. S.; Padgett-Stewart, T.; Hunt, B. E.; Graves, A.

F.; Peel, A. J.; Westcott, D. A.; Rainwater, T. R.; Chumchal, M. M.; Cobb, G. P.;

Altizer, S.; Plowright, R. K.; Boardman, W. S. J. Land Use, Season, and Parasitism

Predict Metal Concentrations in Australian Flying Fox Fur. *Science of The Total*

Environment **2022**, *841*, 156699. https://doi.org/10.1016/j.scitotenv.2022.156699.

(39) Evans, M. N.; Waller, S.; Müller, C. T.; Goossens, B.; Smith, J. A.; Bakar, M. S. A.;

Kille, P. The Price of Persistence: Assessing the Drivers and Health Implications of

Metal Levels in Indicator Carnivores Inhabiting an Agriculturally Fragmented

Landscape. *Environmental Research* **2022**, *207*, 112216.

https://doi.org/10.1016/j.envres.2021.112216.

(40) Draghi, S.; Agradi, S.; Riva, F.; Tarhan, D.; Bilgiç, B.; Dokuzeylül, B.; Ercan, A. M.;

Or, M. E.; Brecchia, G.; Vigo, D.; Arioli, F.; Di Cesare, F.; Curone, G. Roe Deer

(*Capreolus capreolus*) Hair as a Bioindicator for the Environmental Presence of Toxic

and Trace Elements. *Toxics* **2023**, *11* (1), 49. https://doi.org/10.3390/toxics11010049.

(41) Pozebon, D.; Scheffler, G. L.; Dressler, V. L. Elemental Hair Analysis: A Review of

Procedures and Applications. *Analytica Chimica Acta* **2017**, *992*, 1–23.

https://doi.org/10.1016/j.aca.2017.09.017.

(42) Azad, S.; McFadden, K.; Clark, J. D.; Wactor, T.; Jachowski, D. S. Applying Spatially

Explicit Capture–Recapture Models to Estimate Black Bear Density in South Carolina.

Wildlife Society Bulletin **2019**, *43* (3), 500–507. https://doi.org/10.1002/wsb.1007.

(43) Eltayeb, M. A. H.; Van Grieken, RenéE. Iron, Copper, Zinc and Lead in Hair from

Sudanese Populations of Different Age Groups. *Science of The Total Environment* **1990**,

95, 157–165. https://doi.org/10.1016/0048-9697(90)90061-X.

(44) Tehrani, M. W.; Galusha, A. L.; Kannan, A.; Parsons, P. J. Lead Uptake into Calcified

and Keratinized Compartments of Horns from a Convenience Sample of Lead-Dosed

Goats. *Journal of Toxicology and Environmental Health, Part A* **2021**, *84* (18), 729–

742. https://doi.org/10.1080/15287394.2021.1938766.

(45) Specht, A. J.; Kirchner, K. E.; Weisskopf, M. G.; Pokras, M. A. Lead Exposure

Biomarkers in the Common Loon. *Science of the Total Environment* **2019**, *647*, 639–

644. https://doi.org/10.1016/j.scitotenv.2018.08.043.

(46) Specht, A. J.; Parish, C. N.; Wallens, E. K.; Watson, R. T.; Nie, L. H.; Weisskopf, M. G.

Feasibility of a Portable X-Ray Fluorescence Device for Bone Lead Measurements of

Condor Bones. *Science of the Total Environment* **2018**, *615*, 398–403.

https://doi.org/10.1016/j.scitotenv.2017.09.123.

(47) Specht, A. J.; Dickerson, A. S.; Weisskopf, M. G. Comparison of Bone Lead Measured

via Portable X-Ray Fluorescence across and within Bones. *Environmental Research*

2019, *172*, 273–278. https://doi.org/10.1016/j.envres.2019.02.031.

(48) Khuder, A.; Bakir, M. A.; Hasan, R.; Mohammad, A. Determination of Nickel, Copper,

Zinc and Lead in Human Scalp Hair in Syrian Occupationally Exposed Workers by

Total Reflection X-Ray Fluorescence. *Environmental Monitoring and Assessment* **2008**,

143 (1), 67–74. https://doi.org/10.1007/s10661-007-9958-x.

(49) Kempson, I. M.; Skinner, W. M.; Kirkbride, K. P. Advanced Analysis of Metal

Distributions in Human Hair. *Environmental Science & Technology* **2006**, *40* (10),

3423–3428. https://doi.org/10.1021/es052158v.

(50) Śliwiński, M. G.; Latty, C. J.; Spaleta, K. J.; Taylor, R. J.; Severin, K. P. Rapid, Non-

Destructive Analysis of Calcium and Strontium in Eggshells by WD-XRF.

Chemosphere **2020**, *251*, 126253. https://doi.org/10.1016/j.chemosphere.2020.126253.

(51) Rouillon, M.; Kristensen, L. J.; Gore, D. B. Handheld X-Ray Fluorescence

Spectrometers: Radiation Exposure Risks of Matrix-Specific Measurement Scenarios.

Appl Spectrosc **2015**, *69* (7), 815–822. https://doi.org/10.1366/14-07809.

(52) Nzabanita, D.; Hampton, J. O.; Toop, S. D.; Bengsen, A. J.; Specht, A. J.; Flesch, J. S.;

Hufschmid, J.; Nugegoda, D. Expanding the Use of Portable XRF to Monitor Lead

Exposure in an Australian Duck Species Two Decades after a Ban on Lead Shot.

Science of The Total Environment **2023**, *869*, 161803.

https://doi.org/10.1016/j.scitotenv.2023.161803.

(53) Andersen, G. E.; McGregor, H. W.; Johnson, C. N.; Jones, M. E. Activity and Social

Interactions in a Wide-Ranging Specialist Scavenger, the Tasmanian Devil (*Sarcophilus*

harrisii), Revealed by Animal-Borne Video Collars. *PLOS ONE* **2020**, *15* (3),

e0230216. https://doi.org/10.1371/journal.pone.0230216.

(54) Hampton, J. O.; Cobb, M. L.; Toop, S. D.; Flesch, J. S.; Hyndman, T. H. Elevated Lead

Exposure in Australian Hunting Dogs during a Deer Hunting Season. *Environmental*

Pollution **2023**, *323*, 121317. https://doi.org/10.1016/j.envpol.2023.121317.

(55) Specht, A. J.; Kponee, K.; Nkpaa, K. W.; Balcom, P. H.; Weuve, J.; Nie, L. H.;

Weisskopf, M. G. Validation of X-Ray Fluorescence Measurements of Metals in

Toenail Clippings against Inductively Coupled Plasma Mass Spectrometry in a Nigerian

Population. *Physiological Measurement* **2018**, *39* (8), 085007.

https://doi.org/10.1088/1361-6579/aad947.

(56) Kim, R.; Aro, A.; Rotnitzky, A.; Amarasiriwardena, C.; Hu, H. K X-Ray Fluorescence

Measurements of Bone Lead Concentration: The Analysis of Low-Level Data. *Physics*

in Medicine and Biology **1995**, *40* (9), 1475–1485. https://doi.org/10.1088/0031-

9155/40/9/007.

(57) Specht, A. J.; Lin, Y.; Weisskopf, M.; Yan, C.; Hu, H.; Xu, J.; Nie, L. H. XRF-

Measured Bone Lead (Pb) as a Biomarker for Pb Exposure and Toxicity among

Children Diagnosed with Pb Poisoning. *Biomarkers* **2016**, *21* (4), 347–352.

https://doi.org/10.3109/1354750X.2016.1139183.

(58) Kuhn, M. Caret: Classification and Regression Training, 2022.

(59) R Core Team. R: A Language and Environment for Statistical Computing, 2021.

https://www.R-project.org/.

(60) Burnham, K. P.; Anderson, D. R. *Model Selection and Multimodel Inference: A*

Practical Information-Theoretic Approach; Springer New York: USA, 2002.

(61) Hernout, B. V.; McClean, C. J.; Arnold, K. E.; Walls, M.; Baxter, M.; Boxall, A. B. A.

Fur: A Non-Invasive Approach to Monitor Metal Exposure in Bats. *Chemosphere* **2016**,

147, 376–381. https://doi.org/10.1016/j.chemosphere.2015.12.104.

(62) Vermeulen, F.; D'Havé, H.; Mubiana, V. K.; Van den Brink, N. W.; Blust, R.; Bervoets,

L.; De Coen, W. Relevance of Hair and Spines of the European Hedgehog (*Erinaceus*

europaeus) as Biomonitoring Tissues for Arsenic and Metals in Relation to Blood.

Science of The Total Environment **2009**, *407* (5), 1775–1783.

https://doi.org/10.1016/j.scitotenv.2008.10.039.

(63) Jutha, N.; Jardine, C.; Schwantje, H.; Mosbacher, J.; Kinniburgh, D.; Kutz, S.

Evaluating the Use of Hair as a Non-Invasive Indicator of Trace Mineral Status in

Woodland Caribou (*Rangifer Tarandus Caribou*). *PLOS ONE* **2022**, *17* (6), e0269441.

https://doi.org/10.1371/journal.pone.0269441.

(64) Fernández, V.; Caselli, A.; Tammone, A.; Condorí, W. E.; Vanstreels, R. E. T.;

Delaloye, A.; Sosa, C.; Uhart, M. M. Lead Exposure in Dogs Fed Game Meat and Offal

from Culled Invasive Species in El Palmar National Park, Argentina. *Environmental*

Science and Pollution Research **2021**, *28* (33), 45486–45495.

https://doi.org/10.1007/s11356-021-13880-z.

(65) Barbosa, F.; Tanus, -Santos José Eduardo; Gerlach, R. F.; Parsons, P. J. A Critical

Review of Biomarkers Used for Monitoring Human Exposure to Lead: Advantages,

Limitations, and Future Needs. *Environmental Health Perspectives* **2005**, *113* (12),

1669–1674. https://doi.org/10.1289/ehp.7917.

(66) Borghesi, F.; Migani, F.; Andreotti, A.; Baccetti, N.; Bianchi, N.; Birke, M.; Dinelli, E.

Metals and Trace Elements in Feathers: A Geochemical Approach to Avoid

Misinterpretation of Analytical Responses. *Science of The Total Environment* **2016**,

544, 476–494. https://doi.org/10.1016/j.scitotenv.2015.11.115.

(67) Nie, H.; Sánchez, B. N.; Wilker, E.; Weisskopf, M.; Schwartz, J.; Sparrow, D.; Hu, H.

Bone Lead and Endogenous Exposure in an Environmentally Exposed Elderly

Population: The Normative Aging Study. *Journal of occupational and environmental*

medicine / American College of Occupational and Environmental Medicine **2009**, *51*

(7), 848–857. https://doi.org/10.1097/JOM.0b013e3181aa0106.