Portable X-ray fluorescence for identifying lead exposure in wild

mammals through hair analysis

Evie M. Jones [a], Andrew J. Bengsen [b,c], Aaron J. Specht [d], Amelia J. Koch [e], Rodrigo K.

Hamede [a], Menna E. Jones [a], Jordan O. Hampton [f,g]

[a] School of Natural Sciences, University of Tasmania, Private Bag 55, Hobart, TAS, 7001,

Australia

[b] Vertebrate Pest Research Unit, NSW Department of Primary Industries, 4 Marsden Park

Rd, Calala, New South Wales 2340, Australia

[c] Biosphere Environmental Consultants, Tamworth, NSW 2340, Australia

[d] Purdue University, 610 Purdue Mall, West Lafayette, Indiana 47907, United States

[e] Forest Practices Authority, 30 Patrick St, Hobart, TAS, 7001, Australia

[f] Faculty of Science, University of Melbourne, Parkville, Victoria 3052, Australia

[g] Harry Butler Institute, Murdoch University, 90 South Street, Murdoch, Western Australia

6150, Australia

* Corresponding author.

Email address: evie.jones@utas.edu.au

Abstract

Lead toxicity threatens scavenging wildlife globally. To allow inexpensive measurement of

long-term lead exposure from archived wildlife tissues, portable X-ray fluorescence (XRF)[1]

devices have been validated to measure lead concentrations in bone samples from bird

species. In this study, we evaluated the efficacy of portable XRF for hair lead measurement in

a specialist mammalian scavenger, the threatened Tasmanian devil (*Sarcophilus harrisii*). We

attempted to first validate the method by analyzing relatively large (~1.0 g) hair samples from

39 deceased animals initially with XRF, and then with inductively coupled plasma mass

spectrometry (ICP-MS) (validation study). We then measured lead concentrations via XRF

only in a further 61 hair samples taken from live devils trapped in a plantation forestry

landscape, where devils may scavenge on browsing marsupials that are regularly culled using

lead ammunition (plantation study). Portable XRF measures of hair lead levels were

positively correlated with ICP-MS values ($R^2 = 0.95$), although the relationship was weak at

low lead concentrations. In the validation study, 95% of sampled Tasmanian devils had

relatively low lead levels (≤ 3 mg/kg), but the remaining two animals showed considerably

elevated lead levels (> 15 mg/kg), suggesting that harmful long-term lead exposure can occur

in this species. In the plantation study, lead levels were comparable to those found in the hair

of mammals in other studies, though a higher proportion (11%) of devils showed levels > 3

mg/kg than in the validation study, suggesting devils may have higher exposure to lead in a

landscape subjected to regular culling of browsing mammals. Our results suggests that

portable XRF is a promising emerging technology, and after some refinement, may allow

non-invasive real-time measurement of lead concentrations in the hair of mammals.

Keywords: Tasmanian devil, marsupial, scavenger, ecotoxicology

1. Introduction

---

[1] XRF: X-ray fluorescence
[2] DFTD: Devil facial tumour disease

Lead is a widely studied heavy metal with many well-known toxic effects. Contemporary

research has implicated lead-based ammunition as a major source of anthropogenic

environmental lead around the world (Arnemo et al., 2022; Sonne et al., 2023), which is a

threat to scavenging wildlife (Plaza and Lambertucci, 2019). Many scavenging species,

particularly avian species, have been found with elevated lead exposure levels from lead-

based ammunition (Pain et al., 2019), which can cause both lethal and sublethal effects, such

as altered behavior (Ecke et al., 2017). Lead exposure in wildlife scavengers is a global issue,

but relatively little research has been conducted in Australia, beyond studies of lead exposure

in one avian species, the wedge-tailed eagle (*Aquila audax*) (Hampton et al., 2021; Lohr et

al., 2020; Pay et al., 2021). This is concerning, given that culling, hunting and commercial

harvesting of fauna is ubiquitous in Australia, and relies almost exclusively on lead-based

ammunition (Hampton et al., 2022, 2018).

In contrast to avian species, relatively few studies have assessed the effects of lead

pollution from ammunition in wild mammalian scavengers (Chiverton et al., 2022). The

exception to this trend has been studies of bears (*Ursus* spp.) in the Northern Hemisphere

(Boesen et al., 2019; Brown et al., 2023, 2022; Fuchs et al., 2021). One recent study from the

Southern Hemisphere demonstrated harmful lead levels in an Australian marsupial species,

the Tasmanian devil (*Sarcophilus harrisii*), when fed whole shot carcasses in captivity

(Hivert et al., 2018). The Tasmanian devil is the largest marsupial carnivore in the world and

a morphologically specialized scavenger (Jones et al., 2003). They are endangered due to

devil facial tumor disease (DFTD)[2], a transmissible cancer which has caused population

declines of over 80% since 1996 (Cunningham et al., 2021). Due to their endangered status,

the species is the focus of intense conservation efforts, including proactive attempts to

identify threatening processes that may prevent or delay their recovery, such as threats from

anthropogenic activities (Jones et al., 2023, 2007). The aforementioned study of Hivert et al.

(2018) also measured lead exposure in wild devils but restricted their investigations to blood

lead concentrations, reflecting only short-term exposure (Pain et al., 2019). Most of these

wild devils had blood lead concentrations within the normal range, though 6% showed higher

concentrations of lead (0.21 to 0.65 µmol/L).

The low population size of Tasmanian devils and their reliance on scavenging makes the

potential impact of lead contamination from eating culled carcasses particularly concerning,

especially to local devil populations in modified landscapes where culling of browsing

mammals is common (Hutchinson et al., unpublished results). For instance, in Tasmanian

timber plantations (henceforth 'plantations'), culling of the marsupial browsers Bennett's

wallaby (*Notamacropus rufogriseus*), rufous-bellied pademelon (*Thylogale billardierii*) and

brushtail possum (*Trichosurus vulpecula*), which constitute primary prey for devils

(Andersen et al., 2017), is regularly conducted using lead-based ammunition prior to planting

in coupes to protect seedlings (FPA, 2017).

To our knowledge, all past studies of lead exposure in scavenging mammals have utilized

invasive sampling via collection of blood (Boesen et al., 2019; Brown et al., 2023; Fuchs et

al., 2021; Hivert et al., 2018; Rogers et al., 2012), teeth (Brown et al., 2022) or liver (Kelly et

al., 2021). Recent reviews have highlighted the value of non-invasive sampling of heavy

metal exposure in wild mammals (Jota Baptista et al., 2022). One non-invasive approach to

investigating heavy metal exposure in mammals is through hair analysis, a technique used by

some recent studies on non-scavenging mammal species such as fruit bats (*Pteropus* spp.)

(Sánchez et al., 2022), Malay civets (*Viverra tangalunga*) (Evans et al., 2022) and roe deer

(*Capreolus capreolus*) (Draghi et al., 2023). Collecting hair samples from wild animals can

be done remotely using hair snares (Azad et al., 2019) or through minimal handling of

trapped wild animals. Invasive sampling studies typically require *ex situ* analysis via

inductively coupled plasma mass spectrometry (ICP-MS) in a laboratory to analyze lead

concentrations, which is expensive and time-consuming (Hampton et al., 2021).

X-ray fluorescence (XRF) is another option for analyzing lead exposure levels in hair. The

XRF device analyzes the elemental content of samples through the fluorescence produced by

X-rays from a low energy x-ray tube (50 kV). XRF has long been used in human medicine

(Specht et al., 2019a) and has a wide range of uses in wildlife research, such as examining

elemental characteristics of tissues such as antlers (Singh et al., 2022), tracking monotreme

species in the illicit wildlife trade (Brandis et al., 2018), and identifying trafficked animal

parts such as ivory (McLellan et al., 2022). Portable XRF units have recently become

commercially available. As they are handheld and battery-powered, these units are relatively

inexpensive, quick and practical for use in the field, with most functions taking only three

minutes to measure samples.

Advantages of this method over ICP-MS include lower cost and non-destructive analysis

of samples (Hampton et al., 2021). Portable XRF has been validated for measurement of lead

in bones of several avian species, showing strong correlations with ICP-MS values, such as

wedge-tailed eagles (Hampton et al., 2021), California condors (*Gymnogyps californianus*)

(Specht et al., 2018b), Pacific black duck (*Anas superciliosa*) (Nzabanita et al., 2023), and

common loons (*Gavia immer*) (Specht et al., 2019b). However, portable XRF has been

validated for measurement of lead in the hair of few mammalian species other than humans

(Buddhachat et al., 2016).

Here, we investigated hair lead concentrations in wild Tasmanian devils in two stages. The

first stage was a validation study (comparing portable XRF to traditional ICP-MS methods)

and the second stage was a study of lead exposure in a forestry plantation landscape. Our

hypotheses were: 1) portable XRF measures of hair lead levels in Tasmanian devils are

similar enough to ICP-MS results that they can be used as a proxy, and 2) portable XRF can

rapidly identify mammals with markedly elevated lead exposure levels through non-

destructive and non-invasive analysis of hair samples.

2. Materials and methods

2.1 Validation study

Hair from Tasmanian devils (Fig. 1) (n = 39) was collected from archived collections held

in the Tasmanian Museum and Art Gallery (TMAG) and the University of Tasmania

(UTAS). All samples were collected between April 2017 and February 2022 from across

Tasmania but concentrated on urban areas surrounding Hobart (Hutchinson et al.,

unpublished results). Approximately 1 g of hair was shaved from whole frozen devil

carcasses. Hair samples were analyzed via both XRF and ICP-MS analysis, with the same

sample used for each method to minimize the potential for confounding effects of variation in

lead concentrations between hair from different parts of the body (Specht et al., 2019a).

2.2 Forestry plantation landscape study

Relatively small (~0.1 g) hair samples were collected from 61 live wild devils living

within Surrey Hills, a forestry plantation landscape in northwest Tasmania (-41.420S,

145.814E). Surrey Hills is an intensively managed forestry landscape dominated by

Eucalyptus nitens plantations (Jones et al., 2023). Culling, using lead-based ammunition, is

regularly undertaken at the site prior to planting in coupes to protect seedlings from browsing

by three marsupial species: Bennett's wallaby, rufous-bellied pademelon, and brushtail

possum. All three species are common prey of devils (Andersen et al., 2017), and there is

evidence that culled browsers are scavenged by devils (Le Mar and McArthur, 2000).

Between February 2021 and March 2022, we captured devils in custom-made 80 cm by 30

cm PVC pipe traps, shaved a 1 cm^2 patch of fur from their right shoulders, and recorded the

age (from tooth wear) and sex of the animal. These samples were analyzed using XRF only,

as they were not large enough for ICP-MS methods.

2.3 XRF measurements

For XRF analysis of hair samples, we used similar methods as in past studies of lead in

birds (Hampton et al., 2021; Specht et al., 2018b) and humans (Specht et al., 2018a). We used

the Niton XL3t GOLDD+ (Thermo Fisher Scientific, Omaha, USA) portable XRF device,

the same unit as used as in previous related studies (Bhatia et al., 2021; Hampton et al., 2021;

Specht et al., 2018a). The device has a thermoelectric-cooled silicon drift detector (25 mm^2

area, 1 mm thickness). We used the device's customization option to optimize X-ray tube

settings and filtration: 50 kV, 40 µA, and a filter combination of silver and iron (Hampton et

al., 2021; Specht et al., 2019b, 2018b). We used a 3-minute (180 second) read-time for each

sample.

The XRF unit was placed in a stand with hair samples placed on top of it, to ensure that

samples were as close as possible to the beam detector (Hampton et al., 2021; Specht et al.,

2018b). To analyze the measurements, we used the standard procedure from past studies

(Hampton et al., 2021; Specht et al., 2018a), using peak fitting through MATLAB to identify

the net counts of the lead peak and the corresponding Compton scattering peak from the

silver X-ray tube anode. We used the silver Compton scattering peak to normalize the

differences in geometry and mass among samples of different sizes (Hampton et al., 2021;

Specht et al., 2018a).

2.4 ICP-MS measurements

The same hair samples (validation study samples only) were then sent to a commercial

laboratory at Edith Cowan University Analytical Facility (Joondalup, WA, Australia) to

determine their lead concentrations via inductively coupled mass spectrometry (ICP-MS).

Hair samples were freeze-dried and then underwent acid digestion. The hair samples were

homogenized and processed as per Pay et al. (2021): specifically, a 0.4 g aliquot was

transferred into ICP-grade polytetrafluoroethylene vessels which contained 5 mL of 68%

trace analysis-grade nitric acid (Primar Plus™; Thermo☐Fisher Scientific, Omaha, USA), 0.5

mL of 34–37% trace metal-grade hydrochloric acid (Fisher Scientific), and 3 mL of 30%

hydrogen peroxide (Emsure ISO®; Merck, Rahway, USA). The samples were then digested

for 15 minutes in a microwave digestion system at 150°C. Samples were subsequently diluted

with reverse osmosis deionized water (Milli☐Q®; Merck) to a total volume of 50 mL before

being transferred to polypropylene tubes. For ICP–MS analysis, an iCAP™ Q

(Thermo☐Fisher Scientific) unit was coupled to a CETAC™ ASX☐520 AutoSampler

(Thermo☐Fisher Scientific). Qtegra™ (Thermo☐Fisher Scientific, Omaha, USA) software

was used for data acquisition and element quantitation.

A standard curve was created with concentration ranges of iCAP™ Q element standards

(Thermo☐Fisher Scientific) and ICP-MS☐68A solutions (High-Purity Standards, North

Charleston, USA) to calibrate the instruments prior to analysis. We used Bone Ash Standard

Reference Material 1400 (National Institute of Standards and Technology, Gaithersburg,

USA) as a positive control Certified Reference Material (CRM). The CRM was digested

twice, and two ICP-MS readings were made on each digestion. CRM ICP–MS readings had

an average accuracy of 100.2%. We re-analyzed every fifth or sixth sample to create a

duplicate read, with an average relative standard deviation (RSD) of 2.2%. We also

conducted duplicate blind sample digestions on five randomly selected samples (average

RSD = 2.0%). The lead concentrations were measured in mg/kg of dry weight (equivalent to

ppm). Limit of detection (LOD) was 0.0015 mg/kg and limit of quantification (LOQ) was

0.005 mg/kg.

2.5 Statistical analysis

We identified the uncertainty values for XRF measurements as per past studies using the

same equipment (Specht et al., 2019b). We calculated the uncertainty (σ) of each

measurement using the equation:

Equation 1:

$$\sigma = \frac{c \times \sqrt{\frac{BKG}{t}}}{NET}$$

where *c* is concentration, *BKG* is the background counts as estimated by our fitting, *t* is

measurement time, and *NET* is net lead counts from a Gaussian function (Zhang et al., 2021).

We retained the negative values for lead in hair because they represent a point estimate of

hair lead concentration incorporating uncertainty (Kim et al., 1995; Specht et al., 2016).

Linear regression was used to estimate the relationship between ICP-MS-derived and

XRF-derived lead concentrations for sampled tested via both methods (n = 39) with the

following equation:

Equation 2:

$$ICP_i = \beta_0 + \beta_1(XRF_i) + \epsilon_i$$

where β_0 is the intercept, β_1 is the slope of the relationship, and ϵ_i is the residual error. The

predictive competence of the regression model was evaluated with 10-fold cross-validation

repeated three times. Modelling was conducted using the caret package (Kuhn, 2022) in the R

statistical environment (RStudio Team, 2023).

For each estimation method, we calculated the proportion of samples classified into 'high'

or 'low' levels of lead exposure. Such classification can be problematic for hair lead

concentrations (Fernández et al., 2021), because there is no standard classification for

harmful lead concentrations in hair, as there is for blood, liver and bone (Pain et al., 2019).

As such, we defined 'low' lead concentration as ≤ 3 mg/kg where most of our validation data

was concentrated, and 'high' lead concentrations as > 3 mg/kg. We used the same

classification for both XRF values and predicted ICP-MS values and for the plantation

landscape samples. Additionally, we constructed linear models with the predictors of sex and

age (juvenile < 1 year, subadult 1–2 years, adult > 2 years) to look for any influence of these

parameters on predicted ICP-MS lead levels in devils, using Akaike's Information Criterion

(AIC) values to select the best fitting model (Burnham and Anderson, 2002).

3. Results

3.1 Validation of XRF for devil hair samples

We found a positive relationship between lead concentrations in the hair of Tasmanian

devils estimated using portable XRF and ICP-MS (n = 39) (Fig. 2). The regression model

showed a reasonable correlation between portable XRF and ICP-MS values, explaining 95%

of the variability in the data over the range of values observed. The theoretical ICP-MS

estimate of hair lead corresponding with an XRF estimate of 0 mg/kg was -0.564 mg/kg and

the expected ICP-MS lead concentration increased by 0.947 units (SE = 0.034) for every unit

increase estimated by XRF (R^2 = 0.95, t_{37} = 27.825, p < 0.001; RMSE = 1.4). ICP-MS

estimates could therefore be predicted from portable XRF estimates across the range of

sample data (n = 61) using the equation:

Equation 3:

$$ICP_i = \beta_0 + \beta_1 XRF_i + \epsilon_i$$

where β_0 = -0.564 and β_1 = 0.947.

The two highest values exerted a high degree of influence over the form of the regression.

We therefore ran a second model excluding these values. This model showed limited,

although still significant, correlation between XRF and ICP-MS values, explaining only 13%

of the variability in the data ($R^2 = 0.13$, $t_{37} = 2.323$, $p = 0.03$).

Most of the samples (94.87% as measured via either method) showed lead concentrations

of < 3 mg/kg (Table 1) and the relationship between XRF and ICP-MS was weak at these

lower values. However, the two points with the highest lead levels (> 15 mg/kg) measured

consistently high with both XRF and ICP-MS (Fig. 2). XRF estimates tended to be higher

than ICP-MS: the mean lead concentration was 2.33 mg/kg when measured via XRF versus

1.65 mg/kg when measured via ICP-MS. However, the frequency of 'high' lead exposure (>

3 mg/kg) was identical (5.13%) for both methods (Table 1). Given the dissimilarity between

XRF and ICP-MS values, we feel that it is important to transform XRF values into ICP-MS

equivalents prior to analysis, using equation 3.

3.2 Forestry plantation landscape study

Table 2 shows portable XRF hair lead measurements and uncertainties from devils in the

plantation landscape. These data were converted to ICP-MS values using equation 3. Even

using this transformed data, 11% of devils displayed 'high' (> 3 mg/kg) lead levels,

substantially higher than the proportion of devils displaying 'high' levels in the validation

analysis (5%, Table 1). The best fitting model predicting lead levels in devil hair in this

landscape was the null model, indicating no influence of age or sex on lead levels in devils

(Table 3).

4. Discussion

We present the first investigation of portable XRF for measurement of lead levels in hair

in a non-human species, and the first report of lead exposure in a wild-living specialized

mammalian scavenger as measured via hair lead levels. Using regression analysis, we were

able to characterize the relationship between XRF and ICPS-MS values at a coarse scale over

a wide range of data. However, the relationship was highly variable at lower values, so XRF

is not useful for precise predictions of low lead exposure levels in devil hair. We suggest that

portable XRF could be used as an initial screening method for mammalian hair to identify

individuals with the largest burden of lead exposure versus a typical exposure profile.

Lead levels in devil hair, in both the validation study (mean 1.65 ± 4.61 SD) and the

plantation landscape study (mean 1.05 ± 1.02 mg/kg), were comparable to other published

research on lead exposure in mammals (both wild and domestic). The levels we found were

lower than those found in the hair of domestic dogs (*Canis familiaris*) used for hunting in

Argentina (mean 2.37 ± 6.21 mg/kg) (Fernández et al., 2021), similar to those in hair from

fruit bats in eastern Australia (medians for three species: 1.26/1.64/2.26 mg/kg) (Sánchez et

al., 2022) and roe deer in Italy (mean 1.39 ± 1.63 mg/kg) (Draghi et al., 2023), and higher

than those found in hair from Malay civets sampled in Malaysian Borneo (mean 0.52 ± 1.08

mg/kg) (Evans et al., 2022). Similar to several of these studies (Draghi et al., 2023; Evans et

al., 2022; Sánchez et al., 2022), there was no evidence of variations in lead levels in devil hair

among sexes and age classes.

Without a standard classification of harmful lead levels in hair, even apparently low levels

of lead in devil hair may be cause for concern. Two devils, both in the validation study,

showed atypically high hair lead levels: 15.77 and 25.34 mg/kg, when no other devils in

either sample group had > 4.5 mg/kg (none > 2 mg/kg in the validation study). The source of

these unusually high levels is unknown. The generally low hair lead levels are consistent with

other studies showing low lead levels in the blood (Hivert et al., 2018) and liver (Hutchinson

et al., unpublished results) of wild devils. Grey wolves (*Canis lupus*), cougars (*Puma*

concolor) and American black bears (*Ursus americanus*) in the greater Yellowstone

ecosystem likewise showed relatively low lead levels in blood and tissue samples, indicating

that some mammalian scavengers in some areas may not be impacted by lead exposure even

where hunting using lead-based ammunition occurs (Rogers et al., 2012). Brown bears

(*Ursus arctos*) did show elevated lead levels in this landscape, but lead levels did not increase

during the hunting season (Rogers et al., 2012). In contrast, brown bears sampled in northern

Europe have shown markedly elevated lead levels in blood and milk (Boesen et al., 2019),

and lead levels in the teeth of Canadian black bears increase markedly during hunting seasons

(Brown et al., 2022). It is clear that further research is required into risk factors for harmful

lead exposure in free ranging mammalian species (Chiverton et al., 2022).

The proportion of devils displaying 'high' lead levels (> 3 mg/kg) was more than twice as

high in the plantation landscape (11%) than in the validation study (5%), the latter of which

used samples from throughout Tasmania but concentrated in urban areas near Hobart

(Hutchinson et al., unpublished results). This is a high enough discrepancy to suggest a real

difference between these values even when accounting for limited accuracy of the XRF

analysis. This result indicates support for the hypothesis that devils living in modified

landscapes subjected to regular culling may be at elevated risk of lead contamination

compared to devil populations in unmodified habitats (or at least those from urban areas with

minimal shooting activity), as has been found in some other mammalian non-specialist

scavengers in areas where hunting occurs (Brown et al., 2022; Kelly et al., 2021). Notably,

our study results came from only two sample populations with different locations and time

periods, so the discrepancy could be due to other factors that we could not incorporate into

the analysis. Further investigation to confirm causes is needed. Regardless, these results

highlight the importance of investigating lead levels in local scavenger populations in areas

with intensive hunting or culling operations, as the level of risk may not be identified by

studies of broader populations.

5. Conclusions

XRF provides a cost effective and field-friendly method of analyzing lead levels in mammal hair. Though the relationship was variable, we found sufficient correlation between lead levels in devil hair derived via XRF and ICP-MS to distinguish between high and low lead concentrations. While ICP-MS remains the best approach to quantifying low lead exposure levels in non-human mammalian hair with a high degree of accuracy, XRF is a viable non-invasive alternative in situations where ICP-MS is impractical, such as measurements in the field and where non-destructive sampling is preferred, as with archived museum specimens. At the least, our preliminary data suggest the method has promise as a screening test to distinguish animals with marked lead exposure from those animals with background lead levels. Our data suggest that investigating devil lead exposure levels in landscapes where culling is regularly conducted would be worthwhile to identify risks arising from use of lead-based ammunition.

Funding

This work was supported by the Paddy Pallin Foundation [grant number H0029051], the McKenzie Fellowship Program of the University of Melbourne [grant number 000376], the Holsworth Wildlife Research Endowment [Grant No. H0027981], the Dr Eric Guiler Tasmanian Devil Research Grant, and a matched funding scheme through Forests and Wood Products Australia Limited, with funding provided by Forico Pty Ltd, SFM, Sustainable Timber Tasmania, Timberlands, Norske Skog, and Private Forests Tasmania. Dr. Aaron Specht was partially funded by a training grant from the National Institute of Occupational Safety and Health [grant number K01OH012528].

Acknowledgements

We acknowledge the Palawa people as the Traditional Owners of the lands on which this

research was conducted. The live animal trapping and sample collection component of this

study was approved by the University of Tasmania Animal Ethics Committee (AEC),

approval number 23211, and conducted under a permit to take threatened fauna for scientific

purposes (#TFA 21284) from the Tasmanian Department of Natural Resources and

Environment. Sample collection from dead animals was performed under permit #TFA

21284. Samples were shipped interstate for laboratory analysis under a Wildlife Export

Permit (#TFA 21284) from the Tasmanian Department of Natural Resources and

Environment. For access to samples, we acknowledge the following people: Bill Brown and

Sarah Peck from the Department of Natural Resources and Environment Tasmania, and

David Hocking and Nicole Zehntner from the Tasmanian Museum and Art Gallery (TMAG).

For assistance with dissections, we thank Regi Broeren. For laboratory analysis, we thank

Mark Bannister from Edith Cowan University. For fauna licenses needed to import samples

to Victoria, we thank Sue Hadden and Roberta Campbell from the Victorian Department of

Environment, Land and Water. For facilitating XRF hire, we thank Shellby Thomas-Brookes,

Denise Guldemond and Alex Toy from Portable Analytical Solutions. We thank Forico Pty

Ltd for providing accommodation and access to land they manage for fieldwork.

References

Andersen, G.E., Johnson, C.N., Barmuta, L.A., Jones, M.E., 2017. Dietary partitioning of

Australia's two marsupial hypercarnivores, the Tasmanian devil and the spotted-tailed

quoll, across their shared distributional range. PLoS ONE 12, e0188529.

https://doi.org/10.1371/journal.pone.0188529

Arnemo, J.M., Fuchs, B., Sonne, C., Stokke, S., 2022. Hunting with Lead Ammunition: A

One Health Perspective, in: Tryland, M. (Ed.), Arctic One Health: Challenges for

Northern Animals and People. Springer International Publishing, Switzerland, pp.

439–468. https://doi.org/10.1007/978-3-030-87853-5_21

Azad, S., McFadden, K., Clark, J.D., Wactor, T., Jachowski, D.S., 2019. Applying spatially

explicit capture–recapture models to estimate black bear density in South Carolina.

Wildlife Society Bulletin 43, 500–507. https://doi.org/10.1002/wsb.1007

Bhatia, M., Specht, A.J., Ramya, V., Sulaiman, D., Konda, M., Balcom, P., Sunderland,

E.M., Qureshi, A., 2021. Portable X-ray fluorescence as a rapid determination tool to

detect parts per million levels of Ni, Zn, As, Se, and Pb in human toenails: A south

India case study. Environmental Science & Technology 55, 13113–13121.

https://doi.org/10.1021/acs.est.1c00937

Boesen, A.H., Thiel, A., Fuchs, B., Evans, A.L., Bertelsen, M.F., Rodushkin, I., Arnemo,

J.M., 2019. Assessment of the LeadCare® Plus for use on Scandinavian brown bears

(*Ursus arctos*). Frontiers in Veterinary Science 6.

Brandis, K.J., Meagher, P.J.B., Tong, L.J., Shaw, M., Mazumder, D., Gadd, P., Ramp, D.,

2018. Novel detection of provenance in the illegal wildlife trade using elemental data.

Scientific Reports 8, 15380. https://doi.org/10.1038/s41598-018-33786-0

Brown, L., Fuchs, B., Arnemo, J.M., Kindberg, J., Rodushkin, I., Zedrosser, A., Pelletier, F.,

2023. Lead exposure in brown bears is linked to environmental levels and the

distribution of moose kills. Science of the Total Environment 162099.

https://doi.org/10.1016/j.scitotenv.2023.162099

Brown, L., Rosabal, M., Dussault, C., Arnemo, J.M., Fuchs, B., Zedrosser, A., Pelletier, F.,

2022. Lead exposure in American black bears increases with age and big game

harvest density. Environmental Pollution 315, 120427.

https://doi.org/10.1016/j.envpol.2022.120427

Buddhachat, K., Klinhom, S., Siengdee, P., Brown, J.L., Nomsiri, R., Kaewmong, P.,

390   Thitaram, C., Mahakkanukrauh, P., Nganvongpanit, K., 2016. Elemental analysis of

391   bone, teeth, horn and antler in different animal species using non-invasive handheld

392   X-ray fluorescence. PLOS ONE 11, e0155458.

393   https://doi.org/10.1371/journal.pone.0155458

394 Burnham, K.P., Anderson, D.R., 2002. Model selection and multimodel inference: a practical

395   information-theoretic approach. Springer New York, USA.

396 Chiverton, L., Cromie, R., Kock, R., 2022. European mammal exposure to lead from

397   ammunition and fishing weight sources. Heliyon 8, e10014.

398   https://doi.org/10.1016/j.heliyon.2022.e10014

399 Cunningham, C.X., Comte, S., McCallum, H., Hamilton, D.G., Hamede, R., Storfer, A.,

400   Hollings, T., Ruiz☐Aravena, M., Kerlin, D.H., Brook, B.W., Hocking, G., Jones,

401   M.E., 2021. Quantifying 25 years of disease-caused declines in Tasmanian devil

402   populations: host density drives spatial pathogen spread. Ecology Letters 24, 958–

403   969. https://doi.org/10.1111/ele.13703

404 Draghi, S., Agradi, S., Riva, F., Tarhan, D., Bilgiç, B., Dokuzeylül, B., Ercan, A.M., Or,

405   M.E., Brecchia, G., Vigo, D., Arioli, F., Di Cesare, F., Curone, G., 2023. Roe deer

406   (*Capreolus capreolus*) hair as a bioindicator for the environmental presence of toxic

407   and trace elements. Toxics 11, 49. https://doi.org/10.3390/toxics11010049

408 Evans, M.N., Waller, S., Müller, C.T., Goossens, B., Smith, J.A., Bakar, M.S.A., Kille, P.,

409   2022. The price of persistence: Assessing the drivers and health implications of metal

410   levels in indicator carnivores inhabiting an agriculturally fragmented landscape.

411   Environmental Research 207, 112216. https://doi.org/10.1016/j.envres.2021.112216

412 Fernández, V., Caselli, A., Tammone, A., Condorí, W.E., Vanstreels, R.E.T., Delaloye, A.,

413   Sosa, C., Uhart, M.M., 2021. Lead exposure in dogs fed game meat and offal from

culled invasive species in El Palmar National Park, Argentina. Environmental Science

and Pollution Research International 28, 45486–45495.

https://doi.org/10.1007/s11356-021-13880-z

FPA, 2017. State of the forests Tasmania 2017. Forest Practices Authority, Hobart, Tasmania.

Fuchs, B., Thiel, A., Zedrosser, A., Brown, L., Hydeskov, H.B., Rodushkin, I., Evans, A.L.,

Boesen, A.H., Græsli, A.R., Kindberg, J., Arnemo, J.M., 2021. High concentrations of

lead (Pb) in blood and milk of free-ranging brown bears (*Ursus arctos*) in

Scandinavia. Environmental Pollution 287, 117595.

https://doi.org/10.1016/j.envpol.2021.117595

Hampton, J.O., Laidlaw, M., Buenz, E., Arnemo, J.M., 2018. Heads in the sand: public health

and ecological risks of lead-based bullets for wildlife shooting in Australia. Wildlife

Research 45, 287–306. https://doi.org/10.1071/WR17180

Hampton, J.O., Pay, J.M., Katzner, T.E., Arnemo, J.M., Pokras, M.A., Buenz, E., Kanstrup,

N., Thomas, V.G., Uhart, M., Lambertucci, S.A., Krone, O., Singh, N.J., Naidoo, V.,

Ishizuka, M., Saito, K., Helander, B., Green, R.E., 2022. Managing macropods

without poisoning ecosystems. Ecological Management & Restoration 23, 153–157.

https://doi.org/10.1111/emr.12555

Hampton, J.O., Specht, A.J., Pay, J.M., Pokras, M.A., Bengsen, A.J., 2021. Portable X-ray

fluorescence for bone lead measurements of Australian eagles. Science of The Total

Environment 789, 147998. https://doi.org/10.1016/j.scitotenv.2021.147998

Hivert, L.G., Clarke, J.R., Peck, S.J., Lawrence, C., Brown, W.E., Huxtable, S.J., Schaap, D.,

Pemberton, D., Grueber, C.E., 2018. High blood lead concentrations in captive

Tasmanian devils (*Sarcophilus harrisii*): a threat to the conservation of the species?

Australian Veterinary Journal 96, 442–449. https://doi.org/10.1111/avj.12753

Hutchinson, D.J., Jones, E.M., Pay, J.M., Clarke, J.R., Hufschmid, J., Lohr, M.T., Hampton,

J.O., in review. Evidence of minimal lead exposure in a scavenging marsupial species.

Australian Veterinary Journal.

Jones, E.M., Koch, A.J., Mann, D.F., Hamede, R.K., Jones, M.E., 2023. What drives the

abundance of marsupial carnivores in production forest landscapes? Forest Ecology

and Management 529, 120745. https://doi.org/10.1016/j.foreco.2022.120745

Jones, M., Archer, M., Dickman, C., 2003. Predators with pouches: the biology of

carnivorous marsupials. CSIRO Publishing.

Jones, M.E., Jarman, P.J., Lees, C.M., Hesterman, H., Hamede, R.K., Mooney, N.J., Mann,

D., Pukk, C.E., Bergfeld, J., McCallum, H., 2007. Conservation management of

Tasmanian devils in the context of an emerging, extinction-threatening disease: Devil

Facial Tumor Disease. EcoHealth: Conservation Medicine: Human Health:

Ecosystem Sustainability 4, 326. https://doi.org/10.1007/s10393-007-0120-6

Jota Baptista, C., Seixas, F., Gonzalo-Orden, J.M., Oliveira, P.A., 2022. Biomonitoring

metals and metalloids in wild mammals: invasive versus non-invasive sampling.

Environmental Science and Pollution Research International 29, 18398–18407.

https://doi.org/10.1007/s11356-022-18658-5

Kelly, J., Carstensen, M., Hildebrand, E.C., Long, L., 2021. Assessing lead exposure in free-

ranging gray wolves (*Canis lupus)* in Minnesota, USA. Journal of Wildlife Diseases

57, 917–921. https://doi.org/10.7589/JWD-D-20-00185

Kim, R., Aro, A., Rotnitzky, A., Amarasiriwardena, C., Hu, H., 1995. K X-ray fluorescence

measurements of bone lead concentration: the analysis of low-level data. Physics in

Medicine and Biology 40, 1475–1485. https://doi.org/10.1088/0031-9155/40/9/007

Kuhn, M., 2022. caret: Classification and Regression Training.

Le Mar, K., McArthur, C., 2000. Relocating radio-collared targeted marsupials after a 1080-

poisoning operation. Tasforests 12, 155–160.

Lohr, M.T., Hampton, J.O., Cherriman, S., Busetti, F., Lohr, C., 2020. Completing a

worldwide picture: preliminary evidence of lead exposure in a scavenging bird from

mainland Australia. Science of the Total Environment 715, 135913.

https://doi.org/10.1016/j.scitotenv.2019.135913

McLellan, M.J., Carrothers, K.L., Spicer, A.M., 2022. Investigating the utility of a handheld

X-ray fluorescence (XRF) device as a field test for suspected ivory. Forensic Science

International: Animals and Environments 2, 100041.

https://doi.org/10.1016/j.fsiae.2022.100041

Pain, D.J., Mateo, R., Green, R.E., 2019. Effects of lead from ammunition on birds and other

wildlife: A review and update. Ambio 48, 935–953. https://doi.org/10.1007/s13280-

019-01159-0

Pay, J.M., Katzner, T.E., Hawkins, C.E., Kock, A.J., Wiersma, J.M., Brown, W.E., Mooney,

N.J., Cameron, E.Z., 2021. High frequency of lead exposure in the population of an

endangered Australian top predator, the Tasmanian wedge-tailed eagle (*Aquila audax*

fleayi). Environmental Toxicology and Chemistry 40, 219–230.

https://doi.org/10.1002/etc.4914

Plaza, P.I., Lambertucci, S.A., 2019. What do we know about lead contamination in wild

vultures and condors? A review of decades of research. Science of the Total

Environment 654, 409–417. https://doi.org/10.1016/j.scitotenv.2018.11.099

Rogers, T.A., Bedrosian, B., Graham, J., Foresman, K.R., 2012. Lead exposure in large

carnivores in the greater Yellowstone ecosystem. Journal of Wildlife Management 76,

575–582. https://doi.org/10.1002/jwmg.277

RStudio Team, 2023. RStudio: Integrated Development for R.

Sánchez, C.A., Penrose, M.T., Kessler, M.K., Becker, D.J., McKeown, A., Hannappel, M.,

Boyd, V., Camus, M.S., Padgett-Stewart, T., Hunt, B.E., Graves, A.F., Peel, A.J.,

Westcott, D.A., Rainwater, T.R., Chumchal, M.M., Cobb, G.P., Altizer, S., Plowright,

R.K., Boardman, W.S.J., 2022. Land use, season, and parasitism predict metal

concentrations in Australian flying fox fur. Science of The Total Environment 841,

156699. https://doi.org/10.1016/j.scitotenv.2022.156699

Singh, R., Khanna, P., Singh, A., Goyal, S., 2022. Elemental characterization of antlers of

various deer species using X-ray fluorescence (XRF): A tool for forensic examination.

Forensic Science International 332, 111172.

https://doi.org/10.1016/j.forsciint.2022.111172

Sonne, C., Lam, S.S., Kanstrup, N., 2023. Commentary: The environmental threats from lead

ammunition. Eco-Environment & Health. https://doi.org/10.1016/j.eehl.2023.02.001

Specht, A.J., Dickerson, A.S., Weisskopf, M.G., 2019a. Comparison of bone lead measured

via portable x-ray fluorescence across and within bones. Environmental Research 172,

273–278. https://doi.org/10.1016/j.envres.2019.02.031

Specht, A.J., Kirchner, K.E., Weisskopf, M.G., Pokras, M.A., 2019b. Lead exposure

biomarkers in the Common Loon. Science of the Total Environment 647, 639–644.

https://doi.org/10.1016/j.scitotenv.2018.08.043

Specht, A.J., Kponee, K., Nkpaa, K.W., Balcom, P.H., Weuve, J., Nie, L.H., Weisskopf,

M.G., 2018a. Validation of X-ray fluorescence measurements of metals in toenail

clippings against inductively coupled plasma mass spectrometry in a Nigerian

population. Physiological Measurement 39, 085007. https://doi.org/10.1088/1361-

6579/aad947

Specht, A.J., Lin, Y., Weisskopf, M., Yan, C., Hu, H., Xu, J., Nie, L.H., 2016. XRF-

measured bone lead (Pb) as a biomarker for Pb exposure and toxicity among children

diagnosed with Pb poisoning. Biomarkers 21, 347–352.

https://doi.org/10.3109/1354750X.2016.1139183

Specht, A.J., Parish, C.N., Wallens, E.K., Watson, R.T., Nie, L.H., Weisskopf, M.G., 2018b.

Feasibility of a portable X-ray fluorescence device for bone lead measurements of

condor bones. Science of the Total Environment 615, 398–403.

https://doi.org/10.1016/j.scitotenv.2017.09.123

Zhang, X., Specht, A.J., Wells, E., Weisskopf, M.G., Weuve, J., Nie, L.H., 2021. Evaluation

of a portable XRF device for in vivo quantification of lead in bone among a US

population. Science of The Total Environment 753, 142351.

https://doi.org/10.1016/j.scitotenv.2020.142351

Fig. 1. A wild Tasmanian devil (*Sarcophilus harrisii*). Photo: Evie Jones.

Fig. 2. Correlation between portable XRF and ICP-MS lead concentration measurements in

Tasmanian devil (*Sarcophilus harrisii*) hair, including all observed values. Shaded polygon

represents the 95% prediction interval.

531 **Table 1.** Lead concentration in hair samples from 39 Tasmanian devils (*Sarcophilus harrisii*),

532 estimated using inductively-coupled mass spectrometry (ICP-MS) and portable X-ray

533 fluorescence (XRF). Estimates are partitioned into low (≤ 3 mg/kg) and high (> 3 mg/kg)

534 exposure categories.

	Mean	Min	Max	SD	% ≤ 3 mg/kg	% > 3 mg/kg
ICP-MS (mg/kg)	1.65	0.12	25.34	4.61	94.87	5.13
Portable XRF (mg/kg)	2.33	0.08	25.16	4.76	94.87	5.13
Portable XRF uncertainty (mg/kg)	0.58	0.20	1.97	0.39	NA	NA

538 **Table 2.** Lead concentrations in hair derived from portable XRF and predicted ICP-MS levels

539 from regression analysis for live Tasmanian devils (*Sarcophilus harrisii*), sampled during a

540 trapping program in a forestry plantation landscape (n = 61).

	Mean	Min	Max	SD	% ≤ 3 mg/kg	% > 3 mg/kg
Predicted ICP-MS (mg/kg)	1.04	-1.88	5.47	1.45	88.52	11.47
Portable XRF (mg/kg)	1.69	-1.39	6.37	1.53	81.97	18.03
Portable XRF uncertainty (mg/kg)	0.85	0.25	2.02	0.45	NA	NA

544 **Table 3.** Model ranking testing the importance of Tasmanian devil (*Sarcophilus harrisii*) age

545 and sex in predicting lead levels in devil hair in a forestry plantation landscape. df = degrees

546 of freedom, AIC = Akaike information criterion, ΔAIC = Akaike information criterion

547 difference, AICwt = Akaike information criterion weight.

Model	df	AIC	ΔAIC	AICwt
Null	2	221.5	0	0.47
Sex	3	222.8	1.24	0.25
Age	4	223.5	1.96	0.18
Age + sex	5	224.7	3.19	0.10

548
549