Metabolic memory of Δ9-tetrahydrocannabinol exposure in pluripotent stem cells and primordial
germ cells-like cells

Roxane Verdikt[1], Abigail A. Armstrong[2], Jenny Cheng[3], Young Sun Hwang[4], Amander T. Clark[4,5,6], Xia
Yang[7,8], and Patrick Allard[1,9]*

[1]Institute for Society and Genetics, University of California, Los Angeles, Los Angeles, CA 90095, USA
[2]Department of Obstetrics/Gynecology and Reproductive Endocrinology and Infertility, University of California, Los
Angeles, CA, USA
[3]Molecular, Cellular, and Integrative Physiology Graduate Program, University of California, Los Angeles, Los
Angeles, CA 90095, USA
[4]Department of Molecular Cell and Developmental Biology, University of California, Los Angeles, Los Angeles, CA,
90095, USA
[5]Center for Reproductive Science, Health and Education, University of California, Los Angeles, Los Angeles, CA
90095, USA
[6]Eli and Edythe Broad Center of Regenerative Medicine and Stem Cell Research, University of California, Los
Angeles, Los Angeles, CA, 90095, USA
[7]Integrative Biology and Physiology Department, University of California, Los Angeles, CA, 90095, USA
[8]Department of Molecular and Medical Pharmacology, University of California, Los Angeles, Los Angeles, CA 90095,
USA
[9]Molecular Biology Institute, University of California, Los Angeles, Los Angeles, CA 90095, USA

* Corresponding author:
Patrick Allard, Boyer Hall, 611 Charles E Young Dr E, University of California, Los Angeles, Los Angeles,
90095. Email: pallard@ucla.edu

ABSTRACT

Cannabis, the most consumed illicit psychoactive drug in the world, is increasingly used by pregnant
women. However, while cannabinoid receptors are expressed in the early embryo, the impact of
phytocannabinoids exposure on early embryonic processes is lacking. Here, we leverage a stepwise in vitro
differentiation system that captures early embryonic developmental cascade to investigate the impact of
exposure to the most abundant phytocannabinoid, Δ9-tetrahydrocannabinol (Δ9-THC). We demonstrate
that Δ9-THC increases the proliferation of naïve mouse embryonic stem cells (ESCs) but not of their primed
counterpart. Surprisingly, this increased proliferation, dependent on the CB1 receptor binding, is only
associated with moderate transcriptomic changes. Instead, Δ9-THC capitalizes on ESCs' metabolic
bivalence by increasing their glycolytic rates and anabolic capabilities. A memory of this metabolic rewiring
is retained throughout differentiation to Primordial Germ Cell-Like Cells in the absence of direct exposure
and is associated with an alteration of their transcriptional profile. These results represent the first in-
depth molecular characterization of the impact of Δ9-THC exposure on early stages of germline
development.

KEYWORDS
Metabolism, cannabis, Δ9-THC, embryonic stem cells, primordial germ cells.

INTRODUCTION

Cannabis is the most widely used illicit psychoactive drug in the world (U.N. Office on Drugs and Crime, 2022). In the United States, an estimated 49.6 million people, roughly 18% of the population, consumed cannabis at least once in 2020, with indications that these numbers will likely increase in the coming years as attitudes and regulations change (Mennis et al., 2023; Substance Abuse and Mental Health Services Administration, 2020). In particular, between 7-12% of expecting women report cannabis use, predominantly during the first trimester to alleviate the symptoms of morning sickness (Chabarria et al., 2016; Volkow et al., 2019; Young-Wolff et al., 2018). These statistics indicate that a significant number of developing embryos are exposed to cannabis, with limited knowledge of the biological repercussions of such exposure.

Among the several hundred unique phytocannabinoids present in Cannabis sativa, (−)-trans-Δ9-tetrahydrocannabinol (Δ9-THC) is chiefly responsible for the psychoactivity of cannabis (Andre et al., 2016). As a result, the level of Δ9-THC in recreational cannabis has increased over the last 10 years and now commonly accounts for 20% of total compounds (Chandra et al., 2019). The psychoactive effects of Δ9-THC arise from its binding and subsequent activation of the G protein-coupled cannabinoid receptors CB1 largely expressed in the central nervous system (Pacher et al., 2020). In this context, Δ9-THC exposure has been shown to durably alter metabolic, transcriptional and epigenetic programs in the brain (Bénard et al., 2012; Prini et al., 2017; Szutorisz and Hurd, 2018; Watson et al., 2015). While over the last decades, significant attention has been paid to Δ9-THC's neurological effects, there is also evidence, albeit more limited, of its impact on reproductive functions (Lo et al., 2022). Data shows CB1 expression in the male and the female reproductive tracts, in the pre-implantation embryo and in the placenta (Lo et al., 2022; Paria et al., 1995). In animal models as well as in humans, exposure to cannabis is associated with reduced fertility, decreased testis weight and sperm count, and impairment of embryo implantation (Lo et al., 2022). In males, these effects are correlated with an alteration of the sperm transcriptome and epigenome (Murphy et al., 2018; Osborne et al., 2020; Schrott and Murphy, 2020). Epidemiological evidence also indicates that Δ9-THC exposure is associated with long-lasting adverse effects, with exposures in parents affecting the offspring (Smith et al., 2020; Szutorisz and Hurd, 2018). Despite this accumulating evidence, the molecular impact and mechanisms of Δ9-THC exposure at the earliest stages of germ cells development remain to be determined.

Progression through states of pluripotency is controlled by metabolic reprogramming in the early mammalian embryo (Verdikt and Allard, 2021; Zhang et al., 2018). Accordingly, cultured pluripotent stem cells (PSCs) exhibiting different developmental potentials are marked by specific metabolic signatures, similar to the ones displayed by their in vivo counterparts in the embryo. For instance, mouse embryonic stem cells (ESCs) are naïve PSCs that are functionally equivalent to the inner cell mass (ICM) of the E3.5 preimplantation mouse blastocyst (Nichols and Smith, 2009). The extended developmental potential of mouse ESCs is associated with their metabolic bivalence, as these cells rely on both glycolysis and oxidative phosphorylation for energy production. Differentiation of naïve ESCs into primed PSCs such as epiblast-like cells (EpiLCs) is accompanied by an important metabolic shift towards aerobic glycolysis, in link with a highly-proliferative phenotype and a more restricted developmental potential (Hayashi et al., 2017; Verdikt and Allard, 2021; Zhang et al., 2018). Primordial germ cells (PGCs), the embryonic precursors of gametes in metazoans (Kurimoto and Saitou, 2018), are considered dormant totipotent cells because they possess the unique ability to reacquire totipotency upon fertilization (Hayashi et al., 2017). In the mouse

embryo, the precursors to PGCs arise around embryonic day 6.25 (E6.25) from formative pluripotent cells
in the epiblast (Kinoshita et al., 2021; Kurimoto and Saitou, 2018). Progressive increase in oxidative
phosphorylation correlates with the specification and differentiation from epiblast PSCs towards PGCs, a
process that can be replicated in vitro by inducing PGC-like cells (PGCLCs) from EpiLCs (Hayashi et al.,
2011). In particular, the extensive metabolic, transcriptional, and epigenetic reprogramming that PGCs
undergo during their development has been proposed to be uniquely sensitive to environmental insults,
with potential consequences in the offspring (Verdikt et al., 2022).
Here, we deployed this in vitro differentiation system to investigate the impact of Δ9-THC exposure
on early developmental stages. We demonstrate that exposure of ESCs and EpiLCs to Δ9-THC durably alters
their metabolome. We reveal that, in the absence of continuous exposure, metabolic memory of Δ9-THC
is passed onto the PGCLCs stage leading to transcriptional defects in these cells. Together, our findings
highlight the role of metabolic reprogramming as a mechanism for early developmental Δ9-THC exposure.
97

RESULTS
Δ9-THC induces cellular proliferation of mouse embryonic stem cells but not of mouse epiblast-like cells.
To model the impact of early Δ9-THC exposure on early embryonic events, we first tested three
distinct developmental windows: 1) exposure of ESCs, 2) exposure of EpiLCs and 3) combined ESCs+EpiLCs
exposure (Figure 1A). Cells were either exposed to the vehicle (mock) or exposed to Δ9-THC in a wide dose
range of 0.1nM-100μM, corresponding to the reported physiologically-relevant concentrations of Δ9-THC
in cannabis users (Fuchs Weizman et al., 2021; Hunault et al., 2008; Pacifici et al., 2019).
The viability of ESCs exposed to increasing concentrations of Δ9-THC for 48h was not significantly
altered until the maximal dose of 100μM, subsequently serving as a positive control, at which only 13.17%
of cells remained alive (Figure 1B, $p=5.10^{-15}$, unpaired T-test). While no significant changes in viability were
observed between 10nM and 1μM Δ9-THC, the number of viable ESCs significantly increased by 1.69, 1.52
and 1.28-fold, respectively, compared to the mock-treated condition (Figure 1C, p=0.002, p=0.01 and
p=0.03 for 10nM, 100nM and 1μM of Δ9-THC, unpaired T-test). Of note, the number of viable ESCs
significantly increased for doses of Δ9-THC as low as 1nM (Supplementary Figure 1, 1.59-fold increase,
p=0.0005, unpaired T-test), corresponding to doses found in users of "light cannabis" products (Pacifici et
al., 2019). To determine whether the increased number of viable cells recovered after Δ9-THC exposure
was due to higher proliferation, we performed bromodeoxyuridine (BrdU) labeling experiments. Exposed
cells were pulsed with BrdU for 30 minutes, and its incorporation in actively dividing cells was measured
by flow cytometry. The percentage of BrdU-positive ESCs significantly increased between 10nM and 10μM
of Δ9-THC compared to the mock-treated condition (Figure 1D, p=0.01, p=0.001, p=0.05 and p=0.01 for
10nM, 100nM, 1μM and 10μM of Δ9-THC, unpaired T-test). Co-staining with DAPI (4',6-diamidino-2-
phenylindole) to analyze the cell cycle showed that, at 100nM of Δ9-THC, a significantly higher proportion
of cells were in the G2/M phase (Supplementary Figure 2, 2.76-fold increase, p=0.02, unpaired T-test),
consistent with cellular proliferation. Finally, whereas the present data was obtained using female ESCs,
the proliferative effect of Δ9-THC exposure on ESCs was also observed in male ESCs (Supplementary Figure
3), suggesting that proliferation is sex-independent.
Next, we derived EpiLCs from unexposed ESCs and performed the same dose-response experiments.
Akin to ESCs, Δ9-THC exposure in EpiLCs did not significantly alter cellular viability until the dose of 100μM
(Figure 1E, 11.56% of viable cells, $p=1.6^{-13}$, unpaired T-test). However, contrary to ESCs, Δ9-THC exposure

in EpiLCs did not significantly increase the number of viable cells nor the percentage of BrdU-positive cells
(Figure 1F and Figure 1G). For the dose of 10µM of Δ9-THC, viable EpiLCs numbers and BrdU-positive EpiLCs
decreased compared to the mock-treated condition (Figure 1F, 1.53-fold decrease, p=0.0007, Figure 1G,
1.42-fold decrease, p=0.0024, unpaired T-test). When continuously exposing ESCs and EpiLCs, cell viability
was more significantly and negatively impacted, except at the dose of 100nM of Δ9-THC (Figure 1H).
Deriving EpiLCs from exposed ESCs and exposing them to Δ9-THC for 48h did not significantly affect either
their cell number nor their incorporation of BrdU (Figure 1I and Figure 1J), indicating that the increased
proliferation observed at the ESCs stage is not carried through the naïve-to-prime transition.
Finally, we further assessed the impact of Δ9-THC exposure in human pluripotent stem cells. Human
embryonic stem cells (hESCs) differ from their murine counterparts, particularly because they resemble a
primed pluripotent state, akin to mouse EpiLCs (Weinberger et al., 2016). Cell viability of hESCS
continuously exposed to 100nM of Δ9-THC was not significantly impacted (Supplementary Figure 4A).
However, hESCs number was significantly decreased upon Δ9-THC exposure (Supplementary Figure 4B,
1.26-fold decrease, p=0.004, unpaired T-test), in a similar trend as observed with mouse EpiLCs (Figure 1E
and Figure 1I).
Together, the systematic testing of different exposure schemes of Δ9-THC in different pluripotent
stem cell populations (mouse and human ESCs and mouse EpiLCs) revealed that physiologically relevant
doses of Δ9-THC (10nM-1µM) specifically stimulate the proliferation of mouse ESCs, but not of human
ESCs, nor of mouse EpiLCs, whether the latter are derived from exposed ESCs or not.
146
Expression of the CB1 receptor does not explain differences in proliferative outcomes.
We next sought to understand the source of variation in proliferative outcomes in response to Δ9-
THC between naïve mouse embryonic stem cells and primed pluripotent epiblast-like cells. Such
differential effects have been previously reported, with Δ9-THC eliciting the proliferation of neural
progenitors (Galve-Roperh et al., 2013) and of human breast carcinoma cell lines (Takeda et al., 2008) but
suppressing the proliferation of activated CD4[+] T cells (Yang et al., 2016) and of non-small cell lung cancer
cells (Preet et al., 2008). In these studies, the differential expression of cannabinoid receptors at the cell
surface was proposed to primarily mediate the variation in cellular outcomes.
We therefore first tested whether expression levels of CB1 varied between ESCs and EpiLCs.
Western-blot analysis of membrane proteins revealed however that CB1 was expressed at the same levels
at the cell surface of both ESCs and EpiLCs (Figure 2A and 2B). We next determined whether the Δ9-THC-
induced proliferative phenotype in ESCs was due to the engagement of the CB1 cannabinoid receptor. To
do so, ESCs were pretreated for 1h with 1µM of SR141716 (also known as rimonabant, a specific CB1
blocker (Rinaldi-Carmona et al., 1994)) then exposed to 100nM or 100µM of Δ9-THC for 48h. Rimonabant
pre-treatment did not significantly alter the viability of ESCs compared to conditions exposed to Δ9-THC
only (Figure 2C) but abolished Δ9-THC-induced ESCs increased cell number at 100nM Δ9-THC (Figure 2D,
1.53-fold decrease, p=1.66^{-05}, when comparing 100nM of Δ9-THC +/- 1µM of SR141716, unpaired T-test).
Notably, SR141716 pre-treatment, while not altering cell viability, reduced cell number compared to
control, suggesting a basal role for CB1 in promoting proliferation.
Thus, the expression of CB1 at the cell surface does not explain the differential impact of Δ9-THC on
ESC and EpiLC proliferation even if CB1 engagement is a required event for this effect in ESCs.

Δ9-THC exposure increases glycolysis in ESCs and EpiLCs.

In the central nervous system, Δ9-THC is a known metabolic perturbator which increases bioenergetic metabolism (Bartova and Birmingham, 1976; Bénard et al., 2012). As mentioned above, the transition of naïve ESCs into the primed state of EpiLCs is accompanied by a switch to glycolysis for energy production (Hayashi et al., 2017; Verdikt and Allard, 2021). Thus, to capture the impact of Δ9-THC at every point of their transition between metabolic states, we used the continuous exposure scheme of ESCs and EpiLCs outlined in Figure 1H-J. Similar to our other exposure schemes, at lower Δ9-THC doses, the proliferation of ESCs was observed but not of EpiLCs. We performed these exposures in a wide Δ9-THC dose range (10nM-10μM) followed by bioenergetics assessment (Figure 3A).

First, we assessed the global energy metabolism of exposed cells by measuring the nicotinamide adenine dinucleotide (phosphate) couple ratios (NAD(P)+/NAD(P)H) using the WST-1 assay. In ESCs, the ratio of NAD(P)+/NAD(P)H significantly increased 1.57, 1.54, 1.29 and 1.38 -fold, for 10nM, 100nM, 1μM and 10μM of Δ9-THC, respectively, compared to the mock-treated condition (Figure 3B, p=2.87^{-06}, p=8.01^{-05}, p=0.03, and p=0.0003 for 10nM, 100nM, 1μM and 10μM of Δ9-THC, unpaired T-test). In contrast, no significant increase was observed in NAD(P)+/NAD(P)H ratios in exposed EpiLCs (Figure 3B). Consistent with the impact of continuous Δ9-THC exposure on EpiLCs viability (Figure 1H), the NAD(P)+/NAD(P)H ratios significantly decreased at 10μM of Δ9-THC in EpiLCs (Figure 3B, 59% decrease for 10μM of Δ9-THC compared to the mock-treated condition, p=6.55^{-09}, unpaired T-test). Of note, the NAD(P)+/NAD(P)H ratios were slightly but significantly decreased at 100nM of Δ9-THC in hESCs (Supplementary Figure 5, 1.09-fold decrease, p=0.04, unpaired T-test), in agreement with the deleterious effect of Δ9-THC on cell number (Supplementary Figure 4B).

Because the elevated NAD(P)+/NAD(P)H levels in Δ9-THC-exposed mouse ESCs could indicate increased mitochondrial activity in the context of oxidative phosphorylation (Locasale and Cantley, 2011), we next studied changes in mitochondrial membrane potential of exposed cells using the Mitotracker CMXRos fluorescent dye (Pendergrass et al., 2004). A significant increase in mean fluorescence intensity (MFI) associated with the mitochondrial stain was observed at 100nM of Δ9-THC in ESCs (Figure 3C, p=0.02, unpaired T-test), indicating that, at this dose, the observed increase in NAD(P)+/NAD(P)H could be explained by higher mitochondrial membrane potential. By contrast, no change in EpiLCs mitochondrial activity was detected (Figure 3C), consistent with these cells relying on glycolysis for energy production (Hayashi et al., 2017; Verdikt and Allard, 2021).

Changes in mitochondrial activity in ESCs upon Δ9-THC exposure, although significant, remained modest and are unlikely to be the sole contributor to the more significant increase in NAD(P)+/NAD(P)H upon exposure. Thus, we performed an in-depth analysis of the differential impact of Δ9-THC on ESCs and EpiLCs bioenergetics by measuring both glycolysis (extracellular acidification rate, ECAR) and mitochondrial respiration (oxygen consumption rate, OCR) using a Seahorse bioanalyzer. At 100nM of Δ9-THC, the maximal glycolytic capacity of both ESCs and EpiLCs increased significantly (Figure 3D, 15% increase, p=0.03 and 22% increase, p=0.03 for ESCs and EpiLCs, respectively, compared to the mock-treated condition, unpaired T-test). In both cell types, a significant decrease in glycolytic capacity was observed at 10μM of Δ9-THC (Figure 3D, 39.8% reduction, p=0.0006 and 44.8% reduction, p=0.0001, for ESCs and EpiLCs, respectively, compared to the mock-treated condition, unpaired T-test). Of note, the maximal glycolytic capacity of EpiLCs in the untreated condition was higher than the one of ESCs, in agreement with their metabolic shift towards aerobic glycolysis (Figure 3D, 7.88% higher ECAR rate in

mock-treated EpiLCs compared to mock-treated ESCs, p=0.03, unpaired T-test). As a consequence, Δ9-THC
exposure significantly impacted more glycolysis in EpiLCs than ESCs, both in basal capacity and upon
mitochondrial inhibition by oligomycin (Supplementary Figure 6A and Figure 6B). In addition, at 100nM of
Δ9-THC, the maximal respiratory capacity of ESCs was significantly increased compared to the mock-
treated condition (Figure 3E, 21.8% increase, p=0.03, unpaired T-test). This increase was observed only for
the maximal respiratory capacity of ESCs, but not for basal respiration, nor for ATP-linked respiration
(Supplementary Figure 6C), suggesting that Δ9-THC impact on mitochondrial respiration does not support
increased energetic production. In agreement with EpiLCs metabolic shift towards a glycolytic phenotype,
increasing doses of Δ9-THC did not alter their maximal respiratory capacity (Figure 3E), nor their global
oxygen consumption rate (Supplementary Figure 6D). In both cell types, a significant decrease in oxygen
consumption rate was observed at 10µM of Δ9-THC (Figure 3E and Supplementary Figure 6C and Figure
1D).
Together, our analysis of cellular bioenergetics following Δ9-THC exposure showed an increased
glycolytic rate in ESCs that was also observed in EpiLCs. However, the increased oxygen consumption and
the associated increase in mitochondrial activity were observed only in ESCs following exposure to 100nM
of Δ9-THC, likely for the oxidization of the accumulating pyruvate generated from glycolysis.
226
Δ9-THC-induced increase in glycolysis supports anabolism and ESCs proliferation
Because our data indicated that the impact of Δ9-THC exposure on stem cells' bioenergetics did not
result in greater ATP production, we next sought to characterize the global metabolic impact of Δ9-THC in
these cells. ESCs and EpiLCs were continuously exposed to 100nM Δ9-THC and intracellular metabolites
were detected and quantified by mass spectrometry (Figure 4A-E). To explore the metabolic signatures in
the different samples, we performed a global principal component analysis (PCA) (Figure 4B). All samples
clustered in well-defined groups of replicates, both by cell type on the first principal component
(accounting for 65.81% of the variation) and by Δ9-THC exposure on the second principal component
(accounting for 20.83% of the variation). Of the 126 metabolites detected in ESCs, 39 were significantly
upregulated (Figure 4C and Supplementary Figure 7A) and only two metabolites – NAPDH and Adenine –
were significantly downregulated. Of the 138 metabolites detected in EpiLCs, 95 were significantly
upregulated (Figure 4C and Supplementary Figure 7B) and only one metabolite – NAPDH – was significantly
downregulated. In agreement with the PCA, the overlap of over-expressed metabolites in response to Δ9-
THC exposure was important between the two stem cell populations (Figure 4C, accounting for 79.49%
and 32.63% of all upregulated metabolites in ESCs and EpiLCs, respectively). The functional interpretation
of the significantly upregulated metabolites confirmed the Δ9-THC-associated increase in energy
metabolism in the two stem cell populations. Indeed, amongst the 25 metabolic pathways upregulated,
pyruvate metabolism and glycolysis were detected in both ESCs and EpiLCs (Figure 4D and Figure 4E,
respectively). Increased mitochondrial respiration was also seen in ESCs with the enrichment of
(ubi)quinone metabolism, indicating an increased synthesis of ubiquinone that serves as an electron
carrier in oxidative phosphorylation. Of note, metabolite measurements showed that the ratio of
glutathione in its reduced to oxidized form (GSH/GSSG) was unchanged in both stem cell types in response
to Δ9-THC (Supplementary Figure 7C), suggesting that the increased mitochondrial respiration does not
cause an overt elevation of oxidative stress. Importantly, and in agreement with the PCA, in both ESCs and
EpiLCs, Δ9-THC exposure elicited an increase in metabolic pathways that feed anabolic reactions, in

particular contributing to the synthesis of amino acids (tyrosine, tryptophan, arginine, alanine, valine,
(iso)leucine, etc.), nucleotides ("Pyrimidine metabolism", "Purine metabolism"), NAD(P)+ ("Nicotinate and
nicotinamide metabolism") and fatty acids ("Butanoate metabolism") (Figure 4D and Figure 4E).

Extensive metabolic profiling of ESCs and EpiLCs upon Δ9-THC exposure thus indicated that the
increased glycolytic rates in both stem cell populations, rather than provoking an increased production of
energy under the form of ATP, participated in increased anabolism. Such increased anabolism could
explain the proliferation observed in ESCs upon Δ9-THC exposure. To test this hypothesis, we exposed ESCs
to 100nM of Δ9-THC for 48h as above but 24h before the harvest, cells were exposed to 10mM of 2-
deoxyglucose (2-DG), an inhibitor of glycolysis (Barban and Schulze, 1961). Despite increasing the energy
stress (Supplementary Figure 8), inhibition of glycolysis by 2-DG did not significantly impact viability over
this shorter time frame and at this concentration (Figure 4F). Importantly, glycolytic inhibition by 2-DG
abrogated the Δ9-THC-induced increase in both cell number and NAD(P)+/NAD(P)H levels (Figure 4G and
Figure 4H, 1.39-fold reduction and $p=5.56^{-05}$ and 1.68-fold reduction and $p=0.0064$, respectively, when
comparing 100nM of Δ9-THC +/- 10mM 2-DG, unpaired T-test). Thus, exposure to Δ9-THC increases
anabolism in both ESCs and EpiLCs, however, this increased anabolism only supports cellular proliferation
in ESCs.

Δ9-THC exposure is associated with the upregulation of genes involved in anabolic pathways in ESCs but
not in EpiLCs.

Our data shows that Δ9-THC exposure increases anabolic pathways in both ESCs and EpiLCs and that
this causes the proliferation of ESCs but not of EpiLCs. We thus next examined whether this differential
impact of Δ9-THC on ESCs and EpiLCs was mirrored by a change in these cells' transcriptomes. To this aim,
we performed RNA-sequencing (RNA-seq) on ESCs and EpiLCs continuously exposed to 100nM Δ9-THC or
to the vehicle control (Figure 5A).

Unsupervised exploration of the global transcriptome by PCA revealed that the vast majority of data
variation could be attributed to the cell type (PC1, accounting for 98% of the variation) rather than to Δ9-
THC exposure (PC2, accounting for 1% of the variation, Figure 5B). In addition, mining for the expression
of pluripotency markers after Δ9-THC exposure suggested that Δ9-THC does not appear to influence the
differentiation dynamics from ESCs to EpiLCs (Supplementary Figure 9). We identified a low number of
differentially expressed genes (DEGs) in both ESCs and EpiLCs (Figure 5C and Figure 5D, respectively),
indicating that Δ9-THC exposure only moderately impacts ESC and EpiLC transcriptomes. In ESCs, only 12
genes were significantly upregulated with a log2(fold-change)>0.5 and only 9 were significantly
downregulated at the same threshold (Figure 5C, significance corresponds to adjusted p-value≤0.05).
More genes were differentially expressed when looking at lower fold-changes (|log2(fold-change)|>0.25,
Figure 5C), confirming that the magnitude of transcriptional effects due to Δ9-THC exposure is moderate.
This low transcriptional impact following Δ9-THC exposure was also observed in EpiLCs (Figure 5D).
Nevertheless, gene ontologies (GO) associated with Δ9-THC-induced DEGs revealed the biological
significance of these low transcriptional changes (Figure 5E). In particular, GO terms associated with
metabolic pathways involved in anabolism were significantly over-represented for upregulated genes in
ESCs following Δ9-THC exposure (Figure 5E), such as: "Cellular aromatic compound metabolic process",
"Cellular nitrogen compound biosynthetic process", "Organonitrogen compound metabolic process". This
suggests that the glycolytic rewiring elicited by Δ9-THC exposure in ESCs has some transcriptional support.
Indeed, when performing joint pathway integration between our transcriptomics data and our targeted

metabolomics (Pang et al., 2022), we observed that Δ9-THC-induced perturbed genes and metabolites
were associated with the observed anabolic effects (Figure 5F). In contrast, GO terms associated with
metabolism were not found within the upregulated DEGs in EpiLCs. However, several GO terms relating to
alterations in cellular components were enriched in EpiLCs (Figure 5E), such as: "Organelle organization",
"Cellular component organization or biogenesis", "Microtubule-based process". This indicates that Δ9-THC
exposure significantly upregulated genes in EpiLCs that impact organelles structure, integrity and position,
in agreement with several reports in the literature (Lojpur et al., 2019; Miller et al., 2019).

Multiple reports indicate that Δ9-THC exposure alters the epigenome of sperm and brain tissue,
both in terms ofDNA methylation level and histone post-translational modifications (Murphy et al., 2018;
Prini et al., 2017; Schrott and Murphy, 2020; Watson et al., 2015). Despite the low transcriptional impact
of Δ9-THC exposure in ESCs and EpiLCs, we evaluated the expression of more than 100 genes encoding
epigenetic modifiers in our RNA-seq datasets (Supplementary Table 1). Data shows that the expression
levels of multiple epigenetic modifiers were significantly altered, albeit at a low levels, in either ESCS or
EpiLCs upon Δ9-THC exposure (|log2(fold-change)|>0.25, Supplementary Figure 10). In particular, the
expression levels of the DNA dioxygenase *Tet2* were significantly increased in EpiLCs following Δ9-THC
exposure, which would contribute to changes in DNA methylation dynamics and differentiation potential
(Sohni et al., 2015). The expression of the two histone deacetylases *Hdac5* and *Hdac11* were significantly
decreased in ESCs following Δ9-THC exposure, while the expression of the histone phosphorylase *Rps6ka5*
increased under the same conditions. In EpiLCs, the expression of the histone methyltransferase *Kmt2c*
and of the regulators of histone ubiquitination Dzip3 and *Mysm1* were all significantly increased upon Δ9-
THC exposure. Collectively, RNA-seq data suggests the existence of an epigenetic remodelling following
Δ9-THC exposure, although further analysis of the respective epigenetic marks associated with these
modifiers is needed.

Together, our analysis of ESCs and EpiLCs transcriptomes reveals a difference in the response of
these stem cell populations to Δ9-THC exposure: the transcriptional alterations observed in ESCs
supported their increased anabolism and proliferation, whereas changes in EpiLCs gene expression did not
correlate with their metabolic changes.

322 Proliferation of Primordial Germ Cell-Like Cells stemming from prior Δ9-THC exposure.

PGCs display a distinct transcriptomic and metabolic profile compared to their cellular precursors
that are recapitulated in vitro during the differentiation of ESCs into EpiLCs and then of EpiLCs into PGCLCs.
Thus, we asked whether the metabolic alterations observed in ESCs and EpiLCs could lead to an altered
differentiation program in PGCLCs. To this aim, we continuously exposed ESCs and EpiLCs to a Δ9-THC dose
range of 1nM-1µM (or mock control), before changing to a Δ9-THC-free media and inducing PGCLCs
differentiation (Figure 6A). In particular, we took advantage of ESCs that harbor two fluorescent reporters
for germline markers, Blimp1:mVenus and Stella:CFP (Ohinata et al., 2008). Thus, the induction efficiency
of PGCLCs within 5-days embryoid bodies can be detected by monitoring the fluorescence associated with
each cell in flow cytometry, allowing for the determination of a double-negative population (DN), a single-
positive population (SP) wherein Blimp1:mVenus is expressed and a double-positive population (DP)
expressing both Blimp1:mVenus and Stella:CFP, which represents the true specified PGCLC population
(Supplementary Figure 11).

We first measured the impact of ESCs + EpiLCs Δ9-THC exposure on PGCLC induction efficiency. Flow
analyses revealed a dose-dependent increase in the induction efficiency of SP and DP cell populations
(Figure 6B). Specifically, at 100nM Δ9-THC, a significant decrease in DN was observed, with a corresponding
significant increase of 1.14-fold in SP and of 1.64-fold in DP cells (Figure 6C, p=0.0002, p=0.05, and p=1.55^{-06}
for 100nM of Δ9-THC in DN, SP and DP populations respectively compared to the mock-treated
condition, unpaired T-test). The same pattern was observed when male ESCs and EpiLCs exposed to 100nM
of Δ9-THC were differentiated in PGCLCs (Supplementary Figure 12).
To determine if the increased proportion of PGCLCs generated from exposed precursors was due to
higher proliferative kinetics, we performed a proliferation tracing assay (Tempany et al., 2018). The tracing
dye was added to the cells on the day of aggregate formation, and fluorescence attenuation due to cell
division was measured in each subpopulation on day 5. At 100nM Δ9-THC, a smaller proportion of DN cells
underwent two or three mitotic divisions compared to the control (Figure 6D, 1.14-fold fewer cells and
1.12-fold fewer cells, p=0.05 and p=0.04 for 2 divisions and 3 divisions, respectively, unpaired T-test). In
parallel, for the same dose, a significantly higher proportion of SP and DP cells underwent three mitotic
divisions compared to the control (Figure 6D, 1.24-fold and 1.11-fold, p=0.03 and p=0.0035, for 3 divisions
in SP and DP cells, compared to the control, unpaired T-test). These results, therefore, indicate that the
higher number of PGCLCs observed upon Δ9-THC exposure originates from their increased proliferation
during their specification and differentiation.
Finally, we sought to determine if the increased PGCLC proliferation was not due to residual
intracellular Δ9-THC persisting from EpiLCs over the span of PGCLCs differentiation. To do so, intracellular
levels of Δ9-THC were quantified by mass spectrometry in the cells on the day of aggregate formation and
in day 5 embryoid bodies (referred as to "EpiLCs" and "PGCLCs", respectively, in Supplementary Figure 13).
Data shows that no Δ9-THC could be detected in day 5 embryoid bodies, indicating that Δ9-THC does not
persist to levels higher than the limit of detection of mass spectrometry (1ng/mL). This suggests that the
proliferative effects are not due to residual Δ9-THC persisting in the cells during differentiation towards
PGCLCs. Thus, Δ9-THC causes an alteration of the developmental kinetics that PGCLCs normally undergo,
even in the absence of direct continuous exposure.
362

PGCLCs derived from Δ9-THC-exposed cells present an altered metabolism and transcriptome
Since exposure to Δ9-THC prior to their specification increased the number of PGCLCs and ESCs and
PGCLCs share similar metabolic programs (Hayashi et al., 2017; Verdikt and Allard, 2021), we next sought
to characterize their associated metabolic and transcriptional changes. We, therefore, assessed the impact
of exposure of ESCs + EpiLCs to 100nM Δ9-THC on PGCLCs metabolism (Figure 7A).
First, NAD(P)+/NAD(P)H assessment revealed a modest but significant 1.17-fold increase in
NAD(P)+/NAD(P)H ratio in whole day 5 embryoid bodies deriving from exposed ESCs + EpiLCs compared
to those deriving from mock-treated cells (Figure 7B, p=0.01, unpaired T-test). To garner cell type-specific
information on whether these metabolic changes were related to mitochondrial activity and the
differentiation of PGCLCs, we assessed the mitochondrial membrane potential of each subpopulation in
day 5 embryoid bodies. Embryoid bodies were incubated with Mitotracker CMXRos (Pendergrass et al.,
2004), and dissociated and analyzed by flow cytometry. The MFI associated with the mitochondrial stain
was then measured in each subpopulation (Figure 7C). A significant increase in MFI was observed in DN,
SP and DP populations deriving from exposed ESCs +EpiLCs compared to those deriving from mock controls

(Figure 7C, 1.17, 1.16, 1.23 -fold, p=0.006, p=0.05 and p=0.01 for DN, SP and DP, respectively, unpaired T-
test). These results indicate that the metabolic changes induced by Δ9-THC prior to PGCLCs induction and
differentiation are not reset during the profound reprogramming that PGCLCs undergo.

Because our results indicated a sustained impact of Δ9-THC beyond the period of direct exposure,
we further examined PGCLCs by performing a transcriptomic analysis. In particular, day 5 embryoid bodies
deriving from ESCs + EpiLCs, either exposed to 100nM of Δ9-THC or mock-exposed, were sorted and the
total RNA of DP subpopulations, representing true PGCLCs, was analyzed by RNA-seq (Figure 7A).
Unsupervised analysis of the global transcriptome in DP PGCLCs by PCA delineated a transcriptional
signature of prior Δ9-THC exposure (Figure 7D, PC1 accounting for 59% of the variance and PC2 accounting
for 24% of variance). Volcano plot of DEGs between DP PGCLCs deriving from mock- or 100nM Δ9-THC-
exposed ESCs and EpiLCs revealed that most of the significant transcriptional change was towards
downregulation rather than upregulation (Figure 7E, 11 genes were significantly upregulated whereas 97
were significantly downregulated, |log2(fold-change)|>0.25 and adjusted p-value≤0.05). Despite the low
number of upregulated DEGs, the functional annotation of their associated GO terms showed that all
terms enriched corresponded to metabolic processes involved in oxidative phosphorylation (Figure 7F,
"Aerobic electron transport chain", "Mitochondrial respiratory chain complex I assembly", "Electron
transport couple proton transport"). Thus, our data indicate that the metabolic changes induced by
exposure to Δ9-THC prior to PGCLCs specification are retained through transcriptional reprogramming.
Importantly, while our results show that pre-specification Δ9-THC exposure increases PGCLCs number and
mitochondrial activity, the functional annotation of GO terms associated with downregulated DEGs
suggests degradation of PGCLCs quality. Indeed, and reminiscent of GO terms observed in EpiLCs, several
GO terms relating to alterations in structural cellular components ("Anatomical structure morphogenesis",
"Cellular anatomical entity"), and in particular the interface with the extracellular environment ("External
encapsulating structure organization", "Membrane", "Cell periphery", "Extracellular region",
"Extracellular space", "Extracellular matrix structural constituent") were enriched (Figure 7G).
Furthermore, GO terms associated with cell adhesion and junction ("Cell adhesion", "Cell migration",
"Collagen metabolic process", "Cell junction", "Anchoring junction", "Collagen trimer") were also enriched
in downregulated genes.

Together, our data show that Δ9-THC exposure in ESCs and EpiLCs durably alters their metabolome
and that these changes are carried through PGCLCs specification and differentiation, leading to an
alteration of PGCLCs transcriptional program (Figure 8).

DISCUSSION

With greater social acceptance and legalization, cannabis use has increased worldwide (Mennis et
al., 2023; U.N. Office on Drugs and Crime, 2022). Yet, the impact of such heightened use on reproductive
functions, and in particular, on the earliest developmental stages is not well understood. Cannabis use
directly alters adult male fertility and causes abnormal embryo implantation (Lo et al., 2022). Using a well-
characterized in vitro model of early embryonic differentiation events culminating in the differentiation of
PGCLCs, our study is the first to shed light on the impact of Δ9-THC at these stages which unfold during
the first trimester in humans (Chabarria et al., 2016; Volkow et al., 2019; Young-Wolff et al., 2018).

Our data revealed the differential effects of Δ9-THC on naïve and primed pluripotent stem cells,
respectively represented by mouse ESCs and EpiLCs. In particular, exposure to Δ9-THC increased ESCs

proliferation which was in a similar range to what has been previously reported for human breast
carcinoma cell lines (about 30-50% between 10nM and 1μM of Δ9-THC) (Takeda et al., 2008). Differential
expression and use of cannabinoid receptors on the surface of exposed cells have been shown to correlate
with Δ9-THC proliferative phenotypes (Galve-Roperh et al., 2013; Preet et al., 2008; Takeda et al., 2008;
Yang et al., 2016). However, our experiments demonstrated that despite being required for Δ9-THC-
induced proliferation in ESCs, CB1 expression did not significantly differ at the surface of ESCs and EpiLCs.

Because Δ9-THC is a known perturbator of mitochondrial function as previously described in the
central nervous system (Bartova and Birmingham, 1976; Bénard et al., 2012), we studied the metabolic
impact of its exposure in ESCs and EpiLCs. Our data indicate that, at 100nM, Δ9-THC exposure increased
the glycolytic rate in both ESCs and EpiLCs. Bioenergetics analyses and metabolite measurements showed
that this increased glucose metabolism did not support increased energy production in the mitochondria,
but rather, that it led to the accumulation of metabolic intermediates used in anabolic reactions for the
synthesis of amino acids, nucleotides, and lipids. Thus, the metabolic signatures associated with Δ9-THC
exposure are reminiscent of those inherently occurring during naïve-to-prime transition, during which
increased aerobic glycolytic rates feed anabolic reactions ultimately fueling proliferation (Lunt and Vander
Heiden, 2011). We verified this model by testing the requirement of increased glycolysis to support
proliferation and indeed observed that ESCs proliferation upon Δ9-THC exposure is abrogated in the
presence of the glycolytic inhibitor 2-DG.

Transcriptomic analyses revealed that the metabolic reprogramming induced by Δ9-THC exposure
in ESCs was transcriptionally encoded, with increased expression of genes involved in anabolic pathways.
In contrast, functional annotations of DEGs in EpiLCs did not show such transcriptional control of increased
anabolism. Comparing the outputs of the metabolomic and transcriptomic analyses (i.e. PCA plots and
volcano plots), the impact of Δ9-THC at these early stages seems to be primarily metabolic, although the
moderate effects on the transcriptome appear to support the metabolic outcome as revealed by our
integrated analysis (Figure 5F). Together, we propose that Δ9-THC exposure elicits a reprogramming of
ESCs that (1) coaxes them to rely more on aerobic glycolysis, (2) drives anabolic pathways, and therefore
(3) leads to their proliferation. In EpiLCs, the impact of Δ9-THC exposure is not sufficient to override the
cellular and metabolic programs of these already highly proliferative cells that are fully reliant on aerobic
glycolysis (Figure 8).

Finally, we assessed the impact of Δ9-THC exposure in ESCs and EpiLCs on the differentiation of
PGCLCs. Our data indicate that at the physiologically relevant dose of 100nM of Δ9-THC, a significant
increase in PGCLCs was observed. In particular, during PGCLCs differentiation, metabolic reprogramming
and increased oxidative phosphorylation play a critical role in the reacquisition of an extended
developmental potential (Hayashi et al., 2017; Verdikt and Allard, 2021). Thus, we investigated whether
the metabolic alterations observed in ESCs and EpiLCs upon Δ9-THC exposure could be carried through
PGCLCs differentiation. Metabolic characterization revealed that PGCLCs arising from exposed ESCs and
EpiLCs showed increased mitochondrial respiration. Thus, in the absence of direct or indirect continuous
exposure, Δ9-THC still has lasting consequences on the metabolome of embryonic germ cells. A recent
study in drosophila reported that nutrient stress induces oocyte metabolites remodelling that drives the
onset of metabolic diseases in the progeny (Hocaoglu et al., 2021). This indicates that non-DNA-associated
factors, such as germline metabolites, can act as factors of inheritance. Similarly, we show here that
exposure to Δ9-THC remodels ESCs and EpiLCs metabolome and that a metabolic memory of this exposure

is retained during PGCLCs differentiation (Figure 8). In addition to metabolic remodelling, we show that
the PGCLCs transcriptome is also altered. In particular, despite proliferation and a higher number of cells,
the number of DEGs that were downregulated in PGCLCs deriving from Δ9-THC-exposed ESCs and EpiLCs
suggests a general degradation of PGCLCs' homeostasis. Functional annotation further indicated that these
downregulated genes are related to structural cellular components, to the interaction with the
extracellular environment and, specifically, to cell adhesion and junction. During the development of the
central nervous system, perinatal Δ9-THC exposure has also been associated with alteration in cell
adhesion, with an impact on neuronal interactions and morphology (Gómez et al., 2007; Keimpema et al.,
2011; Kittler et al., 2000). Cell-cell adhesion is crucial in PGCs' formation both in cell culture systems
(Okamura et al., 2003) as well as in vivo where it controls PGCs motility during their migration towards the
developing somatic gonad (Barton et al., 2016). Our results thus show that exposure to Δ9-THC prior to
specification affects embryonic germ cells' transcriptome and metabolome. This in turn could have
adverse consequences on cell-cell adhesion with an impact on PGC normal development in vivo.

Despite epidemiological evidence of cannabis exposure in parents being associated with adverse
effects in the offspring (Smith et al., 2020; Szutorisz and Hurd, 2018), the molecular mechanisms involved
in the inheritance of exposure have not been extensively studied. In vitro models for germ cell
development offer a unique opportunity for such studies (Verdikt et al., 2022) and our work in mouse
PGCLCs identifies metabolites as relevant carriers of information across developmental stages. While in
vitro gametogenesis, up to the reconstitution of fully functional spermatozoa and oocytes, is a reality in
mouse (Hikabe et al., 2016; Komeya et al., 2018; Luo and Yu, 2022), transposition to humans has been
hindered by ethical and technical considerations (Luo and Yu, 2022). For instance, protocols for human
PGCLCs have been developed (Gell et al., 2020) but proper induction to viable, fertile offspring cannot be
verified (Luo and Yu, 2022). Nevertheless, our preliminary results in hESCs show that Δ9-THC exposure
negatively impacts their global energy metabolism (Supplementary Figure 5). Future studies will need to
confirm whether such metabolic reprogramming is also carried over developmental stages in human
models for the germline.

Together, our studies reveal a moderate but significant impact of Δ9-THC exposure on early
embryonic processes. Our work also highlights the importance of the metabolic remodelling induced by
Δ9-THC and its potential role as a driver of exposure memory through differentiation stages.

ACKNOWLEDGMENTS
We thank Jessica Scholes, Felicia Codrea and Jeffrey Calimlim from the UCLA BSCRC Flow Cytometry
Core for their assistance in cell sorting. We thank Linsey Stiles (UCLA Mitochondria and Metabolism Core),
Johanna ten Hoeve-Scott and Thomas Graeber (UCLA Metabolomics Center) for their technical support in
the metabolomics analyses. We thank Xinmin Li and his team at the UCLA Technology Center for Genomics
& Bioinformatics for their technical support in high-throughput sequencing. We thank Gazmend Elezi and
Julian Whitelegge (UCLA Pasarow Mass Spectrometry Laboratory) for their help in Δ9-THC quantification.
We thank Luigi Bellocchio and Giovanni Marsicano for their recommendations in Δ9-THC exposure setups.
We would like to acknowledge the UCLA Cannabis Research Initiative (UCRI) and Dr Ziva Cooper for their
continuous support and encouragements.
PA is supported by NIEHS R01 ES027487, the John Templeton Foundation Grant 60742, and the Iris
Cantor-UCLA Women's Health Center and the NCATS UCLA CTSI Grant Number UL1TR001881. RV is a

postdoctoral fellow and acknowledges support from the Belgian-American Educational Foundation (BAEF).
ATC is supported by NIH/NICHD R01HD079546 (ATC).

MATERIAL AND METHODS
Data availability
The RNA sequencing data from this study is made available at the Gene Expression Omnibus (GEO)
under the following accession number GSE226955. All other data are available upon request.

Cell culture and PGCLCs model
Mouse ESCs containing the two fluorescent reporters Blimp1::mVenus and Stella::ECFP (BVSC cells)
were described previously (Ohinata et al., 2008). The female BVSC clone H18 and male clone R8 were
kindly provided by Mitinori Saitou. Cells were seeded on coated plates (Poly-L-ornithine [0.001%; A-004-
C; Sigma-Aldrich] and laminin [300ng/mL; L2020; Sigma-Aldrich]) in 2i+LIF culture medium (N2B27 Media,
CHIR99021 [30μM; NC9785126; Thermo Fisher], PD0325901 [10μM; NC9753132; Thermo Fisher], ESGRO®
Leukemia Inhibitory Factor (LIF) [1,000 U/mL, ESG1106; Sigma-Aldrich]) for 48h. Differentiation of ESCs to
EpiLCs was performed by seeding the cells on Human Plasma Fibronectin (HPF)-coated plates [16.7μg/mL;
33016015; Thermo Fisher] in the presence of EpiLC induction medium (N2B27 medium containing activin
A [20ng/mL; 50-398-465; Thermo Fisher], basic fibroblast growth factor (bFGF) [12ng/mL; 3139FB025; R&D
Systems], and KnockOut Serum Replacement [KSR, 1%; Thermo Fisher]). For PGCLCs induction, 44h EpiLCs
were harvested using TrypLE™ Select (1X) (Thermo Fisher) and seeded either in 96-wells plate (Nunclon
Sphera, Thermo Fisher) or in EZsphere plates for large-scale induction (Nacalai) in the presence of GK15
medium (Glasgow's Minimal Essential Medium [GMEM, 11710035, Thermo Fisher] supplemented with
15% KSR, 0.1 mM Minimal Essential Medium Nonessential Amino Acids [MEM-NEAA], 1 mM sodium
pyruvate, 0.1 mM 2-mercaptoethanol, 100U/mL penicillin, 0.1mg/mL streptomycin, and 2 mM L-
glutamine in the presence of bone morphogenetic protein 4 [BMP4; 500ng/mL; 5020-BP-010/CF; R&D
Systems], LIF, stem cell factor [SCF; 100ng/mL; 50-399-595; R&D Systems], bone morphogenetic protein
8b [BMP8b; 500ng/mL; 7540-BP-025; R&D Systems], and epidermal growth factor [EGF; 50ng/mL;
2028EG200; R&D Systems]. Cells were cultured for 5d before collection, dissociation of embryoid bodies
and downstream experiments.
The hESCs UCLA2 cells (Diaz Perez et al., 2012) were cultured on plates coated with Recombinant
laminin-511 E8 (iMatrix-511 Silk, 892 021, Amsbio) and were maintained under a feeder-free condition in
the StemFit® Basic03 medium (SFB-503, Ajinomoto) containing bFGF (100-18B, Peprotech). Prior to
passaging, hESC cultures were treated with a 1:1 mixture of TrypLE Select (12563011, Thermo Fisher) and
0.5 mM EDTA/PBS for 15 min at 37ºC to dissociate them into single cells. For routine maintenance, hESCs
were plated into a 6-well plate (3516, Corning) at a density of 2×10^3 cells/cm2 with 10μM ROCK inhibitor
(Y-27632; Tocris, 1254) added in culture medium for 1 day after hESCs passaging. hESCs were plated into
24-wells plate (3526, Corning) at 10,000 cells per well for the viability and viable cell count. For WST-1
assays, hESCs were seeded in 96-wells plate (353072, Falcon) at 1,500 cells per well.
All cells were cultured in a humidified environment at 37°C under 5% CO_2. All cells were tested
negative for mycoplasma by a PCR test (ATCC® ISO 9001:2008 and ISO/IEC 17025:2005 quality standards).

Δ9-THC exposures

To assess the impact of Δ9-THC exposure on the developmental trajectory of PGCLCs, three exposure schemes were tested: 1) ESCs exposure only, 2) EpiLCs exposure only, and 3) ESCs+EpiLCs exposure. The stock of Δ9-THC was obtained from the National Institute on Drug Abuse (7370-023 NIDA; Bethesda, MD). The stock was adjusted to a concentration of 200mM diluted in ethanol, aliquoted and stored according to the DEA's recommendations. The dose range of 0-100μM was determined based on Δ9-THC physiological measurements in the blood, plasma, and follicular fluid (Fuchs Weizman et al., 2021; Hunault et al., 2008; Pacifici et al., 2019). For each exposure, new aliquots of Δ9-THC were diluted in ESCs or EpiLCs culture media in coated tubes (Sigmacote, Sigma Aldrich). Exposure was performed for 48h. Solubility tests were performed and ethanol was added to reach the same amount for each Δ9-THC concentration (0.05% ethanol). Vehicle control corresponded to 0.05% ethanol added to the respective culture media for ESCs or EpiLCs. All experiments performed are authorized under DEA registration number RA0546828.

Δ9-THC quantification

Samples were treated with 500μl of 1% Formic Acid (543804, Sigma Aldrich). Labelled Tetrahydrocannabinol (THC-d3), used as an internal standard, was added to every sample to account for compound loss during sample processing. Samples were then mixed vigorously and centrifuged at 16.000g for 5 min at room temperature. The supernatants were loaded into phospholipid removal cartridges (Phenomenex Phree) and the eluents were dried down in a vacuum concentrator. Samples were reconstituted in 30μl of HPLC-grade water, vortexed rigorously, and centrifuged at 16.000g for 5 min at room temperature. The supernatant was transferred to HPLC vials and 15μl were injected for analysis onto a hybrid linear ion trap/orbitrap mass spectrometer (Thermo Scientific LTQ Orbitrap XL, UCLA Pasarow Mass Spectrometry Lab). For specificity and accurate quantitative measurement, the mass spectrometer was set to fragment preselected precursor ions for THC and THC-d3 under standard MS/MS fragmentation conditions in positive ion mode. The mass spectrometer was coupled to a Dionex Ultimate 300 HPLC (Thermo Scientific) with a reversed phase Phenomenex analytical column (Kinetex 1.7μm Polar C18 100 Å 100 x 2.1 mm) equilibrated in eluant A (water/formic acid, 100/0.1, v/v) and eluted (100μl/min) with a linearly increasing concentration of eluant B (acetonitrile/formic acid, 100/0.1, v/v; min/%B, 0/5, 5/5, 8/95, 13/95, 14/5, 25/5). Data was collected and processed with instrument manufacturer-supplied software Xcalibur 2.07. A set of standard curve samples were prepared in cell culture media for each experiment. Samples and standards were prepared in duplicates. The standard curve was made by plotting the known concentration of THC per standard against the ratio of measured chromatographic peak areas corresponding to the THC over that of the IS THC-d3 (analyte/IS). The trendline equation was then utilized to calculate the absolute concentrations of the THC in cell culture.

PGCLCs induction efficiency

Changes in PGCLCs induction were calculated by flow cytometry. Practically, d5 aggregates were harvested, dissociated using TrypLE™ Select, and resuspended in fluorescence-activated cell sorting (FACS) buffer (1×Dulbecco's phosphate buffered saline [DPBS], 1% BSA, 1 mM EDTA, 25 mM HEPES). Quantification of subfractions of double-positive PGCLCs (Blimp1::mVenus+ and Stella::ECFP+), single-positive (Blimp1::mVenus+) and double-negative cells was performed on a BD Biosciences LSRII (UCLA BSCRC Flow Cytometry Core). Cells were initially identified by forward- and side-scatter gating, with back-

gating used to verify the accuracy by which target cell populations were identified. Cell populations of
interest were identified by 2-D plots displaying the parameter of interest, using embryoid bodies cultured
in GK15 medium without added cytokines and BMPs as a negative control. Fluorescent compensation
beads were used as positive controls and to calculate the spectral overlap (Thermo Fisher, A10514 and 01-
2222-42 adsorbed to a CD45 Pacific Blue antibody [OB180026], serving as compensation control for
mVenus and ECFP, respectively). Manually defined gates as well as quadrants were used, as indicated. The
FlowJo software was used to calculate the percentage of induction and generate the associated graphs
(version 10, FlowJo, LLC).

Cell viability and proliferation studies
The viability and viable cell count of ESCs and EpiLCs were calculated using Trypan blue (0.4%,
Thermo Fisher) on a Countess II FL Automated Cell Counter (Thermo Fisher). For BrdU incorporation
studies, cells were permeabilized, fixed, and stained using the BrdU Flow Kit (PerCP-Cy™5.5 Mouse anti-
BrdU, BD Biosciences) before analysis by flow cytometry on a BD Biosciences LSRII (UCLA BSCRC Flow
Cytometry Core). Quantification of PGCLCs proliferation was performed using CellTrace™ Yellow (5µM,
added at the induction, Thermo Fisher), which binds to intracellular amines after diffusing through cell
membranes. The overall fluorescent signal, which gradually decreases as cell division occurs, reflects the
number of cell divisions occurring and was measured on a BD Biosciences LSRII (UCLA BSCRC Flow
Cytometry Core). The FlowJo software was used to calculate the percentage of induction, the number of
cell divisions and generate the associated graphs (version 10, FlowJo, LLC).

CB1 antagonist treatment
To block the effects of Δ9-THC on the cannabinoid receptor CB1, ESCs were plated on 48-well plate
and were pre-treated with 1µM of SR141716/Rimonabant (SML0800, Sigma Aldrich) for 1h before being
exposed to the dose range of Δ9-THC, as above. After 24h incubation, this procedure was repeated and
cells were harvested after 48h total incubation. The viability and viable cell count was calculated using
Trypan blue (0.4%, Thermo Fisher) on a Countess II FL Automated Cell Counter (Thermo Fisher). The
concentration of 1µM of Rimonabant was chosen based on previous experiments (Lojpur et al., 2019) and
did not impact cell viability nor cell number on its own.

Western blotting
Membrane proteins were extracted from cell pellets using the Mem-PER™ Plus Membrane Protein
Extraction Kit (89842, Thermo Fisher) according to the manufacturer's protocol. Western blotting was
performed with 25µg of protein extracts. The immunodetection was assessed using primary antibodies
targeting CB1 (101500, Cayman Chemical) or β-actin (3700, Cell Signaling Technology) as loading control.
Horseradish peroxidase (HRP)-conjugated secondary antibodies were used for chemiluminescence
detection (Amersham).

WST-1 assay
The colorimetric assay WST-1 was used according to the manufacturer's instructions (Roche). The
tetrazolium salt WST-1 is reduced by mitochondrial dehydrogenases to formazan using NAD(P)H as co-
substrates. Thus, the quantity of formazan is directly proportional to NAD(P)$^{+}$.

Mitochondrial activity

Staining for mitochondria was performed by incubating cells at 37°C with 250nM MitoTracker CMXRos (M7512, Thermo Fisher) for 30min (Pendergrass et al., 2004). Cells were washed and analyzed by flow cytometry on a BD Biosciences LSRII (UCLA BSCRC Flow Cytometry Core). The FlowJo software (version 10, FlowJo, LLC) was used to calculate the mean fluorescence intensity (MFI) corresponding to the average fluorescence intensity of each event of the selected cell population within the chosen fluorescence channel associated to MitoTracker CMXRos.

Seahorse experiments

The extracellular acidification rate (ECAR) and the oxygen consumption rate (OCR) are indicative of glycolysis and mitochondrial respiration, respectively. A total of 10×10^3 ESCs and 8×10^3 EpiLCs were seeded on Seahorse XF96 plates (101085-004, Agilent Technologies) and exposed to increasing doses of Δ9-THC for 48h. On the day of the assay, cells were washed with assay medium (unbuffered DMEM assay medium [5030, Sigma Aldrich] supplemented with 31.6mM NaCl, 3mg/L phenol red, 5mM HEPES, 5mM glucose, 2mM glutamine and 1mM sodium pyruvate). For OCR measurement, compounds were injected sequentially during the assay resulting in final concentrations of 2μM oligomycin, 0.75μM and 1.35μM FCCP, 1μM rotenone and 2μM antimycin. ECAR was measured in parallel. The measured quantities were normalized to the protein content as measured by a BCA quantitation (23227, Thermo Fisher).

Mass spectrometry-based metabolomics analysis

To extract intracellular metabolites, cells were rinsed with cold 150mM ammonium acetate (pH 7.3) then incubated with 80% ice-cold methanol supplemented with 10 nmol D/L-norvaline for 1h. Following resuspension, cells were pelleted by centrifugation (15,000g, 4°C for 15min). The supernatant was transferred into a glass vial and metabolites were dried down under vacuum, then resuspended in 70% acetonitrile. Mass spectrometry analysis was performed at the UCLA Metabolomics Center with an UltiMate 3000RSLC (Thermo Scientific) coupled to a Q Exactive mass spectrometer (Thermo Scientific) in polarity-switching mode with positive voltage 3.0 kV and negative voltage 2.25 kV. Separation was achieved using a gradient elution with (A) 5mM NH4AcO (pH 9.9) and (B) acetonitrile. The gradient ran from 15% (A) to 90% (A) over 18 min, followed by an isocratic step for 9 minutes and re-equilibration for 7 minutes. Metabolites were quantified as area under the curve based on retention times and using accurate mass measurements (≤ 3 ppm) with the TraceFinder 3.1 software (Thermo Scientific). For heatmap depiction, the relative amounts of metabolites were normalized to the mean value across all samples for the same condition and to the number of viable cells harvested in parallel on a control plate. Pathway enrichment for up- and downregulated KEGG metabolites (|log2(fold-change)|=0.25) was determined using the MetaboAnalyst 5.0 platform (www.metaboanalyst.ca)(Pang et al., 2022).

RNA-sequencing

Total RNA was extracted from ESCs and EpiLCs pellets using the AllPrep DNA/RNA Micro Kit (Qiagen), according to the manufacturer's protocol. For PGCLCs, d5 embryoid bodies were harvested and cells were dissociated using TrypLE™ Select followed by resuspension in fluorescence-activated cell sorting (FACS) buffer (1×Dulbecco's phosphate buffered saline [DPBS], 1% BSA, 1 mM EDTA, 25 mM HEPES) and cell suspension were passed through a cell strainer (70μm). Cells were sorted on a BD Biosciences FACSAria III (UCLA BSCRC Flow Cytometry Core). Practically, cell populations of interest, being double-positive

661 (Blimp1::mVenus+ and Stella::ECFP+) were sorted and collected in microtubes containing GK15 medium.
662 Total RNA was extracted from double-positive PGCLCs using the AllPrep DNA/RNA Micro Kit (Qiagen). RNA
663 concentration was measured using a NanoDrop™ 2000 UV spectrophotometer (Thermo Fisher). Libraries
664 were prepared with the KAPA mRNA HyperPrep Kit (BioMek) or with the RNA library prep kit (ABClonal)
665 following the manufacturers' protocols. Briefly, poly(A) RNA were selected, fragmented and double-
666 stranded cDNA synthesized using a mixture of random and oligo(dT) priming, followed by end repair to
667 generate blunt ends, adaptor ligation, strand selection, and polymerase chain reaction (PCR) amplification
668 to produce the final library. Different index adaptors were used for multiplexing samples in one sequencing
669 lane. Sequencing was performed on an Illumina NovaSeq 6000 sequencers for paired end (PE), 2×150 base
670 pair (bp) runs. Data quality check was performed using Illumina Sequencing Analysis Viewer (SAV)
671 software. Demultiplexing was performed with Illumina Bcl2fastq2 program (version 2.19.1.403; Illumina
672 Inc.).

Differential gene expression analysis

674 The quality of the reads was verified using FastQC (Andrews, 2010) before reads were aligned to the
675 mm10 reference genome (GRCm39) using STAR (Dobin et al., 2013) with the following arguments: --
676 readFilesCommand zcat --outSAMtype BAM SortedByCoordinate --quantMode GeneCounts --
677 outFilterMismatchNmax 5 --outFilterMultimapNmax 1. The quality of the resulting alignments was
678 assessed using QualiMap (García-Alcalde et al., 2012). The Python package HTseq was used for gene counts
679 (Anders et al., 2015) using the following arguments: --stranded=no --idattr=gene_id --type=exon --
680 mode=union -r pos --format=bam. Output files were filtered to remove genes with low count (≤10) then
681 were used for differential gene expression analysis using DESeq2 (Love et al., 2014). The negative binomial
682 regression model of ComBat-seq was used to correct unwanted batch effects (Zhang et al., 2020). For a
683 gene to be classified as showing differential expression between treated and untreated cells, a threshold
684 of |log2(fold-change)|=0.5 and Benjamini-Hochberg adjusted p-value ≤0.05 had to be met.

Gene Ontology (GO) Analysis

686 Lists of differentially expressed genes were generated from read counts using DESeq2 Bioconductor
687 package (Love et al., 2014). Enrichment of GO terms in lists of up- and downregulated genes (|log2(fold-
688 change)|=0.25) was determined using g:Profiler (Raudvere et al., 2019). Redundant GO terms were
689 removed using reduce + visualize gene ontology (REVIGO) (Supek et al., 2011). Terms were included if the
690 fold enrichment (frequency of DEGs in each GO term to the frequency of total genes in GO terms) was
691 higher than 1.5 and if the Benjamini-Hochberg-adjusted p-value was less than 0.05. Plots for GO terms
692 were generated using a custom R script (Bonnot et al., 2019).

Statistical Methods

694 Statistical analyses, when not otherwise specified, were performed using GraphPad Prism 9
695 software. For significance testing, two-tailed T-tests were performed on pairwise comparisons. In all cases,
696 significance was determined by p-values less than or equal to 0.05. Each figure corresponds to at least
697 three independent biological repeats with three technical replicates (N=3, n=3), unless otherwise
698 specified. Number of asterisks on plots indicate level of statistical significance: *(p<0.05), **(p<0.01),
699 ***(p<0.001), ****(p<0.0001).

700 **REFERENCES**
701

Anders S, Pyl PT, Huber W. 2015. HTSeq-A Python framework to work with high-throughput sequencing
data. *Bioinformatics* **31**:166–169. doi:10.1093/bioinformatics/btu638

Andre CM, Hausman JF, Guerriero G. 2016. Cannabis sativa: The plant of the thousand and one
molecules. *Front Plant Sci* **7**:1–17. doi:10.3389/fpls.2016.00019

Andrews S. 2010. FastQC: a quality control tool for high throughput sequence data.
https://www.bioinformatics.babraham.ac.uk/projects/fastqc/

Barban S, Schulze HO. 1961. The Effects of 2-Deoxyglucose on the Growth and Metabolism of Cultured
Human Cells. *Journal of Biological Chemistry* **236**:1887–1890. doi:10.1016/s0021-9258(18)64100-6

Barton LJ, LeBlanc MG, Lehmann R. 2016. Finding their way: themes in germ cell migration. *Curr Opin Cell*
Biol **42**:128–137. doi:10.1016/j.ceb.2016.07.007

Bartova A, Birmingham MK. 1976. Effect of Δ9 tetrahydrocannabinol on mitochondrial NADH oxidase
activity. *Journal of Biological Chemistry* **251**:5002–5006. doi:10.1016/s0021-9258(17)33213-1

Bénard G, Massa F, Puente N, Lourenço J, Bellocchio L, Soria-Gómez E, Matias I, Delamarre A, Metna-
Laurent M, Cannich A, Hebert-Chatelain E, Mulle C, Ortega-Gutiérrez S, Martín-Fontecha M,
Klugmann M, Guggenhuber S, Lutz B, Gertsch J, Chaouloff F, López-Rodríguez ML, Grandes P,
Rossignol R, Marsicano G. 2012. Mitochondrial CB1 receptors regulate neuronal energy
metabolism. *Nat Neurosci* **15**:558–564. doi:10.1038/nn.3053

Bonnot T, Gillard M, Nagel D. 2019. A Simple Protocol for Informative Visualization of Enriched Gene
Ontology Terms. *Bio Protoc* **9**:1–9. doi:10.21769/bioprotoc.3429

Chabarria KC, Racusin DA, Antony KM, Kahr M, Suter MA, Mastrobattista JM, Aagaard KM. 2016.
Marijuana use and its effects in pregnancy. *Am J Obstet Gynecol* **215**:506.e1-506.e7.
doi:10.1016/j.ajog.2016.05.044

Chandra S, Radwan MM, Majumdar CG, Church JC, Freeman TP, ElSohly MA. 2019. New trends in
cannabis potency in USA and Europe during the last decade (2008–2017). *Eur Arch Psychiatry Clin*
Neurosci **269**:5–15. doi:10.1007/s00406-019-00983-5

Diaz Perez S V., Kim R, Li Z, Marquez VE, Patel S, Plath K, Clark AT. 2012. Derivation of new human
embryonic stem cell lines reveals rapid epigenetic progression in vitro that can be prevented by
chemical modification of chromatin. *Hum Mol Genet* **21**:751–764. doi:10.1093/hmg/ddr506

Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, Batut P, Chaisson M, Gingeras TR. 2013.
STAR: Ultrafast universal RNA-seq aligner. *Bioinformatics* **29**:15–21.
doi:10.1093/bioinformatics/bts635

Fuchs Weizman N, Wyse BA, Szaraz P, Defer M, Jahangiri S, Librach CL. 2021. Cannabis alters epigenetic
integrity and endocannabinoid signalling in the human follicular niche. *Human Reproduction* **00**:1–
10. doi:10.1093/humrep/deab104

Galve-Roperh I, Chiurchiù V, Díaz-Alonso J, Bari M, Guzmán M, Maccarrone M. 2013. Cannabinoid
receptor signaling in progenitor/stem cell proliferation and differentiation. *Prog Lipid Res* **52**:633–
650. doi:10.1016/j.plipres.2013.05.004

García-Alcalde F, Okonechnikov K, Carbonell J, Cruz LM, Götz S, Tarazona S, Dopazo J, Meyer TF, Conesa
A. 2012. Qualimap: Evaluating next-generation sequencing alignment data. *Bioinformatics* **28**:2678–
2679. doi:10.1093/bioinformatics/bts503

Gell JJ, Liu W, Sosa E, Chialastri A, Hancock G, Tao Y, Wamaitha SE, Bower G, Dey SS, Clark AT. 2020. An
Extended Culture System that Supports Human Primordial Germ Cell-like Cell Survival and Initiation
of DNA Methylation Erasure. *Stem Cell Reports* **14**. doi:10.1016/j.stemcr.2020.01.009

Gómez M, Hernández M, Fernández-Ruiz J. 2007. The activation of cannabinoid receptors during early
postnatal development reduces the expression of cell adhesion molecule L1 in the rat brain. *Brain*
Res **1145**:48–55. doi:10.1016/j.brainres.2007.01.102

Hayashi K, Ohta H, Kurimoto K, Aramaki S, Saitou M. 2011. Reconstitution of the Mouse Germ Cell
Specification Pathway in Culture by Pluripotent Stem Cells. *Cell* **146**:519–532.
doi:10.1016/j.cell.2011.06.052

Hayashi Y, Otsuka K, Ebina M, Igarashi Kaori, Takehara A, Matsumoto M, Kanai A, Igarashi Kazuhiko, Soga
T, Matsui Y. 2017. Distinct requirements for energy metabolism in mouse primordial germ cells and
their reprogramming to embryonic germ cells. *Proc Natl Acad Sci U S A* **114**:8289–8294.
doi:10.1073/pnas.1620915114

Hikabe O, Hamazaki N, Nagamatsu G, Obata Y, Hirao Y, Hamada N, Shimamoto S, Imamura T, Nakashima
K, Saitou M, Hayashi K. 2016. Reconstitution in vitro of the entire cycle of the mouse female germ
line. *Nature* **539**:299–303. doi:10.1038/nature20104

Hocaoglu H, Wang L, Yang M, Yue S, Sieber M. 2021. Heritable shifts in redox metabolites during
mitochondrial quiescence reprogramme progeny metabolism. *Nat Metab* **3**:1259–1274.
doi:10.1038/s42255-021-00450-3

Hunault CC, Mensinga TT, Leenders MEC, Meulenbelt J. 2008. Delta-9-tetrahydrocannabinol (THC) serum
concentrations and pharmacological effects in males after smoking a combination of tobacco and
cannabis containing up to 69 mg THC. *Psychopharmacology (Berl)* **201**:171–181.
doi:10.1007/s00213-008-1260-2

Keimpema E, MacKie K, Harkany T. 2011. Molecular model of cannabis sensitivity in developing neuronal
circuits. *Trends Pharmacol Sci* **32**:551–561. doi:10.1016/j.tips.2011.05.004

Kinoshita M, Barber M, Mansfield W, Cui Y, Spindlow D, Stirparo GG, Dietmann S, Nichols J, Smith A.
2021. Capture of Mouse and Human Stem Cells with Features of Formative Pluripotency ll ll
Capture of Mouse and Human Stem Cells with Features of Formative Pluripotency. *Stem Cell*
28:453-471.e8. doi:10.1016/j.stem.2020.11.005

Kittler JT, Grigorenko E V., Clayton C, Zhuang SY, Bundey SC, Trower MM, Wallace D, Hampson R,
Deadwyler S. 2000. Large-scale analysis of gene expression changes during acute and chronic

exposure to Δ9-THC in rats. *Physiol Genomics* **2000**:175–185.
doi:10.1152/physiolgenomics.2000.3.3.175

Komeya M, Sato T, Ogawa T. 2018. In vitro spermatogenesis: A century-long research journey still half
way. *Reprod Med Biol* **17**:407–420. doi:10.1002/rmb2.12225

Kurimoto K, Saitou M. 2018. Epigenome regulation during germ cell specification and development from
pluripotent stem cells. *Curr Opin Genet Dev* **52**:57–64. doi:10.1016/j.gde.2018.06.004

Lo JO, Hedges JC, Girardi G. 2022. Impact of cannabinoids on pregnancy, reproductive health, and
offspring outcomes. *Am J Obstet Gynecol* **227**:571–581. doi:10.1016/j.ajog.2022.05.056

Locasale JW, Cantley LC. 2011. Metabolic flux and the regulation of mammalian cell growth. *Cell Metab*
14:443–451. doi:10.1016/j.cmet.2011.07.014

Lojpur T, Easton Z, Raez-Villanueva S, Laviolette S, Holloway AC, Hardy DB. 2019. Δ9-
Tetrahydrocannabinol leads to endoplasmic reticulum stress and mitochondrial dysfunction in
human BeWo trophoblasts. *Reproductive Toxicology* **87**:21–31. doi:10.1016/j.reprotox.2019.04.008

Love MI, Huber W, Anders S. 2014. Moderated estimation of fold change and dispersion for RNA-seq
data with DESeq2. *Genome Biol* **15**:1–21. doi:10.1186/s13059-014-0550-8

Lunt SY, Vander Heiden MG. 2011. Aerobic glycolysis: Meeting the metabolic requirements of cell
proliferation. *Annu Rev Cell Dev Biol* **27**:441–464. doi:10.1146/annurev-cellbio-092910-154237

Luo Y, Yu Y. 2022. Research Advances in Gametogenesis and Embryogenesis Using Pluripotent Stem
Cells. *Front Cell Dev Biol* **9**:1–14. doi:10.3389/fcell.2021.801468

Mennis J, Stahler GJ, Mason MJ. 2023. Cannabis Legalization and the Decline of Cannabis Use Disorder
(CUD) Treatment Utilization in the US. *Curr Addict Rep*. doi:10.1007/s40429-022-00461-4

Miller ML, Chadwick B, Dickstein DL, Purushothaman I, Egervari G, Rahman T, Tessereau C, Hof PR,
Roussos P, Shen L, Baxter MG, Hurd YL. 2019. Adolescent exposure to Δ 9 -tetrahydrocannabinol
alters the transcriptional trajectory and dendritic architecture of prefrontal pyramidal neurons. *Mol*
Psychiatry **24**:588–600. doi:10.1038/s41380-018-0243-x

Murphy SK, Itchon-Ramos N, Visco Z, Huang Z, Grenier C, Schrott R, Acharya K, Boudreau MH, Price TM,
Raburn DJ, Corcoran DL, Lucas JE, Mitchell JT, McClernon FJ, Cauley M, Hall BJ, Levin ED, Kollins SH.
2018. Cannabinoid exposure and altered DNA methylation in rat and human sperm. *Epigenetics*
13:1208–1221. doi:10.1080/15592294.2018.1554521

Nichols J, Smith A. 2009. Naive and Primed Pluripotent States. *Cell Stem Cell* **4**:487–492.
doi:10.1016/j.stem.2009.05.015

Ohinata Y, Sano M, Shigeta M, Yamanaka K, Saitou M. 2008. A comprehensive, non-invasive visualization
of primordial germ cell development in mice by the Prdm1-mVenus and Dppa3-ECFP double
transgenic reporter. *Reproduction* **136**:503–514. doi:10.1530/REP-08-0053

Okamura D, Kimura T, Nakano T, Matsui Y. 2003. Cadherin-mediated cell interaction regulates germ cell
determination in mice. *Development* **130**:6423–6430. doi:10.1242/dev.00870

Osborne AJ, Pearson JF, Noble AJ, Gemmell NJ, Horwood LJ, Boden JM, Benton MC, Macartney-Coxson
DP, Kennedy MA. 2020. Genome-wide DNA methylation analysis of heavy cannabis exposure in a
New Zealand longitudinal cohort. *Transl Psychiatry* **10**:1–10. doi:10.1038/s41398-020-0800-3

Pacher P, Kogan NM, Mechoulam R. 2020. Beyond THC and Endocannabinoids. *Annu Rev Pharmacol*
Toxicol **60**. doi:10.1146/annurev-pharmtox-010818-021441

Pacifici R, Pichini S, Pellegrini M, Rotolo MC, Giorgetti R, Tagliabracci A, Busardò FP, Huestis MA. 2019.
THC and CBD concentrations in blood , oral fluid and urine following a single and repeated
administration of " light cannabis ." *Clin Chem Lab Med* 1–8.

Pang Z, Zhou G, Ewald J, Chang L, Hacariz O, Basu N, Xia J. 2022. Using MetaboAnalyst 5.0 for LC–HRMS
spectra processing, multi-omics integration and covariate adjustment of global metabolomics data.
Nat Protoc **17**:1735–1761. doi:10.1038/s41596-022-00710-w

Paria BC, Das SK, Dey SK. 1995. The preimplantation mouse embryo is a target for cannabinoid ligand-
receptor signaling. *Proc Natl Acad Sci U S A* **92**:9460–9464. doi:10.1073/pnas.92.21.9460

Pendergrass W, Wolf N, Pool M. 2004. Efficacy of MitoTracker Green™ and CMXRosamine to measure
changes in mitochondrial membrane potentials in living cells and tissues. *Cytometry Part A* **61**:162–
169. doi:10.1002/cyto.a.20033

Preet A, Ganju RK, Groopman JE. 2008. Δ9-Tetrahydrocannabinol inhibits epithelial growth factor-
induced lung cancer cell migration in vitro as well as its growth and metastasis in vivo. *Oncogene*
27:339–346. doi:10.1038/sj.onc.1210641

Prini P, Rusconi F, Zamberletti E, Gabaglio M, Penna F, Fasano M, Battaglioli E, Parolaro D, Rubino T.
2017. Adolescent THC exposure in female rats leads to cognitive deficits through a mechanism
involving chromatin modifications in the prefrontal cortex. *J Psychiatry Neurosci* **43**:170082.
doi:10.1503/jpn.170082

Raudvere U, Kolberg L, Kuzmin I, Arak T, Adler P, Peterson H, Vilo J. 2019. G:Profiler: A web server for
functional enrichment analysis and conversions of gene lists (2019 update). *Nucleic Acids Res*
47:W191–W198. doi:10.1093/nar/gkz369

Rinaldi-Carmona M, Barth F, Héaulme M, Shire D, Calandra B, Congy C, Martinez S, Maruani J, Néliat G,
Caput D, Ferrara P, Soubrié P, Brelière JC, Le Fur G. 1994. SR141716A, a potent and selective
antagonist of the brain cannabinoid receptor. *FEBS Lett* **350**:240–244. doi:10.1016/0014-
5793(94)00773-X

Schrott R, Murphy SK. 2020. Cannabis use and the sperm epigenome: a budding concern? *Environ*
Epigenet **6**:1–10. doi:10.1093/eep/dvaa002

Smith A, Kaufman F, Sandy MS, Cardenas A. 2020. Cannabis Exposure During Critical Windows of
Development: Epigenetic and Molecular Pathways Implicated in Neuropsychiatric Disease. *Curr*
Environ Health Rep **7**:325–342. doi:10.1007/s40572-020-00275-4

Sohni A, Bartoccetti M, Khoueiry R, Spans L, Vande Velde J, De Troyer L, Pulakanti K, Claessens F, Rao S,
Koh KP. 2015. Dynamic Switching of Active Promoter and Enhancer Domains Regulates Tet1 and

Tet2 Expression during Cell State Transitions between Pluripotency and Differentiation . *Mol Cell*
Biol **35**:1026–1042. doi:10.1128/mcb.01172-14

Substance Abuse and Mental Health Services Administration. 2020. Key substance use and mental health
indicators in the United States: Results from the 2019 National Survey on Drug Use and Health.

Supek F, Bošnjak M, Škunca N, Šmuc T. 2011. Revigo summarizes and visualizes long lists of gene
ontology terms. *PLoS One* **6**. doi:10.1371/journal.pone.0021800

Szutorisz H, Hurd YL. 2018. High times for cannabis: Epigenetic imprint and its legacy on brain and
behavior. *Neurosci Biobehav Rev* **85**:93–101. doi:10.1016/j.neubiorev.2017.05.011

Takeda S, Yamaori S, Motoya E, Matsunaga T, Kimura T, Yamamoto I, Watanabe K. 2008. Δ9-
Tetrahydrocannabinol enhances MCF-7 cell proliferation via cannabinoid receptor-independent
signaling. *Toxicology* **245**:141–146. doi:10.1016/j.tox.2007.12.019

Tempany JC, Zhou JHS, Hodgkin PD, Bryant VL. 2018. Superior properties of CellTrace Yellow™ as a
division tracking dye for human and murine lymphocytes. *Immunol Cell Biol* **96**:149–159.
doi:10.1111/imcb.1020

U.N. Office on Drugs and Crime. 2022. 2022 World Drug Report.

Verdikt R, Allard P. 2021. Metabolo-epigenetics: the interplay of metabolism and epigenetics during early
germ cells development. *Biol Reprod* **105**:616–624. doi:10.1093/biolre/ioab118

Verdikt R, Armstrong AA, Allard P. 2022. Transgenerational inheritance and its modulation by
environmental cuesCurrent Topics in Developmental Biology. Elsevier Inc. pp. 1–46.
doi:10.1016/bs.ctdb.2022.10.002

Volkow ND, Han B, Compton WM, McCance-Katz EF. 2019. Self-reported Medical and Nonmedical
Cannabis Use Among Pregnant Women in the United States. *JAMA - Journal of the American*
Medical Association **322**:167–169. doi:10.1001/jama.2018.20391

Watson CT, Szutorisz H, Garg P, Martin Q, Landry JA, Sharp AJ, Hurd YL. 2015. Genome-Wide DNA
Methylation Profiling Reveals Epigenetic Changes in the Rat Nucleus Accumbens Associated with
Cross-Generational Effects of Adolescent THC Exposure. *Neuropsychopharmacology* **40**:2993–3005.
doi:10.1038/npp.2015.155

Weinberger L, Ayyash M, Novershtern N, Hanna JH. 2016. Dynamic stem cell states: Naive to primed
pluripotency in rodents and humans. *Nat Rev Mol Cell Biol* **17**:155–169. doi:10.1038/nrm.2015.28

Yang X, Bam M, Nagarkatti PS, Nagarkatti M. 2016. RNA-seq analysis of δ9-tetrahydrocannabinol-treated
T cells reveals altered gene expression profiles that regulate immune response and cell
proliferation. *Journal of Biological Chemistry* **291**:15460–15472. doi:10.1074/jbc.M116.719179

Young-Wolff KC, Tucker L, Alexeeff S, Anne M, Conway A, Weisner C, Goler N. 2018. Among Pregnant
Females in California From 2009 – 2016. *JAMA - Journal of the American Medical Association*
318:2490–2491. doi:10.1001/jama.2017.17225.Trends

Zhang J, Zhao J, Dahan P, Lu V, Zhang C, Li H, Teitell MA. 2018. Metabolism in Pluripotent Stem Cells and
Early Mammalian Development. *Cell Metab* **27**:332–338. doi:10.1016/j.cmet.2018.01.008

Zhang Y, Parmigiani G, Johnson WE. 2020. ComBat-seq: Batch effect adjustment for RNA-seq count data.
NAR Genom Bioinform **2**:1–10. doi:10.1093/nargab/lqaa078

 FIGURE LEGENDS

Figure 1: Δ9-THC exposure provokes the proliferation of ESCs but not EpiLCs

(A) Diagram illustrating Δ9-THC exposure scheme and experimental strategy. bFGF: basic fibroblast growth

factor, ESCs, embryonic stem cells; EpiLCs, epiblast-like cells; LIF, leukemia inhibitory factor. **(B, E, H)**

Whisker boxplot indicating the median cellular viability of stem cells exposed to the different Δ9-THC doses

and associated errors. **(C, F, I)** Whisker boxplot indicating the median number of viable cells exposed to

the different Δ9-THC doses indicated and associated errors. **(D, G, J)** Whisker boxplot indicating the median

percentage of BrdU-stained cells exposed to the different Δ9-THC doses and associated errors. ESCs

exposed cells are presented in (B, C and D). EpiLCs exposed cells deriving from unexposed ESCs are

presented in (E, F and G). EpiLCs exposed cells deriving from exposed ESCs are presented in (H, I and J).

At least three independent biological repeats with three technical replicates (N=3, n=3). Statistical

significance: *($p<0.05$), **($p<0.01$), ***($p<0.001$), ****($p<0.0001$).

Figure 2: Implication of the CB1 receptor in the proliferative phenotype.

(A) Western blot analysis of transmembrane protein extracts of ESCs or EpiLCs. Antibodies raised against

CB1 or β-actin serving as a loading control were used for immunoblotting. **(B)** Quantification of the gel

presented in (A) was done using Image Studio (version 5.2). **(C)** Whisker boxplot indicating the median

cellular viability of stem cells exposed to the different Δ9-THC and rimonabant doses indicated and their

associated errors. **(D)** The median numbers of viable cells exposed to the different Δ9-THC and rimonabant

doses indicated were normalized to their own control (+/- rimonabant). Median and associated errors

were plotted in whisker boxplots. At least three independent biological repeats with three technical

replicates (N=3, n=3). Statistical significance: **($p<0.01$), ****($p<0.0001$).

Figure 3: Δ9-THC exposure provokes an increase in glycolytic rates in ESCs and EpiLCs.

(A) Diagram illustrating Δ9-THC exposure scheme and experimental strategy. **(B)** The NAD(P)+/NADPH

ratio of stem cells exposed to the different Δ9-THC doses was normalized to the one measured in the

mock-treated condition. Median and associated errors were plotted in whisker boxplots. **(C)** Mean

fluorescence intensity (MFI) associated with the Mitotracker CMXRos stain was normalized to the one

measured in the mock-treated condition. Median and associated errors were plotted in whisker boxplots.

(D) Median and associated error of the maximal extracellular acidification rate (ECAR) measured in cells

exposed to the different Δ9-THC doses and normalized to the protein content was plotted in whisker

boxplots. **(E)** Median and associated error of the maximal oxygen consumption rate (OCR) measured in

cells exposed to the different Δ9-THC doses and normalized to the protein content was plotted in whisker

boxplots. For (B and C), 5 technical repeats of 3 biological repeats (n=15) were plotted. One same

representative experiment out of three independent experiments was used to plot results in (D and E).

Statistical significance: *($p<0.05$), **($p<0.01$), ***($p<0.001$), ****($p<0.0001$).

Figure 4: Δ9-THC-induced glycolysis sustain anabolism and ESCs proliferation

(A) Diagram illustrating Δ9-THC exposure scheme and experimental strategy. **(B)** PCA of the metabolomics

profiling of either ESCs or EpiLCs mock-exposed or exposed to 100nM Δ9-THC. **(C)** Venn diagram showing

the overlap in upregulated metabolites following Δ9-THC exposure in ESCs and EpiLCs. **(D and E)** KEGG

metabolite sets enrichment analysis for upregulated metabolites in ESCs and EpiLCs, respectively,

performed by MetaboAnalyst[38]. KEGG, Kyoto Encyclopedia of Genes and Genomes. **(F)** Whisker boxplot
indicating the median cellular viability of stem cells exposed to 100nM of Δ9-THC and 10mM of 2-DG, as
indicated, and their associated errors. **(G)** The median numbers of viable cells exposed to 100nM of Δ9-
THC and 10mM of 2-DG, as indicated, were normalized to their own control (+/- 2-DG). Median and
associated errors were plotted in whisker boxplots. **(H)** The NAD(P)+/NADPH ratio of stem cells exposed
to 100nM of Δ9-THC and 10mM of 2-DG, as indicated, was normalized to the one measured in the mock-
treated condition (+/- 2-DG). Median and associated errors were plotted in whisker boxplots. At least three
independent biological repeats with three technical replicates (N=3, n=3). Statistical significance:
*(p<0.05), **(p<0.01), ***(p<0.001), ****(p<0.0001).

Figure 5: Metabolic changes following Δ9-THC exposure in ESCs are transcriptionally encoded.
(A) Diagram illustrating Δ9-THC exposure scheme and experimental strategy. **(B)** PCA of the
transcriptomics profiling of either ESCs or EpiLCs mock-exposed or exposed to 100nM Δ9-THC. **(C and D)**
Volcano plot in ESCs and EpiLCs, respectively, showing significance [expressed in $\log_{10}$(adjusted p-value or
false-discovery rate, FDR)] versus fold-change [expressed in log2(fold-change, FC)]. Thresholds for
significance (adjusted p-value≤0.05) and gene expression fold-change [|(log2(FC)|>0.25 or |log2(FC)|>0.5]
are shown as dashed lines. Color code is as follows: log2(FC)>0.5 in red, log2(FC)>0.25 in orange, log2(FC)>0
in light orange, log2(FC)<0 in light blue, log2(FC)>-0.25 in blue, log2(FC)>0.5 in dark blue and p-value<0.01
in pink. **(E)** Gene ontology (GO) terms associated with up- and downregulated DEGs [|(log2(FC)|>0.25 and
p<0.01)] in ESCs and EpiLCs as determined by g:Profiler[55]. **(F)** Joint pathway analysis performed by the
multi-omics integration tool of MetaboAnalyst[38]. The p-values were weighted based on the proportions of
genes and metabolites at the individual pathway level.

Figure 6: PGCLCs deriving from ESCs and EpiLCs exposed to 100nM of Δ9-THC proliferate.
(A) Diagram illustrating Δ9-THC exposure scheme and experimental strategy. **(B)** Representative flow
contour plots showing distribution of live-gated events, gating strategy for Stella:CFP versus
Blimp1:mVenus and percentages of cells in each subpopulations for ESCs and EpiLCs exposed to the
different doses of Δ9-THC indicated. DN: double negative, SP: single positive, DP: double positive
subpopulations. **(C)** The percentage of events in the gates associated to each subpopulation was
normalized to the one measured in the mock-treated condition. Median and associated errors were
plotted in whisker boxplots independently for each subpopulation. **(D)** Representative histograms showing
CellTrace™ Yellow staining profile of cells arising from ESCs and EpiLCs exposed to the different doses of
Δ9-THC indicated. The Y-axis represents the average percentage of cells in each category of subpopulations
undividing (purple), undergoing 1 division (blue), 2 divisions (green) or 3 divisions (orange). One
representative experiment out of three is represented. Statistical significance: *(p<0.05), **(p<0.01),
(p<0.001), *(p<0.0001).

Figure 7: Δ9-THC exposure prior to specification increases mitochondrial respiration in PGCLCs.
(A) Diagram illustrating Δ9-THC exposure scheme and experimental strategy. **(B)** The NAD(P)+/NADPH
ratio of embryoid bodies arising from ESCs and EpiLCs exposed to 100nM of Δ9-THC was normalized to the
one measured in the mock-treated condition. Median and associated errors were plotted in whisker
boxplot. **(C)** Mean fluorescence intensity (MFI) associated with the Mitotracker CMXRos stain in each

subpopulation was normalized to the one measured in the mock-treated condition. Median and associated
errors were plotted in whisker boxplots. **(D)** PCA of the transcriptomics profiling of DP PGCLCs deriving
from ESCs and EpiLCs either mock-exposed or exposed to 100nM Δ9-THC. **(E)** Volcano plot in DP PGCLCs
showing significance [expressed in $\log_{10}$(adjusted p-value or false-discovery rate, FDR)] versus fold-change
[expressed in log2(fold-change, FC)]. Thresholds for significance and different enrichment ratios
[|(log2(FC)|>0.25 or |log2(FC)|>0.5] are shown as dashed lines. Color code is as follows: log2(FC)>0.5 in
red, log2(FC)>0.25 in orange, log2(FC)>0 in light orange, log2(FC)<0 in light blue, log2(FC)>-0.25 in blue,
log2(FC)>0.5 in dark blue and p-value<0.01 in pink. **(F and G)** Gene ontology (GO) terms associated with
up- and downregulated DEGs [|(log2(FC)|>0.25 and p<0.01)], respectively, as determined by g:Profiler[55].
Statistical significance: *(p<0.05), **(p<0.01).
980
981 **Figure 8: Metabolic impact of Δ9-THC exposure in pluripotent stem cells and primordial germ cells-like**
982 **cells.**
983 Diagram illustrating the impact of Δ9-THC exposure on stem cells metabolism.
984
985 **Supplementary Figure 1: Δ9-THC induces ESCs proliferation for as low as 1nM.**
986 **(A)** Whisker boxplot indicating the median cellular viability of ESCs exposed to the different Δ9-THC doses
987 and associated errors. **(B)** Whisker boxplot indicating the median number of viable cells exposed to the
988 different Δ9-THC doses indicated and associated errors. At least three independent biological repeats with
989 three technical replicates (N=3, n=3). Statistical significance: *(p<0.05), ***(p<0.001), ****(p<0.0001).
990
Supplementary Figure 2: Δ9-THC induces alteration in ESCs cell cycle.
(A) Representative flow contour plots showing distribution of BrdU-stained and DAPI-stained cells,
exposed to the different doses of Δ9-THC indicated. The frequency of events in each gate is indicated. **(B)**
The median percentage of events and associated errors for each cell cycle gate were plotted in histograms.
At least three independent biological repeats with three technical replicates (N=3, n=3). Statistical
significance: *(p<0.05), **(p<0.01).
997
Supplementary Figure 3: Δ9-THC exposure in male ESCs also provokes cell proliferation.
(A) Whisker boxplot indicating the median cellular viability of male ESCs (the R8 cell line, see Material and
Methods section) exposed to the different Δ9-THC doses and associated errors. **(B)** Whisker boxplot
indicating the median number of viable cells exposed to the different Δ9-THC doses indicated and
associated errors. At least three independent biological repeats with three technical replicates (N=3, n=3).
Statistical significance: *(p<0.05), **(p<0.01), ****(p<0.0001).
1004
1005 **Supplementary Figure 4: hESCs cell number decreases upon Δ9-THC exposure.**
1006 **(A)** Whisker boxplot indicating the median cellular viability of human embryonic stem cells continuously
1007 exposed to 100nM Δ9-THC doses over 6 days and associated errors. **(B)** Whisker boxplot indicating the
1008 median number of viable cells exposed to 100nM of Δ9-THC doses indicated and associated errors. For (A
1009 and B), 6 technical repeats of 2 biological repeats (n=12) were plotted. Statistical significance: **(p<0.01).
1010

Supplementary Figure 5: hESCS metabolism is slightly but significantly impacted by Δ9-THC exposure.
The NAD(P)+/NADPH ratio of hESCs exposed to 100nM of Δ9-THC was normalized to the one measured in the mock-treated condition. Median and associated errors were plotted in whisker boxplots. One representative experiment out of two independent experiments was used to plot results. Statistical significance: *($p<0.05$).

Supplementary Figure 6: Extracellular acidification rates and oxygen consumption rates in ESCs and EpiLCs upon Δ9-THC exposure.
(A and B) Traces were plotted for the extracellular acidification rate (ECAR) measurements in ESCs and EpiLCs, respectively, exposed to the different Δ9-THC doses indicated and normalized to the protein content. The oligomycin injection time is indicated by an arrow and allows to differentiate basal glycolytic rate from maximal glycolytic rate (when mitochondria are inhibited). The datapoints used in the main figure correspond to the first timepoint in the maximal glycolytic capacity section. **(C and D)** Traces were plotted for the oxygen consumption rate (OCR) measurements in ESCs and EpiLCs, respectively, exposed to the different Δ9-THC doses indicated and normalized to the protein content. The oligomycin, FCCP and AntimycinA/Rotenone injection times are indicated by arrows and allow to differentiate basal respiration from ATP-coupled respiration and maximal respiratory capacity. The datapoints used in the main figure correspond to the second timepoint in the maximal respiratory capacity section. FCCP: Carbonyl cyanide-p-trifluoromethoxyphenylhydrazone. Statistical significance: *($p<0.05$), **($p<0.01$), ****($p<0.0001$).

Supplementary Figure 7: Metabolite profiling in ESCs and EpiLCs upon Δ9-THC exposure.
(A and B) Heatmaps showing the log2 of the amount of each metabolite upregulated in ESCs and EpiLCs upon exposure to 100nM of Δ9-THC. The relative amounts of metabolites were normalized to the mean value across all samples for one same condition and to the number of viable cells harvested in parallel on a control plate. **(C)** Histograms showing the ratio of reduced to oxidized glutathione (GSH/GSSG) based on the amounts measured in the metabolomics profiling.

Supplementary Figure 8: Extracellular acidification rates and oxygen consumption rates in ESCs upon Δ9-THC and 2-DG exposure.
(A) Traces were plotted for the extracellular acidification rate (ECAR) measurements in ESCs exposed to 100nM of Δ9-THC and 10mM of 2-DG, as indicated, and normalized to the protein content. The oligomycin injection time is indicated by an arrow and allows to differentiate basal glycolytic rate from maximal glycolytic rate (when mitochondria are inhibited). **(B)** Traces were plotted for the oxygen consumption rate (OCR) measurements in ESCs exposed to 100nM of Δ9-THC and 10mM of 2-DG, as indicated, and normalized to the protein content. The oligomycin, FCCP and AntimycinA/Rotenone injection times are indicated by arrows and allow to differentiate basal respiration from ATP-coupled respiration and maximal respiratory capacity. FCCP: Carbonyl cyanide-p-trifluoromethoxyphenylhydrazone. Statistical significance: *($p<0.05$), **($p<0.01$).

Supplementary Figure 9: Δ9-THC exposure does not alter markers of pluripotency.
Gene expression profiles of markers for the inner cell mass (ICM) and epiblast. Histograms show the median and associated errors of normalized gene counts in each condition, as indicated.

1052

Supplementary Figure 10: Δ9-THC exposure alter the expression of some epigenetic modifiers.

Histograms show the median and associated errors of normalized gene counts in each condition, as indicated. Only genes with |(log2(FC)|>0.25 and p-value<0.01 from Supplementary Table 1 were plotted. Statistical significance: **(p<0.01), ***(p<0.001), ****(p<0.0001).

Supplementary Figure 11: PGCLCs gating and sorting strategy.

(A) Representative flow contour plots showing distribution of events and gating based on embryoid bodies dissociation. **(B and C)** Representative flow contour plots to isolate singlets based on width to height ratios on the side scatter and front scatter, respectively. **(D)** Gating strategy for Stella:CFP versus Blimp1:mVenus on the negative control, corresponding to embryoid bodies obtained in an induction medium without cytokines and BMPs (GK15 only). **(E)** Gating strategy for Stella:CFP versus Blimp1:mVenus on mock-treated cells, corresponding to embryoid bodies obtained in an induction medium complemented with cytokines and BMPs. DN: double negative, SP: single positive, DP: double positive subpopulations.

Supplementary Figure 12: Male PGCLCs deriving from ESCs and EpiLCs exposed to 100nM of Δ9-THC proliferate.

(A) Diagram illustrating Δ9-THC exposure scheme and experimental strategy. **(B)** Representative flow contour plots showing distribution of live-gated events, gating strategy for Stella:CFP versus Blimp1:mVenus and percentages of cells in each subpopulations for ESCs and EpiLCs exposed to 100nM of Δ9-THC. DN: double negative, SP: single positive, DP: double positive subpopulations. **(C)** The percentage of events in the gates associated to each subpopulation was normalized to the one measured in the mock-treated condition. Median and associated errors were plotted in whisker boxplots independently for each subpopulation. At least three independent biological repeats with three technical replicates (N=3, n=3). Statistical significance: *(p<0.05).

Supplementary Figure 13: No residual Δ9-THC is detected in day 5 embryoid bodies.

Intracellular levels of Δ9-THC were quantified by mass spectrometry in EpiLCs on the day of aggregate formation and in day 5 embryoid bodies (referred as to "EpiLCs" and "PGCLCs"). Histograms show the median and associated errors of two independent quantifications. Statistical significance: ****(p<0.0001).

1082

 Figure 2

1086   **Figure 3**

 Figure 4

Figure 6

1091   **Figure 7**

 Figure 8

A.

B.

1098   **Supplementary Figure 5**

 Supplementary Figure 6

A.

B.

     Supplementary Figure 9

 Supplementary Figure 10

DNA methylation

Histone deacetylation

Histone methylation

Histone phosphorylation

Histone ubiquitination

1105   **Supplementary Figure 12**

1106

 Supplementary Figure 13

1108