Metabolic memory of Δ9-tetrahydrocannabinol exposure in pluripotent stem cells and primordial germ cells-like cells

Roxane Verdikt[1], Abigail A. Armstrong[2], Jenny Cheng[3], Xia Yang[4,5], and Patrick Allard[1,6]*

[1]Institute for Society and Genetics, University of California, Los Angeles, Los Angeles, CA 90095, USA
[2]Department of Obstetrics/Gynecology and Reproductive Endocrinology and Infertility, University of California, Los Angeles, CA, USA
[3]Molecular, Cellular, and Integrative Physiology Graduate Program, University of California, Los Angeles, Los Angeles, CA 90095, USA
[4]Integrative Biology and Physiology Department, University of California, Los Angeles, CA, 90095, USA
[5] Department of Molecular and Medical Pharmacology, University of California, Los Angeles, Los Angeles, CA 90095, USA
[6] Molecular Biology Institute, University of California, Los Angeles, Los Angeles, CA 90095, USA

* Corresponding author:
Patrick Allard, Boyer Hall, 611 Charles E Young Dr E, University of California, Los Angeles, Los Angeles, 90095. Email: pallard@ucla.edu

ABSTRACT

Cannabis, the most consumed psychoactive drug in the world, is increasingly used by pregnant women. However, while cannabinoid receptors are expressed in the early embryo, the impact of phytocannabinoids exposure on early embryonic processes is lacking. Here, we leverage a stepwise in vitro differentiation system that captures early embryonic developmental cascade to investigate the impact of exposure to the most abundant phytocannabinoid, Δ9-tetrahydrocannabinol (Δ9-THC). We demonstrate that Δ9-THC increases the proliferation of naïve mouse embryonic stem cells (ESCs) but not of their primed counterpart. Surprisingly, this increased proliferation, dependent on the CB1 receptor binding, is only associated with moderate transcriptomic changes. Instead, Δ9-THC capitalizes on ESCs' metabolic bivalence by increasing their glycolytic rates and anabolic capabilities. A memory of this metabolic rewiring is retained throughout differentiation to Primordial Germ Cell-Like Cells in the absence of direct exposure and is associated with an alteration of their transcriptional profile. These results represent the first in-depth molecular characterization of the impact of Δ9-THC exposure on early developmental stages.

KEYWORDS
cannabis, Δ9-THC, metabolism, embryonic stem cells, primordial germ cells.

INTRODUCTION

Cannabis is the most widely used psychoactive drug in the world[1]. In the United States, an estimated 49.6 million people, roughly 18% of the population, consumed cannabis at least once in 2020, with indications that these numbers will likely increase in the coming years as attitudes and regulations change[2,3]. In particular, between 7-12% of expecting women report cannabis use, predominantly during the first trimester to alleviate the symptoms of morning sickness[4–6]. These statistics indicate that a significant number of developing embryos are exposed to cannabis, with limited knowledge of the biological repercussions of such exposure.

Among the several hundred unique phytocannabinoids present in Cannabis sativa, (−)-trans-Δ9-tetrahydrocannabinol (Δ9-THC) is chiefly responsible for the psychoactivity of cannabis[7]. As a result, the level of Δ9-THC in recreational cannabis has increased over the last 10 years and now commonly accounts for 20% of total compounds[8]. The psychoactive effects of Δ9-THC arise from its binding and subsequent activation of the G protein-coupled cannabinoid receptors CB1 largely expressed in the central nervous system[9]. In this context, Δ9-THC exposure has been shown to durably alter metabolic, transcriptional and epigenetic programs in the brain[10–13]. While over the last decades, significant attention has been paid to Δ9-THC's neurological effects, there is also evidence, albeit more limited, of its impact on reproductive functions[14]. Data shows CB1 expression in the male and the female reproductive tracts, in the pre-implantation embryo and in the placenta[14,15]. In animal models as well as in humans, exposure to cannabis is associated with reduced fertility, decreased testis weight and sperm count, and impairment of embryo implantation[14]. In males, these effects are correlated with an alteration of the sperm transcriptome and epigenome[16–18]. Epidemiological evidence also indicates that Δ9-THC exposure is associated with long-lasting adverse effects, with exposures in parents affecting the offspring[13,19]. Despite this accumulating evidence, the molecular impacts and mechanisms of Δ9-THC exposure at the earliest stages of development remain to be determined.

Progression through states of pluripotency is controlled by metabolic reprogramming in the early mammalian embryo[20,21]. Accordingly, cultured pluripotent stem cells (PSCs) exhibiting different developmental potentials are marked by specific metabolic signatures, similar to the ones displayed by their in vivo counterparts in the embryo. For instance, mouse embryonic stem cells (ESCs) are naïve PSCs that are functionally equivalent to the inner cell mass (ICM) of the E3.5 preimplantation mouse blastocyst[22]. The extended developmental potential of mouse ESCs is associated with their metabolic bivalence, as these cells rely on both glycolysis and oxidative phosphorylation for energy production. Differentiation of naïve ESCs into primed PSCs such as epiblast-like cells (EpiLCs) is accompanied by an important metabolic shift towards aerobic glycolysis, in link with a highly-proliferative phenotype and a more restricted developmental potential[20,21,23]. Primordial germ cells (PGCs), the embryonic precursors of gametes in metazoans[24], are considered dormant totipotent cells because they possess the unique ability to reacquire totipotency upon fertilization[23]. In the mouse embryo, PGCs arise around embryonic day 7.5 (E7.5) from a subset of primed PSCs in the epiblast[24]. Progressive increase in oxidative phosphorylation correlates with the specification and differentiation from epiblast PSCs towards PGCs, a process that can be replicated in vitro by inducing PGC-like cells (PGCLCs) from EpiLCs[25]. In particular, the extensive metabolic, transcriptional, and epigenetic reprogramming that PGCs undergo during their development has been proposed to be uniquely sensitive to environmental insults, with potential consequences in the offspring[26].

Here, we deployed this in vitro differentiation system to investigate the impact of Δ9-THC
exposure on early developmental stages. We demonstrate that exposure of ESCs and EpiLCs to
Δ9-THC durably alters their metabolome. We reveal that, in the absence of continuous exposure,
metabolic memory of Δ9-THC is passed onto the PGCLCs stage leading to transcriptional defects
in these cells. Together, our findings highlight the role of metabolic reprogramming as a
mechanism for early developmental Δ9-THC exposure.
90
RESULTS
Δ9-THC induces cellular proliferation of mouse embryonic stem cells but not of mouse
epiblast-like cells.
To model the impact of early Δ9-THC exposure on early embryonic events, we first tested
three distinct developmental windows: 1) exposure of ESCs, 2) exposure of EpiLCs and 3)
combined ESCs+EpiLCs exposure (Figure 1A). Cells were either exposed to the vehicle (mock)
or exposed to Δ9-THC in a wide dose range of 10nM-100μM, corresponding to the reported
physiologically-relevant concentrations of Δ9-THC in cannabis users[26–28].
The viability of ESCs exposed to increasing concentrations of Δ9-THC for 48h was not
significantly altered until the maximal dose of 100μM, subsequently serving as a positive control,
at which only 13.17% of cells remained alive (Figure 1B, p<0.0001, unpaired T-test). While no
significant changes in viability were observed between 10nM and 1μM Δ9-THC, the number of
viable ESCs significantly increased by 1.69, 1.52 and 1.28-fold, respectively, compared to the
mock-treated condition (Figure 1C, p=0.002, p=0.01 and p=0.03 for 10nM, 100nM and 1μM of Δ9-
THC, unpaired T-test). To determine whether this increased number of viable cells recovered after
Δ9-THC exposure was due to higher proliferation, we performed bromodeoxyuridine (BrdU)
labeling experiments. Exposed cells were pulsed with BrdU for 30 minutes, and its incorporation
in actively dividing cells was measured by flow cytometry. The percentage of BrdU-positive ESCs
significantly increased between 10nM and 10μM of Δ9-THC compared to the mock-treated
condition (Figure 1D, p=0.01, p=0.001, p=0.05 and p=0.01 for 10nM, 100nM, 1μM and 10μM of
Δ9-THC, unpaired T-test).
Next, we derived EpiLCs from unexposed ESCs and performed the same dose-response
experiments. Akin to ESCs, Δ9-THC exposure in EpiLCs did not significantly alter cellular viability
until the dose of 100μM (Figure 1E, 11.56% of viable cells, p<0.0001, unpaired T-test). However,
contrary to ESCs, Δ9-THC exposure in EpiLCs did not significantly increase the number of viable
cells nor the percentage of BrdU-positive cells (Figure 1F and Figure 1G). For the dose of 10μM
of Δ9-THC, viable EpiLCs numbers and BrdU-positive EpiLCs decreased compared to the mock-
treated condition (Figure 1F, 1.53-fold decrease, p=0.0007, Figure 1G, 1.42-fold decrease,
p=0.0024, unpaired T-test).
Finally, when continuously exposing ESCs and EpiLCs, cell viability was more significantly
and negatively impacted, except at the dose of 100nM of Δ9-THC (Figure 1H). Deriving EpiLCs
from exposed ESCs and exposing them to Δ9-THC for 48h did not significantly affect either their
cell number nor their incorporation of BrdU (Figure 1I and Figure 1J), indicating that the increased
proliferation observed at the ESCs stage is not carried through the naïve-to-prime transition.
Together, the systematic testing of different exposure schemes of Δ9-THC in ESCs and
EpiLCs revealed that physiologically relevant doses of Δ9-THC (10nM-1μM) specifically stimulate

the proliferation of ESCs, but not of EpiLCs, whether the latter are derived from exposed ESCs or
not.
129
Expression of the CB1 receptor does not explain differences in proliferative outcomes.
We next sought to understand the source of variation in proliferative outcomes in response
to Δ9-THC between naïve mouse embryonic stem cells and primed pluripotent epiblast-like cells.
Such differential effects have been previously reported with Δ9-THC eliciting the proliferation of
neural progenitors[29] and of human breast carcinoma cell lines[30] but suppressing the proliferation
of activated CD4[+] T cells[31] and of non-small cell lung cancer cells[32]. In these studies, the
differential expression of cannabinoid receptors at the cell surface was proposed to primarily
mediate the variation in cellular outcomes.
We therefore first tested whether expression levels of CB1 varied between ESCs and
EpiLCs. Western-blot analysis of membrane proteins revealed however that CB1 was expressed
at the same levels at the cell surface of both ESCs and EpiLCs (Figure 2A and 2B). We next
determined whether the Δ9-THC-induced proliferative phenotype in ESCs was due to the
engagement of the CB1 cannabinoid receptor. To do so, ESCs were pretreated for 1h with 1µM
of SR141716 (also known as rimonabant, a specific CB1 blocker[33]) then exposed to 100nM or
100µM of Δ9-THC for 48h. Rimonabant pre-treatment did not significantly alter the viability of
ESCs compared to conditions exposed to Δ9-THC only (Figure 2C) but abolished Δ9-THC-
induced ESCs increased cell number at 100nM Δ9-THC (Figure 2D, 1.53-fold decrease,
p<0.0001, when comparing 100nM of Δ9-THC +/- 1µM of SR141716, unpaired T-test). Notably,
SR141716 pre-treatment, while not altering cell viability, reduced cell number compared to control,
suggesting a basal role for CB1 in promoting proliferation.
Thus, the expression of CB1 at the cell surface does not explain the differential impact of
Δ9-THC on ESC and EpiLC proliferation even if CB1 engagement is a required event for this effect
in ESCs.
153
Δ9-THC exposure increases glycolysis in ESCs and EpiLCs.
In the central nervous system, Δ9-THC is a known metabolic perturbator which increases
bioenergetic metabolism[12,34]. As mentioned above, the transition of naïve ESCs into the primed
state of EpiLCs is accompanied by a switch to glycolysis for energy production [20,23]. Thus, to
capture the impact of Δ9-THC at every point of their transition between metabolic states, we used
the continuous exposure scheme of ESCs and EpiLCs outlined in Figure 1H-J. Similar to our other
exposure schemes, at lower Δ9-THC doses, the proliferation of ESCs was observed but not of
EpiLCs. We performed these exposures in a wide Δ9-THC dose range (10nM-10µM) followed by
bioenergetics assessment (Figure 3A).
First, we assessed the global energy metabolism of exposed cells by measuring the
nicotinamide adenine dinucleotide (phosphate) couple ratios (NAD(P)+/NAD(P)H) using the WST-
1 assay. In ESCs, the ratio of NAD(P)+/NAD(P)H significantly increased 1.57, 1.54, 1.29 and 1.38
-fold, for 10nM, 100nM, 1µM and 10µM of Δ9-THC, respectively, compared to the mock-treated
condition (Figure 3B, p<0.0001, p<0.0001, p=0.03, and p=0.0003 for 10nM, 100nM, 1µM and
10µM of Δ9-THC, unpaired T-test). In contrast, no significant increase was observed in
NAD(P)+/NAD(P)H ratios in exposed EpiLCs (Figure 3B). Consistent with the impact of continuous
Δ9-THC exposure on EpiLCs viability (Figure 1H), the NAD(P)+/NAD(P)H ratios significantly

decreased at 10µM of Δ9-THC in EpiLCs (Figure 3B, 59% decrease for 10µM of Δ9-THC
compared to the mock-treated condition, p<0.0001, unpaired T-test).

Because the elevated NAD(P)+/NAD(P)H levels in Δ9-THC-exposed ESCs could indicate
increased mitochondrial activity in the context of oxidative phosphorylation[35], we next studied
changes in mitochondrial membrane potential of exposed cells using the Mitotracker CMXRos
fluorescent dye[36]. A significant increase in mean fluorescence intensity (MFI) associated with the
mitochondrial stain was observed at 100nM of Δ9-THC in ESCs (Figure 3C, p=0.02, unpaired T-
test), indicating that, at this dose, the observed increase in NAD(P)+/NAD(P)H could be explained
by higher mitochondrial membrane potential. By contrast, no change in EpiLCs mitochondrial
activity was detected (Figure 3C), consistent with these cells relying on glycolysis for energy
production[20,23].

Changes in mitochondrial activity in ESCs upon Δ9-THC exposure, although significant,
remained modest and are unlikely to be the sole contributor to the more significant increase in
NAD(P)+/NAD(P)H upon exposure. Thus, we performed an in-depth analysis of the differential
impact of Δ9-THC on ESCs and EpiLCs bioenergetics by measuring both glycolysis (extracellular
acidification rate, ECAR) and mitochondrial respiration (oxygen consumption rate, OCR) using a
Seahorse bioanalyzer. At 100nM of Δ9-THC, the maximal glycolytic capacity of both ESCs and
EpiLCs increased significantly (Figure 3D, 15% increase, p=0.03 and 22% increase, p=0.03 for
ESCs and EpiLCs, respectively, compared to the mock-treated condition, unpaired T-test). In both
cell types, a significant decrease in glycolytic capacity was observed at 10µM of Δ9-THC (Figure
3D, 39.8% reduction, p=0.0006 and 44.8% reduction, p=0.0001, for ESCs and EpiLCs,
respectively, compared to the mock-treated condition, unpaired T-test). Of note, the maximal
glycolytic capacity of EpiLCs in the untreated condition was higher than the one of ESCs, in
agreement with their metabolic shift towards aerobic glycolysis (Figure 3D, 7.88% higher ECAR
rate in mock-treated EpiLCs compared to mock-treated ESCs, p=0.03, unpaired T-test). As a
consequence, Δ9-THC exposure significantly impacted more glycolysis in EpiLCs than ESCs,
both in basal capacity and upon mitochondrial inhibition by oligomycin (Supplementary Figure 1A
and Figure 1B). In addition, at 100nM of Δ9-THC, the maximal respiratory capacity of ESCs was
significantly increased compared to the mock-treated condition (Figure 3E, 21.8% increase,
p=0.03, unpaired T-test). This increase was observed only for the maximal respiratory capacity of
ESCs, but not for basal respiration, nor for ATP-linked respiration (Supplementary Figure 1C),
suggesting that Δ9-THC impact on mitochondrial respiration does not support increased energetic
production. In agreement with EpiLCs metabolic shift towards a glycolytic phenotype, increasing
doses of Δ9-THC did not alter their maximal respiratory capacity (Figure 3E), nor their global
oxygen consumption rate (Supplementary Figure 1D). In both cell types, a significant decrease in
oxygen consumption rate was observed at 10µM of Δ9-THC (Figure 3E and Supplementary Figure
1C and Figure 1D).

Together, our analysis of cellular bioenergetics following Δ9-THC exposure showed an
increased glycolytic rate in ESCs that was also observed in EpiLCs. However, the increased
oxygen consumption and the associated increase in mitochondrial activity were observed only in
ESCs following exposure to 100nM of Δ9-THC, likely for the oxidization of the accumulating
pyruvate generated from glycolysis.

Δ9-THC-induced increase in glycolysis supports anabolism and ESCs proliferation

Because our data indicated that the impact of Δ9-THC exposure on stem cells' bioenergetics did not result in greater ATP production, we next sought to characterize the global metabolic impact of Δ9-THC in these cells. ESCs and EpiLCs were continuously exposed to 100nM Δ9-THC and intracellular metabolites were detected and quantified by mass spectrometry (Figure 4A-E). To explore the metabolic signatures in the different samples, we performed a global principal component analysis (PCA) (Figure 4B). All samples clustered in well-defined groups of replicates, both by cell type on the first principal component (accounting for 65.81% of the variation) and by Δ9-THC exposure on the second principal component (accounting for 20.83% of the variation). Of the 126 metabolites detected in ESCs, 39 were significantly upregulated (Figure 4C and Supplementary Figure 2A) and only two metabolites – NAPDH and Adenine – were significantly downregulated. Of the 138 metabolites detected in EpiLCs, 95 were significantly upregulated (Figure 4C and Supplementary Figure 2B) and only one metabolite – NAPDH – was significantly downregulated. In agreement with the PCA, the overlap of over-expressed metabolites in response to Δ9-THC exposure was important between the two stem cell populations (Figure 4C, accounting for 79.49% and 32.63% of all upregulated metabolites in ESCs and EpiLCs, respectively). The functional interpretation of the significantly upregulated metabolites confirmed the Δ9-THC-associated increase in energy metabolism in the two stem cell populations. Indeed, amongst the 25 metabolic pathways upregulated, pyruvate metabolism and glycolysis were detected in both ESCs and EpiLCs (Figure 4D and Figure 4E, respectively). Increased mitochondrial respiration was also seen in ESCs with the enrichment of (ubi)quinone metabolism, indicating an increased synthesis of ubiquinone that serves as an electron carrier in oxidative phosphorylation. Of note, metabolite measurements showed that the ratio of glutathione in its reduced to oxidated form (GSH/GSSG) was unchanged in both stem cell types in response to Δ9-THC (Supplementary Figure 2C), suggesting that the increased mitochondrial respiration does not cause an overt elevation of oxidative stress. Importantly, and in agreement with the PCA, in both ESCs and EpiLCs, Δ9-THC exposure elicited an increase in metabolic pathways that feed anabolic reactions, in particular contributing to the synthesis of amino acids (tyrosine, tryptophan, arginine, alanine, valine, (iso)leucine, etc.), nucleotides ("Pyrimidine metabolism", "Purine metabolism"), NAD(P)+ ("Nicotinate and nicotinamide metabolism") and fatty acids ("Butanoate metabolism") (Figure 4D and Figure 4E).

Extensive metabolic profiling of ESCs and EpiLCs upon Δ9-THC exposure thus indicated that the increased glycolytic rates in both stem cell populations, rather than provoking an increased production of energy under the form of ATP, participated in increased anabolism. Such increased anabolism could explain the proliferation observed in ESCs upon Δ9-THC exposure. To test this hypothesis, we exposed ESCs to 100nM of Δ9-THC for 48h as above but 24h before the harvest, cells were exposed to 10mM of 2-deoxyglucose (2-DG), an inhibitor of glycolysis[37]. Despite increasing the energy stress (Supplementary Figure 3), inhibition of glycolysis by 2-DG did not significantly impact viability over this shorter time frame and at this concentration (Figure 4F). Importantly, glycolytic inhibition by 2-DG abrogated the Δ9-THC-induced increase in both cell number and NAD(P)+/NAD(P)H levels (Figure 4G and Figure 4H, 1.39-fold reduction and $p < 0.0001$ and 1.68-fold reduction and $p = 0.0064$, respectively, when comparing 100nM of Δ9-THC +/- 10mM 2-DG, unpaired T-test). Thus, exposure to Δ9-THC increases anabolism in both ESCs and EpiLCs, however, this increased anabolism only supports cellular proliferation in ESCs.

Δ9-THC exposure is associated with the upregulation of genes involved in anabolic pathways in ESCs but not in EpiLCs.

Our data shows that Δ9-THC exposure increases anabolic pathways in both ESCs and EpiLCs and that this causes the proliferation of ESCs but not of EpiLCs. We thus next examined whether this differential impact of Δ9-THC on ESCs and EpiLCs was mirrored by a change in these cells' transcriptomes. To this aim, we performed RNA-sequencing (RNA-seq) on ESCs and EpiLCs continuously exposed to 100nM Δ9-THC or to the vehicle control (Figure 5A).

Unsupervised exploration of the global transcriptome by PCA revealed that the vast majority of data variation could be attributed to the cell type (PC1, accounting for 98% of the variation) rather than to Δ9-THC exposure (PC2, accounting for 1% of the variation, Figure 5B). This suggests that Δ9-THC exposure only moderately impacts ESC and EpiLC transcriptomes. In agreement, we identified a low number of differentially expressed genes (DEGs) in both ESCs and EpiLCs (Figure 5C and Figure 5D, respectively). In ESCs, only 12 genes were significantly upregulated with a log2(fold-change)>0.5 and only 9 were significantly downregulated at the same threshold (Figure 5C, significance corresponds to adjusted p-value≤0.05). More genes were differentially expressed when looking at lower fold-changes (|log2(fold-change)|>0.25, Figure 5C), confirming that the magnitude of transcriptional effects due to Δ9-THC exposure is moderate. This low transcriptional impact following Δ9-THC exposure was also observed in EpiLCs (Figure 5D). Nevertheless, gene ontologies (GO) associated with Δ9-THC-induced DEGs revealed the biological significance of these low transcriptional changes (Figure 5E). In particular, GO terms associated with metabolic pathways involved in anabolism were significantly over-represented for upregulated genes in ESCs following Δ9-THC exposure (Figure 5E), such as: "Cellular aromatic compound metabolic process", "Cellular nitrogen compound biosynthetic process", "Organonitrogen compound metabolic process". This suggests that the glycolytic rewiring elicited by Δ9-THC exposure in ESCs has some transcriptional support. Indeed, when performing joint pathway integration between our transcriptomics data and our targeted metabolomics[38], we observed that Δ9-THC-induced perturbed genes and metabolites were associated with the observed anabolic effects (Figure 5F). In contrast, GO terms associated with metabolism were not found within the upregulated DEGs in EpiLCs. However, several GO terms relating to alterations in cellular components were enriched in EpiLCs (Figure 3E), such as: "Organelle organization", "Cellular component organization or biogenesis", "Microtubule-based process". This indicates that Δ9-THC exposure significantly upregulated genes in EpiLCs that impact organelles structure, integrity and position, in agreement with several reports in the literature[39,40].

Together, our analysis of ESCs and EpiLCs transcriptomes reveals a difference in the response of these stem cell populations to Δ9-THC exposure: the transcriptional alterations observed in ESCs supported their increased anabolism and proliferation, whereas changes in EpiLCs gene expression did not correlate with their metabolic changes.

Proliferation of Primordial Germ Cell-Like Cells stemming from prior Δ9-THC exposure.

PGCs display a distinct transcriptomic and metabolic profile compared to their cellular precursors that are recapitulated in vitro during the differentiation of ESCs into EpiLCs and then of EpiLCs into PGCLCs. Thus, we asked whether the metabolic alterations observed in ESCs and EpiLCs could lead to an altered differentiation program in PGCLCs. To this aim, we continuously exposed ESCs and EpiLCs to a Δ9-THC dose range of 10nM-1μM (or mock control), before

changing to a Δ9-THC-free media and inducing PGCLCs differentiation (Figure 6A). In particular,
we took advantage of ESCs that harbor two fluorescent reporters for germline markers,
Blimp1:mVenus and Stella:CFP[41]. Thus, the induction efficiency of PGCLCs within 5-days old
embryoid bodies can be detected by monitoring the fluorescence associated with each cell in flow
cytometry, allowing for the determination of a double-negative population (DN), a single-positive
population (SP) wherein Blimp1:mVenus is expressed and a double-positive population (DP)
expressing both Blimp1:mVenus and Stella:CFP, which represents the true specified PGCLC
population.
We first measured the impact of ESCs + EpiLCs Δ9-THC exposure on PGCLC induction
efficiency. Flow analyses revealed a dose-dependent increase in the induction efficiency of SP
and DP cell populations (Figure 6B). Specifically, at 100nM Δ9-THC, a significant decrease in DN
was observed, with a corresponding significant increase of 1.14-fold in SP and of 1.64-fold in DP
cells (Figure 6C, p=0.0002, p=0.05, and p<0.0001 for 100nM of Δ9-THC in DN, SP and DP
populations respectively compared to the mock-treated condition, unpaired T-test). To determine
if the increased proportion of PGCLCs generated from exposed precursors was due to higher
proliferative kinetics, we performed a proliferation tracing assay[42]. The tracing dye was added to
the cells on the day of aggregates formation, and fluorescence attenuation due to cell division was
measured in each subpopulation on day 5. At 100nM Δ9-THC, a smaller proportion of DN cells
underwent two or three mitotic divisions compared to the control (Figure 6D, 1.14-fold fewer cells
and 1.12-fold fewer cells, p=0.05 and p=0.04 for 2 divisions and 3 divisions, respectively, unpaired
T-test). In parallel, for the same dose, a significantly higher proportion of SP and DP cells
underwent three mitotic divisions compared to the control (Figure 6D, 1.24-fold and 1.11-fold,
p=0.03 and p=0.0035, for 3 divisions in SP and DP cells, compared to the control, unpaired T-
test). These results, therefore, indicate that the higher number of PGCLCs observed upon Δ9-
THC exposure originates from their increased proliferation during their specification and
differentiation.
Thus, Δ9-THC causes an alteration of the developmental kinetics that PGCLCs normally
undergo, even in the absence of direct exposure.
329
PGCLCs derived from Δ9-THC-exposed cells present an altered metabolism and
transcriptome
Since exposure to Δ9-THC prior to their specification increased the number of PGCLCs and
ESCs and PGCLCs share similar metabolic programs[20,23], we next sought to characterize their
associated metabolic and transcriptional changes. We, therefore, assessed the impact of
exposure of ESCs + EpiLCs to 100nM Δ9-THC on PGCLCs metabolism (Figure 7A).
First, NAD(P)+/NAD(P)H assessment revealed a modest but significant 1.17-fold increase
in NAD(P)+/NAD(P)H ratio in whole day 5 embryoid bodies deriving from exposed ESCs + EpiLCs
compared to those deriving from mock-treated cells (Figure 7B, p=0.01, unpaired T-test). To
garner cell type-specific information on whether these metabolic changes were related to
mitochondrial activity and the differentiation of PGCLCs, we assessed the mitochondrial
membrane potential of each subpopulation in day 5 embryoid bodies. Embryoid bodies were
incubated with Mitotracker CMXRos[36], and dissociated and analyzed by flow cytometry. The MFI
associated with the mitochondrial stain was then measured in each subpopulation (Figure 7C). A
significant increase in MFI was observed in DN, SP and DP populations deriving from exposed

ESCs +EpiLCs compared to those deriving from mock controls (Figure 7C, 1.17, 1.16, 1.23 -fold, p=0.006, p=0.05 and p=0.01 for DN, SP and DP, respectively, unpaired T-test). These results indicate that the metabolic changes induced by Δ9-THC prior to PGCLCs induction and differentiation are not reset during the profound reprogramming that PGCLCs undergo.

Because our results indicated a sustained impact of Δ9-THC beyond the period of direct exposure, we further examined PGCLCs by performing a transcriptomic analysis. In particular, day 5 embryoid bodies deriving from ESCs + EpiLCs, either exposed to 100nM of Δ9-THC or mock-exposed, were sorted and the total RNA of DP subpopulations, representing true PGCLCs, was analyzed by RNA-seq (Figure 7A). Unsupervised analysis of the global transcriptome in DP PGCLCs by PCA delineated a transcriptional signature of prior Δ9-THC exposure (Figure 7D, PC1 accounting for 59% of the variance and PC2 accounting for 24% of variance). Volcano plot of DEGs between DP PGCLCs deriving from mock- or 100nM Δ9-THC-exposed ESCs and EpiLCs revealed that most of the significant transcriptional change was towards downregulation rather than upregulation (Figure 7E, 11 genes were significantly upregulated whereas 97 were significantly downregulated, |log2(fold-change)|>0.25 and adjusted p-value≤0.05). Despite the low number of upregulated DEGs, the functional annotation of their associated GO terms showed that all terms enriched corresponded to metabolic processes involved in oxidative phosphorylation (Figure 7F, "Aerobic electron transport chain", "Mitochondrial respiratory chain complex I assembly", "Electron transport couple proton transport"). Thus, our data indicate that the metabolic changes induced by exposure to Δ9-THC prior to PGCLCs specification are retained through transcriptional reprogramming. Importantly, while our results show that pre-specification Δ9-THC exposure increases PGCLCs number and mitochondrial activity, the functional annotation of GO terms associated with downregulated DEGs suggests degradation of PGCLCs quality. Indeed, and reminiscent of GO terms observed in EpiLCs, several GO terms relating to alterations in structural cellular components ("Anatomical structure morphogenesis", "Cellular anatomical entity"), and in particular the interface with the extracellular environment ("External encapsulating structure organization", "Membrane", "Cell periphery", "Extracellular region", "Extracellular space", "Extracellular matrix structural constituent") were enriched (Figure 7G). Furthermore, GO terms associated with cell adhesion and junction ("Cell adhesion", "Cell migration", "Collagen metabolic process", "Cell junction", "Anchoring junction", "Collagen trimer") were also enriched in downregulated genes.

Together, our data show that Δ9-THC exposure in ESCs and EpiLCs durably alters their metabolome and that these changes are carried through PGCLCs specification and differentiation, leading to an alteration of PGCLCs transcriptional program (Figure 8).

DISCUSSION

With greater social acceptance and legalization, cannabis use has increased worldwide[1,3]. Yet, the impact of such heightened use on reproductive functions, and in particular, on the earliest developmental stages are not well understood. Cannabis use directly alters adult male fertility and causes abnormal embryo implantation[14]. Using a well-characterized in vitro model of early embryonic differentiation events culminating into the differentiation of PGCLCs, our study is the first to shed light on the impact of Δ9-THC at these stages which unfold during the first trimester in humans[4–6].

Our data revealed the differential effects of Δ9-THC on naïve and primed pluripotent stem
cells, respectively represented by ESCs and EpiLCs. In particular, exposure to Δ9-THC increased
ESCs proliferation which was in a similar range to what has been previously reported for human
breast carcinoma cell lines (about 30-50% between 10nM and 1μM of Δ9-THC)[30]. Differential
expression and use of cannabinoid receptors on the surface of exposed cells have been shown
to correlate with Δ9-THC proliferative phenotypes[29–32]. However, our experiments demonstrated
that despite being required for Δ9-THC-induced proliferation in ESCs, CB1 expression did not
significantly differ at the surface of ESCs and EpiLCs.
Because Δ9-THC is a known perturbator of mitochondrial function as previously described
in the central nervous system[12,34], we studied the metabolic impact of its exposure in ESCs and
EpiLCs. Our data indicate that, at 100nM, Δ9-THC exposure increased the glycolytic rate in both
ESCs and EpiLCs. Bioenergetics analyses and metabolite measurements showed that this
increased glucose metabolism did not support increased energy production in the mitochondria,
but rather, that it led to the accumulation of metabolic intermediates used in anabolic reactions for
the synthesis of amino acids, nucleotides, and lipids. Thus, the metabolic signatures associated
with Δ9-THC exposure are reminiscent of those inherently occurring during naïve-to-prime
transition, during which increased aerobic glycolytic rates feed anabolic reactions ultimately
fueling proliferation[24]. We verified this model by testing the requirement of increased glycolysis to
support proliferation and indeed observed that ESCs proliferation upon Δ9-THC exposure is
abrogated in the presence of the glycolytic inhibitor 2-DG.
Transcriptomic analyses revealed that the metabolic reprogramming induced by Δ9-THC
exposure in ESCs was transcriptionally encoded, with increased expression of genes involved in
anabolic pathways. In contrast, functional annotations of DEGs in EpiLCs did not show such
transcriptional control of increased anabolism. Comparing the outputs of the metabolomic and
transcriptomic analyses (i.e. PCA plots and volcano plots), the impact of Δ9-THC at these early
stages seems to be primarily metabolic, although the moderate effects on the transcriptome
appear to support the metabolic outcome as revealed by our integrated analysis (Figure 5F).
Together, we propose that Δ9-THC exposure elicits a reprogramming of ESCs that (1) coaxes
them to rely more on aerobic glycolysis, (2) drives anabolic pathways, and therefore (3) leads to
their proliferation. In EpiLCs, the impact of Δ9-THC exposure is not sufficient to override the
cellular and metabolic programs of these already highly proliferative cells that are fully reliant on
aerobic glycolysis (Figure 8).
Finally, we assessed the impact of Δ9-THC exposure in ESCs and EpiLCs on the
differentiation of PGCLCs. Our data indicate that at the physiologically relevant dose of 100nM of
Δ9-THC, a significant increase in PGCLCs was observed. In particular, during PGCLCs
differentiation, metabolic reprogramming and increased oxidative phosphorylation play a critical
role in the reacquisition of an extended developmental potential[20,23]. Thus, we investigated
whether the metabolic alterations observed in ESCs and EpiLCs upon Δ9-THC exposure could
be carried through PGCLCs differentiation. Metabolic characterization revealed that PGCLCs
arising from exposed ESCs and EpiLCs showed increased mitochondrial respiration. Thus, in the
absence of continuous exposure, Δ9-THC still has lasting consequences on the metabolome of
embryonic germ cells. A recent study in drosophila reported that nutrient stress induces oocytes
metabolites remodeling that drives the onset of metabolic diseases in the progeny[43]. This indicates
that non-DNA-associated factors, such as germline metabolites, can act as factors of inheritance.

Similarly, we show here that exposure to Δ9-THC remodels ESCs and EpiLCs metabolome and
that a metabolic memory of this exposure is retained during PGCLCs differentiation (Figure 8). In
addition to metabolic remodeling, we show that the PGCLCs transcriptome is also altered. In
particular, despite proliferation and a higher number of cells, the number of DEGs that were
downregulated in PGCLCs deriving from Δ9-THC-exposed ESCs and EpiLCs suggests a general
degradation of PGCLCs' homeostasis. Functional annotation further indicated that these
downregulated genes are related to structural cellular components, to the interaction with the
extracellular environment and, specifically, to cell adhesion and junction. During the development
of the central nervous system, perinatal Δ9-THC exposure has also been associated with
alteration in cell adhesion, with an impact on neuronal interactions and morphology[44–46]. Cell-cell
adhesion is crucial in PGCs' formation both in cell culture systems[47] as well as in vivo where it
controls PGCs motility during their migration towards the developing somatic gonad[48]. Our results
thus suggest that exposure to Δ9-THC prior to specification affects embryonic germ cells'
transcriptome and metabolome, with potentially adverse consequences on cell-cell adhesion that
could impact their normal development in vivo.
Together, our studies reveal a moderate but significant impact of Δ9-THC exposure on early
embryonic processes. Our work also highlights the importance of the metabolic remodeling
induced by Δ9-THC and its potential role as a driver of exposure memory through differentiation
stages.

METHODS

Data availability

The RNA sequencing data from this study is made available at the Gene Expression Omnibus (GEO) under the following accession number GSE226955. All other data are available upon request.

Cell culture and PGCLCs model

Mouse ESCs containing the two fluorescent reporters Blimp1::mVenus and Stella::ECFP (BVSC cells) were described previously [41]. The female BVSC clone H18 was kindly provided by Mitinori Saitou and cells were seeded on coated plates (Poly-L-ornithine [0.001%; A-004-C; Sigma-Aldrich] and laminin [300ng/mL; L2020; Sigma-Aldrich]) in 2i+LIF culture medium (N2B27 Media, CHIR99021 [30μM; NC9785126; Thermo Fisher], PD0325901 [10μM; NC9753132; Thermo Fisher], ESGRO® Leukemia Inhibitory Factor (LIF) [1,000 U/mL, ESG1106; Sigma-Aldrich]) for 48h. Differentiation of ESCs to EpiLCs was performed by seeding the cells on Human Plasma Fibronectin (HPF)-coated plates [16.7μg/mL; 33016015; Thermo Fisher] in the presence of EpiLC induction medium (N2B27 medium containing activin A [20ng/mL; 50-398-465; Thermo Fisher], basic fibroblast growth factor (bFGF) [12ng/mL; 3139FB025; R&D Systems], and KnockOut Serum Replacement [KSR, 1%; Thermo Fisher]). For PGCLCs induction, 44h EpiLCs were harvested using TrypLE™ Select (1X) (Thermo Fisher) and seeded either in 96-wells plate (Nunclon Sphera, Thermo Fisher) or in EZsphere plates for large-scale induction (Nacalai) in the presence of GK15 medium (Glasgow's Minimal Essential Medium [GMEM, 11710035, Thermo Fisher] supplemented with 15% KSR, 0.1 mM Minimal Essential Medium Nonessential Amino Acids [MEM-NEAA], 1 mM sodium pyruvate, 0.1 mM 2-mercaptoethanol, 100U/mL penicillin, 0.1mg/mL streptomycin, and 2 mM L-glutamine in the presence of bone morphogenetic protein 4 [BMP4; 500ng/mL; 5020-BP-010/CF; R&D Systems], LIF, stem cell factor [SCF; 100ng/mL; 50-399-595; R&D Systems], bone morphogenetic protein 8b [BMP8b; 500ng/mL; 7540-BP-025; R&D Systems], and epidermal growth factor [EGF; 50ng/mL; 2028EG200; R&D Systems]. Cells were culture for 5d before collection, dissociation of embryoid bodies and downstream experiments. All cells were cultured in a humidified environment at 37°C under 5% CO_2.

Δ9-THC exposures

To assess the impact of Δ9-THC exposure on the developmental trajectory of PGCLCs, three exposure schemes were tested: 1) ESCs exposure only, 2) EpiLCs exposure only, and 3) ESCs+EpiLCs exposure. The stock of Δ9-THC was obtained from the National Institute on Drug Abuse (7370-023 NIDA; Bethesda, MD). The stock was adjusted to a concentration of 200mM diluted in ethanol, aliquoted and stored according to the DEA's recommendations. The dose range of 0-100μM was determined based on Δ9-THC physiological measurements in the blood, plasma, and follicular fluid[26–28]. For each exposure, new aliquots of Δ9-THC were diluted in ESCs or EpiLCs culture media in coated tubes (Sigmacote, Sigma Aldrich). Exposure was performed for 48h. Solubility tests were performed and ethanol was added to reach the same amount for each Δ9-THC concentration (0.05% ethanol). Vehicle control corresponded to 0.05% ethanol added to the respective culture media for ESCs or EpiLCs. All experiments performed are authorized under DEA registration number RA0546828.

PGCLCs induction efficiency

Changes in PGCLCs induction were calculated by flow cytometry. Practically, d5 aggregates were harvested, dissociated using TrypLE™ Select, and resuspended in fluorescence-activated cell sorting (FACS) buffer (1×Dulbecco's phosphate buffered saline [DPBS], 1% BSA, 1 mM EDTA, 25 mM HEPES). Quantification of subfractions of double-positive PGCLCs (Blimp1::mVenus+ and Stella::ECFP+), single-positive (Blimp1::mVenus+) and double-negative cells was performed on a BD Biosciences LSRII (UCLA BSCRC Flow Cytometry Core). Cells were initially identified by forward- and side-scatter gating, with back-gating used to verify the accuracy by which target cell populations were identified. Cell populations of interest were identified by 2-D plots displaying the parameter of interest, using embryoid bodies cultured in GK15 medium without added cytokines and BMPs as a negative control. Manually defined gates as well as quadrants were used, as indicated. The FlowJo software was used to calculate percent of induction and generate the associated graphs (version 10, FlowJo, LLC).

Cell viability and proliferation studies

The viability and viable cell count of ESCs and EpiLCs was calculated using Trypan blue (0.4%, Thermo Fisher) on a Countess II FL Automated Cell Counter (Thermo Fisher). For BrdU incorporation studies, cells were permeabilized, fixed, and stained using the BrdU Flow Kit (PerCP-Cy™5.5 Mouse anti-BrdU, BD Biosciences) before analysis by flow cytometry on a BD Biosciences LSRII (UCLA BSCRC Flow Cytometry Core). Quantification of PGCLCs proliferation was performed using CellTrace™ Yellow (5µM, added at the induction, Thermo Fisher), which binds to intracellular amines after diffusing through cell membranes. The overall fluorescent signal, that gradually decreases as cell division occurs, reflects the number of cell divisions occurring and was measured on a BD Biosciences LSRII (UCLA BSCRC Flow Cytometry Core). The FlowJo software was used to calculate percent of induction, numbers of cell division and generate the associated graphs (version 10, FlowJo, LLC).

CB1 antagonist treatment

To block the effects of Δ9-THC on the cannabinoid receptor CB1, ESCs were plated on 48-well plate and were pre-treated with 1µM of SR141716/Rimonabant (SML0800, Sigma Aldrich) for 1h before being exposed to the dose range of Δ9-THC, as above. After 24h incubation, this procedure was repeated and cells were harvested after 48h total incubation. The viability and viable cell count was calculated using Trypan blue (0.4%, Thermo Fisher) on a Countess II FL Automated Cell Counter (Thermo Fisher). The concentration of 1µM of Rimonabant was chosen based on previous experiments[39] and did not impact cell viability nor cell number on its own.

Western blotting

Membrane proteins were extracted from cell pellets using the Mem-PER™ Plus Membrane Protein Extraction Kit (89842, Thermo Fisher) according to the manufacturer's protocol. Western blotting was performed with 25µg of protein extracts. The immunodetection was assessed using primary antibodies targeting CB1 (101500, Cayman Chemical) or β-actin (3700, Cell Signaling Technology) as loading control. Horseradish peroxidase (HRP)-conjugated secondary antibodies were used for chemiluminescence detection (Amersham).

WST-1 assay

The colorimetric assay WST-1 was used according to the manufacturer's instructions (Roche). The tetrazolium salt WST-1 is reduced by mitochondrial dehydrogenases to formazan using NAD(P)H as co-substrates. Thus, the quantity of formazan is directly proportional to NAD(P)$^+$.

Mitochondrial activity

Staining for mitochondria was performed by incubating cells at 37°C with 250nM MitoTracker CMXRos (M7512, Thermo Fisher) for 30min [36]. Cells were washed and analyzed by flow cytometry on a BD Biosciences LSRII (UCLA BSCRC Flow Cytometry Core). The FlowJo software (version 10, FlowJo, LLC) was used to calculate the mean fluorescence intensity (MFI) corresponding to the average fluorescence intensity of each event of the selected cell population within the chosen fluorescence channel associated to MitoTracker CMXRos.

Seahorse experiments

The extracellular acidification rate (ECAR) and the oxygen consumption rate (OCR) are indicative of glycolysis and mitochondrial respiration, respectively. A total of 10×10^3 ESCs and 8×10^3 EpiLCs were seeded on Seahorse XF96 plates (101085-004, Agilent Technologies) and exposed to increasing doses of Δ9-THC for 48h. On the day of the assay, cells were washed with assay medium (unbuffered DMEM assay medium [5030, Sigma Aldrich] supplemented with 31.6mM NaCl, 3mg/L phenol red, 5mM HEPES, 5mM glucose, 2mM glutamine and 1mM sodium pyruvate). For OCR measurement, compounds were injected sequentially during the assay resulting in final concentrations of 2µM oligomycin, 0.75µM and 1.35µM FCCP, 1µM rotenone and 2µM antimycin. ECAR was measured in parallel. The measured quantities were normalized to the protein content as measured by a BCA quantitation (23227, Thermo Fisher).

Mass spectrometry-based metabolomics analysis

To extract intracellular metabolites, cells were rinsed with cold 150mM ammonium acetate (pH7.3) then incubated with 80% ice-cold methanol supplemented with 10 nmol D/L-norvaline for 1h. Following resuspension, cells were pelleted by centrifugation (15,000g, 4°C for 15min). The supernatant was transferred into a glass vial and metabolites were dried down under vacuum then resuspended in 70% acetonitrile. Mass spectrometry analysis was performed at the UCLA Metabolomics Center with an UltiMate 3000RSLC (Thermo Scientific) coupled to a Q Exactive mass spectrometer (Thermo Scientific) in polarity-switching mode with positive voltage 3.0 kV and negative voltage 2.25 kV. Separation was achieved using a gradient elution with (A) 5mM NH4AcO (pH 9.9) and (B) acetonitrile. The gradient ran from 15% (A) to 90% (A) over 18 min, followed by an isocratic step for 9 minutes and re-equilibration for 7 minutes. Metabolites were quantified as area under the curve based on retention times and using accurate mass measurements (≤ 3 ppm) with the TraceFinder 3.1 software (Thermo Scientific). For heatmap depiction, the relative amounts of metabolites were normalized to the mean value across all samples for one same condition and to the number of viable cells harvested in parallel on a control plate. Pathway enrichment for up- and downregulated KEGG metabolites (|log2(fold-change)|=0.25) was determined using the MetaboAnalyst 5.0 platform (www.metaboanalyst.ca)[38].

RNA-sequencing

Total RNA was extracted from ESCs and EpiLCs pellets using the AllPrep DNA/RNA Micro Kit (Qiagen), according to the manufacturer's protocol. For PGCLCs, d5 embryonic bodies were harvested and cells were dissociated using TrypLE™ Select followed by resuspension in fluorescence-activated cell sorting (FACS) buffer (1×Dulbecco's phosphate buffered saline [DPBS], 1% BSA, 1 mM EDTA, 25 mM HEPES) and cell suspension were passed through a cell strainer (70µm). Cells were sorted on a BD Biosciences FACSAria III (UCLA BSCRC Flow Cytometry Core). Practically, cell populations of interest, being double-positive (Blimp1::mVenus+ and Stella::ECFP+) were sorted and collected in microtubes containing GK15 medium. Total RNA was extracted from double-positive PGCLCs using the AllPrep DNA/RNA Micro Kit (Qiagen). RNA concentration was measured using a NanoDrop™ 2000 UV spectrophotometer (Thermo Fisher). Libraries were prepared with the KAPA mRNA HyperPrep Kit (BioMek) or with the RNA library prep kit (ABClonal) following the manufacturers' protocols. Briefly, poly(A) RNA were selected, fragmented and double-stranded cDNA synthesized using a mixture of random and oligo(dT) priming, followed by end repair to generate blunt ends, adaptor ligation, strand selection, and polymerase chain reaction (PCR) amplification to produce the final library. Different index adaptors were used for multiplexing samples in one sequencing lane. Sequencing was performed on an Illumina NovaSeq 6000 sequencers for paired end (PE), 2×150 base pair (bp) runs. Data quality check was performed using Illumina Sequencing Analysis Viewer (SAV) software. Demultiplexing was performed with Illumina Bcl2fastq2 program (version 2.19.1.403; Illumina Inc.).

Differential gene expression analysis

The quality of the reads was verified using FastQC[49] before reads were aligned to the mm10 reference genome (GRCm39) using STAR [50] with the following arguments: --readFilesCommand zcat --outSAMtype BAM SortedByCoordinate --quantMode GeneCounts --outFilterMismatchNmax 5 --outFilterMultimapNmax 1. The quality of the resulting alignments was assessed using QualiMap[51]. The Python package HTseq was used for gene counts [52] using the following arguments: --stranded=no --idattr=gene_id --type=exon --mode=union -r pos --format=bam. Output files were filtered to remove genes with low count (≤10) then were used for differential gene expression analysis using DESeq2[53]. The negative binomial regression model of ComBat-seq was used to correct unwanted batch effects[54]. For a gene to be classified as showing differential expression between treated and untreated cells, a threshold of |log2(fold-change)|=0.5 and Benjamini-Hochberg adjusted p-value ≤0.05 had to be met.

Gene Ontology (GO) Analysis

Lists of differentially expressed genes were generated from read counts using DESeq2 Bioconductor package[53]. Enrichment of GO terms in lists of up- and downregulated genes (|log2(fold-change)|=0.25) was determined using g:Profiler[55]. Redundant GO terms were removed using reduce + visualize gene ontology (REVIGO)[56]. Terms were included if the fold enrichment (frequency of DEGs in each GO term to the frequency of total genes in GO terms) was higher than 1.5 and if the Benjamini-Hochberg-adjusted p-value was less than 0.05. Plots for GO terms were generated using a custom R script[57].

Statistical Methods

Statistical analyses, when not otherwise specified, were performed using GraphPad Prism 9 software. For significance testing, two-tailed T-tests were performed on pairwise comparisons. In all cases, significance was determined by p-values less than or equal to 0.05. Each figure corresponds to at least three independent biological repeats with three technical replicates (N=3, n=3), unless otherwise specified. Number of asterisks on plots indicate level of statistical significance: *($p<0.05$), **($p<0.01$), ***($p<0.001$) and ****($p<0.0001$).

ACKNOWLEDGMENTS

We thank Jessica Scholes, Felicia Codrea and Jeffrey Calimlim from the UCLA BSCRC Flow Cytometry Core for their assistance in cell sorting. We thank Linsey Stiles (UCLA Mitochondria and Metabolism Core), Johanna ten Hoeve-Scott and Thomas Graeber (UCLA Metabolomics Center) for their technical support in the metabolomics analyses. We thank Xinmin Li and his team at the UCLA Technology Center for Genomics & Bioinformatics for their technical support in the high-throughput sequencing. We thank Luigi Bellocchio and Giovanni Marsicano for their recommendations in Δ9-THC exposure setups. We would like to acknowledge the UCLA Cannabis Research Initiative (UCRI) and Dr Ziva Cooper for their continuous support and encouragements.

PA is supported by NIEHS R01 ES027487, the John Templeton Foundation Grant 60742, and the Iris Cantor-UCLA Women's Health Center and the NCATS UCLA CTSI Grant Number UL1TR001881. RV is a postdoctoral fellow and acknowledges support from the Belgian-American Educational Foundation (BAEF).

 Figure 2

649 **Figure 3**

Figure 5

 Figure 6

A.

ESCs   EpiLCs   PGCLCs

BMP4+BMP8a+SCF+EGF+LIF

bFGF+ActivinA

2i+LIF

Day 0   1   2   3   4   5   6   7   8   9

Δ9-THC exposure

Metabolic profiling
Transcriptome profiling
(DP PGCLCs)

B.

NAD(P)$^+$/NAD(P)H

mock   100nM Δ9-THC

C.

DN

MFI:Mitotracker
(normalized to the mock)

mock   100nM Δ9-THC

SP

MFI:Mitotracker
(normalized to the mock)

mock   100nM Δ9-THC

DP

MFI:Mitotracker
(normalized to the mock)

mock   100nM Δ9-THC

D.

PC2

PC1

● DP PGCLCs mock
● DP PGCLCs 100nM Δ9-THC

E.

DP PGCLCs

$-\log_{10}$(FDR)

$\log_2$(FC)

F.

Biosynthetic process (GO:0009058)
Aerobic electron transport chain (GO:0019646)
Mitochondrial respiratory chain complex I assembly (GO:0032981)
Electron transport coupled proton transport (GO:0015990)
Respirasome (GO:0070469)
Respiratory chain complex (GO:0098803)
Electron transfer activity (GO:0009055)
Oxidoreduction-driven active transmembrane transporter activity (GO:0015453)

$-\log_{10}$(FDR)

GO:BP
GO:CC
GO:MF

G.

Biological regulation (GO:0065007)
Multicellular organismal process (GO:0032501)
Developmental process (GO:0032502)
External encapsulating structure organization (GO:0045229)
Anatomical structure morphogenesis (GO:0009653)
Cell adhesion (GO:0007155)
Locomotion (GO:0040011)
Cell death (GO:0008219)
Apoptotic process (GO:0006915)
Positive regulation of developmental process (GO:0051094)
Cell migration (GO:0016477)
Cell population proliferation (GO:0008283)
Ossification (GO:0001503)
Collagen metabolic process (GO:0032963)
Cellular anatomical entity (GO:0110165)
Membrane (GO:0016020)
Cell periphery (GO:0071944)
Endomembrane system (GO:0012505)
Extracellular region (GO:0005576)
Bounding membrane of organelle (GO:0098588)
Cell junction (GO:0030054)
External encapsulating structure (GO:0030312)
Extracellular space (GO:0005615)
Anchoring junction (GO:0070161)
Cell surface (GO:0009986)
Perinuclear region of cytoplasm (GO:0048471)
Collagen trimer (GO:0005581)
Growth factor binding (GO:0019838)
Extracellular matrix structural constituent (GO:0005201)
Protein binding (GO:0005515)

$-\log_{10}$(FDR)

GO:BP
GO:CC
GO:MF

Figure 8

656 **Supplementary Figure 1**

 Supplementary Figure 3

FIGURE LEGENDS

Figure 1: Δ9-THC exposure provokes the proliferation of ESCs but not EpiLCs

(A) Diagram illustrating Δ9-THC exposure scheme and experimental strategy. bFGF: basic fibroblast growth factor, ESCs, embryonic stem cells; EpiLCs, epiblast-like cells; LIF, leukemia inhibitory factor. **(B, E, H)** Whisker boxplot indicating the median cellular viability of stem cells exposed to the different Δ9-THC doses and associated errors. **(C, F, I)** Whisker boxplot indicating the median number of viable cells exposed to the different Δ9-THC doses indicated and associated errors. **(D, G, J)** Whisker boxplot indicating the median percentage of BrdU-stained cells exposed to the different Δ9-THC doses and associated errors. ESCs exposed cells are presented in (B, C and D). EpiLCs exposed cells deriving from unexposed ESCs are presented in (E, F and G). EpiLCs exposed cells deriving from exposed ESCs are presented in (H, I and J).

Figure 2: Implication of the CB1 receptor in the proliferative phenotype.

(A) Western blot analysis of transmembrane protein extracts of ESCs or EpiLCs. Antibodies raised against CB1 or β-actin serving as a loading control were used for immunoblotting. **(B)** Quantification of the gel presented in (A) was done using Image Studio (version 5.2). **(C)** Whisker boxplot indicating the median cellular viability of stem cells exposed to the different Δ9-THC and rimonabant doses indicated and their associated errors. **(D)** The median numbers of viable cells exposed to the different Δ9-THC and rimonabant doses indicated were normalized to their own control (+/- rimonabant). Median and associated errors were plotted in whisker boxplots.

Figure 3: Δ9-THC exposure provokes an increase in glycolytic rates in ESCs and EpiLCs.

(A) Diagram illustrating Δ9-THC exposure scheme and experimental strategy. **(B)** The NAD(P)+/NADPH ratio of stem cells exposed to the different Δ9-THC doses was normalized to the one measured in the mock-treated condition. Median and associated errors were plotted in whisker boxplots. **(C)** Mean fluorescence intensity (MFI) associated with the Mitotracker CMXRos stain was normalized to the one measured in the mock-treated condition. Median and associated errors were plotted in whisker boxplots. **(D)** Median and associated error of the maximal extracellular acidification rate (ECAR) measured in cells exposed to the different Δ9-THC doses and normalized to the protein content was plotted in whisker boxplots. **(E)** Median and associated error of the maximal oxygen consumption rate (OCR) measured in cells exposed to the different Δ9-THC doses and normalized to the protein content was plotted in whisker boxplots. For (B and C), 5 technical repeats of 3 biological repeats (n=15) were plotted. One same representative experiment out of three independent experiments was used to plot results in (D and E).

Figure 4: Δ9-THC-induced glycolysis sustain anabolism and ESCs proliferation

(A) Diagram illustrating Δ9-THC exposure scheme and experimental strategy. **(B)** PCA of the metabolomics profiling of either ESCs or EpiLCs mock-exposed or exposed to 100nM Δ9-THC. **(C)** Venn diagram showing the overlap in upregulated metabolites following Δ9-THC exposure in ESCs and EpiLCs. **(D and E)** KEGG metabolite sets enrichment analysis for upregulated metabolites in ESCs and EpiLCs, respectively, performed by MetaboAnalyst[38]. KEGG, Kyoto Encyclopedia of Genes and Genomes. **(F)** Whisker boxplot indicating the median cellular viability of stem cells exposed to 100nM of Δ9-THC and 10mM of 2-DG, as indicated, and their associated errors. **(G)** The median numbers of viable cells exposed to 100nM of Δ9-THC and 10mM of 2-DG,

703   as indicated, were normalized to their own control (+/- 2-DG). Median and associated errors were
704   plotted in whisker boxplots. **(H)** The NAD(P)+/NADPH ratio of stem cells exposed to 100nM of Δ9-
705   THC and 10mM of 2-DG, as indicated, was normalized to the one measured in the mock-treated
706   condition (+/- 2-DG). Median and associated errors were plotted in whisker boxplots.
707
Figure 5: Metabolic changes following Δ9-THC exposure in ESCs are transcriptionally
encoded.
(A) Diagram illustrating Δ9-THC exposure scheme and experimental strategy. **(B)** PCA of the
transcriptomics profiling of either ESCs or EpiLCs mock-exposed or exposed to 100nM Δ9-THC.
(C and D) Volcano plot in ESCs and EpiLCs, respectively, showing significance [expressed in
log_{10}(adjusted p-value or false-discovery rate, FDR)] versus fold-change [expressed in log2(fold-
change, FC)]. Thresholds for significance (adjusted p-value≤0.05) and gene expression fold-
change [|(log2(FC)|>0.25 or |log2(FC)|>0.5] are shown as dashed lines. Color code is as follows:
log2(FC)>0.5 in red, log2(FC)>0.25 in orange, log2(FC)>0 in light orange, log2(FC)<0 in light blue,
log2(FC)>-0.25 in blue, log2(FC)>0.5 in dark blue and p-value<0.01 in pink. **(E)** Gene ontology
(GO) terms associated with up- and downregulated DEGs [|(log2(FC)|>0.25 and p<0.01)] in ESCs
and EpiLCs as determined by g:Profiler[55]. **(F)** Joint pathway analysis performed by the multi-omics
integration tool of MetaboAnalyst[38]. The p-values were weighted based on the proportions of
genes and metabolites at the individual pathway level.
722
Figure 6: PGCLCs deriving from ESCs and EpiLCs exposed to 100nM of Δ9-THC
proliferate.
(A) Diagram illustrating Δ9-THC exposure scheme and experimental strategy. **(B)** Representative
flow contour plots showing distribution of live-gated events, gating strategy for Stella:CFP versus
Blimp1:mVenus and percentages of cells in each subpopulations for ESCs and EpiLCs exposed
to the different doses of Δ9-THC indicated. DN: double negative, SP: single positive, DP: double
positive subpopulations. **(C)** The percentage of events in the gates associated to each
subpopulation was normalized to the one measured in the mock-treated condition. Median and
associated errors were plotted in whisker boxplots independently for each subpopulation. **(D)**
Representative histograms showing CellTrace™ Yellow staining profile of cells arising from ESCs
and EpiLCs exposed to the different doses of Δ9-THC indicated. The Y-axis represents the
average percentage of cells in each category of subpopulations undividing (purple), undergoing 1
division (blue), 2 divisions (green) or 3 divisions (orange). One representative experiment out of
three is represented.
737
Figure 7: Δ9-THC exposure prior to specification increases mitochondrial respiration in
PGCLCs.
(A) Diagram illustrating Δ9-THC exposure scheme and experimental strategy. **(B)** The
NAD(P)+/NADPH ratio of embryoid bodies arising from ESCs and EpiLCs exposed to 100nM of
Δ9-THC was normalized to the one measured in the mock-treated condition. Median and
associated errors were plotted in whisker boxplots. **(C)** Mean fluorescence intensity (MFI)
associated with the Mitotracker CMXRos stain in each subpopulation was normalized to the one
measured in the mock-treated condition. Median and associated errors were plotted in whisker
boxplots. **(D)** PCA of the transcriptomics profiling of DP PGCLCs deriving from ESCs and EpiLCs

either mock-exposed or exposed to 100nM Δ9-THC. **(E)** Volcano plot in DP PGCLCs showing
significance [expressed in $\log_{10}$(adjusted p-value or false-discovery rate, FDR)] versus fold-
change [expressed in log2(fold-change, FC)]. Thresholds for significance and different enrichment
ratios [|(log2(FC)|>0.25 or |log2(FC)|>0.5] are shown as dashed lines. Color code is as follows:
log2(FC)>0.5 in red, log2(FC)>0.25 in orange, log2(FC)>0 in light orange, log2(FC)<0 in light blue,
log2(FC)>-0.25 in blue, log2(FC)>0.5 in dark blue and p-value<0.01 in pink. **(F and G)** Gene
ontology (GO) terms associated with up- and downregulated DEGs [|(log2(FC)|>0.25 and
p<0.01)], respectively, as determined by g:Profiler[55].

Figure 8: Metabolic impact of Δ9-THC exposure in pluripotent stem cells and primordial
germ cells-like cells.
Diagram illustrating the impact of Δ9-THC exposure on stem cells metabolism.

Supplementary Figure 1: Extracellular acidification rates and oxygen consumption rates
in ESCs and EpiLCs upon Δ9-THC exposure.
(A and B) Traces were plotted for the extracellular acidification rate (ECAR) measurements in
ESCs and EpiLCs, respectively, exposed to the different Δ9-THC doses indicated and normalized
to the protein content. The oligomycin injection time is indicated by an arrow and allows to
differentiate basal glycolytic rate from maximal glycolytic rate (when mitochondria are inhibited).
The datapoints used in the main figure correspond to the first timepoint in the maximal glycolytic
capacity section. **(C and D)** Traces were plotted for the oxygen consumption rate (OCR)
measurements in ESCs and EpiLCs, respectively, exposed to the different Δ9-THC doses
indicated and normalized to the protein content. The oligomycin, FCCP and AntimycinA/Rotenone
injection times are indicated by arrows and allow to differentiate basal respiration from ATP-
coupled respiration and maximal respiratory capacity. The datapoints used in the main figure
correspond to the second timepoint in the maximal respiratory capacity section. FCCP: Carbonyl
cyanide-p-trifluoromethoxyphenylhydrazone.

Supplementary Figure 2: Metabolite profiling in ESCs and EpiLCs upon Δ9-THC exposure.
(A and B) Heatmaps showing the log2 of the amount of each metabolite upregulated in ESCs and
EpiLCs upon exposure to 100nM of Δ9-THC. The relative amounts of metabolites were normalized
to the mean value across all samples for one same condition and to the number of viable cells
harvested in parallel on a control plate. **(C)** Histograms showing the ratio of reduced to oxidized
glutathione (GSH/GSSG) based on the amounts measured in the metabolomics profiling.

Supplementary Figure 3: Extracellular acidification rates and oxygen consumption rates
in ESCs upon Δ9-THC and 2-DG exposure.
(A) Traces were plotted for the extracellular acidification rate (ECAR) measurements in ESCs
exposed to 100nM of Δ9-THC and 10mM of 2-DG, as indicated, and normalized to the protein
content. The oligomycin injection time is indicated by an arrow and allows to differentiate basal
glycolytic rate from maximal glycolytic rate (when mitochondria are inhibited). **(B)** Traces were
plotted for the oxygen consumption rate (OCR) measurements in ESCs exposed to 100nM of Δ9-
THC and 10mM of 2-DG, as indicated, and normalized to the protein content. The oligomycin,
FCCP and AntimycinA/Rotenone injection times are indicated by arrows and allow to differentiate

basal respiration from ATP-coupled respiration and maximal respiratory capacity. FCCP: Carbonyl
cyanide-p-trifluoromethoxyphenylhydrazone.

REFERENCES

1. U.N. Office on Drugs and Crime. 2022 World Drug Report. (2022).
2. Substance Abuse and Mental Health Services Administration. Key substance use and mental health indicators in the United States: Results from the 2019 National Survey on Drug Use and Health. (2020).
3. Mennis, J., Stahler, G. J. & Mason, M. J. Cannabis Legalization and the Decline of Cannabis Use Disorder (CUD) Treatment Utilization in the US. Curr. Addict. Reports (2023). doi:10.1007/s40429-022-00461-4
4. Chabarria, K. C. et al. Marijuana use and its effects in pregnancy. Am. J. Obstet. Gynecol. **215**, 506.e1-506.e7 (2016).
5. Volkow, N. D., Han, B., Compton, W. M. & McCance-Katz, E. F. Self-reported Medical and Nonmedical Cannabis Use Among Pregnant Women in the United States. JAMA - J. Am. Med. Assoc. **322**, 167–169 (2019).
6. Young-Wolff, K. C. et al. Among Pregnant Females in California From 2009 – 2016. JAMA - J. Am. Med. Assoc. **318**, 2490–2491 (2018).
7. Andre, C. M., Hausman, J. F. & Guerriero, G. Cannabis sativa: The plant of the thousand and one molecules. Front. Plant Sci. **7**, 1–17 (2016).
8. Chandra, S. et al. New trends in cannabis potency in USA and Europe during the last decade (2008–2017). Eur. Arch. Psychiatry Clin. Neurosci. **269**, 5–15 (2019).
9. Pacher, P., Kogan, N. M. & Mechoulam, R. Beyond THC and Endocannabinoids. Annu. Rev. Pharmacol. Toxicol. **60**, (2020).
10. Watson, C. T. et al. Genome-Wide DNA Methylation Profiling Reveals Epigenetic Changes in the Rat Nucleus Accumbens Associated with Cross-Generational Effects of Adolescent THC Exposure. Neuropsychopharmacology **40**, 2993–3005 (2015).
11. Prini, P. et al. Adolescent THC exposure in female rats leads to cognitive deficits through a mechanism involving chromatin modifications in the prefrontal cortex. J. Psychiatry Neurosci. **43**, 170082 (2017).
12. Bénard, G. et al. Mitochondrial CB1 receptors regulate neuronal energy metabolism. Nat. Neurosci. **15**, 558–564 (2012).
13. Szutorisz, H. & Hurd, Y. L. High times for cannabis: Epigenetic imprint and its legacy on brain and behavior. Neurosci. Biobehav. Rev. **85**, 93–101 (2018).
14. Lo, J. O., Hedges, J. C. & Girardi, G. Impact of cannabinoids on pregnancy, reproductive health, and offspring outcomes. Am. J. Obstet. Gynecol. **227**, 571–581 (2022).
15. Paria, B. C., Das, S. K. & Dey, S. K. The preimplantation mouse embryo is a target for cannabinoid ligand-receptor signaling. Proc. Natl. Acad. Sci. U. S. A. **92**, 9460–9464 (1995).
16. Murphy, S. K. et al. Cannabinoid exposure and altered DNA methylation in rat and human sperm. Epigenetics **13**, 1208–1221 (2018).
17. Osborne, A. J. et al. Genome-wide DNA methylation analysis of heavy cannabis exposure in a New Zealand longitudinal cohort. Transl. Psychiatry **10**, 1–10 (2020).
18. Schrott, R. & Murphy, S. K. Cannabis use and the sperm epigenome: a budding concern? Environ. Epigenetics **6**, 1–10 (2020).
19. Smith, A., Kaufman, F., Sandy, M. S. & Cardenas, A. Cannabis Exposure During Critical Windows of Development: Epigenetic and Molecular Pathways Implicated in Neuropsychiatric Disease. Curr. Environ. Heal. Reports **7**, 325–342 (2020).
20. Verdikt, R. & Allard, P. Metabolo-epigenetics: the interplay of metabolism and epigenetics during early germ cells development. Biol. Reprod. **105**, 616–624 (2021).
21. Kurimoto, K. & Saitou, M. Epigenome regulation during germ cell specification and

development from pluripotent stem cells. Curr. Opin. Genet. Dev. **52**, 57–64 (2018).

22.   Verdikt, R., Armstrong, A. A. & Allard, P. Transgenerational inheritance and its modulation
by environmental cues. in Current Topics in Developmental Biology 1–46 (Elsevier Inc.,
2022). doi:10.1016/bs.ctdb.2022.10.002

23.   Hayashi, Y. et al. Distinct requirements for energy metabolism in mouse primordial germ
cells and their reprogramming to embryonic germ cells. Proc. Natl. Acad. Sci. U. S. A.
114, 8289–8294 (2017).

24.   Lunt, S. Y. & Vander Heiden, M. G. Aerobic glycolysis: Meeting the metabolic
requirements of cell proliferation. Annu. Rev. Cell Dev. Biol. **27**, 441–464 (2011).

25.   Hayashi, K., Ohta, H., Kurimoto, K., Aramaki, S. & Saitou, M. Reconstitution of the Mouse
Germ Cell Specification Pathway in Culture by Pluripotent Stem Cells. Cell **146**, 519–532
(2011).

26.   Hunault, C. C., Mensinga, T. T., Leenders, M. E. C. & Meulenbelt, J. Delta-9-
tetrahydrocannabinol (THC) serum concentrations and pharmacological effects in males
after smoking a combination of tobacco and cannabis containing up to 69 mg THC.
Psychopharmacology (Berl). **201**, 171–181 (2008).

27.   Pacifici, R. et al. THC and CBD concentrations in blood , oral fluid and urine following a
single and repeated administration of " light cannabis ". Clin. Chem. Lab. Med. 1–8
(2019).

28.   Fuchs Weizman, N. et al. Cannabis alters epigenetic integrity and endocannabinoid
signalling in the human follicular niche. Hum. Reprod. **00**, 1–10 (2021).

29.   Galve-Roperh, I. et al. Cannabinoid receptor signaling in progenitor/stem cell proliferation
and differentiation. Prog. Lipid Res. **52**, 633–650 (2013).

30.   Takeda, S. et al. Δ9-Tetrahydrocannabinol enhances MCF-7 cell proliferation via
cannabinoid receptor-independent signaling. Toxicology **245**, 141–146 (2008).

31.   Yang, X., Bam, M., Nagarkatti, P. S. & Nagarkatti, M. RNA-seq analysis of δ9-
tetrahydrocannabinol-treated T cells reveals altered gene expression profiles that regulate
immune response and cell proliferation. J. Biol. Chem. **291**, 15460–15472 (2016).

32.   Preet, A., Ganju, R. K. & Groopman, J. E. Δ9-Tetrahydrocannabinol inhibits epithelial
growth factor-induced lung cancer cell migration in vitro as well as its growth and
metastasis in vivo. Oncogene **27**, 339–346 (2008).

33.   Rinaldi-Carmona, M. et al. SR141716A, a potent and selective antagonist of the brain
cannabinoid receptor. FEBS Lett. **350**, 240–244 (1994).

34.   Bartova, A. & Birmingham, M. K. Effect of Δ9 tetrahydrocannabinol on mitochondrial
NADH oxidase activity. J. Biol. Chem. **251**, 5002–5006 (1976).

35.   Locasale, J. W. & Cantley, L. C. Metabolic flux and the regulation of mammalian cell
growth. Cell Metab. **14**, 443–451 (2011).

36.   Pendergrass, W., Wolf, N. & Pool, M. Efficacy of MitoTracker Green™ and CMXRosamine
to measure changes in mitochondrial membrane potentials in living cells and tissues.
Cytom. Part A **61**, 162–169 (2004).

37.   Barban, S. & Schulze, H. O. The Effects of 2-Deoxyglucose on the Growth and
Metabolism of Cultured Human Cells. J. Biol. Chem. **236**, 1887–1890 (1961).

38.   Pang, Z. et al. Using MetaboAnalyst 5.0 for LC–HRMS spectra processing, multi-omics
integration and covariate adjustment of global metabolomics data. Nat. Protoc. **17**, 1735–
1761 (2022).

39.   Lojpur, T. et al. Δ9-Tetrahydrocannabinol leads to endoplasmic reticulum stress and
mitochondrial dysfunction in human BeWo trophoblasts. Reprod. Toxicol. **87**, 21–31
(2019).

40.   Miller, M. L. et al. Adolescent exposure to Δ 9 -tetrahydrocannabinol alters the
transcriptional trajectory and dendritic architecture of prefrontal pyramidal neurons. Mol.
Psychiatry **24**, 588–600 (2019).

41.   Ohinata, Y., Sano, M., Shigeta, M., Yamanaka, K. & Saitou, M. A comprehensive, non-
invasive visualization of primordial germ cell development in mice by the Prdm1-mVenus
and Dppa3-ECFP double transgenic reporter. Reproduction **136**, 503–514 (2008).

42.   Tempany, J. C., Zhou, J. H. S., Hodgkin, P. D. & Bryant, V. L. Superior properties of
CellTrace Yellow^{TM} as a division tracking dye for human and murine lymphocytes.
Immunol. Cell Biol. **96**, 149–159 (2018).

43.   Hocaoglu, H., Wang, L., Yang, M., Yue, S. & Sieber, M. Heritable shifts in redox
metabolites during mitochondrial quiescence reprogramme progeny metabolism. Nat.
Metab. **3**, 1259–1274 (2021).

44.   Kittler, J. T. et al. Large-scale analysis of gene expression changes during acute and
chronic exposure to Δ9-THC in rats. Physiol. Genomics **2000**, 175–185 (2000).

45.   Gómez, M., Hernández, M. & Fernández-Ruiz, J. The activation of cannabinoid receptors
during early postnatal development reduces the expression of cell adhesion molecule L1
in the rat brain. Brain Res. **1145**, 48–55 (2007).

46.   Keimpema, E., MacKie, K. & Harkany, T. Molecular model of cannabis sensitivity in
developing neuronal circuits. Trends Pharmacol. Sci. **32**, 551–561 (2011).

47.   Okamura, D., Kimura, T., Nakano, T. & Matsui, Y. Cadherin-mediated cell interaction
regulates germ cell determination in mice. Development **130**, 6423–6430 (2003).

48.   Barton, L. J., LeBlanc, M. G. & Lehmann, R. Finding their way: themes in germ cell
migration. Curr. Opin. Cell Biol. **42**, 128–137 (2016).

49.   Andrews, S. FastQC: a quality control tool for high throughput sequence data. (2010).
Available at: https://www.bioinformatics.babraham.ac.uk/projects/fastqc/.

50.   Dobin, A. et al. STAR: Ultrafast universal RNA-seq aligner. Bioinformatics **29**, 15–21
(2013).

51.   García-Alcalde, F. et al. Qualimap: Evaluating next-generation sequencing alignment
data. Bioinformatics **28**, 2678–2679 (2012).

52.   Anders, S., Pyl, P. T. & Huber, W. HTSeq-A Python framework to work with high-
throughput sequencing data. Bioinformatics **31**, 166–169 (2015).

53.   Love, M. I., Huber, W. & Anders, S. Moderated estimation of fold change and dispersion
for RNA-seq data with DESeq2. Genome Biol. **15**, 1–21 (2014).

54.   Zhang, Y., Parmigiani, G. & Johnson, W. E. ComBat-seq: Batch effect adjustment for
RNA-seq count data. NAR Genomics Bioinforma. **2**, 1–10 (2020).

55.   Raudvere, U. et al. G:Profiler: A web server for functional enrichment analysis and
conversions of gene lists (2019 update). Nucleic Acids Res. **47**, W191–W198 (2019).

56.   Supek, F., Bošnjak, M., Škunca, N. & Šmuc, T. Revigo summarizes and visualizes long
lists of gene ontology terms. PLoS One **6**, (2011).

57.   Bonnot, T., Gillard, M. & Nagel, D. A Simple Protocol for Informative Visualization of
Enriched Gene Ontology Terms. Bio-Protocol **9**, 1–9 (2019).

930