Using Neural Style Transfer to study the evolution of animal signal design: A case study in an ornamented fish

Yseult Héjja-Brichard[1,2], Kara Million[1], Julien P Renoult[2] *, Tamra C Mendelson[1] *

[1] Department of Biological Sciences, University of Maryland, Baltimore County, Baltimore, Maryland, United States; [2] CEFE, Univ Montpellier, CNRS, EPHE, Univ Paul-Valery Montpellier, Montpellier, Occitanie, France

* denotes equal contribution

Abstract

The sensory drive hypothesis of animal signal evolution describes how animal communication signals and preferences evolve as adaptations to local environments. While classical approaches to testing this hypothesis often focus on preference for one aspect of a signal, deep learning techniques like generative models can create and manipulate stimuli without targeting a specific feature. Here, we used an artificial intelligence technique called neural style transfer to experimentally test preferences for color patterns in a fish. Findings in empirical aesthetics show that humans tend to prefer images with the visual statistics of the environment because the visual system is adapted to process them efficiently, making those images easier to process. Whether this is the case in other species remains to be tested. We therefore manipulated how similar or dissimilar male body patterns were to their habitats using the Neural Style Transfer (NST) algorithm. We predicted that males whose body patterns are more similar to their native habitats will be preferred by conspecifics. Our findings suggest that both males and females are sensitive to habitat congruence in their preferences, but to different extents, requiring additional investigation. Nonetheless, this study demonstrates the potential of artificial intelligence for testing hypotheses about animal communication signals.

Keywords: *neural style transfer - processing bias - sensory drive - mate choice - communication - Etheostoma*

Highlights

- Neural style transfer was used to test preferences for patterns in a colorful fish
- Results trend in support of the processing bias hypothesis, a component of sensory drive
- This study demonstrates the use of generative AI for animal behavioral studies

Introduction

Animals are characterised by a vast array of colours and patterns that evolve due to both natural and sexual selection: animals need to be able to hide from predators and to be visible to potential mates. In both contexts, the habitat plays a crucial role in the evolution of colour patterns. In the case of predation, avoidance may involve background-matching (Merilaita & Lind, 2005) or disruptive colouration (Cuthill et al., 2005; Stevens et al., 2008), two camouflage strategies that rely on matching or contrasting characteristics (colours, patterns) of the visual environment, respectively (see e.g., Cuthill et al., 2016). In the case of mating, for an animal to successfully convey a courtship message, it must be detectable by the receiver, meaning an individual must stand out from its background. However, based on theoretical and empirical results in human visual preferences (Menzel et al., 2015), and correlative studies in animals (Hulse et al., 2020), authors have proposed that sexual signals could also be more attractive if their patterning matches that of the background (Renoult & Mendelson, 2019). This hypothesis remains to be tested empirically in animals.

For attracting a mate, conspicuous signalling can be achieved with a broad diversity of colours and patterns, and understanding the drivers of this diversity in sexual signals has been a central focus of evolutionary biology. One of the most influential hypotheses is sensory drive (Endler, 1992; Endler & Basolo, 1998; Seehausen et al., 2008). Based largely on signal detection theory, sensory drive suggests that the need to be optimally conspicuous leads to the evolution of different signals in different environments. Most evidence supporting this hypothesis comes from studies that investigate the link between one feature of an animal signal design, such as colouration, and the corresponding physical characteristic in the environment, like the colour of ambient light. The main prediction of sensory drive is that the detectability of a signal should be favoured to have a higher signal-to-noise ratio. A compelling example is the case of the colourful dewlap of Anolis lizards, which is used for courtship and competition signalling. Anolis species in bright habitats tend to have red or orange dewlaps, while species in darker-lit habitats tend to have white or yellow dewlaps, both maximizing visibility in their respective habitats (Fitch & Hillis, 1984; Fleishman, 1992; Fleishman et al., 2022).

However, to attract a mate, a signal must be more than detected. A robust finding in empirical aesthetics is that the subjective ease with which a signal is processed by the brain is a strong determinant of how attractive that signal will be; this is the fluency theory of aesthetics, which predicts that a stimulus that is easy to process (depending on both the stimulus' intrinsic properties and the observer's perceptual processing of that stimulus) will trigger positive affect for its observer (Fantz, 1957; Munar et al., 2015; Reber et al., 2004; Renoult et al., 2016). In addition, neuroscientists have extensively documented that perceptual systems are adapted to efficiently process natural scenes and their spatial statistics (Olshausen & Field, 1997; Simoncelli & Olshausen, 2001). As a consequence, signal designs that imitate the spatial statistics of natural scenes should be processed more efficiently, because they match the sensitivity, or tuning, of neurons that are adapted to those habitats. Combined with the fluency theory of aesthetics, this prediction suggests that signal designs that imitate the spatial statistics of natural scenes should also generate a pleasant feeling of fluency (Geller et al.,

82 2022) and thus they are likely attractive. Combining the fluency theory of aesthetics with
83 sensory drive, Renoult & Mendelson (2019) proposed the concept of a processing bias and
84 suggested that this mechanism could explain the attractiveness and evolution of complex
85 signal designs in both human and animal communication.
86
One way to test the attractiveness of sexual signals that mimic the patterns of natural habitats
is to use the Fourier slope, which corresponds to the distribution of spatial frequencies (i.e.
luminance distribution) in pattern images. In camouflage studies, the Fourier slope has been
shown to reliably predict the degree of background matching in cuttlefish and octopus (Zylinski
et al., 2011; Josef et al., 2012). Similarly, the Fourier slope can be used to determine whether
background matching plays a role in attractiveness. For instance, jumping spiders prefer
abstract patterns that more closely match the Fourier sope of terrestrial habitats (Hardenbicker
& Tedore, 2023). However, the Fourier slope only captures one aspect of a visual pattern: the
scale invariance in the distribution of luminance contrasts. Other features could be equally or
even more important in explaining how background matching can increase visual
attractiveness.
98
Testing preferences for complex designs and for designs that do not yet exist requires new
approaches. Tools from artificial intelligence and in particular from deep learning represent a
valuable complementary approach. Generative models, such as variational autoencoders
(VAEs; Kingma & Welling, 2019) and generative adversarial networks (GANs, Goodfellow et
al., 2014; 2020) can be used to create or manipulate stimuli and study animal preferences
without the need to target a specific feature. Exploiting the architecture of GANs, (Talas et al.,
2020) developed CamoGAN to study the evolutionary arms race between the camouflage of
a synthetic prey and its predator. Another example is CycleGAN (Zhu et al., 2020). This GAN-
based algorithm can be trained to transform one image into a different category and could be
used to model an intermediate between two phenotypes or between two species to study
conspecific preferences. A complementary method, StyleGAN3 (Karras et al., 2021), can be
used to modify specific features of an image. Another type of generative model is the (latent)
diffusion model (Sohl-Dickstein et al., 2015; Ho et al., 2020; Song & Ermon, 2020). Such a
model can take a sentence as input and turn it into an image (see e.g. (Ramesh et al., 2021;
Nichol et al., 2022) and was made famous with the release of the DALL-E application by
OpenAI. Diffusion models can thus be used to synthesize rich and controlled images to test
specific hypotheses about signal preferences. Although they have interesting properties and
can generate high-quality images, such models require a large number of images and
extensive computer power to be trained, and they have a somewhat complex architecture.
The use of APIs such as DALL-E can be employed as an alternative but offers little control to
the user.
120
One tool of artificial intelligence that allows a high level of control over stimulus generation is
the Neural Style Transfer (NST) algorithm developed by Gatys and colleagues (2015, 2017).
The algorithm requires two input images, one for content and one for style, and aims to render
the content image in the style of the style image. Using the architecture of a convolutional
neural network (CNN), the NST algorithm extracts the content, namely the global arrangement
of an image, such as the objects that constitute it, and the style, computing the Gram matrix
of an image which corresponds to the texture information and local variations of luminance, of

two images, respectively. The algorithm creates a third image that combines the content of
one image with the style of the other (Figure 1). Importantly, the user can determine how much
weight to put on the style or on the content to modulate the output image as desired. Moreover,
the algorithm only requires two input images and no extra training on the user end, and several
accurate implementations are ready to use along with tutorials.
133
Here we used NST to study visual preferences in the Rainbow Darter (*Etheostoma*
caeruleum), a colourful freshwater fish species native to eastern North America. The Rainbow
Darter inhabits creeks and small to medium rivers with gravel substrate (Kuehne & Barbour,
2014; Page & Burr, 2011). As for most species in this genus, Rainbow Darters are
characterized by visually complex nuptial color patterns, which are expressed only by males,
suggesting that these patterns are evolving due to sexual selection. Analyzing the body
patterns of ten species in the same genus, including the Rainbow Darter, Hulse and
colleagues (2020) found a correlation between the Fourier slope of male body patterns and
that of their respective habitats. Female patterns did not show this correlation, suggesting that
only the nuptial ornamentation mimics the spatial statistics of the species' habitat.
144
We, therefore, aimed to test whether the correlation between the spatial statistics of male
luminance patterns and their preferred habitats could be driven by sexual selection. The
processing bias hypothesis stipulates that patterns that mimic the spatial statistics of natural
scenes are easier to process and thus preferred by conspecifics (Renoult & Mendelson, 2019).
Using the style transfer technique, we manipulated the luminance patterns of video animations
of male Rainbow Darters to be either more or less similar to their typical gravel habitat ("native
habitat" hereafter). This method allowed us to investigate the pattern of a signal holistically,
that is, without arbitrarily focusing on an isolated characteristic (e.g. the Fourier slope) as is
more traditionally done. We then tested the preference of male and female darters for images
of conspecifics that do and do not exhibit the "style" of their native habitat. We predicted that
animations of males whose body patterns are the most similar to their habitat's statistics, thus
being easier to process, will be preferred over animations that are less similar to their habitats.
157

Methods

Fish collection and maintenance

We collected male and female Rainbow Darters from Israel Creek (39°29'35.1492' N,
77°19'28.0488'' W) and Tuscarora Creek (39°15'6.948'' N, 77°28'48.288'' W) in Frederick
County, Maryland, between March 15th and April 21st 2022. Fish were transported to the
University of Maryland, Baltimore County in aerated coolers and individually housed in a
recirculating aquarium system (Aquaneering, Inc.). Tanks contained no substrate to avoid
confounding effects but were enriched with a small artificial tree and a bag of crushed coral.
Fish were provided natural lighting matching their photoperiod with a constant temperature of
19°C and fed a diet of frozen bloodworms every other day. Finally, males were visually isolated
from each other to avoid aggressive behaviours and from females to prevent courtship
behaviours but females could see each other. Permission to collect fish was granted by the
Maryland Department of Natural Resources Fishing and Boating Services Scientific Collection

(permit number SCP202246A). Housing and experimental procedures were approved by the
Institutional Animal Care and Use Committee of the University of Maryland, Baltimore County
(protocol 623).

Stimuli creation

We used the Neural Style Transfer (NST), a CNN-based algorithm developed by Gatys and
colleagues (2017), to alter the spatial statistics of photographs of male Rainbow Darters.
Photographs were taken of males from Middle Fork Red River, Trammel Fork, and Salt Fork
in Powell County, KY, Allen County, KY, and Vermilion County, IL, respectively, according to
methods in Hulse et al., (2020), using a Canon EOS 5D Mark IV digital camera under standard
lighting conditions and Zerene Stacker to combine each stack into an image. Habitat videos
were taken using an Ikelite 200DL underwater housing for a Canon EOS 5D Mark IV digital
camera equipped with a Sigma 24 mm f/1.4 lens (see Hulse et al., 2020 for details). Rainbow
Darters are found almost exclusively in stream habitats with abundant gravel or boulder, and
very rarely in open sandy habitats (Page et al. 2011; Kuehne and Barbour 2014). Thus, we
considered images of gravel-bottomed streams to represent the "native" habitat and sandy-
bottomed streams to represent the "foreign" habitat. Notably, sand habitats are also the most
different from gravel habitats when comparing the Fourier slope of freshwater habitat types
(Hulse et al., 2020).
189
A VGG-19 pre-trained network (Simonyan & Zisserman, 2015) with average pooling instead
of max pooling and the L-BFGS algorithm for optimisation was chosen for the NST
architecture. The content information corresponds to the fourth block of feature layers (out of
five) of the CNN while the style information is extracted via a Gram matrix (Figure 1). Briefly,
the gram matrix G is the inner product of vectorised (flattened) feature maps F extracted for
each layer separately, such that $G = F^T F$, and corresponds to the degree of correlations
between the feature maps, that is, how often those features co-occur in an image. Such
features can be orientations, variations in brightness, patterns, or shapes, and they
correspond to the texture of an image.
199
200 To render one content image in the style of another image, the algorithm computes and
201 minimises the total loss of the generated image, which corresponds to the weighted sum of
202 the content loss and the style loss such that
203
204 $$\mathcal{L}_{\text{total}} = \alpha\ \mathcal{L}_{\text{content}} + \beta\ \mathcal{L}_{\text{style}},$$
205
where α and β are the weighting factors for content and style reconstruction, respectively. The
content loss is the squared Euclidean distance between the respective intermediate higher-
level feature representation F of the generated image (x) and content image (x_c) at layer ℓ:
$$\mathcal{L}_{\text{content}} = \sum_{i,j}\ \ (F_l(x) - F_l(x_c))^2_{i,j}.$$
The style function is the sum of the squared Euclidean distance between the gram matrices
of the generated image (x) and the style reference image (x_s) at layer ℓ:
$$\mathcal{L}_{\text{style}} = \sum_l\ \ w_l E_l,$$
with w_l as a scaling factor that equals 1 divided by the number of active layers, and E
calculated as:

215
$$E_l = \frac{1}{4N_l^2}\sum_{i,j} \quad (G_l(x) - G_l(x_s))_{i,j}^2,$$

216
217 with

218
$$G_l(x) = \frac{1}{H_l(x)*W_l(x)}F_l(x)^T F_l(x),$$

219
220 and $H_l(x)*W_l(x)$ corresponding to the height and width of each feature map at each layer.
221 This last formula corresponds to the simplified version G = F^{T}F described above.
222
For our study purposes, a photograph of an original fish and a photograph of the habitat (either
gravel or sand) were fed to the NST algorithm as content and style images, respectively.
Content and style have adaptable weights to improve the final result. We kept them constant
across stimuli, using the recommended values (content weight = 1; style weight = 1e3). For
the style, we only considered the two first blocks of convolutional layers (out of five) as they
tend to represent the local statistics of a given image. This means that the scaling factor w_l
was equal to 0.5 for layers 1 and 2 and to 0 for layers 3 to 5. For the content, we kept the
original parameters and used the feature maps from the convolutional layers of the fourth
block.
232

233
234 *Figure 1. Neural Style Transfer. Panel A: The algorithm produces a new image by minimising the*

The original colours of the fish (i.e. the content image) were put back on the final image, using
the colour control option of the NST (luminance-only transfer), given that color is known to
affect preferences in darter fish (Williams et al., 2013; Williams & Mendelson, 2013). Briefly,
the style transfer manipulates the luminance channel only, by first converting both images
from the BGR space to the YCbCr space, a space that separates luminance information from
colour information. The loss is computed on the luminance channel of that space for both
images. The luminance of the output image is then combined with the CbCr information of the
input style image, so that the output image keeps the colours of the style image instead of the
content image. An example of such output images is shown in Figure 1B.

The PyTorch (Paszke et al., 2019) implementation of the algorithm
(https://pytorch.org/tutorials/advanced/neural_style_tutorial.html) with our modifications, as
well as the detailed procedure used to create the stimuli, are available on a GitHub repository:
https://github.com/yseulthb/NST_fish.

To control for potential biases, we used three images per habitat category (gravel or sand)
and styled five different male individuals. Stylized fish were then animated using Blender
(Blender Online Community, 2018). The path followed by the animated fish reproduced the
path of a live fish, whose video was taken as a representative example of natural motion
(Blender files provided on the GitHub project folder). Animations were displayed in underwater
videos depicting either gravel (species-specific habitat) or sand (foreign habitat). We selected
three different background videos per habitat. For each pair of video animations presented to
test individuals, we matched their luminance values (both in terms of background and
animated fish). Pairs of videos also presented different animated individuals. Beyond that
constraint, all possible combinations of replicates of individual male image, habitat image, and
habitat video were randomized across trials and focal individuals.

In summary, we transferred the style of habitats to the body of male Rainbow Darters (one
habitat image per fish). Animated images of artificially stylized and nonstylized fish were then
presented to live conspecifics, either a male or a female. In addition to manipulating the fish
patterns, we also alternated the video background of the fish animations to determine whether
individuals prefer the style of fish that is more similar to the background against which it is
displayed, regardless of the habitat type (i.e. native vs foreign), i.e., testing a preference for
background matching per se. We compared the amount of time fish spent with the animated
fish exhibiting the native habitat style (and with a foreign habitat style) when shown in a native
habitat vs in a foreign habitat video background. Finally, we explored the influence of style
transfer by comparing preferences for stylized vs nonstylized animated fish.

Dichotomous choice experiments

We ran a series of six experiments to determine whether Rainbow Darters showed a preference for animations stylized with their native habitat statistics. All trials took place during the known breeding season of the species. To avoid habituation, fish were never tested two days in a row, with at least a day between two test sessions. Tested individuals ("focal fish" hereafter) were presented with a pair of stimuli that displayed different animated fish but the same video background. This dichotomous choice experimental paradigm is classically used in the study of mate preferences and has been extensively used in darters to study both male and female mate preferences (Mattson et al., 2020; Roberts et al., 2017; Roberts & Mendelson, 2017). Dichotomous choice tests also show that male darters strongly prefer the nuptial color patterns of other conspecific males, suggesting that male-male competition is also driven by pattern preferences (Williams & Mendelson 2013). The different experimental conditions, their characteristics and sample sizes are summarised in Table 1. In addition to estimating preference for differently styled fish, we estimated in a seventh experiment their preference for their habitat, irrespective of the animated fish (i.e. while showing the same animated, unstyled fish on both sides). This experiment allowed us to test whether individuals have a preference for their native habitats, which is a fundamental assumption of our hypothesis.

Trials took place in a 37.9L tank flanked by two monitors along the short sides displaying the video animations. The long sides of the tank were covered on the inside with nonreflective black plexiglass to inhibit fish from seeing and interacting with their own reflection. Water was systematically changed between trials and tanks were filled up to 75% of their capacity. Trials were video-recorded using an overhead video camera (Logitech). A 12cm wide zone in the middle of the tank was delineated with transparent removable dividers and served as an acclimation area. Association zones were defined as the 5cm-wide area in front of the video animations (see Figure 2). This distance corresponds to the usual courting distance between males and females (darters are pair spawners) and is in the range of an estimated visual acuity for that species (Caves et al., 2017).

To control for experimental side bias, each trial alternated on which side of the tank the stimulus conditions (e.g. gravel-styled or sand-styled) were shown. Most fish took part in all seven trials, while the experiment order was randomised between individuals to control for potential order effects. Furthermore, using five individuals as animated fish combined with the different modification styles limits any recognition bias that could influence our results. Additionally, those five animated fish were paired with each other in all possible combinations to avoid a confounding effect of size difference. It should be noted, however, that fish size difference does not necessarily predict the outcome of male fights in that species (*unpublished experiment*).

For the first five minutes of a trial, focal fish were contained in the acclimation area while opaque dividers blocked the video animations. Those opaque dividers were then removed and the fish remained in the acclimation area for an additional five minutes. This protocol gave the focal fish time to visually assess both sides. Finally, the transparent dividers were removed and the focal fish could freely move in the tank for 15 minutes. We identified the preferred stimulus by comparing the amount of time spent in the two association zones during the first 12 minutes after the transparent dividers were removed. Time spent with a stimulus in a dichotomous mate choice trial is a reliable proxy for preference in freshwater fishes (Aspbury & Basolo, 2002; Brooks & Endler, 2001; Lehtonen & Lindström, 2008). Trial videos were scored using BORIS, an open-source event logging software for video and audio files (Friard & Gamba, 2016), by four persons who were blind to experimental conditions. Three mutually exclusive behaviors were defined: the focal fish was either in the central zone ('center') or in one of the association zones ('left' or 'right'). An event would start when the fish entered one of the zones and would end when it left that zone to enter another zone. At the end of the video, the durations for all three events (i.e., time spent in each zone) were summed up separately and the data entered into a shared, online table. Once all videos were scored, left and right zones were associated to their respective conditions (e.g., sand or gravel). In order to ensure all four scorers reached consensus on association times, interobserver reliability was first tested on six trials. Times reported for the three zones were compared across the observers and estimated to be 90% similar between scorers (i.e. 90% of reported times did not differ by more than half a second from each other). Trials were then assigned evenly across three authors for scoring.

Table 1: Experimental conditions and sample sizes for the animated fish (Experiments 1 to 6) and the habitat (Experiment 7) preference experiments. Grey cells indicate column names.

Experiment	Background video	Style of fish 1	Style of fish 2	Sample size
Experiment 1	native	native	foreign	M: 20 F: 17
Experiment 2	foreign	native	foreign	M: 17 F: 16

Experiment	Fish stimulus	Habitat video 1	Habitat video 2	Sample size
Experiment 3	native	native	nonstylized	M: 19 F: 13
Experiment 4	native	foreign	nonstylized	M: 16 F: 15
Experiment 5	foreign	native	nonstylized	M: 20 F: 17
Experiment 6	foreign	foreign	nonstylized	M: 19 F: 16
Experiment 7	nonstylized	native	foreign	M: 16 F: 13

Statistical analyses

For all seven experiments separately, we first checked for a potential side bias by comparing times spent in the left vs right association zones irrespective of the stimuli shown. We used either t-tests or Mann-Whitney Wilcoxon tests depending on the data distribution.

To estimate habitat or animation preference, we compared the averaged time spent in the association zone for each condition as well as the averaged time spent in the central zone. This latter measure informs us of overall interest in the stimuli. To compare times spent in those three areas and as data followed a normal distribution (as indicated by normality assessing Shapiro-Wilks tests), we ran one-way ANOVAs. If ANOVA results were significant, we ran posthoc tests (Tukey HSD tests) to identify which pairs of stimuli were significantly different. We also report the number of entries in each association zone, a proxy for mobility. Finally, we counted the number of times the focal fish entered the association zones and used chi-square tests to identify any difference. To determine whether preferences might vary depending on the video background type (native or foreign habitat), we ran t-tests with unequal variance between pairs of experiments where the only difference was the background video type (experiment 1 vs 2, experiment 3 vs 5, and experiment 4 vs 6). All analyses are reported for males and females separately. Statistical analyses were performed with the software R (R Core Team, 2021) using the car package (Fox and Weisberg, 2019).

Results

Experiments 1 to 6: Fish preference

We analysed the data of 13 to 17 females and 16 to 20 males (see Table 1) for animation preference. We initially tested a minimum of 15 and a maximum of 20 individuals per experiment. However, individuals that either froze or stayed in the same location without further moving for the entire duration of the experiment were discarded to avoid inflating the time spent in the central area, as these fish provide no information about time spent in the association zones and therefore no information about pattern preferences. We found no side

380　bias for any of the six experiments testing fish preference, neither for females nor males.
381　Detailed test results are provided in the supplementary material.
382
Experiment 1 tested the preference for a fish animation stylized with the statistics of a Rainbow
Darter's native habitat versus a foreign habitat, with a background video of a native habitat
(gravel). According to the processing bias hypothesis, fish should prefer the animation with
the statistics of their habitat, as their visual system processes those more easily. Results were
significant for focal females only ($F(2,48) = 4.75$ $p = 0.013$ for females; $F(2,57) = 1.89$ $p = 0.16$
for males; Fig. 3A). Post-hoc tests found that the amount of time females spent in the
association zones of the animation stylized with the native habitat was significantly greater
than the amount of time spent in the central zone ($t = -3.03$, $p = 0.011$).
391
392　On average, males entered the association zone of the fish stylized with the native habitat 6.5
393　times and of the fish stylized with the foreign habitat 7.8 times. For females, they entered 6.35
394　times and 5.53 times, respectively. No significant difference was found for either group ($X^2 =$
395　0.1182, df = 1, $p = 0.731$ for males; $X^2 = 0.05708$, df = 1, $p = 0.811$ for females).
396
Experiment 2 used the same animation types as Experiment 1 but a background video of a
foreign habitat (sand) instead of a native habitat to explore whether the background style
influences preferences, as in Menzel et al. (2015). Again, the time difference was significant
for females only ($F(2,45) = 10.3$ $p < 0.001$ for females; $F(2,45) = 10.3$ $p < 0.001$ for males; Fig.
3B). Post-hoc tests revealed that females spent more time in the two association zones (native
habitat (NH) style: $t = -4.54$, $p < 0.001$; foreign habitat (FH) style: $t = 2.57$, $p = 0.035$), compared
to the central zone.
There was no significant difference in the number of entries between the two association
zones, for either sex (males: mean NH = 7.1, mean FH = 8.1; $X^2 = 0.07387$, df = 1, p = 0.786:
females: mean RDH = 4.4, mean FH = 4.2; $X^2 = 0.004106$, df = 1, p = 0.949).
407
Experiment 3 compared fish stylized with a native habitat versus a nonstylised (NS) fish,
displayed in front of a video of a native habitat. This experiment aimed to determine whether
the style transfer procedure influences fish's preferences. A preference for the styled
animation, when it corresponds to the native habitat statistics, should remain present when
compared to the natural version of the fish. Results were significant for both males ($F(2,54) =$
13.7 $p < 0.001$) and females ($F(2,36) = 7.94$, $p = 0.0014$; Fig. 3C). Post-hoc tests found that
the amount of time females spent in the association zone of the fish stylized with the native
habitat was significantly greater than the amount of time spent in the central zone ($t = -3.96$, p
< 0.001). Males spent significantly more time in the association zone of the fish stylized with
the native habitat compared to the central zone ($t = -5.17$, $p < 0.001$) and more time in the
association zone of the fish stylized with the native habitat compared to the nonstylized fish (t
$= -3.27$, $p = 0.0052$). Thus males preferred the fish that was stylized with the native habitat
over the natural version of the fish.
421
422　There was no significant difference in the number of entries between the two association
423　zones, for either sex (males: mean NH = 6.3, mean NS = 5.2; $X^2 = 0.07387$, df = 1, p = 0.786:
424　females: mean NH = 4.9, mean NS = 4.4; $X^2 = 0.004106$, df = 1, p = 0.949).
425

Experiment 4 tested a preference for fish stylized with a foreign habitat versus a nonstylized fish, with a background video of a native habitat. A stronger preference for the nonstylized fish (natural phenotype) is expected, as the animation with the foreign habitat statistics should not be better processed by the visual system. Results were significant for both males ($F(2,45)$ = 3.45 p = 0.04) and females ($F(2,42)$ = 6.9, p =0.0026; Fig. 3D). Post-hoc tests revealed that females spent significantly more time in the association zone of the fish stylized with the foreign habitat (t = 3.65, p = 0.0021) and of the nonstylized fish (t = 2.44, p = 0.0487) compared to the central zone. Males spent significantly more time in the association zones of the nonstylized fish compared to the central zone (t = 2.48, p = 0.044).

There was no significant difference in the number of entries between the two association zones, for either sex (males: mean FH = 6.3, mean NS = 6.7; X^2 = 0.07387, df = 1, p = 0.786: females: mean FH = 6.0, mean NS = 6.1; X^2 = 0.004106, df = 1, p = 0.949).

Experiment 5 tested a preference for the animation stylized with the native habitat versus a nonstylized animation, with a background video of a foreign habitat. Similarly to Experiment 3, a preference for the styled animation is expected. Results were significant for both males ($F(2,57)$ = 3.52 p = 0.036) and females ($F(2,48)$ = 10.2, p < 0.001; Fig. 3E). Post-hoc tests revealed that females spent significantly more time with the fish stylized with a native habitat (t = -3.86, p = 0.00099) and with the nonstylized fish (t = 3.97, p = 0.00073) compared to the central zone. Males spent significantly more time in the association zone of the fish stylized with native habitat compared to the central zone (t = -2.43, p = 0.048).

There was no significant difference in the number of entries between the two association zones, for either sex (males: mean NH = 7.8, mean NS = 8.0; X^2 = 0.07387, df = 1, p = 0.786: females: mean NH = 4.1, mean NS = 3.9; X^2 = 0.004106, df = 1, p = 0.949).

Experiment 6 tested a preference for the animation stylized with a foreign habitat versus a nonstylized animation, with a background video of a foreign habitat. Similarly to Experiment 4, a preference for the nonstylized fish (natural phenotype) is expected. Results were significant for females only ($F(2,45)$ = 7.24 p = 0.0019 for females; $F(2,54)$ = 3.01, p = 0.058 for males, Fig. 3F). Post-hoc tests revealed that the amount of time females spent in the association zone of the animation stylized with a foreign habitat was significantly greater than the amount of time spent in the central zone (t = 3.81, p = 0.0013).

There was no significant difference in the number of entries between the two association zones, for either sex (males: mean FH = 8.1, mean NS = 8.7; X^2 = 0.07387, df = 1, p = 0.786: females: mean FH = 6.6, mean NS = 5.9; X^2 = 0.004106, df = 1, p = 0.949).

Background comparisons

A potential influence of the background video on preferences was investigated by showing animations in front of two different backgrounds: one that corresponded to the native habitat of the focal fish and one that showed a foreign habitat. There was no significant difference in time spent in either association zone between experiments that differ only in the type of video background (native or foreign habitat), that is, between experiment 1 and experiment 2 (native habitat style: t = -0.481, df = 33.505, p = 0.6337; foreign habitat style: t = 0.662, df = 34.942, p = 0.512), between experiment 3 and experiment 5 (native habitat style: t = 0.66602, df =

25.447, *p* = 0.5114; nonstylized: *t* = -0.854, df = 26.827, *p* = 0.4006) and between experiment 4 and experiment 6 (foreign habitat style: *t* = 0.242, df = 24.599, *p* = 0.8105; nonstylized: *t* = 0.194, df = 29.886, *p* = 0.8475). This suggests that background videos did not influence the preferences of focal individuals.

Experiment 7. Habitat preference

A processing bias hypothesis assumes that animals prefer the visual statistics of their own habitat. To estimate preference for native habitat (NH) versus foreign habitat (FH), we analysed the data of 13 females (out of 15 tested) and 16 males (out of 20 tested). Discarded individuals either froze or stayed in the same location without further moving for the entire duration of the experiment. T-tests for the remaining males and females showed no difference between times spent on either side of the tank (females: *t* = -1.0095, df = 23.992, *p* = 0.3228; males: *t* = 0.79333, df = 29.307, *p* = 0.434); thus we did not exclude any individuals due to side bias.

Results were significant for both males (*F*(2,45) = 6.35 *p* = 0.0037) and females (*F*(2,36) = 17.3, *p* < 0.001; Fig 3G). Tukey's HSD Test for multiple comparisons found that the amount of time males spent in the association zone of their native habitat was significantly greater than the amount of time spent in the central zone (*t* = -3.54, *p* = 0.0026). Females spent significantly more time associating with their habitat compared to the central zone (*t* = -5.85, *p* <0.001) and more time associating with their habitat compared to the less preferred habitat (*t* = -3.46, *p* = 0.0039). Thus females preferred their "native" habitat over the "foreign" habitat.

There was no significant difference in the number of entries between the two association zones, for either sex (males: mean NH = 6.8, mean FH = 6.5; X² = 0.07387, df = 1, p = 0.786: females: mean NH = 5.4, mean FH = 4.0; X² = 0.004106, df = 1, p = 0.949).

Table 2: Summary of significant results (2-way ANOVA, *p* < 0.05)

	Significant results	Shown in
Experiment 1 *Native vs foreign,* *native background*	Native zone > central zone	Females
Experiment 2 *Native vs foreign,* *Foreign background*	Native zone > central zone Foreign zone > central zone	Females
Experiment 3 *Native vs nonstylized,* *Native background*	Native zone > central zone Native zone > nonstylized zone	Females and Males Males
Experiment 4 *Foreign vs nonstylized,* *Native background*	Native zone > central zone Nonstylized zone > central zone	Females Females and Males
Experiment 5	Native zone > central zone	Females and Males

Native vs nonstylized, foreign background	Nonstylized zone > central zone	Females
Experiment 6 *Foreign vs nonstylized, foreign background*	Foreign zone > central zone	Females
Experiment 7 *Nonstylized, native vs foreign background*	Native zone > central zone Native zone > foreign zone	Females and Males Females

Figure 3: Time (ms) spent in the association or central zones for the seven experiments. Fish animations stylized to reflect the statistics of the species' native habitat (gravel) are represented in blue, a foreign habitat (sand) in orange, nonstylized animations in green, and the central zone in grey. Asterisks and horizontal bars indicate statistically significant differences between conditions.

Discussion

Using the neural style transfer algorithm, we modified animations of male Rainbow Darter fish
to reflect the style of (1) their preferred habitat (gravel) and (2) a less preferred ("foreign")
habitat (sand). We then investigated female and male preferences for stylized animations as
well as nonstylized animation. We found that, across all experiments, females spent more time
in the association zones than in the central zone, but they showed no significant preference
for any one pattern over another (native habitat style, foreign habitat style, and nonstylized).
In contrast, males spent more time in the association zones in only half of the experiments
(Experiments 3, 4, 5); however, when they showed a significant preference, males preferred
animated fish stylized to reflect their preferred habitat over the nonstylized animated fish,
spending more time in the corresponding association zone than in the two other zones
(Experiment 3). As for habitat preferences alone (Experiment 7), females preferred the video
of their habitat, spending more time in the corresponding association zone than in the two
other zones. Males, on the other hand, only preferred their habitat over the central zone; no
difference was found between times spent associating with videos of gravel versus sand
habitat.
526

Our finding that males showed a preference for the fish stylized with the native habitat over a
nonstylized fish (Experiment 3) provides some support for sensory drive and the processing
bias hypothesis (Endler, 1992; Endler & Basolo, 1998; Seehausen et al., 2008; Renoult &
Mendelson, 2019). Sensory drive predicts that animals will have habitat-specific preferences,
and processing bias suggests that those preferences are determined by redundant patterns
in the environment. As such, processing bias is an example of a pre-existing preference that
derives from adaptations of the receiver's sensory and cognitive systems in other contexts,
which can be exploited by signalers to increase mating success (Ryan & Cummings, 2013;
Fuller & Endler, 2018). For example, pioneering studies on the Túngara frog revealed that
females' preferences for chucks, a specific component of males' calls, are influenced by the
tuning properties of their auditory system (see e.g. Ryan et al., 1990). In that case, sexual
selection favored the evolution of male traits, here the chuck calls, that exploit pre-existing
female biases emerging from the property of their auditory system, shaped by their
environment (Ryan & Keddy-Hector, 1992). The majority of examples of pre-existing
preferences that have been explained in the context of sensory drive have been limited mostly
to simple features. Thus, pre-existing preferences for complex patterns, such as the
preference for artificial crests in Australian finches (Burley & Symanski, 1998), are often
assumed to emerge incidentally or are not well explained. The processing bias hypothesis
provides an explanation for pre-existing preferences for complex patterns, and the attraction
of males in our study towards fish stylized with the statistics of their native habitat supports
that hypothesis. However, it should be noted that this preference does not appear consistently
across experiments (e.g., no such effect in experiment 5), nor was it present in females.
Nonetheless, other results are at least consistent with expectations of processing bias, with
fish spending significantly more time with the native habitat style as compared to the neutral
zone. One solution may be to increase the sample size. Studies in humans that demonstrate
a pre-existing preference for the spatial statistics of natural scenes typically have sample sizes

of up to 50 participants (see e.g. Menzel et al., 2015; Spehar et al., 2015). Reaching such a
sample size with wild animals is a challenge, but not an insurmountable one.
555
Across our experiments, we tested the preferences of both males and females, as both sexes
can play a role in sexual selection, and previous research suggests that male and female
preferences for a given trait can be either similar or opposed (Candolin, 2004; Qvarnstrom &
Forsgren, 1998; Wong & Candolin, 2005). In darters, males compete physically and vigorously
with other males for access to females, defending their territory, challenging courting males,
and mate guarding, and both sexes are choosy when it comes to selecting a mate. Females
and males prefer the same conspecific male color and pattern in at least two species of
darters, *Etheostoma barrenense* and *E. zonale* (Williams & Mendelson, 2013), but whether
they share a similar preference for the style of the pattern might depend on its function. For
example, a male pattern might signal dominance to other males and predict the likelihood of
winning a fight with that individual (see e.g. Guerrera et al., 2022); a male might prefer to
interact longer with a male against which it would win. Therefore, it would be interesting to
compare the pattern of the observing fish to the displayed fish to determine whether certain
visual characteristics could predict the preferences of focal individuals. For females, the same
pattern could indicate a male's fitness or paternal quality; females would presumably spend
more time associating with a higher-quality male. Only males in our study showed a significant
preference for animations with the style of their preferred habitat over a foreign habitat;
however, our results do not allow us to determine whether male and female preferences are
significantly aligned or opposed. Whether intra- or intersexual processes play a more
important role in the design of darter sexual signals therefore remains an open and interesting
question.
577
In addition to manipulating the fish patterns, we also alternated the video background of the
animations to determine whether individuals prefer the style of fish that is more similar to the
background against which it is displayed, regardless of habitat type (i.e. native vs foreign).
This design allowed us to test for a preference for background matching per se. Comparing
experiments that differed only in background video (Experiment 1 vs Experiment 2,
Experiment 3 vs Experiment 5, and Experiment 4 vs Experiment 6; Figure 4) reveals no effect
of background on association time. For each relevant comparison, there were no significant
differences in the amount of time spent in each of the three zones, indicating no effect of the
background, as that is the only difference between each pair of experiments. This result was
somewhat surprising, as a previous study in humans that manipulated the Fourier slope of the
background against which faces were presented found that the background slope influenced
the attractiveness of the faces (Menzel et al 2015). Sample sizes were higher in the human
study than in ours, but an additional explanation is that the Fourier slope has a different effect
on preferences than does the style of an image.
592
593 **Limitations and future directions**
594
The camera viewpoint is an important factor to consider when choosing style images. We
chose wide-angle images from the perspective of a benthic fish, keeping the camera close to
the stream floor. Wide-angle images that include an entire visual scene are usually the go-to
when testing the adaptation of the visual system to natural scene statistics (e.g. Gupta et al.,

2023). Indeed, the efficient coding hypothesis states that the visual system is adapted to such
statistics (Olshausen & Field, 1996; Simoncelli & Olshausen, 2001), but it does not necessarily
specify other aspects of visual perception, such as visual acuity or the breadth of the visual
field of a given species. However, at least one study, of mice, was able to predict visual
perception with cameras filming the environment from the perspective of those animals (Qiu
et al., 2021). We assume that similar retinal adaptation takes place in darters and thus we
used a similar approach, using images taken from the perspective of the animal.
606
The scale of the style images might also affect relevant statistics. For instance, habitat images
are not scale invariant; thus, the scale of the image affects which image properties are
transferred (e.g., more or less fine-grained details, more or less repetitive patterns) and thus
the preference of the observer fish. Experimenting with different scales would enable direct
testing of how scale changes the properties of the output image and the preferences of the
fish. In addition, investigating the image properties of different stylized images could help
disentangle which properties influence fish preferences, from their Fourier slope (as
mentioned on Figure S3) to their contrast value. In the current study, only the mean luminance
(i.e. brightness) difference was minimized between pairs of stimuli; thus, it remains possible
that the observer fish noticed differences in an unknown image property that influenced its
visual preference. Follow-up studies should consider aspects such as image scaling and visual
acuity.
619
In our experiments, we kept the color of the animated fish constant. One important future
direction will be to test the interaction between colors and patterns while modulating spatial
statistics of signal designs and ask how this influences preferences. Another future research
direction will be to focus on specific body parts of the fish rather than the body as a whole, to
assess whether some parts are more susceptible to sexual selection. For instance, Gumm &
Mendelson (2011) found no correlation between body colors and fin colors of several darter
species, suggesting that patterns on these different body parts may be evolving independently.
These differences make sense since dorsal fins are often erected for courtship or male-male
fights (see e.g. Johnston, 1994) and concealed at rest. Fin patterns therefore may be evolving
due to sexual selection alone and be less constrained by camouflage. Conveniently, the most
recent implementation of the NST algorithm (Gatys et al., 2017) has two control options that
make it useful for addressing such questions: spatial and color controls. The spatial control
lets the user predefine on which part of an image to transfer the style. This is useful for
modifying subparts of a body and identifying potential differences in their effect on visual
preferences or communication. With color control, the model allows the user to either retain
the original color of the content image (i.e., to "transfer" only achromatic information) or to
transfer both chromatic and achromatic information. This option is convenient for testing the
influence of color on visual preferences.
638
Other methods could also be used instead of the current implementation. For instance, the
Laplacian Pyramid method (Burt & Adelson, 1983; Sunkavalli et al., 2010) enables the transfer
of texture information (the "style" of an image) from one image to another and represents a
less computationally intensive alternative. We also recommend exploring alternative and ever-
evolving Neural Style Transfer methods. For instance, the comparative work of Wright &

Ommer (2022) could guide the reader in determining which method would lead to better results
both in terms of qualitative output and required computational power.
646
Our implementation, based on a PyTorch tutorial, makes it convenient to style any new image
even for an audience not familiar with the language or the concept of neural style transfer. The
code, in the form of a notebook, only requires a few adaptations, such as changing the weights
put on the style and the content image depending on the desired output. Comments in the
code should help to run the script easily even without prior knowledge of deep learning
algorithms. Finally, because the algorithm is based on the VGG-19 architecture pretrained on
ImageNet, a generalist database containing images of diverse objects, people and animals,
no additional training is required for the algorithm, making it more straightforward to implement
for any new pair of images.
656
Our study and suggestions for future research demonstrate how the style transfer algorithm
and other AI-based techniques can be used to study the evolution of signal design. For
instance, the ability of style transfer via the Gram matrix to separate texture information makes
it particularly suitable to study camouflage, as highlighted in Hulse et al. (2022). Specifically,
using style transfer, one can distinguish between two camouflage strategies: background
matching, which depends on texture (i.e., patterns and variations in color or shading) to blend
in with the environment, and disruptive coloration, which hinders individual recognition by
breaking up a visual outline, thereby creating visual confusion. Using a similar approach in
fish, Hulse et al. (2022) compared the gram matrices of images of male and female darters to
habitat images of different categories to determine whether fish species were more visually
similar to the habitat in which they occur as compared to habitats in which they are typically
not found. Female patterns were found more similar to their habitats than were nuptial male
patterns, highlighting a potential trade-off between sexual and natural selection (i.e.,
camouflage) for males during the mating season. Interestingly, that study also found
differences between network layers, revealing how those can be used to estimate the
influence of spatial scales on the studied trait.

Conclusion

Using artificial intelligence methods to test the effect of low-level spatial patterning on the
visual preferences of fishes, our study provides one of the first attempts to use rapidly
advancing AI techniques to directly query animal behavior. Our results bring no clear support
to the prediction that male nuptial patterns matching the visual statistics of their habitat are
preferred over patterns matching a foreign habitat or a nonstylized fish. However, the
hypothesis of a processing bias is also not rejected. Indeed, the significant results we obtained
are either consistent with or support the hypothesis. In particular males significantly preferred
animations whose patterns match the native habitat over nonstylized animations. Importantly,
our study demonstrates how artificial neural networks can be used to test hypotheses about
animal signal evolution, by creating or manipulating stimuli without the need to target a specific
feature. Indeed, such generative models will allow us to move toward a more holistic
perceptual representation of complex signals, paving the way to exciting new questions.

Data Availability

The PyTorch implementation of the algorithm with our modifications, as well as the detailed procedure used to create the stimuli, are available on a GitHub repository: https://github.com/yseulthb/NST_fish, as recommended by Huettmann & Arhonditsis (2023).

Acknowledgements

We would like to thank the students of the DART lab who helped with trials and video scoring. This work was supported by the National Science Foundation grant NSF IOS 2026334.

Supplementary material

Supplementary table: Assessing potential side bias in the seven experiments

	Males	Females
Experiment 1	t = 1.765, df = 38, p = 0.0855	t = -1.758, df = 31.81, p = 0.0883
Experiment 2	t = 1.952, df = 26.82, p = 0.0614	W = 94, p = 0.207
Experiment 3	t = 0.9881, df = 35.97, p = 0.33	t = 0.07073, df = 23.98, p = 0.944
Experiment 4	t = -0.3773, df = 26.28, p = 0.709	t = -1.626, df = 27.84, p = 0.115
Experiment 5	t = 1.071, df = 37.92, p = 0.291	W = 135, p = 0.76
Experiment 6	t = 0.2342, df = 34.42, p = 0.816	t = 0.834, df = 29.92, p = 0.411
Experiment 7	t = 0.7933, df = 29.31, p = 0.434	t = -1.01, df = 23.99, p = 0.323

Figure S1: Fish images used as content images

Fish 1

Fish 6

Fish 3

Fish 7

Fish 4

704
705
706
707 Figure S2: Underwater images used as style images

Gravel 1

Sand 1

Gravel 3

Sand 2

Gravel 4

Sand 3

Gravel 5

Sand 4

Figure S3: Stylized fish

Text S1: Fourier slope of stylized images

Although pairs of stylized fish that were presented to fish observers were luminance-matched,
they differed in their Fourier slopes. Fourier slopes of fish stylized with the gravel habitat
pictures (-2.74 ±0.08) were significantly different from Fourier slopes of fish stylized with the
sand habitat pictures (-3.00 ±0.10; t-test: t = -8.819, p < 0.0001).

References

https://doi.org/10.1086/720267

Fox J, Weisberg S (2019). An R Companion to Applied Regression, Third edition. Sage, Thousand Oaks CA. https://socialsciences.mcmaster.ca/jfox/Books/Companion/

Aspbury, A. S., & Basolo, A. L. (2002). Repeatable female preferences, mating order and mating success in the poeciliid fish, Heterandria formosa. *Behavioral Ecology and Sociobiology*, *51*(3), 238–244. https://doi.org/10.1007/s00265-001-0443-1

Brooks, R., & Endler, J. A. (2001). Female Guppies Agree to Differ: Phenotypic and Genetic Variation in Mate-Choice Behavior and the Consequences for Sexual Selection. *Evolution*, *55*(8), 1644–1655. https://doi.org/10.1111/j.0014-3820.2001.tb00684.x

Burley, N. T., & Symanski, Richard. (1998). "A Taste for the Beautiful": Latent Aesthetic Mate Preferences for White Crests in Two Species of Australian Grassfinches. *The American Naturalist*, *152*(6), 792–802. https://doi.org/10.1086/286209

Burt, P., & Adelson, E. (1983). The Laplacian Pyramid as a Compact Image Code. *IEEE Transactions on Communications*, *31*(4), 532–540. IEEE Transactions on Communications. https://doi.org/10.1109/TCOM.1983.1095851

Candolin, U. (2004). Opposing Selection on a Sexually Dimorphic Trait through Female Choice and Male Competition in a Water Boatman. *Evolution*, *58*(8), 1861–1864.

Caves, E. M., Sutton, T. T., & Johnsen, S. (2017). Visual acuity in ray-finned fishes correlates with eye size and habitat. *Journal of Experimental Biology*, *220*(9), 1586–1596. https://doi.org/10.1242/jeb.151183

Cuthill, I. C., Sanghera, N. S., Penacchio, O., Lovell, P. G., Ruxton, G. D., & Harris, J. M. (2016). Optimizing countershading camouflage. *Proceedings of the National Academy of Sciences*, *113*(46), 13093–13097. https://doi.org/10.1073/pnas.1611589113

Cuthill, I. C., Stevens, M., Sheppard, J., Maddocks, T., Párraga, C. A., & Troscianko, T. S.

(2005). Disruptive coloration and background pattern matching. *Nature*, *434*(7029),

72–74. https://doi.org/10.1038/nature03312

Endler, J. A. (1992). Signals, Signal Conditions, and the Direction of Evolution. *The*

American Naturalist, *139*, S125–S153. https://doi.org/10.1086/285308

Endler, J. A., & Basolo, A. L. (1998). Sensory ecology, receiver biases and sexual selection.

Trends in Ecology & Evolution, *13*(10), 415–420. https://doi.org/10.1016/s0169-

5347(98)01471-2

Fantz, R. L. (1957). Form preferences in newly hatched chicks. *Journal of Comparative and*

Physiological Psychology, *50*(5), 422–430. https://doi.org/10.1037/h0044973

Fitch, H. S., & Hillis, D. M. (1984). The Anolis Dewlap: Interspecific Variability and

Morphological Associations with Habitat. *Copeia*, *1984*(2), 315–323.

https://doi.org/10.2307/1445187

Fleishman, L. J. (1992). The Influence of the Sensory System and the Environment on

Motion Patterns in the Visual Displays of Anoline Lizards and Other Vertebrates. *The*

American Naturalist, *139*, S36–S61. https://doi.org/10.1086/285304

Fleishman, L. J., Perez-Martinez, C. A., & Leal, M. (2022). Can Sensory Drive Explain the

Evolution of Visual Signal Diversity in Terrestrial Species? A Test with Anolis Lizards.

The American Naturalist, 000–000. https://doi.org/10.1086/720267

Friard, O., & Gamba, M. (2016). BORIS: A free, versatile open-source event-logging

software for video/audio coding and live observations. *Methods in Ecology and*

Evolution, *7*(11), 1325–1330. https://doi.org/10.1111/2041-210X.12584

Fuller, R. C., & Endler, J. A. (2018). A perspective on sensory drive. *Current Zoology*, *64*(4),

465–470. https://doi.org/10.1093/cz/zoy052

Gatys, L. A., Ecker, A. S., & Bethge, M. (2015). A Neural Algorithm of Artistic Style.

arXiv:1508.06576 [Cs, q-Bio]. http://arxiv.org/abs/1508.06576

Gatys, L. A., Ecker, A. S., Bethge, M., Hertzmann, A., & Shechtman, E. (2017). *Controlling*

Perceptual Factors in Neural Style Transfer. 3985–3993.

https://openaccess.thecvf.com/content_cvpr_2017/html/Gatys_Controlling_Perceptual_Factors_CVPR_2017_paper.html

Geller, H. A., Bartho, R., Thömmes, K., & Redies, C. (2022). Statistical image properties predict aesthetic ratings in abstract paintings created by neural style transfer. *Frontiers in Neuroscience*, *16*, 999720. https://doi.org/10.3389/fnins.2022.999720

Godfrey, D., Lythgoe, J. N., & RUMBALL, D. A. (1987). Zebra stripes and tiger stripes: The spatial frequency distribution of the pattern compared to that of the background is significant in display and crypsis. *Biological Journal of the Linnean Society*, *32*(4), 427–433. https://doi.org/10.1111/j.1095-8312.1987.tb00442.x

Goodfellow, I. J., Pouget-Abadie, J., Mirza, M., Xu, B., Warde-Farley, D., Ozair, S., Courville, A., & Bengio, Y. (2014). Generative Adversarial Networks. *arXiv:1406.2661 [Cs, Stat]*. http://arxiv.org/abs/1406.2661

Goodfellow, I., Pouget-Abadie, J., Mirza, M., Xu, B., Warde-Farley, D., Ozair, S., Courville, A., & Bengio, Y. (2020). Generative adversarial networks. *Communications of the ACM*, *63*(11), 139–144. https://doi.org/10.1145/3422622

Guerrera, A. G., Daniel, M. J., & Hughes, K. A. (2022). Black and orange coloration predict success during male–male competition in the guppy. *Behavioral Ecology*, *33*(6), 1196–1206. https://doi.org/10.1093/beheco/arac093

Gumm, J. M., & Mendelson, T. C. (2011). The evolution of multi-component visual signals in darters (genus Etheostoma). *Current Zoology*, *57*(2), 125–139. https://doi.org/10.1093/czoolo/51.2.125

Gupta, D., Młynarski, W., Sumser, A., Symonova, O., Svatoň, J., & Joesch, M. (2023). Panoramic visual statistics shape retina-wide organization of receptive fields. *Nature Neuroscience*, *26*(4), 606–614. https://doi.org/10.1038/s41593-023-01280-0

Hardenbicker, M.-C., & Tedore, C. (2023). Peacock spiders prefer image statistics of average natural scenes over those of male ornamentation. *Behavioral Ecology*, arad042. https://doi.org/10.1093/beheco/arad042

Ho, J., Jain, A., & Abbeel, P. (2020). *Denoising Diffusion Probabilistic Models*

(arXiv:2006.11239). arXiv. https://doi.org/10.48550/arXiv.2006.11239

Huettmann, F., & Arhonditsis, G. (2023). Towards an ecological informatics scholarship that

is reflective, repeatable, transparent, and sharable! *Ecological Informatics*, *76*,

102132. https://doi.org/10.1016/j.ecoinf.2023.102132

Hulse, S. V., Renoult, J. P., & Mendelson, T. C. (2020). Sexual signaling pattern correlates

with habitat pattern in visually ornamented fishes. *Nature Communications*, *11*(1),

Article 1. https://doi.org/10.1038/s41467-020-16389-0

Hulse, S. V., Renoult, J. P., & Mendelson, T. C. (2022). Using deep neural networks to

model similarity between visual patterns: Application to fish sexual signals. *Ecological*

Informatics, *67*, 101486. https://doi.org/10.1016/j.ecoinf.2021.101486

Johnston, C. E. (1994). Spawning Behavior of the Goldstripe Darter (Etheostoma parvipinne

Gilbert and Swain) (Percidae). *Copeia*, *1994*(3), 823–825.

https://doi.org/10.2307/1447204

Josef, N., Amodio, P., Fiorito, G., & Shashar, N. (2012). Camouflaging in a Complex

Environment—Octopuses Use Specific Features of Their Surroundings for

Background Matching. *PLOS ONE*, *7*(5), e37579.

https://doi.org/10.1371/journal.pone.0037579

Karras, T., Aittala, M., Laine, S., Härkönen, E., Hellsten, J., Lehtinen, J., & Aila, T. (2021).

Alias-Free Generative Adversarial Networks. *Advances in Neural Information*

Processing Systems, *34*, 852–863.

https://proceedings.neurips.cc/paper/2021/hash/076ccd93ad68be51f23707988e9349

06-Abstract.html

Kingma, D. P., & Welling, M. (2019). An Introduction to Variational Autoencoders.

Foundations and Trends® in Machine Learning, *12*(4), 307–392.

https://doi.org/10.1561/2200000056

Kuehne, R. A., & Barbour, R. W. (2014). *The American Darters* (Reprint edition). University

Press of Kentucky.

Lehtonen, T. K., & Lindström, K. (2008). Repeatability of mating preferences in the sand

goby. *Animal Behaviour*, *75*(1), 55–61. https://doi.org/10.1016/j.anbehav.2007.04.011

Mattson, C. L., Roberts, N. S., & Mendelson, T. C. (2020). Male preference for conspecific

females depends on male size in the splendid darter, Etheostoma barrenense.

Animal Behaviour, *165*, 89–96. https://doi.org/10.1016/j.anbehav.2020.04.022

Menzel, C., Hayn-Leichsenring, G. U., Langner, O., Wiese, H., & Redies, C. (2015). Fourier

Power Spectrum Characteristics of Face Photographs: Attractiveness Perception

Depends on Low-Level Image Properties. *PLOS ONE*, *10*(4), e0122801.

https://doi.org/10.1371/journal.pone.0122801

Merilaita, S., & Lind, J. (2005). Background-matching and disruptive coloration, and the

evolution of cryptic coloration. *Proceedings of the Royal Society B: Biological*

Sciences, *272*(1563), 665–670. https://doi.org/10.1098/rspb.2004.3000

Munar, E., Gómez-Puerto, G., Call, J., & Nadal, M. (2015). Common Visual Preference for

Curved Contours in Humans and Great Apes. *PLOS ONE*, *10*(11), e0141106.

https://doi.org/10.1371/journal.pone.0141106

Nichol, A., Dhariwal, P., Ramesh, A., Shyam, P., Mishkin, P., McGrew, B., Sutskever, I., &

Chen, M. (2022). *GLIDE: Towards Photorealistic Image Generation and Editing with*

Text-Guided Diffusion Models (arXiv:2112.10741). arXiv.

https://doi.org/10.48550/arXiv.2112.10741

Olshausen, B. A., & Field, D. J. (1996). Emergence of simple-cell receptive field properties

by learning a sparse code for natural images. *Nature*, *381*(6583), 607–609.

https://doi.org/10.1038/381607a0

Olshausen, B. A., & Field, D. J. (1997). Sparse coding with an overcomplete basis set: A

strategy employed by V1? *Vision Research*, *37*(23), 3311–3325.

https://doi.org/10.1016/S0042-6989(97)00169-7

Page, L. M., & Burr, B. M. (2011). *Peterson field guide to freshwater fishes of North America*

north of Mexico (2nd ed). Houghton Mifflin Harcourt.

Paszke, A., Gross, S., Massa, F., Lerer, A., Bradbury, J., Chanan, G., Killeen, T., Lin, Z.,

Gimelshein, N., Antiga, L., Desmaison, A., Köpf, A., Yang, E., DeVito, Z., Raison, M.,

Tejani, A., Chilamkurthy, S., Steiner, B., Fang, L., … Chintala, S. (2019). *PyTorch:*

An Imperative Style, High-Performance Deep Learning Library (arXiv:1912.01703).

arXiv. https://doi.org/10.48550/arXiv.1912.01703

Qiu, Y., Zhao, Z., Klindt, D., Kautzky, M., Szatko, K. P., Schaeffel, F., Rifai, K., Franke, K.,

Busse, L., & Euler, T. (2021). Natural environment statistics in the upper and lower

visual field are reflected in mouse retinal specializations. *Current Biology*, *31*(15),

3233-3247.e6. https://doi.org/10.1016/j.cub.2021.05.017

Qvarnstrom, A., & Forsgren, E. (1998). Should females prefer dominant males? *Trends in*

Ecology & Evolution.

Ramesh, A., Pavlov, M., Goh, G., Gray, S., Voss, C., Radford, A., Chen, M., & Sutskever, I.

(2021). *Zero-Shot Text-to-Image Generation* (arXiv:2102.12092). arXiv.

http://arxiv.org/abs/2102.12092

Reber, R., Schwarz, N., & Winkielman, P. (2004). Processing Fluency and Aesthetic

Pleasure: Is Beauty in the Perceiver's Processing Experience?: *Personality and*

Social Psychology Review. https://doi.org/10.1207/s15327957pspr0804_3

Renoult, J. P., Bovet, J., & Raymond, M. (2016). Beauty is in the efficient coding of the

beholder. *Royal Society Open Science*, *3*(3), 160027.

https://doi.org/10.1098/rsos.160027

Renoult, J. P., & Mendelson, T. C. (2019a). Processing bias: Extending sensory drive to

include efficacy and efficiency in information processing. *Proceedings of the Royal*

Society B: Biological Sciences, *286*(1900), 20190165.

https://doi.org/10.1098/rspb.2019.0165

Renoult, J. P., & Mendelson, T. C. (2019b). Processing bias: Extending sensory drive to

include efficacy and efficiency in information processing. *Proceedings of the Royal*

880 *Society B: Biological Sciences, 286*(1900), 20190165.

881 https://doi.org/10.1098/rspb.2019.0165

882 Roberts, N. S., Gumm, J. M., & Mendelson, T. C. (2017). Darter (Percidae: Etheostoma)

883 species differ in their response to video stimuli. *Animal Behaviour, 131*, 107–114.

884 https://doi.org/10.1016/j.anbehav.2017.07.013

885 Roberts, N. S., & Mendelson, T. C. (2017). Male mate choice contributes to behavioural

886 isolation in sexually dimorphic fish with traditional sex roles. *Animal Behaviour, 130*,

887 1–7. https://doi.org/10.1016/j.anbehav.2017.06.005

888 Ryan, M. J., & Cummings, M. E. (2013). Perceptual Biases and Mate Choice. *Annual*

889 *Review of Ecology, Evolution, and Systematics, 44*(1), 437–459.

890 https://doi.org/10.1146/annurev-ecolsys-110512-135901

891 Ryan, M. J., Fox, J. H., Wilczynski, W., & Rand, A. S. (1990). Sexual selection for sensory

892 exploitation in the frog Physalaemus pustulosus. *Nature, 343*(6253), Article 6253.

893 https://doi.org/10.1038/343066a0

894 Ryan, M. J., & Keddy-Hector, A. (1992). Directional Patterns of Female Mate Choice and the

895 Role of Sensory Biases. *The American Naturalist, 139*, S4–S35.

896 https://doi.org/10.1086/285303

897 Seehausen, O., Terai, Y., Magalhaes, I. S., Carleton, K. L., Mrosso, H. D. J., Miyagi, R., van

898 der Sluijs, I., Schneider, M. V., Maan, M. E., Tachida, H., Imai, H., & Okada, N.

899 (2008). Speciation through sensory drive in cichlid fish. *Nature, 455*(7213), Article

900 7213. https://doi.org/10.1038/nature07285

901 Simoncelli, E., & Olshausen, B. (2001). Natural image statistics and neural representation.

902 *Annual Review of Neuroscience, 24*(1), 1193–1216.

903 https://doi.org/10.1146/annurev.neuro.24.1.1193

904 Simonyan, K., & Zisserman, A. (2015). Very Deep Convolutional Networks for Large-Scale

905 Image Recognition. *arXiv:1409.1556 [Cs]*. http://arxiv.org/abs/1409.1556

906 Sohl-Dickstein, J., Weiss, E. A., Maheswaranathan, N., & Ganguli, S. (2015). *Deep*

Unsupervised Learning using Nonequilibrium Thermodynamics (arXiv:1503.03585).

arXiv. https://doi.org/10.48550/arXiv.1503.03585

Song, Y., & Ermon, S. (2020). *Generative Modeling by Estimating Gradients of the Data*

Distribution (arXiv:1907.05600). arXiv. https://doi.org/10.48550/arXiv.1907.05600

Spehar, B., Wong, S., van de Klundert, S., Lui, J., Clifford, C., & Taylor, R. (2015). Beauty

and the beholder: The role of visual sensitivity in visual preference. *Frontiers in*

Human Neuroscience, *9*.

https://www.frontiersin.org/articles/10.3389/fnhum.2015.00514

Stevens, M., Winney, I. S., Cantor, A., & Graham, J. (2008). Outline and surface disruption

in animal camouflage. *Proceedings of the Royal Society B: Biological Sciences*,

276(1657), 781–786. https://doi.org/10.1098/rspb.2008.1450

Sunkavalli, K., Johnson, M. K., Matusik, W., & Pfister, H. (2010). Multi-scale image

harmonization. *ACM Trans. Graph.*, *29*(4), 125:1-125:10.

https://doi.org/10.1145/1778765.1778862

Talas, L., Fennell, J. G., Kjernsmo, K., Cuthill, I. C., Scott-Samuel, N. E., & Baddeley, R. J.

(2020). CamoGAN: Evolving optimum camouflage with Generative Adversarial

Networks. *Methods in Ecology and Evolution*, *11*(2), 240–247.

https://doi.org/10.1111/2041-210X.13334

Williams, T. H., Gumm, J. M., & Mendelson, T. C. (2013). Sexual selection acting on a

speciation trait in darters (Percidae: Etheostoma). *Behavioral Ecology*, *24*(6), 1407–

1414. https://doi.org/10.1093/beheco/art080

Williams, T. H., & Mendelson, T. C. (2013). Male and female responses to species-specific

coloration in darters (Percidae: Etheostoma). *Animal Behaviour*, *85*(6), 1251–1259.

https://doi.org/10.1016/j.anbehav.2013.03.012

Wong, B. B. M., & Candolin, U. (2005). How is female mate choice affected by male

competition? *Biological Reviews*, *80*(4), 559–571.

https://doi.org/10.1017/S1464793105006809

Wright, M., & Ommer, B. (2022). ArtFID: Quantitative Evaluation of Neural Style Transfer. In

B. Andres, F. Bernard, D. Cremers, S. Frintrop, B. Goldlücke, & I. Ihrke (Eds.),

Pattern Recognition (pp. 560–576). Springer International Publishing.

https://doi.org/10.1007/978-3-031-16788-1_34

Zhu, J.-Y., Park, T., Isola, P., & Efros, A. A. (2020). Unpaired Image-to-Image Translation

using Cycle-Consistent Adversarial Networks. *arXiv:1703.10593 [Cs]*.

http://arxiv.org/abs/1703.10593

Zylinski, S., How, M. J., Osorio, D., Hanlon, R. T., & Marshall, N. J. (2011). To Be Seen or to

Hide: Visual Characteristics of Body Patterns for Camouflage and Communication in

the Australian Giant Cuttlefish Sepia apama. *The American Naturalist*, *177*(5), 681–

690. https://doi.org/10.1086/659626

Community, B. O. (2018). *Blender - a 3D modelling and rendering package*. Stichting

Blender Foundation, Amsterdam. Retrieved from http://www.blender.org