Ubiquitous, B_{12}-dependent virioplankton utilizing ribonucleotide triphosphate reductase demonstrate interseasonal dynamics and associate with a diverse range of bacterial hosts in the pelagic ocean

Ling-Yi Wu[1], Gonçalo J. Piedade[2,3], Ryan M. Moore[4], Amelia O. Harrison[4], Ana M. Martins[3], Kay D. Bidle[5], Shawn W. Polson[4], Eric Sakowski[6], Jozef I. Nissimov[7], Jacob T. Dums[4,8], Barbra D. Ferrell[4], K. Eric Wommack[4,*]

[1] Theoretical Biology and Bioinformatics, Science4Life, Utrecht University, Padualaan 8, Utrecht, 3584 CH, the Netherlands

[2] Department of Marine Microbiology and Biogeochemistry, NIOZ Royal Netherlands Institute for Sea Research, 1797 SZ t'Horntje, The Netherlands

[3] Department of Oceanography and Fisheries and Ocean Sciences Institute-OKEANOS, University of the Azores, 9901-862 Horta, Faial, Azores, Portugal

[4] Delaware Biotechnology Institute, University of Delaware, 590 Avenue 1743, Newark, DE, 19713, United States

[5] Department of Marine and Coastal Sciences, Rutgers University, 71 Dudley Rd., New Brunswick, NJ, 08901, United States

[6] Johns Hopkins University, Department of Earth Sciences, Baltimore MD, United States

[7] Department of Biology, University of Waterloo, 200 University Ave. West, Waterloo, Ontario, N2L 3G1, Canada

[8] Biotechnology Program, North Carolina State University, 2800 Faucette Dr. Raleigh, NC, 27695, United States

Keywords: oceanic carbon cycle, compositional data, metagenomic amplicons, phage-host interactions, virus phylogeny/oligotrophic microbial community, long-read sequencing

* To whom correspondence should be addressed. Tel: +01 302-831-4362; Fax: +44 000 0000000; Email: wommack@udel.edu

ABSTRACT

Through infection and lysis of their coexisting bacterial hosts, viruses impact the biogeochemical

cycles sustaining globally significant pelagic oceanic ecosystems. Currently, little is known of the

ecological interactions between lytic viruses and their bacterial hosts underlying these biogeochemical

impacts at ecosystem scales. This study focused on populations of lytic viruses carrying the B_{12}-

dependent Class II monomeric ribonucleotide reductase (RNR) gene, ribonucleotide triphosphate

reductase (RTPR), documenting seasonal changes in pelagic virioplankton and bacterioplankton using

amplicon sequences of RTPR and the 16S rRNA gene, respectively. Amplicon sequence libraries were

analyzed using compositional data analysis tools that account for the compositional nature of these data.

Both virio- and bacterioplankton communities responded to environmental changes typically seen across

seasonal cycles as well as shorter term upwelling–downwelling events. Defining RTPR-carrying viral

populations according to major phylogenetic clades proved a more robust means of exploring

virioplankton ecology than operational taxonomic units defined by percent sequence homology.

Virioplankton RTPR populations showed positive associations with a broad phylogenetic diversity of

bacterioplankton including dominant taxa within pelagic oceanic ecosystems such as *Prochlorococcus*

and SAR11. Temporal changes in RTPR-virioplankton, occurring as both free viruses and within infected

cells, indicated possible viral–host pairs undergoing sustained infection and lysis cycles throughout the

seasonal study. Phylogenetic relationships inferred from RTPR sequences mirrored ecological patterns

in virio- and bacterioplankton populations demonstrating possible genome to phenome associations for

an essential viral replication gene.

INTRODUCTION

The subtropical pelagic ocean gyres are extensive, coherent regions that collectively represent the

largest ecosystems on Earth, accounting for 40% of the earth's surface [1]. Circulation within the upper

kilometer of subtropical gyres is primarily wind driven [2]. Due to their size, the horizontal and vertical

motion of water within this upper layer of the gyres plays an outsized role in regulating global nutrient

cycles and the size of the atmospheric carbon pool. Both of these issues are critical concerns for

understanding anthropogenic impacts on fisheries, oceanic general circulation, and global climate.

Ultimately, the activities of and the interactions between oceanic microbial communities—composed of

protists, phytoplankton, bacterioplankton, and virioplankton—regulate the flow of carbon and nutrient

elements within the pelagic ocean gyres [3,4]. Truly appreciating how oceans will respond to environmental

changes brought on by anthropogenic activities requires detailed, mechanistic understanding of oceanic

microbial communities [5–8]. This study focused on specific interactions between important and ubiquitous

populations of lytic viruses [9,10] and bacterioplankton, the most abundant microbial host group within the

low nutrient ecosystems that dominate much of the pelagic ocean.

Viral cell lysis is among the microbial community interactions most directly responsible for the

conversion of cellular biomass into dissolved and particulate organic carbon [11]. This process is one of the

major top-down controls on bacterioplankton productivity, responsible for between 10 and 40% of daily

bacterioplankton mortality in the ocean [12]. Upon lysis, cell contents not consumed by viral replication are

released, contributing to dissolved nutrient pools within marine ecosystems. Bacterioplankton cell lysis

comprises the largest known carbon flux process between biomass and the dissolved organic matter

(DOM) pool amounting to ~150 gigatons C/yr [6,13]. The conversion of biomass to DOM from viral lysis is

five times greater than that performed by other biological mechanisms such as grazing and programmed

cell death [6,13]. This flux from biomass to DOM sustains nutrient cycles, providing essential growth

substrates for bacterioplankton and phytoplankton productivity. The availability and concentrations of

different nutrients within the DOM pool comprise the bottom-up controls on bacterioplankton productivity

[6,13,14]. Besides influencing the composition, diversity, and productivity of marine bacterial communities

through top-down and bottom-up controls, viruses can influence bacterial physiology, improving fitness in

the face of environmental changes [15,16] by altering metabolic pathways [17,18] through horizontal gene

transfer; and through the expression of metabolic genes carried within viral genomes during viral infection

[19,20]. However, the behavior of individual virioplankton populations within ecosystems and their detailed

interactions with bacterioplankton host populations remain largely unknown. This study examined both

virioplankton and bacterioplankton dynamics within the North Atlantic Subtropical Gyre across time with

the objective of observing interactions at the community and population levels.

For some time, microbial ecologists have investigated bacterial populations and communities utilizing

16S rRNA gene amplicon sequences, a universal gene marker for bacterial diversity [21]. Due to their

polyphyletic origins, viruses do not have a universally conserved marker gene that could be similarly used

for tracing viral evolutionary history and community ecology investigations. Nevertheless, there are a few

widespread genes exhibiting robust evolutionary histories that can be used as markers for viral

community ecology studies within natural environments. Genes such as the DNA polymerase A gene

(encoding DNA polymerase I) [22,23], terminase gp20 gene [24–27], major capsid protein gp23 gene of T4-like

phages [28–30], and ribonucleotide reductase (RNR) genes [9] have all informed studies assessing the

diversity of abundant viral populations. This study used amplicons of the ribonucleotide triphosphate

reductase (RTPR) gene for following the dynamics of virioplankton populations.

RNRs are the only known enzymes capable of reducing ribonucleotides to deoxyribonucleotides,

essential substrates for DNA synthesis [31,32]. RNRs are ancient enzymes that were essential to the

emergence of DNA [33]. As a consequence, RNR genes are under rigorous evolutionary selection

pressure, and among the most abundantly identified genes in marine virome libraries [34]. Importantly,

RNR genes are present in tailed phages within the class Caudoviricetes and have been identified within

the genomes of viruses infecting hosts within all three domains of life. Generally, lytic viruses carry RNR

genes, which ensures a steady supply of deoxyribonucleotides for DNA synthesis [35]. Therefore, RNRs

are most commonly observed within the genomes of lytic phages.

RNRs are biologically informative as this group of enzymes consists of three physiological classes

according to reactivity with oxygen. Class I RNRs are oxygen-dependent. Class II RNRs are oxygen-

independent and rely upon an adenosylcobalamin (vitamin B_{12}) cofactor for ribonucleotide reduction.

Class II RNRs are further divided according to structure and substrate utilization. Monomeric Class II

RTPR reduces ribonucleotide triphosphate to deoxyribonucleotide triphosphate (dNTPs), the substrate for

DNA synthesis. Dimeric Class II reduces ribonucleotide diphosphate to deoxyribonucleotide diphosphate

thus requiring a dinucleotide kinase for the final step in forming dNTPs. Class III RNRs are sensitive to

oxygen. As such, Class I and Class II RNRs, not Class III RNRs, are commonly present within

virioplankton communities found in oxygenated ocean water [10,32]. This study leveraged the predominance

of RTPR genes within viruses for examining the ecological dynamics of virioplankton populations. RTPR

and 16S rRNA amplicon sequence data were analyzed against a backdrop of microbial and

oceanographic observations using the latest compositional data analysis (CoDA) tools [36,37] [38,39] [40,41] for

assessing how lytic, B_{12}-dependent viral populations responded to seasonal dynamic changes within the

euphotic zone of a subtropical oceanic gyre, a region critically important within the global carbon cycle.

MATERIALS AND METHODS

Study Site and Sample Collection

Seawater samples were collected from a fixed station at South of Channel Faial-Pico islands (SCFP;

38°23'11.67"N, 28°35'6.14"W), located in the northeastern Atlantic, 6.4 nautical miles off São Mateus

(Pico Island) in the Azores archipelago (ICES subdivision Xa2). The station is representative of a larger

open ocean ecosystem, the Mid-Atlantic Ridge ecoregion, corresponding to the International Council for

the Exploration of the Seas (ICES) areas X and XII, which includes a high proportion of high seas [42]. The

Azores are located along the northern boundary of the North Atlantic Subtropical Gyre, which constitutes

a barrier between superficial cold waters of Nordic provenance and warmer waters of the gyre. Due to

the influence of large scale currents such as the Gulf Stream, the North Atlantic Current, and the Azores

Current, the islands are mostly exposed to the transport of warm waters of equatorial and tropical origin.

These currents result in high salinity, sharp horizontal temperature gradients, and high sea surface

temperatures (on average 15–16 °C in winter and 22–24 °C in summer). These physical conditions result

in low nutrient regimes, and correspond with low ecosystem productivity [43]. Nevertheless, frontal regions

and underwater features (seamounts, eddies, upwelling areas) can enhance local productivity [44,45].

Water samples were collected in acid-rinsed (0.1 M HCl) and seawater-flushed carboys between 9:30

and 11:00 local time from 0, 5, 25, 50, and 75 m depths during seven sampling events on May 12th, June

8th, June 22nd, July 11th, July 27th, August 5th, and September 8th of 2016. Surface seawater (0 m

depth) was collected by directly submerging a carboy. Seawater at other depths was collected using a

2.5 L Niskin bottle and subsequently transferred to a 5 L carboy. Seawater samples were transported

inside a light-tight cooler and laboratory-processed within 2 h of collection. Subsamples for flow

cytometric enumeration of phytoplankton were fixed immediately using glutaraldehyde (0.5% final

concentration). Flow cytometry samples were transported on ice (1–2 h) and stored at -80 °C until further

processing.

Water column conductivity, temperature, and depth (CTD) data was collected down to 150 m using a

CTD instrument (MIDAS Valeport, Totnes, United Kingdom) (**Supplementary Table S1)**. Salinity

(unitless, on the Practical Salinity Scale PSS-78) was expressed as a function of conductivity. Density

(kg/m³) was expressed as a function of pressure. CTD profiles were visualized using the cmocean colour

palettes [46] on Ocean Data View 5.3.0 [47] (https://odv.awi.de). Temperature and salinity values between

sampling days were interpolated using the weighted average grid method [47]. Temperature, salinity, and

density were determined by averaging all values 0.5 m above and below the targeted sampling depths.

Equipment malfunction prevented CTD data collection on September 8th, thus, temperature, salinity, and

density metadata were estimated from the nearest time, pixel, and depth from the Marine Copernicus

Global Ocean Physics model [48] (GLOBAL_ANALYSISFORECAST_PHY_CPL_001_015,

http://marine.copernicus.eu/). Average values for chlorophyll a (mg/m^3), sea surface temperature (SST,

°C), particulate organic carbon (POC, mg/m^3), and photosynthetically active radiation (PAR, Einstein/m^{2}d)

data in 2016 were obtained from the NASA Ocean Color website (http://oceancolor.gsfc.nasa.gov/) at

1 km resolution. Satellite data was mapped using SeaDAS 7 [49] for the 9 km^2 area surrounding sampling

station SCFP.

Phytoplankton Enumeration using Flow Cytometry

Phytoplankton cells were enumerated using a BD Influx Mariner 209s Flow Cytometer (BD

Biosciences, San Jose, CA) equipped with a 488 nm argon laser and 525 ± 15, 542 ± 27, 585 ± 40, and

692 ± 40 nm emission filters with a nozzle size of 70 μm. After thawing and gentle vortexing, samples

were run for 2 min [50]. A blank consisting of 0.22 μm filtered Milli-Q water was run before and after

counting. The flow rate was determined by running Milli-Q water for 2 min and dividing the mass

difference (determined with a scale) by time; mass was converted to volume using the density of 1 kg/L.

Phytoplankton cells were identified based on chlorophyll fluorescence (692 ± 40 nm) from 488 nm

excitation (**Supplementary Table S1**).

Virio- and Bacterioplankton Biomass Collection

The cellular (>0.22 μm) and subcellular (0.02–0.22 μm) fractions, biologically defined in this study as

bacterioplankton and virioplankton, respectively, were separated through two consecutive syringe

filtrations using 0.22 and 0.02 μm filters. All tubing and containers were acid-washed and deionized

water-rinsed prior to sample filtration. The bacterioplankton fraction was collected by filtering 5 L of

seawater through a 0.22 μm pore-size Sterivex filter (Merck Millipore, Darmstadt, Germany) under a

vacuum pressure system linked to a filtration ramp (Merck Millipore, Burlington, MA). The sample

collection flask was rinsed twice using the first 200 mL of 0.22 μm filtrate. Subsequently, the virioplankton

fraction was collected by filtering 1 L of 0.22 μm filtrate through a 0.02 μm pore size Anotop filter

(Whatman, Maidstone, United Kingdom) [51,52] connected to a 60 mL syringe, using a caulk gun which

generated a steady pressure on the syringe. Both 0.22 μm and 0.02 μm filters with bacterio- or

virioplankton biomass were parafilm sealed and stored at -80 °C until DNA extraction.

DNA Extraction from the Virioplankton Fraction and Tag-Encoded RTPR Amplicon Sequencing

In brief, DNA was extracted from the 0.02 μm Anotop filters (virioplankton fraction) and quantified

using previously reported protocols [52]. RTPR PCR amplicons of ~750 bp were obtained using

virioplankton template DNA and degenerate primers designed using sequence alignments derived from

uncultivated virus populations observed within marine viromes. After magnetic bead purification,

amplicon molecules were tagged with a unique barcode sequence using ligation. Purified, barcoded

amplicons were then subjected to a limited cycle PCR with barcode sequences as template targets.

Enriched barcoded amplicons from each sample were then purified, pooled in equal proportions, and

sequenced using a PacBio RSII sequencer (Pacific Biosciences, Menlo Park, CA). Detailed methods

describing DNA extraction, amplification of RTPR target genes, and amplicon sequencing are provided in

Supplementary Methods.

DNA Extraction from the Bacterioplankton Fraction, and 16S rRNA Amplicon and Tag-Encoded

RTPR Sequencing

DNA was extracted from the 0.22 μm Sterivex filters (bacterioplankton fraction) using the

phenol/chloroform method [53]. The V3–V4 hypervariable region of the 16S rRNA gene was PCR-amplified

and sequenced from the bacterioplankton fraction on the Illumina MiSeq (Illumina, San Diego, CA) using

a Nextera XT DNA Library Preparation Kit (Illumina), which exploits a dual-indexing strategy for

multiplexed sequencing [54]. RTPR genes within the bacterioplankton fraction were PCR-amplified,

barcoded, and enriched using the same protocol as described for virioplankton. Detailed methods

describing DNA extraction, amplification of 16S rRNA and RTPR target genes, and amplicon sequencing

are provided in **Supplementary Methods**.

RTPR amplicon quality control

General and detailed bioinformatics analysis pipelines are summarized in **Fig. 1** and **Supplementary**

Fig. S1, respectively. Briefly, RTPR gene amplicon sequences from the virioplankton fraction (PacBio

RSII sequencer) and the bacterioplankton fraction (PacBio Sequel sequencer) were initially screened for

low quality bases and read length. Circular consensus sequence (CCS) reads were generated. Reads

with less than three full passes, less than 98% minimum predicted accuracy, and with a length shorter

than 250 bp or longer than 5,000 bp were excluded. Subsequently, sequence reads were demultiplexed,

and primer and barcode sequences were detected and removed. Reads were translated into predicted

amino acid sequences using a custom frameshift polishing pipeline

(https://github.com/dnasko/frameshift_polisher). Key catalytic residues within translated amplicons were

used for validating sequences as true RTPRs using the Protein Active Site Validation (PASV)

(https://github.com/mooreryan/pasv) pipeline [55]. Detailed methods describing bioinformatic processing of

RTPR amplicon sequences are provided in **Supplementary Methods**.

Virioplankton RTPR 98% OTUs and phylogenetic clades

Quality-controlled virioplankton RTPR amino acid sequences were run through 11 *de novo* operational taxonomic unit (OTUs) clustering assessments varying the percent sequence identity (90– 100% amino acid sequence identity) using the *cluster_fast* command with default settings in USEARCH (version 11.0) [56]. This heuristic process identified 98% OTU clusters as a balance between optimal lumping and splitting of OTUs (**Supplementary Fig. S2**). Virioplankton RTPR 98% OTU amino acid sequences were taxonomically divided into clades using a phylotyping method. OTUs were aligned using MAFFT (version 7.450) with the ginsi method. A phylogenetic tree of virioplankton RTPR 98% OTUs was built using FastTree (version 2.1.11) [57] and visualized using Iroki [58]. Virioplankton RTPR phylogenetic clades were defined manually.

16S quality control and classification

General and detailed analysis pipelines are summarized in **Fig. 1** and **Supplementary Fig. S1.** Demultiplexed, paired-end 16S rRNA reads from the bacterioplankton fraction were imported into QIIME2 (QIIME2 2019.1) [59] for quality filtering and generation of amplicon sequence variants (ASVs) using DADA2 [60]. Forward and reverse reads were truncated at positions 35 and 300 and at positions 35 and 295, respectively. 16S ASVs were taxonomically identified (**Supplementary Table S2**) using the Silva 132 QIIME-compatible release (full length, seven-level taxonomy) [61] as a reference database in QIIME2. Sequences taxonomically assigned to Archaea (22 ASVs) and chloroplasts (230 ASVs) were removed prior to downstream analysis.

For proportionality tests with bacterioplankton 16S ASVs and virioplankton RTPR 98% OTUs, ASVs were restricted to only those that occurred in the 29 samples with virioplankton RTPR amplicons. From that subset of 29 samples, only ASVs with a total count greater than 100 and with at least 10 counts in at least 10 samples were retained for downstream analysis. This avoided the risk of spurious associations based on rare populations observed in few samples.

RTPR and 16S community and population analyses

Virioplankton RTPR 98% OTUs, virioplankton RTPR phylogenetic clades, and bacterioplankton 16S ASVs were used in community and population analyses such as alpha and beta diversity, and population proportionality. The influence of environmental conditions on virio- and bacterioplankton communities were determined based on complete-linkage hierarchical clustering of RTPR 98% OTUs, RTPR phylogenetic clades, or 16S ASV community profiles using the hclust method in R. Tested environmental features include sampling day, depth, temperature, density, current velocity, salinity, nanoeukaryote abundance, picoeukaryote abundance, *Prochlorococcus* abundance, *Synechococcus* abundance, total autotroph abundance, viral abundance, and bacterial abundance. Clustering of virio- and bacterioplankton communities was based on the Aitchison distance between the May 12th 5 m sample and every other sample. Resulting dendrograms were visualized with Iroki. Correlations between virio- or bacterioplankton communities and associated environmental features were tested using the *qiime*

diversity mantel [65] command in QIIME2 (permutation times: 999). Significant correlations were defined as

p <0.05. Associated environmental features that showed significant correlations with sample community

composition were plotted next to the sample community composition dendrogram.

Alpha diversity of the virio- or bacterioplankton communities in each sample was estimated based on

the Shannon index using the R package DivNet (version 0.3.2) [66] and displayed using ggplot2 (version

3.3.0) [67]. Significance of alpha diversity differences was tested using betta function in the R package

DivNet breakaway (version 4.6.14) [68]. Estimated Shannon indices were converted to the effective

number of species (ENS, Hill numbers) [69]. Correlations of ENS estimates based on 98% OTUs or RTPR

phylogenetic clades were tested using the cor.test method in R. Beta diversity was visualized through

principal component analysis (PCA) plots of sample community compositions using the R package

compositions (version 1.40.3) [70]. Significance of beta diversity differences between sampling day or

depth groups was tested using *beta-group-significance* command (permanova) method [71] in QIIME2.

Associations within or between virioplankton RTPR phylogenetic clades and bacterioplankton 16S

ASV populations were explored using proportionality (*rho*, ρ) tests, with -1 < ρ < 1 [72]. Only positive ρ

values reflecting a positive association between two populations were considered, as negative ρ values

cannot be conclusively explained [73]. Only those associations with false discovery rates (FDRs) lower

than 0.05 were retained for further analysis. Proportionality and FDR tests were performed using the R

package propr (version 4.2.8). Associations with FDR <0.05 were extracted and plotted in a heatmap

using ComplexHeatmap [74]. Columns and rows of the heatmap were clustered based on phylogenetic

relations of virio- and bacterioplankton, respectively. Virioplankton RTPR clades were arranged by their

topological order on the 98% OTU phylogenetic tree. Bacterioplankton 16S ASVs were arranged by their

16S phylogenetic placement or association patterns.

`

Virio- and bacterioplankton population analysis using common RTPR 98% OTUs

In 27 of 35 samples, RTPR amplicons were obtained from both virio- and bacterioplankton fractions

yielding 47,997 quality-controlled RTPR amino acid sequences. Quality-controlled virio- and

bacterioplankton RTPR amino acid sequences were run through 11 *de novo* operational taxonomic unit

(OTU) clustering assessments varying the percent sequence identity (90–100% amino acid sequence

identity) using the *cluster_fast* command with default settings in USEARCH. This heuristic process

identified 5,411 98% OTU clusters as a balance between optimal lumping and splitting of OTUs

(**Supplementary Fig. S3**). OTUs with less than ten counts were removed prior to downstream analysis.

The COUNTIF function in Excel was used for binning the remaining 254 RTPR 98% OTUs into those

containing sequences from both fractions, only the virioplankton fraction, or only the bacterioplankton

fraction. RTPR 98% OTU counts were calculated using the SUM function in Excel prior to zero

replacement and clr transformation. An arbitrary clr value of 5 or -5 was assigned to those common 98%

OTUs found exclusively in the virio- or bacterioplankton fraction, respectively. The difference in clr

abundance between fractions of each RTPR 98% OTU was calculated using the subtraction formula in

Excel. Venn diagrams were plotted to show the total clr abundance of each OTU and the distribution of

each OTU (difference in clr abundance) in the virio- and/or bacterioplankton fraction.

Only the August 5th sampling date yielded RTPR amplicons from virio- and bacterioplankton

fractions at all depths. Thus, this date was chosen for assessing depth-resolved distribution patterns of

RTPR virioplankton populations within the virio- and bacterioplankton fractions. Calculation of clr

abundance values for the August 5th OTUs was performed by removing any OTU not present in at least

one August 5th sample, leaving 1,477 of the 5,411 OTUs observed in the combined virio- and

bacterioplankton datasets. Twenty-nine August 5th OTUs contained more than 50 sequences and had

two or fewer zero replacements, and were retained for further analysis. Similarity between the

phylogenetic distance of these 29 RTPR 98% OTUs and their depth-resolved distribution patterns within

the virioplankton fraction, the bacterioplankton fraction, and the combined virio- and bacterioplankton

fraction were tested by plotting the cophenetic distance versus Aithison distance for all combinations of

29 OTUs. Combinations rather than permutations of OTUs were used, as the distances used are

symmetric (e.g., d(A, B) = d(B, A), where d is the distance function, and A and B are OTUs). Self-

distances (e.g., d(A, A), d(B, B)) were not included. Linear models and significance tests were performed

with the lm function in R.

Data transformation

Compositional data analysis (CoDA) methods were applied to all analyses of community diversity and

population interactions. Zero values in the feature tables [virioplankton RTPR 98% OTUs

(**Supplementary Table S3**); RTPR phylogenetic clade (**Supplementary Table S4**); bacterioplankton 16S

ASVs (**Supplementary Table S5**); bacterioplankton 16S ASVs retained in the 29 samples with RTPR

amplicons (**Supplementary Table S6**); and common RTPR 98% OTUs found in both the bacterio- and

virioplankton fractions (**Supplementary Tables S7 & S8**)] were replaced by imputed values using the

count zero multiplicative method from the R package zCompositions (version 1.3.3.1) [62] in R (version

3.5.3) [63]. The centered log-ratio (clr) transform was applied to the zero-replaced data set to produce clr

abundance using the R package CoDaSeq (version 0.99.4) [64]. Within the August 5th subset (sampling

date with RTPR amplicons in both fractions at all depths) of common RTPR 98% OTUs (**Supplementary**

Table S9), zero count values were replaced using *cmultRepl* with the method count zero multiplicative

(CZM) in the zCompositions package (version 1.3.4) of R (version 4.0.2). Subsequently, the clr

transformation was performed using base 2 in the R package CoDaSeq.

RTPR 98% OTUs within global RNR context

RTPR 98% OTUs from both virio- and bacterioplankton fractions, RTPRs identified through stringent

homology search of the *Tara* Ocean dataset [75], and RNRdb [76] databases were aligned using Geneious

10.0.9 MAFFT alignment FFT-NS-2 and trimmed to the region of interest (H346 to S643 in *Lactobacillus*

leichmannii monomeric class 2 adenosylcobalamin-dependent ribonucleotide-triphosphate reductase). A

phylogenetic tree of trimmed sequences was built using FastTree and visualized with Iroki.

RESULTS

Water Column Environment

Temperature of surface waters at station SCFP increased from May 12th (16.4 °C) to August 5th

(22.0 °C) (**Fig. 2A**), agreeing with broader sea surface temperature (SST) data derived from satellite

imagery (**Supplementary Fig. S4A**). Surface salinity (using the Practical Salinity Scale) increased from

May 12th (35.9) to July 11th (36.1), decreased by 0.1 on July 27th, and increased again to 36.1 on

August 5th (**Fig. 2B**). Surface waters were well mixed down to 50 m as indicated by temperature and

salinity contours prior to June (**Fig. 2**). Water column stratification intensified throughout the sampling

period, accompanying the mixed layer depth rising from 43 m (May 12th) to 5 m (August 5th). Salinity

showed distinct sub-surface increases during the study period, indicating substantial water movement in

the sampling region. Salinity changes in the sub-surface were centered at the 75 m depth, from June 8th

to July 27th, especially on July 11th. The observations suggested the presence of an anticyclonic

(downwelling) vortex at the sampling location around July 11th. Uplifts of isotherms and isohalines found

in temperature and salinity profiles between sampling dates suggested the presence of cyclonic

(upwelling) vortices, on June 22nd (more pronounced) and on July 27th (less pronounced) extending to

August 5th. Satellite-derived data in 2016 (**Supplementary Fig. S4A**) showed one phytoplankton bloom,

based on chlorophyll a and particulate organic carbon (POC), that occurred between February and April,

with its maximum in March.

The sampling period started after the seasonal spring phytoplankton bloom and spanned the

environmental and biological conditions observed from late spring to late summer. Observed changes in

phytoplankton abundance corresponded with seasonal environmental changes during the sampling

period (**Fig. 2A** and **Supplementary Fig. S4B**). High phytoplankton abundance (19–61 x 10^3 cells/mL)

was observed throughout the water column from 0 to 75 m depth on May 12th, indicating the end of the

annual spring bloom. Subsequently, phytoplankton abundance fell below 2 x 10^3 cells/mL with the

exception of a sub-surface peak of ~55 x 10^3 cells/mL at 50 m on June 22nd, coinciding with the

observed cyclonic upwelling event (**Supplementary Table S1**). The lowest phytoplankton abundance

was observed on July 11th, coinciding with the anticyclonic downwelling event. Another increase in

phytoplankton abundance coincided with the second, albeit less pronounced, cyclonic upwelling event

during the subsequent two sampling dates, July 27th and August 5th.

Changing environmental conditions and virioplankton and bacterioplankton community dynamics

For unknown reasons, only 29 of 35 virioplankton samples yielded RTPR amplicons (**Supplementary Tables S3 & S10**). From a total of 30,473 virioplankton RTPR amino acid sequences, 3,697 RTPR 98% OTUs were identified (**Supplementary Fig. S1 and S2**). The RTPR 98% OTUs clustered into 17 phylogenetic clades as manually determined by phylogenetic tree topology (**Fig. 3**). The three clades topologically located between clades 12 and 13 (gray color branches in **Fig. 3**) were excluded from downstream analyses due to their low abundance (7 OTUs, 12 amino acid sequences) and distant phylogenetic relations to other clades, leaving 3,690 98% OTUs (30,461 amino acid sequences, **Supplementary Table S3**) in 17 clades (**Supplementary Table S4**). Phylogenetic clade 5 was the most abundant based on both amino acid sequence counts (3,860) and clr abundance (1.07). The 17 RTPR phylogenetic clades were used in subsequent compositional data analyses, including community diversity and population interactions analyses, as clades of related OTUs provided clearer ecological signals than the larger collection of more highly resolved RTPR 98% OTUs. However, as shown in supplemental analyses, RTPR 98% OTU community diversity within and between samples was also determined for comparison with the phylogenetic approach.

The V3–V4 region of the 16S rRNA gene was PCR-amplified from 35 bacterioplankton fraction samples and sequenced (**Supplementary Tables S2 & S10**). After sequence quality filtering, 1,779 amplicon sequence variants (ASVs) were identified from a total of 918,792 sequences across all samples (**Supplementary Table S5**). Complete-linkage hierarchical clustering based on the Aitchison distance of clr-transformed sequence abundance between the first (May 12th 5 m) water sample and every other sample was performed for each of three datasets (virioplankton RTPR 98% OTUs, RTPR phylogenetic clades, or bacterioplankton 16S ASVs). Mantel tests assessed each Aitchison distance matrix with distance matrices of all collected environmental metadata (**Supplementary Table S11**) to identify environmental factors correlated with and potentially driving the clustering of virio- or bacterioplankton communities. Sample depth, collection day (displayed as Julian day), temperature, and water density were significantly (Spearman and Pearson *p*-value <0.02) correlated with virioplankton community composition (RTPR phylogenetic clades and RTPR 98% OTUs) and bacterioplankton community composition. For the significant environmental variables, correlation coefficients with virioplankton populations were generally higher when using the RTPR phylogenetic clades.

Hierarchical clustering of virioplankton communities based on RTPR phylogenetic clades formed four large groups (vA, vB, vC, and vD) (**Fig. 4A**) based on their distance from the May 12th 5 m sample. Community groups vA/vB and groups vC/vD were organized into equally distant supergroups. Within each group, communities had similar profiles for depth/density and for temperature/day. Groups vA/vB contained communities generally occurring at deeper depths with lower water temperatures, whereas groups vC/vD contained communities generally collected at shallower depths with higher water temperatures. However, for all groups, there were outliers likely reflecting some of the low rank

correlation coefficients (0.19–0.51) between community structure and environmental metadata
(**Supplementary Table S11**). Bacterioplankton communities formed three groups (bA, bB, and bC)
(**Fig. 4B**) based on their distance to the May 12th 5 m sample. Communities in groups bA and bB were
observed at lower water temperatures, whereas group bC communities were found in higher water
temperatures at shallow depths (<25 m). The division between groups bA and bB was more subtle and
may have been forced by depth as group bB contained communities in only 50 and 75 m samples.
The effects of sampling date (**Fig. 5A**) and depth (**Supplementary Fig. S5A**) on viral and bacterial
ENS were significant (betta <0.05; data not shown); therefore, these environmental parameters were
selected for analysis of their interactions with alpha and beta diversity of the virio- and bacterioplankton
communities. Over the interseasonal sampling period, virioplankton alpha diversity according to RTPR
phylogenetic clades (**Fig. 5A)** fluctuated slightly reaching a maximum around the July 11th anticyclonic
downwelling event (**Fig. 2)**. This peak was shouldered by minimum diversity estimates on June 8th and
July 27th corresponding with cyclonic upwelling events. Virioplankton alpha diversity fluctuated slightly
with depth (**Supplementary Fig. S5A)**. Maximal and minimal diversity estimates occurred at 25 and
75 m, respectively. ENS estimates of alpha diversity based on 98% OTUs (**Supplementary Fig. S6A
and B**) were three to six-fold higher than ENS estimates for virioplankton based on RTPR phylogenetic
clades (**Fig. 5A and Supplementary Fig. S5A**) as there were thousands of 98% OTUs versus only 17
RTPR phylogenetic clades. The general trends of ENS estimates based on 98% OTUs and RTPR
phylogenetic clades were consistent (Spearman's rank correlation: ρ = 0.82, *p* value = 0.034; Pearson's
product-moment correlation: *r* =0.78, *p* value = 0.037). Increases in RTPR 98% OTU ENS
(**Supplementary Fig. S6A**) were observed on June 22nd and July 11th, corresponding with the observed
cyclonic upwelling event. The maximum ENS was observed on September 8th; unfortunately its
association with an upwelling event is unknown as equipment failure prevented temperature
measurements on this date. The maximum and minimum virioplankton RTPR 98% OTU ENS in samples
by depth were 5 and 50 m, respectively (**Supplementary Fig. S6B**). Bacterioplankton alpha diversity
steadily declined from its May 12th peak to a minimum value on July 11th followed by a slight increase
over the remaining sampling dates (**Fig. 5A**). In contrast to virioplankton, bacterioplankton alpha diversity
did change dramatically with depth demonstrating steady increases from minimal diversity estimates in
surface waters to the maximal diversity estimates observed at 75 m (**Supplementary Fig. S5A**).
Relationships between virioplankton RTPR phylogenetic clade communities were examined using
principal components analysis (**Fig. 5B**). Sixty-five percent of the variability between communities was
explained by the first two principal components (PC1 48.9%, PC2 16.0%), and sampling date (**Fig. 5B)**
seemed to explain this variability more than sample depth (**Supplementary Fig. S5B**). Using the 98%
OTU populations, ~35% of the variability between communities was explained by the first two principal
components (PC1 22.8%, PC2 12.1%), and was associated with both sampling date and depth
(**Supplementary Fig. S6C and D**). The variability in virioplankton communities explained by the RTPR
98% OTUs was approximately half of that explained by phylogenetic clades. This trend was attributed to

the higher number and granularity of RTPR 98% OTU populations in describing virioplankton community

structure. For either means of defining virioplankton communities, PCA analyses showed that the May

12th virioplankton communities grouped closer to the origins of both the PC1 and PC2 axes, apart from

most other sampling dates, and the August 5th samples tended to place toward the ends of PC1 and PC2

axes. Virioplankton communities as defined by RTPR phylogenetic clades demonstrated significant

differences based on date and depth (PERMANOVA *p* value <0.05; **Supplementary Table S12A & D)**,

largely agreeing with observations of 98% OTUs **(Supplementary Table S12B & E**). May 12th

virioplankton communities differed from most other timepoints. June 8th communities differed from July

27th, and August 5th communities differed from those on June 8th, June 22nd, and July 11th. Depth-

based virioplankton community structure (**Supplementary Fig. S5B)** did not group as strongly on the

PCA plots as date-based communities (**Fig. 5B**). Nevertheless, surface communities (0 m) were different

from those at 50 and 75 m, and 5 m communities differed from 75 m communities.

Relationships between bacterioplankton 16S ASV communities were examined using principal

components analysis as well (**Fig. 5C)**. Approximately 38% of the variability between communities was

explained by the first two principal components (PC1 25.2%, PC2 12.4%), and was associated with both

sampling date (**Fig. 5C**) and depth (**Supplementary Fig. S5C**). As seen with viral communities defined

by either RTPR phylogenetic clades or 98% OTUs, the May 12th bacterioplankton samples grouped

toward the origin of both PC1 and PC2, apart from most other sampling dates. The August 5th

bacterioplankton communities showed a greater spread across PC1 than the virioplankton communities

for this sampling date. These observations were statistically confirmed. Bacterioplankton communities

on May 12th and June 8th were different from those observed on all other sampling dates, except the pair

of samples from June 8th and June 22nd (PERMANOVA *p* value and/or *q* value <0.05; **Supplementary**

Table S12C). Bacterial 16S ASV communities were also separated by depth, with 50 m and 75 m

communities occurring along the left side of PC1 (**Supplementary Fig. S5C**). Surface bacterioplankton

communities at 0 and 5 m significantly diverged from the deep communities at 50 and 75 m; 25 m

communities diverged from the 75 m communities; and 50 m communities diverged from the 75 m

communities (PERMANOVA *p* value and/or *q* value <0.05; **Supplementary Table S12F**).

Associations between virioplankton RTPR phylogenetic clades and bacterioplankton 16S ASVs

The proportionality test was used for comparing changes in the clr abundance of 17 virioplankton

RTPR populations (as defined by phylogenetic clades on **Fig. 3**) and the subset of 221 bacterioplankton

16S ASV populations in 27 samples with both virioplankton RTPR and bacterioplankton 16S amplicons

(**Supplementary Table S13**). Virio- and bacterioplankton population pairs with ρ values ≥0.48 had

positive associations as the false discovery rates were ≤0.05 (**Supplementary Fig. S7**). Nine of the 17

virioplankton RTPR phylogenetic clades had positive associations (ρ ≥.48) with 39 of 221

bacterioplankton ASV populations (**Fig. 6**). Associating virioplankton clades were phylogenetically

diverse and were observed at both high (clade 2 and 5; clr 1.03 and 1.07, respectively) and low

abundance (clade 3 and 11; clr -0.90 and -2.57, respectively) (**Fig. 3**). Virio- and bacterioplankton

abundance did not appear to drive these associations as low-abundance RTPR virioplankton associated
with high-abundance bacterioplankton and vice versa (**Supplementary Fig. S8)**. The phylogenetic
diversity of positively-associating bacterioplankton populations was broad, encompassing seven bacterial
phyla observed among the included 16S ASV populations. The bacterioplankton 16S ASV populations
ranged in clr abundance from the most abundant taxon 3.65 (Prochlorococcus_MIT9313) to one of the
least abundant taxon -1.99 (SAR116_clade).
Virioplankton RTPR phylogenetic clades 1 and 3 were found to be associated with the most abundant
bacterioplankton 16S ASV, Prochlorococcus_MIT9313, and several 16S ASVs from Proteobacteria, such
as SAR86, SAR11, and the Aegan169 marine group, well known to be abundant within oceanic
ecosystems. Virioplankton clade 2 showed associations with multiple bacterioplankton ASVs from
Proteobacteria and Bacteroidetes. Interestingly, RTPR phylogenetic clade 2 was not associated with any
of the same 16S ASVs as clades 1 and 3, despite its close phylogenetic relationship with these clades
(**Fig. 3**). Phylogenetic clades 3 and 4 were both associated with SAR11_clade_IV_1, and clade 4 was
also associated with SAR11_clade_IV_2. Virioplankton clades 11 and 15 were each only associated with
a single bacterioplankton 16S ASV, uncultured_Thiotrichaceae and SAR86_clade_4, respectively.
Virioplankton clade 5, the most abundant RTPR phylogenetic clade (clr 1.07, **Fig. 3**), showed the most
positive associations across a wide phylogenetic breadth of bacterioplankton populations, encompassing
four of the nine bacterial phyla and 15 of the 39 total positively-associated 16S ASVs, including
SAR11_clade_II.
Fourteen out of 136 non-self comparisons between virioplankton RTPR populations showed a
positive association (**Supplementary Fig. S9**, **Supplementary Table S13**). These associations between
virioplankton clades did not follow a pattern according to phylogenetic distance as only three of the 14
associations were between neighboring RTPR clades (clades 2 and 3; clades 5 and 6; and clades 8 and
9).
Of the 221 bacterioplankton 16S ASV populations, 36 showed positive, non-self associations with
other ASV populations (**Supplementary Fig. S10**, **Supplementary Table S13**). Overall, there were 390
positive non-self associations among the 36 ASVs, ~5% of all the non-self comparisons (1260).
Positively associating bacterioplankton 16S ASV populations were clearly divided by their association
with either SAR11_clade_II or Prochlorococcus_MIT9313 (red boxes in **Supplementary Fig. S10**). No
16S ASVs associated with both SAR11_clade_II and Prochlorococcus_MIT9313. The SAR_clade_II
population was associated with only other low abundance (clr <0) ASVs, whereas
Prochlorococcus_MIT9313 was associated with mostly high abundance (clr >0) ASVs. Beyond the
delineation of SAR11_clade_II from Prochlorococcus_MIT9313, positively associating bacterioplankton
ASV populations could be divided into three larger groups (**Supplementary Fig. S10** x-axis). Nearly all of
the ASV populations within group 1 all showed positive associations with SAR11_clade_II and nine other
taxa (clr values ranging from -1.38 to -0.35) across five phyla (Marinimicrobia:
Marinimicrobia_SAR406_clades 1 (clr -1.22) & 2 (clr -1.13); Actinobacteria: Candidatus_Actinomarina (clr

-1.38); Planctomycetes: uncultured_Rubinisphaeraceae (clr -1.11); Verrucomicrobia: Roseibacillus (clr -

0.35); Proteobacteria: UBA10353_marine_group (clr -1.12), Ga0077536 (clr -1.28), OM60_NOR5_clade

(clr -1.03), and Rhodobacteraceae (clr -1.02)). ASV populations within group 2 were defined by their

consistent association with the most dominant ASV taxa Cyanobacteria: Prochlorococcus_MIT9313 (clr

3.65). In addition to this singular representative of the Cyanobacteria, group 2 consisted of 13

Proteobacteria (clr values ranging from -0.45 to 2.15) and three *Bacteriodetes* (clr values ranging from -

1.45 to 0.19). Several of the ASV taxa in group 2, such as Proteobacteria:

uncultured_Ectothiorhodospiraceae_2 (clr 1.72), Proteobacteria: SAR11_clade_IV_1 (clr 2.15) & 2 (clr

1.95), and Proteobacteria: SAR86_clade 3 (clr 1.51) were the most abundant bacterioplankton

populations observed in the study. ASV populations within group 3, which consisted of only three

Proteobacteria taxa, showed no consistent associations within the group and few associations overall.

Depth- and time-resolved viral–host associations

Spatiotemporal patterns in bacterioplankton associations with clade 5, the most abundant RTPR

virioplankton population, were explored by plotting changes in clr abundance at each sampling depth over

time (**Supplementary Fig. S11**). Clade 5 RTPR viral populations showed their highest abundances at

sampling depths of 25 m and deeper, with consistently high abundances occurring at 75 m. Deeper

samples also contained a greater number and diversity of bacterioplankton ASV taxa showing

associations with clade 5 RTPR virioplankton. Among the sampling dates, July 11th was noteworthy as

several of the ASV taxa associated with clade 5 RTPR virioplankton showed dramatic reductions in clr

abundance corresponding with the anticyclonic downwelling event. Correspondingly, the clr abundance

of clade 5 virioplankton held steady or declined on the two subsequent sampling dates (July 27th and

August 5th). A similar spatiotemporal analysis of associations between virio- and bacterioplankton

populations was conducted for abundant bacterioplankton populations, Prochlorococcus_MIT9313 and

three SAR11 clades, and four associating RTPR virioplankton populations (**Supplementary Fig. S12**).

Clades 1 and 3 RTPR virioplankton populations showed associations with Prochlorococcus_MIT9313

(**Fig. 6**). Within a sampled depth across time, the clr abundance of clade 1 and 3 viruses was relatively

steady, however, clr abundance of these two populations decreased in the 75 m samples as did the

abundance of their associated Prochlorococcus_MIT9313 population. RTPR clade 3 was also associated

with SAR11_clade_IV_1 which, like Prochlorococcus_MIT9313, was most abundant in surface waters.

RTPR clade 4 showed positive associations with SAR11_clade_IV_1 and _2 **(Fig. 6),** which were the

second and third most abundant bacterioplankton taxa showing associations with RTPR virioplankton.

While these SAR11 populations were most abundant at the surface (clr values of around 3.75), they were

also abundant throughout the water column and changed only modestly with time. The behavior of RTPR

clade 4 showed a similar pattern as it was most abundant in surface waters (clr values ≥ 1) and then

lower abundance at 50 and 75 m (clr values ~ 0). RTPR clade 5 associated with SAR11_clade_II which

was the rarest of the SAR11 taxa showing associations with an RTPR virioplankton population.

SAR11_clade_II populations increased with depth, showing their highest abundances (clr values ~ 0) at

75 m. Similarly, RTPR clade 5 was less abundant at shallow depths (clr values < 1) and then increased

to its highest abundances at 75 m (clr values > 1).

Free and actively replicating viral populations: RTPR 98% OTUs occurring in virio- and bacterioplankton fractions

Twenty-nine virioplankton and 31 bacterioplankton samples provided RTPR amplicons with an

overlap of 27 samples providing amplicons from both. These 27 paired samples provided 47,997 RTPR

quality-filtered amino acid sequences, which were clustered *de novo* into 5,411 RTPR common 98%

OTUs (**Supplementary Table S7**). After removing 98% RTPR OTUs with less than ten sequences, 21

OTUs were observed only in the virioplankton fraction, 40 only in the bacterioplankton fraction, and 193 in

both fractions (**Fig. 7** and **Supplementary Table S8**). An encompassing phylogenetic analysis of these

RTPR 98% OTUs, sequences from the RNR database, and viromes from the *Tara* Oceans expedition

showed that the RTPR 98% OTUs from the virio- and bacterioplankton fractions occurred within a section

of the tree containing few sequences from known bacteria or viruses (RNRdb) but several sequences

from *Tara* Oceans viromes (**Supplementary Fig. S13**). The most abundant OTUs were primarily found in

the shared fraction (**Fig. 7A**). Less abundant OTUs were largely found in either virio- or bacterioplankton

fractions only. Examining the relative contribution of virio- or bacterioplankton amplicon sequences to

RTPR 98% OTUs required assignment of an arbitrary clr of 5 and -5 to those OTUs found only within the

virio- or bacterioplankton fractions, respectively. The difference in clr abundance within each RTPR 98%

OTU (clr(virioplankton)-clr(bacterioplankton), **Fig. 7B**) found that within most of the shared OTUs,

virioplankton contributed a greater number of amplicons (66% of all shared OTUs (128 of 193), red

colored OTUs above the dashed line in **Fig. 7B**).

All samples (fractions and depths) from the August 5th sampling date provided RTPR amplicon

sequences, enabling coordinate examination of virioplankton populations observed both as free viruses

(i.e., within the virioplankton fraction) and viruses replicating within cells (i.e., within the bacterioplankton

fraction). Twenty-nine RTPR 98% OTU populations were selected, each containing greater than 50 total

sequences with two or fewer zero-replaced samples (**Supplementary Table S9**). These heuristics

prevented the inclusion of rare and inconsistently observed virioplankton populations. Phylogenetic

analysis of these 29 RTPR 98% OTUs alongside observations of clr abundance of each OTU within the

virio- and bacterioplankton fractions addressed the hypothesis that the evolutionary distance between

virioplankton populations reflects similarity in their ecological behavior (**Fig. 8**). Statistical testing

comparing the cophenetic distance of the OTUs with the Aitchison distance of the OTUs according to

their clr abundances across the August 5th samples demonstrated a significant (p = 1.6 e^{-5}), weak (R^2 =

0.045), positive correlation (correlation r = 0.212) between evolutionary distance and clr abundance in the

bacterioplankton fraction (**Supplementary Fig. S14**). There was no significant relationship between

cophenetic distance and clr abundance in the virioplankton fraction or when combining the virio- and

bacterioplankton fractions (data not shown). Thus, more closely related RTPR virioplankton populations

demonstrated similar ecological behavior in terms of abundance as viruses replicating within cells (i.e.,

the bacterioplankton fraction), but not as free viruses.

The 29 August 5th RTPR 98% OTUs occurred across nearly all of the major RTPR clades identified

in **Fig. 3**, with the exception of clades 10 and 11 which were among the rarest clades observed in the

study (clr values -0.95 and -2.57, respectively). Clade 7 was the most represented among the August 5th

OTUs. These OTUs demonstrated a distinctive evolutionary split with a group of closely related OTUs (9,

15, 7, 16 and 26) clustering away from OTU 25 (**Fig. 8**). The clr abundance heat maps of the closely

related group were similar with the majority of sequences occurring in the virioplankton fraction above 25

meters. In contrast, OTU 25 sequences were more abundant below 25 m demonstrating high abundance

in the bacterioplankton fraction at 50 m. By and large, closely related August 5th OTUs showed similar

clr abundance patterns supporting the finding of a significant relationship between evolutionary distance

of RTPR virioplankton populations and Aithison distance.

DISCUSSION

Ribonucleotide triphosphate reductase: a hallmark gene within viruses

As the only enzyme capable of reducing ribonucleotides to deoxyribonucleotides, RNR effectively

controls the rate of DNA synthesis [33]. Thus, it makes intuitive sense that RNR genes would be common

within the genomes of lytic viruses as speed and efficiency in genome replication is under positive

selection pressure for viruses exhibiting a virulent lifecycle. Indeed, RNR genes are among the most

commonly observed genes within dsDNA viral genomes and viromes, lending support to this hypothesis

[34,77–79]. RNRs are ancient proteins as enzymatic ribonucleotide reduction was a critical step from the RNA

world to the emergent dominance of DNA as the principle information-carrying molecule for life on Earth

[80–82]. Thus, it is possible that RNRs existed within viral genomes not long after the emergence of DNA.

Phylogenetic analysis of known and environmental RTPRs (**Supplementary Fig. S13**) indicated that viral

RTPR proteins diverged from cellular RTPRs and are as deeply branching, reflecting both the unique

selective pressure on viral RTPRs and their long evolutionary history. However, the tree also shows the

emergence of cellular RTPRs from clades dominated by viral sequences and vice versa. These

instances may represent viral to cellular (and cellular to viral) gene transfer events. Throughout the long

evolutionary history of RNRs there is evidence of gene transfer between cellular kingdoms, most notably

the transfer of Class I RNRs from bacteria to archaea and ultimately eukaryotes [31] where Class I RNRs

predominate (**Supplementary Fig. S15A**). In the case of nucleotide metabolism proteins, such as RNRs,

it is possible that greater levels of biochemical innovation occur within lytic viruses than within cells, as

viruses typically carry one allele of a particular nucleotide metabolism gene which is under intense

selective pressure for fast and efficient DNA replication. Certainly, at the level of RNR class, it is clear

that selective processes may influence the distribution of RNR classes across the tree of life, where

viruses demonstrate a distinctly high frequency of Class II RTPRs in comparison with cellular kingdoms.

Among known viruses, Class II RTPRs demonstrate a distinctive skew in taxonomic distribution favoring

tailed phages having a myoviridae or siphoviridae morphology (**Supplementary Figs. S15A & B**).

Interestingly, a breadth of microbial diversity was associated with several of the RTPR viral

phylogenetic clades (**Fig. 6**). There are a few possible explanations for this. One possibility is that

physiochemical factors were stronger drivers of microbial dynamics than viral-induced mortality, resulting

in positive associations between RTPR viral clades and a sub-community of similarly adapted organisms,

both host and non-host alike. However, it is also possible that the diverse taxa positively associated with

several of the RTPR viral clades are a reflection of the hosts for viruses within these clades. Abundant

marine phages, like 37-F6, have been predicted to infect hosts spanning different phyla [83], while direct

linkages between viral RNR genes and 16S rRNA genes from diverse bacterial taxa have been observed

[84]. In the latter case, viruses associated with diverse taxa were less efficient at infecting these hosts than

viruses with narrow host ranges [84], but even infrequent cross-taxa infections could provide ample

opportunities for horizontal gene transfer (HGT) of RNR genes between viral populations, resulting in

similar RNR sequences across a diverse group of viruses infecting different hosts. Furthermore,

horizontal gene transfer of RNRs across all three domains of life [31], as well as between phage

populations [34], is well-established, suggesting that HGT has been critical to shaping viral ecology and

cobalamin cycling in marine environments. Answering questions surrounding cellular–viral gene

exchanges and their contribution towards the emergence of biochemical innovation will improve

understanding of the evolutionary mechanisms behind adaptation in microbial communities.

Biochemistry constrains RTPR virioplankton population ecology

RNR is an essential gene for cellular life as all cells contain dsDNA genomes. However, the

frequency of RNR types varies substantially among known cellular organisms and viruses

(**Supplementary Fig. S15A**). Meta-analysis of RNR protein sequences within the RNRdb [76] along with

sequences in the UniProt knowledgebase [85] showed the high frequency of RTPR (i.e., Class II

monomeric RNR) within known viruses (~22%) as compared to cellular kingdoms (2–3%). Within the

Caudovirales most RTPR genes occurred within siphophages (**Supplementary Fig. S15B**). In contrast,

Archaea and Bacteria favor Class II dimeric or Class III RNRs for anaerobic ribonucleotide reduction, a

biochemical strategy strikingly different from viruses. In the case of RTPR, this monomeric Class II RNR

requires a adensyolcobalamin (B_{12}) co-factor (like all Class II RNRs), can perform ribonucleotide

reduction under aerobic or anaerobic conditions, and, along with Class III RNRs, utilizes a ribonucleotide

triphosphate (NTP) substrate [86]. The B_{12} requirement and use of NTP substrates have intriguing

implications for the ecology of viruses carrying RTPR. Since cobamide nutrient (including B_{12}) synthesis

is complex, specialized, and energy intensive, few marine bacterial and archaeal phyla possess the entire

synthesis pathway [87,88]. Most bacteria and phytoplankton are auxotrophic for B_{12} and are thus dependent

on the few taxa capable of B_{12} production. RTPR virioplankton populations must infect a host that can

either synthesize B_{12} or assimilate and subsequently modify a B_{12} precursor containing the corrin ring.

Moreover, B_{12} has several chemical analogs depending on the combination of ligands at the α and β sites

of the corrin ring [89,90].

There is a growing recognition that selectivity in B_{12} analogs may drive interaction networks within

microbial communities as cobamide specificity is commonly observed in microorganisms [91]. For example,

pseudocobalamin, produced by cyanobacteria such as *Prochlorococcus* (the dominant bacterioplankton

taxa observed in this study), is poorly utilized by B_{12} auxotrophic microalgae [90,92]. Specificity in B_{12}

requirements among bacterioplankton taxa may explain the separation observed between

bacterioplankton populations associating with either SAR11_clade_II populations (**Supplementary Fig.**

S10, Group 1) or Prochlorococcus_MIT9313 (**Supplementary Fig. S10**, Group 2). Positive association

patterns between virioplankton RTPR phylotype populations and bacterioplankton 16S ASV populations

reflected the split seen between SAR11 and *Prochlorococcus* associations. RTPR virioplankton clades 5

and 6 associated with SAR11_clade_II and 14 other bacterioplankton populations (**Fig. 6**), three of which

(Magnetospiraceae, Rhodobacteraceae, Planktomarina) may be capable of B_{12} production [90]. Eleven of

these bacterioplankton populations are associated with SAR11_clade_II (**Supplementary Fig. S10**,

Group 1). Similarly, RTPR virioplankton clades 1 and 3 associated with Prochlorococcus_MIT9313, and

six other bacterioplankton populations that also demonstrated associations with *Prochlorococcus*

(**Supplementary Fig. S9**, Group 2). Three of these seven (AEGEAN_169_marine_group_1,

AEGEAN_169_marine_group_2, and Prochlorococcus_MIT9313) occur in an order, family, or genera

containing species capable of B_{12} production (**Fig. 6**).

Time- and depth-resolved observations of RTPR virioplankton clades 1, 3, and 5 (**Supplementary**

Figs. S11 and **S12**) indicate that changes in the abundance of these virioplankton populations followed

those of their associated bacterioplankton populations throughout the euphotic zone. It seems unlikely

that the parallels in association patterns observed between bacterioplankton populations

(**Supplementary Fig. S10**) and those observed between RTPR virioplankton and bacterioplankton

populations (**Fig. 6**) are happenstance. It is likely that these positive virio- to bacterioplankton association

patterns reflect the B_{12}-driven association patterns among their bacterioplankton hosts [91]. It is likely that a

few heterotrophic bacterial taxa and *Prochlorococcus* were the principal B_{12}-producing populations in the

euphotic surface waters of the Azores. Although Thaumarchaeota are also important oceanic B_{12}

producers, metagenomic biogeographic surveys indicate that these archaeal populations predominate at

high latitudes or in mesopelagic waters [88].

It has been hypothesized that virioplankton influence oceanic primary and secondary productivity by

controlling the flux of limiting trace nutrients, such as iron [93], through cell lysis. The common presence

and dynamic nature of RTPR virioplankton populations now places viral–host interactions as another

important component within the vitamin traffic pathways of this trace nutrient in the oligotrophic ocean [94].

While technically B_{12} is not a nutrient as it is not incorporated into cellular biomass, it displays oceanic

distribution patterns reminiscent of nutrients such as nitrogen. Within surface waters of the oligotrophic

oceanic gyres, B_{12} occurs in vanishingly low concentrations of less than 1.75 picomolar [90,95]. However,

deeper waters typically demonstrate higher B_{12} concentrations [96]. Depth-resolved dissolved B_{12}

concentrations occurring at our study site were likely highest below 25 m, similar to those recorded for in

April and August at the western-most station of a recent study of eastern north Atlantic central waters off

the Iberian peninsula [95]. While RTPR virioplankton populations did show differences in their depth and

time distribution (**Fig. 8, Supplementary Figs. S11** and **S12**), the abundance of these populations as a

fraction of overall virioplankton abundance could not be discerned using an amplicon approach.

Nevertheless, as RTPR viral populations depend on B_{12} for a steady supply of deoxyribonucleotides, it is

possible that these virioplankton regulate or enhance B_{12} uptake or production during infection.

Subsequently, lysis by RTPR-utilizing virioplankton could increase the flux of B_{12} pools disproportionately

over virioplankton populations that do not utilize B_{12}-dependent proteins. New quantitative approaches

for observations of viral–host interactions based on detecting single viral genes using polonies [97–99] could

provide data on the *in situ* frequency and production rate of RTPR virioplankton populations, as has been

recently demonstrated for pelagic cyanophage populations [100].

Unlike Class I and dimeric Class II RNRs that use ribonucleotide diphosphate (NDP) substrates,

RTPR uses ribonucleotide triphosphate (NTP) substrates. This difference in substrates may correlate

with significant differences in viral population ecology. Virioplankton populations utilizing a Class I or

dimeric Class II RNR would be dependent on NDP production from RNA digestion [101] as well as an NDP

kinase for producing dNTPs [102]. In contrast, RTPR virioplankton populations have direct and immediate

access to the cellular pool of NTPs (which is substantially larger than the NDP pool [103]) for DNA

synthesis. These biochemical features of RTPR would favor a rapid lytic cycle. It is possible that RTPR-

carrying virioplankton are highly lytic viruses specifically infecting actively growing bacterioplankton

populations containing large intercellular pools of rNTP substrates.

However, there are possible costs to an RTPR strategy in terms of DNA synthesis regulation.

Improved control of dNTP substrate levels for DNA synthesis and allosteric regulation of RNR enzyme

activity have been proposed as possible advantages of NDP versus NTP reduction [82] . Thus, RTPR-

carrying viruses may pay a cost for faster DNA synthesis with lower fidelity as imbalances or excesses in

dNTP levels from runaway RTPR activity can increase mutagenesis rates [102,104]. The fact that RTPRs are

common within the virioplankton [34,77,78] and are proportionally more represented within viruses than within

cellular genomes (**Supplementary Fig. S15**) indicates that the possible fitness costs of RTPR do not

outweigh the benefits of utilizing this gene for dNTP synthesis. Nevertheless, it is possible that the

frequency of RTPR-carrying viruses may depend on environmental context as the majority of known

viruses carry a Class I or Class II dimeric RNR (**Supplementary Fig. S15**).

RTPR virioplankton phylotypes demonstrated dynamic responses to ecosystem change

At the community and population scale, RTPR-carrying virioplankton demonstrated dynamic behavior

throughout the four month study, responding to seasonal and oceanographic changes influencing the

activity of co-occurring phyto- and bacterioplankton communities. In this sense, RTPR virioplankton

populations demonstrated community dynamic behavior reminiscent of that seen from studies utilizing

amplicons of structural genes such as T4 major capsid protein [28–30], portal protein [24,105], and terminase [25–

27]. A unique aspect of this study was that primer sets targeting viral RTPR genes were used for

amplifying RTPR from both the virioplankton (0.02–0.22 µm) and bacterioplankton (>0.22 µm) fractions.

While viral genes have been amplified and sequenced from bacterioplankton fractions in past studies [29],

to our knowledge none have examined coordinate occurrence of OTUs across extracellular (i.e.,

virioplankton fraction) and intracellular (i.e., bacterioplankton fraction) virioplankton populations. The

majority of RTPR OTUs observed in both fractions showed a greater abundance in the virioplankton

fraction (**Fig. 7**) indicating host populations undergoing active lysis. These OTUs had the highest clr

abundance of all RTPR virioplankton populations observed in the study. This observation makes sense

as each viral particle released contains an RTPR gene copy and dozens of viral particles are released for

each lysed cell. RTPR OTUs observed solely within either the virio- or bacterioplankton fraction were less

frequent (8% virioplankton only; 16% bacterioplankton only) and typically showed low clr abundance.

Virioplankton-only RTPR OTUs likely represented viral populations either no longer actively infecting

hosts or somehow displaced from their host populations. Bacterioplankton-only RTPR OTUs may have

represented lysogenic viruses existing as prophages, free viral genomic DNA within the >0.22 µm size

fraction, some lytic viral populations yet to lyse infected cells, or inadvertent amplification of a bacterial

RTPR gene. Distinguishing between these four possibilities would be difficult; nevertheless, it is clear

that the majority of observed RTPR OTUs (84%) occurred within the virioplankton.

Individual RTPR virioplankton populations were actively infecting, replicating, and subsequently

disappearing from the pelagic euphotic zone. Using an amplicon approach provided the advantage of

deeply sampling RTPR virioplankton; it was less clear how these data should be partitioned into

ecologically meaningful populations. Community and population ecology analyses conducted using either

a heuristic approach (binning amplicons into 98% amino acid identity OTUs) or an evolutionary approach

(defining major phylogenetic clades of RTPR amplicons) demonstrated the superiority of the evolutionary

approach. Community beta diversity analyses based on clr-transformed OTU or clade abundance

showed that the evolutionary approach (clades, **Fig. 5B** and **Supplementary Fig. S5B**) captured a

greater proportion of variability in virioplankton communities than the heuristic approach (98% OTUs

Supplementary Fig. S6C and D). The evolutionary approach lumped RTPR virioplankton populations

into 17 clades as opposed to thousands of 98% OTUs, likely providing a greater signal to noise ratio for

observing virioplankton ecological behavior. Indeed, virioplankton evolutionary relationships gleaned

from RTPR phylogeny demonstrated a significant connection with the ecological behavior of virioplankton

populations on the August 5th sampling date (**Fig. 8**). The RTPR genes carried within each clade

population likely occurred within a diversity of genomic backgrounds. Thus, it was surprising that this

single gene showed a statistically significant positive relationship between evolutionary distance (RTPR

phylogenetic tree) and patterns of clr abundance within the bacterioplankton fraction (Aitchison distance)

(Supplementary Fig. S14). The significant correlation of cophenetic distance (RTPR phylogeny) and

ecological behavior (Aitchison distance) of RTPR OTUs identified within the bacterioplankton and not

within free viruses is interesting as these populations represented those RTPR viral populations

undergoing active replication.

Examinations of connections between gene-based viral phylogeny and viral ecology have been

limited, in large part due to the widely acknowledged mosaicism within viral genomes [106]. Nevertheless,

when considering the influence of enzyme biochemistry or the presence/absence of particular auxiliary

metabolic genes [107], it is clear that single genes have a dramatic influence on a virus' ecology. Single-

gene amplicon approaches provide not only an evolutionary framework for understanding viral ecology,

but also have practical advantages over metagenomic approaches in terms of cost, depth and breadth of

sampling, and computational complexity [108]. Prior work examining genetic connections to phage host

range phenotypes found that while whole genome networks provided better predictive capability, single

gene groups could also show strong connections [109]. Interestingly, DNA replication and phage replication

were among those Rapid Annotation Subsystem gene groups [110] showing a stronger predictive ability for

host range. This might not be surprising when considering that phage genome and replication processes

may require intimate interaction with host anabolic systems. Examining possible connections between

single genes and whole genomes and other important phage phenotypes, especially those describing

infection dynamics, should ultimately improve the predictive ability of metagenome data for ecosystem

modeling of phage impacts on ecosystem processes.

AUTHOR CONTRIBUTIONS

GP designed the study, performed processing of water samples and DNA extraction following water

samples collection and part of the RTPR library preparation. LW conducted the majority of the RTPR

library preparation and all of the 16S rDNA library preparation, performed all of the bioinformatics and

biostatistical analyses following sequence processing, and wrote and edited the manuscript. RM, CL, AH,

JD, BF, AM, KB, SP, JN and EW contributed to the study design, protocol development, data analysis

and interpretation, and manuscript preparation. All authors read, edited, and approved the final

manuscript.

ACKNOWLEDGEMENTS

We would like to thank, by order of involvement, Clara M. Loureiro, Catharina Pieper, Ricardo Fernandes,

Christien Laber, Prasanna Joglekar, Joana Botelho, Ana Pavon, and Marilia Olio for the help collecting

and processing the samples. We would like to thank Dr. Daniel Nasko for input on library preparation and

statistical analysis. We would like to acknowledge the skipper Renato Bettencourt, and all laboratory and

facility management personnel from Departamento de Oceanografia e P, OKEANOS Institute of Marine

Research, and IMAR-Instituto do Mar from the University of the Azores, DMCS-RU and DBI-UD.

FUNDING

This research was supported by the Estagiar-L fellowship; EEA grant funded BIOMETOREzores

(PT02_Aviso2_0001) project; the mobility program FLAD-UA Crossing the Atlantic; Chinese Scholarship

Council (CSC), and National Science Foundation grant number 1736030. Support from the University of

Delaware Center for Bioinformatics and Computational Biology (CBCB) Core Facility, the University of

Delaware Sequencing and Genotyping Center, and use of the BIOMIX compute cluster was made

possible through funding from Delaware INBRE (NIGMS P20GM103446), the State of Delaware, and the

Delaware Biotechnology Institute.

Conflict of interest statement. None declared.

FIGURES

Figure 1. Overview of experimental methods and analysis pipeline. The 0.22 µm filter captured

bacterioplankton cells. The 0.02 μm filter captured virioplankton particles. Hexagons represent molecular

genetic experimental methods. QC stands for a collection of quality control steps specific to the type of

next-generation sequencing technology used in sequencing amplicon libraries. Circles represent

conventional data analysis methods. Squares represent compositional data analysis (CoDA) methods.

Figure 2. Vertical profile contours of **(A)** temperature and phytoplankton cell numbers, and **(B)** salinity at

station South of Channel Faial-Pico islands (SCFP) from May 12th to August 5th, 2016. Data was not

collected on September 8th due to equipment malfunction. Vertical gray lines indicate the sampling date

and the maximum depth of data collection. Temperature and salinity data between sampling dates was

interpolated using the weighted average grid method. Circle sizes indicate phytoplankton abundances at

each sampled depth (surface, 25, 50 and 75 m). Surface values were the average abundance at the 0

and 5 m depths.

Figure 3. Approximate maximum likelihood tree of virioplankton RTPR 98% OTU representative

sequences. Shared branch, node, and clade number colors indicate RTPR phylogenetic clade manual

assignment. Seven OTUs (between clades 12 and 13) with low abundance and distant relationships with

other phylogenetic clades are represented by gray branches and were excluded from downstream

analysis. Node sizes indicate 98% OTU counts. Each phylogenetic clade's total amino acid sequence

counts and clr abundance are indicated in rounded and squared parentheses, respectively. Scale bar

indicates the average number of amino acid substitutions per site.

Figure 4. Phylogenetic clustering of virio- and bacterioplankton communities (dendrograms) based on **(A)** virioplankton RTPR phylogenetic clades and **(B)** bacterioplankton 16S ASVs, alongside environmental variables (columns of horizontal bars) demonstrating significant correlations (Spearman and Pearson tests, $p < .05$). Dendrograms are based on complete-linkage hierarchical clustering of each sample's Aitchison distance to the sample collected May 12th at 5 m (May12_05m) based on centered-log ratio (clr) abundance. Scale bar indicates Aitchison distance between samples. Major groups of virio- or bacterioplankton communities are labeled to the left of each dendrogram. Environmental variables are represented in columns, with horizontal bars indicating the measurement for each sample both by color gradient and by bar length compared to the numerical scale along each column's horizontal axis [Depth (blue), Day (green), Temperature (red), and Density (gray)]. Sample labels with asterisks (*) indicate samples where temperature and density were estimated using the Marine Copernicus Global Ocean Physics model.

Figure 5. Sampling date influenced virioplankton (RTPR phylogenetic clade) and bacterioplankton (16S

ASV) community alpha and beta diversity. **(A)** Effective number of species (ENS, based on Shannon

indices) of virioplankton RTPR phylogenetic clades and bacterioplankton 16S communities by sampling

date. Error bars represent two standard deviations of the estimates. Principal component analysis (PCA)

plots of beta diversity based on centered-log ratio (clr) transformed abundance of **(B)** virioplankton RPTR

phylogenetic clade communities or **(C)** bacterioplankton 16S ASV communities. Samples (circles) are

colored according to sampling date.

Figure 6. Matrix (ρ proportionality) of significantly associated ($\rho \geq .48$) virioplankton RTPR phylogenetic

clades (x-axis) and bacterioplankton 16S ASVs (y-axis) (**Supplementary Table S13**). Virioplankton

RTPR clades are colored and arranged by their topological order on the 98% OTU phylogenetic tree (**Fig.**

3). Bacterioplankton 16S ASVs are arranged according to phylogeny. Bacterioplankton 16S ASV phylum

(colored block), raw sequence count, clr abundance, and taxonomic assignment are shown on the right.

Asterisk indicates those ASVs within orders, families, or genera containing species capable of B_{12}

synthesis according to a pangenomic survey by Heal *et al*. 2017 [90].

Figure 7. RTPR 98% OTUs (n=254) identified from amino acid sequences present in 27 paired samples with amplicons from the virio- and bacterioplankton fractions. RTPR 98% OTUs (filled circles) were observed in either only the bacterioplankton fraction (blue circle, n=40), only the virioplankton fraction (red circle, n=21), or both (Shared, n=193). The dashed line in the overlap region divides the shared OTUs more abundant in the virioplankton fraction (above the line, n=128) from those more abundant in the bacterioplankton fraction (below the line, n=65). **(A)** RTPR 98% OTUs are colored by their total centered

log ratio (clr) abundance in both fractions (clr[virioplankton]+clr[bacterioplankton]). **(B)** RTPR 98% OTUs

are colored by the difference between clr abundance in the virioplankton and bacterioplankton fractions

(clr[virioplankton] - clr[bacterioplankton]). Red colors indicate greater abundance of that OTU in the

virioplankton fraction (values > 0, where 5 is an arbitrary maximum representing OTU presence in the

virioplankton only). Blue colors indicate greater abundance of that OTU in the bacterioplankton fraction

(values < 0, where -5 is an arbitrary minimum representing OTU presence in the bacterioplankton only).

Figure 8. Phylogram of 29 RTPR 98% OTUs occurring within both the bacterioplankton (B) and

virioplankton (V) fractions on the August 5th sampling date, with heat maps at nodes representing OTU

centered-log ratio (clr) abundance in each sample (fraction and depth). Node labels indicate OTU

number. Branches are colored and labeled according to membership of the OTUs in the RTPR clades

(C1–C9; C12–C17) identified in the inset tree (**Fig. 3**). Scale bar indicates the average number of amino

acid substitutions per site.

REFERENCES

1.   McClain, C. R., Signorini, S. R. & Christian, J. R. Subtropical gyre variability observed by ocean-color

satellites. *Deep Sea Res. Part 2 Top. Stud. Oceanogr.* **51**, 281–301 (2004).

2.   Huang, R. X. & Russell, S. Ventilation of the Subtropical North Pacific. *J. Phys. Oceanogr.* **24**, 2589–
2605 (1994).

3.   Sherr, E. & Sherr, B. Understanding Roles of Microbes in Marine Pelagic Food Webs: A Brief History.
in *Microbial Ecology of the Oceans* (ed. Kirchman, D. L.) 27–44 (John Wiley & Sons, Inc., 2008).
doi:10.1002/9780470281840.ch2.

4.   Goldman, J. C. Conceptual Role for Microaggregates in Pelagic Waters. *Bull. Mar. Sci.* **35**, 462–476
(1984).

5.   Díez, B., Pedrós-Alió, C. & Massana, R. Study of Genetic Diversity of Eukaryotic Picoplankton in
Different Oceanic Regions by Small-Subunit rRNA Gene Cloning and Sequencing. *Appl. Environ.*
Microbiol. **67**, 2932 (2001).

6.   Wilhelm, S. W. & Suttle, C. A. Viruses and Nutrient Cycles in the Sea: Viruses play critical roles in the
structure and function of aquatic food webs. *Bioscience* **49**, 781–788 (1999).

7.   Ban, N. & Alder, J. How wild is the ocean? Assessing the intensity of anthropogenic marine activities
in British Columbia, Canada. *Aquat. Conserv.* **18**, 55–85 (2008).

8.   Li, D. & Daler, D. Ocean pollution from land-based sources: East China Sea, China. *Ambio* **33**, 107–
113 (2004).

9.   Sakowski, E. G. *et al.* Ribonucleotide reductases reveal novel viral diversity and predict biological and
ecological features of unknown marine viruses. *Proceedings of the National Academy of Sciences*
111, 15786–15791 (2014).

10. Harrison, A. O., Moore, R. M., Polson, S. W. & Wommack, K. E. Reannotation of the Ribonucleotide
Reductase in a Cyanophage Reveals Life History Strategies Within the Virioplankton. *Front.*
Microbiol. **10**, 134 (2019).

11. Lara, E. *et al.* Unveiling the role and life strategies of viruses from the surface to the dark ocean. *Sci*
Adv **3**, e1602565 (2017).

12. Fuhrman, J. A. Marine viruses and their biogeochemical and ecological effects. *Nature* **399**, 541–548
(1999).

13. Suttle, C. A. Viruses in the sea. *Nature* **437**, 356–361 (2005).

14. Wommack, K. E. & Colwell, R. R. Virioplankton: viruses in aquatic ecosystems. *Microbiol. Mol. Biol.*
Rev. **64**, 69–114 (2000).

15. Jiang, S. C. & Paul, J. H. Gene transfer by transduction in the marine environment. *Appl. Environ.*
Microbiol. **64**, 2780–2787 (1998).

16. A Schwartz, D. & Lindell, D. Genetic hurdles limit the arms race between Prochlorococcus and the
T7-like podoviruses infecting them. *ISME J.* **11**, (2017).

17. Ochman, H., Lawrence, J. G. & Groisman, E. A. Lateral gene transfer and the nature of bacterial
innovation. *Nature* **405**, 299–304 (2000).

18. Rosenwasser, S., Ziv, C., van Creveld, S. G. & Vardi, A. Virocell Metabolism: Metabolic Innovations

During Host–Virus Interactions in the Ocean. *Trends Microbiol.* **24**, 821–832 (2016).

19. Crummett, L. T., Puxty, R. J., Weihe, C., Marston, M. F. & Martiny, J. B. H. The genomic content and

context of auxiliary metabolic genes in marine cyanomyoviruses. *Virology* **499**, 219–229 (2016).

20. Thompson, L. R. *et al.* Phage auxiliary metabolic genes and the redirection of cyanobacterial host

carbon metabolism. *Proceedings of the National Academy of Sciences* **108**, E757–E764 (2011).

21. Weisburg, W. G., Barns, S. M., Pelletier, D. A. & Lane, D. J. 16S ribosomal DNA amplification for

phylogenetic study. *J. Bacteriol.* **173**, 697–703 (1991).

22. Schmidt, H. F., Sakowski, E. G., Williamson, S. J., Polson, S. W. & Wommack, K. E. Shotgun

metagenomics indicates novel family A DNA polymerases predominate within marine virioplankton.

ISME J. **8**, 103–114 (2014).

23. Brussaard, C. P. D., Short, S. M., Frederickson, C. M. & Suttle, C. A. Isolation and phylogenetic

analysis of novel viruses infecting the phytoplankton Phaeocystis globosa (Prymnesiophyceae). *Appl.*

Environ. Microbiol. **70**, 3700–3705 (2004).

24. Zhong, Y., Chen, F., Wilhelm, S. W., Poorvin, L. & Hodson, R. E. Phylogenetic diversity of marine

cyanophage isolates and natural virus communities as revealed by sequences of viral capsid

assembly protein gene g20. *Appl. Environ. Microbiol.* **68**, 1576–1584 (2002).

25. Short, C. M. & Suttle, C. A. Nearly identical bacteriophage structural gene sequences are widely

distributed in both marine and freshwater environments. *Appl. Environ. Microbiol.* **71**, 480–486

(2005).

26. Marston, M. F. & Sallee, J. L. Genetic diversity and temporal variation in the cyanophage community

infecting marine Synechococcus species in Rhode Island's coastal waters. *Appl. Environ. Microbiol.*

69, 4639–4647 (2003).

27. Goldin, S., Hulata, Y., Baran, N. & Lindell, D. Quantification of T4-Like and T7-Like Cyanophages

Using the Polony Method Show They Are Significant Members of the Virioplankton in the North

Pacific Subtropical Gyre. *Front. Microbiol.* **11**, 1210 (2020).

28. Comeau, A. M. & Krisch, H. M. The Capsid of the T4 Phage Superfamily: The Evolution, Diversity,

and Structure of Some of the Most Prevalent Proteins in the Biosphere. *Mol. Biol. Evol.* **25**, 1321–

1332 (2008).

29. Needham, D. M., Sachdeva, R. & Fuhrman, J. A. Ecological dynamics and co-occurrence among

marine phytoplankton, bacteria and myoviruses shows microdiversity matters. *ISME J.* **11**, 1614–

1629 (2017).

30. Liu, L., Cai, L. & Zhang, R. Co-existence of freshwater and marine T4-like myoviruses in a typical

subtropical estuary. *FEMS Microbiol. Ecol.* **93**, (2017).

31. Lundin, D., Gribaldo, S., Torrents, E., Sjöberg, B.-M. & Poole, A. M. Ribonucleotide reduction -

horizontal transfer of a required function spans all three domains. *BMC Evol. Biol.* **10**, 383 (2010).

32. Nordlund, P. & Reichard, P. Ribonucleotide Reductases. *Annu. Rev. Biochem.* **75**, 681–706 (2006).

33. Herrick, J. & Sclavi, B. Ribonucleotide reductase and the regulation of DNA replication: an old story

and an ancient heritage. *Mol. Microbiol.* **63**, 22–34 (2007).

34. Dwivedi, B., Xue, B., Lundin, D., Edwards, R. A. & Breitbart, M. A bioinformatic analysis of

ribonucleotide reductase genes in phage genomes and metagenomes. *BMC Evol. Biol.* **13**, 33 (2013).

35. Sullivan, M. B., Coleman, M. L., Weigele, P., Rohwer, F. & Chisholm, S. W. Three Prochlorococcus

cyanophage genomes: signature features and ecological interpretations. *PLoS Biol.* **3**, e144–e144

(2005).

36. McMurdie, P. J. & Holmes, S. Waste Not, Want Not: Why Rarefying Microbiome Data Is Inadmissible.

PLoS Comput. Biol. **10**, 1–12 (2014).

37. Weiss, S. *et al.* Normalization and microbial differential abundance strategies depend upon data

characteristics. *Microbiome* **5**, 27 (2017).

38. Gloor, G. B., Macklaim, J. M., Pawlowsky-Glahn, V. & Egozcue, J. J. Microbiome Datasets Are

Compositional: And This Is Not Optional. *Front. Microbiol.* **8**, 2224 (2017).

39. McLaren, M. R. *et al.* Consistent and correctable bias in metagenomic sequencing experiments. *Elife*

8, e46923 (2019).

40. Bian, G. *et al.* The Gut Microbiota of Healthy Aged Chinese Is Similar to That of the Healthy Young.

mSphere **2**, (2017).

41. Fernandes, A. D. *et al.* Unifying the analysis of high-throughput sequencing datasets: characterizing

RNA-seq, 16S rRNA gene sequencing and selective growth experiments by compositional data

analysis. *Microbiome* **2**, 15 (2014).

42. Publication Reports - EcosystemOverview_Azores_2019.pdf.

http://www.ices.dk/sites/pub/Publication%20Reports/Forms/DispForm.aspx?ID=36442

doi:10.17895/ices.advice.5753.

43. Bashmachnikov, I., Belonenko, T. V. & Koldunov, A. V. Intra-annual and interannual non-stationary

cycles of chlorophyll concentration in the Northeast Atlantic. *Remote Sens. Environ.* **137**, 55–68

(2013).

44. Arístegui, J. *et al.* Plankton metabolic balance at two North Atlantic seamounts. *Deep Sea Res. Part 2*

Top. Stud. Oceanogr. **56**, 2646–2655 (2009).

45. Mendonça, A. *et al.* Is there a seamount effect on microbial community structure and biomass? The

case study of Seine and Sedlo seamounts (northeast Atlantic). *PLoS One* **7**, e29526 (2012).

46. True Colors of Oceanography: Guidelines for Effective and Accurate Colormap Selection |

Oceanography. https://tos.org/oceanography/article/true-colors-of-oceanography-guidelines-for-

effective-and-accurate-colormap doi:10.5670/oceanog.2016.66.

47. Schlitzer, R. Interactive analysis and visualization of geoscience data with Ocean Data View. *Comput.*

Geosci. **28**, 1211–1218 (2002).

48. Sotillo, M. G. *et al.* The MyOcean IBI Ocean Forecast and Reanalysis Systems: operational products

and roadmap to the future Copernicus Service. *Journal of Operational Oceanography* **8**, 63–79

(2015).

49. Baith, K., Lindsay, R., Fu, G. & McClain, C. R. Data analysis system developed for ocean color

satellite sensors. *Eos Trans. AGU* **82**, 202–202 (2001).

50. Marie, D., Partensky, F., Vaulot, D. & Brussaard, C. Enumeration of phytoplankton, bacteria, and

viruses in marine samples. *Curr. Protoc. Cytom.* **Chapter 11**, Unit 11.11 (2001).

51. Iker, B. C., Kitajima, M. & Gerba, C. P. Extraction and Purification of Viral Nucleic Acids from

Environmental Samples. in *Sample Preparation Techniques for Soil, Plant, and Animal Samples* (ed.

Micic, M.) 315–324 (Springer New York, 2016). doi:10.1007/978-1-4939-3185-9_22.

52. Mueller, J. A., Culley, A. I. & Steward, G. F. Variables influencing extraction of nucleic acids from

microbial plankton (viruses, bacteria, and protists) collected on nanoporous aluminum oxide filters.

Appl. Environ. Microbiol. **80**, 3930–3942 (2014).

53. Fuhrman, J. A., Comeau, D. E., Hagström, A. & Chan, A. M. Extraction from natural planktonic

microorganisms of DNA suitable for molecular biological studies. *Appl. Environ. Microbiol.* **54**, 1426–

1429 (1988).

54. Fadrosh, D. W. *et al.* An improved dual-indexing approach for multiplexed 16S rRNA gene

sequencing on the Illumina MiSeq platform. *Microbiome* **2**, 6 (2014).

55. Moore, R. M. *et al.* PASV: Automatic protein partitioning and validation using conserved residues.

bioRxiv 2021.01.20.427478 (2021) doi:10.1101/2021.01.20.427478.

56. Edgar, R. C. Search and clustering orders of magnitude faster than BLAST. *Bioinformatics* **26**, 2460–

2461 (2010).

57. Price, M. N., Dehal, P. S. & Arkin, A. P. FastTree 2 -- Approximately Maximum-Likelihood Trees for

Large Alignments. *PLoS One* **5**, 1–10 (2010).

58. Moore, R. M., Harrison, A. O., McAllister, S. M., Polson, S. W. & Wommack, K. E. Iroki: automatic

customization and visualization of phylogenetic trees. *PeerJ* **8**, e8584 (2020).

59. Bolyen, E. *et al.* Reproducible, interactive, scalable and extensible microbiome data science using

QIIME 2. *Nat. Biotechnol.* **37**, 852–857 (2019).

60. Callahan, B. J. *et al.* DADA2: High-resolution sample inference from Illumina amplicon data. *Nat.*

Methods **13**, 581 EP– (2016).

61. Quast, C. *et al.* The SILVA ribosomal RNA gene database project: improved data processing and

web-based tools. *Nucleic Acids Res.* **41**, D590–D596 (2012).

62. Palarea-Albaladejo, J. & Martin-Fernandez, J. A. zCompositions -- R package for multivariate

imputation of left-censored data under a compositional approach. *Chemometrics Intellig. Lab. Syst.*

143, 85–96 (2015).

63. Team, R. C. *R: A Language and Environment for Statistical Computing.* (2013).

64. Gloor, G. B. & Reid, G. Compositional analysis: a valid approach to analyze microbiome high-

throughput sequencing data. *Can. J. Microbiol.* **62**, 692–703 (2016).

65. Mantel, N. The Detection of Disease Clustering and a Generalized Regression Approach. *Cancer*

Res. **27**, 209–220 (1967).

66. Willis, A. D. & Martin, B. D. DivNet: Estimating diversity in networked communities. *bioRxiv* (2018)
doi:10.1101/305045.

67. Wickham, H. *ggplot2: Elegant Graphics for Data Analysis*. (Springer, 2016).

68. Willis, A. & Bunge, J. Estimating diversity via frequency ratios. *Biometrics* **71**, 1042–1049 (2015).

69. Kang, S., Rodrigues, J. L. M., Ng, J. P. & Gentry, T. J. Hill number as a bacterial diversity measure
framework with high-throughput sequence data. *Sci. Rep.* **6**, 38263 (2016).

70. van den Boogaart, K. G., Tolosana-Delgado, R. & Bren, M. compositions: Compositional Data
Analysis. *Austral Ecol.* **26**, 32–46 (2018).

71. Anderson, M. J. A new method for non-parametric multivariate analysis of variance. *Austral Ecol.* **26**,
32–46 (2001).

72. Quinn, T. P., Richardson, M. F., Lovell, D. & Crowley, T. M. propr: An R-package for Identifying
Proportionally Abundant Features Using Compositional Data Analysis. *Sci. Rep.* **7**, 16252 (2017).

73. Erb, I. & Notredame, C. How should we measure proportionality on relative gene expression data?
Theory Biosci. **135**, 21–36 (2016).

74. Gu, Z., Eils, R. & Schlesner, M. Complex heatmaps reveal patterns and correlations in
multidimensional genomic data. *Bioinformatics* **32**, 2847–2849 (2016).

75. Alberti, A. *et al.* Viral to metazoan marine plankton nucleotide sequences from the Tara Oceans
expedition. *Sci Data* **4**, 170093 (2017).

76. Lundin, D., Torrents, E., Poole, A. M. & Sjöberg, B.-M. RNRdb, a curated database of the universal
enzyme family ribonucleotide reductase, reveals a high level of misannotation in sequences
deposited to Genbank. *BMC Genomics* vol. 10 Preprint at https://doi.org/10.1186/1471-2164-10-589
(2009).

77. Sakowski, E. G. *et al.* Ribonucleotide reductases reveal novel viral diversity and predict biological and
ecological features of unknown marine viruses. *Proceedings of the National Academy of Sciences*
vol. 111 15786–15791 Preprint at https://doi.org/10.1073/pnas.1401322111 (2014).

78. Wommack, K. E., Eri Wommack, K., Nasko, D. J., Chopyk, J. & Sakowski, E. G. Counts and
sequences, observations that continue to change our understanding of viruses in nature. *Journal of*
Microbiology vol. 53 181–192 Preprint at https://doi.org/10.1007/s12275-015-5068-6 (2015).

79. Bidle, K. D. Elucidating marine virus ecology through a unified heartbeat. *Proceedings of the National*
Academy of Sciences of the United States of America vol. 111 15606–15607 (2014).

80. Stubbe, J. Ribonucleotide reductases: the link between an RNA and a DNA world? *Curr. Opin. Struct.*
Biol. **10**, 731–736 (2000).

81. Forterre, P. The two ages of the RNA world, and the transition to the DNA world: a story of viruses
and cells. *Biochimie* **87**, 793–803 (2005).

82. Lundin, D., Berggren, G., Logan, D. T. & Sjöberg, B.-M. The origin and evolution of ribonucleotide
reduction. *Life* **5**, 604–636 (2015).

83. Martinez-Hernandez, F. *et al.* Single-virus genomics reveals hidden cosmopolitan and abundant

viruses. *Nat. Commun.* **8**, 15892 (2017).

84. Sakowski, E. G. *et al.* Interaction dynamics and virus-host range for estuarine actinophages captured

by epicPCR. *Nat Microbiol* **6**, 630–642 (2021).

85. UniProt: the universal protein knowledgebase in 2021. *Nucleic Acids Res.* **49**, D480–D489 (2021).

86. Torrents, E. Ribonucleotide reductases: essential enzymes for bacterial life. *Frontiers in Cellular and*

Infection Microbiology vol. 4 Preprint at https://doi.org/10.3389/fcimb.2014.00052 (2014).

87. Sañudo-Wilhelmy, S. A., Gómez-Consarnau, L., Suffridge, C. & Webb, E. A. The Role of B Vitamins

in Marine Biogeochemistry. *Annual Review of Marine Science* vol. 6 339–367 Preprint at

https://doi.org/10.1146/annurev-marine-120710-100912 (2014).

88. Doxey, A. C., Kurtz, D. A., Lynch, M. D. J., Sauder, L. A. & Neufeld, J. D. Aquatic metagenomes

implicate Thaumarchaeota in global cobalamin production. *ISME J.* **9**, 461–471 (2015).

89. Grossman, A. Nutrient Acquisition: The Generation of Bioactive Vitamin B12 by Microalgae. *Current*

biology: CB vol. 26 R319–21 (2016).

90. Heal, K. R. *et al.* Two distinct pools of B $_{12}$ analogs reveal community interdependencies in the ocean.

Proceedings of the National Academy of Sciences vol. 114 364–369 Preprint at

https://doi.org/10.1073/pnas.1608462114 (2017).

91. Sokolovskaya, O. M., Shelton, A. N. & Taga, M. E. Sharing vitamins: Cobamides unveil microbial

interactions. *Science* **369**, eaba0165 (2020).

92. Helliwell, K. E. *et al.* Cyanobacteria and Eukaryotic Algae Use Different Chemical Variants of Vitamin

B12. *Current Biology* vol. 26 999–1008 Preprint at https://doi.org/10.1016/j.cub.2016.02.041 (2016).

93. Poorvin, L., Rinta-Kanto, J. M., Hutchins, D. A. & Wilhelm, S. W. Viral release of iron and its

bioavailability to marine plankton. *Limnol. Oceanogr.* **49**, 1734–1741 (2004).

94. Giovannoni, S. J. Vitamins in the sea. *Proceedings of the National Academy of Sciences of the*

United States of America vol. 109 13888–13889 (2012).

95. Joglar, V. *et al.* Cobalamin and microbial plankton dynamics along a coastal to offshore transect in

the Eastern North Atlantic Ocean. *Environmental Microbiology* vol. 23 1559–1583 Preprint at

https://doi.org/10.1111/1462-2920.15367 (2021).

96. Sañudo-Wilhelmy, S. A. *et al.* Multiple B-vitamin depletion in large areas of the coastal ocean. *Proc.*

Natl. Acad. Sci. U. S. A. **109**, 14041–14045 (2012).

97. Mitra, R. D., Shendure, J., Olejnik, J., Edyta-Krzymanska-Olejnik & Church, G. M. Fluorescent in situ

sequencing on polymerase colonies. *Anal. Biochem.* **320**, 55–65 (2003).

98. Mruwat, N. *et al.* A single-cell polony method reveals low levels of infected Prochlorococcus in

oligotrophic waters despite high cyanophage abundances. *ISME J.* **15**, 41–54 (2021).

99. Baran, N., Goldin, S., Maidanik, I. & Lindell, D. Quantification of diverse virus populations in the

environment using the polony method. *Nat Microbiol* **3**, 62–72 (2018).

100.   Carlson, M. C. G. *et al.* Viruses affect picocyanobacterial abundance and biogeography in the

North Pacific Ocean. *Nat Microbiol* **7**, 570–580 (2022).

101.   Zuo, Y. & Deutscher, M. P. Exoribonuclease superfamilies: structural analysis and phylogenetic
1095   distribution. *Nucleic Acids Res.* **29**, 1017–1026 (2001).

1096   102.   Mathews, C. K. Deoxyribonucleotides as genetic and metabolic regulators. *FASEB J.* **28**, 3832–
1097   3840 (2014).

1098   103.   Varik, V., Oliveira, S. R. A., Hauryliuk, V. & Tenson, T. HPLC-based quantification of bacterial
1099   housekeeping nucleotides and alarmone messengers ppGpp and pppGpp. *Sci. Rep.* **7**, 11022 (2017).

1100   104.   Wheeler, L. J., Rajagopal, I. & Mathews, C. K. Stimulation of mutagenesis by proportional
1101   deoxyribonucleoside triphosphate accumulation in Escherichia coli. *DNA Repair* **4**, 1450–1456
1102   (2005).

1103   105.   Dedeo, C. L., Cingolani, G. & Teschke, C. M. Portal Protein: The Orchestrator of Capsid
1104   Assembly for the dsDNA Tailed Bacteriophages and Herpesviruses. *Annu Rev Virol* **6**, 141–160
1105   (2019).

1106   106.   Hatfull, G. F. Bacteriophage genomics. *Curr. Opin. Microbiol.* **11**, 447–453 (2008).

1107   107.   Hurwitz, B. L. & U'Ren, J. M. Viral metabolic reprogramming in marine ecosystems. *Curr. Opin.*
1108   *Microbiol.* **31**, 161–168 (2016).

1109   108.   Adriaenssens, E. M. & Cowan, D. A. Using signature genes as tools to assess environmental viral
1110   ecology and diversity. *Appl. Environ. Microbiol.* **80**, 4470–4480 (2014).

1111   109.   Shapiro, J. W. & Putonti, C. Gene Co-occurrence Networks Reflect Bacteriophage Ecology and
1112   Evolution. *MBio* **9**, (2018).

1113   110.   Overbeek, R. *et al.* The SEED and the Rapid Annotation of microbial genomes using Subsystems
1114   Technology (RAST). *Nucleic Acids Res.* **42**, D206–14 (2014).

1115   111. Katoh, K., Misawa, K., Kuma, K.-I. & Miyata, T. MAFFT: a novel method for rapid multiple sequence
1116   alignment based on fast Fourier transform. *Nucleic Acids Res.* **30**, 3059–3066 (2002).

1117   112. Kearse, M. *et al.* Geneious Basic: an integrated and extendable desktop software platform for the
1118   organization and analysis of sequence data. *Bioinformatics* **28**, 1647–1649 (2012).

1119   113. Caporaso, J. G. *et al.* QIIME allows analysis of high-throughput community sequencing data. *Nat.*
1120   *Methods* **7**, (2010).

1121   114. Altschul, S. F. *et al.* Gapped BLAST and PSI-BLAST: a new generation of protein database search
1122   programs. *Nucleic Acids Res.* **25**, 3389–3402 (1997).

1123   115. Katoh, K. & Standley, D. M. MAFFT multiple sequence alignment software version 7: improvements
1124   in performance and usability. *Mol. Biol. Evol.* **30**, 772–780 (2013).