# Quick-quick-slow: the foxtrot migration and dynamic non-breeding range

Ivan Pokrovsky[1] *, Teja Curk[2], Andreas Dietz[3], Ivan Fufachev[4], Olga Kulikova[5], Sebastian Rößler[3], Martin Wikelski[1,6]

[1] *Department of Migration, Max Planck Institute of Animal Behavior, Radolfzell, Germany*

[2] *Leibniz Institute for Zoo and Wildlife Research, Berlin, Germany*

[3] *Deutsches Zentrum für Luft- und Raumfahrt e.V. (DLR), Wessling, Germany*

[4] *Institute of Plant and Animal Ecology, Yekaterinburg, Russia*

[5] *Institute of Biological Problems of the North, Magadan, Russia*

[6] *Centre for the Advanced Study of Collective Behaviour, University of Konstanz, 78457 Konstanz, Germany*

\* *Corresponding author*

*E-mail: ipokrovsky@ab.mpg.de*

## ABSTRACT

Advancements in tracking technologies have revolutionized our understanding of bird migrations, revealing a diverse array of migratory behaviors. We propose a new pattern of migratory behavior termed 'foxtrot migration,' characterized by alternating quick and slow phases during the non-breeding period. This behavior involves directional and continuous movements, distinct from traditional seasonal itinerancy or forage-on-fly migration. Using the Rough-legged buzzard as a model species, we confirmed the presence of foxtrot migration and 'dynamic non-breeding ranges', driven by environmental factors such as snow cover dynamics. We advocate for accurate representation of dynamic ranges on maps and emphasize the need to consider range dynamics when assessing species conservation status. Our findings underscore the importance of understanding complex migratory behaviors in the face of environmental change, facilitated by advancements in tracking technologies. This knowledge is crucial for effective conservation strategies amid ongoing global environmental challenges.

Keywords: *migration patterns, bird movement ecology, bird migration, types of migration, conservation, climate change, species ranges, non-breeding range, snow cover, dynamic range, foxtrot migration.*

## INTRODUCTION

In recent decades, advanced tracking technology has greatly improved our knowledge of bird migrations (Wikelski *et al.* 2007; Jetz *et al.* 2022; Kays & Wikelski 2023). In the 19th century, the finding of a White stork *(Ciconia ciconia)* in Germany with an African arrow led to the discovery that birds did not hibernate at the bottom of water bodies during the winter but rather made extensive seasonal migrations (Richter & Bick 2018). Subsequent studies of bird migration in the 20th century using ringing techniques revealed a diversity of seasonal migratory behaviors, including 1) propensity to migrate (migrants, residents, partial migrants), 2) differences among populations (leapfrog, chain, parallel migrations), 3) connectivity and philopatry (constancy of breeding and non-breeding locations), 4) length of migration routes (long-distance and short-distance migrants), 5) variations in flight paces during migration (with stopovers, forage-on-fly, nonstop), and 6) tendencies to move within a single season (breeding and non-breeding itinerancy) (Alerstam, Hedenström & Åkesson 2003; Newton 2008; Alerstam 2011; Berthold, Gwinner & Sonnenschein 2013). Different combinations of these behaviors contribute to a wide variety of bird migration patterns (Chapman *et al.* 2014; Alerstam & Bäckman 2018; Lislevand *et al.* 2020). However, the advent of 21st-century technology, such as GPS transmitters, provides a higher resolution of movement and the ability to observe more details of migratory behavior (Wikelski *et al.* 2007; Jetz *et al.* 2022). Just as advances in image resolution allow us to see finer details in images, improvements in bird tracking technology now allow us to identify previously undiscovered migratory behaviors, revealing new patterns.

Current technologies allow observing forms of migratory behavior that are intermediate or integrative to those described above. Itinerancy was initially defined as the movements of Afro-Palearctic migrants during the non-breeding season in the Sahel, with a combination of stopovers and flights in between (Moreau 1972). Subsequently, researchers investigated this behavior for multiple species wintering in Africa, attributing their movements during the non-breeding season to changes in food availability predicted by NDVI (Zwarts *et al.* 2012; Schlaich *et al.* 2016; Thorup *et al.* 2017; Verhoeven *et al.* 2021). In 1998, Garcia and Arroyo hypothesized that certain species, like the Montagu's harrier *(Circus pygargus)*, might not only cover tens to hundreds of kilometers between various sites in their winter range but could also demonstrate directional, continuous movement throughout the entire non-breeding season. This movement may span a distance similar to their migration distance (ca. 1000 km), essentially indicating a prolonged, slower-paced continuation of migration (García & Arroyo 1998). Because this behavior is intermediate between seasonal movement (itinerancy) and the actual

migration process, we can describe it as either 'persistent directional non-breeding itinerancy' or

'very slow forage-on-fly migration in the non-breeding area'. Both options are clumsy and can

confuse. We suggest naming this behavior with a new term to avoid confusion with established

terms such as 'itinerancy' for any seasonal movement and 'forage-on-fly' denoting movement

from breeding to non-breeding sites (Alerstam, Hake & Kjellén 2006; Alerstam 2011; Lopez-

Ricaurte *et al.* 2024). Considering this behavior as an extension of migration and looking at the

complete life cycle of such species, it includes a quick phase during the transition between

breeding and non-breeding areas, followed by a slow phase of directed and seasonal movement

within the non-breeding area, and finally another quick phase towards the breeding area.

Analogous to the alternating fast and slow movements of the foxtrot dance, we propose the term

'foxtrot migration' for this seasonal movement to provide a concise and easily understandable

description (Figure 1a). Consequently, we suggest to term the non-breeding range of species

exhibiting this migration type as 'dynamic range' (Figure 1b).

**Figure 1. Foxtrot migration and dynamic winter range scheme.** a) Foxtrot migration. The
color bar along the y-axis corresponds to the color coding of the habitats in panel b of the chart
on the right. b) Dynamic winter range.

The concept of dynamic range, derived from foxtrot migration, is critical to

understanding the current range declines and shifts observed in many species. Typically, a

species' range is divided into areas of common and uncommon, with temporal heterogeneity

limited to breeding and non-breeding distinctions (Keller *et al.* 2020; Billerman *et al.* 2024).

However, suppose a species exhibits a foxtrot migration, moving gradually in one direction and

back again during the non-breeding season. In this case, it becomes inaccurate to represent its

non-breeding range as static. This discrepancy is particularly pronounced when ranges are

mapped using mid-winter surveys, such as Christmas counts. During this period, the species may

be present at the outer limits of its non-breeding range, leading to a false classification as

common in that area when, in fact, it is rare if the entire non-breeding season is considered.

Similarly, extended negative population dynamics of a species in a particular part of its non-

breeding range could be misinterpreted as a threat to the entire population. Indeed, if this species

has a foxtrot migration, range dynamics may have changed, possibly resulting in the species

ceasing to migrate to that particular part of its range. Advanced technologies now make it

possible the identification of foxtrot migration and range dynamics. This, in turn, facilitates

accurate range mapping and proper assessment of threats to the species.

This pattern of behavior has been proposed in species that overwinter in Africa, where

non-breeding movements are primarily related to the distribution of insect abundance (García &

Arroyo 1998). At the same time, it is reasonable to assume that this behavior may also be

characteristic of species that spend the non-breeding season in mid-latitudes, where food

availability may be linked to snow cover dynamics. These dynamics have changed particularly

rapidly in recent decades due to global warming (Davy & Outten 2020; Previdi, Smith & Polvani

2021). As a result, global climate change may significantly affect birds that breed in the Arctic

and winter in mid-latitudes. The Arctic is experiencing more pronounced effects of global

change than other regions, with declines in the abundance and range of several species and

populations (Post *et al.* 2009; Gilg *et al.* 2012; Schmidt *et al.* 2012). Therefore, understanding

how snow cover dynamics affect Arctic species across habitats throughout the annual cycle is

essential today. Furthermore, elucidating the foxtrot migration and dynamic non-breeding range

of mid-latitude wintering Arctic birds may provide valuable information on expected future

changes in their non-breeding range.

We used the Rough-legged buzzard *(Buteo lagopus)* as a model species to investigate this

phenomenon. The Rough-legged buzzard is an Arctic breeding and mid-latitude wintering raptor

(Ferguson-Lees & Christie 2001; Bechard & Swem 2002). Rough-legged buzzards feed mainly

on small rodents in the Arctic and its wintering grounds (Tast, Kaikusalo & Lagerström 2010;

Pokrovsky *et al.* 2014). They prefer open areas for hunting, and trees and tall bushes or uplands

for resting. In the Arctic, such areas are the southern and typical tundra (Walker *et al.* 2005), and

in the mid-latitudes, areas with fields and patches of forests (wooded fields). The taiga zone,

where there is little open space, is unsuitable for them, although they may nest in the northern

taiga zone on the border with the tundra zone (Sundell *et al.* 2004). Snow cover and day length

play an important role in their life (Terraube *et al.* 2015; Curk *et al.* 2020; Pokrovsky *et al.*

2021). Rough-legged buzzards can only hunt during the day (Pokrovsky *et al.* 2021), and heavy

snow cover makes hunting for small rodents problematic (Sonerud 1986; Vansteelant, Faveyts &

Buckens 2011). These two environmental factors, therefore, affect the availability of prey for

Rough-legged buzzards. At the same time, these factors vary considerably in mid-latitudes

during winter. Thus, prey availability for Rough-legged buzzards in the mid-latitudes increases

until mid-winter if they migrate southwards and after mid-winter if they migrate northwards. We,

therefore, assume that Rough-legged buzzards could track prey availability and experience a

foxtrot migration and dynamic non-breeding range.

For this study, we hypothesized that Rough-legged buzzards undertake a combined

foxtrot migration and dynamic non-breeding range in response to seasonal changes in mid-

latitude environmental factors. We made the following predictions. 1) Buzzards would exhibit a

directional and seasonal movement pattern during the non-breeding period, moving from the

northeast to the southwest and back again. This would result in a dynamic non-breeding range

that would continue to move geographically throughout the season. 2) Non-breeding movements

would occur in suitable open habitats, whereas fall and spring migrations would occur in

unfitting forested areas. 3) Non-breeding movements would differ from fall and spring

migrations in duration, extent, speed, and direction, however would continue throughout the

season. 4) During non-breeding movements, Rough-legged buzzards will experience less snow

cover than if they had stayed where they arrived at the end of the fall migration.

In the following, we will refer to the spring and fall migrations as the quick phase, the

non-breeding movement to the lowest point of latitude as the $1^{st}$ part of the slow phase, and the

movement from the lowest point of latitude to the starting point of the spring migration as the $2^{nd}$

part of the slow phase (Figure 1).

## MATERIAL AND METHODS

### Dataset

For this study, we tracked 43 adult Rough-legged buzzards (35 females and eight males) with the

solar GPS-GSM loggers (e-obs GmbH and UKn – University of Konstanz). The fieldwork was

carried out in the Russian Arctic in 2013-2019 at four study sites: Kolguev Island (69°16′N,

48°87′E), Nenetsky Nature Reserve (68°20′N, 53°18′E), Vaigach Island (69°43′N, 60°08′E), and

Yamal Peninsula (68°12′N, 68°59′E). For details on capture methods and tag parameters, see

Curk *et al.* (2022); for detailed study descriptions, see Pokrovsky *et al.* (2015) for Kolguev and

Pokrovsky *et al.* (2019) for Yamal and Nenetsky.

During data pre-processing, we estimated the date of death using the accelerometer or

GPS data and removed the tracking data if the bird was dead. We then removed duplicated

timestamps and calculated the mean daily positions of each individual. We partitioned the

resulting dataset into several periods: 1) breeding, 2) fall migration, 3)1st part of winter, 4) 2nd

part of winter, and 5) spring migration. We estimated the migration dates – the start and stop

dates of the spring and fall migrations – using an iterative search procedure for piecewise

regression described by Crawley (2007). We estimated the date between winter's first and

second parts as the day when the mean daily latitude was minimum.

## Data analysis

First, we used linear mixed-effects models (R function 'lmer' in the library 'lme4' (Bates *et al.*

2015)) to investigate whether or not Rough-legged buzzards migrated during winter. Latitude

was the response variable, day of the year was a fixed effect, and individuals and year were

included as random effects. Analyses were conducted separately for each migration period (fall,

first phase of winter, second phase of winter, and spring). For both phases of the winter

migration, we analyzed two additional models with longitude as the response variable instead of

latitude. Likelihood ratio tests were used to compare candidate models. The year was not a

calendar year but a year between two consecutive breeding seasons. Thus, fall migration,

consecutive winter, and consecutive spring have the same value for the year. The day of the year

was recalculated consecutively.

Second, we used linear mixed-effects models (R function 'lmer' in the library 'lme4') to

investigate whether migrations' parameters differ between the migration periods. We analyzed

four migration parameters: distance, duration, speed, and direction. The distance was calculated

as the distance between two coordinates (start and end of migration) using the R function 'distm'

in the library 'geosphere' (Hijmans 2016). The duration was calculated as the number of days

between the start and end of migration. Speed was calculated as the ratio of distance to duration.

The direction was calculated as the bearing from the start of the migration coordinates to the end

of the migration coordinates using the R function 'bearing' in the library 'geosphere' (Hijmans

2016). The migration parameter was used as the response variable, the type of migration as a

fixed factor, and individuals as a random factor. Likelihood ratio tests were used to compare

candidate models. We considered four different parameters of migration (distance, duration,

speed, and direction) and four types of migration (fall, first phase of winter, second phase of

winter, and spring). The analysis was done separately for each of the migration parameters.

Then, we used post hoc comparisons using the R function 'emmeans' in the library 'emmeans'

(Lenth *et al.* 2019) to compare the estimated means. In some raptor species, adult females

disperse further than males (Mearns & Newton 1984; Serrano *et al.* 2001; Bildstein 2006;

Whitfield *et al.* 2009). Therefore, we conducted an additional analysis on the effect of sex on

migration length using linear mixed-effects models (R function 'lmer' in the library 'lme4'). The

migration distance was used as the response variable, sex as a fixed factor, and individuals as a

random factor.

Third, we investigated whether vegetation land cover differed between areas crossed

during the quick (fall and spring) and slow (winter) migrations. We used the combined Terra and

Aqua Moderate Resolution Imaging Spectroradiometer (MODIS) Land Cover Climate

Modelling Grid (CMG) (MCD12C1) version 6 dataset (Friedl & Sulla-Menashe 2015). We used

a modified Leaf Area Index (LAI) as a classification scheme. We combined all four forest types

and savannas into one category (forest) and excluded the categories: water bodies, and

unclassified. We, therefore, had five types of vegetation cover: forest, grassland, cropland,

shrubland, and urban. We annotated the mean daily positions with the vegetation cover type

using the Env-DATA tool (Dodge *et al.* 2013). We used general linear mixed effects models

with a binomial distribution (R function 'glmer' in the library 'lme4') to investigate whether

vegetation cover types differ between migration periods. Presence/absence of the studied

vegetation land cover type was used as a response variable, migration type as a fixed factor, and

individuals as a random factor. The analysis was done separately for each of the vegetation land

cover types.

Fourth, we investigated whether snow cover could drive the slow migration phenomenon.

We then compared the snow cover conditions the birds experienced during the winter with two

hypothetical snow cover conditions that the birds would have experienced if they had not

migrated during the winter. The first hypothetical snow cover condition would have happened if

the birds had stayed where they arrived from the north (i.e., where their fall migration ended). To

estimate this parameter, we calculated the winter dynamics of the average snow cover at the

minimum convex polygons (MCP) occupied by the birds in October and April (northeast of their

winter range). A second hypothetical snow cover condition would be if the birds flew

immediately to the southwest and spent the whole winter there. To evaluate this, we calculated

the winter dynamics of average snow cover on the MCPs occupied by the birds in January and

February (southwest of their winter range). We then compared the values obtained for the real

snow cover and two hypothetical snow covers using general linear mixed effects models with a

binomial distribution (R function 'glmer' in the 'lme4' library). Presence/absence of snow cover

was used as a response variable, type of snow cover (real, 1st hypothetical, or 2nd hypothetical) as

a fixed factor, and years as a random factor. The analysis was done separately for each month.

We then used post hoc comparisons to compare the estimated means, using the R function

'emmeans' in the 'emmeans' library (Lenth *et al.* 2019).

We obtained monthly snow cover data with a spatial resolution of ca 500 meters (Global

SnowPack MODIS) from the German Aerospace Center (DLR). This product is based on the

Moderate Resolution Imaging Spectroradiometer (MODIS) daily snow cover products

MOD10A1 and MYD10A1 (version 6 as provided by the National Snow and Ice Data Center

NSIDC), which have been processed to remove the gaps due to cloud cover and polar darkness

(Dietz, Kuenzer & Dech 2015). These processing steps include a combination of data available

from different satellites (Aqua and Terra), 3-day temporal moving window filtering, a regional

snow line elevation interpolation relying on a Digital Elevation Model (DEM), and a seasonal

filter running through the time series for the whole hydrological year (1st of September through

August 31st). The proportion of days in which one pixel is snow-covered per month is referred to

here as fractional snow cover and is derived from these daily gap-filled rasters. Five MODIS

tiles (h19v03, h20v03, h20v04, h21v03 and h21v04) were mosaicked and re-projected to

WGS84. Then, for each month from October to April, we calculated 95% minimum convex

polygons (MCPs) for the distribution of Rough-legged buzzards using the R function 'mcp' in

library 'adehabitatHR' (Calenge 2006). We extracted mean snow cover values from each MCP

from every monthly snow cover raster separately, using the R library 'raster' (Hijmans & van

Etten 2023).

All calculations were performed using R version 4.2.2 'Innocent and Trusting' (R

Development Core Team 2022) and RStudio version 353 'Elsbeth Geranium' (Posit team 2022).

## RESULTS

### Foxtrot migration – always on the move

Except during the breeding season, Rough-legged buzzard migration continues throughout the

year, even after the birds´ arrival at their traditionally recognized ´wintering grounds´ (Figure 2a,

b). For both quick and slow phases of the foxtrot migration, linear mixed-effects models with the

season as a fixed factor received higher support from the likelihood ratio test (p < 0.001, Tables

S1-S3). Rough-legged buzzards started their fall migration (quick phase) on 28 September

(hereafter mean±sd for the day of the year: 271±11, n=31) and ended on 12 October (285±11,
n=33). The mean latitude/longitude where the birds ended their fall migration was
55.57±1.92°/49.35±5.63° (Figure 2a). During the winter, birds continued to migrate at a slower
pace down to 49.53±2.01° latitude (on 5 February, 36±40, n=23) and 34.29±5.11° longitude (on
24 January, 24±47, n=23). Afterward, during the second part of the slow phase, the birds
returned to 55.52±2.63° latitude and 49.79±8.24° longitude to start the spring migration (Figure
2a). Rough-legged buzzards started their spring migration to the Arctic on 27 April (117±7,
n=27) and arrived at the breeding grounds on 15 May (135±8, n=18).

**Figure 2. Foxtrot migration of Rough-legged buzzards.** Q – quick phase. Qf – Quick fall
phase (orange), Qs – Quick spring phase (yellow), S1 – Slow phase, 1$^{st}$ part (light blue), S2 –
Slow phase, 2$^{nd}$ part (dark blue). a) Change in the latitude of 43 Rough-legged buzzards during
the year, red line – mean latitude of all birds, black vertical lines – mean dates of start and end of
the migration phases, blue vertical line – mean date of the minimum latitude. Grey, sky blue, and

piggy pink shaded areas – standard deviation of the means. b) Migration map. c) Difference in
the migration parameters between the migration phases. Lines on the direction plot (down, right)
represent the mean value for each bird; arrows represent the mean direction.

### Quick-slow phase features comparison

During quick phase, individual birds flew greater distances in a shorter time, i.e., at a
faster rate, than during slow phase. After arriving at what is traditionally known as the wintering
grounds, the direction of migration changed, so the direction of quick and slow phases also
differed (Table 1, Figure 2c). The quick phase was 1415±50 km long, whereas the slow phase
(one part) was 1026±55 km, i.e., 389±60 km shorter (p<0.001, Table S4, Figure 2c). During
quick phase, birds flew for 15±3 days, and one part of slow phase lasted 100±4 days, i.e., 85±5
days longer (p<0.001, Table S5, Figure 2c). At the same time, the second part of slow phase was
54±7 days shorter than the first (p<0.001, Table S5, Figure 2c). The migration speed was 104±6
km/day during the quick phase and 12±7 km/day during the slow phase, i.e., about eight times
higher (p<0.001, Table S6, Figure 2c). During the fall migration, birds moved in the SSW
direction (7±2 deg), then turned 50±3 deg (p<0.001, Table S7, Figure 2c) to the west and started
their 1$^{st}$ slow phase until mid-winter. After that, they turned back to the NEE direction (57±2
deg) and performed their 2$^{nd}$ slow phase for several months until they turned 54±3 deg (p<0.001,
Table S7, Figure 2c) to the north and started their spring migration. As a result of additional
analysis of the effect of sex on migration length, we found no significant difference between the
migration distances of males and females (Table S8).

**Table 1.** Parameters of the rough-legged buzzards' migration.

| Phase of migration | Sub-phase of migration | Distance (km) | Duration (days) | Speed (km/day) | Direction (deg) |
|---|---|---|---|---|---|
| Quick | | 1415±50 | 15±3 | 104±6 | |
| | Spring | 1544±72 | 18±4 | 100±8 | 7±2 |
| | Fall | 1334±59 | 14±4 | 107±7 | 198±3 |
| Slow | | 1026±55 | 100±4 | 12±7 | |
| | 1$^{st}$ phase | 1016±71 | 121±4 | 8±7 | 251±3 |

| | 2$^{nd}$ phase | 1042±71 | 66±6 | 17±9 | 57±2 |
|---|---|---|---|---|---|

## Vegetation land cover during migration

During quick phases, Rough-legged buzzards cross the forest zone, while during slow phase,
they migrate within the grassland and cropland zone (Figure 3). Rough-legged buzzards migrated
fast across the tundra zone on the north in the Arctic and then through the taiga zone. Therefore,
during quick phase, the three most common vegetation land cover types were forest (44.5±2.9 %,
hereafter, percentage of all mean daily positions annotated with the given vegetation type),
shrublands (29.9±3 %), and grasslands (24.7±2.8 %, Figure 3a). During slow phase, the three
most common vegetation land cover types were grasslands (65.1±6.2 %), croplands (26.9±6.3
%), and forests (4.9±1.4 %, Figure 3b). According to the linear mixed-effects models, the
percentage of all vegetation land cover types differed between the slow and quick phases
(p<0.001, Table S9), except for the urban lands. Urban lands were more common during the
slow than quick phase (Figure 3). However, this type has been annotated for too few birds to
make an adequate comparison.

**Figure 3. Vegetation land cover during quick and slow phases.** a) Quick phase (spring and
fall periods together). b) Slow phase ($1^{st}$ and $2^{nd}$ parts together). c) Migration map.

## Snow cover – the main reason for the dynamic winter range

During the slow phase of the foxtrot migration, Rough-legged buzzards experienced snow cover
ranging from 4.8±1.0% in October to 85.2±4.6% in February (Figure 4). If birds spent the winter
in the place where they arrived after the fall migration, they would experience snow cover
conditions ranging from 4.6±0.6% in October to 99.5±0.1% in February (Figure 4). And if birds
fly directly to the southeast and stay there for the whole winter, they would experience snow
cover conditions ranging from 1.4±0.2 % in October to 81.1±5.0 % in January (Figure 4). Thus,
if birds fly immediately to the southwest and stay there until the end of the winter, they will find
conditions with less snow cover in spring ($p < 0.001$, Table S10). And if birds stay where they
ended the fall migration, they will find themselves in situations with more snow cover ($p < 0.001$,
Table S10). In the latter case, the difference between real and hypothetical situations is not as
pronounced (85.2% vs. 99.5%), but more means that snow cover will be close to 100% for
several months in this hypothetical situation (Figure 4b).

**Figure 4. Snow cover conditions during the slow phase of the foxtrot migration.** a) 95%
Minimum convex polygons (MCPs) of Rough-legged buzzards during winter. Arrows indicate
the direction of the movement across months. OCT – October, NOV – November, DEC –
December, JAN – January, FEB – February, MAR – March, APR – April. b) Snow cover
conditions for the real situation (black) and two hypothetical situations – if birds spend the
winter in the place where they arrived after the fall migration (green) and if birds fly directly to
the southwest and stay there all winter (red). Dots represent mean values, error lines – standard
errors.

In some raptor species, adult females disperse further than males (Mearns & Newton
1984; Serrano *et al.* 2001; Bildstein 2006; Whitfield *et al.* 2009). While our analysis does not
support this (Table S8), it should be acknowledged that our dataset comprises 35 females, eight
males, and no juveniles. The unbalanced sex ratio and absence of immature individuals in our
sample may have potential influences towards the observed movement patterns in this study.

## DISCUSSION

Our study confirmed the existence of a previously hypothesized bird migration pattern (García & Arroyo 1998) characterized by alternating quick and slow phases (Figure 1). Following the breeding season, birds quickly traverse unfavorable habitats and proceed slowly through winter in favorable environments influenced by external factors. Subsequently, they quickly cross unfavorable terrain once more to reach the breeding region in spring (Figure 2). This pattern, termed 'foxtrot migration', distinguishes itself from 'itinerancy' by featuring directional and continuous movements. Additionally, it differs from 'forage-on-fly migration' as it occurs during the non-breeding period. We propose referring to the non-breeding area as 'dynamic area', reflecting its temporal heterogeneity, where the region with the highest bird density continually changes throughout the non-breeding period.

Our study affirmed the presence of foxtrot migration and a dynamic non-breeding range in Rough-legged Buzzards. The taiga zone posed as unfavorable habitat during the quick phase of foxtrot migration (Figure 3), given the difficulty for Rough-legged Buzzards to locate open areas for hunting. Conversely, the grassland and cropland zone served as favorable habitats during the slow phase throughout the entire non-breeding period (Figure 3), offering numerous open areas for hunting. Snow cover was the external factor driving their continual 1000 km southwest movement during winter (Figure 4). Our analysis revealed that if Rough-legged Buzzards remained at their fall migration endpoint without moving southwest, they would encounter 14.4% more snow cover (99.5% vs. 85.1%, Figure 4). Although this difference may seem small (14.4%), it holds significance for rodent-hunting birds, distinguishing between complete and patchy snow cover. Simultaneously, if Rough-legged Buzzards immediately flew to the southwest and stayed there throughout winter, they would experience 25.7% less snow cover (57.3% vs. 31.6%, Figure 4). Despite a greater difference than in the first case, it doesn't compel them to adopt this strategy, as it represents the difference between various degrees of landscape openness from snow cover. This indicates ecological tipping points in the environment that are not predicted by a slight linear increase in effect size.

This pattern is likely to be observed in many migratory species with distinct seasonal cycles in their non-breeding range. In our study, we focused on an Arctic species that resides in the mid-latitudes during the non-breeding period. In general, foxtrot migration is likely to be a common feature among species where snow cover significantly influences food availability, especially those specializing in rodents. Originally proposed for birds in Africa, where seasonal changes regulate food availability, it's likely that many insectivorous birds that spend the non-

378 breeding season in Africa also exhibit this form of migration. This type of migration and
379 dynamic non-breeding range is widespread both geographically and taxonomically, with obvious
380 conservation implications.

## The conservation implications

The conservation implications of our study are twofold: 1) the use of mid-winter (Christmas)
bird surveys to determine non-breeding range may yield inaccurate results for species with
foxtrot migrations (Figure 5a), and 2) declines in abundance within a particular segment of the
non-breeding range may indicate changes in range dynamics rather than widespread declines in
species abundance (Figure. 5b).

**Figure 5. The use of foxtrot migration research for the assessment of the non-breeding**
**range and conservation status of a particular species.** a) Christmas counts, often used to
determine winter range, may give a misleading representation of non-breeding range for species
with foxtrot migration. b) Changing the conservation status of a species based on reduced
abundance in a particular area may not be appropriate for species with foxtrot migrations. Red –
breeding range, blue – winter range. Latin numbers indicate the month the birds are present in
the area.

Christmas Bird Counts provide insight into the mid-winter distribution of a species. For

example, based on our study of the Rough-legged Buzzard, the species is predominantly present

in the southwestern portion of its non-breeding range during this period, with only a small

proportion present in the rest of the region. As a result, a map of the non-breeding range may

inaccurately depict the species as common in the southwest and extremely rare in the northeast.

This map would be inaccurate because, during the entire non-breeding period, Rough-legged

Buzzards spend both fall and spring in the northeastern part and only mid-winter in the

southwesternmost part.

To address this, continuous year-round GPS tracking of the species provides a means to

track bird locations throughout the non-breeding season, facilitating the creation of accurate

distribution maps, particularly for species that exhibit foxtrot migrations and dynamic ranges.

We advocate representing temporal heterogeneity (range dynamics) on maps as distinct zones,

denoting periods when the species is abundant in a given area. To distinguish temporal from

spatial heterogeneity, we recommend using lines to delineate the boundaries of these zones,

rather than color shading, and incorporating numbers to denote the months of species abundance

in a given zone (Figure 5a). We suggest ecologists include dynamic non-breeding ranges in

descriptions and range maps for foxtrot migratory bird species.

Population counts for a species are often limited to a portion of its range. Therefore,

conclusions about conservation status drawn from such counts may be misleading. A decline in

abundance within a particular portion of the non-breeding range may indicate changes in range

dynamics rather than a general decline in the species. For example, climate change may affect

snow cover dynamics, reducing its intensity in northern regions. As a result, species whose range

dynamics depend on snow cover may choose to remain in the northern areas and not migrate as

far south as they traditionally have. Despite this shift in range dynamics, overall species

abundance may remain unchanged. Similar patterns have affected Rough-legged Buzzards in

some areas of the European non-breeding range.

A 2022 Dutch study found a decline in wintering Rough-legged buzzards over the last 40

years (Hornman, Boele & van Winden 2022). On the one hand, this may represent a

conservation concern. On the other hand, applying the rationale of the foxtrot migration, the

apparent local decline may simply be attributed to climate change, resulting in less

comprehensive snow coverage in the northeastern wintering areas of Rough-legged buzzards

relative to the Netherlands. Such a shift in snow coverage makes it less probable for the birds to

migrate to the Netherlands for overwintering. This proposition is further supported by a study of

the winter population dynamics of Rough-legged buzzards in the Netherlands in 2011, showing
that the main winter population peak occurred in late December, with many birds migrating
(Vansteelant, Faveyts & Buckens 2011). The authors also found that the main migration
occurred after heavy snowfall in northern Europe, supporting our foxtrot migration explanation
for this decline. Therefore, investigating the dynamic range and foxtrot migration is critical to
understanding a species' range and effectively assessing its conservation status.

Significant environmental changes, including changes in snowpack dynamics, have
occurred in recent decades that may have significant impacts on bird migration (Tucker *et al.*
2018; Sumasgutner *et al.* 2021). In order to understand the effects of global environmental
change on bird migration behavior, there is an urgent need for in-depth investigations into the
intricate relationship between migration patterns and evolving environmental factors. The advent
of new technologies in recent decades has expanded our understanding of bird migration,
enabling the identification of novel behavioral patterns and facilitating the study of their
susceptibility to various environmental influences. This knowledge is becoming increasingly
urgent in the face of rapid climate change and anthropogenic habitat alteration, which pose
threats to both wildlife and human populations.

## CONCLUSIONS

Our study has identified and characterized a new pattern of migratory behavior, the 'foxtrot
migration', along with the associated concept of 'dynamic range'. This discovery has significant
implications for conservation strategies and adequate representation of non-breeding habitats. As
animal tracking technology advances, we expect to discover more new details about animal
migrations and, consequently, new patterns of migratory behavior. With these technological
developments, we are gaining a clearer understanding of the complex patterns and behaviors that
drive animal migrations, providing valuable insights for conservation efforts and deepening our
understanding of the natural world.

## ACKNOWLEDGMENTS

454 We are very grateful to everyone who helped us collect data in the field. This study was funded
455 by the Max-Planck Institute of Animal Behavior and the German Air and Space Administration
456 (DLR). We also acknowledge partial funding by the Deutsche Forschungsgemeinschaft (DFG,
457 German Research Foundation) under Germany's Excellence Strategy – EXC 2117 – 422037984.

## REFERENCES

Alerstam, T. (2011) Optimal bird migration revisited. *Journal of Ornithology,* **152,** 5-23.

Alerstam, T. & Bäckman, J. (2018) Ecology of animal migration. *Current Biology,* **28,** R968-R972.

Alerstam, T., Hake, M. & Kjellén, N. (2006) Temporal and spatial patterns of repeated migratory journeys by ospreys. *Journal of Animal Behaviour,* **71,** 555-566.

Alerstam, T., Hedenström, A. & Åkesson, S. (2003) Long-distance migration: evolution and determinants. *Oikos,* **103,** 247-260.

Bates, D., Mächler, M., Bolker, B. & Walker, S. (2015) Fitting Linear Mixed-Effects Models Using lme4. *Journal of Statistical Software,* **067**.

Bechard, M.J. & Swem, T.R. (2002) Rough-legged Hawk (Buteo lagopus), The Birds of North America Online (A. Poole, Ed.). Ithaca: Cornell Lab of Ornithology.

Berthold, P., Gwinner, E. & Sonnenschein, E. (2013) *Avian migration*. Springer Science & Business Media.

Bildstein, K.L. (2006) *Migrating raptors of the world: their ecology & conservation*. Cornell University Press.

Billerman, S.M., Keeney, B.K., Rodewald, P.G. & Schulenberg, T.S. (2024) Birds of the World. Cornell Laboratory of Ornithology, Ithaca, NY, USA.

Calenge, C. (2006) The package "adehabitat" for the R software: A tool for the analysis of space and habitat use by animals. *Ecological Modelling,* **197,** 516-519.

Chapman, B.B., Hulthén, K., Wellenreuther, M., Hansson, L.-A., Nilsson, J.-Å. & Brönmark, C. (2014) Patterns of animal migration. *Animal movement across scales* (eds L.-A. Hansson & Å. Susanne), pp. 11-35. Oxford University Press.

Crawley, M.J. (2007) *The R Book*. Wiley Publishing.

Curk, T., Kulikova, O., Fufachev, I., Wikelski, M., Safi, K. & Pokrovsky, I. (2022) Arctic Migratory Raptor Selects Nesting Area During the Previous Breeding Season. *Frontiers in Ecology and Evolution,* **10**.

Curk, T., Pokrovsky, I., Lecomte, N., Aarvak, T., Brinker, D.F., Burnham, K., Dietz, A., Dixon, A., Franke, A., Gauthier, G., Jacobsen, K.-O., Kidd, J., Lewis, S.B., Øien, I.J., Sokolov, A., Sokolov, V., Solheim, R., Weidensaul, S., Wiebe, K., Wikelski, M., Therrien, J.-F. & Safi, K. (2020) Arctic avian predators synchronise their spring migration with the northern progression of snowmelt. *Scientific Reports,* **10,** 7220.

Davy, R. & Outten, S. (2020) The Arctic surface climate in CMIP6: status and developments since CMIP5. *Journal of Climate,* **33,** 8047-8068.

Dietz, A.J., Kuenzer, C. & Dech, S. (2015) Global SnowPack: a new set of snow cover parameters for studying status and dynamics of the planetary snow cover extent. *Remote Sensing Letters,* **6,** 844-853.

Dodge, S., Bohrer, G., Weinzierl, R., Davidson, S.C., Kays, R., Douglas, D., Cruz, S., Han, J., Brandes, D. & Wikelski, M. (2013) The environmental-data automated track annotation (Env-DATA) system: linking animal tracks with environmental data. *Movement Ecology,* **1,** 3.

Ferguson-Lees, J. & Christie, D. (2001) *Raptors of the World*. Princeton University Press.

Friedl, M. & Sulla-Menashe, D. (2015) MCD12C1 MODIS/Terra+Aqua Land Cover Type Yearly L3 Global 0.05Deg CMG V006.

García, J.T. & Arroyo, B.E. (1998) Migratory movements of western European Montagu's Harrier Circus pygargus: a review. *Bird Study,* **45,** 188-194.

Gilg, O., Kovacs, K.M., Aars, J., Fort, J., Gauthier, G., Gremillet, D., Ims, R.A., Meltofte, H., Moreau, J., Post, E., Schmidt, N.M., Yannic, G. & Bollache, L. (2012) Climate change

and the ecology and evolution of Arctic vertebrates. *Annals of the New York Academy of*
*Sciences,* **1249,** 166-190.

Hijmans, R.J. (2016) geosphere: Spherical Trigonometry. R package version 1.5-5.

Hijmans, R.J. & van Etten, J. (2023) raster: Geographic data analysis and modeling.

Hornman, M., Boele, A. & van Winden, E. (2022) Ruigpootbuizerd: steeds schaarser, behalve in
piekjaren. Nature Today.

Jetz, W., Tertitski, G., Kays, R., Mueller, U., Wikelski, M., Åkesson, S., Anisimov, Y., Antonov,
A., Arnold, W., Bairlein, F., Baltà, O., Baum, D., Beck, M., Belonovich, O., Belyaev, M.,
Berger, M., Berthold, P., Bittner, S., Blake, S., Block, B., Bloche, D., Boehning-Gaese,
K., Bohrer, G., Bojarinova, J., Bommas, G., Bourski, O., Bragin, A., Bragin, A., Bristol,
R., Brlík, V., Bulyuk, V., Cagnacci, F., Carlson, B., Chapple, T.K., Chefira, K.F., Cheng,
Y., Chernetsov, N., Cierlik, G., Christiansen, S.S., Clarabuch, O., Cochran, W.,
Cornelius, J.M., Couzin, I., Crofoot, M.C., Cruz, S., Davydov, A., Davidson, S., Dech,
S., Dechmann, D., Demidova, E., Dettmann, J., Dittmar, S., Dorofeev, D., Drenckhahn,
D., Dubyanskiy, V., Egorov, N., Ehnbom, S., Ellis-Soto, D., Ewald, R., Feare, C.,
Fefelov, I., Fehérvári, P., Fiedler, W., Flack, A., Froböse, M., Fufachev, I., Futoran, P.,
Gabyshev, V., Gagliardo, A., Garthe, S., Gashkov, S., Gibson, L., Goymann, W., Gruppe,
G., Guglielmo, C., Hartl, P., Hedenström, A., Hegemann, A., Heine, G., Ruiz, M.H.,
Hofer, H., Huber, F., Hurme, E., Iannarilli, F., Illa, M., Isaev, A., Jakobsen, B., Jenni, L.,
Jenni-Eiermann, S., Jesmer, B., Jiguet, F., Karimova, T., Kasdin, N.J., Kazansky, F.,
Kirillin, R., Klinner, T., Knopp, A., Kölzsch, A., Kondratyev, A., Krondorf, M., Ktitorov,
P., Kulikova, O., Kumar, R.S., Künzer, C., Larionov, A., Larose, C., Liechti, F., Linek,
N., Lohr, A., Lushchekina, A., Mansfield, K., Matantseva, M., Markovets, M., Marra, P.,
Masello, J.F., Melzheimer, J., Menz, M.H.M., Menzie, S., Meshcheryagina, S., Miquelle,
D., Morozov, V., Mukhin, A., Müller, I., Mueller, T., Navedo, J.G., Nathan, R., Nelson,
L., Németh, Z., Newman, S., Norris, R., Nsengimana, O., Okhlopkov, I., Oleś, W.,
Oliver, R., O'Mara, T., Palatitz, P., Partecke, J., Pavlick, R., Pedenko, A., Perry, A.,
Pham, J., Piechowski, D., Pierce, A., Piersma, T., Pitz, W., Plettemeier, D., Pokrovskaya,
I., Pokrovskaya, L., Pokrovsky, I., Pot, M., Procházka, P., Quillfeldt, P., Rakhimberdiev,
E., Ramenofsky, M., Ranipeta, A., Rapczyński, J., Remisiewicz, M., Rozhnov, V.,
Rienks, F., Rozhnov, V., Rutz, C., Sakhvon, V., Sapir, N., Safi, K., Schäuffelhut, F.,
Schimel, D., Schmidt, A., Shamoun-Baranes, J., Sharikov, A., Shearer, L., Shemyakin,
E., Sherub, S., Shipley, R., Sica, Y., Smith, T.B., Simonov, S., Snell, K., Sokolov, A.,
Sokolov, V., Solomina, O., Soloviev, M., Spina, F., Spoelstra, K., Storhas, M., Sviridova,
T., Swenson, G., Taylor, P., Thorup, K., Tsvey, A., Tucker, M., Tuppen, S., Turner, W.,
Twizeyimana, I., van der Jeugd, H., van Schalkwyk, L., van Toor, M., Viljoen, P., Visser,
M.E., Volkmer, T., Volkov, A., Volkov, S., Volkov, O., von Rönn, J.A.C., Vorneweg, B.,
Wachter, B., Waldenström, J., Weber, N., Wegmann, M., Wehr, A., Weinzierl, R.,
Weppler, J., Wilcove, D., Wild, T., Williams, H.J., Wilshire, J., Wingfield, J., Wunder,
M., Yachmennikova, A., Yanco, S., Yohannes, E., Zeller, A., Ziegler, C., Zięcik, A. &
Zook, C. (2022) Biological Earth observation with animal sensors. *Trends in Ecology &*
*Evolution,* **37,** 293-298.

Kays, R. & Wikelski, M. (2023) The Internet of Animals: what it is, what it could be. *Trends in*
*Ecology & Evolution.*

Keller, V., Herrando, S., Voříšek, P., Franch, M., Kipson, M., Milanesi, P., Martí, D., Anton, M.,
Klvaňová, A., Kalyakin, M.V., Bauer, H.-G. & Foppen, R.P.B. (2020) *European*
*Breeding Bird Atlas 2: Distribution, Abundance and Change.* European Bird Census
Council & Lynx Edicions, Barcelona.

Lenth, R., Singmann, H., Love, J., Buerkner, P. & Herve, M. (2019) Package 'emmeans'.

Lislevand, T., Hahn, S., Rislaa, S. & Briedis, M. (2020) First records of complete annual cycles
in water rails Rallus aquaticus show evidence of itinerant breeding and a complex
migration system. *Journal of Avian Biology,* **51**.

Lopez-Ricaurte, L., Vansteelant, W.M.G., Hernández-Pliego, J., García-Silveira, D., Casado, S.,
Garcés-Toledano, F., Martínez-Dalmau, J., Ortega, A., Rodríguez-Moreno, B. &
Bustamante, J. (2024) Itinerant lifestyle and congregation of lesser kestrels in West
Africa. *Journal of Avian Biology,* **2024,** e03063.

Mearns, R. & Newton, I. (1984) Turnover and dispersal in a peregrine Falco peregrinus
population. *Ibis,* **126,** 347-355.

Moreau, R.E. (1972) *The Palaearctic-African Bird Migration Systems*. Academic Press.

Newton, I. (2008) *The Migration Ecology of Birds*. Elsevier, Oxford, UK.

Pokrovsky, I., Ehrich, D., Fufachev, I., Ims, R.A., Kulikova, O., Sokolov, A., Sokolova, N.,
Sokolov, V. & Yoccoz, N.G. (2019) Nest association between two predators as a
behavioral response to the low density of rodents. *The Auk,* **137**.

Pokrovsky, I., Ehrich, D., Ims, R., Kulikova, O., Lecomte, N. & Yoccoz, N. (2014) Diet, nesting
density, and breeding success of rough-legged buzzards (Buteo lagopus) on the Nenetsky
Ridge, Arctic Russia. *Polar Biology,* **37,** 447-457.

Pokrovsky, I., Ehrich, D., Ims, R.A., Kondratyev, A.V., Kruckenberg, H., Kulikova, O.,
Mihnevich, J., Pokrovskaya, L. & Shienok, A. (2015) Rough-Legged Buzzards, Arctic
Foxes and Red Foxes in a Tundra Ecosystem without Rodents. *Plos One,* **10,** e0118740.

Pokrovsky, I., Kölzsch, A., Sherub, S., Fiedler, W., Glazov, P., Kulikova, O., Wikelski, M. &
Flack, A. (2021) Longer days enable higher diurnal activity for migratory birds. *Journal
of Animal Ecology,* **90,** 2161-2171.

Posit team (2022) RStudio: Integrated Development Environment for R. Posit Software, PBC,
Boston, MA.

Post, E., Forchhammer, M.C., Bret-Harte, M.S., Callaghan, T.V., Christensen, T.R., Elberling,
B., Fox, A.D., Gilg, O., Hik, D.S., Hoye, T.T., Ims, R.A., Jeppesen, E., Klein, D.R.,
Madsen, J., McGuire, A.D., Rysgaard, S., Schindler, D.E., Stirling, I., Tamstorf, M.P.,
Tyler, N.J.C., van der Wal, R., Welker, J., Wookey, P.A., Schmidt, N.M. & Aastrup, P.
(2009) Ecological Dynamics Across the Arctic Associated with Recent Climate Change.
*Science,* **325,** 1355-1358.

Previdi, M., Smith, K.L. & Polvani, L.M. (2021) Arctic amplification of climate change: a
review of underlying mechanisms. *Environmental Research Letters,* **16,** 093003.

R Development Core Team (2022) R: A language and environment for statistical computing. R
Foundation for Statistical Computing, Vienna, Austria.

Richter, S. & Bick, A. (2018) ROSTOCK: The Zoological Collection of the University of
Rostock. *Zoological Collections of Germany: The Animal Kingdom in its Amazing Plenty
at Museums and Universities* (ed. L.A. Beck), pp. 583-589. Springer International
Publishing, Cham.

Schlaich, A.E., Klaassen, R.H.G., Bouten, W., Bretagnolle, V., Koks, B.J., Villers, A. & Both, C.
(2016) How individual Montagu's Harriers cope with Moreau's Paradox during the
Sahelian winter. *Journal of Animal Ecology,* **85,** 1491-1501.

Schmidt, N.M., Ims, R.A., Hoye, T.T., Gilg, O., Hansen, L.H., Hansen, J., Lund, M., Fuglei, E.,
Forchhammer, M.C. & Sittler, B. (2012) Response of an arctic predator guild to
collapsing lemming cycles. *Proceedings of the Royal Society B-Biological Sciences,* **279,**
4417-4422.

Serrano, D., Tella, J.L., Forero, M.G. & Donazar, J.A. (2001) Factors affecting breeding
dispersal in the facultatively colonial lesser kestrel: individual experience vs. conspecific
cues. *Journal of Animal Ecology,* **70,** 568-578.

Sonerud, G.A. (1986) Effect of snow cover on seasonal changes in diet, habitat, and regional
distribution of raptors that prey on small mammals in boreal zones of Fennoscandia.
*Ecography,* **9,** 33-47.
Sumasgutner, P., Buij, R., McClure, C.J.W., Shaw, P., Dykstra, C.R., Kumar, N. & Rutz, C.
(2021) Raptor research during the COVID-19 pandemic provides invaluable
opportunities for conservation biology. *Biological Conservation,* **260,** 109149.
Sundell, J., Huitu, O., Henttonen, H., Kaikusalo, A., Korpimaki, E., Pietiainen, H., Saurola, P. &
Hanski, I. (2004) Large-scale spatial dynamics of vole populations in Finland revealed by
the breeding success of vole-eating avian predators. *Journal of Animal Ecology,* **73,** 167-
178.
Tast, J., Kaikusalo, A. & Lagerström, M. (2010) Diet composition of breeding rough-legged
buzzards Buteo lagopus at Kilpisjärvi, NW Finnish Lapland. *Kilpisjärvi Notes,* **22,** 10-18.
Terraube, J., Villers, A., Ruffino, L., Iso-Iivari, L., Henttonen, H., Oksanen, T. & Korpimäki, E.
(2015) Coping with fast climate change in northern ecosystems: mechanisms underlying
the population-level response of a specialist avian predator. *Ecography,* **38,** 690-699.
Thorup, K., Tøttrup, A.P., Willemoes, M., Klaassen, R.H.G., Strandberg, R., Vega, M.L., Dasari,
H.P., Araújo, M.B., Wikelski, M. & Rahbek, C. (2017) Resource tracking within and
across continents in long-distance bird migrants. *Science advances,* **3,** e1601360.
Tucker, M.A., Böhning-Gaese, K., Fagan, W.F., Fryxell, J.M., Van Moorter, B., Alberts, S.C.,
Ali, A.H., Allen, A.M., Attias, N., Avgar, T., Bartlam-Brooks, H., Bayarbaatar, B.,
Belant, J.L., Bertassoni, A., Beyer, D., Bidner, L., van Beest, F.M., Blake, S., Blaum, N.,
Bracis, C., Brown, D., de Bruyn, P.J.N., Cagnacci, F., Calabrese, J.M., Camilo-Alves, C.,
Chamaillé-Jammes, S., Chiaradia, A., Davidson, S.C., Dennis, T., DeStefano, S.,
Diefenbach, D., Douglas-Hamilton, I., Fennessy, J., Fichtel, C., Fiedler, W., Fischer, C.,
Fischhoff, I., Fleming, C.H., Ford, A.T., Fritz, S.A., Gehr, B., Goheen, J.R., Gurarie, E.,
Hebblewhite, M., Heurich, M., Hewison, A.J.M., Hof, C., Hurme, E., Isbell, L.A.,
Janssen, R., Jeltsch, F., Kaczensky, P., Kane, A., Kappeler, P.M., Kauffman, M., Kays,
R., Kimuyu, D., Koch, F., Kranstauber, B., LaPoint, S., Leimgruber, P., Linnell, J.D.C.,
López-López, P., Markham, A.C., Mattisson, J., Medici, E.P., Mellone, U., Merrill, E., de
Miranda Mourão, G., Morato, R.G., Morellet, N., Morrison, T.A., Díaz-Muñoz, S.L.,
Mysterud, A., Nandintsetseg, D., Nathan, R., Niamir, A., Odden, J., O'Hara, R.B.,
Oliveira-Santos, L.G.R., Olson, K.A., Patterson, B.D., Cunha de Paula, R., Pedrotti, L.,
Reineking, B., Rimmler, M., Rogers, T.L., Rolandsen, C.M., Rosenberry, C.S.,
Rubenstein, D.I., Safi, K., Saïd, S., Sapir, N., Sawyer, H., Schmidt, N.M., Selva, N.,
Sergiel, A., Shiilegdamba, E., Silva, J.P., Singh, N., Solberg, E.J., Spiegel, O., Strand, O.,
Sundaresan, S., Ullmann, W., Voigt, U., Wall, J., Wattles, D., Wikelski, M., Wilmers,
C.C., Wilson, J.W., Wittemyer, G., Zięba, F., Zwijacz-Kozica, T. & Mueller, T. (2018)
Moving in the Anthropocene: Global reductions in terrestrial mammalian movements.
*Science,* **359,** 466-469.
Vansteelant, W., Faveyts, W. & Buckens, J. (2011) Opmerkelijke Ruigpootbuizerdinvasie in de
winter 2010-2011: oorzaken in een historische en Europese context. *Natuur.oriolus,* **77,**
87-98.
Verhoeven, M.A., Loonstra, A.H.J., McBride, A.D., Both, C., Senner, N.R. & Piersma, T. (2021)
Migration route, stopping sites, and non-breeding destinations of adult Black-tailed
Godwits breeding in southwest Fryslân, The Netherlands. *Journal of Ornithology,* **162,**
61-76.
Walker, D.A., Raynolds, M.K., Daniels, F.J.A., Einarsson, E., Elvebakk, A., Gould, W.A.,
Katenin, A.E., Kholod, S.S., Markon, C.J., Melnikov, E.S., Moskalenko, N.G., Talbot,
S.S., Yurtsev, B.A. & Team, C. (2005) The Circumpolar Arctic vegetation map. *Journal
of Vegetation Science,* **16,** 267-282.

654 Whitfield, D.P., Duffy, K., McLeod, D.R., Evans, R.J., MacLennan, A.M., Reid, R., Sexton, D.,
655   Wilson, J.D. & Douse, A. (2009) Juvenile dispersal of white-tailed eagles in western
656   Scotland. *Journal of Raptor Research,* **43,** 110-120.
657 Wikelski, M., Kays, R.W., Kasdin, N.J., Thorup, K., Smith, J.A. & Swenson, G.W. (2007)
658   Going wild: what a global small-animal tracking system could do for experimental
659   biologists. *Journal of Experimental Biology,* **210,** 181-186.
660 Zwarts, L., Bijlsma, R.G., van der Kamp, J. & Wymenga, E. (2012) *Living on the Edge:*
661   *Wetlands and Birds in a Changing Sahel*. KNNV Publishing, Leiden, The Netherlands.

**a)**

Slow phase part 1

OCT
NOV
DEC
JAN

Slow phase part 2

APR
MAR
FEB

**b)**

Snow cover, %

hypothetical snow cover if birds winter in the place where they have arrived from the North

real snow cover that birds experience during the winter

hypothetical snow cover if birds fly directly to the Southwest and stay there all winter

Oct Nov Dec Jan Feb Mar Apr

**a)**

For species with **foxtrot migration**, when mid-winter counts are used to determine winter range (Christmas counts), temporal heterogeneity can be mistaken with spatial heterogeneity

For species with **foxtrot migration**, when GPS tracking is used to determine winter range, temporal heterogeneity becomes evident and should be presented as a **dynamic range**

uncommon

common

Numbers indicate months the birds are in the area

**b)**

**abundance decline** within a specific part of the winter range

Species abundance

Years

Extinct   Threatened   Least Concern

EX  EW  CR  EN  VU  NT  LC

Years

For species with **foxtrot migration**, a decline in abundance within a specific part of the winter range may indicate changes in a range **dynamics** rather than an overall population decrease