Quick-quick-slow: the foxtrot migration and dynamic non-breeding range of the Arctic raptor

Ivan Pokrovsky[1] *, Teja Curk[2], Andreas Dietz[3], Ivan Fufachev[4], Olga Kulikova[5], Sebastian Rößler[3], Martin Wikelski[1,6]

[1] *Department of Migration, Max Planck Institute of Animal Behavior, Radolfzell, Germany*

[2] *Leibniz Institute for Zoo and Wildlife Research, Berlin, Germany*

[3] *Deutsches Zentrum für Luft- und Raumfahrt e.V. (DLR), Wessling, Germany*

[4] *Institute of Plant and Animal Ecology, Yekaterinburg, Russia*

[5] *Institute of Biological Problems of the North, Magadan, Russia*

[6] *Centre for the Advanced Study of Collective Behaviour, University of Konstanz, 78457 Konstanz, Germany*

* *Corresponding author*

E-mail: ipokrovsky@ab.mpg.de

ABSTRACT

Advances in tracking technologies have revealed the diverse migration patterns of birds, which are critical for range mapping and population estimation. While population trends are typically estimated in breeding ranges where birds remain stationary, trends are commonly assessed in non-breeding ranges for species that breed in inaccessible regions such as the Arctic. This task is challenging due to the extensive movements of birds in these ranges, which require a deep understanding of the dynamics of these movements. However, these movements remain understudied, particularly in the mid-latitudes, where many Arctic breeders winter, increasing the risk of misestimating their ranges and numbers. Here, we show that the Arctic breeding raptor Rough-legged buzzard, which overwinters in the mid-latitudes, has a specific wintering strategy: during the entire non-breeding season, they move about 1000 km back and forth under the influence of the advancing snowline, making their non-breeding area dynamic. In essence, this movement represents an extension of the quick migration process, albeit at a slower pace, and we have termed this migration pattern 'foxtrot migration,' drawing an analogy to the alternating fast and slow movements of the foxtrot dance. These results highlight the potential errors in range mapping from single mid-winter surveys and emphasize the importance of this migration pattern in assessing the conservation status of bird species. Understanding this migration pattern helps to correctly estimate bird populations in non-breeding areas, which is especially important for species that nest in hard-to-reach regions such as the Arctic.

Keywords: *migration patterns, bird movement ecology, bird migration, types of migration, conservation, species ranges, snow cover, dynamic range, foxtrot migration.*

INTRODUCTION

In recent years, advances in tracking technology have greatly improved our understanding of bird migration (Wikelski et al. 2007, Kays et al. 2015, Jetz et al. 2022, Kays and Wikelski 2023). By tracking tagged birds, we can study various migratory behaviors, such as their propensity to migrate, connectivity and philopatry, length of migration routes, variations in flight speed, and movements within a single season (Alerstam et al. 2003, Newton 2008, Alerstam 2011, Berthold et al. 2013, Trierweiler et al. 2013, Schlaich et al. 2023). Different combinations of these behaviors contribute to various bird migration patterns (Chapman et al. 2014, Alerstam and Bäckman 2018, Lislevand et al. 2020). Understanding the migratory patterns of birds is critical to accurately mapping their ranges and estimating population trends. Typically, population trends are assessed in breeding areas, where birds are linked to their nesting sites, allowing reliable estimates of breeding numbers (Reif 2013). However, access to some breeding areas, such as the Arctic, presents logistical and political challenges (Gallo-Cajiao et al. 2023, Koivurova and Shibata 2023). As a result, population trends for these species are often estimated in non-breeding areas where birds aren't tied to specific locations (Robinson et al. 2005, JIa et al. 2016). This challenges accurately censusing birds and delineating their ranges, especially when birds move extensively in non-breeding ranges during the season, and censuses are conducted only once per non-breeding season. Therefore, understanding bird movements in non-breeding areas is essential for understanding the ecology of birds in these regions and has practical implications for accurately assessing their ranges and trends in their populations.

Many bird species experience significant seasonal changes in food availability within their non-breeding range due to environmental conditions. This often triggers movements across these areas. For Palearctic migrants wintering in Africa, these movements have been studied extensively (Moreau 1972, Trierweiler et al. 2013, Thorup et al. 2017, Schlaich et al. 2023). However, the non-breeding movements of mid-latitude birds remain poorly understood. At the same time, many Arctic birds winter in the mid-latitudes, and understanding their movement patterns could be crucial for interpreting their population trends. In the mid-latitudes, food availability is strongly influenced by environmental factors such as snow cover dynamics. The progressive movement of the snow cover line from north (-east) to south (-west) and back again between October and May in many mid-latitude regions, particularly those exposed to northwesterly wind systems, significantly affects food availability for numerous bird species (Sonerud 1986, Vansteelant et al. 2011). We propose that species dependent on this environmental factor gradually move away from snow-covered areas during winter and then

gradually move back in the opposite direction, resulting in a directed and continuous

displacement of their non-breeding range.

The Rough-legged buzzard (Buteo lagopus) is an Arctic breeding and mid-latitude

wintering raptor (Ferguson-Lees and Christie 2001, Bechard and Swem 2002). Rough-legged

buzzards feed mainly on small rodents in the Arctic and its wintering grounds (Tast et al. 2010,

Pokrovsky et al. 2014). They prefer open areas for hunting, and trees and tall bushes or uplands

for resting. In the Arctic, such areas are the southern and typical tundra (Walker et al. 2005), and

in the mid-latitudes, areas with fields and patches of forests (wooded fields). The taiga zone,

where there is little open space, is unsuitable for them, although they may nest in the northern

taiga zone on the border with the tundra zone (Sundell et al. 2004). Snow cover and day length

are important in their life (Terraube et al. 2015, Curk et al. 2020, Pokrovsky et al. 2021). Rough-

legged buzzards can only hunt during the day (Pokrovsky et al. 2021), and heavy snow cover

makes hunting for small rodents problematic (Sonerud 1986, Vansteelant et al. 2011). These two

environmental factors, therefore, affect the availability of prey for Rough-legged buzzards. At

the same time, these factors vary considerably in mid-latitudes during winter. Thus, prey

availability for Rough-legged buzzards in the mid-latitudes increases until mid-winter if they

migrate southwards and after mid-winter if they migrate northwards. We, therefore, assume that

Rough-legged buzzards could track prey availability and experience a dynamic non-breeding

range.

In this study, we tracked GPS-tagged Rough-legged buzzards for ten years and analyzed

their movements in the non-breeding mid-latitude range in relation to snow cover. For this study,

we made the following predictions. 1) Rough-legged buzzards would exhibit a directional and

seasonal movement pattern during the non-breeding period, moving from the northeast to the

southwest and back again. This would result in a dynamic non-breeding range that would

continue to move geographically throughout the season. 2) The non-breeding movements would

differ from the fall and spring migrations in duration, extent, speed, and direction but would

continue throughout the season. Thus, over the entire annual cycle, Rough-legged buzzards

would be characterized by a migratory pattern consisting of an alternation of quick (fall and

spring migrations) and slow (non-breeding movements) phases. 3) Non-breeding movements

would occur in suitable open habitats, while fall and spring migrations would occur in unsuitable

forested areas. Thus, vegetation land cover will determine the extent of quick and slow migration

phases. 4) During non-breeding movements, Rough-legged buzzards will experience less snow

cover than if they had stayed where they arrived at the end of the fall migration. Thus, snow

cover dynamics will determine the dynamics of non-breeding movements.

MATERIAL AND METHODS

Dataset

For this study, we tracked 43 adult Rough-legged buzzards (35 females and eight males) with the

solar GPS-GSM loggers (e-obs GmbH and UKn – University of Konstanz). Twenty-eight birds

were fitted with E-obs loggers, 13 birds were fitted with UKn loggers, and two birds were

initially fitted with UKn loggers that were later replaced with E-obs loggers. E-obs loggers

weighed 45 g, or 3.6% of the bird's weight, while UKn loggers weighed 15 g, or 1.2% of the

bird's weight. E-obs loggers recorded GPS positions every hour in full battery mode and every 5

hours in normal battery mode. UKn loggers recorded GPS positions every hour with a full

battery and every 12 hours with a normal battery. Both models operated continuously 24 hours a

day. More detailed information about this dataset can be found in Curk et al. (2022).

The fieldwork was carried out in the Russian Arctic in 2013-2019 at four study sites:

Kolguev Island (69°16′N, 48°87′E), Nenetsky Nature Reserve (68°20′N, 53°18′E), Vaigach

Island (69°43′N, 60°08′E), and Yamal Peninsula (68°12′N, 68°59′E). For details on capture

methods, see Curk et al. (2022); for detailed study descriptions, see Pokrovsky et al. (2015) for

Kolguev and Pokrovsky et al. (2019) for Yamal and Nenetsky.

During data pre-processing, we estimated the date of death using an accelerometer and

GPS data and excluded tracking data corresponding to the period after the bird's death. We then

removed duplicated timestamps and calculated the mean daily positions of each individual. We

partitioned the resulting dataset into several periods: 1) breeding, 2) fall migration, 3)1^{st} part of

winter, 4) 2^{nd} part of winter, and 5) spring migration. We estimated the migration dates – the

start and stop dates of the spring and fall migrations – using an iterative search procedure for

piecewise regression described by Crawley (2007). We created two models for each of the four

dates: a simple model and a piecewise one. The simple model was linear, with latitude as the

response variable and day of the year as a fixed effect. The piecewise model was also linear, with

latitude as the response variable and day of the year as a fixed effect, and included two logical

statements in its formula because we expected two linear segments in the fit. Specifically, it used

"Day Of the Year < BREAK" to define the left regression (before the estimated date) and "Day

Of the Year ≥ BREAK" for the right regression (after the estimated date). We explored different

BREAK values to fit the piecewise model, looking for the value that minimized the residual

standard error. The BREAK values examined included those associated with the start and end of
the fall migration (September 5 to October 5 and October 5 to November 5, respectively), as well
as those associated with the start and end of the spring migration (April 10 to May 10 and May
10 to June 10, respectively). We then validated (using the R function 'anova') that the piecewise
model significantly improved the fit compared to the simple model ($p<0.001$). We estimated the
date between winter's first and second parts as the day when the mean daily latitude was
minimum.

Data analysis

First, we used linear mixed-effects models (R function 'lmer' in the library 'lme4' (Bates et al.
2015)) to investigate whether or not Rough-legged buzzards migrated during winter. Latitude
was the response variable, day of the year was a fixed effect, and individuals and year were
included as random effects. Analyses were conducted separately for each migration period (fall,
first phase of winter, second phase of winter, and spring). For both phases of the winter
migration, we analyzed two additional models with longitude as the response variable instead of
latitude. Likelihood ratio tests were used to compare candidate models. The year was not a
calendar year but a year between two consecutive breeding seasons. Thus, fall migration,
consecutive winter, and consecutive spring have the same value for the year. The day of the year
was recalculated consecutively.

Second, we used linear mixed-effects models (R function 'lmer' in the library 'lme4') to
investigate whether migrations' parameters differ between the migration periods. We analyzed
four migration parameters: distance, duration, speed, and direction. The distance was calculated
as the distance between two coordinates (start and end of migration) using the R function 'distm'
in the library 'geosphere' (Hijmans 2016). The duration was calculated as the number of days
between the start and end of migration. Speed was calculated as the ratio of distance to duration.
The direction was calculated as the bearing from the start of the migration coordinates to the end
of the migration coordinates using the R function 'bearing' in the library 'geosphere' (Hijmans
2016). The migration parameter was used as the response variable, the type of migration as a
fixed factor, and individuals as a random factor. Likelihood ratio tests were used to compare
candidate models. We considered four different parameters of migration (distance, duration,
speed, and direction) and four types of migration (fall, first phase of winter, second phase of
winter, and spring). The analysis was done separately for each of the migration parameters.
Then, we used post hoc comparisons using the R function 'emmeans' in the library 'emmeans'
(Lenth et al. 2019) to compare the estimated means.

In some raptor species, adult females disperse further than males (Mearns and Newton

1984, Serrano et al. 2001, Bildstein 2006, Whitfield et al. 2009). Therefore, we conducted an

additional analysis on the effect of sex on migration length using linear mixed-effects models (R

function 'lmer' in the library 'lme4'). The migration distance was used as the response variable,

sex as a fixed factor, and individuals as a random factor.

Third, we investigated whether vegetation land cover differed between areas crossed

during the quick (fall and spring) and slow (winter) migrations. We used the combined Terra and

Aqua Moderate Resolution Imaging Spectroradiometer (MODIS) Land Cover Climate

Modelling Grid (CMG) (MCD12C1) version 6 dataset (Friedl and Sulla-Menashe 2015). We

used a modified Leaf Area Index (LAI) as a classification scheme. We combined all four forest

types and savannas into one category (forest) and excluded the categories: water bodies, and

unclassified. We, therefore, had five types of vegetation cover: forest, grassland, cropland,

shrubland, and urban. We annotated the mean daily positions with the vegetation cover type

using the Env-DATA tool (Dodge et al. 2013). We used general linear mixed effects models with

a binomial distribution (R function 'glmer' in the library 'lme4') to investigate whether

vegetation cover types differ between migration periods. Presence/absence of the studied

vegetation land cover type was used as a response variable, migration type as a fixed factor, and

individuals as a random factor. The analysis was done separately for each of the vegetation land

cover types.

Fourth, we investigated whether snow cover could drive the slow migration phenomenon.

We then compared the snow cover conditions the birds experienced during the winter with two

hypothetical snow cover conditions that the birds would have experienced if they had not

migrated during the winter. The first hypothetical snow cover condition would have happened if

the birds had stayed where they arrived from the north (i.e., where their fall migration ended). To

estimate this parameter, we calculated the winter dynamics of the average snow cover at the

minimum convex polygons (MCP) occupied by the birds in October and April (northeast of their

winter range). A second hypothetical snow cover condition would be if the birds flew

immediately to the southwest and spent the whole winter there. To evaluate this, we calculated

the winter dynamics of average snow cover on the MCPs occupied by the birds in January and

February (southwest of their winter range). We then compared the values obtained for the real

snow cover and two hypothetical snow covers using general linear mixed effects models with a

binomial distribution (R function 'glmer' in the 'lme4' library). Presence/absence of snow cover

was used as a response variable, type of snow cover (real, 1[st] hypothetical, or 2[nd] hypothetical) as

a fixed factor, and years as a random factor. The analysis was done separately for each month.

We then used post hoc comparisons to compare the estimated means, using the R function

'emmeans' in the 'emmeans' library (Lenth et al. 2019).

We obtained monthly snow cover data with a spatial resolution of ca 500 meters (Global

SnowPack MODIS) from the German Aerospace Center (DLR). This product is based on the

Moderate Resolution Imaging Spectroradiometer (MODIS) daily snow cover products

MOD10A1 and MYD10A1 (version 6 as provided by the National Snow and Ice Data Center

NSIDC), which have been processed to remove the gaps due to cloud cover and polar darkness

(Dietz et al. 2015). These processing steps include a combination of data available from different

satellites (Aqua and Terra), 3-day temporal moving window filtering, a regional snow line

elevation interpolation relying on a Digital Elevation Model (DEM), and a seasonal filter

running through the time series for the whole hydrological year (1st of September through

August 31st). The proportion of days in which one pixel is snow-covered per month is referred to

here as fractional snow cover and is derived from these daily gap-filled rasters. Five MODIS

tiles (h19v03, h20v03, h20v04, h21v03 and h21v04) were mosaicked and re-projected to

WGS84. Then, for each month from October to April, we calculated 95% minimum convex

polygons (MCPs) for the distribution of Rough-legged buzzards using the R function 'mcp' in

library 'adehabitatHR' (Calenge 2006). We extracted mean snow cover values from each MCP

from every monthly snow cover raster separately, using the R library 'raster' (Hijmans and van

Etten 2023).

All calculations were performed using R version 4.2.2 'Innocent and Trusting' (R

Development Core Team 2022) and RStudio version 353 'Elsbeth Geranium' (Posit team 2022).

RESULTS

Migration of Rough-legged buzzards – always on the move

Except during the breeding season, Rough-legged buzzard migration continues throughout the

year, even after the birds´ arrival at their traditionally recognized ´wintering grounds´ (Figure 1a,

b). Rough-legged buzzards started their fall migration on 28 September (hereafter mean±sd for the

day of the year: 271±11, n=31) and ended on 12 October (285±11, n=33). The mean

latitude/longitude where the birds ended their fall migration was $55.57\pm1.92°/49.35\pm5.63°$ (Figure

1a). During the winter, birds continued to migrate at a slower pace down to $49.53\pm2.01°$ latitude

(on 5 February, 36±40, n=23) and $34.29\pm5.11°$ longitude (on 24 January, 24±47, n=23).

Afterward, during the second part of the winter, birds returned to $55.52\pm2.63°$ latitude and

49.79±8.24° longitude to start the spring migration (Figure 1a). Rough-legged buzzards started
their spring migration to the Arctic on 27 April (117±7, n=27) and arrived at the breeding grounds
on 15 May (135±8, n=18).

In the following, we will refer to the spring and fall migrations as the quick phase, the non-
breeding movement to the lowest point of latitude as the 1^{st} part of the slow phase, and the
movement from the lowest point of latitude to the starting point of the spring migration as the 2^{nd}
part of the slow phase (Figure 1a). For both quick and slow phases of the migration, linear mixed-
effects models with the season as a fixed factor received higher support from the likelihood ratio
test (p < 0.001, Tables S1-S3).

Figure 1. Migration of Rough-legged buzzards. Q – quick phase. Qf – Quick fall phase
(orange), Qs – Quick spring phase (yellow), S1 – Slow phase, 1^{st} part (light blue), S2 – Slow
phase, 2^{nd} part (dark blue). a) Change in the latitude of 43 Rough-legged buzzards during the
year, red line – mean latitude of all birds, black vertical lines – mean dates of start and end of the
migration phases, blue vertical line – mean date of the minimum latitude. Grey, sky blue, and

piggy pink shaded areas – standard deviation of the means. b) Migration map. c) Difference in
the migration parameters between the migration phases. Lines on the direction plot (down, right)
represent the mean value for each bird; arrows represent the mean direction for each phase.
Boxes on the boxplots show the interquartile range, the whiskers – maximum and minimum
values.

Quick-slow phase features comparison

During the quick phase, individual birds flew greater distances in a shorter time, i.e., at a
faster rate, than during the slow phase. After arriving at what is traditionally known as the
wintering grounds, the direction of migration changed, so the direction of quick and slow phases
also differed (Table 1, Figure 1c). The quick phase was 1415±50 km long (hereafter mean±sd),
whereas the slow phase (one part) was 1026±55 km, i.e., 389±60 km shorter (p<0.001, Table S4,
Figure 1c). During the quick phase, birds flew for 15±3 days, and one part of the slow phase
lasted 100±4 days, i.e., 85±5 days longer (p<0.001, Table S5, Figure 1c). At the same time, the
second part of the slow phase was 54±7 days shorter than the first (p<0.001, Table S5, Figure
1c). The migration speed was 104±6 km/day during the quick phase and 12±7 km/day during the
slow phase, i.e., about eight times higher (p<0.001, Table S6, Figure 1c). During the fall
migration, birds moved in the SSW direction (7±2 deg), then turned 50±3 deg (p<0.001, Table
S7, Figure 1c) to the west and started their 1st slow phase until mid-winter. After that, they
turned back to the NEE direction (57±2 deg) and performed their 2nd slow phase for several
months until they turned 54±3 deg (p<0.001, Table S7, Figure 1c) to the north and started their
spring migration. As a result of additional analysis of the effect of sex on migration length, we
found no significant difference between the migration distances of males and females (Table S8).

Table 1. Parameters of the Rough-legged buzzards' migration (mean±sd).

Phase of migration	Sub-phase of migration	Distance (km)	Duration (days)	Speed (km/day)	Direction (deg)
Quick		1415±50	15±3	104±6	
	Spring	1544±72	18±4	100±8	7±2
	Fall	1334±59	14±4	107±7	198±3
Slow		1026±55	100±4	12±7	
	1st phase	1016±71	121±4	8±7	251±3
	2nd phase	1042±71	66±6	17±9	57±2

Vegetation land cover during migration

During quick phases, Rough-legged buzzards cross the forest zone, while during the slow phase, they migrate within the grassland and cropland zone (Figure 2). Rough-legged buzzards migrated fast across the tundra zone on the north in the Arctic and then through the taiga zone. Therefore, during the quick phase, the three most common vegetation land cover types were forest (44.5±2.9 %, hereafter, percentage of all mean daily positions annotated with the given vegetation type ± sd), shrublands (29.9±3 %), and grasslands (24.7±2.8 %, Figure 2a). During the slow phase, the three most common vegetation land cover types were grasslands (65.1±6.2 %), croplands (26.9±6.3 %), and forests (4.9±1.4 %, Figure 2b). According to the linear mixed-effects models, the percentage of all vegetation land cover types differed between the slow and quick phases (p<0.001, Table S9), except for the urban lands. Urban lands were more common during the slow than quick phase (Figure 2). However, this type has been annotated for too few birds to make an adequate comparison.

Figure 2. Vegetation land cover during quick and slow phases of the migration. a) Quick phase (spring and fall periods together). b) Slow phase (1st and 2nd parts together). On both a) and b), the bars show the percentage of all mean daily positions annotated with the given vegetation type ± sd. c) Migration map.

Snow cover – the main reason for the dynamic winter range

During the slow phase of the migration, Rough-legged buzzards experienced snow cover ranging from 4.8±1.0% in October (hereafter mean±sd) to 85.2±4.6% in February (Figure 3). If birds spent the winter in the place where they arrived after the fall migration, they would experience snow cover conditions ranging from 4.6±0.6% in October to 99.5±0.1% in February (Figure 3b, green line). If birds fly directly to the southeast and stay there for the whole winter, they would experience snow cover conditions ranging from 1.4±0.2 % in October to 81.1±5.0 % in January (Figure 3b, red line). Thus, if birds fly immediately to the southwest and stay there until the end of the winter, they will find conditions with less snow cover in spring (p<0.001, Table S10). And if birds stay where they ended the fall migration, they will find themselves in situations with more snow cover (p<0.001, Table S10). In the latter case, the difference between real and hypothetical situations is not as pronounced (85.2% vs. 99.5%), but more means that snow cover will be close to 100% for several months in this hypothetical situation (Figure 3b).

Figure 4. Snow cover conditions during the slow phase of the migration. a) 95% Minimum
convex polygons (MCPs) of Rough-legged buzzards during winter. Arrows indicate the direction
of the movement across months. OCT – October, NOV – November, DEC – December, JAN –
January, FEB – February, MAR – March, APR – April. b) Snow cover conditions for the real
situation (black) and two hypothetical situations – if birds spend the winter in the place where
they arrived after the fall migration (green) and if birds fly directly to the southwest and stay
there all winter (red). Dots represent mean values, error lines – standard deviations.

DISCUSSION

Our study identified and characterized a bird migration pattern consisting of an alternation of
quick (fall and spring migrations) and slow (non-breeding movements) phases. This migration
pattern causes the non-breeding range of birds to shift and become dynamic throughout the
season. This has obvious implications for range delineation and assessment of bird population
trends.

Non-breeding movements and migratory patterns

Different bird species employ various wintering strategies, leading to diverse migration patterns.
One strategy is residency, where birds arriving at non-breeding grounds remain there throughout
the season, undertaking only minor foraging flights (Figure 4a). Their home ranges in non-
breeding areas are compatible with those in breeding areas (Kjellén et al. 1997, Alerstam et al.
2006, Newton 2008). Another strategy, initially identified by Moreau (Moreau 1972), is
itinerancy, wherein birds fly between several sites separated by distances ranging from ten to
several hundred kilometers throughout the non-breeding period, spending 1-2 months at each site
(Figure 4b). Most Palearctic-African species adopt this itinerant behavior (Trierweiler et al.
2013, Thorup et al. 2017, Schlaich et al. 2023). The third strategy investigated here involves
birds moving slowly and steadily throughout the non-breeding season, initially in one direction
and then in the opposite direction. In 2009, Strandberg et al. used GPS transmitters to track
Common buzzards *(Buteo buteo)*. Despite some transmitters in their study ceasing to function
mid-winter, the authors observed a phenomenon they termed "prolonged autumn migration,"
wherein birds arriving at their wintering grounds did not halt migration but continued moving at
a slower pace (Strandberg et al. 2009). In our study, we observed this phenomenon throughout
the annual cycle. We demonstrated that, unlike itinerancy, movements in the non-breeding area
are essentially a continuation of migration at a slower pace, influenced by external factors such
as snow cover dynamics. Considering this behavior as an extension of migration and examining
the complete life cycle of such species, it involves a quick phase during the transition between
breeding and non-breeding areas, followed by a slow phase of directed and seasonal movement

347   within the non-breeding area, and finally, another quick phase towards the breeding area.

348   Drawing an analogy to the alternating quick and slow movements of the foxtrot dance, we

349   propose the term 'foxtrot migration' for this seasonal movement to offer a concise and easily

350   understandable description (Figure 4c). Consequently, we suggest referring to the non-breeding

351   range of species exhibiting itinerancy or foxtrot migration as the 'dynamic range' (Figure 4b, c).

Figure 4. Winter strategies and non-breeding range scheme. a) Residency, b) Itinerancy, c)
Foxtrot migration. The color bar along the y-axis corresponds to the color coding of the habitats
of the chart on the right.

Debates may arise regarding introducing a new term for this phenomenon and whether it
should fall under the general term "nomadism" or be categorized as a modified form of another
migratory pattern, such as "slow directional itinerancy." We assert that behavioral pattern names
should aid in understanding movement strategies and avoid confusion. One might ask whether it
is necessary to distinguish between migration and wintering at all, given that migration is
essentially a matter of tracking spatial and temporal changes in favorable conditions. Authors
studying Montagu's harriers *(Circus pygargus)* argue that such generalization hampers
understanding of wintering strategies (Schlaich et al. 2023). Instead, Schleich et al. (2023)
propose explicitly considering the functions of animal-used sites, particularly their contributions
to "wintering" and "refueling for migration." For instance, cuckoos *(Canorus sp.)* utilize
stopovers for both peak foraging conditions and refueling for long flights to subsequent sites
(Thorup et al. 2017). Conversely, Montagu's harriers exhibit negligible refueling between sites
due to short distances and energy-efficient soaring flight (Trierweiler et al. 2013, Schlaich et al.
2023). In our case, Rough-legged buzzards do not make several long stopovers, unlike species
experiencing itinerancy, but move gradually in one direction and then return during the whole
season (Figure 1a). As Schleich et al. (2023) suggested, examining the functions of animal-used
sites reveals another difference between these strategies. The difference is that itinerant birds
wait for better conditions at subsequent sites. In contrast, in our case, the southeastern areas that
birds reach after several months are suitable and offer better conditions than the northeastern
areas, as they have less snow throughout the whole non-breeding season (Figure 3). All these
differences highlight the discrepancy between the wintering strategies referred to as itinerancy
and those described in this study. Therefore, using another term - foxtrot migration - for this
phenomenon is more appropriate.

The dynamic of the foxtrot migration

Our study affirmed the presence of foxtrot migration and a dynamic non-breeding range in
Rough-legged buzzards. The taiga zone posed as an unfavorable habitat during the quick phase
of foxtrot migration (Figure 2), given the difficulty of Rough-legged buzzards locating open
areas for hunting in this habitat. Conversely, the grassland and cropland zone served as favorable
habitats during the slow phase throughout the entire non-breeding period (Figure 2), offering
numerous open areas for hunting. Snow cover was the external factor driving their continual

1000 km southwest movement during winter (Figure 3). Our analysis revealed that if Rough-

legged buzzards remained at their fall migration endpoint without moving southwest, they would

encounter 14.4% more snow cover (99.5% vs. 85.1%, Figure 3). Although this difference may

seem small (14.4%), it holds significance for rodent-hunting birds, distinguishing between

complete and patchy snow cover. Simultaneously, if Rough-legged buzzards immediately flew to

the southwest and stayed there throughout winter, they would experience 25.7% less snow cover

(57.3% vs. 31.6%, Figure 3b). Despite a greater difference than in the first case, it doesn't

compel them to adopt this strategy, as it represents the difference between various degrees of

landscape openness from snow cover. However, it is noteworthy that one bird followed this

strategy by flying directly to the southwest and remaining there throughout the winter (Figure

S1), exhibiting a residency strategy. This observation suggests that the distinction between

winter strategies is flexible.

African-Palearctic migrants do not strictly adhere to a binary division between residency

and itinerancy but demonstrate a continuum between these two strategies. Many species in

Africa exhibit a mixture of both strategies in varying proportions; see Schlaich et al. (2023) and

reference therein. Similarly, in the mid-latitudes, species may not only show any of the three

strategies but also a considerable degree of individual and regional variation. For example,

Rough-legged buzzards from western Arctic regions that overwinter in mid-latitude areas where

snow cover dynamics are less pronounced than in eastern regions are likely to have a higher

proportion of birds with a resident strategy. At the same time, Rough-legged buzzards in North

America also exhibit a foxtrot migration pattern, although with variations. There, birds make a

direct and rapid migration, usually across the boreal forest, followed by slower, more facultative

movements once birds get south of the boreal. However, there is a lot of variation among

individuals and even regions; there are more non-breeding slow movements in eastern North

America but less in western North America (Neil Paprocki pers. comm.). Thus, Rough-legged

buzzards in North America lie along a gradient between foxtrot migration and residency, likely

influenced by different snow cover dynamics on the east and west coasts.

The foxtrot migration pattern is expected to be observed in many migratory species that

exhibit distinct seasonal cycles in their non-breeding range. This migration pattern is expected to

be prevalent for species living in regions where snow cover is a significant determinant of food

availability, which is the case in large parts of the mid-latitudes. Therefore, understanding bird

migration patterns is critical to accurately mapping ranges and assessing population trends in this

region.

Mapping ranges and assessing population trends

The implementation of our study is twofold: 1) the use of mid-winter bird surveys to determine non-breeding range may yield inaccurate results for species with dynamic range (Figure 5a), and 2) declines in abundance within a particular segment of the non-breeding range may indicate changes in range dynamics rather than widespread declines in species abundance (Figure. 5b).

Figure 5. Mapping ranges and assessing population trends for the species with the dynamic non-breeding ranges. a) Mid-winter counts, often used to determine winter range, may give a misleading representation of the non-breeding range for species with dynamic range. b) Changing the conservation status of a species based on reduced abundance in a particular area may not be appropriate for species with dynamic range. Red – breeding range, blue – winter range.

In North America and Europe, the number of non-breeding birds is typically estimated once a season in mid-winter. In North America, these estimates are made during the Christmas counts, usually once in late December (National Audubon Society 2024), while in Europe, they are made during the IWC counts, usually once in January or February (Wetlands International 2024). This approach can lead to errors in range estimation for birds with dynamic ranges. For such species, habitats occupied by birds in fall and spring will be listed as "uncommon" at best,

while habitats occupied by birds only in mid-winter will be listed as "common." However, the
situation is the opposite if we consider the time birds spend in these habitats (Figure 5a). For
example, based on our study of the Rough-legged buzzard, during mid-winter, the species is
predominantly present in the southwestern portion of its non-breeding range, with only a tiny
proportion present in the rest of the region. As a result, a map of the non-breeding range may
show the species as "common" in the southwest and "uncommon" in the northeast. This map
would be inaccurate because, during the entire non-breeding period, Rough-legged buzzards
spend both fall and spring in the northeastern part and only mid-winter in the southwestern part.

To address this, continuous year-round GPS tracking of the species provides a means to
track bird locations throughout the non-breeding season, facilitating the creation of accurate
distribution maps, particularly for species that exhibit dynamic ranges. We advocate representing
temporal heterogeneity (range dynamics) on maps as distinct zones, denoting periods when the
species is abundant in a given area. To distinguish temporal from spatial heterogeneity, we
recommend using lines to delineate the boundaries of these zones, rather than color shading, and
incorporating numbers to denote the months of species abundance in a given zone (Figure 5a).
We suggest ecologists include dynamic non-breeding ranges in descriptions and range maps for
foxtrot migratory and itinerant bird species.

Population counts for a species are often limited to a portion of its range. Therefore,
conclusions about conservation status drawn from such counts may be misleading. A decline in
abundance within a particular portion of the non-breeding range may indicate changes in range
dynamics rather than a general decline in the species (Figure 5b). For example, climate change
may affect snow cover dynamics, reducing its intensity in northern regions. As a result, species
whose range dynamics depend on snow cover may choose to remain in the northern areas and
not migrate as far south as they traditionally have. Despite this shift in range dynamics, overall
species abundance may remain unchanged. Similar patterns have affected Rough-legged
buzzards in some areas of the European non-breeding range.

A 2022 Dutch study found a decline in wintering Rough-legged buzzards over the last 40
years (Hornman et al. 2022). On the one hand, this may represent a conservation concern. On the
other hand, applying the rationale of the dynamic range, the apparent local decline may be
attributed to climate change, resulting in less comprehensive snow coverage in the northeastern
wintering areas of Rough-legged buzzards relative to the Netherlands. Such a shift in snow
coverage makes it less probable for the birds to migrate to the Netherlands for overwintering.
This proposition is further supported by a study of the winter population dynamics of Rough-

legged buzzards in the Netherlands in 2011, showing that the main winter population peak

occurred in late December, with many birds migrating (Vansteelant et al. 2011). Vansteelant et

al. (2011) also found that the main migration occurred after heavy snowfall in northern Europe,

supporting our foxtrot migration explanation for this decline. Therefore, investigating the

dynamic range is critical to understanding a species' range and effectively assessing its

conservation status.

CONCLUSION

Our study sheds light on the overwintering strategy of the Rough-legged buzzard, an Arctic

breeder that migrates to mid-latitudes during the non-breeding season. We have observed a

migration pattern in which these birds travel about 1000 km round trip in response to changing

snowline conditions, which we call the 'foxtrot migration' for its similarity to the alternating

quick and slow movements of the foxtrot dance. Our results highlight the importance of this

migration pattern and the concept of 'dynamic range' in revealing potential inaccuracies in range

mapping from single midwinter surveys and suggest that species distribution maps should

represent temporal variations in range dynamics as separate zones. Understanding and

accounting for this migratory behavior is critical for accurately assessing bird populations in

non-breeding areas. This is especially important for species such as Arctic nesting birds, where

population assessments are primarily conducted in non-breeding ranges due to the logistical and

political challenges of the Arctic.

ACKNOWLEDGMENTS

We are very grateful to everyone who helped us collect data in the field. We would like to thank

Dr Theunis Piersma and an anonymous reviewer for their valuable feedback and constructive

criticism, which significantly improved the quality of this publication. This study was funded by

the Max-Planck Institute of Animal Behavior and the German Air and Space Administration

(DLR). We also acknowledge partial funding by the Deutsche Forschungsgemeinschaft (DFG,

German Research Foundation) under Germany's Excellence Strategy – EXC 2117 – 422037984.

REFERENCES

Alerstam, T. 2011. Optimal bird migration revisited. Journal of Ornithology **152**:5-23.

Alerstam, T., and J. Bäckman. 2018. Ecology of animal migration. Current Biology **28**:R968-
R972.

Alerstam, T., M. Hake, and N. Kjellén. 2006. Temporal and spatial patterns of repeated
migratory journeys by ospreys. Animal Behaviour **71**:555-566.

Alerstam, T., A. Hedenström, and S. Åkesson. 2003. Long-distance migration: evolution and
determinants. Oikos **103**:247-260.
Bates, D., M. Mächler, B. Bolker, and S. Walker. 2015. Fitting Linear Mixed-Effects Models
Using lme4. Journal of Statistical Software **067**.
Bechard, M. J., and T. R. Swem. 2002. Rough-legged Hawk (Buteo lagopus), The Birds of North
America Online (A. Poole, Ed.). Ithaca: Cornell Lab of Ornithology.
Berthold, P., E. Gwinner, and E. Sonnenschein. 2013. Avian migration. Springer Science &
Business Media.
Bildstein, K. L. 2006. Migrating raptors of the world: their ecology & conservation. Cornell
University Press.
Calenge, C. 2006. The package "adehabitat" for the R software: A tool for the analysis of space
and habitat use by animals. Ecological Modelling **197**:516-519.
Chapman, B. B., K. Hulthén, M. Wellenreuther, L.-A. Hansson, J.-Å. Nilsson, and C. Brönmark.
2014. Patterns of animal migration. Pages 11-35 in L.-A. Hansson and Å. Susanne,
editors. Animal movement across scales. Oxford University Press.
Crawley, M. J. 2007. The R Book. Wiley Publishing.
Curk, T., O. Kulikova, I. Fufachev, M. Wikelski, K. Safi, and I. Pokrovsky. 2022. Arctic
Migratory Raptor Selects Nesting Area During the Previous Breeding Season. Frontiers
in Ecology and Evolution **10**.
Curk, T., I. Pokrovsky, N. Lecomte, T. Aarvak, D. F. Brinker, K. Burnham, A. Dietz, A. Dixon,
A. Franke, G. Gauthier, K.-O. Jacobsen, J. Kidd, S. B. Lewis, I. J. Øien, A. Sokolov, V.
Sokolov, R. Solheim, S. Weidensaul, K. Wiebe, M. Wikelski, J.-F. Therrien, and K. Safi.
2020. Arctic avian predators synchronise their spring migration with the northern
progression of snowmelt. Scientific Reports **10**:7220.
Dietz, A. J., C. Kuenzer, and S. Dech. 2015. Global SnowPack: a new set of snow cover
parameters for studying status and dynamics of the planetary snow cover extent. Remote
Sensing Letters **6**:844-853.
Dodge, S., G. Bohrer, R. Weinzierl, S. C. Davidson, R. Kays, D. Douglas, S. Cruz, J. Han, D.
Brandes, and M. Wikelski. 2013. The environmental-data automated track annotation
(Env-DATA) system: linking animal tracks with environmental data. Movement Ecology
1:3.
Ferguson-Lees, J., and D. Christie. 2001. Raptors of the World. Princeton University Press.
Friedl, M., and D. Sulla-Menashe. 2015. MCD12C1 MODIS/Terra+Aqua Land Cover Type
Yearly L3 Global 0.05Deg CMG V006.in N. E. L. P. DAAC, editor.
Gallo-Cajiao, E., N. Dolšak, A. Prakash, T. Mundkur, P. G. Harris, R. B. Mitchell, N. Davidson,
B. Hansen, B. K. Woodworth, R. A. Fuller, M. Price, N. Petkov, V. Mauerhofer, T. H.
Morrison, J. E. M. Watson, S. U. Chowdhury, C. Zöckler, O. Widerberg, D. L. Yong, D.
Klich, V. Smagol, J. Piccolo, and D. Biggs. 2023. Implications of Russia's invasion of
Ukraine for the governance of biodiversity conservation. **4**.
Hijmans, R. J. 2016. geosphere: Spherical Trigonometry. R package version 1.5-5.
Hijmans, R. J., and J. van Etten. 2023. raster: Geographic data analysis and modeling.
Hornman, M., A. Boele, and E. van Winden. 2022. Ruigpootbuizerd: steeds schaarser, behalve in
piekjaren. Nature Today.
Jetz, W., G. Tertitski, R. Kays, U. Mueller, M. Wikelski, S. Åkesson, Y. Anisimov, A. Antonov,
W. Arnold, F. Bairlein, O. Baltà, D. Baum, M. Beck, O. Belonovich, M. Belyaev, M.
Berger, P. Berthold, S. Bittner, S. Blake, B. Block, D. Bloche, K. Boehning-Gaese, G.
Bohrer, J. Bojarinova, G. Bommas, O. Bourski, A. Bragin, A. Bragin, R. Bristol, V.
Brlík, V. Bulyuk, F. Cagnacci, B. Carlson, T. K. Chapple, K. F. Chefira, Y. Cheng, N.
Chernetsov, G. Cierlik, S. S. Christiansen, O. Clarabuch, W. Cochran, J. M. Cornelius, I.
Couzin, M. C. Crofoot, S. Cruz, A. Davydov, S. Davidson, S. Dech, D. Dechmann, E.

Demidova, J. Dettmann, S. Dittmar, D. Dorofeev, D. Drenckhahn, V. Dubyanskiy, N. Egorov, S. Ehnbom, D. Ellis-Soto, R. Ewald, C. Feare, I. Fefelov, P. Fehérvári, W. Fiedler, A. Flack, M. Fröböse, I. Fufachev, P. Futoran, V. Gabyshev, A. Gagliardo, S. Garthe, S. Gashkov, L. Gibson, W. Goymann, G. Gruppe, C. Guglielmo, P. Hartl, A. Hedenström, A. Hegemann, G. Heine, M. H. Ruiz, H. Hofer, F. Huber, E. Hurme, F. Iannarilli, M. Illa, A. Isaev, B. Jakobsen, L. Jenni, S. Jenni-Eiermann, B. Jesmer, F. Jiguet, T. Karimova, N. J. Kasdin, F. Kazansky, R. Kirillin, T. Klinner, A. Knopp, A. Kölzsch, A. Kondratyev, M. Krondorf, P. Ktitorov, O. Kulikova, R. S. Kumar, C. Künzer, A. Larionov, C. Larose, F. Liechti, N. Linek, A. Lohr, A. Lushchekina, K. Mansfield, M. Matantseva, M. Markovets, P. Marra, J. F. Masello, J. Melzheimer, M. H. M. Menz, S. Menzie, S. Meshcheryagina, D. Miquelle, V. Morozov, A. Mukhin, I. Müller, T. Mueller, J. G. Navedo, R. Nathan, L. Nelson, Z. Németh, S. Newman, R. Norris, O. Nsengimana, I. Okhlopkov, W. Oleś, R. Oliver, T. O'Mara, P. Palatitz, J. Partecke, R. Pavlick, A. Pedenko, A. Perry, J. Pham, D. Piechowski, A. Pierce, T. Piersma, W. Pitz, D. Plettemeier, I. Pokrovskaya, L. Pokrovskaya, I. Pokrovsky, M. Pot, P. Procházka, P. Quillfeldt, E. Rakhimberdiev, M. Ramenofsky, A. Ranipeta, J. Rapczyński, M. Remisiewicz, V. Rozhnov, F. Rienks, V. Rozhnov, C. Rutz, V. Sakhvon, N. Sapir, K. Safi, F. Schäuffelhut, D. Schimel, A. Schmidt, J. Shamoun-Baranes, A. Sharikov, L. Shearer, E. Shemyakin, S. Sherub, R. Shipley, Y. Sica, T. B. Smith, S. Simonov, K. Snell, A. Sokolov, V. Sokolov, O. Solomina, M. Soloviev, F. Spina, K. Spoelstra, M. Storhas, T. Sviridova, G. Swenson, P. Taylor, K. Thorup, A. Tsvey, M. Tucker, S. Tuppen, W. Turner, I. Twizeyimana, H. van der Jeugd, L. van Schalkwyk, M. van Toor, P. Viljoen, M. E. Visser, T. Volkmer, A. Volkov, S. Volkov, O. Volkov, J. A. C. von Rönn, B. Vorneweg, B. Wachter, J. Waldenström, N. Weber, M. Wegmann, A. Wehr, R. Weinzierl, J. Weppler, D. Wilcove, T. Wild, H. J. Williams, J. Wilshire, J. Wingfield, M. Wunder, A. Yachmennikova, S. Yanco, E. Yohannes, A. Zeller, C. Ziegler, A. Zięcik, and C. Zook. 2022. Biological Earth observation with animal sensors. Trends in Ecology & Evolution **37**:293-298.

JIa, Q., K. Koyama, C.-Y. Choi, H.-J. Kim, L. Cao, D. Gao, G. Liu, and A. D. Fox. 2016. Population estimates and geographical distributions of swans and geese in East Asia based on counts during the non-breeding season. Bird Conservation International **26**:397-417.

Kays, R., M. C. Crofoot, W. Jetz, and M. Wikelski. 2015. Terrestrial animal tracking as an eye on life and planet. Science **348**.

Kays, R., and M. Wikelski. 2023. The Internet of Animals: what it is, what it could be. Trends in Ecology & Evolution.

Kjellén, N., M. Hake, and T. Alerstam. 1997. Strategies of two Ospreys Pandion haliaetus migrating between Sweden and tropical Africa as revealed by satellite tracking. Journal of Avian Biology:15-23.

Koivurova, T., and A. Shibata. 2023. After Russia's invasion of Ukraine in 2022: Can we still cooperate with Russia in the Arctic? Polar Record **59**:e12.

Lenth, R., H. Singmann, J. Love, P. Buerkner, and M. Herve. 2019. Package 'emmeans'.

Lislevand, T., S. Hahn, S. Rislaa, and M. Briedis. 2020. First records of complete annual cycles in water rails Rallus aquaticus show evidence of itinerant breeding and a complex migration system. Journal of Avian Biology **51**.

Mearns, R., and I. Newton. 1984. Turnover and dispersal in a peregrine Falco peregrinus population. Ibis **126**:347-355.

Moreau, R. E. 1972. The Palaearctic-African Bird Migration Systems. Academic Press.

National Audubon Society. 2024. Audubon Christmas Bird Count. National Audubon Society.

Newton, I. 2008. The Migration Ecology of Birds. Elsevier, Oxford, UK.

Pokrovsky, I., D. Ehrich, I. Fufachev, R. A. Ims, O. Kulikova, A. Sokolov, N. Sokolova, V.
Sokolov, and N. G. Yoccoz. 2019. Nest association between two predators as a
behavioral response to the low density of rodents. The Auk **137**.
Pokrovsky, I., D. Ehrich, R. Ims, O. Kulikova, N. Lecomte, and N. Yoccoz. 2014. Diet, nesting
density, and breeding success of rough-legged buzzards (Buteo lagopus) on the Nenetsky
Ridge, Arctic Russia. Polar Biology **37**:447-457.
Pokrovsky, I., D. Ehrich, R. A. Ims, A. V. Kondratyev, H. Kruckenberg, O. Kulikova, J.
Mihnevich, L. Pokrovskaya, and A. Shienok. 2015. Rough-Legged Buzzards, Arctic
Foxes and Red Foxes in a Tundra Ecosystem without Rodents. Plos One **10**:e0118740.
Pokrovsky, I., A. Kölzsch, S. Sherub, W. Fiedler, P. Glazov, O. Kulikova, M. Wikelski, and A.
Flack. 2021. Longer days enable higher diurnal activity for migratory birds. Journal of
Animal Ecology **90**:2161-2171.
Posit team. 2022. RStudio: Integrated Development Environment for R. Posit Software, PBC,
Boston, MA.
R Development Core Team. 2022. R: A language and environment for statistical computing. R
Foundation for Statistical Computing, Vienna, Austria.
Reif, J. 2013. Long-term trends in bird populations: a review of patterns and potential drivers in
North America and Europe. Acta ornithologica **48**:1-16.
Robinson, R. A., N. A. Clark, R. Lanctot, S. Nebel, B. Harrington, J. A. Clark, J. A. Gill, H.
Meltofte, D. I. Rogers, and K. G. Rogers. 2005. Long term demographic monitoring of
wader populations in non-breeding areas. Wader Study Group Bulletin **106**:17-29.
Schlaich, A. E., V. Bretagnolle, C. Both, B. J. Koks, and R. H. Klaassen. 2023. On the Wintering
Ecology of Montagu's Harriers in West Africa: Itinerancy in Relation to Varying Annual
Environmental Conditions. Ardea **111**:321-342.
Serrano, D., J. L. Tella, M. G. Forero, and J. A. Donazar. 2001. Factors affecting breeding
dispersal in the facultatively colonial lesser kestrel: individual experience vs. conspecific
cues. Journal of Animal Ecology **70**:568-578.
Sonerud, G. A. 1986. Effect of snow cover on seasonal changes in diet, habitat, and regional
distribution of raptors that prey on small mammals in boreal zones of Fennoscandia.
Ecography **9**:33-47.
Strandberg, R., T. Alerstam, M. Hake, and N. Kjellen. 2009. Short-distance migration of the
Common Buzzard Buteo buteo recorded by satellite tracking. Ibis **151**:200-206.
Sundell, J., O. Huitu, H. Henttonen, A. Kaikusalo, E. Korpimaki, H. Pietiainen, P. Saurola, and I.
Hanski. 2004. Large-scale spatial dynamics of vole populations in Finland revealed by
the breeding success of vole-eating avian predators. Journal of Animal Ecology **73**:167-
178.
Tast, J., A. Kaikusalo, and M. Lagerström. 2010. Diet composition of breeding rough-legged
buzzards Buteo lagopus at Kilpisjärvi, NW Finnish Lapland. Kilpisjärvi Notes **22**:10-18.
Terraube, J., A. Villers, L. Ruffino, L. Iso-Iivari, H. Henttonen, T. Oksanen, and E. Korpimäki.
2015. Coping with fast climate change in northern ecosystems: mechanisms underlying
the population-level response of a specialist avian predator. Ecography **38**:690-699.
Thorup, K., A. P. Tøttrup, M. Willemoes, R. H. G. Klaassen, R. Strandberg, M. L. Vega, H. P.
Dasari, M. B. Araújo, M. Wikelski, and C. Rahbek. 2017. Resource tracking within and
across continents in long-distance bird migrants. Science advances **3**:e1601360.
Trierweiler, C., W. C. Mullié, R. H. Drent, K. M. Exo, J. Komdeur, F. Bairlein, A. Harouna, M.
de Bakker, and B. J. Koks. 2013. A Palaearctic migratory raptor species tracks shifting
prey availability within its wintering range in the Sahel. Journal of Animal Ecology
82:107-120.

Vansteelant, W., W. Faveyts, and J. Buckens. 2011. Opmerkelijke Ruigpootbuizerdinvasie in de
winter 2010-2011: oorzaken in een historische en Europese context. Natuur.oriolus
77:87-98.
Walker, D. A., M. K. Raynolds, F. J. A. Daniels, E. Einarsson, A. Elvebakk, W. A. Gould, A. E.
Katenin, S. S. Kholod, C. J. Markon, E. S. Melnikov, N. G. Moskalenko, S. S. Talbot, B.
A. Yurtsev, and C. Team. 2005. The Circumpolar Arctic vegetation map. Journal of
Vegetation Science **16**:267-282.
Wetlands International. 2024. International Waterbird Census.
Whitfield, D. P., K. Duffy, D. R. McLeod, R. J. Evans, A. M. MacLennan, R. Reid, D. Sexton, J.
D. Wilson, and A. Douse. 2009. Juvenile dispersal of white-tailed eagles in western
Scotland. Journal of Raptor Research **43**:110-120.
Wikelski, M., R. W. Kays, N. J. Kasdin, K. Thorup, J. A. Smith, and G. W. Swenson. 2007.
Going wild: what a global small-animal tracking system could do for experimental
biologists. Journal of Experimental Biology **210**:181-186.

a)

Vegetation cover (%)

Quick phase

Grass-lands, Crop-lands, Forests, Urban, Shrub-lands

b)

Vegetation cover (%)

Slow phase

Grass-lands, Crop-lands, Forest, Urban, Shrub-lands

c)

Quick phase
- spring
- fall

Slow phase
- part 1
- part 2

a)

Slow phase part 1

OCT
NOV
DEC
JAN

Slow phase part 2

APR
MAR
FEB

b)

hypothetical snow cover
if birds winter in the place
where they have arrived
from the North

real snow cover that birds
experience during the
winter

hypothetical snow cover
if birds fly directly to the
Southwest and stay
there all winter

Snow cover, %

Oct Nov Dec Jan Feb Mar Apr

a)

For species with **dynamic winter range**, when mid-winter counts are used to determine winter range temporal heterogeneity can be mistaken with spatial heterogeneity

uncommon

common

For species with **dynamic winter range**, when GPS tracking is used to determine winter range temporal heterogeneity becomes evident and should be shown on the map

X, IV

XI, II, III

XII, I

Numbers indicate months the birds are in the area

b)

Species abundance

abundance decline within a specific part of the winter range

Years

Extinct

EX EW

Threatened

CR EN VU

Least Concern

NT LC

Years

For species with **dynamic winter range**, a decline in abundance within a specific part of the winter range may indicate changes in a range dynamics rather than an overall population decrease