Foxtrot migration and dynamic over-wintering range of an arctic raptor

Ivan Pokrovsky[1] *, Teja Curk[2], Andreas Dietz[3], Ivan Fufachev[4], Olga Kulikova[5], Sebastian Rößler[3], Martin Wikelski[1,6]

[1] *Department of Migration, Max Planck Institute of Animal Behavior, Radolfzell, Germany*

[2] *Leibniz Institute for Zoo and Wildlife Research, Berlin, Germany*

[3] *Deutsches Zentrum für Luft- und Raumfahrt e.V. (DLR), Wessling, Germany*

[4] *Institute of Plant and Animal Ecology, Yekaterinburg, Russia*

[5] *Institute of Biological Problems of the North, Magadan, Russia*

[6] *Centre for the Advanced Study of Collective Behaviour, University of Konstanz, 78457 Konstanz, Germany*

* Corresponding author

E-mail: ipokrovsky@ab.mpg.de

ABSTRACT

Advances in tracking technologies have revealed the diverse migration patterns of birds, which are critical for range mapping and population estimation. Population trends are usually estimated in breeding ranges where birds remain stationary, but for species that breed in remote areas like the Arctic, these trends are often assessed in over-wintering ranges. Evaluating population trends during the wintering season is challenging due to the extensive movements of birds in these ranges, which require a deep understanding of the movement dynamics. However, these movements remain understudied, particularly in the mid-latitudes, where many Arctic breeders overwinter, increasing uncertainty in their ranges and numbers. Here, we show that the Arctic breeding raptor Rough-legged buzzard, which overwinters in the mid-latitudes, has a specific wintering strategy. After migrating ca 1,500 km from the Arctic to mid-latitudes, the birds continue to move throughout the entire over-wintering period, covering an additional 1,000 km southwestward and then back northeastward as the snowline advances. This ongoing movement makes their wintering range dynamic over the course of the season. In essence, this movement represents an extension of the quick migration process, albeit at a slower pace, and we have termed this migration pattern 'foxtrot migration,' drawing an analogy to the alternating fast and slow movements of the foxtrot dance. These results highlight the potential errors in range mapping from single mid-winter surveys and emphasize the importance of this migration pattern in assessing the conservation status of bird species. Understanding this migration pattern could help to correctly estimate bird populations in over-wintering ranges, which is especially important for species that nest in hard-to-reach regions such as the Arctic.

Keywords: *migration patterns, bird movement ecology, bird migration, types of migration, conservation, species ranges, snow cover, dynamic range, foxtrot migration.*

INTRODUCTION

In recent years, advances in tracking technology have greatly improved our understanding of bird migration (Wikelski et al. 2007, Kays et al. 2015, Jetz et al. 2022, Kays and Wikelski 2023). By tracking tagged birds, we can study various migratory behaviors, such as their propensity to migrate, connectivity and philopatry, length of migration routes, variations in flight speed, and movements within a single season (Alerstam et al. 2003, Newton 2008, Alerstam 2011, Berthold et al. 2013, Trierweiler et al. 2013, Schlaich et al. 2023). Different combinations of these behaviors contribute to various bird migration patterns (Chapman et al. 2014, Alerstam and Bäckman 2018, Lislevand et al. 2020). Understanding the migratory patterns of birds is critical to accurately map their ranges and estimate their population trends. Typically, population trends are assessed in breeding areas, where birds are linked to their nesting sites, allowing reliable estimates of breeding numbers (Reif 2013). However, access to some breeding areas, such as the Arctic, presents logistical and political challenges (Gallo-Cajiao et al. 2023, Koivurova and Shibata 2023). As a result, population trends for these species are often estimated in over-wintering ranges where birds aren't tied to specific locations (Robinson et al. 2005, JIa et al. 2016). This challenges accurately censusing birds and delineating their ranges, especially when birds move extensively in over-wintering ranges during the season, and censuses are conducted only once per over-wintering season. Therefore, understanding bird movements in over-wintering ranges is essential for understanding the ecology of birds in these regions and has practical implications for accurately assessing their ranges and trends in their populations.

Many bird species experience significant seasonal changes in food availability within their over-wintering range due to environmental conditions. This often triggers movements across these areas. For Palearctic migrants wintering in Africa, these movements have been studied extensively (Moreau 1972, Trierweiler et al. 2013, Thorup et al. 2017, Schlaich et al. 2023). However, the over-wintering movements of mid-latitude birds remain poorly understood. At the same time, many Arctic birds overwinter in the mid-latitudes, and understanding their movement patterns could be crucial for interpreting their population trends. In the mid-latitudes, food availability is strongly influenced by environmental factors such as snow cover dynamics. The progressive movement of the snow cover line from north (-east) to south (-west) and back again between October and May in many mid-latitude regions, particularly those exposed to northwesterly wind systems, significantly affects food availability for numerous bird species (Sonerud 1986, Vansteelant et al. 2011). We propose that species dependent on this environmental factor gradually move away from snow-covered areas during winter and then

gradually move back in the opposite direction, resulting in a directed and continuous

displacement of their over-wintering range.

The Rough-legged buzzard (*Buteo lagopus*) is an Arctic breeding and mid-latitude

wintering raptor (Ferguson-Lees and Christie 2001, Bechard and Swem 2002). Rough-legged

buzzards feed mainly on small rodents in the Arctic and its wintering grounds (Tast et al. 2010,

Pokrovsky et al. 2014). They prefer open areas for hunting, and trees and tall bushes or uplands

for resting and nesting. In the Arctic, where they nest, these habitats are in the southern and

typical tundra zone (Walker et al. 2005). In the mid-latitudes, where they overwinter, these

habitats are in the zone of fields with scattered patches of forests (wooded fields). The taiga

zone, located between the tundra and wooded fields zones, offers limited open space, and is

generally unsuitable for the species, although they may nest in the northern taiga near the tundra

zone (Sundell et al. 2004). Thus, Rough-legged buzzards migrate from the Arctic through the

taiga zone to the wooded fields zone for wintering. However, environmental conditions in the

mid-latitudes vary, potentially influencing their migratory behavior.

The spatio-temporal latitudinal and longitudinal gradients in environmental conditions

impact bird populations (Kiat et al. 2021). Deep snow covers the northeastern regions of suitable

wintering habitats (wooded fields) during mid-winter, significantly hindering the Rough-legged

buzzards' ability to hunt small rodents (Sonerud 1986, Vansteelant et al. 2011, Terraube et al.

2015, Curk et al. 2020). Simultaneously, many Arctic migrants prefer winter as close to the

Arctic as possible to shorten their return to breeding grounds and extend the breeding season

(Berthold et al. 2013). Therefore, we can anticipate three potential non-exclusive migration

strategies for Rough-legged buzzards: 1) birds may migrate through the taiga and northeastern

wooded fields zones to the southwestern areas with less snow cover during winter, 2) birds may

stop in the wooded fields zone after crossing the taiga and move southwest in mid-winter when

snow cover makes hunting rodents impossible, eventually migrating back to the breeding

grounds from southwestern regions, 3) alternatively, similar to the second strategy, birds may

move southwest but return to the northeastern wooded fields zone after mid-winter to be closer

to the breeding grounds, and migrating to the Arctic from there.

In this study, we tracked GPS-tagged Rough-legged buzzards for ten years and analyzed

their movements in the over-wintering mid-latitude range in relation to snow cover. For this

study, we made the following predictions. 1) Rough-legged buzzards could exhibit a directional

and seasonal movement pattern during the over-wintering period, moving from the northeast to

the southwest and back again. This would result in a dynamic over-wintering range that would

continue to move geographically throughout the season. 2) The over-wintering movements

would differ from the fall and spring migrations in duration, extent, speed, and direction but

would continue throughout the season. Thus, over the entire annual cycle, Rough-legged

buzzards would be characterized by a migratory pattern consisting of an alternation of quick (fall

and spring migrations) and slow (over-wintering movements) phases. 3) Over-wintering

movements would occur in suitable open habitats, while fall and spring migrations would occur

in unsuitable forested areas. Thus, vegetation land cover will determine the extent of quick and

slow migration phases. 4) During over-wintering movements, Rough-legged buzzards will

experience less snow cover than if they had stayed where they arrived at the end of the fall

migration. Thus, snow cover dynamics will determine the dynamics of over-wintering

movements.

125   MATERIAL AND METHODS

126   Dataset

For this study, we tracked 43 adult Rough-legged buzzards (35 females and eight males) with the

solar GPS-GSM loggers (E-obs GmbH and UKn – University of Konstanz). Twenty-eight birds

were fitted with E-obs loggers, 13 birds were fitted with UKn loggers, and two birds were

initially fitted with UKn loggers that were later replaced with E-obs loggers. E-obs loggers

weighed 45 g, or 3.6% of the bird's weight, while UKn loggers weighed 15 g, or 1.2% of the

bird's weight. E-obs loggers recorded GPS positions every hour in full battery mode and every 5

hours in normal battery mode. UKn loggers recorded GPS positions every hour with a full

battery and every 12 hours with a normal battery. Both models operated continuously 24 hours a

day. More detailed information about this dataset can be found in Curk et al. (2022).

The fieldwork was carried out in the Russian Arctic in 2013-2019 at four study sites:

Kolguev Island (69°16′N, 48°87′E), Nenetsky Nature Reserve (68°20′N, 53°18′E), Vaigach

Island (69°43′N, 60°08′E), and Yamal Peninsula (68°12′N, 68°59′E). For details on capture

methods, see Curk et al. (2022); for detailed descriptions of study sites, see Pokrovsky et al.

(2015) for Kolguev Island and Pokrovsky et al. (2019) for Yamal and Nenetsky.

During data pre-processing, we removed duplicated timestamps and calculated the mean

daily positions of each individual. We partitioned the resulting dataset into several periods: 1)

breeding, 2) fall migration, 3)1st part of winter, 4) 2nd part of winter, and 5) spring migration. We

estimated the migration dates – the start and stop dates of the spring and fall migrations – using

an iterative search procedure for piecewise regression described by Crawley (2007). We created

146 two models for each of the four dates: a simple model and a piecewise one. The simple model

147 was linear, with latitude as the response variable and day of the year as a fixed effect. The

148 piecewise model was also linear, with latitude as the response variable and day of the year as a

149 fixed effect, and included two logical statements in its formula because we expected two linear

150 segments in the fit. Specifically, it used "Day Of the Year < BREAK" to define the left

151 regression (before the estimated date) and "Day Of the Year ≥ BREAK" for the right regression

152 (after the estimated date). We explored different BREAK values to fit the piecewise model,

153 looking for the value that minimized the residual standard error. The BREAK values examined

154 included those associated with the start and end of the fall migration (September 5 to October 5

155 and October 5 to November 5, respectively), as well as those associated with the start and end of

156 the spring migration (April 10 to May 10 and May 10 to June 10, respectively). We then

157 validated (using the R function 'anova') that the piecewise model significantly improved the fit

158 compared to the simple model (p<0.001). We estimated the date between winter's first and

159 second parts as the day when the mean daily latitude was the lowest.

Data analysis

161 First, we used linear mixed-effects models (R function 'lmer' in the library 'lme4' (Bates et al.

162 2015)) to investigate whether or not Rough-legged buzzards migrated during winter. Latitude

163 was the response variable, day of the year was a fixed effect, and individuals and year were

164 included as random effects. Analyses were conducted separately for each migration period (fall,

165 first phase of winter, second phase of winter, and spring). For both phases of the winter

166 migration, we analyzed two additional models with longitude as the response variable instead of

167 latitude. Likelihood ratio tests were used to compare candidate models. The year was not a

168 calendar year but a year between two consecutive breeding seasons. Thus, fall migration,

169 consecutive winter, and consecutive spring have the same value for the year. The day of the year

170 was recalculated consecutively.

171  Second, we used linear mixed-effects models (R function 'lmer' in the library 'lme4') to

172 investigate whether migrations' parameters differ between the migration periods. We analyzed

173 four migration parameters: distance, duration, speed, and direction. The distance was calculated

174 as the distance between two coordinates (start and end of migration) using the R function 'distm'

175 in the library 'geosphere' (Hijmans 2016). The duration was calculated as the number of days

176 between the start and end of migration. Speed was calculated as the ratio of distance to duration.

177 The direction was calculated as the bearing from the start of the migration coordinates to the end

178 of the migration coordinates using the R function 'bearing' in the library 'geosphere' (Hijmans

2016). Specific migration parameters (see below) were the response variables, the type of

migration was a fixed factor, and individuals were a random factor. Likelihood ratio tests were

used to compare candidate models. We considered four different parameters of migration

(distance, duration, speed, and direction) and four types of migration (fall, first phase of winter,

second phase of winter, and spring). The analysis was done separately for each of the migration

parameters. Then, we used post hoc comparisons using the R function 'emmeans' in the library

'emmeans' (Lenth et al. 2019) to compare the estimated means.

In some raptor species, adult females disperse further than males (Mearns and Newton

1984, Serrano et al. 2001, Bildstein 2006, Whitfield et al. 2009). Therefore, we conducted an

additional analysis on the effect of sex on migration length using linear mixed-effects models (R

function 'lmer' in the library 'lme4'). The migration distance was used as the response variable,

sex as a fixed factor, and individuals as a random factor.

Third, we investigated whether vegetation land cover differed between areas crossed

during the quick (fall and spring) and slow (winter) migrations. We used the combined Terra and

Aqua Moderate Resolution Imaging Spectroradiometer (MODIS) Land Cover Climate

Modelling Grid (CMG) (MCD12C1) version 6 dataset (Friedl and Sulla-Menashe 2015). We

used a modified Leaf Area Index (LAI) as a classification scheme. We combined all four forest

types and savannas into one category (forest) and excluded the categories: water bodies, and

unclassified. We, therefore, had five types of vegetation cover: forest, grassland, cropland,

shrubland, and urban. We annotated the mean daily positions with the vegetation cover type

using the Env-DATA tool (Dodge et al. 2013). We used general linear mixed effects models with

a binomial distribution (R function 'glmer' in the library 'lme4') to investigate whether

vegetation cover types differ between migration periods. Presence/absence of the studied

vegetation land cover type was used as a response variable, migration type as a fixed factor, and

individuals as a random factor. The analysis was done separately for each of the vegetation land

cover types.

Fourth, we investigated whether snow cover could drive the slow migration phenomenon.

We then compared the snow cover conditions the birds experienced during the winter with two

hypothetical snow cover conditions that the birds would have experienced if they had not

migrated during the winter. The first hypothetical snow cover condition would have happened if

the birds had stayed where they arrived from the north (i.e., where their fall migration ended). To

estimate this parameter, we calculated the winter dynamics of the average snow cover at the

minimum convex polygons (MCP) occupied by the birds in October and April (northeast of their

winter range). A second hypothetical snow cover condition would be if the birds flew
immediately to the southwest and spent the whole winter there. To evaluate this, we calculated
the winter dynamics of average snow cover on the MCPs occupied by the birds in January and
February (southwest of their winter range). We then compared the values obtained for the real
snow cover and two hypothetical snow covers using general linear mixed effects models with a
binomial distribution (R function 'glmer' in the 'lme4' library). Presence/absence of snow cover
was used as a response variable, type of snow cover (real, 1st hypothetical, or 2nd hypothetical) as
a fixed factor, and years as a random factor. The analysis was done separately for each month.
We then used post hoc comparisons to compare the estimated means, using the R function
'emmeans' in the 'emmeans' library (Lenth et al. 2019). We also tested whether the distance
between two consecutive monthly MCPs during the over-wintering period is influenced by snow
cover extent or snow cover difference. To investigate this, we employed a linear model, with the
distance between two consecutive monthly MCPs as the response variable. Snow cover at the
first MCP and the difference in snow cover between two consecutive MCPs were included as
fixed factors.

We obtained monthly snow cover data with a spatial resolution of ca 500 meters (Global
SnowPack MODIS) from the German Aerospace Center (DLR). This product is based on the
Moderate Resolution Imaging Spectroradiometer (MODIS) daily snow cover products
MOD10A1 and MYD10A1 (version 6 as provided by the National Snow and Ice Data Center
NSIDC), which have been processed to remove the gaps due to cloud cover and polar darkness
(Dietz et al. 2015). These processing steps include a combination of data available from different
satellites (Aqua and Terra), 3-day temporal moving window filtering, a regional snow line
elevation interpolation relying on a Digital Elevation Model (DEM), and a seasonal filter
running through the time series for the whole hydrological year (1st of September through
August 31st). The proportion of days in which one pixel is snow-covered per month is referred to
here as fractional snow cover and is derived from these daily gap-filled rasters. Five MODIS
tiles (h19v03, h20v03, h20v04, h21v03 and h21v04) were mosaicked and re-projected to
WGS84. Then, for each month from October to April, we calculated 95% minimum convex
polygons (MCPs) for the distribution of Rough-legged buzzards using the R function 'mcp' in
library 'adehabitatHR' (Calenge 2006). We extracted mean snow cover values from each MCP
from every monthly snow cover raster separately, using the R library 'raster' (Hijmans and van
Etten 2023).

All calculations were performed using R version 4.2.2 'Innocent and Trusting' (R

Development Core Team 2022) and RStudio version 353 'Elsbeth Geranium' (Posit team 2022).

RESULTS

Migration of Rough-legged buzzards – always on the move

Except during the breeding season, Rough-legged buzzard migration continues throughout the

year, even after the birds´ arrival at their traditionally recognized ´wintering grounds´ (Figure 1a,

b). Rough-legged buzzards started their fall migration on 28 September (hereafter mean±sd for

the day of the year: 271±11, n=31) and ended on 12 October (285±11, n=33). The mean

latitude/longitude where the birds ended their fall migration was 55.57±1.92°/49.35±5.63°

(Figure 1a). During the winter, birds continued to migrate at a slower pace down to 49.53±2.01°

latitude (on 5 February, 36±40, n=23) and 34.29±5.11° longitude (on 24 January, 24±47, n=23).

Afterward, during the second part of the winter, birds returned to 55.52±2.63° latitude and

49.79±8.24° longitude to start the spring migration (Figure 1a). Rough-legged buzzards started

their spring migration to the Arctic on 27 April (117±7, n=27) and arrived at the breeding

grounds on 15 May (135±8, n=18).

In the following, we will refer to the spring and fall migrations as the quick phase, the

over-wintering movement to the lowest point of latitude as the 1[st] part of the slow phase, and the

movement from the lowest point of latitude to the starting point of the spring migration as the 2[nd]

part of the slow phase (Figure 1a). For both quick and slow phases of the migration, linear

mixed-effects models with the season as a fixed factor received higher support from the

likelihood ratio test ($p < 0.001$, Tables S1-S3).

Figure 1. Migration of Rough-legged buzzards. Q – quick phase. Qf – Quick fall phase (orange), Qs – Quick spring phase (yellow), S1 – Slow phase, 1st part (light blue), S2 – Slow phase, 2nd part (dark blue). a) Change in the latitude of 43 Rough-legged buzzards during the year, red line – mean latitude of all birds, black vertical lines – mean dates of start and end of the migration phases, blue vertical line – mean date of the minimum latitude. Grey, sky blue, and piggy pink shaded areas – standard deviation of the means. b) Migration map. c) Difference in the migration parameters between the migration phases. Lines on the direction plot (down, right) represent the mean value for each bird; arrows represent the mean direction for each phase. Boxes on the boxplots show the interquartile range, the whiskers – maximum and minimum values.

Quick-slow phase features comparison

During the quick phase, individual birds flew greater distances in a shorter time, i.e., at a faster rate, than during the slow phase. After arriving at what is traditionally known as the wintering grounds, the direction of migration changed, so the direction of quick and slow phases also differed (Table 1, Figure 1c). The quick phase (one part) was 1415±50 km long (hereafter mean±sd), whereas the slow phase (one part) was 1026±55 km, i.e., 389±60 km shorter

(p<0.001, Table S4, Figure 1c). During the quick phase, birds flew for 15±3 days, and one part

of the slow phase lasted 100±4 days, i.e., 85±5 days longer (p<0.001, Table S5, Figure 1c). At

the same time, the second part of the slow phase was 54±7 days shorter than the first (p<0.001,

Table S5, Figure 1c). The migration speed was 104±6 km/day during the quick phase and 12±7

km/day during the slow phase, i.e., about eight times higher (p<0.001, Table S6, Figure 1c).

During the fall migration, birds moved in the SSW direction (198±3 deg), then turned 50±3 deg

(p<0.001, Table S7, Figure 1c) to the west and started their 1st slow phase until mid-winter. After

that, they turned back to the NEE direction (57±2 deg) and performed their 2nd slow phase for

several months until they turned 54±3 deg (p<0.001, Table S7, Figure 1c) to the north and started

their spring migration. We found no significant difference between the migration distances of

males and females (Table S8).

Table 1. Parameters of the Rough-legged buzzards' migration (mean±sd).

Phase of migration	Sub-phase of migration	Distance (km)	Duration (days)	Speed (km/day)	Direction (deg)
Quick		1415±50	15±3	104±6	
	Spring	1544±72	18±4	100±8	7±2
	Fall	1334±59	14±4	107±7	198±3
Slow		1026±55	100±4	12±7	
	1st phase	1016±71	121±4	8±7	251±3
	2nd phase	1042±71	66±6	17±9	57±2

Vegetation land cover during migration

During quick phases, Rough-legged buzzards crossed the forest zone, while during the slow

phase, they migrated within the grassland and cropland zones (Figure 2). Rough-legged buzzards

migrated fast across the tundra zone on the north in the Arctic and then through the taiga zone.

Therefore, during the quick phase, the three most common vegetation land cover types were

forest (44.5±2.9 %, hereafter, percentage of all mean daily positions annotated with the given

vegetation type ± sd), shrublands (29.9±3 %), and grasslands (24.7±2.8 %, Figure 2a). During

the slow phase, the three most common vegetation land cover types were grasslands (65.1±6.2

%), croplands (26.9±6.3 %), and forests (4.9±1.4 %, Figure 2b). According to the linear mixed-

effects models, the percentage of all vegetation land cover types differed between the slow and

quick phases (p<0.001, Table S9), except for the urban lands. Urban lands were more common

during the slow than quick phase (Figure 2). However, this type has been annotated for too few

birds to make an adequate comparison.

Figure 2. Vegetation land cover during quick and slow phases of the migration. a) Quick
phase (spring and fall periods together). b) Slow phase (1st and 2nd parts together). On both a)
and b), the bars show the percentage of all mean daily positions annotated with the given
vegetation type ± sd. c) Migration map.

Dynamic winter range

During the slow phase of the migration, Rough-legged buzzards experienced snow cover ranging

from 4.8±1.0% in October (hereafter mean±sd) to 85.2±4.6% in February (Figure 3). If birds

spent the winter in the place where they arrived after the fall migration, they would experience

snow cover conditions ranging from 4.6±0.6% in October to 99.5±0.1% in February (Figure 3b,

green line). If birds fly directly to the southwest and stay there for the whole winter, they would

experience snow cover conditions ranging from 1.4±0.2 % in October to 81.1±5.0 % in January

(Figure 3b, red line). Thus, if birds fly immediately to the southwest and stay there until the end

of the winter, they will find conditions with less snow cover in spring (p<0.001, Table S10). And

if birds stay where they ended the fall migration, they will find themselves in situations with

more snow cover (p<0.001, Table S10). In the latter case, the difference between real and

hypothetical situations is not as pronounced (85.2% vs. 99.5%), but it means that snow cover

will be close to 100% for several months in this hypothetical situation (Figure 3b). The distance
between two consecutive monthly MCPs during the overwintering period did not depend on the
snow cover extent (p=0.45) or the difference in snow cover (p=0.36, Figure S1).

Figure 3. Snow cover conditions during the slow phase of the migration. a) 95% Minimum
convex polygons (MCPs) of Rough-legged buzzards during winter. Arrows indicate the direction
of the movement across months. OCT – October, NOV – November, DEC – December, JAN –
January, FEB – February, MAR – March, APR – April. b) Snow cover conditions for the real
situation (black) and two hypothetical situations – if birds spend the winter in the place where
they arrived after the fall migration (green) and if birds fly directly to the southwest and stay
there all winter (red). Dots represent mean values, error lines – standard deviations.

DISCUSSION

336 Our study identified and characterized a bird migration pattern consisting of an alternation of
337 quick (fall and spring migrations) and slow (over-wintering movements) phases. This migration
338 pattern causes the over-wintering range of birds to shift and become dynamic throughout the

339 season. This has obvious implications for range delineation and assessment of bird population
340 trends.

Over-wintering movements and migratory patterns

Different bird species employ various wintering strategies, leading to diverse migration patterns.

One strategy is 'residency,' where birds arriving at over-wintering grounds remain there

throughout the season, undertaking only minor foraging flights (Figure 4a). Their home ranges in

over-wintering ranges are compatible with those in breeding ranges (Kjellén et al. 1997,

Alerstam et al. 2006, Newton 2008). Another strategy, initially identified by Moreau (Moreau

1972), is 'itinerancy,' wherein birds fly between several sites separated by distances ranging

from ten to several hundred kilometers throughout the over-wintering period, spending 1-2

months at each site (Figure 4b). Most Palearctic-African species adopt this itinerant behavior

(Trierweiler et al. 2013, Thorup et al. 2017, Schlaich et al. 2023). The third strategy investigated

here involves birds moving slowly and steadily throughout the over-wintering season, initially in

one direction and then in the opposite direction. In 2009, Roine Strandberg and his colleagues

used GPS transmitters to track Common buzzards *(Buteo buteo)*. Despite some transmitters in

their study ceasing to function mid-winter, the authors observed a phenomenon they termed

'prolonged autumn migration,' wherein birds arriving at their wintering grounds did not halt

migration but continued moving at a slower pace (Strandberg et al. 2009). In our study, we

observed this phenomenon throughout the annual cycle. We demonstrated that, unlike itinerancy,

movements in the over-wintering range are essentially a continuation of migration at a slower

pace, influenced by external factors such as snow cover dynamics. Considering this behavior as

an extension of migration and examining the complete life cycle of such species, it involves a

quick phase during the transition between breeding and over-wintering ranges, followed by a

slow phase of directed and seasonal movement within the over-wintering range, and finally,

another quick phase towards the breeding range. Drawing an analogy to the alternating quick and

slow movements of the foxtrot dance, we propose the term 'foxtrot migration' for this seasonal

movement to offer a concise and easily understandable description (Figure 4c). Consequently,

we suggest referring to the over-wintering range of species exhibiting itinerancy or foxtrot

migration as the 'dynamic range' (Figure 4b, c).

Figure 4. Winter strategies and over-wintering range scheme. a) Residency, b) Itinerancy, c) Foxtrot migration. The color bar along the y-axis corresponds to the color coding of the habitats of the chart on the right.

Debates may arise regarding introducing a new term for this phenomenon and whether it should fall under the general term 'nomadism' or be categorized as a modified form of another migratory pattern, such as 'slow directional itinerancy.' We assert that behavioral pattern names should aid in understanding movement strategies and avoid confusion. One might ask whether it

is necessary to distinguish between migration and wintering at all, given that migration is
essentially a matter of tracking spatial and temporal changes in favorable conditions. Authors
studying Montagu's harriers *(Circus pygargus)* argue that such generalization hampers
understanding of wintering strategies (Schlaich et al. 2023). Instead, Schleich et al. (2023)
propose explicitly considering the functions of animal-used sites, particularly their contributions
to 'wintering' and 'refueling for migration.' For instance, cuckoos *(Canorus sp.)* utilize
stopovers for both peak foraging conditions and refueling for long flights to subsequent sites
(Thorup et al. 2017). Conversely, Montagu's harriers exhibit negligible refueling between sites
due to short distances and energy-efficient soaring flight (Trierweiler et al. 2013, Schlaich et al.
2023). In our case, Rough-legged buzzards do not make several long stopovers, unlike species
experiencing itinerancy, but move gradually in one direction and then return during the whole
season (Figure 1a). As Schleich et al. (2023) suggested, examining the functions of animal-used
sites reveals another difference between these strategies. The difference is that itinerant birds
wait for better conditions at subsequent sites. In contrast, in our case, the southwestern areas that
birds reach after several months are suitable and offer better conditions than the northeastern
areas, as they have less snow throughout the whole over-wintering season (Figure 3). All these
differences highlight the discrepancy between the wintering strategies referred to as itinerancy
and those described in this study. Therefore, using another term - foxtrot migration - for this
phenomenon is more appropriate.

The dynamic of the foxtrot migration

Our study affirmed the presence of foxtrot migration and a dynamic over-wintering range in
Rough-legged buzzards. The taiga zone proved to be an unfavorable habitat during the quick
phase of foxtrot migration (Figure 2), given the difficulty of Rough-legged buzzards locating
open areas for hunting in this habitat. Conversely, the grassland and cropland zones served as
favorable habitats during the slow phase throughout the entire over-wintering period (Figure 2),
offering numerous open areas for hunting. Within the grassland and cropland zones, the birds
moved southwest and back over 1,000 km throughout the winter (Figure 1). We did not observe
a tendency for birds to leave over-wintering sites when snow cover reached a certain threshold
(Figure S1a). Similarly, we found no evidence that birds stayed in areas with a certain amount of
snow cover (Figure 3), nor did they leave sites when snow cover increased by a certain amount
(Figure S1b).

We speculate that the studied birds, being experienced, anticipated that northeastern areas
would become inaccessible later in the winter, prompting them to leave early without waiting for

significant snow accumulation. Additionally, other factors, such as brief heavy snowfalls, might

have triggered movement, even if these did not result in sustained increases in snow cover

(Vansteelant et al. 2011). Alternatively, this phenomenon could have been more complex, with

multiple factors possibly acting asynchronously, influencing their overwintering movements

(Yanco et al. 2024). While the ultimate reasons behind the over-wintering movements require

further investigation, it is clear that snow cover was a proximate factor driving the birds'

continual 1,000 km southwest movement during winter (Figure 3). Our analysis revealed that if

Rough-legged buzzards remained at their fall migration endpoint without moving southwest,

they would encounter 14.4% more snow cover (99.5% vs. 85.1%, Figure 3). Although this

difference may seem small (14.4%), it holds significance for rodent-hunting birds, distinguishing

between complete and patchy snow cover. Complete snow cover makes hunting impossible,

rendering these regions unsuitable for Rough-legged buzzards.

Unlike African-Palaearctic migrants exhibiting itinerant migratory behavior – waiting for

suitable conditions in more southerly regions – the southern parts of the over-wintering ranges

for Rough-legged buzzards remain favorable throughout the winter. If Rough-legged buzzards

were to immediately fly to the southwest and stay there for the entire winter, they would

experience 25.7% less snow cover (57.3% vs. 31.6%, Figure 3b). Despite this, it doesn't compel

them to adopt this strategy. Thus, the observed phenomenon appears to be a trade-off between

the desire to remain as close as possible to their breeding ranges and the search for suitable

wintering conditions. Notably, one bird did follow this strategy by flying directly to the

southwest and remaining there throughout the winter (Figure S2), exhibiting a residency

strategy. This observation suggests that the distinction between winter strategies is flexible.

African-Palearctic migrants do not strictly adhere to a binary division between residency

and itinerancy but demonstrate a continuum between these two strategies. Many species in

Africa exhibit a mixture of both strategies in varying proportions; see Schlaich et al. (2023) and

reference therein. Similarly, in the mid-latitudes, species may not only show any of the three

strategies but also a considerable degree of individual and regional variation. For example,

Rough-legged buzzards from western Palearctic regions that overwinter in mid-latitude areas

where snow cover dynamics are less pronounced than in eastern regions are likely to have a

higher proportion of birds with a resident strategy. At the same time, Rough-legged buzzards in

North America also exhibit a foxtrot migration pattern, although with variations. There, birds

make a direct and rapid migration, usually across the boreal forest, followed by slower, more

facultative movements once birds get south of the boreal. However, there is a lot of variation

among individuals and even regions; there are more over-wintering slow movements in eastern

North America but less in western North America (Neil Paprocki pers. comm.). Thus, Rough-

legged buzzards in North America lie along a gradient between foxtrot migration and residency,

likely influenced by different snow cover dynamics on the east and west coasts.

447   The foxtrot migration pattern is expected to be observed in many migratory species that

448   exhibit distinct seasonal cycles in their over-wintering range. This migration pattern is expected

449   to be prevalent for species living in regions where snow cover is a significant determinant of

450   food availability, which is the case in large parts of the mid-latitudes. Therefore, understanding

451   bird migration patterns is critical to accurately mapping ranges and assessing population trends

452   in this region.

453   Mapping ranges and assessing population trends

454   The implementation of our study is twofold: 1) the use of mid-winter bird surveys to determine

455   over-wintering range may yield inaccurate results for species with dynamic range (Figure 5a),

456   and 2) declines in abundance within a particular segment of the over-wintering range may

457   indicate changes in range dynamics rather than widespread declines in species abundance

458   (Figure. 5b).

Figure 5. Mapping ranges and assessing population trends for the species with the dynamic over-wintering ranges. a) Mid-winter counts, often used to determine winter range, may give a misleading representation of the over-wintering range for species with dynamic range. b) Changing the conservation status of a species based on reduced abundance in a particular area may not be appropriate for species with dynamic range. Red – breeding range, blue – winter range.

In North America and Europe, the number of over-wintering birds is typically estimated once a season in mid-winter. In North America, these estimates are made during the Christmas counts, usually once in late December (National Audubon Society 2024), while in Europe, they are made during the IWC counts, usually once in January or February (Wetlands International 2024). This approach can lead to errors in range estimation for birds with dynamic ranges. For such species, habitats occupied by birds in fall and spring will be listed as "uncommon" at best, while habitats occupied by birds only in mid-winter will be listed as "common." However, the situation is the opposite if we consider the time birds spend in these habitats (Figure 5a). For example, based on our study of the Rough-legged buzzard, during mid-winter, the species is predominantly present in the southwestern portion of its over-wintering range, with only a tiny proportion present in the rest of the region. As a result, a map of the over-wintering range may show the species as "common" in the southwest and "uncommon" in the northeast. This map

478 would be inaccurate because, during the entire over-wintering period, Rough-legged buzzards

479 spend both fall and spring in the northeastern part and only mid-winter in the southwestern part.

480   To address this, continuous year-round GPS tracking of the species provides a means to

481 track bird locations throughout the over-wintering season, facilitating the creation of accurate

482 distribution maps, particularly for species that exhibit dynamic ranges. We advocate representing

483 temporal heterogeneity (range dynamics) on maps as distinct zones, denoting periods when the

484 species is abundant in a given area. To distinguish temporal from spatial heterogeneity, we

485 recommend using lines to delineate the boundaries of these zones, rather than color shading, and

486 incorporating numbers to denote the months of species abundance in a given zone (Figure 5a).

487 We suggest ecologists include dynamic over-wintering ranges in descriptions and range maps for

488 foxtrot migratory and itinerant bird species.

489   Population counts for a species are often limited to a portion of its range. Therefore,

490 conclusions about conservation status drawn from such counts may be misleading. A decline in

491 abundance within a particular portion of the over-wintering range may indicate changes in range

492 dynamics rather than a general decline in the species (Figure 5b). For example, climate change

493 may affect snow cover dynamics, reducing its intensity in northern regions. As a result, species

494 whose range dynamics depend on snow cover may choose to remain in the northern areas and

495 not migrate as far south as they traditionally have. Despite this shift in range dynamics, overall

496 species abundance may remain unchanged. Similar patterns have affected Rough-legged

497 buzzards in some areas of the European over-wintering range.

498   A 2022 Dutch study found a decline in wintering Rough-legged buzzards over the last 40

499 years (Hornman et al. 2022). On the one hand, this may represent a conservation concern. On the

500 other hand, applying the rationale of the dynamic range, the apparent local decline may be

501 attributed to climate change, resulting in less comprehensive snow coverage in the northeastern

502 wintering areas of Rough-legged buzzards relative to the Netherlands. Such a shift in snow

503 coverage makes it less probable for the birds to migrate to the Netherlands for overwintering.

504 This proposition is further supported by a study of the winter population dynamics of Rough-

505 legged buzzards in the Netherlands in 2011, showing that the main winter population peak

506 occurred in late December, with many birds migrating (Vansteelant et al. 2011). Vansteelant et

507 al. (2011) also found that the main migration occurred after heavy snowfall in northern Europe,

508 supporting our foxtrot migration explanation for this decline. Therefore, investigating the

509 dynamic range is critical to understanding a species' range and effectively assessing its

510 conservation status.

CONCLUSION

Our study sheds light on the overwintering strategy of the Rough-legged buzzard, an Arctic breeder that migrates to mid-latitudes during the non-breeding season. We identified a distinct migration pattern where, following an initial 1,500 km journey to mid-latitudes, the birds continue to move slowly southwestward and then back northeastward throughout the entire over-wintering season. This movement, covering 1,000 km in one direction, makes their over-wintering range dynamic. We call this migration pattern 'foxtrot migration,' drawing an analogy with the alternating quick and slow steps of the foxtrot dance. Our results highlight the importance of this migration pattern and the concept of 'dynamic range' in revealing potential inaccuracies in range mapping from single midwinter surveys and suggest that species distribution maps should represent temporal variations in range dynamics as separate zones. Understanding and accounting for this migratory behavior is critical for accurately assessing bird populations in over-wintering ranges. This is especially important for species such as Arctic nesting birds, where population assessments are primarily conducted in over-wintering ranges due to the logistical and political challenges of the Arctic.

ACKNOWLEDGMENTS

We are very grateful to everyone who helped us collect data in the field. We would like to thank Dr Theunis Piersma and two anonymous reviewers for their valuable feedback and constructive criticism, which significantly improved the quality of this publication. This study was funded by the Max-Planck Institute of Animal Behavior and the German Air and Space Administration (DLR). We also acknowledge partial funding by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) under Germany's Excellence Strategy – EXC 2117 – 422037984.

REFERENCES

Alerstam, T. 2011. Optimal bird migration revisited. Journal of Ornithology **152**:5-23.

Alerstam, T., and J. Bäckman. 2018. Ecology of animal migration. Current Biology **28**:R968-R972.

Alerstam, T., M. Hake, and N. Kjellén. 2006. Temporal and spatial patterns of repeated migratory journeys by ospreys. Animal Behaviour **71**:555-566.

Alerstam, T., A. Hedenström, and S. Åkesson. 2003. Long-distance migration: evolution and determinants. Oikos **103**:247-260.

Bates, D., M. Mächler, B. Bolker, and S. Walker. 2015. Fitting Linear Mixed-Effects Models Using lme4. Journal of Statistical Software **067**.

Bechard, M. J., and T. R. Swem. 2002. Rough-legged Hawk (Buteo lagopus), The Birds of North America Online (A. Poole, Ed.). Ithaca: Cornell Lab of Ornithology.

Berthold, P., E. Gwinner, and E. Sonnenschein. 2013. Avian migration. Springer Science &
Business Media.
Bildstein, K. L. 2006. Migrating raptors of the world: their ecology & conservation. Cornell
University Press.
Calenge, C. 2006. The package "adehabitat" for the R software: A tool for the analysis of space
and habitat use by animals. Ecological Modelling **197**:516-519.
Chapman, B. B., K. Hulthén, M. Wellenreuther, L.-A. Hansson, J.-Å. Nilsson, and C. Brönmark.
2014. Patterns of animal migration. Pages 11-35 *in* L.-A. Hansson and Å. Susanne,
editors. Animal movement across scales. Oxford University Press.
Crawley, M. J. 2007. The R Book. Wiley Publishing.
Curk, T., O. Kulikova, I. Fufachev, M. Wikelski, K. Safi, and I. Pokrovsky. 2022. Arctic
Migratory Raptor Selects Nesting Area During the Previous Breeding Season. Frontiers
in Ecology and Evolution **10**.
Curk, T., I. Pokrovsky, N. Lecomte, T. Aarvak, D. F. Brinker, K. Burnham, A. Dietz, A. Dixon,
A. Franke, G. Gauthier, K.-O. Jacobsen, J. Kidd, S. B. Lewis, I. J. Øien, A. Sokolov, V.
Sokolov, R. Solheim, S. Weidensaul, K. Wiebe, ..., and K. Safi. 2020. Arctic avian
predators synchronise their spring migration with the northern progression of snowmelt.
Scientific Reports **10**:7220.
Dietz, A. J., C. Kuenzer, and S. Dech. 2015. Global SnowPack: a new set of snow cover
parameters for studying status and dynamics of the planetary snow cover extent. Remote
Sensing Letters **6**:844-853.
Dodge, S., G. Bohrer, R. Weinzierl, S. C. Davidson, R. Kays, D. Douglas, S. Cruz, J. Han, D.
Brandes, and M. Wikelski. 2013. The environmental-data automated track annotation
(Env-DATA) system: linking animal tracks with environmental data. Movement Ecology
1:3.
Ferguson-Lees, J., and D. Christie. 2001. Raptors of the World. Princeton University Press.
Friedl, M., and D. Sulla-Menashe. 2015. MCD12C1 MODIS/Terra+Aqua Land Cover Type
Yearly L3 Global 0.05Deg CMG V006.*in* N. E. L. P. DAAC, editor.
Gallo-Cajiao, E., N. Dolšak, A. Prakash, T. Mundkur, P. G. Harris, R. B. Mitchell, N. Davidson,
B. Hansen, B. K. Woodworth, R. A. Fuller, M. Price, N. Petkov, V. Mauerhofer, T. H.
Morrison, J. E. M. Watson, S. U. Chowdhury, C. Zöckler, O. Widerberg, D. L. Yong, D.
Klich, V. Smagol, J. Piccolo, and D. Biggs. 2023. Implications of Russia's invasion of
Ukraine for the governance of biodiversity conservation. **4**.
Hijmans, R. J. 2016. geosphere: Spherical Trigonometry. R package version 1.5-5.
Hijmans, R. J., and J. van Etten. 2023. raster: Geographic data analysis and modeling.
Hornman, M., A. Boele, and E. van Winden. 2022. Ruigpootbuizerd: steeds schaarser, behalve in
piekjaren. Nature Today.
Jetz, W., G. Tertitski, R. Kays, U. Mueller, M. Wikelski, S. Åkesson, Y. Anisimov, A. Antonov,
W. Arnold, F. Bairlein, O. Baltà, D. Baum, M. Beck, O. Belonovich, M. Belyaev, M.
Berger, P. Berthold, S. Bittner, S. Blake, B. Block, D. Bloche, K. Boehning-Gaese, G.
Bohrer, J. Bojarinova, G. Bommas, O. Bourski, A. Bragin, A. Bragin, R. Bristol, V.
Brlík, V. Bulyuk, F. Cagnacci, B. Carlson, T. K. Chapple, K. F. Chefira, Y. Cheng, N.
Chernetsov, G. Cierlik, S. S. Christiansen, O. Clarabuch, W. Cochran, J. M. Cornelius, I.
Couzin, M. C. Crofoot, S. Cruz, A. Davydov, S. Davidson, S. Dech, D. Dechmann, E.
Demidova, J. Dettmann, S. Dittmar, D. Dorofeev, D. Drenckhahn, V. Dubyanskiy, N.
Egorov, S. Ehnbom, D. Ellis-Soto, R. Ewald, C. Feare, I. Fefelov, P. Fehérvári, W.
Fiedler, A. Flack, M. Froböse, I. Fufachev, P. Futoran, V. Gabyshev, A. Gagliardo, S.
Garthe, S. Gashkov, L. Gibson, W. Goymann, G. Gruppe, C. Guglielmo, P. Hartl, A.
Hedenström, A. Hegemann, G. Heine, M. H. Ruiz, H. Hofer, F. Huber, E. Hurme, F.
Iannarilli, M. Illa, A. Isaev, B. Jakobsen, L. Jenni, S. Jenni-Eiermann, B. Jesmer, F.

Jiguet, T. Karimova, N. J. Kasdin, F. Kazansky, R. Kirillin, T. Klinner, A. Knopp, A.
Kölzsch, A. Kondratyev, M. Krondorf, P. Ktitorov, O. Kulikova, R. S. Kumar, C.
Künzer, A. Larionov, C. Larose, F. Liechti, N. Linek, A. Lohr, A. Lushchekina, K.
Mansfield, M. Matantseva, M. Markovets, P. Marra, J. F. Masello, J. Melzheimer, M. H.
M. Menz, S. Menzie, S. Meshcheryagina, D. Miquelle, V. Morozov, A. Mukhin, I.
Müller, T. Mueller, J. G. Navedo, R. Nathan, L. Nelson, Z. Németh, S. Newman, R.
Norris, O. Nsengimana, I. Okhlopkov, W. Oleś, R. Oliver, T. O'Mara, P. Palatitz, J.
Partecke, R. Pavlick, A. Pedenko, A. Perry, J. Pham, D. Piechowski, A. Pierce, T.
Piersma, W. Pitz, D. Plettemeier, I. Pokrovskaya, L. Pokrovskaya, I. Pokrovsky, M. Pot,
P. Procházka, P. Quillfeldt, E. Rakhimberdiev, M. Ramenofsky, A. Ranipeta, J.
Rapczyński, M. Remisiewicz, V. Rozhnov, F. Rienks, V. Rozhnov, C. Rutz, V. Sakhvon,
N. Sapir, K. Safi, F. Schäuffelhut, D. Schimel, A. Schmidt, J. Shamoun-Baranes, A.
Sharikov, L. Shearer, E. Shemyakin, S. Sherub, R. Shipley, Y. Sica, T. B. Smith, S.
Simonov, K. Snell, A. Sokolov, V. Sokolov, O. Solomina, M. Soloviev, F. Spina, K.
Spoelstra, M. Storhas, T. Sviridova, G. Swenson, P. Taylor, K. Thorup, A. Tsvey, M.
Tucker, S. Tuppen, W. Turner, I. Twizeyimana, H. van der Jeugd, L. van Schalkwyk, M.
van Toor, P. Viljoen, M. E. Visser, T. Volkmer, A. Volkov, S. Volkov, O. Volkov, J. A.
C. von Rönn, B. Vorneweg, B. Wachter, J. Waldenström, N. Weber, M. Wegmann, A.
Wehr, R. Weinzierl, J. Weppler, D. Wilcove, T. Wild, H. J. Williams, J. Wilshire, J.
Wingfield, M. Wunder, A. Yachmennikova, S. Yanco, E. Yohannes, A. Zeller, C.
Ziegler, A. Zięcik, and C. Zook. 2022. Biological Earth observation with animal sensors.
Trends in Ecology & Evolution **37**:293-298.
JIa, Q., K. Koyama, C.-Y. Choi, H.-J. Kim, L. Cao, D. Gao, G. Liu, and A. D. Fox. 2016.
Population estimates and geographical distributions of swans and geese in East Asia
based on counts during the non-breeding season. Bird Conservation International **26**:397-
417.
Kays, R., M. C. Crofoot, W. Jetz, and M. Wikelski. 2015. Terrestrial animal tracking as an eye
on life and planet. Science **348**.
Kays, R., and M. Wikelski. 2023. The Internet of Animals: what it is, what it could be. Trends in
Ecology & Evolution.
Kiat, Y., B. Davaasuren, T. Erdenechimeg, D. Troupin, and N. Sapir. 2021. Large-scale
longitudinal climate gradient across the Palearctic region affects passerine feather moult
extent. Ecography **44**:124-133.
Kjellén, N., M. Hake, and T. Alerstam. 1997. Strategies of two Ospreys Pandion haliaetus
migrating between Sweden and tropical Africa as revealed by satellite tracking. Journal
of Avian Biology:15-23.
Koivurova, T., and A. Shibata. 2023. After Russia's invasion of Ukraine in 2022: Can we still
cooperate with Russia in the Arctic? Polar Record **59**:e12.
Lenth, R., H. Singmann, J. Love, P. Buerkner, and M. Herve. 2019. Package 'emmeans'.
Lislevand, T., S. Hahn, S. Rislaa, and M. Briedis. 2020. First records of complete annual cycles
in water rails Rallus aquaticus show evidence of itinerant breeding and a complex
migration system. Journal of Avian Biology **51**.
Mearns, R., and I. Newton. 1984. Turnover and dispersal in a peregrine Falco peregrinus
population. Ibis **126**:347-355.
Moreau, R. E. 1972. The Palaearctic-African Bird Migration Systems. Academic Press.
National Audubon Society. 2024. Audubon Christmas Bird Count. National Audubon Society.
Newton, I. 2008. The Migration Ecology of Birds. Elsevier, Oxford, UK.
Pokrovsky, I., D. Ehrich, I. Fufachev, R. A. Ims, O. Kulikova, A. Sokolov, N. Sokolova, V.
Sokolov, and N. G. Yoccoz. 2019. Nest association between two predators as a
behavioral response to the low density of rodents. The Auk **137**.

Pokrovsky, I., D. Ehrich, R. Ims, O. Kulikova, N. Lecomte, and N. Yoccoz. 2014. Diet, nesting
density, and breeding success of rough-legged buzzards (Buteo lagopus) on the Nenetsky
Ridge, Arctic Russia. Polar Biology **37**:447-457.
Pokrovsky, I., D. Ehrich, R. A. Ims, A. V. Kondratyev, H. Kruckenberg, O. Kulikova, J.
Mihnevich, L. Pokrovskaya, and A. Shienok. 2015. Rough-Legged Buzzards, Arctic
Foxes and Red Foxes in a Tundra Ecosystem without Rodents. PLoS ONE **10**:e0118740.
Pokrovsky, I., A. Kölzsch, S. Sherub, W. Fiedler, P. Glazov, O. Kulikova, M. Wikelski, and A.
Flack. 2021. Longer days enable higher diurnal activity for migratory birds. Journal of
Animal Ecology **90**:2161-2171.
Posit team. 2022. RStudio: Integrated Development Environment for R. Posit Software, PBC,
Boston, MA.
R Development Core Team. 2022. R: A language and environment for statistical computing. R
Foundation for Statistical Computing, Vienna, Austria.
Reif, J. 2013. Long-term trends in bird populations: a review of patterns and potential drivers in
North America and Europe. Acta ornithologica **48**:1-16.
Robinson, R. A., N. A. Clark, R. Lanctot, S. Nebel, B. Harrington, J. A. Clark, J. A. Gill, H.
Meltofte, D. I. Rogers, and K. G. Rogers. 2005. Long term demographic monitoring of
wader populations in non-breeding areas. Wader Study Group Bulletin **106**:17-29.
Schlaich, A. E., V. Bretagnolle, C. Both, B. J. Koks, and R. H. Klaassen. 2023. On the Wintering
Ecology of Montagu's Harriers in West Africa: Itinerancy in Relation to Varying Annual
Environmental Conditions. Ardea **111**:321-342.
Serrano, D., J. L. Tella, M. G. Forero, and J. A. Donazar. 2001. Factors affecting breeding
dispersal in the facultatively colonial lesser kestrel: individual experience vs. conspecific
cues. Journal of Animal Ecology **70**:568-578.
Sonerud, G. A. 1986. Effect of snow cover on seasonal changes in diet, habitat, and regional
distribution of raptors that prey on small mammals in boreal zones of Fennoscandia.
Ecography **9**:33-47.
Strandberg, R., T. Alerstam, M. Hake, and N. Kjellen. 2009. Short-distance migration of the
Common Buzzard Buteo buteo recorded by satellite tracking. Ibis **151**:200-206.
Sundell, J., O. Huitu, H. Henttonen, A. Kaikusalo, E. Korpimaki, H. Pietiainen, P. Saurola, and I.
Hanski. 2004. Large-scale spatial dynamics of vole populations in Finland revealed by
the breeding success of vole-eating avian predators. Journal of Animal Ecology **73**:167-
178.
Tast, J., A. Kaikusalo, and M. Lagerström. 2010. Diet composition of breeding rough-legged
buzzards Buteo lagopus at Kilpisjärvi, NW Finnish Lapland. Kilpisjärvi Notes **22**:10-18.
Terraube, J., A. Villers, L. Ruffino, L. Iso-Iivari, H. Henttonen, T. Oksanen, and E. Korpimäki.
2015. Coping with fast climate change in northern ecosystems: mechanisms underlying
the population-level response of a specialist avian predator. Ecography **38**:690-699.
Thorup, K., A. P. Tøttrup, M. Willemoes, R. H. G. Klaassen, R. Strandberg, M. L. Vega, H. P.
Dasari, M. B. Araújo, M. Wikelski, and C. Rahbek. 2017. Resource tracking within and
across continents in long-distance bird migrants. Science advances **3**:e1601360.
Trierweiler, C., W. C. Mullié, R. H. Drent, K. M. Exo, J. Komdeur, F. Bairlein, A. Harouna, M.
de Bakker, and B. J. Koks. 2013. A Palaearctic migratory raptor species tracks shifting
prey availability within its wintering range in the Sahel. Journal of Animal Ecology
82:107-120.
Vansteelant, W., W. Faveyts, and J. Buckens. 2011. Opmerkelijke Ruigpootbuizerdinvasie in de
winter 2010-2011: oorzaken in een historische en Europese context. Natuur.oriolus
77:87-98.
Walker, D. A., M. K. Raynolds, F. J. A. Daniels, E. Einarsson, A. Elvebakk, W. A. Gould, A. E.
Katenin, S. S. Kholod, C. J. Markon, E. S. Melnikov, N. G. Moskalenko, S. S. Talbot, B.

A. Yurtsev, and C. Team. 2005. The Circumpolar Arctic vegetation map. Journal of
Vegetation Science **16**:267-282.
Wetlands International. 2024. International Waterbird Census.
Whitfield, D. P., K. Duffy, D. R. McLeod, R. J. Evans, A. M. MacLennan, R. Reid, D. Sexton, J.
D. Wilson, and A. Douse. 2009. Juvenile dispersal of white-tailed eagles in western
Scotland. Journal of Raptor Research **43**:110-120.
Wikelski, M., R. W. Kays, N. J. Kasdin, K. Thorup, J. A. Smith, and G. W. Swenson. 2007.
Going wild: what a global small-animal tracking system could do for experimental
biologists. Journal of Experimental Biology **210**:181-186.
Yanco, S. W., R. Y. Oliver, F. Iannarilli, B. S. Carlson, G. Heine, U. Mueller, N. Richter, B.
Vorneweg, Y. Andryushchenko, N. Batbayar, M. Dagys, M. Desholm, B. Galtbalt, A. E.
Gavrilov, O. A. Goroshko, E. I. Ilyashenko, V. Y. Ilyashenko, J. Månsson, E. A. Mudrik,
T. Natsagdorj, L. Nilsson, S. Sherub, H. Skov, T. Sukhbaatar, R. Zydelis, M. Wikelski,
W. Jetz, and I. Pokrovsky. 2024. Migratory birds modulate niche tradeoffs in rhythm
with seasons and life history. Proceedings of the National Academy of Sciences.

a)
Breeding period

Q Slow phase **Q**

Latitude

Jul Sep Nov Jan Mar May

Qf S1 S2 Qs

1st part 2nd part

b)

Quick phase   **Slow** phase
spring —yellow—   part 1 —blue—
fall —orange—   part 2 —dark blue—

c)

Distance (km*1000)   Duration (days)   Speed (km/day)

Qs Qf S1 S2
Migration phase

Qs Qf S1 S2
Migration phase

Qs Qf S1 S2
Migration phase

N W E S

a)

Vegetation cover (%)

Quick phase

Grass-lands | Crop-lands | Forests | Urban | Shrub-lands

b)

Vegetation cover (%)

Slow phase

Grass-lands | Crop-lands | Forest | Urban | Shrub-lands

c)

Quick phase
- spring
- fall

Slow phase
- part 1
- part 2

a)

breeding period — migration — **residency** — migration — breeding period

Latitude

VIII IX X XI XII I II III IV V VI VII

breeding range

unsuitable habitat
migration

static winter range
residency

b)

breeding period — migration — **itinerancy** — migration — breeding period

Latitude

VIII IX X XI XII I II III IV V VI VII

breeding range

unsuitable habitat
migration

XI, XII

I, II

III, IV

dynamic winter range
itinerancy

Numbers indicate months the birds are in the area

c)

breeding period — **foxtrot migration** — breeding period

quick phase | **slow** phase | quick phase

1st part | 2nd part

Latitude

VIII IX X XI XII I II III IV V VI VII

breeding range

unsuitable habitat
foxtrot migration | **quick** phases

X, IV

XI, II, III

XII, I

dynamic winter range
foxtrot migration | **slow** phase

Numbers indicate months the birds are in the area

a)

For species with **dynamic winter range**, when mid-winter counts are used to determine winter range temporal heterogeneity can be mistaken with spatial heterogeneity

uncommon
common

For species with **dynamic winter range**, when GPS tracking is used to determine winter range temporal heterogeneity becomes evident and should be shown on the map

X, IV
XI, II, III
XII, I

Numbers indicate months the birds are in the area

b)

abundance decline within a specific part of the winter range

Species abundance

Years

Extinct   Threatened   Least Concern
EX EW   CR EN VU   NT LC

Years

For species with **dynamic winter range**, a decline in abundance within a specific part of the winter range may indicate changes in a range dynamics rather than an overall population decrease