Predictive reward-prediction errors of climbing fiber inputs integrate modular reinforcement learning with supervised learning

Huu Hoang[1*#], Shinichiro Tsutsumi[2*], Masanori Matsuzaki[3], Masanobu Kano[4,5], Keisuke Toyama[1], Kazuo Kitamura[6#], Mitsuo Kawato[7#]

[1] ATR Neural Information Analysis Laboratories, Japan

[2] RIKEN Center for Brain Science, Japan

[3] Department of Physiology, Graduate School of Medicine, The University of Tokyo, Japan

[4] Department of Neurophysiology, Graduate School of Medicine, The University of Tokyo, Japan

[5] International Research Center for Neurointelligence (WPI-IRCN), The University of Tokyo, Japan

[6] Department of Neurophysiology, Graduate School of Interdisciplinary Research, University of Yamanashi, Japan

[7] ATR Computational Neuroscience Laboratories, Japan

* These authors contributed equally to this work

Correspondence: Huu Hoang (hoang@atr.jp), Kazuo Kitamura (kitamurak@yamanashi.ac.jp) and Mitsuo Kawato (kawato@atr.jp)

Abstract

Although the cerebellum is typically linked to supervised learning algorithms, it also exhibits extensive connections to reward processing. In this study, we investigated the cerebellum's role in executing reinforcement learning algorithms, with a particular emphasis on essential reward-prediction errors. We employed the Q-learning model to accurately reproduce the licking responses of mice in a Go/No-go auditory-discrimination task. This method enabled the calculation of reinforcement learning variables, such as reward, predicted reward, and reward-prediction errors in each learning trial. By tensor component analysis of two-photon Ca^{2+} imaging data, we found that climbing fiber inputs of the two distinct components, which were specifically activated during Go and No-go cues in the learning process, showed an inverse relationship with predictive reward-prediction errors. Given the hypothesis of bidirectional parallel-fiber Purkinje-cell synaptic plasticity, Purkinje cells in these components could develop specific motor commands for their respective auditory cues, guided by the predictive reward-prediction errors from their climbing fiber inputs. These results indicate a possible role of context-specific actors in modular reinforcement learning, integrating with cerebellar supervised learning capabilities.

Introduction

Historically, the cerebellum has been viewed as implementing supervised learning algorithms, a perspective provided by seminal works (Marr, 1969; Albus, 1971; Ito ,1969; Kawato et al., 1987; Kawato and Gomi, 1992; Wolpert et al. 1998; Kawato, 1999). Simultaneously, the basal ganglia's dopaminergic neurons have been associated with reinforcement learning algorithms (Schultz et al. 1997; Bayer and Glimcher 2005; Haruno and Kawato, 2006; Pessiglione et al., 2006; Akiti et al. 2022; Amo et al., 2022). This understanding has been bolstered by a substantial body of evidence (Shidara et al., 1993; Medina et al., 2000; Medina and Lisberger, 2008; Haruno et al. 2004; Doya 2000; Kaplan et al., 2020; Dabney et al. 2020; Starkweather and Uchida, 2020; Kawato and Samejima, 2007; Raymond and Medina, 2018; Kawato et al., 2020). However, recent research has identified reward-related variables in both the mossy-fiber-granule cell pathway (Wagner et al., 2017; Sendhilnathan et al., 2020) and climbing fibers (Heffley et al., 2018; Heffley and Hull 2019; Kostandinov et al., 2019; Larry et al., 2019; Sendhilnathan et al., 2021), distinct from movement-related information (Heffley et al., 2018; Larry et al., 2019). Intriguingly, the association between cerebellar activities and reward variables is dependent on the cerebellar cortex's zonal organization (Kostandinov et al., 2019; Tsutsumi et al., 2019; Hoang et al., 2023). Anatomical studies further underscore reciprocal connectivity between the cerebellum and the reward circuitry (Bostan and Strick, 2018; Carta et al., 2019; Wagner et al., 2019; Chabrol et al., 2019). These developments point to a potential role of the cerebellum in reward processing and reinforcement learning tasks driven by reward and penalty, as reviewed in recent literature (Wagner and Luo, 2020; Kawato et al., 2021; Kostadinov and Häusser, 2022). Yet, it remains unclear if the cerebellum encodes essential reinforcement learning variables, such as reward-prediction errors (Dabney et al. 2020; Amo et al., 2022; Starkweather and Uchida, 2020), or if its unique zonal structure plays a role in modular reinforcement learning algorithms (Doya et al., 2002; Bentivegna et al., 2004; Kawato et al., 2007, 2021; Sugimoto, Haruno et al., 2012; Sugimoto, Morimoto et al., 2012).

In the present study, we revisited two-photon Ca^{2+} imaging data of climbing fibers from eight aldolase-C zones in the Crus II region (Sugihara and Shinoda, 2004, 2007) of the mouse cerebellum during a Go/No-go auditory-discrimination licking task (Tsutsumi et al., 2019). Given the nature of this task as a reinforcement-learning driven by rewards and penalties, we applied a formal reinforcement learning algorithm, Q-learning (Watkins and Dayan, 1992), to reproduce licking responses, as well as to compute reinforcement learning variables such as rewards, predicted rewards, and reward-prediction errors in each learning trial. For the analysis of the two-photon Ca^{2+} imaging data of climbing fibers, we employed a hyper-resolution spike timing estimation algorithm (Hoang et al., 2020) and tensor component analysis (Williams et al., 2019) to identify four key components impacting the learning process (Hoang et al., 2023). Our results indicated that the inputs from climbing fibers to the first two components, which show bi-directional modulation during learning, strongly and negatively correlated with predictive reward-prediction errors. Notably, this correlation was specific to the cue-response condition in which each component was engaged. Considering the bidirectional synaptic plasticity of parallel-fiber Purkinje cell synapses (Ito and Kano, 1982; Hirano 1990; Ito 2001; Linden 1999; Hirano, 2013), the Purkinje cells in these two components could work as context-specific actors (Barto et al., 1983) within modular reinforcement learning algorithms. Essentially, the functional modules of Purkinje cells aligned roughly with aldolase-C zones could develop distinct and context-specific motor commands based on predictive reward-prediction errors relayed by climbing fiber inputs in the framework combining cerebellar supervised learning with modular reinforcement learning (Kawato et al., 2021).

Results

Q-learning model of licking behavior

We employed a Q-learning algorithm to model licking behavior of mice in the Go/No-go experiment (Fig 1A). The Q-learning model was selected because it is one of the simplest reinforcement learning algorithms, based on state-action value functions rather than state value functions, with a small number of

hyper-parameters. Briefly, mice were trained to distinguish two auditory cues (high versus low frequency tones) by either licking or not licking during a response period of 1s after cue onset to obtain a liquid reward. The reinforcement-learning-algorithm state s is determined by the two auditory cues (s = Go / No-go), and the reinforcement-learning-algorithm action a is determined by licking during a time window of [0, 1] s after the auditory cue (a = Lick / No-lick). A reward value R-P was assigned according to cue-response conditions, including HIT trials (lick after Go cue) in Go task (R-P = 1), false alarm trials (FA, lick after No-go cue, R-P $= -\xi$, $0 \leq \xi \leq 1$), correct rejection trials (CR, no lick after No-go cue, R-P = 0) and MISS trials (no lick after Go cue, R-P = 0). Note that the R-P value for FA trials was constrained to be negative because a lick after a No-go cue was punished with a timeout of 4.5 s, while that of CR and MISS trials were 0 because neither reward nor penalty was given in those trials. The Q-learning algorithm assumed that a reward prediction Q was computed as a function of the two variables s and a; *state and action*. In each trial, a reward prediction error δQ, a difference between reward R-P and Q, was used to update computation of Q in the next trial with a learning rate α. For an action policy, we used the softmax function with a single-temperature parameter τ to convert Q values into a probability distribution over the two possible actions; Lick and No-lick. Since mice had undergone pre-training sessions lasting 3 days, when they were always rewarded for a lick within 1 second for both cues, initial Q values for both Go and No-go cues were positive ($0 \leq q_1 \leq 1$, $0 \leq q_2 \leq 1$, for Go and No-go cues respectively, see Methods for details). It is important to note that we did not explicitly model the time course within trial or temporal difference errors as in previous studies (e.g., Schultz et al., 1997). Instead, we assumed a single timing of cue representation for computation of Q and δQ, which precedes the timing of reward delivery by about 500~2,000 ms. Thus, the reward-prediction error δQ in this study is a predictive signal estimated before the actual delivery of liquid rewards. We will later discuss implications of this assumption on timing for possible neural computations. Briefly, we show that the cerebellar reinforcement learning in this study could acquire the actor but not the critique through temporal difference error.

Behavioral data indicated that the lick rate in HIT trials was high in the early period (average and standard deviation of lick latency, 0.25 ± 0.15 s, Fig 1B) and extended over the reward delivery period (liquid rewards were delivered three times at 0.41, 0.82, and 1.23 s after the first lick, thus 0.66, 1.07 and 1,48 s after the cue on average). In contrast, the lick rate in FA trials was also high in the early period (average and standard deviation of lick latency, 0.31 ± 0.23 s), but it gradually reduced to baseline because no reward was given for No-go cues. We fitted the Q-learning model to the licking behavior of individual mice (n = 17) in 26,517 trials. Fitting performance was good for both Go and No-go trials, which showed an increase in fraction correct and a decrease in fraction incorrect, respectively, at both the population (Fig 1C) and individual (Fig 1D) levels (average and standard deviation of coefficients of determination of 17 mice, 0.87 ± 0.15 for Go cue and 0.61 ± 0.18 for No-go cue, respectively, see Methods for details). The hyper-parameters estimated for individual mice were broadly distributed (average and standard deviation; 0.002 ± 0.002 for α, 0.12 ± 0.07 for q_1, 0.24 ± 0.17 for q_2, 0.14 ± 0.07 for τ, and 0.84 ± 0.24 for ξ, Fig 1E), indicating that each animal utilized a distinct strategy for optimally learning to obtain the reward. As temporal evolution of the reward-related variables with discrimination learning, Q values for both cues and lick (s=Go / No-go and a=Lick) were positive and intermediate initially, then increased for HIT trials (s=Go and a=Lick) and decreased toward negative values for FA trials (s=No-go and a=Lick) as learning progressed (Fig 1F). Note that the negative Q value for FA trials at the later stage of learning accurately reflected the negative R-P value assigned for those trials by the Q-learning model. As a consequence, δQ values converge to zero for both cue-response conditions (Fig 1G). More specifically, the δQ values for HIT trials (s=Go and a=Lick) were positive initially and monotonically decreased during learning. By contrast, the initial δQ value for FA trials (s=No-go and a=Lick) was negative and large, because of a large difference in negative reward R-P and the initial positive value of reward prediction $q_2 > 0$. Throughout the course of learning, this δQ value monotonically increased (decreased its magnitude) to zero, indicating a better agreement between negative R-P and Q values. For CR and MISS trials, both Q and δQ remained constant at zero (Fig 1F, G).

Figure 1: Q-learning model of a Go/No-go experiment. A: schematic of the Q-learning model. B: the averaged lick rate for the four cue-response conditions (blue, red, orange and magenta for HIT, FA, CR and MISS trials, respectively: see also inset for color codes). Solid and dashed vertical lines indicate the cue onset and response window (1s after cue), respectively. Light cyan shading represents the window for possible reward delivery (0.5 - 2 s after cue). C-D: fraction correct for Go cue (blue) and fraction incorrect for No-go cue (red) for experimental data (solid lines) and Q-learning model (dashed lines), averaged for 17 mice in 7 training sessions (C) and for individual mice (D). Vertical bars in C show standard errors. E: hyperparameter values of the Q-learning model estimated for individual mice (1~17); and from left to right and top to bottom, learning rate α, initial Q values for Go and No-go cues, q_1 and q_2, respectively, temperature τ, and punishment value for FA trials ξ. F-G: evolution of Q (F) and δQ (G) of a representative mouse for the four state-action combinations (HIT, FA, CR, MISS) during the course of learning.

Zonal complex spike firings and their correlations with reward prediction error

While mice learned Go/No-go discrimination tasks, we conducted two-photon recordings of climbing fibers (sampling rate, 7.8 Hz) from 6,445 Purkinje cells in eight cerebellar zones (from 7+ to 4b-, see Methods for details). Please note that the experimental data analyzed in this study is identical to those of the previous work (Hoang et al. 2022), and contains those of (Tsutsumi et al. 2019) as a part. A hyper-resolution algorithm (HA_time, Hoang et al. 2020) was applied to estimate timing of complex spikes (CSs) at a resolution of 100 Hz. Similar to the previous work (Hoang et al. 2022), we studied CS activity as population peri-stimulus time histograms (PSTHs) sampled in three learning stages (from top to bottom, 1st, 2nd, and 3rd stages with fraction correct <0.6, 0.6 - 0.8, 0.8<, respectively, Fig 2A) for the four cue-response conditions, or the corresponding four state-action combinations. Briefly, CSs in HIT trials (n = 3,788) were large and distributed across the entire medial hemisphere at the initial learning stage, but later on (2nd and 3rd stages), they became stronger and compartmentally focused on positive zones. In contrast, PSTHs in FA trials (n = 1,757) were distributed across almost the entire Crus II and gradually decreased and were more confined to lateral zones along with learning. PSTHs for CR trials (n = 2,229) were mainly distributed within zones 6- and 6+, and there was only spontaneous CS activity in MISS trials (n = 201).

Our previous study demonstrated that CSs in the lateral hemisphere represent cognitive error signals related to No-go cues (Hoang et al., 2022). Therefore, in the present study, we analyzed trial-by-trial correlations between CSs in zones 6-, 6+ and 5- with the reward prediction error δQ for FA trials. As expected, on a trial basis, we found that CSs, defined as the mean firing rate in 0 - 0.5 s after cue onset, in these three zones were negatively correlated with δQ, with the strongest correlation found for zone 5- (slope = -0.53, $p < 0.01$ for 6-, slope = -0.60, $p<0.001$ for 6+ and slope = -0.76, $p < 0.0001$ for 5-, Fig 2B-D). The temporal window of 0 - 0.5 s for calculation of the mean firing rate was selected because CSs in FA trials of these three zones were mostly confined within 0 - 0.5 s after the cue onset (Fig 2A).

Figure 2: CS activities to cues and their correlation with reward prediction error. A: Heat-maps showed PSTHs of climbing fiber firings in 8 Ald-C zones (7+ to 4b-) in the four cue-response conditions (columns) and three learning stages (rows, 1st, 2nd, and 3rd stages with fraction correct <0.6, 0.6 - 0.8, 0.8<, respectively). The blue line indicates cue onset. B-D: correlation of firing activity of Ald-C zones 6- (B), 6+ (C) and 5- (D) with δQ in the FA trials. Each dot represents a trial of 17 mice. Figure 2A was reproduced from (Hoang et al. 2023).

Partial Least-Squares regression of zonal activity with reinforcement-learning and sensorimotor-control explanatory variables

The previous analysis was constrained to only three zones and δQ. Here, we systematically studied correlations between the firing activity of neurons in all eight cerebellar zones with rewards, as well as sensorimotor variables, by partial least-squares regression (PLS regression) to find meaningful correlations under the situation of multicollinearity between explanatory variables (range of correlations between explanatory variables, 0.12 ± 0.23 with the highest correlations found for R vs. Q and R vs. δQ as 0.82 and 0.78, respectively). For this purpose, neuronal activity in each trial was defined as the mean firing rate in [-0.5, 2] s after cue onset of neurons in the same Ald-C zone. For explanatory variables, we included R, Q, δQ as reward variables, and lick count in the three response windows ([0, 0.5] s for early lick - ELick, [0.5, 2] s for reward lick - RLick and [2, 4] s for late lick - LLick) for Go and No-go cues, separately, as sensorimotor variables. Here, we prepared 6 sensorimotor variables (2 ⬜ 3=6; Go vs No-go multiplied by three response windows of licks) (Fig. 3, inset). Note that these three response windows correspond well with licking behavior of mice, as well as reward delivery period (Fig 1B). Because these sensorimotor variables span large temporal windows of 0 - 4 s after the cue, the mean firing rate was computed for the comparable large temporal window of [-0.5, 2] s after the cue onset. We further note that the physical reward R (R=1 for HIT trials, R=0 otherwise), which was used as one of explanatory variables of PLS regression, is different from the reward-punishment R-P used in the Q-learning model (see Methods for details).

We calculated the variable importance in projection scores (VIP scores) to quantitatively estimate the importance of a given explanatory variable to the PLS regression model. The VIP score does not provide the sign (positive or negative) of correlations, but explanatory variables, whose VIP scores are larger than 1, are generally considered important in PLS regression (see Methods for details). We found that neurons in the lateral hemisphere were associated with Q and δQ, but not the reward R per se (Fig 3). More specifically, 6- and 6+ were strongly associated with Q (VIP score, 1.8 and 1.7 for 6- and 6+,

respectively), while 6- and 5- were strongly associated with δQ (VIP score = 1.6 and 2.1 for 6- and 5-, respectively). Note that these results are consistent with results of simple correlation analysis (Fig 2B-D). In contrast, neurons in the medial hemisphere (5+ to 4b-) were strongly associated with the reward R (VIP score > 1.2). Notably, we also found a complicated association between zonal activity and sensorimotor variables, i.e., licking following two cues. That is, medial zones were associated with the lick number in the reward period (Go ⍰ RLick and No-go ⍰ RLick, VIP score > 1) while those in the lateral hemisphere (7+ and 5-) were associated with the lick number in the reward period following No-go cues (No-go ⍰ RLick, VIP score = 1.1 for both 7+ and 5-, respectively). Only zone 7+ was associated with late lick following Go cues (Go ⍰ LLick, VIP score = 1.1).

Figure 3: Partial least squares regression of spiking activity and reinforcement-learning and sensorimotor-control variables. Bars showed the variable-importance-in-prediction (VIP) scores of 9 reinforcement-learning and sensorimotor-control variables (from left to right, R, Q, δQ, Go ⍰ ELick, No-go ⍰ ELick, Go ⍰ RLick, No-go ⍰ RLick, Go ⍰ LLick, No-go ⍰ LLick) for spiking activity of neurons in 8 Ald-C zones. Dashed lines indicated VIP score = 1, which is considered a threshold of importance. See the inset for color codes of the 9 explanatory variables.

The generative model of spiking activity at a trial basis by tensor component analysis

The PLS regression shown in Fig 3 suggested a functional organization of CSs, moderately constrained by the zonal structure, with respect to sensorimotor and reward processing. In our previous study, we conducted tensor component analysis (TCA) for CS PSTHs of >6,000 PCs and found four well separated components (TC1-4) that explained more than 50% of variance of PSTHs (Fig 4A, see Hoang et al. 2023 for details). One of the reasons that relatively moderate anatomical distributions of different functions are found in Fig 3 is that each zone and even each neuron contain multiple functional components, as demonstrated in Hoang et al. (2023) as well as in previous studies demonstrating multiplexed representations (Markanday et al. 2021; Ikezoe et al. 2023). In order to overcome this difficulty due to multiplexing for revealing precise functions of each component, we next examined functional representations of each tensor component utilizing Q-learning and trial-based analyses, while incorporating timing information of each spike instead of broadly computing the average spike rate over a wide temporal window.

In this study, we used TCs as a generative model of spiking activity as a trial basis to reveal associations of functionally organized CSs and reinforcement-learning and sensorimotor-control variables. Briefly, the TC score of a neuron in a particular trial was estimated by filtering the spike train of that neuron by the temporal profile of the corresponding TC, weighted by neuronal and cue-response condition coefficients (Fig 4B, see Methods for details). This computation incorporated trial-to-trial variability of spiking activity while maintaining fundamental properties of TCs. For example, neurons, whose TC1 and TC2 coefficients were highest among all neurons recorded, had high TC1 and TC2 activities in HIT and FA trials, respectively (Fig 4C-D). Note that TC1 and TC2 were selectively activated in HIT and FA trials, respectively (Fig 4A). Following such a computation, the TC score, averaged TC activity of all neurons in the same recording session, shared a similar structure of zonal distribution and cue-response condition with those of the TCs (compare Fig 4A and Fig 4E, but note that abscissae are learning stages and cue-

response conditions, respectively). Specifically, TC1 scores were high in HIT trials and for positive zones. In contrast, TC2 scores were high in FA trials and distributed in the lateral hemisphere. TC3 scores were high in HIT trials and distributed in the medial hemisphere. TC4 scores had similar zonal distribution with TC2 scores, except that they were non-zero only for CR trials.

Figure 4: Tensor-component analysis (TCA) and computation of tensor score at a trial-basis. A: TCA was conducted for PSTHs in 4 cue-response conditions of n=6,445 neurons and the resulting four tensor components (TC1-4) explained more than 50% of variance. B: for the i-th single neuron, its activity for the r-th TC (y^r) in the particular j-th trial was computed by filtering spike timings with temporal profile of the r-th TC b^r_t, multiplying corresponding coefficients w^r_i of the i-th neuron and a^r_c of the cue-response condition c. C-D: PSTHs (C) of two representative neurons, which have the highest coefficients of TC1 and TC2, respectively, and their TC1 and TC2 scores, respectively, computed for all trials in their corresponding sessions (D). E: Heatmaps showed TC1-4 scores averaged for all neurons in each of the eight zones distinctively for the four cue-response conditions

Sparse canonical correlation analysis

To find the variables that contribute the most to each TC score, we conducted sparse canonical correlation analysis (sCCA) of TC scores and the same 9 reinforcement-learning and sensorimotor-control variables used in PLS regression (see Methods for details). As a result, TC1 and TC3 were associated with only reward variables while TC2 and TC4 were associated with both reward and sensorimotor variables in No-go trials (Fig 5A). More specifically, TC1 and TC3 were positively correlated with reward R, reward prediction Q and also reward prediction error δQ, with high coefficients of R and Q for TC1 (0.66 and 0.62) and R for TC3 (0.78). We can safely state that TC1 is mainly related to reward and its prediction, and that TC3 is mainly related to reward. Remarkably, TC2 was negatively correlated with δQ (coefficient, -0.83) but it was positively correlated with the early lick count in No-go trials (0.42). Similarly, TC4 was negatively correlated with both R (coefficient, -0.65) and early lick count in No-go trials (-0.59). We note that those associations revealed by sCCA can also be found by PLS regression (Fig S1); thus, they are not dependent on specific analysis methods. We further confirmed significant correlation for each TC component by linear regression of TC1 vs. Q (slope = 0.46, p < 0.0001, Fig 5B), TC1 vs. δQ (slope = 0.35, p < 0.0001, Fig 5C), TC2 vs. δQ (slope = -0.42, p < 0.0001, Fig 5D), TC3 vs. R (slope = 0.30, p < 0.0001, Fig 5E) and TC4 vs. No-go ⬚ ELick (slope = -0.24, p < 0.0001, Fig 5F). Notably, these correlations were also significant (p < 0.0001) with comparable slopes, even when using only trials of cue-response condition with which each TC is primarily associated (slope = 0.32 for TC1-HIT, slope = -1.24 for TC2-FA, and slope = -0.36 for TC4-CR trials, Figs 5B, D, F). The most interesting exception was obtained for TC1 vs. δQ for HIT trials. The correlation was strongly significant (slope = -0.65, p < 0.0001, Fig 5C), but the correlation was negative for only HIT data (slope = -0.65), while the overall correlation was positive as shown above (slope = 0.35).

Figure 5: Sparse canonical-correlation analysis (sparse CCA) of TC scores with reinforcement-learning and sensorimotor-control variables. A: Bars show coefficients of reinforcement-learning and sensorimotor-control variables corresponding to TC1-4 scores. B-F: the scatter plots of trials show correlations of TC1 with Q (B), TC1 with δQ (C), TC2 with δQ (D), TC3 with R (E) and TC4 with No-go ⬚ ELick (F) at trial-basis. Black and gray lines indicate regression between variables when using all trials and trials of the cue-response condition with which each TC is primarily associated, i.e., TC1-HIT, TC2-FA and TC4-CR, respectively. Panel E shows the boxplot with gray lines indicating the median and the bottom and top edges of the box the 25th and 75th percentiles, respectively. All correlations in B-F are significant (p<0.0001). Color convention of trials is the same as Figure 1. The inset of A shows color codes of the selected 6 reward and sensorimotor variables among 9 according to sCCA.

Discussion

In this study, we conducted three correlation analyses to elucidate contributions of CSs to behavior during learning of Go/No-go auditory-discrimination licking tasks. The first analysis showed that lateral zones (6-, 6+, and 5-) are associated with reward-prediction error, confirming our earlier findings of cognitive

error signals in these zones (Hoang et al., 2023). Next, building upon distinct contributions of zonal-organized CF inputs to behavioral variables (Tsutsumi et al., 2019), we examined the firing rate of neurons across all eight cerebellar zones against nine reinforcement-learning and sensorimotor-control variables, using partial least-squares regression to handle multicollinearity. This analysis indicated a moderate distribution of variables across zones, reflecting the multifunctional nature of each zone and neuron (Markanday et al., 2021; Ikezoe et al., 2023; Hoang et al., 2023). The third analysis used tensor component analysis to decompose spiking activity into four tensor-components, examining their trial-based functional representations. We employed different complex spike measures across the analyses, ranging from mean firing rates to precise spike timings. Our methods evolved from focusing on a few zones to covering all eight, and from basic linear regression to advanced techniques like sparse canonical correlation analysis. Despite these methodological differences, our findings were consistent. We found that TC1 neurons, distributed most densely in 6+, were positively correlated with reward R, reward prediction Q, and reward prediction error δQ (Figs 2C, 3(6+), 5B&C). However, we note that TC1 is negatively correlated with δQ if constrained only for HIT trials (Fig. 5C) and the negative correlation was also demonstrated in Fig 2C for 6+ zone. In contrast, TC2 neurons in zones 6- and 5- were negatively correlated with reward prediction error δQ (Figs 2B, 2D, 3(6- and 5-) and 5D). TC3 neurons in medial zones were positively correlated with reward R and licking in reward delivery period (Figs 3(5+,5a-,5a+,4b-) and 5E). Finally, TC4 neurons, distributed most densely in 6-, were negatively correlated with both reward R and early licking following No-go cues (Figs 3(6-) and 5A and F). Although we excluded the reward-penalty variable (R-P) from regression analyses due to high correlation with other reward variables, including it in the analysis did not alter our overall findings (Fig S2).

Notably, these results were also in good agreement with our recent work. In Hoang et al. (2023), we found that TC1-4 corresponds to timing control of the first lick, cognitive error signals, reward-related signals and action inhibition, respectively. Here, we showed that TC2 was negatively correlated with reward prediction error δQ; thus, cognitive error signals with a positive sign can be computed as sign-

reversed reward prediction errors with a negative sign by reinforcement learning algorithms. That is, during learning, cognitive error signals decrease due to an increase (with the same magnitude) of negative reward prediction errors. Similarly, our previous work indicated TC4 neurons inhibit licks in No-go trials. This is reinforced by sparse CCA's findings of TC4's negative correlation with No-go ⬚ ELick. Since R and Q values for No-go cues were below average and declined during learning, the increase in TC4 activity is consistent with its negative correlations with reward R and prediction Q (Fig S3). Both studies corroborate TC3's correlations with reward R and reward licking behavior. However, a discrepancy exists in TC1's role: Hoang et al. (2023) found TC1's synchronized spikes within [0, 0.5] sec positively correlated with the precise timing of the first lick in HIT trials, which wasn't observed here. There are two possible reasons for this inconsistency. First, in this study, we investigated correlations in all cue-response conditions and not specifically in HIT trials, while our previous study examined only HIT trials for TC1. Second, we did not utilize synchronization of CSs in this study, which is important for timing control (Welsh et al., 1995; Tsutsumi et al., 2020; Wagner et al., 2021), while our previous study examined only synchronized spikes. Because R and Q for Go cues were higher than their averages and increased during learning, positive correlations of TC1 with these two reward-related variables are expected due to an increase of TC1 activity (Fig S3).

Modular reinforcement learning characteristics

In a supplementary analysis, our objective was to separate the roles of complex spikes (CSs) in cue-cognition and motor functions. We included an auditory cue variable (Cue=1 for Go, Cue=0 for No-go) distinct from motor-related variables (Fig S4). While PLS analysis of zonal activity yielded results similar to previous findings (Fig S4A), a notable shift was observed in TC4's response: a reduced negative correlation with early licks following No-go cues, but a stronger negative correlation with auditory cues (Fig S4B). This suggests that TC4 neurons primarily inhibit early licks in response to No-go cues, rather than acting as general motor commands for all auditory cues. These findings, along with previous research (Heffley et al. 2018; Larry et al. 2019), indicate that neither a single zone nor a tensor component

(TC) solely encodes basic licking motor commands without regard to auditory cues. If CSs in any zone or TC were purely associated with motor control of licking, CF responses would be similar between HIT and FA trials, and between MISS and CR trials, but our results show significant differences (Figs. 2A, 4A, and E). Thus, the cerebellar reinforcement learning algorithm investigated in this study functions as a modular system: TC1 and TC3 for Go cues, and TC2 and TC4 for No-go cues, rather than a simplistic, straightforward algorithm. In line with our proposed modular framework, CF inputs in TC1 and TC2 distinctly mediate reward-prediction errors for Go and No-go cues, respectively. TC2 consistently demonstrates a strong negative correlation with reward prediction error δQ in false alarm (FA) trials across all analyses (Figs 2B, 2D, 3(6- and 5-), 5D, and S4). Similarly, TC1 neurons' representation of reward-prediction error is significant (Fig 5A, C), especially in 36% of these neurons (Fig S6). When focusing on HIT trials, TC1 shows robust negative correlations with reward-prediction errors (Fig 5C), a finding further supported by statistical analysis (Fig S7).

A critical question arises regarding the timing of TC1 and TC2 computations of reward-prediction errors. Their activity peaks within 0.2 seconds post-cue onset (Fig S3), whereas rewards are delivered between 0.5 and 2 seconds. Intriguingly, TC1 and TC2 activities begin to rise even before the cue (Fig. S3), suggesting the involvement of internal forward models in preemptively calculating reward prediction errors. These forward models, which may include a network loop involving the inferior olive nucleus, cerebral cortex, basal ganglia, and cerebellum, function anticipatorily (Bostan and Strick, 2018; Wagner and Luo, 2020; Watabe-Uchida et al., 2017; Kostadinov and Häusser, 2022). We propose that in this task, the acquisition of reinforcement learning critic, crucial for calculating predictive reward prediction error at cue onset, does not take place in Crus II. Should temporal-difference-error learning be active in Crus II, we would expect to observe complex spike (CS) activities correlating with expected liquid reward delivery [0.5, 2] s, showing contrasting patterns between HIT and FA trials during early learning. However, among the eight zones and four tensor components (TCs) analyzed, only TC3 neurons displayed firing patterns consistent with this temporal condition, but their activity patterns were not

contrastive between HIT and FA for that temporal window, and correlation with δQ was relatively weak (Fig 5A). Therefore, it appears that the learning of temporal-difference-error and/or critic primarily occurs outside Crus II. Our findings suggest that TC1 and TC2 neurons function as context-specific actors within a modular reinforcement learning algorithm for Go and No-go cues, respectively, as depicted in Fig. 6. Within this integrated system, the cerebral cortex, basal ganglia, and cerebellum, which are interconnected, collectively support modular reinforcement learning. In this model, each functional module, represented by Purkinje cells, engages in supervised learning designed for specific actors (see Fig 6A).

Figure 6: The modular reinforcement learning of the Go/No-go task and bi-directional plasticity at the PF-PC synapses. A: The state (s) and action (a) are conveyed to the critic by upstream networks and afferent copies from the actors, respectively. The critic then computes a temporal-difference (TD) prediction error δQ as comparing the observed reward-penalty value R-P with the predicted Q value. The prediction error δQ is used to train the state-value prediction in the critic as well as policy of the actors (red arrows). In the Go/No-go task, subsets of Purkinje cells act as context-dependent actors for Go (gray

shade) and No-go (blue shade) cues separately. Here, we postulated that TC1 and TC2 neurons acquire necessary motor commands by utilizing negative reward-prediction error –δQ, relayed by their CF inputs, in a supervised learning framework. B: bidirectional PF-PC plasticity may occur depending on the magnitude of CSs. Consequently, modulation of SSs was in the same direction with change of CS activities during learning (black arrows). Note that CSs of TC1 and TC2 neurons were negatively correlated with reward-prediction errors in Go (blue line) and No-go (red line) trials, respectively. Horizontal dashed lines indicate the threshold $\bar{CS}$, which determines LTD or LTP at the PF-PC synapses (see Discussions). MFs – mossy fibers, PFs – parallel fibers, CF – climbing fiber, GCs – Granule cells, PCs – Purkinje cells, LTP – long-term potentiation, LTD – long-term depression, CS – complex spike, SS – simple spike.

The most striking computational finding of our study is the negative correlation between TC1 and TC2 neurons and the reward prediction error δQ. We observed an increase in TC1 firings in HIT trials and a decrease in TC2 firings in FA trials, correlating with changes in early licking behaviors in these conditions (Fig S3, Hoang et al., 2023). This supports the integration of supervised learning theories of internal models and reinforcement learning within the cerebellum. According to our hypothesis, TC1 and TC2 neurons develop necessary motor commands through reward prediction errors transmitted by their climbing fibers. During initial learning stages, substantial negative reward prediction errors activate TC2 neurons' complex spikes (CSs) strongly. If simple spikes (SSs) of these neurons also fire within 0-500 ms at a baseline rate of 50-100 Hz, this leads to numerous inappropriate licks. The concurrent activation of parallel fiber and climbing fiber inputs at this stage likely induces long-term depression (LTD) at parallel-fiber-to-Purkinje-cell (PF-PC) synapses, reducing SS modulation and mirroring CS modulation. This aligns with simulations by Yamamoto et al. (2002) regarding adaptive control of ocular following responses. As learning progresses, negative reward prediction errors diminish, and CSs activate less, with SSs of these neurons facilitating suppression of licks following No-go cues. In this scenario, TC2 neurons turn into TC4 neurons as suggested in Hoang et al. (2023). In contrast, TC1 neurons show lower CS

activity early on (Fig S3), indicating a tendency for long-term potentiation (LTP) at their PF-PC synapses. Thus, their SSs likely increase over time, enabling more precise licking responses to Go cues, as detailed in Hoang et al. (2023). Intriguingly, the observed bidirectional plasticity at PF-PC synapses, potentially driven by climbing fibers, seems to follow a unified learning rule:

$$\Delta w_i = \alpha \times \delta Q \times x_i \ \approx -\alpha \times (CS - \bar{CS}) \times x_i$$

Here, α, x_i, and w_i represent a constant learning rate, PF input, and PF-PC synaptic weight of the *i*-th granule cell, respectively. The parameter $\bar{CS}$ denotes the threshold amplitude determining plasticity, with LTP requiring a smaller value than LTD (Coesmans et al., 2004). This learning rule suggests that LTP (positive Δw_i, synaptic weight increases) occurs when $CS - \bar{CS} < 0$, whereas LTD (negative Δw_i, synaptic weight decreases) happens when $CS - \bar{CS} > 0$ (Fig 6B). It's worth noting that synaptic plasticity could be also zonal-dependent, with TC1 neurons, primarily distributed in Aldolase-C positive zones (Fig 4A), are more likely to undergo LTP (de Zeeuw and ten Brinke, 2015; de Zeeuw, 2021). Future simultaneous recordings of SSs and CSs may reveal neural mechanisms of these modes of synaptic plasticity.

Technical advances and conclusions

CFs show associations with both sensorimotor variables and a wide range of reward contingencies, suggesting the cerebellum's role in reinforcement learning across various tasks (Wagner and Luo, 2020; Kostadinov and Häusser, 2022). Despite this, prior has not clearly defined the underlying algorithms, specifically how reward variables are implemented and computed in reinforcement learning. The specific contribution of reward-related CF inputs to behavior learning is also not fully understood. To bridge these gaps, we applied the Q-learning algorithm to analyze the licking behavior of mice in Go/No-go tasks and to estimate reward variables. In our computational approach, we differentiated between the state value function V(*s*), dependent on the state *s* alone, and the state-action value function Q(*s*,*a*), reliant on both the state *s* and action *a*. Two-photon Ca^{2+} imaging data highlighted the dominance of the state-action

value function, with significant differences observed in responses (lick) in HIT vs MISS and FA vs CR trials (Fig 2A). We thus employed a state-action Q-value function for expected reward estimation and used the softmax function for action selection based on the learned Q values. Our Q-learning model, with only five hyper-parameters, effectively captured individual mice's learning patterns in licking behavior (Figs 1C and D). Statistical analysis confirmed the importance of all five hyper-parameters, as models with fewer parameters showed reduced performance (Fig S5), emphasizing their relevance in our Go/No-go task. Analysis of the estimated hyper-parameters revealed unique learning strategies among individual mice, with variations in learning rates, penalty values, initial Q values, and individual strategies (temperature) for optimal reward acquisition (Fig 1E).

This study introduces an innovative technical approach by using tensor component analysis (TCA) as a generative model for analyzing spiking activity in single trials, despite the complexity of CFs conveying multiple types of information (Markanday et al. 2021; Ikezoe et al., 2023; Hoang et al., 2023). Typically, unsupervised statistical methods like TCA, while effective in decomposing spiking activity into meaningful components, struggle with noise sensitivity, making them less reliable for single-trial analysis. To overcome this, we applied TCA to peristimulus time histograms (PSTHs) of complex spikes (CSs). This approach reduces trial-to-trial variability, enabling us to identify low-dimensional tensor components (TCs) that represent the underlying dynamics of spiking activity. We then used the temporal profiles of these TCs to refine spike activity analysis in individual trials. As a result, the derived TC scores effectively reflect the spiking dynamics corresponding to the four TCs identified by TCA. This method offers a powerful means to analyze CF activities, providing valuable insights into a range of learning behaviors.

In summary, our study employs a computational approach integrating a Q-learning model, a high-resolution spike detection algorithm, and tensor component analysis (TCA) to explore reward processing in the cerebellum. The results indicate that inputs from climbing fibers to two distinct Purkinje cell

ensembles encode predictive reward-prediction errors. This finding situates within an integrated framework that merges modular reinforcement learning with supervised learning. The significance of this framework lies in its potential to enhance the development of new reinforcement learning algorithms, which could efficiently master complex tasks in a limited number of trials (Kawato et al., 2021). It's important to note some limitations: the complex spikes (CSs) we studied were recorded from Purkinje cells in separate sessions, potentially introducing variability. Also, our research did not track the CSs of individual Purkinje cells throughout the learning process. Consequently, the proposed modular reinforcement learning algorithm in the cerebellum needs further validation through continuous monitoring of individual neurons during learning, coupled with causal analysis of neuronal responses and corresponding behavioral changes.

Methods

Q-learning model

We adopted a Q-learning model, a reinforcement learning algorithm, to model licking behavior of mice in a Go/No-go experiment. The algorithm works by learning a state-action value function, commonly referred to as the Q-function. The Q-function is a mapping from a state-action pair to a scalar value that represents the expected reward for taking a particular action in a particular state. The Q-function is updated over time based on the observed rewards and transitions to new states.

In the Go/No-go auditory-discrimination licking task, the Q-function of a state s (s = Go / No-go cue) and an action a (a = Lick / No-lick) were updated at trial t as follows

$\delta Q_t = \text{R-P}_t - Q_t$

$Q_{t+1} = Q_t + \alpha \cdot \delta Q_t$

where δQ is the reward prediction error, α is the learning rate and R-P is the reward-penalty function

R-P = 1 for s = Go, a = Lick (HIT)

R-P = -ξ for s = No-go, a = Lick (FA)

R-P = 0, otherwise (CR and MISS)

The probability of selecting a given action a in state s is determined by the softmax function comparing Q-function values of an action to all others:

$$Prob(s, a) = \frac{exp\{Q(s, a)/\tau\}}{\sum_a exp\{Q(s, a)/\tau\}}$$

where τ is the temperature parameter, representing a trade-off between exploitation and exploration. Because mice underwent pre-training sessions for 3 days, during which they were rewarded by licking after 1 second of both cues, initial Q values for Go and No-go cues should be positive: $Q_0(s=Go, a=Lick)$ = q_1 and $Q_0(s=No\text{-}go, a=Lick)$ = q_2. In total, the Q-learning model contains 5 hyper-parameters: learning rate α (range, $0.001 \leq \alpha \leq 0.1$), ξ ($0 \leq \xi \leq 1$), temperature τ ($0.01 \leq \tau \leq 0.5$), initial Q values for Go and No-go cues, q_1 and q_2 ($0 \leq q_1 \leq 1, 0 \leq q_2 \leq 1$). We estimated these parameters for individual animals by maximizing the likelihood defined as the sum of Prob(s,a) for all trials.

It is important to emphasize that we fitted the Q-learning model to behavioral data on a trial basis with trials from different sessions concatenated. To evaluate fitting, for each session, we computed the fraction correct for Go cues (number of HIT trials / total number of Go trials) and the fraction incorrect for No-go cues (number of FA trials / total number of No-go trials) from the data, while the probability Prob(s,a) that the last HIT and FA trials were selected from the model (Figs 1CD). The coefficient of determination (R^2) was calculated to measure goodness-of-fit between the data and model probabilities across sessions of individual animals.

$$R^2 = 1 - \frac{SS_{res}}{SS_{tot}}$$
$$SS_{res} = \sum_i (h_i - f_i)^2$$
$$SS_{tot} = \sum_i (h_i - \bar{h})^2$$

, where h_i and f_i are the probabilities of the data and the model at the i-th session, respectively, and $\bar{h}$ is the mean of the data probability across sessions.

Estimation of complex spike activity from two-photon recordings

Ca signals in Purkinje cell dendrites were evaluated for regions-of-interest (ROIs) of the two-photon images extracted by Suite2p software (Pachitariu et al., 2017) and manually selected. Spike trains were reconstructed for 6,445 Purkinje cells sampled in 17 mice, using hyperacuity software (HA_time, Hoang et al. 2020) that detected complex spike (CS) activities for calcium signals of two-photon imaging with a temporal resolution of 100 Hz (see Hoang et al. 2022 for details).

Complex spike to cue stimulus

To evaluate the CS of a single neuron to the cue stimulus, we constructed a peri-stimulus time histogram (PSTH) of CSs in [-0.5, 2] s with a time bin of 50 ms for the four cue-response conditions. They include HIT, FA, CR, or MISS, according to licking behavior within a response period of 1 s to the two cues, i.e., correct lick in response to the go cue, unwarranted lick in response to the No-go cue, correct response rejection to the No-go cue, or response failure to the Go cue, respectively. Each PSTH was subtracted from its baseline activity, defined as the mean value of the firing rate in the range of [-2, -1] s before cue onset.

Partial least-squares regression analysis

We sought to reveal correlations between zonal activity and behavior variables on a single-trial basis (Fig 3). For each trial, neuronal activity was calculated as the mean firing rate of the neurons in the same Ald-C zone, in [-0.5, 2] s after cue onset. Behavior variables include physical reward R (R=1 for HIT trials, R=0 otherwise), Q, δQ, number of licks in 0-0.5 s for Go and No-go cues (Go ⮕ ELick and No-go ⮕ ELick), the number of licks in the reward period (0.5 - 2 s after cue, Go ⮕ RLick and No-go ⮕ RLick) and the number of licks in the late period (2 - 4 s after cue, Go ⮕ LLick and No-go ⮕ LLick).

We conducted partial least-squares (PLS) regression to resolve multi-collinearity of behavioral variables, e.g. between R, Q and δQ. PLS regression searches for a set of low-dimensional components that performs a simultaneous decomposition of dependent and explanatory variables with the constraint that these components explain as much as possible the covariance between the variables. The VIP (Variable Importance in Projection) score is a measure of the importance of each explanatory variable in a PLS regression model, with higher scores indicating greater importance. The VIP score for the j-th variable is given as:

$$VIP_j = \sqrt{\frac{\sum_{f=1}^{F} w_{jf}^2 \times SSY_f \times J}{SSY_{total} \times F}}$$

, where w_{jf} is the weight of the j-th variable and the f-th PLS component, SSY_f is the sum of squares of explained variance for the f-th PLS component and J is the number of explanatory variables (J=9). SSY_{total} is the total sum of squares explained for the dependent variable, and F is the total number of PLS components. In our analysis, the number of PLS components was optimized by 10-fold cross-validation.

Note that the VIP score does not provide sign information (positive or negative) of correlations between explanatory and dependent variables. Although there was no known threshold for a systematic evaluation, VIP score > 1 was typically used as an indicator for a variable to be considered significant. PLS was conducted using the MATLAB function, *plsregress*.

Tensor component analysis

Let x_{ntk} denote the PSTH of neuron n at time step t within cue-response condition k. TCA yields the decomposition

$$x_{ntk} \approx \hat{x}_{ntk} = \sum_{r=1}^{R} \lambda_r w_n^r b_t^r a_k^r$$

, where R is the number of tensor components, w_n^r, b_t^r and a_k^r are the coefficients of the neuron, temporal, and response condition factors, respectively. Those coefficients were scaled to be unit length with the rescaling value λ_r for each component r. We introduced a non-negative constraint of those coefficients ($w_n^r \geq 0$, $b_t^r \geq 0$ and $a_k^r \geq 0$ for all r, n, t and k). In the previous study, we optimized the number of components R = 4 for which solutions were most stable and fitting scores were high (Hoang et al., 2023).

Computation of tensor component score for a single trial

TCA was efficient for decomposing PSTHs into biologically-meaningful components (Hoang et al. 2023), but for systematic analysis of associations of CSs and variables on a trial basis, it is crucial to estimate activity of such components for individual neurons in a single trial. Note that TCA was carried out for PSTHs computed over multiple trials in a given session, but we needed a firing index for each trial for Q-learning analysis. For that purpose, we employed a novel approach to utilize time series of multiple spikes decomposed by TCA as generative models of CSs.

The tensor-based activity of the i-th neuron in the j-th trial (corresponding to the cue-response condition c) with respect to the component r-th (r = 1, .. , 4) was evaluated as:

$$y_{i,j}^r = w_i^r \times a_c^r \times \sum_{s=1}^{S} \delta(t - t_s)b_t^r$$

, where $\sum_{s=1}^{S} \delta(t - t_s)$ represents the timing of CS firings sampled from [-0.5, 2] s after cue onset as the summation of Dirac delta functions δ.

For the j-th trial, the TC score of the r-th component was calculated as the averaged $y_{i,j}^r$ across all neurons in the same recording session. As a result, each trial has four TC scores corresponding to the four TCs.

$$TCscore_j^r = \frac{1}{N}\sum_{i=1}^{N} y_{i,j}^r$$

Sparse canonical correlation analysis

Because CSs may multiplex different information, we conducted sparse Canonical Correlation Analysis (sCCA) to analyze the relationship between TC scores and behavior variables. The goal of sCCA is to find a set of linear combinations (known as "canonical variates") of the variables in each set, such that the correlation between the two sets of canonical variates is maximized. The sCCA includes a sparsity constraint, which promotes solutions in which only a small number of variables are selected in calculation of the canonical variates. This can result in more interpretable and biologically relevant solutions, as it reduces the amount of noise in the data.

In our analysis, sCCA was conducted using the *PMA* package of R, with a LASSO penalty applied to enforce sparsity. L1 bounds were set 0.4 and 0.6 for TC scores and behavior variables, respectively (larger L1 bound corresponds to less penalization). Explanatory and dependent variables were standardized to have mean zero and standard deviation 1 before performing the analysis. The number of canonical variables was 4. The resulting coefficient of TC scores was either 1 or -1. For better interpretation, coefficient vectors of behavioral variables (reported in Fig 5A) were re-signed so that coefficients of TC scores were all 1.

Statistics

All statistical analyses were performed using MATLAB software. Unless otherwise stated, data are presented as means ± standard deviation. For evaluation of correlations between neuronal response and single behavior variables (Figs 2B-D and Figs 5B-F), we fitted a linear mixed-effects model with fixed effect for behavior variables and mouse index as random intercept. The analysis was conducted using MATLAB function *fitlme*, with the slope and its significance p-value were reported. Significance level: n.s, $p > 0.05$; * $p < 0.05$; ** $p < 0.01$; *** $p < 0.001$; **** $p < 0.0001$.

Acknowledgments

HH, KK and KT were supported by Grants-in-Aid for Scientific Research in Innovative Areas (17H06313) and for Transformative Research Areas (22H05160 to M.M., 22H05156 to M Kawato, and 22H05161 to KK). HH and KT were partially supported by JST ERATO (JPMJER1801, "Brain-AI hybrid"). HH, M Kawato, and KK were partially supported by the Grant Number JP21dm0307002, JP21dm0307008, and JP19dm0207080, respectively, Japan Agency for Medical Research and Development (AMED). M Kawato was partially supported by Innovative Science and Technology Initiative for Security Grant Number JP004596, Acquisition, Technology & Logistics Agency (ATLA), Japan. M Kano and KK were partially supported by Grants-in-Aid for Scientific Research (JP18H04012, JP20H05915, JP21H04785 to M.Kano and JP22H00460 to KK) from the Japan Society for the Promotion of Science (JSPS).

References

Akiti K, Tsutsui-Kimura I, Xie Y, Mathis A, Markowitz JE, Anyoha R, Datta SR, Mathis MW, Uchida N, Watabe-Uchida M. Striatal dopamine explains novelty-induced behavioral dynamics and individual variability in threat prediction. *Neuron* **110**(22), 3789–3804 (2022).

Albus, J. S. A theory of cerebellar function. *Math Biosci* **10**, 25–61 (1971).

Amo R, Matias S, Yamanaka A, Tanaka KF, Uchida N, Watabe-Uchida M. A gradual temporal shift of dopamine responses mirrors the progression of temporal difference error in machine learning. *Nat Neurosci.* **25**(8),1082–1092 (2022).

Barto A., Sutton R., Anderson C. Neuron-like elements that can solve difficult learning control problems *IEEE Trans Syst Man Cybern* **13**(5), 835–846 (1983).

Bayer H., Glimcher P. Midbrain dopamine neurons encode a quantitative reward prediction error signal. *Neuron* **47**(1), 129–141 (2005).

DC. Bentivegna, CG. Atkeson, A. Ude, & G. Cheng. Learning to act from observation and practice. *Int. J. Humanoid Robotics*. **1**, 585-611 (2004).

Bostan, A. C. & Strick, P. L. The basal ganglia and the cerebellum: nodes in an integrated network. *Nat Rev Neurosci* **19**, 338–350 (2018).

Carta I., Chen C., Schott A., Dorizan S., Khodakhah K. Cerebellar modulation of the reward circuitry and social behavior. *Science* **363**(6424), eaav0581 (2019).

Chabrol F., Blot A., Mrsic-Flogel T. Cerebellar contribution to preparatory activity in motor neocortex. *Neuron* **103**, 506–519 (2019).

Coesmans M, Weber JT, De Zeeuw CI, Hansel C. Bidirectional parallel fiber plasticity in the cerebellum under climbing fiber control. *Neuron* **44**(4), 691–700 (2004).

Dabney W, Kurth-Nelson Z, Uchida N, Starkweather CK, Hassabis D, Munos R, Botvinick M. A distributional code for value in dopamine-based reinforcement learning. *Nature* **577**(7792), 671–675 (2020).

Doya K. Complementary roles of basal ganglia and cerebellum in learning and motor control. *Curr Opin Neurobiol.* **10**(6), 732–739 (2000).

K. Doya, K. Samejima, K. Katagiri, M. Kawato, Multiple model-based reinforcement learning. *Neural Comput*. **14**(6), 1347-69 (2002).

Haruno M., Kawato M. Different neural correlates of reward expectation and reward expectation error in the putamen and caudate nucleus during stimulus-action-reward association learning. *J Neurophysiol.* **95**(2), 948–959 (2006).

Haruno M, Kuroda T, Doya K, Toyama K, Kimura M, Samejima K, Imamizu H, Kawato M. A neural correlate of reward-based behavioral learning in caudate nucleus: a functional magnetic resonance imaging study of a stochastic decision task. *J Neurosci.* **24**(7), 1660–1665 (2004).

Heffley, W. & Hull, C. Classical conditioning drives learned reward prediction signals in climbing fibers across the lateral cerebellum. *Elife* **8**, (2019).

Heffley, W. *et al.* Coordinated cerebellar climbing fiber activity signals learned sensorimotor predictions. *Nat Neurosci* **21**, 1431–1441 (2018).

Hirano T. Depression and potentiation of the synaptic transmission between a granule cell and a Purkinje cell in rat cerebellar culture. *Neurosci Lett* **119**, 141–144. (1990).

Hirano T. Long-term depression and other synaptic plasticity in the cerebellum. *Proc Jpn Acad Ser B Phys Biol Sci* **89**(5), 183–195 (2013).

Hoang, H. *et al.* Improved hyperacuity estimation of spike timing from calcium imaging. *Sci Rep* **10**, 17844 (2020).

Hoang, H., Tsutsumi, S., Matsuzaki, M., Kano, M., Kawato, M., Kitamura, K., Toyama, K. Dynamic organization of cerebellar climbing fiber response and synchrony in multiple functional components reduces dimensions for reinforcement learning, *eLife* 12:e86340 (2023).

Ikezoe, K. *et al.* Cerebellar climbing fibers multiplex movement and reward signals during a voluntary movement task in mice. *Communications Biology* 6:924 (2023).

Ito M. Cerebellar long-term depression: characterization, signal transduction, and functional roles. *Physiol Rev* **81**(3), 1143–1195 (2001).

Ito, M. Neurophysiological aspects of the cerebellar motor control system. *Int J Neurol Neurother* **7**, 162–176 (1970).

Ito M., Kano M. Long-lasting depression of parallel fiber-Purkinje cell transmission induced by conjunctive stimulation of parallel fibers and climbing fibers in the cerebellar cortex. *Neurosci Lett* **33**(3), 253–258 (1982).

Kaplan A, Mizrahi-Kliger AD, Israel Z, Adler A, Bergman H. Dissociable roles of ventral pallidum neurons in the basal ganglia reinforcement learning network. *Nat Neurosci.* **23**(4), 556–564 (2020).

Kawato M. Internal models for motor control and trajectory planning. *Curr Opin Neurobiol* **9**(6), 718–727 (1999).

Kawato M., Furukawa K., Suzuki R. A hierarchical neural-network model for control and learning of voluntary movement. *Biol Cybern* **57**(3), 169–85 (1987).

Kawato M., Gomi H. A computational model of four regions of the cerebellum based on feedback-error learning. *Biol Cybern* **68**, 95–103 (1992).

Kawato, M., Ohmae, S., Hoang, H., Sanger, T. 50 Years Since the Marr, Ito, and Albus Models of the Cerebellum. *Neurosci*., **462**, 151–174 (2021).

Kawato M, Samejima K. Efficient reinforcement learning: computational theories, neuroscience and robotics. *Curr Opin Neurobiol.* **17**(2), 205–212 (2007).

Kostadinov, D., Beau, M., Blanco-Pozo, M. & Häusser, M. Predictive and reactive reward signals conveyed by climbing fiber inputs to cerebellar Purkinje cells. *Nat Neurosci* **22**, 950–962 (2019).

Kostadinov, D. & Häusser, M. Reward signals in the cerebellum: Origins, targets, and functional implications. *Neuron* **110**, 1290–1303 (2022).

Larry, N., Yarkoni, M., Lixenberg, A., Joshua, M. Cerebellar climbing fibers encode expected reward size. *Elife* 8:e46870 (2019).

Linden D. The return of the spike: postsynaptic action potentials and the induction of LTP and LTD. *Neuron* **22**, 661–666 (1999).

Markanday, A., Inoue, J., Dicke, P. W. & Thier, P. Cerebellar complex spikes multiplex complementary behavioral information. *PLoS Biol* **19**, e3001400 (2021).

Marr, D. A Theory of Cerebellar Cortex. *J. Physiol* **202**, 437–470 (1969).

Medina JF, Lisberger SG. Links from complex spikes to local plasticity and motor learning in the cerebellum of awake-behaving monkeys. *Nat Neurosci.* **11**(10):1185–1192 (2008).

Medina JF, Nores WL, Ohyama T, Mauk MD. Mechanisms of cerebellar learning suggested by eyelid conditioning. *Curr Opin Neurobiol.* **10**(6), 717–724 (2000).

Pachitariu, M. et al. Suite2p: beyond 10,000 neurons with standard two-photon microscopy. *bioRxiv* (2017).

Pessiglione M, Seymour B, Flandin G, Dolan RJ, Frith CD. Dopamine-dependent prediction errors underpin reward-seeking behaviour in humans. *Nature* **442**(7106), 1042–1045 (2006).

Raymond JL, Medina JF. Computational Principles of Supervised Learning in the Cerebellum. *Annu Rev Neurosci.* **41**, 233–253 (2018).

Schultz W., Dayan P., Montague P. A neural substrate of prediction and reward. *Science*. **275**(5306), 1593–1599 (1997).

Sendhilnathan, N., Ipata, A. & Goldberg, M. E. Mid-lateral cerebellar complex spikes encode multiple independent reward-related signals during reinforcement learning. *Nat Commun* **12**, 6475 (2021).

Sendhilnathan, N., Semework, M., Goldberg, M., Ipata, A. Neural Correlates of Reinforcement Learning in Mid-lateral Cerebellum. *Neuron* 106(1):188–198.e5 (2020).

Shidara M., Kawano K., Gomi H., Kawato M. Inverse-dynamics model eye movement control by Purkinje cells in the cerebellum. *Nature* **365**, 50–52 (1993).

Starkweather CK, Uchida N. Dopamine signals as temporal difference errors: recent advances. *Curr Opin Neurobiol.* **67**, 95–105 (2021).

Sugihara I, Shinoda Y. Molecular, topographic, and functional organization of the cerebellar cortex: a study with combined aldolase C and olivocerebellar labeling. *J Neurosci.* **24**(40), 8771–8785 (2004).

Sugihara I, Shinoda Y. Molecular, topographic, and functional organization of the cerebellar nuclei: analysis by three-dimensional mapping of the olivo-nuclear projection and aldolase C labeling. *J Neurosci.* **27**(36), 9696–9710 (2007).

N. Sugimoto, M. Haruno, K. Doya, M. Kawato, MOSAIC for multiple-reward environments. *Neural Comput.* **24**(3) 577–606 (2012).

N. Sugimoto, J. Morimoto, SH. Hyon, M. Kawato, The eMOSAIC model for humanoid robot control. *Neural Netw.* **29-30**, 8-19 (2012).

Sutton, R. S. & Barto, A. G. Reinforcement Learning: An Introduction 2nd edn, *MIT Press* (2018).

Tsutsumi, S. *et al.* Modular organization of cerebellar climbing fiber inputs during goal-directed behavior. *Elife* **8**, (2019).

Tsutsumi, S. et al. Purkinje Cell Activity Determines the Timing of Sensory-Evoked Motor Initiation. *Cell Rep* **33**, 108537 (2020).

Wagner, M. J., Kim, T. H., Savall, J., Schnitzer, M. J. & Luo, L. Cerebellar granule cells encode the expectation of reward. *Nature* **544**, 96–100 (2017).

Wagner, M. J. & Luo, L. Neocortex-Cerebellum Circuits for Cognitive Processing. *Trends Neurosci* **43**, 42–54 (2020).

Wagner, M. J. *et al.* Shared Cortex-Cerebellum Dynamics in the Execution and Learning of a Motor Task. *Cell* **177**, 669–682.e24 (2019).

Wagner, M. J. et al. A neural circuit state change underlying skilled movements. *Cell* **184**, 3731–3747.e21 (2021).

Watabe-Uchida, M., Eshel, N. & Uchida, N. Neural Circuitry of Reward Prediction Error. *Annu Rev Neurosci* **40**, 373–394 (2017).

Watkins, C., Dayan, P. Q-learning. *Mach. Learn.* **8**, 279–292 (1992).

Welsh, J. P., Lang, E. J., Suglhara, I. & Llinás, R. Dynamic organization of motor control within the olivocerebellar system. *Nature* **374**, 453–457 (1995).

Williams, A. H. *et al.* Unsupervised Discovery of Demixed, Low-Dimensional Neural Dynamics across Multiple Timescales through Tensor Component Analysis. *Neuron* **98**, 1099–1115.e8 (2018).

Wolpert D., Miall C., Kawato M. Internal models in the cerebellum. *Trends in Cognitive Sciences* **2**, 338–347 (1998).

de Zeeuw, C. I. Bidirectional learning in upbound and downbound microzones of the cerebellum. *Nat Rev Neurosci* **22**, 92–110 (2021).

de Zeeuw, C. I. & ten Brinke, M. M. Motor learning and the cerebellum. Cold Spring Harb Perspect Biol 7, (2015).

Supplemental Figures

Figure S1: PLS regression of TC activity and behavior variables. Color convention is the same as Figure

3.

Figure S2: PLS regression for zonal activity (A) and sparse CCA for TC1-4 scores (B) with an additional reward-penalty R-P variable. Color convention is the same as Figure 3 with a dark blue column for R-P.

Figure S3: At each learning stage, we first sampled 300 neurons whose coefficients were highest for each tensor component, TC1-4. Then we excluded those that appeared in at least two samples. e.g TC1 and TC3. This figure plots PSTHs in the four cue-response conditions of selected neurons for the four tensor components. Blue, green and red traces are for 1st, 2nd and 3rd learning stages, respectively.

Figure S4: PLS regression for zonal activity (A) and sparse CCA for TC1-4 scores (B) with auditory cue and motor variables independent. In these analyses, the auditory cue (Cue=1 for Go and Cue=0 for No-go cues) was made independent from motor (licking) variables, resulting in a total of 7 exploratory variables (Cue, R, Q, δQ, ELick, RLick, and LLick). Color convention is the same as Figure 3 with a new black column corresponding to the auditory cue.

Model #1: all hyper-parameters varied
Model #2: varied: α, τ, ξ, q1 - fixed: q2 = q1
Model #3: varied: α, q1, q2 - fixed: $\tau = 0.1$, $\xi = 0.5$
Model #4: varied: α, q1 - fixed: q2 = q1, $\tau = 0.1$, $\xi = 0.5$
Model #5: varied: q1 - fixed: $\alpha = 0.05$, q2 = q1, $\tau = 0.1$, $\xi = 0.5$

Figure S5: BIC score, estimated for 17 mice, of the five Q-learning models with different numbers of hyper-parameters. The total BIC score was 21,235; 21,291; 21,414; 21,532 and 69,325 for Model #1-5, respectively.

Figure S6: PLSR for the top TC neurons. A: we sampled the top 300 neurons for each TC at each learning stage, then removed overlapping ones (see Hoang et al., 2023 for details). As a result, we selected 2,096 neurons (termed "topTC neurons") and individually conducted PLSR of their TC score with the top four explanatory variables, including R, Q, δQ and No-go ⊠ ELick (Fig 5A). For individual neurons, we selected the explanatory variable which has the highest VIP score. The pie-charts indicated fraction of the explanatory variable selected for each of the topTC1-4 neurons. B: the averaged VIP score of PLSR for topTC1-4 neurons. The horizontal line indicates VIP score = 1.

Fig S7: Shuffled correlations between TC1 and TC2 score with reward-prediction errors δQ. A: For each animal, we shuffled the pairs of TC1 score vs. reward-prediction errors δQ in HIT trials and conducted the regression analysis similar to Fig 5C. The bar-plots indicated frequency of the slope (left) and p-value (right) of the regression for 1000 shuffled times. B: similar to A but for TC2 score vs. δQ in FA trials. Note that, for the original data, the slope of regression was -0.65 for TC1-HIT and -1.24 for TC2-FA (vertical red lines) with p-value < 0.00001 (reported in Fig 5C&D).