The origin and structural evolution of *de novo* genes in *Drosophila*

Junhui Peng, Li Zhao

Laboratory of Evolutionary Genetics and Genomics, The Rockefeller University, New York, NY 10065, USA

*Correspondence to: lzhao@rockefeller.edu

Abstract

Although previously thought to be unlikely, recent studies have shown that *de novo* gene origination from previously non-genic sequences is a common mechanism for gene innovation in many species and taxa. These young genes provide a unique set of candidates to study the structural and functional origination of proteins. However, our understanding of their protein structures and how these structures originate and evolve are still limited, due to a lack of systematic and case studies. Here, we combined high-quality base-level whole genome alignments, bioinformatic analysis, and computational structure modeling to study the origination, evolution, and protein structure of lineage-specific *de novo* genes. We identified 589 *de novo* gene candidates in *D. melanogaster* that originated within the *Drosophilinae* lineage. We found a gradual shift in sequence composition, evolutionary rates, and expression patterns with their gene ages, indicating possible gradual shifts or adaptations of their functions. Surprisingly, we found little overall protein structural changes for *de novo* genes in the *Drosophilinae* lineage. We identified a number of *de novo* gene candidates with protein products that are potentially well-folded, many of which are more likely to contain transmembrane and signal proteins compared to other annotated protein-coding genes. Using ancestral sequence reconstruction, we found that most potentially well-folded proteins are often born folded. Interestingly, we observed one case where disordered ancestral proteins become ordered within a relatively short evolutionary time. Single-cell RNA-seq analysis in testis showed that several young *de novo* genes are biased in the early spermatogenesis stage, indicating unneglectable roles of early germline cells in the *de novo* gene origination in testis. This study provides a systematic overview of the origin, evolution, and structural changes of *Drosophilinae*-specific *de novo* genes.

Key words: *De novo* genes; Drosophila; origin; structures; fold; ancestral sequence reconstruction; molecular dynamics

Introduction

De novo genes are novel genes born from scratch through previously non-genic DNA sequences (Begun et al., 2006; Levine et al., 2006; McLysaght & Hurst, 2016). Recent works support the existence of a considerable number of young *de novo* genes in various species and taxa, including humans and *Drosophila* (Begun et al., 2007; Cai et al., 2008; Chen et al., 2010; Heinen et al., 2009; Knowles & McLysaght, 2009; D. Li et al., 2010; Van Oss & Carvunis, 2019; L. Zhang et al., 2019; Zhao et al., 2014; Zheng & Zhao, 2022). Some studies proposed that *de novo* genes tend to be highly disordered (Wilson et al., 2017), which may help prevent misfolding or aggregation. This is important because the accumulation of misfolded or aggregated proteins can be neurotoxic in complex

eukaryotes (Wilson et al., 2017). However, other studies suggested that *de novo* genes may not necessarily be disordered (Bornberg-Bauer et al., 2021a; Bungard et al., 2017; Lange et al., 2021) and instead suggest that their structures could be highly conserved after their origination.

Despite the aforementioned advancements, our understanding of the protein structures of *de novo* genes remains very limited. It is still unclear whether *de novo* genes are capable of being well-folded, how frequently they are well-folded, and if they possess novel structural folds. The main obstacle was the lack of accurate, efficient, and scalable structural characterization tools that could be applied to a large amount of *de novo* genes. Here, we applied AlphaFold2 (Jumper et al., 2021) computational predictions to evaluate the foldability of *de novo* genes. With the rapid development in genome sequencing and alignment and deep learning techniques, AlphaFold2, along with other neural network approaches, e.g., trRosetta (J. Yang et al., 2020) and RoseTTAFold (Baek et al., 2021a), have demonstrated the ability to predict protein structures with near-atomic accuracy. AlphaFold2 has also been applied at genomic scales to predict protein structures of the human proteome and the proteomes of several other species (Tunyasuvunakool et al., 2021; Varadi et al., 2021). In addition, the increasing power of bioinformatic and computational approaches have made it possible to obtain highly accurate general structural properties of genes, including intrinsic structural disorder (Necci et al., 2021), relative solvent accessibility (S. Wang et al., 2016), and the probability of being transmembrane proteins (S. Wang et al., 2016) or signal proteins (Almagro Armenteros et al., 2019a).

In addition to the structure of *de novo* genes, the evolution of their sequences and structures after origination remains unclear. To address this question, it is necessary to identify and compare lineage-specific *de novo* genes ranging from very young and old within a relatively diverged lineage. However, due to low genome sequencing quality and high genome recombination rates in some species, this process could be very difficult, especially with increased divergence time (Vakirlis, Carvunis, et al., 2020; Van Oss & Carvunis, 2019). This limitation greatly hampered our understanding of the origination of *de novo* genes and how their sequences and structures evolve after origination. In addition, it has been historically difficult to distinguish between rapidly evolving genes and *de novo* originated genes (Moyers & Zhang, 2016a; Weisman et al., 2020). However, with the recent advancement in whole-genome sequence alignments and its ability to progressively align the genomes of an entire phylogenetic tree, including diverged species with high accuracy (Armstrong et al., 2020), we can now identify *de novo* protein-coding gene candidates with high confidence through the support of synteny-based alignments and non-coding sequences in outgroup species.

In this work, we utilized progressive whole-genome syntenic alignments and all-vs-all BLASTP analysis (Altschul et al., 1990), to identify 589 *de novo* protein-coding genes in *D. melanogaster* that were born at least ~67 million years since the *Drosophilinae* lineage (Kumar et al., 2017) with the support of ortholog non-coding DNA sequences in their corresponding outgroup species. We performed bioinformatic analysis and computational structural predictions to characterize the structural and functional properties of each *de novo* gene. Additionally, we employed ancestral sequence reconstruction (ASR) to examine the sequence and structural evolution of these genes throughout their evolutionary history. Our results provided a systematic overview of the foldability of

proteins encoded by *de novo* genes and the patterns in which their sequences and structures evolve after origination.

Results

Identification of *Drosophilinae* lineage-specific *de novo* gene candidates in *D. melanogaster*

We built the alignment of 20 *Acalyptratae* fly species (Figure 1A) using Progressive Cactus Aligner (Armstrong et al., 2020). A summary of the alignments can be found in Figure S1. Overall, the protein-coding bases in *D. melanogaster* can be aligned to closely related species at a high coverage (97% with *D. sechellia*), while the coverage dropped significantly to around 50% in relatively distant species (49% with *B. dorsalis*). For the 13968 annotated protein-coding genes in *D. melanogaster* investigated in our study, we observed 13798 (98.8%) of them were covered in cactus alignments. Among these 13798 annotated protein-coding genes in *D. melanogaster*, we found 1285 of them were aligned to previously unannotated regions in their most distant aligned lineages *Li* (Figure 1B). We further removed genes with their corresponding unannotated hits being potentially genic (Material and Methods) in lineage *Li* or have homologous protein sequences in more outgroup species (Figure 1B, Material and Methods). Because of that, we did not observe *de novo* genes that are *D. melanogaster* specific. However, we observed 73 genes with unannotated hits in *D. simulans*, *D. sechellia,* or other species in more distant lineages. For these 73 genes, 47 of them were considered not to be *de novo* genes since the unannotated aligned regions were predicted to be unannotated putative orthologs (Material and Methods) with high confidence based on gene structure predictions, while other 26 genes were identified as *de novo* gene candidates but were inferred to be originated from more distant lineages. We also removed cases where homologous genes are translocated to other syntenic blocks or chromosome arms, using homology and copy numbers inferred from OrthoMCL (L. Li et al., 2003). To this step, we obtained 686 *de novo* gene candidates. As a last filtering step, we searched against UniProt Knowledgebase sequence database (UniprotKB) (Bateman et al., 2022) to filter candidates that have homologs in species that are more distant to their inferred origination lineage (Material and Methods, Figure 1B). Finally, combined with homology and synteny, we identified 589 lineage-specific *de novo* protein-coding gene candidates in *D. melanogaster* that are potentially originated within *Drosophilinae* lineage (supplementary File S1: The list of *de novo* gene candidates and their properties). All these *de novo* gene candidates were supported by evidence of possible ancestral noncoding DNA sequences in *Li*. Based on the location of ancestral noncoding DNA sequences, we found that 421 of them were born from intergenic regions, and 168 of them from intragenic regions.

Figure 1. Workflow to identify *Drosophillinae* lineage-specific *de novo* genes in *D. melanogaster*. (A) Phylogenetic tree of the *Drosophilinae*, *Acalyptratae*, and *Arthropods* species. Progressive cactus genome alignment was performed among the 20 *Acalyptratae* genomes. (B) The workflow and pipeline to identify *Drosophilinae* lineage-specific *de novo* genes and track the origination within the *Drosophilinae* lineage for the current protein-coding genes in *D. melanogaster* (see Material and Methods).

The origin of *Drosophilinae*-specific *de novo* genes is more likely to be associated with open chromatins than transposable elements

Our analysis shows that most *de novo* gene candidates had biased expressions in testis, head, and ovary. Thus, to understand their relationships with open chromatin regions, which are enriched with regulatory sequences, we examined ATAC-seq data from the three tissues (Witt et al., 2021). We found that both intergenic and intragenic *de novo* gene candidates have more peaks in their nearby ±500 bp regions than putative random ORFs in intergenic regions. In these regions, the peaks of the *de novo* gene candidates also have higher peak intensities. When considering broader regions, we found that *de novo* gene candidates, regardless of being intergenic or intragenic, were closer to their nearest peaks. This suggest that *de novo* genes are partly associated with open chromatin conformation changes. To assess the potential involvement of TEs in de novo gene origination, we searched for DNA repeat signals (see Material and Methods) in these regions. However, we did not find evidence of association between *de novo* gene candidates and TEs in *Drosophila*, contrasting with previous reports of up to 20% of *de novo* transcripts being associated with TEs in primates, including humans (Ruiz-Orera et al., 2015). One possible explanation that is the primate genomes have much higher TE contents than insect, thus the potential regulatory roles of TEs in *Drosophila* new genes is limited. Our findings suggest that in *Drosophila*, lineage-

specific *de novo* genes are associated with open chromatin regions but not TEs (Figure S2), a different pattern from what has been observed in primates.

De novo gene candidates are mostly adaptive and are shaped by both adaptive and non-adaptive changes

We found, compared to other annotated genes, *de novo* genes have different properties in sequence composition (GC content), sequence evolution (ω, ω_a, ω_{na}, and α), structural properties, and expression patterns (Figure 2). Specifically, *de novo* genes exhibits a lower GC content (Figure 2A), and this trend applies to each of the amino acids that contains G or C in their codons (Figure 2B). This suggests that *de novo* genes might be using immature codons and that translational selection on codon usage plays an important a role in their evolution.

Compared to other protein-coding genes, *de novo* genes tend to be more disordered (p=3e-86), more exposed (1e-91), and are more likely to be transmembrane proteins (2e-6) or secretory proteins (p=8e-12) (Figure 2C).

Compared to other proteins, *de novo* genes have higher male specificity (p=2e-85), higher tissue specificity (p=5e-93), relatively lower expression levels in females (p=3e-79), and slightly higher expression levels in males (p=0.02). These patterns are consistent with observations in other lineages, further indicating that de novo genes exhibit some universal patterns that are not dependent on their lineage or identification method.

By analyzing comparative and population genomics data, we found that *de novo* genes are under faster sequence evolution compared to other protein-coding genes (Figure 2E). The adaptation rates, nonadaptation rates, and proportions of adaptive changes of *de novo* genes are higher than other genes (Figure 2E), indicating that both adaptive evolution and relaxation of purifying selection contribute to the elevated evolutionary rates of *de novo* genes. The results that *de novo* genes are more likely to be transmembrane proteins or secretory proteins (Figure 2C) and are more likely to be male specific or tissue specific (Figure 2E) suggest that some *de novo* genes might have specific molecular or cellular functions (Vakirlis, Acar, et al., 2020).

Figure 2. Special properties of *de novo* gene candidates. (A) GC contents of *de novo* gene candidates (denoted as "*de novo*") were significantly lower than other annotated protein-coding genes (denoted as "other"). (B) GC contents of each amino acids in *de novo* gene candidates were significantly lower than amino acids in other annotated protein-coding genes. *De novo* gene candidates also differed in structural properties (C), expression patterns (D), and evolutionary patterns (E). (F) Evolutionary rates of *de novo* gene candidates with different gene ages. Due to insufficient data, L1, L2, and L3 were not shown. (G) Correlations between different properties of *de novo* gene candidates and their gene ages. Positive correlations have P-values greater than 0, while negative correlations have P-values smaller than 0. Expression patterns and evolutionary patterns of *de novo* gene candidates were significantly correlated with gene ages, while structural properties were not. (H) Structural disorder (ISD) and solvent accessibility of de novo gene candidates with different gene ages. (I) The probability of containing transmembrane or signal proteins for *de novo* gene candidates with different gene ages.

De novo gene candidates undergo gradual sequence/function changes without significant structural changes

To understand whether and how structures change after the fixation of a *de novo* protein-coding gene, we then studied structural differences in *de novo* genes among different origination lineages or gene ages. We compared the differences in protein properties of *de novo* genes to their origination lineages (L1 to L9, Figure 1A) by applying Kendall tau and Spearman's rank correlation analysis. We observed significant sequence changes among different origination lineages. For example, the sequence evolutionary rates (ω) of *de novo* genes are significantly negatively correlated with origination lineage (Kendall tau P = 2e-9, Spearman's rank P = 2e-9, Figure 2F and

2G). The significant changes of ω might be dominated by the decrease of non-adaptive changes (ω_{na}) since non-adaptation rates had significant negative correlations with origination lineages as well (Kendall tau P = 4e-3, Spearman's rank P = 3e-3, Figure 2F and 2G), while the negative correlations between adaptation rates and origination lineages were not significant (Kendall tau P = 0.2, Spearman's rank P = 0.2, Figure 2F and 2G). The decrease of non-adaptive changes (ω_{na}) might indicate an increase in the strength of purifying selection, suggesting the functional importance of *de novo* genes increase over limited evolutionary time.

We found that GC contents are higher for *de novo* genes originated in younger lineages and lower in older lineages (Kendall tau P = 2e-6, Spearman's rank P = 1e-6, Figure 2G), which might suggest an important role of translational selection in the evolution of *de novo* genes. We observed significant correlations between male specificity and tissue specificity and origination lineages (Kendall tau P = 4e-6 and 5e-7, Spearman's rank P = 4e-6 and 3e-7, Figure 2G). Similarly, we observed weaker but significant correlations for male and female expression levels (Kendall tau P = 6e-3 and 2e-3, Spearman's rank P = 5e-3 and 2e-3, Figure 2G). The correlations between expression patterns and origination lineages might indicate a gradual change in protein functions, which might further suggest that these *de novo* genes are under certain degrees of selection.

Interestingly, we did not observe significant changes in structural properties among different origination lineages, including structural disorder (ISD), solvent accessibility, probability of being transmembrane proteins, and probability of being signal proteins with the P-values of both Kendall tau and Spearman's rank correlations greater than 0.5 (Figure 2G, 2H, and 2I). This indicates that upon origination, the overall structural properties of *de novo* genes might remain similar in *Drosophilinae* lineage. The correlations between *de novo* gene properties and their gene ages indicate that these *de novo* gene candidates might undergo gradual sequential and functional changes without significant structural changes.

A small subset of *de novo* genes is potentially well-folded with complex structural folds

We used AlphaFold2 (Jumper et al., 2021) to predict the structural models of *de novo* gene candidates. We further used the per-residue confidence score (pLDDT) from AlphaFold2 predictions to estimate the foldability of these candidates as high pLDDT scores often indicate accurate protein folding, and low pLDDT scores highly correlate with protein disorder (Jumper et al., 2021). We showed that pLDDT from AlphaFold2 predictions strongly correlated with pLDDT from RoseTTAFold (Baek et al., 2021b) predictions (Pearson correlation R = 0.79, P = 4e-68) and the convergence of trRosetta (J. Yang et al., 2020) predictions (Pearson correlation R = 0.64, P = 4e-37) (Figure 3A, Material and Methods), suggesting consistency among the three state-of-the-art prediction methods. We categorized *de novo* gene candidates into three structural groups, each for 1) potentially well-folded, 2) partially folded, and 3) not folded. We defined genes to be potentially well folded if their average per-residue confidence score (pLDDT) were greater than 0.8 and the percent of confidently predicted residues (pLDDT > 0.7) greater than 90%; genes to be potentially partially folded if more than 30% of their residues or more than 50 consecutive residues being confidently predicted (pLDDT > 0.7); and the remaining genes to be potentially not folded. We found that most of the *de novo* gene candidates might only be partially folded (318/589) or not folded

(236/589) (Figure 3B, supplementary File S2: The list of *de novo* gene candidates and their properties). Interestingly, we found 35 *de novo* gene candidates that are potentially well-folded. Among these 35 *de novo* genes, 16 of them might only fold into simple folds containing a single or two α-helices, while another 19 might fold into complex folds (Table S1). We further performed 200 ns MD simulations for each of the 19 structural models to refine the structural models and characterize the stabilities of the structural folds (see Material and Methods, Table S2). The structural models all retained highly similar structural folds during the MD simulations, with pairwise TM-scores of representative conformations larger than 0.70 (Figure 3D). For example, CG42590 and CG43070 had averaged pLDDT scores of 0.90 and 0.89 (Figure 3C, Table S1). Their structural folds remained highly stable during MD simulations with averaged pairwise TM-score values of 0.98 and 0.92 (Table S2).

Figure 3. AlphaFold2 structure predictions of *de novo* gene candidates. (A) AlphaFold2 predictions are strongly correlated with RoseTTAFold and trRosetta predictions. (B) Most *de novo* gene candidates might not have the ability to be well-folded according to pLDDT, per-residue confidence score, and P(confident), the percent of residues that were confidently predicted. (C) Two examples of potentially well-folded *de novo* gene candidates: CG42590 and Cg43070. In both cases, the core regions were predicted with very high confidence (blue). (D) The predicted structural models of the 19 potentially well-folded *de novo* genes retained highly similar structural folds during MD simulations with pairwise TM-score of representative structures larger than 0.70.

Potentially well-folded *de novo* genes might be novel structural folds

To check if potentially well-folded *de novo* genes have novel structural folds, we compared their MD refined structures with all experimentally determined protein structures in protein data bank (PDB). To do this, we searched through protein databank (PDB) to check whether the potentially well-folded genes have novel structural

folds using RUPEE (Ayoub & Lee, 2019). Interestingly, we found that most of them (16/19) have similar structural folds in PDB (Table S1). These similar structural folds were not due to sequence similarity. In contrast, the sequence identity between the de novo gene candidates and their similar structures were mostly less than 10% (Table S1). For example, the structural model of CG43195 is similar to the A chain of PDB structure 1U89 with a TM-score of 0.70. However, their sequence identity is only 4% (Table S1). Notably, we also found 3 *de novo* gene candidates, Eig71Ei, Eig71Ej, and CG43251, with maximum TM-score against protein structures in PDB smaller than 0.5 (Table S1, Figure S3), suggesting that the 3 candidates might adopt novel structural folds that have not been identified before. The results indicated that *de novo* genes might be well-folded with novel structural folds.

Most potentially well-folded *de novo* genes are likely to be born well-folded

We showed that *de novo* gene candidates could undergo fast sequence adaptation without significant structural changes. We further investigate the structural changes of *de novo* genes after origination using the 19 potentially well folded *de novo* genes as examples. For all the 19 *de novo* genes, we reconstructed their ancestral states, including most ancestral states and intermediate states in other lineages, by ancestral sequence reconstruction (ASR). We then used AlphaFold2 to predict the 3D structures of these ancestral states. We found that the ancestral states of the 19 *de novo* genes were all predicted to be potentially folded at high confidence (average pLDDT > 0.7, Table S1). We further compared the structural models of ancestral states to current states. We found that they share similar structural folds with all pairwise TM-scores greater than 0.56 (Figure 4A, top panel), suggesting that these potentially well-folded *de novo* genes are likely to be born with similar structural folds to the current forms. To refine the structural models and characterize the stabilities of the structural folds, we further conducted 200 ns MD simulations starting from the ancestral structural models (see Materials and Methods, summarized in supplementary File S3: MD simulations of ancestral state structural models). We found that most of the ancestral state structural models retained similar structural folds to their current D. *melanogaster* forms during MD simulations, with almost all pairwise TM-scores of representative conformations close to or larger than 0.70 (Figure 4A, middle panel and bottom panel). In addition, we found that the structural fold stabilities and their similarities to their current forms did not change with the age of ancestral states (Figure 4B). Taking CG42590 as an example, MD simulations of CG42590 in the current form revealed that it contains a highly stable structural fold (Figure 4C, left column). Meanwhile, MD refined structural models of CG42590 ancestral states revealed that they are highly similar to CG42590 current form (Figure 4D, right column), with average TM-score of 0.90. Interestingly, we also observed one exception, CG43251. We found that CG43251 might undergo substantial global structural changes from ancestral states to their current state (Figure 4A, last column), with average TM-score of 0.32 to their current form. In the MD refined structure of the current form, CG43251 forms a stable fold with an alpha-helix in the N-terminal and a β-hairpin in the C-terminal (Figure 4D, left column), while in its ancestral states, the structures were disordered (Figure 4D, right column). Altogether, the

results suggested that after origination, most potentially well folded *de novo* genes might preserve a similar fold as they were born, while in some rare cases, the candidates undergo substantial structural changes.

Figure 4. Structural evolution of 19 potentially well-folded *de novo* gene candidates. (A) Ancestral states have similar structural folds to their current forms as predicted by AlphaFold2 ("Pred", top panel). The structural folds of ancestral structural models were stable ("AncMD", middle panel) and similar to their current forms ("ToDmel", bottom panel) during MD simulations, except for the case of CG43251 (last column). (B) Structural fold stability of ancestral states ("AncMD", top panel), and current forms ("melMD", middle panel) as a function of the ancestral state ages. Structural models of ancestral states were similar to their current forms, regardless of their ancestral branches ("ToDmel", bottom panel). (C and D) Two examples of the structural evolution of potentially well-folded *de novo* gene candidates. AlphaFold2 predicted structural model (blue), MD refined structural model (red) and other representative structural models (gray) of CG42590 (C, left panel) and CG43251 (D, left panel). MD refined structural models of different ancestral states of CG42590 (C, right panel) and CG43251 (D, right panel) colored by residue index, with N-terminus being blue and C-terminus being red. Ancestral states of CG42590 have similar structural folds while those of GC43251 are different.

Early germline cells in testis are unneglectable in *de novo* gene origination

Among the 589 *de novo* gene candidates identified, more than half of them (301) were biased expressed toward testis. To investigate how *de novo* genes originated and evolved in testis, we analyzed the expression patterns of testis-biased *de novo* genes using testis single-cell RNA-sequencing data (Witt et al., 2019). We used the expression patterns of all annotated testis-biased *Drosophila melanogaster* genes to cluster these genes into four different clusters (Figure 5A). We found that genes in cluster #1 were highly expressed in early spermatogenesis stage, genes in cluster #2 showed average expression in spermatogonia and spermatocytes stages, genes in cluster

#3 showed average expression in spermatocytes and spermatids stages, and genes in cluster #4 showed peak expression in spermatids stage. We found that testis-biased *de novo* genes' expression profile is similar to other testis-biased genes (Figure 5B). We then asked whether *de novo* genes in different clusters showed different structure/sequence characteristics. We found that *de novo* gene candidates in cluster #1 were significantly different from other genes. For example, these *de novo* gene candidates were less likely to be transmembrane or secretory proteins, and they were more likely to be disordered or exposed proteins than *de novo* gene candidates in other clusters (Figure 5C). We also found that, compared to other clusters, *de novo* genes in cluster #1 evolve faster, and the faster evolutionary rates were mostly contributed by their faster adaptation rates (Figure 5D). To further correlate the evolution of *de novo* genes with different developmental stages of the testis, we calculated the correlations between the developmental stages and gene ages of testis biased *de novo* gene candidates. We also plotted the developmental stages and gene ages of testis-biased *de novo* genes using the first two principal components defined by all testis-biased *D. melanogaster* genes (Figure S4). Interestingly, we found that for testis-biased *de novo* genes, gene ages significantly correlate with their peak developmental stages (Figure S5, Kendall tau rank correlation, p = 1e-3 and Spearman rank correlation, p = 1e-3). In agreement with this result, we found that young *de novo* genes tend to be enriched in early germline stages (cluster #1) while old *de novo* genes tend to be in mid-to-late stages (cluster #2-#4, Table S3) and the proportions of young de novo genes slightly decline from early stage (cluster #1) to late stage (cluster #4, Table S4). While previous studies emphasized the importance of mid-to-late spermatogenesis stages in *de novo* gene origination (Kaessmann, 2010; Soumillon et al., 2013; Van Oss & Carvunis, 2019; Witt et al., 2019), our results indicate that the early spermatogenesis stages might also play an unneglectable role in *de novo* gene origination.

Figure 5. Expression pattern of testis-biased *de novo* gene candidates. (A) Clustering of all *D. melanogaster* testis-biased genes. (B) Expression patterns of testis-specific (left) and testis-biased (right) *de novo* gene candidates in different clusters. (C) *de novo* gene candidates in Cluster #1 are less likely to be transmembrane (TM probability) or signal (Signal probability) proteins but are more likely to have a higher structural disorder (ISD) and higher solvent accessibility (RSA). (D) *de novo* gene candidates in Cluster #1 have higher evolutionary rates (ω), adaptation rates (ω_a), and proportions of adaptive changes (α), but not higher nonadaptation rates (ω_{na}).

Discussion

In the past decade, various synteny-based methods have been applied to determine the *de novo* emergence of *de novo* genes (Van Oss & Carvunis, 2019). However, many of them suffered from divergence between genomes in longer timescales, fragmented genome assemblies, and high genome recombination rates (Van Oss & Carvunis, 2019). Thus, in this study, we used whole-genome alignments instead of synteny mapping to determine the ortholog regions of *D. melanogaster* protein-coding genes. Specifically, we used the base-level progressive cactus aligner, which has been proved to be highly accurate in both pairwise genome alignments (Paten et al., 2011) and progressive genome alignments of an entire phylogenetic tree (Armstrong et al., 2020). With the high-quality base-level whole genome alignments, we can identify *de novo* genes with the support of alignments against non-coding sequences in outgroup species. The aligned non-coding sequences could also facilitate the discovery of

possible origination events of *de novo* genes. Despite these merits, there are still some inherent drawbacks to this method. The first is that we did not include taxonomy restricted genes (TRG) or orphan genes that cannot be aligned in progressive cactus alignments, which could underestimate the number of true *de novo* originated genes. However, we should also be aware that, although TRG or orphan genes are likely to be *de novo* emerged, without the support of noncoding alignments, we cannot rule out the possibility of other mechanisms, such as horizontal gene transfer (HGT), transposition, etc. The second is that we used simulated data to distinguish unannotated orthologs and unannotated noncoding regions. For genes with unannotated ortholog regions, we define orthology if their gene structure prediction scores were significantly beyond random expectations. In this case, unannotated ortholog regions with frameshift indels but high prediction scores might also be considered true orthologs. Thus, it is likely to assign a number of *de novo* genes to a more ancestral branch than their true origin dates. However, we argue that, according to the high average genomic mutation rates in *Drosophila* species (Haag-Liautard et al., 2007), an unannotated ortholog region with high prediction scores might be under strong purifying selection or might just be dead recently, otherwise this region would be completely random after random neutral mutations for millions of years. This compromise might be necessary as misidentification of old genes as young genes may lead to biased patterns (Moyers & Zhang, 2015, 2016b; Weisman et al., 2020). We further found gene expression evidence from five different *Drosophila* species for many of these unannotated regions (Figure S6). Note that it is still likely that we might have excluded potential genic sequences that were too diverged to have meaningful gene structure predictions (Iwata & Gotoh, 2012a). Another drawback was that the estimation of origination lineage might be affected by the unavailability of high-quality genomic data for some other species in the lineages of interest. With the great efforts being made to improve the genome sequencing and annotation qualities, we will be able to better study *de novo* gene emergence events in the future.

There were many case studies trying to characterize the foldability and structure of proteins encoded by some specific *de novo* originated genes (Bornberg-Bauer et al., 2021b; Bungard et al., 2017; Lange et al., 2021). However, it remains unclear whether *de novo* genes could be well-folded with complex structural folds, how often *de novo* genes could be well-folded and whether *de novo* genes could have novel structural folds. Here, we explored the foldability of *de novo* genes/proteins by highly accurate and efficient AlphaFold2 predictions. In general, the AlphaFold2 predictions correlated significantly with other state-of-the-art neural network approaches, such as trRosetta and RoseTTAFold, and were comparable to high resolution structure determination techniques, such as solution NMR and x-ray crystallography, especially for small proteins (Tejero et al., 2022). We found that most *de novo* genes might be partially folded or not folded, and only very few of them have the potential to be well-folded with complex structural folds (19/589), among which several (3/19) might have novel structural folds. The complex structural folds of these *de novo* gene candidates remained stable during 200 ns MD simulations (Figure 3D), which further suggested the high accuracy of AlphaFold2 predictions. Our results provided a systematic overview of the foldability of *Drosophilinae* lineage-specific *de novo* genes and highlighted the possibility that *de novo* proteins can be well-folded and even folded into novel protein folds.

The fact that *de novo* genes were preferentially expressed in testis has led to the hypothesis that *de novo* genes were "out-of-testis". The "out-of-testis" hypothesis was thought to be related to the facilitating role of the permissive chromatin state in germline cells (Majic & Payne, 2020). Similar findings were observed in our study - we found that *de novo* genes were associated with open chromatins in testis (Figure S2), and that more than half of the identified *de novo* gene candidates were biased toward testis. These findings highlight the important role of testis and chromatin state in the origination of *de novo* gene origination. In addition, an examination of another two hotspot-tissues, head, and ovary, suggested that *de novo* gene candidates were also associated with open chromatins in these two tissues (Figure S2), highlighting the non-neglectable role of open chromatins in the emergence of *de novo* genes. In agreement with previous findings that spermatocytes and spermatids were the "hotspots" for *de novo* genes (Witt et al., 2019), we found that most testis-biased *de novo* gene candidates identified here have their peak expression in late spermatogenesis stages, including spermatocytes and spermatids. Notably, we further found that gene ages of *de novo* genes significantly correlated with their developmental stages in spermatogenesis (Figure S5). Specifically, young *de novo* genes were more likely to be expressed in early spermatogenesis stages, while old *de novo* genes were more likely to be expressed in late spermatogenesis stages (Figure S5, Table S3-S4). The results might indicate that *de novo* genes that originated in late spermatogenesis stages may have higher probabilities of being retained due to a stronger fitness effect in sperm competition, while those that originated in early spermatogenesis stages were more likely to be lost during evolution. However, it could also be possible that testis biased *de novo* genes might undergo a gradual transition from early to late spermatogenesis stages during their evolutionary trajectories. In all, our results indicate an unneglectable role of early spermatogenesis stages in *de novo* gene origination. To further address the precise mechanisms, it would be necessary to conduct functional assays to examine the fitness effect of young and old *de novo* genes, which is beyond the current scope of this study.

It has been debated whether *de novo* genes change their protein structures after origination, and it is still unclear how de novo genes evolve within a limited evolutionary time scale. Some studies found that protein disorder increases with the age of *de novo* genes in *Lachancea* (Vakirlis et al., 2018), and some other studies suggested *de novo* genes evolved or adapted gradually to avoid disorder (Wilson et al., 2017). While some case studies (Bungard et al., 2017; Lange et al., 2021) and a recent study focusing on *de novo* ORFs (Schmitz et al., 2018) suggested that protein disorder and other properties hardly change after origination. Here, by comparing *de novo* genes originated from different lineages, our results supported that protein disorder and other structural properties change little after origin. Interestingly, other properties, such as GC content, evolutionary rate, and expression pattern change with the age of *de novo* genes. Specifically, compared to younger *de novo* genes, older *de novo* genes have higher GC content, slower evolutionary rate, and are more broadly expressed in various tissues. The results also indicated that GC content was not the necessary cause of high protein disorder in *de novo* genes, as protein disorder hardly changes while GC content decreases with the age of *de novo* genes. The changes in GC content might be contributed by mechanisms such as the gradual maturation of codons, which increase the translational efficiency of *de novo* genes. We noticed that *de novo* genes have increased strength of purifying

selection (decreased ω_{na}) with *de novo* gene ages, while the adaptation rates change little, which might be related to more important functions. In agreement with this result, it was reported that, at a larger evolutionary timescale, the number of PPIs and genetic interactions increased gradually with the age of genes (Abrusán, 2013a). Taken together, we proposed that, upon origination, *de novo* genes were involved in certain molecular functions with strong fitness effects, partly due to the tendency to encode signal proteins or transmembrane proteins (Vakirlis, Acar, et al., 2020). They tend to be in the periphery of cellular networks (Abrusán, 2013b), thus were more likely to be tolerated by the host. The gradual shift of sequence and expression patterns of *de novo* genes might increase their chances to be involved in protein-protein interactions (PPI), genetic interactions or other intermolecular interactions through extensive adaptive evolution (Peng et al., 2022), which can further facilitate the integration of these genes into cellular networks. On the other hand, *de novo* genes might retain their basic molecular functions by constraining similar overall structures (see section *Most potentially well-folded de novo genes are likely to be born well-folded* in Results) or key sequence motifs along their evolutionary trajectories.

Overall, our study provided a general and easy-to-use pipeline to identify lineage-specific *de novo* gene candidates. We also provided a systematic overview of the protein foldability of *Drosophilinae* lineage-specific *de novo* gene candidates. We proposed that *de novo* genes undergo gradual sequence and functional adaptation without major protein structure changes. With more high-quality genome sequencing data, and transcription and translational data for more related species in the future, we will be able to identify *de novo* genes with higher confidence and uncover more about their emergence and how they were incorporated into the genomes and interactomes.

Material and Methods

Identification of de novo gene candidates in D. melanogaster

To infer synteny, we used Progressive Cactus Genome Aligner (Armstrong et al., 2020) to align the genomes of 20 *Drosophilinae* species and 2 *Acalyptratae* species (Figure 1A). To infer homology between different protein-coding genes, we also run all-vs-all *blastp* using all the protein sequences of 20 *Acalyptratae* species along with another 8 *Arthropods* species (Figure 1A). After obtaining the synteny and homology information, we used the following workflow (Figure 1B) to identify possible candidates of *de novo* genes.

1. We used progressive cactus aligner to align the 20 genomes of species in *Acalyptratae* lineages (L1-L9, Figure 1A).
2. We used *halLiftover* to determine the syntenic ortholog region for each query protein-coding gene.
3. If the ortholog region is an annotated coding gene, we assign it to the ortholog of the query when the annotated gene has a *blastp* E-value smaller than 0.05 to the query. Annotated coding genes that do not meet this criterion might be involved in other evolutionary mechanisms (e.g., transposition, genome rearrangement, etc.) and are not considered in this work.
4. If the ortholog region is not annotated as a protein-coding gene, we assign it as an unannotated hit. For each unannotated hit, we use *Genewise* (Birney et al., 2004) and *Spaln* (Iwata & Gotoh, 2012b) to predict

gene structure from the unannotated hit using the query *D. melanogaster* gene as reference. For the predicted cDNA sequence, we align its translated protein sequence to the query using MAFFT-LINSI (Katoh et al., 2002). We assign the unannotated hit as an unannotated putative ortholog if the prediction score was significantly beyond random simulations (see next section, *Random simulations of Geneise/Spaln*, for details), otherwise we assign it as a noncoding hit.

5. Repeat step 1 to 3 for each of the orthologs identified in step 2 to search for any possible noncoding hits.

6. We assigned the query as a possible *de novo* gene candidate and inferred the origination lineage to be Li $(i \in \{0...9\})$ only when it has: (1) unannotated putative orthologs and homologous protein-coding genes in lineages Li (Fig. 1A), (2) unannotated hits, but neither uannotated putative orthologs nor homologous in more distant lineages, and (3) no homologs in the 8 distant *Arthropods* species as shown in Fig. 1A. These candidates then are subjected for further searches in all other species in UniprotKB (see below).

7. Finally, we used *jackhmmer* (S. R. Eddy, 2011) and *blastp* to search against UniProtKB sequence database to further filter out candidates that have homologs in species more distant to the inferred origination lineage defined in step 6. To control for the possibility of false positives, which were quite frequent in iterative profile searches, we conducted three independent searches as follows:

    i.   *blastp* search with E-value cutoff of 0.05.

    ii. Iterative *jackhmmer* with options "--incE 1e-5 -E 10".

    iii. Iterative *jackhmmer* with options "--incE 1e-5 -E 10". To control for false positives, after each iteration, we manually built hmm profile for next iteration by removing possible false positives with best 1 domain E-value larger than 1e-5 (S. Eddy, 1992). For each *de novo* gene candidate, we stopped the search once the search converged or reached 5 iterations.

    To further control for possible false positives, for each *de novo* gene candidate, we required a reliable homolog to appear in at least two of the above searches with a E-value cutoff of 0.001. At last, we removed candidates that have reliable homologs in species that are more distant than their inferred origination lineage obtained in step 6. The removed candidates, along with representative reliable homologs can be found in *supplementary File S4: reliable_homologs*.

8. As an example, if a protein coding gene was identified to be *de novo* and originated from L3, the gene has: (1) homologs or putative orthologs in lineage L3, (2) unannotated hits, but neither putative orthologs or homologs in species from L4 to L9, (3) no homologs in 8 distant Arthropods species, and (4) no reliable homologs in other species that are more distant than L3 in UniprotKB.

For each *de novo* gene candidate identified here, we used the longest protein isoforms as the protein sequence for further analysis.

Random simulations of Genewise/Spaln

We used random simulations to calculate the random expectations of Genewise/Spaln gene structure predictions. The random simulations were carried out as following:

1. We simulated situations with different protein length, ranging from 15, 20, 25, 50, 75, 100, 200, 500 to 1000.

2. For each protein length (N), we ran 10,000 random simulations. For each simulation, we generated a random protein sequence according to the ratio of each amino acid in *D. melanogaster*, and a random DNA sequence with 3*(N+300) base pairs according to the genome-wide GC content of *D. melanogaster* protein-coding DNA sequences. We performed Genewise or Spaln gene structure predictions using the protein as reference and the DNA sequence as target. To assess the results of the predictions, we used the score reported by spliced alignment, as the score includes the penalty for introducing splicing sites. We calculated the mean (Mean) and standard deviation (SD) of the 10,000 random simulations.

3. We fitted Mean and SD as functions of protein length (N) respectively, using a two-phase decay function using Lmfit (Newville & Stensitzki, 2018)

$$S = S0 + SpanFast * exp(-KFast * N) + SpanSlow * exp(-KSlow * N)$$

where

$$SpanFast = (S0 - Smin) * PercentFast * 0.01$$
$$SpanFast = (S0 - Smin) * (1 - PercentFast) * 0.01$$

S0 and Smin were the expected maximum and minimum spliced alignment scores. We then can predict the random expectation scores for Genewise/Spaln gene structure prediction score for any protein references with length of N by the two-phase decay function (Figure S7).

4. We computed the probability of the protein reference and DNA target being completely unrelated or non-homologous using one-tailed gaussian distribution with the inferred mean and standard deviations of Genewise/Spaln prediction scores. We inferred the target DNA sequences being potentially genic and homologous to the reference protein at a P-value of 10^{-6}.

Bioinformatic analysis and evolutionary analysis
We used RepeatMasker (Smit AF et al.,) and Dfam (Storer et al., 2021) to search for DNA repeats, including transposable elements (TEs) and simple repeats. We used ORFfinder to detect all putative ORFs that are intergenic and longer than 75 nt (coded for 25 amino acids). We included nested ORFs to better understand possible residue composition of these intergenic putative ORFs. For protein property predictions, we used deepcnf (S. Wang et al., 2016) to predict per residue probability of helix, sheet, coil, structural disorder, solvent accessibility and transmembrane probability. These properties were further normalized by protein length. We used signalp-5.0 (Almagro Armenteros et al., 2019b) to determine the probability of a protein containing a signal peptide. Sequence evolutionary rate (ω), adaptation rate (ω_a), and nonadaptation rates (ω_{na}) of *de novo* gene candidates were obtained from a previous study (Peng et al., 2022).

Gene expression patterns.

We downloaded gene expression profiles from FlyAtlas2 (Leader et al., 2018). We converted FPKM to TPM by normalizing FPKM against the summation of all FPKMs as follows:

$$TPM_i = \frac{FPKM_i}{\sum FPKM_j} \times 10^6$$

After TPM conversion, we only retained genes with expression levels larger than 0.1 TPM for further analysis. We treated male and female whole-body TPM as male and female expression levels. To describe male specificities of *D. melanogaster* genes, we first calculated Z-score by:

$$zscore = \frac{TPM(male\ expression) - TPM(female\ expression)}{\sqrt{sd^2(male\ expression) + sd^2(female\ expression)}}$$

We then calculated a normalized Z-score by:

$$zscore(norm) = \frac{zscore - \min(zscore)}{\max(zscore) - \min(zscore)}$$

To characterize tissue specificities, we calculated tau values (Yanai et al., 2005) based on the expression profiles of 27 different tissues.

Structural prediction for de novo gene candidates

We combined AlphaFold2 (Jumper et al., 2021), RoseTTAFold (Baek et al., 2021b) and trRosetta (J. Yang et al., 2020) predictions to characterize the foldability of the *de novo* gene candidates. Specifically, for AlphaFold2 predictions, if the *de novo* gene was present in AlphaFold Protein Structure Database (AlphaFold DB), we downloaded the predicted structure from the database, otherwise we ran AlphaFold v2.0.0 scripts (https://github.com/deepmind/alphafold/tree/v2.0.0) to predict the protein structure. For RoseTTAFold predictions, we used the script *run_e2e_ver.sh* to predict the protein structures. We used the per-residue confidence score (pLDDT) from AlphaFold2 and RoseTTAFold predictions to characterize the foldability of the *de novo* genes. For trRosetta predictions, we followed similar procedures as DeepMSA (C. Zhang et al., 2020) and iteratively searched through Uniclust30 (Mirdita et al., 2017), UniRef90 (Suzek et al., 2015), metaclust (Steinegger & Söding, 2018) and tara(Y. Wang et al., 2019) protein sequence databases at default E-value cutoff of 1e-3 to collect homologous or distantly related homologous sequences. Specifically, at the first stage (Uniclust30), we used hhblits (Remmert et al., 2012) to search against Uniclust30. At each stage from stage 2 to stage 4 (UniRef90, metaclust, and tara), we first used jackhmmer (S. R. Eddy, 2011) to search against the database and then used hhblits against a custom database built from the sequences generated by jackhmmer. After each stage of searching, we used MAFFT to construct a multiple sequence alignment (MSA) to ensure that the MSA is of high quality. With the final MSA as input, we generated 100 parallel structural models by trRosetta (J. Yang et al., 2020). To characterize the foldability of the *de novo* genes and the convergence of the predictions, we selected an ensemble of 20 structural models with the lowest potential energies and calculated the average pairwise TM-score (Y. Zhang & Skolnick, 2004) by TM-align (Y. Zhang & Skolnick, 2005a).

Ancestral sequence reconstruction

For each *de novo* gene, we extracted their orthologs and unannotated putative orthologs (if there are any) from the above analysis (see section *Identification of de novo gene candidates in D. melanogaster* in Material and Methods). We used MAFFT-LINSI (Katoh et al., 2002) to align the orthologous protein sequences. An initial guide tree was built based on the phylogeny relationships between the species that had orthologs. We then used the AAML module in PAML 4.9 (Z. Yang, 2007) to reconstruct the most probable ancestral sequences based on the alignments and the initial guide tree. For each lineage, we only considered amino acids that had more than 50% coverage among the species in this lineage, as the reconstruction can be unreliable at sites with alignment gaps. We further used AlphaFold2 (Jumper et al., 2021) to predict the structures of the most probable ancestral sequences.

Molecular dynamics simulations

Starting from the predicted structural models of the 19 potentially well-folded *de novo* gene candidates and their most probable ancestral states, we set up the molecular dynamics simulations using the GROMACS-2019 (Abraham et al., 2015) and the Amber ff99SB-ILDN force field (Lindorff-Larsen et al., 2010). For each system, we placed the protein in a dodecahedral box, with a minimum distance between the solute and the box boundary of 1.2 nm. We filled the simulation box with TIP3P water molecules and additional Na^+ and Cl^- ions to neutralize the system and reach the 0.15 M salt concentration. We further minimized the potential energy of the system by the steepest descent method followed by conjugate gradient method until the maximum force was smaller than $200 \text{ kJ mol}^{-1} \text{ nm}^{-1}$. For each simulation system, we performed 200 ns MD simulations. We clustered the conformations from the last 150 ns simulations into 5 clusters by the density peaks algorithm (Rodriguez & Laio, 2014). To obtain structural fold similarity during MD simulations, we used TMalign (Y. Zhang & Skolnick, 2005b) to compute the pairwise TM-score of the centers of the 5 clusters. We also referred to the center of the 5 clusters as representative structural models and the center of the top cluster as the MD refined structural model during MD simulations.

Clustering of testis-biased genes

For each testis-biased gene, we obtained the scaled expression levels at different germline cell development stages, including 1) Early spermatogonia, 2) Late spermatogonia, 3) Early spermatocytes, 4) Late spermatocytes, 5) Early spermatids, and 6) Late spermatids. We further applied principal components analysis (PCA) on the scaled expression dataset to reduce the dimensionality. We then applied a k-means clustering algorithm to cluster these genes using the first three PCs, which could explain 82.0%, 12.2%, and 3.2% of the variants, respectively. The testis-biased genes were finally clustered into four clusters (Figure S3).

Data Availability

Whole genome alignments used in this study are available at 10.6084/m9.figshare.19989395, along with the raw alignments and spliced alignments of de novo gene candidates, which include noncoding hits in outgroup species. Code and scripts for this study are available on GitHub: https://github.com/LiZhaoLab/DrosophilaDenovoGene.

References

Abraham, M. J., Murtola, T., Schulz, R., Páll, S., Smith, J. C., Hess, B., & Lindah, E. (2015). Gromacs: High performance molecular simulations through multi-level parallelism from laptops to supercomputers. *SoftwareX*, *1–2*, 19–25. https://doi.org/10.1016/j.softx.2015.06.001

Abrusán, G. (2013a). Integration of new genes into cellular networks, and their structural maturation. *Genetics*, *195*(4), 1407–1417. https://doi.org/10.1534/genetics.113.152256

Abrusán, G. (2013b). Integration of new genes into cellular networks, and their structural maturation. *Genetics*, *195*(4), 1407–1417. https://doi.org/10.1534/genetics.113.152256

Almagro Armenteros, J. J., Tsirigos, K. D., Sønderby, C. K., Petersen, T. N., Winther, O., Brunak, S., von Heijne, G., & Nielsen, H. (2019a). SignalP 5.0 improves signal peptide predictions using deep neural networks. *Nature Biotechnology*, *37*(4), 420–423. https://doi.org/10.1038/s41587-019-0036-z

Almagro Armenteros, J. J., Tsirigos, K. D., Sønderby, C. K., Petersen, T. N., Winther, O., Brunak, S., von Heijne, G., & Nielsen, H. (2019b). SignalP 5.0 improves signal peptide predictions using deep neural networks. *Nature Biotechnology*, *37*(4), 420–423. https://doi.org/10.1038/s41587-019-0036-z

Altschul, S. F., Gish, W., Miller, W., Myers, E. W., & Lipman, D. J. (1990). Basic local alignment search tool. *Journal of Molecular Biology*, *215*(3), 403–410. https://doi.org/10.1016/S0022-2836(05)80360-2

Armstrong, J., Hickey, G., Diekhans, M., Fiddes, I. T., Novak, A. M., Deran, A., Fang, Q., Xie, D., Feng, S., Stiller, J., Genereux, D., Johnson, J., Marinescu, V. D., Alföldi, J., Harris, R. S., Lindblad-Toh, K., Haussler, D., Karlsson, E., Jarvis, E. D., … Paten, B. (2020). Progressive Cactus is a multiple-genome aligner for the thousand-genome era. *Nature*, *587*(7833), 246–251. https://doi.org/10.1038/s41586-020-2871-y

Ayoub, R., & Lee, Y. (2019). Rupee: A fast and accurate purely geometric protein structure search. *PLoS ONE*, *14*(3), e0213712. https://doi.org/10.1371/journal.pone.0213712

Baek, M., DiMaio, F., Anishchenko, I., Dauparas, J., Ovchinnikov, S., Lee, G. R., Wang, J., Cong, Q., Kinch, L. N., Dustin Schaeffer, R., Millán, C., Park, H., Adams, C., Glassman, C. R., DeGiovanni, A., Pereira, J. H., Rodrigues, A. V., Van Dijk, A. A., Ebrecht, A. C., … Baker, D. (2021a). Accurate prediction of protein structures and interactions using a three-track neural network. *Science*, *373*(6557), 871–876. https://doi.org/10.1126/science.abj8754

Baek, M., DiMaio, F., Anishchenko, I., Dauparas, J., Ovchinnikov, S., Lee, G. R., Wang, J., Cong, Q., Kinch, L. N., Dustin Schaeffer, R., Millán, C., Park, H., Adams, C., Glassman, C. R., DeGiovanni, A., Pereira, J. H., Rodrigues, A. V., Van Dijk, A. A., Ebrecht, A. C., … Baker, D. (2021b). Accurate prediction of protein structures and interactions using a three-track neural network. *Science*, *373*(6557), 871–876. https://doi.org/10.1126/science.abj8754

Bateman, A., Martin, M.-J., Orchard, S., Magrane, M., Ahmad, S., Alpi, E., Bowler-Barnett, E. H., Britto, R., Bye-A-Jee, H., Cukura, A., Denny, P., Dogan, T., Ebenezer, T., Fan, J., Garmiri, P., da Costa Gonzales, L. J., Hatton-Ellis, E., Hussein, A., Ignatchenko, A., … Zhang, J. (2022). UniProt: the Universal Protein Knowledgebase in 2023. *Nucleic Acids Research*. https://doi.org/10.1093/nar/gkac1052

Begun, D. J., Lindfors, H. A., Kern, A. D., & Jones, C. D. (2007). Evidence for de Novo Evolution of Testis-Expressed Genes in the Drosophila yakuba/Drosophila erecta Clade. *Genetics*, *176*(2), 1131–1137. https://doi.org/10.1534/genetics.106.069245

Begun, D. J., Lindfors, H. A., Thompson, M. E., & Holloway, A. K. (2006). Recently Evolved Genes Identified From Drosophila yakuba and D. erecta Accessory Gland Expressed Sequence Tags. *Genetics*, *172*(3), 1675–1681.

Birney, E., Clamp, M., & Durbin, R. (2004). GeneWise and Genomewise. *Genome Research*, *14*(5), 988–995. https://doi.org/10.1101/gr.1865504

Bornberg-Bauer, E., Hlouchova, K., & Lange, A. (2021a). Structure and function of naturally evolved de novo proteins. In *Current Opinion in Structural Biology* (Vol. 68, pp. 175–183). https://doi.org/10.1016/j.sbi.2020.11.010

Bornberg-Bauer, E., Hlouchova, K., & Lange, A. (2021b). Structure and function of naturally evolved de novo proteins. In *Current Opinion in Structural Biology* (Vol. 68, pp. 175–183). https://doi.org/10.1016/j.sbi.2020.11.010

Bungard, D., Copple, J. S., Yan, J., Chhun, J. J., Kumirov, V. K., Foy, S. G., Masel, J., Wysocki, V. H., & Cordes, M. H. J. (2017). Foldability of a Natural De Novo Evolved Protein. *Structure*, *25*(11), 1687-1696.e4. https://doi.org/10.1016/j.str.2017.09.006

Cai, J., Zhao, R., Jiang, H., & Wang, W. (2008). De novo origination of a new protein-coding gene in Saccharomyces cerevisiae. *Genetics*, *179*(1), 487–496. https://doi.org/10.1534/genetics.107.084491

Chen, S., Zhang, Y. E., & Long, M. (2010). New genes in *Drosophila* quickly become essential. *Science (New York, N.Y.)*, *330*(6011), 1682–1685. https://doi.org/10.1126/science.1196380

Eddy, S. (1992). HMMER user's guide. *Department of Genetics, Washington University School of Medicine*, *2*(1), 13.

Eddy, S. R. (2011). Accelerated profile HMM searches. *PLoS Computational Biology*, *7*(10). https://doi.org/10.1371/journal.pcbi.1002195

Haag-Liautard, C., Dorris, M., Maside, X., Macaskill, S., Halligan, D. L., Charlesworth, B., & Keightley, P. D. (2007). Direct estimation of per nucleotide and genomic deleterious mutation rates in Drosophila. *Nature*, *445*, 82–85.

Heinen, T. J. A. J., Staubach, F., Häming, D., & Tautz, D. (2009). Emergence of a new gene from an intergenic region. *Current Biology : CB*, *19*(18), 1527–1531.

Iwata, H., & Gotoh, O. (2012a). Benchmarking spliced alignment programs including Spaln2, an extended version of Spaln that incorporates additional species-specific features. *Nucleic Acids Research*, *40*(20), e161. https://doi.org/10.1093/nar/gks708

Iwata, H., & Gotoh, O. (2012b). Benchmarking spliced alignment programs including Spaln2, an extended version of Spaln that incorporates additional species-specific features. *Nucleic Acids Research*, *40*(20), e161. https://doi.org/10.1093/nar/gks708

Jumper, J., Evans, R., Pritzel, A., Green, T., Figurnov, M., Ronneberger, O., Tunyasuvunakool, K., Bates, R., Žídek, A., Potapenko, A., Bridgland, A., Meyer, C., Kohl, S. A. A., Ballard, A. J., Cowie, A., Romera-Paredes, B., Nikolov, S., Jain, R., Adler, J., … Hassabis, D. (2021). Highly accurate protein structure prediction with AlphaFold. *Nature*, *596*(7873), 583–589. https://doi.org/10.1038/s41586-021-03819-2

Kaessmann, H. (2010). Origins, evolution, and phenotypic impact of new genes. *Genome Research*, *20*(10), 1313–1326.

Katoh, K., Misawa, K., Kuma, K., & Miyata, T. (2002). MAFFT: a novel method for rapid multiple sequence alignment based on fast Fourier transform. *Nucleic Acids Research*, *30*(14), 3059–3066. https://doi.org/10.1093/nar/gkf436

Knowles, D. G., & McLysaght, A. (2009). Recent de novo origin of human protein-coding genes. *Genome Research*, *19*(10), 1752–1759. https://doi.org/10.1101/gr.095026.109

Kumar, S., Stecher, G., Suleski, M., & Hedges, S. B. (2017). TimeTree: A Resource for Timelines, Timetrees, and Divergence Times. *Molecular Biology and Evolution*, *34*(7), 1812–1819. https://doi.org/10.1093/molbev/msx116

Lange, A., Patel, P. H., Heames, B., Damry, A. M., Saenger, T., Jackson, C. J., Findlay, G. D., & Bornberg-Bauer, E. (2021). Structural and functional characterization of a putative de novo gene in Drosophila. *Nature Communications*, *12*(1), 1667. https://doi.org/10.1038/s41467-021-21667-6

Leader, D. P., Krause, S. A., Pandit, A., Davies, S. A., & Dow, J. A. T. (2018). FlyAtlas 2: A new version of the Drosophila melanogaster expression atlas with RNA-Seq, miRNA-Seq and sex-specific data. *Nucleic Acids Research*, *46*(D1), D809–D815. https://doi.org/10.1093/nar/gkx976

Levine, M. T., Jones, C. D., Kern, A. D., Lindfors, H. A., & Begun, D. J. (2006). Novel genes derived from noncoding DNA in Drosophila melanogaster are frequently X-linked and exhibit testis-biased expression. *Proceedings of the National Academy of Sciences of the United States of America*, *103*(26), 9935–9939. https://doi.org/10.1073/pnas.0509809103

Li, D., Dong, Y., Jiang, Y., Jiang, H., Cai, J., & Wang, W. (2010). A de novo originated gene depresses budding yeast mating pathway and is repressed by the protein encoded by its antisense strand. *Cell Research*. https://doi.org/10.1038/cr.2010.31

Li, L., Stoeckert, C. J., & Roos, D. S. (2003). OrthoMCL: identification of ortholog groups for eukaryotic genomes. *Genome Research*, *13*(9), 2178–2189. https://doi.org/10.1101/gr.1224503

Lindorff-Larsen, K., Piana, S., Palmo, K., Maragakis, P., Klepeis, J. L., Dror, R. O., & Shaw, D. E. (2010). Improved side-chain torsion potentials for the Amber ff99SB protein force field. *Proteins: Structure, Function and Bioinformatics*, *78*(8), 1950–1958. https://doi.org/10.1002/prot.22711

Majic, P., & Payne, J. L. (2020). Enhancers facilitate the birth of de novo genes and gene integration into regulatory networks. *Molecular Biology and Evolution*, *37*(4), 1165–1178. https://doi.org/10.1093/molbev/msz300

McLysaght, A., & Hurst, L. D. (2016). Open questions in the study of de novo genes: what, how and why. *Nature Reviews Genetics*, *17*(9), 567–578. https://doi.org/10.1038/nrg.2016.78

Mirdita, M., Von Den Driesch, L., Galiez, C., Martin, M. J., Soding, J., & Steinegger, M. (2017). Uniclust databases of clustered and deeply annotated protein sequences and alignments. *Nucleic Acids Research*, *45*(D1), D170–D176. https://doi.org/10.1093/nar/gkw1081

Moyers, B. A., & Zhang, J. (2015). Phylostratigraphic bias creates spurious patterns of genome evolution. *Molecular Biology and Evolution*, *32*(1), 258–267. https://doi.org/10.1093/molbev/msu286

Moyers, B. A., & Zhang, J. (2016a). Evaluating Phylostratigraphic Evidence for Widespread de Novo Gene Birth in Genome Evolution. *Molecular Biology and Evolution*, *33*(5), 1245–1256. https://doi.org/10.1093/molbev/msw008

Moyers, B. A., & Zhang, J. (2016b). Evaluating Phylostratigraphic Evidence for Widespread de Novo Gene Birth in Genome Evolution. *Molecular Biology and Evolution*, *33*(5), 1245–1256. https://doi.org/10.1093/molbev/msw008

Necci, M., Piovesan, D., Hoque, M. T., Walsh, I., Iqbal, S., Vendruscolo, M., Sormanni, P., Wang, C., Raimondi, D., Sharma, R., Zhou, Y., Litfin, T., Galzitskaya, O. V., Lobanov, M. Y., Vranken, W., Wallner, B., Mirabello, C., Malhis, N., Dosztányi, Z., … Tosatto, S. C. E. (2021). Critical assessment of protein intrinsic disorder prediction. *Nature Methods*, *18*(5), 472–481. https://doi.org/10.1038/s41592-021-01117-3

Newville, M., & Stensitzki, T. (2018). Non-Linear Least-Squares Minimization and Curve-Fitting for Python. *Non-Linear Least-Squares Minimization and Curve-Fitting for Python*, https://doi.org/10.5281/zenodo.11813. https://doi.org/10.5281/ZENODO.11813

Paten, B., Earl, D., Nguyen, N., Diekhans, M., Zerbino, D., & Haussler, D. (2011). Cactus: Algorithms for genome multiple sequence alignment. *Genome Research*, *21*(9), 1512–1528. https://doi.org/10.1101/gr.123356.111

Peng, J., Svetec, N., & Zhao, L. (2022). Intermolecular Interactions Drive Protein Adaptive and Coadaptive Evolution at Both Species and Population Levels. *Molecular Biology and Evolution*, *39*(1), msab350. https://doi.org/10.1093/molbev/msab350

Remmert, M., Biegert, A., Hauser, A., & Söding, J. (2012). HHblits: Lightning-fast iterative protein sequence searching by HMM-HMM alignment. *Nature Methods*, *9*(2), 173–175. https://doi.org/10.1038/nmeth.1818

Rodriguez, A., & Laio, A. (2014). Clustering by fast search and find of density peaks. *Science*, *344*(6191), 1492–1496. https://doi.org/10.1126/science.1242072

Ruiz-Orera, J., Hernandez-Rodriguez, J., Chiva, C., Sabidó, E., Kondova, I., Bontrop, R., Marqués-Bonet, T., & Albà, M. M. (2015). Origins of De Novo Genes in Human and Chimpanzee. *PLOS Genetics*, *11*(12), e1005721. https://doi.org/10.1371/journal.pgen.1005721

Schmitz, J. F., Ullrich, K. K., & Bornberg-Bauer, E. (2018). Incipient de novo genes can evolve from frozen accidents that escaped rapid transcript turnover. *Nature Ecology and Evolution*, *2*(10), 1626–1632. https://doi.org/10.1038/s41559-018-0639-7

Smit AF, Hubley R, & Green P. (n.d.). *RepeatMasker Open-4.0. 2013-2015*. http://www.repeatmasker.org

Soumillon, M., Necsulea, A., Weier, M., Brawand, D., Zhang, X., Gu, H., Barthès, P., Kokkinaki, M., Nef, S., Gnirke, A., Dym, M., deMassy, B., Mikkelsen, T. S., & Kaessmann, H. (2013). Cellular Source and Mechanisms of High Transcriptome Complexity in the Mammalian Testis. *Cell Reports*, *3*(6), 2179–2190. https://doi.org/10.1016/j.celrep.2013.05.031

Steinegger, M., & Söding, J. (2018). Clustering huge protein sequence sets in linear time. *Nature Communications*, *9*(1), 2542. https://doi.org/10.1038/s41467-018-04964-5

Storer, J., Hubley, R., Rosen, J., Wheeler, T. J., & Smit, A. F. (2021). The Dfam community resource of transposable element families, sequence models, and genome annotations. *Mobile DNA*, *12*(1), 2. https://doi.org/10.1186/s13100-020-00230-y

Suzek, B. E., Wang, Y., Huang, H., McGarvey, P. B., & Wu, C. H. (2015). UniRef clusters: A comprehensive and scalable alternative for improving sequence similarity searches. *Bioinformatics*, *31*(6), 926–932. https://doi.org/10.1093/bioinformatics/btu739

Tejero, R., Huang, Y. J., Ramelot, T. A., & Montelione, G. T. (2022). AlphaFold Models of Small Proteins Rival the Accuracy of Solution NMR Structures. *BioRxiv*, 2022.03.09.483701. https://doi.org/10.1101/2022.03.09.483701

Tunyasuvunakool, K., Adler, J., Wu, Z., Green, T., Zielinski, M., Žídek, A., Bridgland, A., Cowie, A., Meyer, C., Laydon, A., Velankar, S., Kleywegt, G. J., Bateman, A., Evans, R., Pritzel, A., Figurnov, M., Ronneberger, O., Bates, R., Kohl, S. A. A., … Hassabis, D. (2021). Highly accurate protein structure prediction for the human proteome. *Nature*, *596*(7873), 590–596. https://doi.org/10.1038/s41586-021-03828-1

Vakirlis, N., Acar, O., Hsu, B., Castilho Coelho, N., Van Oss, S. B., Wacholder, A., Medetgul-Ernar, K., Bowman, R. W., Hines, C. P., Iannotta, J., Parikh, S. B., McLysaght, A., Camacho, C. J., O'Donnell, A. F., Ideker, T., & Carvunis, A.-R. (2020). De novo emergence of adaptive membrane proteins from thymine-rich genomic sequences. *Nature Communications*, *11*(1), 781. https://doi.org/10.1038/s41467-020-14500-z

Vakirlis, N., Carvunis, A.-R., & McLysaght, A. (2020). Synteny-based analyses indicate that sequence divergence is not the main source of orphan genes. *ELife*, *9*, e53500. https://doi.org/10.7554/eLife.53500

Vakirlis, N., Hebert, A. S., Opulente, D. A., Achaz, G., Hittinger, C. T., Fischer, G., Coon, J. J., & Lafontaine, I. (2018). A Molecular Portrait of De Novo Genes in Yeasts. *Molecular Biology and Evolution*, *35*(3), 631–645. https://doi.org/10.1093/molbev/msx315

Van Oss, S. B., & Carvunis, A.-R. (2019). De novo gene birth. *PLOS Genetics*, *15*(5), e1008160. https://doi.org/10.1371/journal.pgen.1008160

Varadi, M., Anyango, S., Deshpande, M., Nair, S., Natassia, C., Yordanova, G., Yuan, D., Stroe, O., Wood, G., Laydon, A., Žídek, A., Green, T., Tunyasuvunakool, K., Petersen, S., Jumper, J., Clancy, E., Green, R., Vora, A., Lutfi, M., … Velankar, S. (2021). AlphaFold Protein Structure Database: massively expanding the structural coverage of protein-sequence space with high-accuracy models. *Nucleic Acids Research*. https://doi.org/10.1093/nar/gkab1061

Wang, S., Li, W., Liu, S., & Xu, J. (2016). RaptorX-Property: a web server for protein structure property prediction. *Nucleic Acids Research*, *44*(W1), W430–W435. https://doi.org/10.1093/nar/gkw306

Wang, Y., Shi, Q., Yang, P., Zhang, C., Mortuza, S. M., Xue, Z., Ning, K., & Zhang, Y. (2019). Fueling ab initio folding with marine metagenomics enables structure and function predictions of new protein families. *Genome Biology*, *20*(1), 229. https://doi.org/10.1186/s13059-019-1823-z

Weisman, C. M., Murray, A. W., & Eddy, S. R. (2020). Many, but not all, lineage-specific genes can be explained by homology detection failure. *PLoS Biology*, *18*(11), e3000862. https://doi.org/10.1371/journal.pbio.3000862

Wilson, B. A., Foy, S. G., Neme, R., & Masel, J. (2017). Young Genes are Highly Disordered as Predicted by the Preadaptation Hypothesis of De Novo Gene Birth. *Nature Ecology & Evolution*, *1*(6), 0146–146. https://doi.org/10.1038/s41559-017-0146

Witt, E., Benjamin, S., Svetec, N., & Zhao, L. (2019). Testis single-cell RNA-seq reveals the dynamics of de novo gene transcription and germline mutational bias in Drosophila. *ELife*, *8*, e47138. https://doi.org/10.7554/eLife.47138

Witt, E., Svetec, N., Benjamin, S., & Zhao, L. (2021). Transcription Factors Drive Opposite Relationships between Gene Age and Tissue Specificity in Male and Female *Drosophila* Gonads. *Molecular Biology and Evolution*, *38*(5), 2104–2115. https://doi.org/10.1093/molbev/msab011

Yanai, I., Benjamin, H., Shmoish, M., Chalifa-Caspi, V., Shklar, M., Ophir, R., Bar-Even, A., Horn-Saban, S., Safran, M., Domany, E., Lancet, D., & Shmueli, O. (2005). Genome-wide midrange transcription profiles reveal expression level relationships in human tissue specification. *Bioinformatics*, *21*(5), 650–659. https://doi.org/10.1093/bioinformatics/bti042

Yang, J., Anishchenko, I., Park, H., Peng, Z., Ovchinnikov, S., & Baker, D. (2020). Improved protein structure prediction using predicted interresidue orientations. *Proceedings of the National Academy of Sciences of the United States of America*, *117*(3), 1496–1503. https://doi.org/10.1073/pnas.1914677117

Yang, Z. (2007). PAML 4: Phylogenetic analysis by maximum likelihood. *Molecular Biology and Evolution*, *24*(8), 1586–1591. https://doi.org/10.1093/molbev/msm088

Zhang, C., Zheng, W., Mortuza, S. M., Li, Y., & Zhang, Y. (2020). DeepMSA: Constructing deep multiple sequence alignment to improve contact prediction and fold-recognition for distant-homology proteins. *Bioinformatics*, *36*(7), 2105–2112. https://doi.org/10.1093/bioinformatics/btz863

Zhang, L., Ren, Y., Yang, T., Li, G., Chen, J., Gschwend, A. R., Yu, Y., Hou, G., Zi, J., Zhou, R., Wen, B., Zhang, J., Chougule, K., Wang, M., Copetti, D., Peng, Z., Zhang, C., Zhang, Y., Ouyang, Y., … Long, M. (2019). Rapid evolution of protein diversity by de novo origination in Oryza. *Nature Ecology & Evolution*, *3*(4), 679–690. https://doi.org/10.1038/s41559-019-0822-5

Zhang, Y., & Skolnick, J. (2004). Scoring function for automated assessment of protein structure template quality. *Proteins: Structure, Function and Genetics*, *57*(4), 702–710. https://doi.org/10.1002/prot.20264

Zhang, Y., & Skolnick, J. (2005a). TM-align: A protein structure alignment algorithm based on the TM-score. *Nucleic Acids Research*, *33*(7), 2302–2309. https://doi.org/10.1093/nar/gki524

Zhang, Y., & Skolnick, J. (2005b). TM-align: A protein structure alignment algorithm based on the TM-score. *Nucleic Acids Research*, *33*(7), 2302–2309. https://doi.org/10.1093/nar/gki524

Zhao, L., Saelao, P., Jones, C. D., & Begun, D. J. (2014). Origin and spread of de novo genes in Drosophila melanogaster populations. *Science*, *343*(6172), 769–772. https://doi.org/10.1126/science.1248286

Zheng, E. B., & Zhao, L. (2022). Protein evidence of unannotated ORFs in Drosophila reveals diversity in the evolution and properties of young proteins. *ELife*, *11*. https://doi.org/10.7554/eLife.78772