1 **Title**
2 Aversive experience drives offline ensemble reactivation to link memories across days
3
4
5 **Authors**

6 Yosif Zaki[1], Zachary T. Pennington[1], Denisse Morales-Rodriguez[1], Taylor R. Francisco[1], Alexa R.
7 LaBanca[1], Zhe Dong[1], Sophia Lamsifer[1], Simón Carrillo Segura[2], Hung-Tu Chen[4], Zoé Christenson
8 Wick[1], Alcino J. Silva[3], Matthijs van der Meer[4], Tristan Shuman[1], André Fenton[5,6], Kanaka Rajan[1],
9 Denise J. Cai[1*]
10
[1]Nash Department of Neuroscience, Icahn School of Medicine at Mount Sinai, New York, NY, 10029
[2]Graduate Program in Mechanical and Aerospace Engineering, Tandon School of Engineering, New
York University, Brooklyn, NY, 11201
[3]Department of Neurobiology, Psychiatry & Biobehavioral Sciences, and Psychology, Integrative Center
for Learning and Memory, Brain Research Institute, UCLA, Los Angeles, CA 90095
[4]Department of Psychological & Brain Sciences, Dartmouth College, Hanover, NH, 03755
[5]Center for Neural Science, New York University, New York, NY, 10003
[6]Neuroscience Institute at the NYU Langone Medical Center, New York, NY, 10016
19
20
Correspondence:
Denise J. Cai
Nash Department of Neuroscience, Icahn School of Medicine at Mount Sinai
One Gustave L. Levy Place
Box 1639
New York, NY 10029
Email: denisecai@gmail.com

28
29
30
31 **Keywords**
32 hippocampus, memory integration, memory-linking, ensemble, reactivation, co-firing, offline periods,
33 stress, PTSD

Abstract

Memories are encoded in neural ensembles during learning and stabilized by post-learning reactivation. Integrating recent experiences into existing memories ensures that memories contain the most recently available information, but how the brain accomplishes this critical process remains unknown. Here we show that in mice, a strong aversive experience drives the offline ensemble reactivation of not only the recent aversive memory but also a neutral memory formed two days prior, linking the fear from the recent aversive memory to the previous neutral memory. We find that fear specifically links retrospectively, but not prospectively, to neutral memories across days. Consistent with prior studies, we find reactivation of the recent aversive memory ensemble during the offline period following learning. However, a strong aversive experience also increases co-reactivation of the aversive and neutral memory ensembles during the offline period. Finally, the expression of fear in the neutral context is associated with reactivation of the shared ensemble between the aversive and neutral memories. Taken together, these results demonstrate that strong aversive experience can drive retrospective memory-linking through the offline co-reactivation of recent memory ensembles with memory ensembles formed days prior, providing a neural mechanism by which memories can be integrated across days.

Main Text

Individual memories are initially encoded by ensembles of cells active during a learning event[1-5] and are stabilized during offline periods following learning through reactivation of those ensembles[6-17]. These reactivations often occur in brief synchronous bursts, which are necessary to drive memory consolidation[18-20]. Most research on episodic memory has focused on how the brain maintains stable representations of discrete memories; however, animals are constantly aggregating new memories and updating past memories as new, relevant information is learned[21]. Moreover, most studies of associative learning have focused on cues that directly precede or occur with an outcome. However, oftentimes in nature, a predictor may not immediately precede an outcome but animals are nonetheless capable of learning to make an inference about the association (e.g., conditioned taste aversion)[22]. It is unclear the environmental variables that could promote memories to be linked across long periods (i.e.,

days), and the neural mechanisms of memory integration across such disparate time periods are poorly

understood. In addition, while it has been shown that offline periods support memory consolidation,

recent studies have suggested that offline periods following learning may be important for memory

integration processes as well[23-26].

Strong aversive experience drives retrospective memory-linking

To investigate how memories are integrated across days, we first designed a behavioral

experiment to test whether mice would spread fear from an aversive memory to a neutral memory

formed two days prior (Retrospective memory-linking) or two days after (Prospective memory-linking)

(Figure 1A). In the Retrospective group, mice first experienced a Neutral context followed by an

Aversive context paired with a foot shock two days later. In the Prospective group, mice experienced an

Aversive context followed by a Neutral context two days later. Both groups were then tested in the

Aversive context to test for recall of the aversive memory, followed by testing in the previously

experienced Neutral context or an unfamiliar Novel context to test for non-specific fear generalization.

Memory-linking was defined as a selective increase in fear in the Neutral context compared to the

Novel context, both contexts in which they had never been shocked. Notably, this definition

distinguishes memory-linking from a broader generalization of fear across contexts. Mice froze no

differently in the Aversive context in either group, suggesting that the perceived negative valence of the

Aversive context was not different between groups (Figure 1B). Interestingly, in the Retrospective

group, mice froze more in the Neutral context compared to the Novel context, suggesting that fear

spread retrospectively from the Aversive context to the Neutral context experienced two days prior.

However, in the Prospective group, there was no difference in freezing between the Neutral and Novel

contexts, suggesting that memory-linking between the Aversive and Neutral contexts did not occur

prospectively across days (Figure 1C). Consistent with prior studies, mice froze in the Neutral context in

both Prospective and Retrospective conditions when the Neutral and Aversive contexts were

experienced within a day (5h apart, Extended Figure 1A)[27,28]. However, when the contexts were

separated by more than one day, mice froze in the Neutral context only in the Retrospective and not the

Prospective condition (Extended Figure 1B).

We next asked what environmental variables drove two memories to be linked retrospectively

across days. It has previously been suggested that the emotional salience of an experience enhances

its storage into memory[29,30], as well as its likelihood of altering past neutral memories in humans[31].

Thus, we hypothesized that the more aversive the experience, the more likely that fear would be

retrospectively linked to a previous neutral memory. To test this, we manipulated the shock intensity

during aversive encoding to test if stronger shock would drive retrospective memory-linking (Figure 1D).

Mice were exposed to a Neutral context followed by an Aversive context paired with a low or high shock

two days later (Low Shock group & High Shock group). Mice were then tested in the Aversive, Neutral,

and a Novel context in the subsequent three days. As expected, mice in the High Shock group froze

more than mice in the Low Shock group during recall in the Aversive context (Figure 1E). We found that

only High Shock mice exhibited a selective increase in freezing in the previously experienced Neutral

context relative to the Novel context during recall (Figure 1F; Extended Figure 1C-E). If the perceived

aversiveness of an experience affects the likelihood of retrospective memory-linking, we hypothesized

that levels of freezing during Aversive memory recall would positively correlate with memory-linking—

defined as the difference between freezing in the Neutral context and in the Novel context. Indeed, in

the High Shock mice, the freezing during Aversive context recall positively correlated with the degree of

memory-linking (Figure 1G).

We next investigated how the brain links recent aversive memories with past neutral memories

formed days prior. It has been well established in rodents and humans that memories are reactivated

during restful periods following learning (i.e., offline periods) to promote the storage of recently learned

information[17,32-34]. However, recent work in humans has shown that offline periods can drive the

integration of discrete memories as well[23,35,36]. Thus, we hypothesized that following an aversive

experience (High Shock group), the offline period may be serving not only to support the consolidation

of the aversive memory, but also to link the recent aversive memory with the prior neutral memory, thus

increasing freezing during recall of the Neutral context. A major site of memory formation in the brain is

the hippocampus, where rapid plasticity following an experience promotes the formation of a memory

for that experience and reflects memory expression thereafter[18,27,37-39]. Thus, we asked whether

hippocampal activity during the offline period following Aversive encoding was necessary to drive

retrospective memory-linking. To do this, we used a chemogenetic manipulation system to disrupt

endogenous hippocampal activity during the offline period following Aversive encoding paired with a

strong shock (Extended Figure 2). We predicted that this would disrupt retrospective memory-linking.

Prior studies have shown that PSAM4-GlyR (PSAM) is an inhibitory ionotropic receptor with no

endogenous ligand, and binding of the PSEM ligand with the PSAM receptor causes robust

hyperpolarization in neurons[40]. We injected mice with a pan-neuronal, PSAM4-GlyR-expressing virus

bilaterally in hippocampus and during the offline period immediately following Aversive encoding, we

administered either PSEM to manipulate offline hippocampal activity, or injected saline as a control. We

found that mice that received saline during the offline period exhibited a selective increase in freezing in

the Neutral over the Novel context, demonstrating retrospective memory-linking. In contrast, mice that

received PSEM no longer showed this selective increase in freezing in the Neutral context (Extended

Figure 2A-C). To ensure that this effect on retrospective memory-linking was not due to a disrupted

memory for the Aversive context, we repeated the experiment, administering PSEM or saline during the

offline period, and then tested mice in the Aversive context. We found that mice that received PSEM

froze no differently compared to saline controls during Aversive memory recall, suggesting that the

strong aversive memory was left intact (Extended Figure 2D,E). These results suggest that

hippocampal activity during the offline period is necessary to drive retrospective memory-linking.

 Figure 1

Prospective vs. Retrospective Memory-Linking

Low Shock vs. High Shock Retrospective Memory-Linking

Figure 1. Strong aversive experience drives retrospective memory-linking to a neutral context learned days ago. A) Schematic of prospective vs retrospective memory-linking behavior experiment. Mice either received a Neutral experience followed by an Aversive experience two days later (Retrospective) or the Aversive experience followed by Neutral (Prospective). One day after the second experience, mice were tested in the Aversive context they were shocked in. The following day, mice were tested in either the previously experienced Neutral context or a Novel context.

B) Freezing during Aversive recall in Prospective vs Retrospective groups. There was no difference in Aversive recall freezing between Prospective & Retrospective conditions (t_{34} = 0.36, p = 0.72) (*Retrospective, N = 16 mice; Prospective, N = 20 mice*).

C) Freezing during Neutral vs Novel recall in Prospective vs Retrospective groups. There was a significant interaction between freezing in Neutral vs Novel recall in the Retrospective vs Prospective groups, suggesting the Aversive experience retrospectively linked to the Neutral memory, but not prospectively. Significant interaction between Direction (Prospective vs Retrospective) and Context (Neutral vs Novel), ($F_{1,32}$ = 4.90, p = 0.034) (*Retrospective Neutral, N = 8 mice; Retrospective Novel, N = 8 mice; Prospective Neutral, N = 12 mice, Prospective Novel, N = 8 mice*). Post-hoc, *Retrospective (t_{32} = 2.586, p = 0.029), Prospective (t_{32} = 0.452, p = 0.6546).*

D) Schematic of Low Shock vs High Shock retrospective memory-linking experiment. Mice received a Neutral experience followed by a 1hr offline session in their homecage. Two days later, they received either 3 low shocks (0.25mA) or 3 high shocks (1.5mA, same amplitude as in Figure 1A) in an Aversive context, followed by another 1hr offline session in their homecage. The following day they were tested in the Aversive context, and for the following two days they were tested in the Neutral and Novel contexts, counterbalanced. Calcium imaging was performed during all the sessions.

E) Freezing during Aversive recall in Low vs High Shock mice. Mice froze more in the Aversive context after receiving a high shock vs low shock ($t_{18.8}$ = 5.877, p = 0.000012) (*Low Shock, N = 10 mice; High Shock, N = 12 mice*).

F) Freezing during Neutral vs Novel recall in Low vs High Shock mice. Mice only displayed enhanced freezing in Neutral vs Novel (i.e., retrospective memory-linking) after High Shock and not Low Shock. Significant effect of Context (Neutral vs Novel) ($F_{1,20}$ = 17.32, p = 0.000048) and significant interaction between Context and Amplitude ($F_{1,20}$ = 4.99, p = 0.037) (*Low Shock, N = 10 mice; High Shock, N = 12 mice*). High Shock mice froze more in the Neutral vs Novel contexts (t_{11} = 4.37, p = 0.002) while Low Shock mice froze no differently in the two contexts (t_9 = 1.23, p = 0.249).

G) Correlation between Aversive recall freezing and memory-linking strength. The strength of the aversive memory was correlated with the degree of retrospective memory-linking in High Shock mice (R^2 = 0.45, p = 0.016), but not in Low Shock mice (R^2 = 0.0003, p = 0.963) (*Low Shock, N = 10 mice; High Shock, N = 12 mice*).

Strong aversive learning drives offline reactivation of a past neutral ensemble

Previous work has suggested that memory reactivation during offline periods following learning could promote not only the consolidation of recently formed memories, but also support the integration of memories[23,25,26,35,36,41]. Consistent with previous studies, we expected that during the offline period following Aversive encoding (while mice are in their homecage), the ensemble active during Aversive encoding would be reactivated to drive consolidation of the recently learned aversive memory. However, we also hypothesized that if the aversive experience was strong enough, the ensemble active during the neutral experience (from two days prior) would be reactivated as well, integrating the neutral and aversive memories.

We first validated that we could detect ensemble reactivation after a salient experience using calcium imaging. To do this, we conducted a contextual fear conditioning experiment, recording hippocampal CA1 calcium dynamics using the open-source UCLA Miniscopes[27] (Extended Figure 3A,B). We recorded during Aversive encoding, the first hour offline following Aversive encoding, and during recall of the Aversive context and exposure to a Novel context. Consistent with previous literature, we found that the ensemble of cells active during Aversive encoding was reactivated offline and preferentially reactivated during Aversive memory recall, suggesting a stable neural memory ensemble (Extended Figure 3C-K).

To next investigate whether a strong aversive experience was driving offline reactivation of ensembles representing both the aversive and neutral memories, we performed calcium imaging recordings in CA1 during the offline periods following the initial Neutral experience (Offline1) and subsequent Aversive experience (Offline2) in both Low and High Shock groups (Figure 2; Extended Figure 4; same experiment as in Figure 1D). Consistent with the literature[18,20] and with our previous experiment (Extended Figure 3), following the initial Neutral encoding, the cells that were active during that experience (Neutral ensemble) were more active compared with cells not active during Neutral encoding (Remaining ensemble) in both Low and High Shock groups (Figure 2B, line graphs). There was no difference in the fraction of cells that made up the Neutral ensemble in the Low vs High Shock groups (Figure 2B, pie charts). To measure ensemble reactivation during the offline period after Aversive encoding, we separated cells that were active during the offline period into four ensembles

based on when those cells were previously active: Neutral ensemble represented cells active during the

initial Neutral encoding and not Aversive encoding; Aversive ensemble represented cells active during

Aversive encoding and not Neutral encoding; Neutral ∩ Aversive ensemble represented cells that were

active during both Neutral and Aversive encoding; and Remaining ensemble represented cells not

observed to be active prior to the offline period (Figure 2C). There was no difference in the fraction of

cells that made up each ensemble across Low and High Shock groups (Figure 2C, pie charts). In the

Low Shock group, consistent with prior literature[14], we found the Aversive ensemble, the Neutral

ensemble, and the Neutral ∩ Aversive ensemble had higher calcium activity than the Remaining

ensemble. And the Neutral ensemble was less active than the Aversive and Neutral ∩ Aversive

ensembles (Figure 2C, line graphs, left side). These results are consistent with prior studies

demonstrating offline reactivation of neuronal ensembles that were recently active during learning[7-9]. In

contrast, in the High Shock group, the Neutral ensemble was no differently active than the Aversive and

Neutral ∩ Aversive ensembles (Figure 2C, line graphs, right side), indicating that the high shock

increased reactivation of the Neutral ensemble.

Since the Neutral ensemble was more highly reactivated after high shock, we next investigated

whether the Neutral, Aversive, and Neutral ∩ Aversive ensembles might be firing together on a finer

temporal scale. Hippocampal activity is known to exhibit organized bursts, oftentimes accompanied by

sharp-wave ripples in the local field potential, during which cells active during learning are preferentially

reactivated[18]. These events have been found to support memory consolidation[18-20]. Although calcium

dynamics are of a coarser timescale than sharp-wave ripples, we observed that during the offline

recordings, hippocampal calcium activity periodically exhibited brief bursts of activity during which

numerous cells were co-active (Extended Figure 5A,B, from our validation study in Extended Figure 3),

consistent with previous reports[42,43]. We found that these bursts were unlikely to occur from shuffled

neuronal activities, suggesting that these events were organized events during which groups of

hippocampal neurons were synchronously active (Extended Figure 5C-F). We isolated these brief burst

periods to ask whether ensembles that were previously active during encoding were selectively

participating in these brief burst events (Figure 2D-I; Extended Figure 5A,B; see Methods). We first

measured these burst events after a single Aversive learning experience and found that a larger

fraction of Aversive ensemble cells participated in these events than the Remaining ensemble cells

(Extended Figure 5L). Interestingly, these burst events coincided with the mouse briefly slowing down

about 1 second prior to the event, and about 1 second after its onset resuming its locomotion,

suggesting that these bursts occurred during periods of brief quiescence (Extended Figure 5I,J)[18].

We then asked whether a strong shock paired with an Aversive context would drive the Neutral

ensemble to also participate within these bursts after Aversive encoding (experiment from Figure 1D).

In both Low and High Shock mice and after both Neutral and Aversive encoding, frequencies of burst

events (defined by periods when the mean activity of the entire recorded population reached above a

required threshold; see Methods) were comparable across groups and decreased across the hour

(Extended Figure 4G,H). As expected, after Neutral encoding, both Low and High Shock groups had a

larger fraction of the Neutral ensemble participating in these burst events than the Remaining ensemble

(Figure 2D,F). After Aversive encoding, both groups again showed selective participation of the

Aversive ensemble that was most recently active (Figure 2G) as well as of the Neutral ∩ Aversive

ensemble that was previously active during both learning events (Figure 2H). However, only in the High

Shock group (and not the Low Shock group) the Neutral ensemble selectively participated in these

burst events as well (Figure 2I), suggesting that a strong aversive experience drove the recruitment of

the Neutral ensemble into these burst events.

 Figure 2

Figure 2. Strong aversive experience drives reactivation of a past neutral ensemble.

A) Representative histology (left) of GCaMP6f expression in hippocampal CA1, imaged with a confocal microscope. Green represents AAV1-Syn-GCaMP6f expression, while blue represents a cellular DAPI stain. Maximum intensity projection of an example mouse across one recording session, imaged with a Miniscope (middle), with the spatial footprints of all recorded cells during that session (right) randomly color-coded.

B) During Offline1 after Neutral encoding, cells that were active during Neutral encoding (Neutral ensemble) made up ~25-30% of the offline cell population (pie charts) ($X^2 = 0.122$, $df = 1$, $p = 0.73$). The Neutral ensemble was more highly active than the Remaining ensemble during the offline period (line graphs; A.U.). There was a main effect of Ensemble ($F_{1,159} = 59.19$, $p = 1.4e-12$), no effect of Amplitude ($F_{1,13} = 0.039$, $p = 0.85$), and an effect of Time ($F_{1,159} = 4.33$, $p = 0.039$), and all interactions $p > 0.05$ (*Low Shock, N = 7 mice; High Shock, N = 8 mice*).

C) During Offline2 after Aversive encoding, similar proportions of previously active cells were reactivated across Low and High shock groups (pie charts) ($X^2 = 0.326$, $df = 3$, $p = 0.955$). However, ensembles were differentially reactivated based upon the amplitude of the Aversive experience (*Ensemble x Amplitude: $F_{3,331} = 5.36$, $p = 0.0013$*) (line graphs; A.U.). In Low Shock mice, the Neutral, Aversive, and Neutral ∩ Aversive ensembles were more highly active than the Remaining ensemble (*contrast, $t_{18} = 4.22$, $p = 0.0005$*). Additionally, these ensembles were differentially active relative to one another ($F_{2,12} = 4.03$, $p = 0.046$). This was driven by the Neutral ensemble being less active. The Neutral ensemble was less active than the Aversive and Neutral ∩ Aversive ensembles ($t_{12} = 2.83$, $p = 0.03$) while the Aversive ensemble was no differently active than the Neutral ∩ Aversive ensemble ($t_{12} = 0.19$, $p = 0.85$). In High Shock mice, the Neutral, Aversive, and Neutral ∩ Aversive ensembles were all more highly active than the Remaining ensemble ($t_{21} = 4.36$, $p = 0.0003$), but these three ensembles were no differently active from each other ($F_{2,14} = 1.52$, $p = 0.25$) (*Low Shock, N = 7 mice; High Shock, N = 8 mice*).

D) During the offline periods, hippocampal activity displayed brief bursts of neural activity. To detect these bursts, we computed the z-scored mean activity of the entire recorded population and applied a threshold of z=2 and defined burst periods as all the timepoints above this threshold. The left raster represents an example burst period during Offline1, during which mean population activity briefly reached above threshold. Each row of the raster represents the activity of every recorded neuron, color-coded based on the ensemble it was a part of (blue represents Neutral ensemble and grey represents Remaining ensemble; see legend in Figure 2B). The top black trace represents the z-scored mean population activity. The right raster represents an example non-burst period.

E) Same as D but an example burst and non-burst period for Offline2. Each row of the raster again is color-coded based on the ensemble it was a part of (Aversive in red, Neutral ∩ Aversive in purple, Neutral in blue, and Remaining in grey; see legend in Figure 2C).

F) During Offline1 in both Low and High Shock groups, a larger fraction of the Neutral ensemble participated in bursts than the Remaining ensemble did (*Ensemble: $F_{1,13} = 16.33$, $p = 0.001$; Amplitude: $F_{1,13} = 0.009$, $p = 0.925$; Ensemble x Amplitude: $F_{1,13} = 0.0058$, $p = 0.940$*) (*Low Shock, N = 7 mice; High Shock, N = 8 mice*).

G) During Offline2 in both Low and High Shock groups, a larger fraction of the Aversive ensemble participated in bursts than the Remaining ensemble (*Ensemble: $F_{1,13} = 13.57$, $p = 0.0028$; Amplitude: $F_{1,13} = 0.000078$, $p = 0.99$; Ensemble x Amplitude: $F_{1,13} = 0.16$, $p = 0.69$*) (*Low Shock, N = 7 mice; High Shock, N = 8 mice*).

H) During Offline2 in both Low and High Shock groups, a larger fraction of the Neutral ∩ Aversive ensemble participated in bursts than the Remaining ensemble (*Ensemble: $F_{1,13} = 13.95$, $p = 0.0025$; Amplitude: $F_{1,13} = 0.014$, $p = 0.91$; Ensemble x Amplitude: $F_{1,13} = 0.31$, $p = 0.58$*) (*Low Shock, N = 7 mice; High Shock, N = 8 mice*).

I) During Offline2, Neutral and Remaining ensembles differentially participated in bursts in High and Low Shock groups (*Ensemble x Amplitude: $F_{1,13} = 5.186$, $p = 0.040$*). High Shock mice showed higher participation of the Neutral ensemble relative to Remaining ensemble ($t_7 = 4.88$, $p = 0.0036$), whereas Low Shock mice showed no different participation between the two ensembles ($t_6 = 1.33$, $p = 0.23$) (*Low Shock, N = 7 mice; High Shock, N = 8 mice*).

Strong aversive experience drives co-bursting of the Neutral ∩ Aversive ensemble with the Neutral ensemble

Since after High Shock, the Neutral and Aversive ensembles were both participating in burst events, we next asked whether the two ensembles co-participated within the same bursts, or whether they participated separately in different bursts. Co-bursting between the Neutral ensemble and Aversive ensemble could suggest a process through which the two ensembles can become integrated into a cell assembly likely to co-fire during memory recall thereafter. This process could occur through Hebbian plasticity[44] or through behavioral timescale synaptic plasticity, which has been proposed to drive the formation of place fields in hippocampal neurons[37]. Previous work has shown that hippocampal neurons become highly co-active during recall of an aversive memory but not during initial learning[45], that co-activity relationships among hippocampal neurons can distinguish between contexts that a mouse has experienced[46], and that ensembles that are highly co-active during an offline period following learning are more likely to be reactivated during memory recall than non-co-active neurons[15].

To ask whether the Neutral, Aversive, and Neutral ∩ Aversive ensembles were co-bursting after Aversive encoding, we measured the fraction of burst events that each ensemble participated in independently of each other (Figure 3A) and the fraction that the ensembles co-participated in (Figure 3D) during the offline period following the Aversive experience (Extended Figure 4I; see Methods). Previously, we had found that the Neutral ∩ Aversive cells (those active during both Neutral and Aversive encoding) were the most highly active during the offline period (Figure 2C). Highly active subpopulations of neurons have been proposed to form a 'hub-like' population of neurons that may orchestrate the activity of other neurons in a larger network[47,48]. Therefore, these highly active neurons could be organizing the activity of other neurons in the hippocampus to drive activity during this offline period. Thus, we hypothesized that co-participation between the highly active Neutral ∩ Aversive ensemble and the Neutral ensemble would be enhanced after a strong aversive experience.

We found that during burst events, the Neutral ∩ Aversive ensemble participated independently more frequently than the Neutral and Aversive ensembles did, but there was no difference between Low and High Shock mice (Figure 3B). Notably, during non-burst periods, independent ensemble bursting did not vary between any of the ensembles (Figure 3C). We next measured co-participation of

the ensembles in all combinations (Figure 3D). We found that in the Low Shock group, co-participation

between the three ensembles was less likely to occur than the other combinations; however,

surprisingly, in the High Shock group, co-participation between the three ensembles was no different

from the other combinations (Figure 3E). Additionally, in the High Shock group, the Neutral ∩ Aversive

ensemble co-participated with the Neutral ensemble more than it did with the Aversive ensemble,

whereas in the Low Shock group, the Neutral ∩ Aversive ensemble co-participated no differently with

the Neutral and Aversive ensembles (Figure 3E). Importantly, there were no differences in ensemble

co-bursting between Low and High Shock groups during non-burst periods (Figure 3F), suggesting that

the ensemble co-participation was confined to periods when the hippocampus was synchronously

active. These results suggested that after a strong aversive experience, the Neutral ∩ Aversive

ensemble was preferentially co-bursting with the Neutral ensemble. To confirm that this was the case,

we used cross-correlations as another measure of co-activity to measure how co-active the Neutral ∩

Aversive ensemble was with the Neutral and the Aversive ensembles. Indeed, only in the High Shock

group, the Neutral ∩ Aversive ensemble was preferentially correlated with the Neutral ensemble

compared with the Aversive ensemble during the offline period (Extended Figure 4K). Collectively,

these results suggest that a strong aversive experience increases the co-bursting of the Neutral ∩

Aversive ensemble with the Neutral ensemble, perhaps to link fear of the recent aversive experience

with the past neutral memory.

350 **Figure 3**

Figure 3. Strong aversive experience drives co-reactivation of the Neutral ensemble with the Neutral ∩ Aversive ensemble.

A) Representation of the quantification of independent participation during bursts versus non-bursting periods. Burst events were defined by the whole recorded population, as in Figure 2E (outlined by yellow rectangles). However, now the z-scored mean population activity of the Aversive, Neutral, and Neutral ∩ Aversive ensembles was computed to ask how frequently each ensemble participated in whole population bursts independently of one another. Independent participation meant one ensemble participated while the other two did not.

B) During burst periods, the Neutral ∩ Aversive ensemble participated independently in more bursts than the Aversive ensemble ($t_{14} = 7.95, p = 0.000002$) and more than the Neutral ensemble ($t_{14} = 5.59, p = 0.0001$) but there was no difference in participation across Low vs High Shock mice ($F_{1,13} = 1.43, p = 0.25$) and no interaction ($F_{2,26} = 2.49, p = 0.10$) (*Low Shock, N = 7 mice; High Shock, N = 8 mice*).

C) During non-burst periods, there was no difference in participation across ensembles ($F_{2,26} = 0.38, p = 0.69$) or between Low and High Shock mice ($F_{1,13} = 0.73, p = 0.41$), and no interaction ($F_{2,26} = 0.36, p = 0.70$) (*Low Shock, N = 7 mice; High Shock, N = 8 mice*).

D) Representation of the quantification of co-participation during bursts vs non-bursting periods. As in Figure 3B, the whole population was used to define bursts and the z-scored mean population activities were used to define participation of each ensemble. Co-participation was defined as a whole population burst (outlined by yellow rectangles) during which multiple ensembles participated simultaneously. There were four possible combinations (from left to right: N∩A x N, N∩A x A, N x A, N∩A x N x A) (N∩A = Neutral ∩ Aversive; N = Neutral; A = Aversive).

E) During burst periods, there was a significant interaction between Ensemble Combination and Low vs High Shock ($p = 0.01$), suggesting that the patterns of co-bursting varied in Low vs High Shock mice. Post-hoc tests revealed that in Low Shock mice, co-participation between all 3 ensembles was less likely to occur than the other combinations ($t_{18} = 4.73, p = 0.0003$), while in High Shock mice, co-participation between all 3 ensembles occurred no differently than the other combinations ($t_{21} = 0.358, p = 0.72$). Additionally, in the High Shock group, the N∩A ensemble preferentially co-participated with the Neutral ensemble compared to with the Aversive ensemble ($t_{21} = 2.373, p = 0.05$), whereas in the Low Shock group, the N∩A ensemble participated no differently with the Neutral and Aversive ensembles ($t_{18} = 1.196, p = 0.25$) (*Low Shock, N = 7 mice; High Shock, N = 8 mice*).

F) During non-burst periods, co-participation between all 3 ensembles was less likely than the other combinations ($t_{39} = 10.92, p = 1.98e\text{-}13$); however, there was no effect of Low vs High Shock ($F_{1,13} = 0.038, p = 0.847$) and no interaction ($F_{3,39} = 0.198, p = 0.897$) (*Low Shock, N = 7 mice; High Shock, N = 8 mice*).

Strong aversive experience drives co-reactivation of the Neutral & Aversive and Neutral

ensembles during Neutral context recall

Finally, we asked whether hippocampal ensemble reactivation could support the freezing observed in the Neutral context during recall after a high shock and not low shock (as shown in Figure 1F). To do this, we measured hippocampal ensemble activity while mice recalled the Neutral context after the offline period, compared with ensemble activity when they were placed in a Novel context as a control (Figure 4A). Since High Shock mice froze significantly more in the Neutral vs Novel contexts during recall (Figure 1F), we hypothesized that Neutral context recall would drive the aversive memory representation to be reactivated, whereas exposure to a Novel context would not provoke the reactivation of the aversive memory representation. Previously, we found that during the offline period, the Neutral ∩ Aversive ensemble specifically co-reactivated with the Neutral ensemble (Figure 3E, Extended Figure 4K), perhaps forming an integrated ensemble of neurons that is more likely to fire together in the future. If this were the case, when High Shock mice recalled the Neutral context and reactivated the Neutral ensemble, we predicted they might also reactivate the Neutral ∩ Aversive ensemble, perhaps through a process of pattern completion[49], thereby driving freezing in the Neutral context. Importantly, we expected this not to occur in Low Shock mice, where Neutral and Neutral ∩ Aversive ensemble co-reactivation was not observed, or in High Shock mice during Novel context exposure, since fear did not selectively spread to the Novel context (Figure 1F).

During recall of the Neutral context and exposure to a Novel context, we measured the fraction of cells active during that session which were previously active during encoding of the Neutral or Aversive contexts or active during both Neutral and Aversive encoding (Figure 4A; Extended Figure 4E,F). We previously observed that during the offline period, the Neutral ensemble co-fired with the Neutral ∩ Aversive after high shock but not after low shock (Figure 3D,E; Extended Figure 3K), potentially forming an integrated ensemble that is more likely to fire together later. Thus, we hypothesized that after high shock, during Neutral context recall, the Neutral ensemble (representing the Neutral context) would be reactivated, and this would, in turn, trigger reactivation of the Neutral ∩ Aversive ensemble. As expected, cells exclusively active during Neutral encoding and not Aversive encoding were more likely to be reactivated during Neutral recall than during Novel context exposure in

both Low Shock and High Shock groups, suggesting a stable and selective neural population

representing the neutral memory (Figure 4B). The cells exclusively active during Aversive encoding

were not selectively reactivated during Neutral or Novel contexts in either group (Figure 4C).

Interestingly, the cells active during both Neutral and Aversive encoding (Neutral ∩ Aversive ensemble)

were more reactivated during Neutral recall than during Novel context exposure in the High Shock but

not the Low Shock group (Figure 4D). This suggests that after ensemble co-reactivation during the

offline period following high shock, the Neutral ensemble and the Neutral ∩ Aversive ensembles were

more likely to reactivate together during Neutral recall.

The high shock aversive experience prompted an ensemble from days ago to be reactivated

offline. During subsequent Neutral recall, mice exhibited increased freezing despite never having been

shocked in that context. Therefore, the memory of the Neutral context had been modified to become

perceived as negative in High Shock mice. If this offline reactivation of the Neutral ensemble was

indeed modifying the neutral memory representation, we hypothesized that during Neutral recall, the

activity patterns observed would be different from the activity patterns observed during Neutral

encoding in the High Shock mice, compared to in Low Shock mice, and perhaps compared to the

change observed from Aversive encoding to Aversive recall. To test this, we computed a mean

population activity vector during Neutral encoding and correlated it with 30-second population vectors

across Neutral recall, to measure the similarity between activity patterns during encoding and recall

(see Methods)[50]. We repeated this for Aversive encoding and correlated it with activity patterns during

Aversive recall. Consistent with our hypothesis, Neutral encoding-to-recall correlations were lower in

High Shock mice compared to Low Shock mice. In High Shock mice, the Neutral encoding-to-recall

correlations were also lower than Aversive encoding-to-recall correlations, suggesting that the neutral

memory representation was significantly altered from encoding to recall in High Shock mice (Figure

4E). These results collectively suggest that a strong aversive experience drove the Neutral ∩ Aversive

and Neutral ensembles to co-fire during the offline period, altering the neutral memory representation.

And during Neutral recall, these ensembles were again co-reactivated, leading to the enhanced

freezing observed in the Neutral context.

 Figure 4

A

Encoding

Low Shock

| Day 1 | | Day 3 | |
| Neutral | Offline1 | Aversive (Low Shock) | Offline2 |

Recall

| Day 4 | Day 5 | Day 6 |
| Aversive (No Shock) | Novel | Neutral |

Counterbalanced

High Shock

| Day 1 | | Day 3 | |
| Neutral | Offline1 | Aversive (High Shock) | Offline2 |

| Day 4 | Day 5 | Day 6 |
| Aversive (No Shock) | Novel | Neutral |

Counterbalanced

B Ensembles Exclusive to Neutral Encoding

Low Shock   *High Shock*

% Recall Cells Active vs Recall Context (Neutral, Novel)

*** (Low Shock)   *** (High Shock)

C Ensembles Exclusive to Aversive Encoding

Low Shock   *High Shock*

% Recall Cells Active vs Recall Context (Neutral, Novel)

D Ensembles Present in Both

Low Shock   *High Shock*

% Recall Cells Active vs Recall Context (Neutral, Novel)

**** (High Shock)

Reactivation of Encoding Population Vectors During Recall

E

Low Shock   *High Shock*

Kendal Tau Correlation (Encoding to Recall) vs (Aversive, Neutral)

* (between Low Shock and High Shock)
** (High Shock)

Figure 4. Strong aversive experience drives Neutral ∩ Aversive ensemble reactivation during Neutral context recall.

A) Behavioral schematic of calcium imaging experiment, as in Figure 1D. Here, we focused on hippocampal activity during the Aversive, Neutral, and Novel recall sessions.

B) Cells active only during the Neutral experience and not the Aversive experience were more likely to be reactivated when mice were placed back in the Neutral context, compared to when they were placed in a Novel context ($F_{1,12} = 24.44$, $p = 0.0003$). There was no effect of shock amplitude ($F_{1,12} = 3.08$, $p = 0.10$) (*Low Shock, N = 6 mice; High Shock, N = 8 mice*).

C) Cells active during the Aversive experience and not the Neutral experience were no differently reactivated in Neutral vs Novel contexts. (*Amplitude: $F_{1,12} = 0.029$, $p = 0.869$; Context: $F_{1,12} = 1.39$, $p = 0.261$; Amplitude x Context: $F_{1,12} = 0.14$, $p = 0.71$*) (*Low Shock, N = 6 mice; High Shock, N = 8 mice*).

D) Cells active during both the initial Neutral and Aversive experiences were subsequently more likely to be reactivated in the Neutral context compared to Novel context in High Shock mice ($t_7 = 8.53$, $p = 0.00012$), but not Low Shock mice (*$t_5 = 0.55$, $p = 0.61$; Context x Amplitude: $F_{1,12} = 10.33$, $p = 0.007$*) (*Low Shock, N = 6 mice; High Shock, N = 8 mice*).

E) In High Shock mice, population activity patterns in the Neutral context changed significantly from Neutral encoding to Neutral recall (*Amplitude: $F_{1,12} = 5.65$; SessionPair: $F_{1,12} = 10.42$; Amplitude x SessionPair: $F_{1,12} = 6.22$*). During Neutral recall in High Shock mice, population activity vectors were less correlated with the average Neutral encoding population vector than Aversive recall activity was with the average Aversive encoding population vector ($t_7 = 4.10$, $p = 0.009$). Neutral encoding-to-recall correlations were also lower in High vs Low Shock mice ($t_{6.92} = 2.98$, $p = 0.042$). Aversive encoding-to-recall correlations were no different in High vs Low Shock mice ($t_{6.11} = 1.13$, $p = 0.30$). In Low Shock mice, Neutral and Aversive encoding-to-recall correlations were no different ($t_5 = 0.23$, $p = 0.83$) (*Low Shock, N = 6 mice; High Shock, N = 8 mice*).

Discussion

How animals actively update memories as they encounter new information remains a fundamental question in neuroscience[21]. Past work has shown that individual experiences are encoded by subpopulations of neurons across the brain that are highly active during learning[51,52]. These neuronal ensembles undergo synaptic modifications after learning to support memory storage[53-56]. After learning, activity of these ensembles is necessary[38,57] and sufficient[2] to drive memory recall, and their reactivation during memory recall is correlated with the strength of memory recall[1]. How memories encoded across time are integrated remains a critical and unanswered question in neuroscience. The memory allocation hypothesis suggests that neurons with high intrinsic excitability at the time of learning are likely to be allocated to a memory trace[5,58]. Prior studies suggest that two memories encoded within a day are likely to be linked because they share an overlapping population of highly excitable neurons during the initial learning. This shared neural ensemble links the two temporally related memories, such that the recall of one memory is more likely to trigger the recall of another memory that was encoded close in time[4,27,28,59]. Here we demonstrate that memories can be dynamically updated even days after they have been encoded and consolidated, and that this process is driven by ensemble co-reactivation during a post-learning period.

Whether linking memories across days is an adaptive or maladaptive process may depend on the environmental conditions. Under everyday circumstances, memories that are encoded far apart in time and which share no features in common may typically not need to be linked, and memories must also be segregated to allow for proper recall of distinct memories. Notably, the hippocampus has been shown to successfully discriminate between distinct memories[60,61]. However, after a potentially life-threatening experience, especially one where the source of the aversive outcome is ambiguous (as in the aversive experience employed here), it could benefit an animal to link fear from that aversive experience to prior events, particularly if the event is rare and novel as seen in conditioned taste aversion[22]. Our results suggest that a highly aversive experience is more likely to drive memory-linking than a mild aversive experience (Figure 1D-G), consistent with this intuition. Moreover, our results suggest that fear is more likely to be linked retrospectively to past events rather than prospectively to future events (Figure 1A-C). This is consistent with the notion that cues that occurred before an

outcome can predict that outcome. On a shorter timescale, it has been well established that when a

neutral cue directly precedes a foot shock by seconds, this drives associative learning between the cue

and the foot shock to drive cue-elicited freezing[62,63]. Interestingly, however, if the cue instead occurs

directly *after* the foot shock, the animal no longer freezes in response to cue presentation thereafter,

presumably because the cue predicts the ensuing absence of the aversive event[64]. Though the

difference in timescale suggests that different mechanisms are likely at play in these two scenarios, our

results are consistent with the idea that cues occurring prior to an outcome can be interpreted as

predictive cues to the animal. A recent review has also suggested that animals use "retrospective

cognitive maps" to infer the states that precede an outcome, to draw causal associations between

those stimuli[65]. Our results suggest that offline periods are responsible for driving this retrospective

inference (Figure 5).

Offline periods offer an opportunity for the brain to draw inferences about relationships that were

not necessarily formed at the time of learning. In humans, it has been shown that an emotional

experience can retrospectively increase memory for previously experienced neutral objects, only after a

period of consolidation[31]. A separate study demonstrated that this retrospective memory enhancement

coincided with increased functional hippocampal-cortical coupling and fMRI BOLD activity in the ventral

tegmental area[35]. Moreover, a recent study in mice showed that two contexts with strongly shared

geometrical features can be integrated immediately after learning (i.e., 15min after learning), whereas

two contexts with subtly shared geometrical features require an offline period after learning (i.e., 1 day)

to drive their integration. During this offline period, cortical ensemble co-reactivation drives this memory

integration[66]. Our study demonstrates that a highly aversive experience can alter the likelihood of

retrospective memory-linking, that this is dependent upon post-learning hippocampal activity, and is

accompanied by co-reactivation of the ensembles for the two memories.

Past studies have shown that ensemble reactivation occurs during both sleep (NREM and REM

sleep) and wake states. Reactivation during different states have been proposed to support different

memory processes. For instance, classical studies demonstrated that following a salient experience,

the patterns of neuronal activity that were present during learning are replayed in the same sequential

order offline, and this replay has been observed during both NREM[9] and REM[8] sleep. The replay

observed during sleep was proposed to support memory consolidation, and indeed, disruption of sharp-wave ripples (during which most of these replay events occur) disrupts the storage of memories such that memory recall is disrupted thereafter[16,19]. Remarkably, one study found that prolonging sharp wave ripple durations benefited memory while cutting them short impaired memory[67]. In addition to sleep, it has also been observed that hippocampal replay occurs while animals are awake and engaged in an experimental task, and it can occur in a forward or reverse direction[10,12,68,69]. This has led to the idea that different forms of replay may serve different functions, from memory consolidation to planning and decision-making[18,39], though this remains a debate[70]. More generally, sleep has been shown to strongly benefit learning in both rodents[17,33,34,71] and in humans[32,72-74], and neurophysiological events during sleep, such as sharp wave ripples and sleep spindles, have been suggested to support memory consolidation[16,19,71]. Whether ensemble *co-reactivation* supporting memory integration is a sleep state specific phenomenon and whether distinct sleep/wake states differentially support memory consolidation versus integration has yet to be answered. Our results suggest that the transient population bursts during which we observed ensemble co-reactivation occurs during quiet wake, since locomotion decreased about one-second prior to each burst and resumed one-second following it (Extended Figure 5I,J). However, this study did not explicitly measure ensemble reactivation during distinct sleep states – thus, it remains unclear whether ensemble co-reactivation may occur in a sleep state specific manner to drive memory-linking. A recent study demonstrated that in a neural network model with autonomous offline reactivation, interleaved periods of NREM and REM sleep were critical for the integration of memories[25]. However, a previous study in rats suggested that offline reactivation and modification of a past neutral memory occurred during wake periods[24]. Thus, resolving whether and how different sleep states support memory integration processes will be an important future direction.

Finally, these results have implications for the interpretation of the clinical manifestation of memory-related conditions such as post-traumatic stress disorder (PTSD). PTSD transpires from one or multiple traumatic events and is hallmarked by uncontrollable fear in non-life-threatening contexts[75]. A common form of behavioral treatment for PTSD is exposure therapy, whereby the patient is carefully re-exposed to the trauma-associated conditioned stimuli, seeking to detach the association between

those stimuli and fear. In many cases, exposure therapy successfully decreases fear, but patients are

often prone to relapse thereafter[76]. Our results suggest that highly salient aversive experiences can

drive fear to be associated with seemingly unrelated stimuli that were not present at the time of the

aversive experience, and that this scales with the perceived aversiveness of the experience (Figure

1G). This predicts that while exposure therapy may successfully inhibit fear to the trauma stimuli, the

fear from the trauma may have spread to other stimuli that were not directly targeted by the therapy.

Thus, it may be useful to consider stimuli that were experienced across time that may have insidiously

become linked with the trauma. Ultimately, our results point to the offline period after an aversive event

as a potential intervention timepoint to unlink memories separated across days.

Figure 5

Figure 5. Offline ensemble reactivation drives retrospective memory-linking across days. After single
experiences, the cells active during learning are reactivated to support their consolidation. After a strong aversive
experience, memories are linked retrospectively across days by the co-reactivation of the ensembles representing
both the recent and the past neutral memory ensembles. During recall of the neutral memory, many of the cells
that were active during both the neutral and aversive experiences are reactivated to drive fear in the neutral
context.

References

Reijmers, L. G., Perkins, B. L., Matsuo, N. & Mayford, M. Localization of a stable neural correlate of associative memory. *Science* **317**, 1230-1233 (2007). https://doi.org:10.1126/science.1143839

Liu, X. *et al.* Optogenetic stimulation of a hippocampal engram activates fear memory recall. *Nature* **484**, 381-385 (2012). https://doi.org:10.1038/nature11028

Han, J. H. *et al.* Neuronal competition and selection during memory formation. *Science* **316**, 457-460 (2007). https://doi.org:10.1126/science.1139438

Rogerson, T. *et al.* Synaptic tagging during memory allocation. *Nat Rev Neurosci* **15**, 157-169 (2014). https://doi.org:10.1038/nrn3667

Josselyn, S. A. & Frankland, P. W. Memory Allocation: Mechanisms and Function. *Annu Rev Neurosci* **41**, 389-413 (2018). https://doi.org:10.1146/annurev-neuro-080317-061956

Pavlides, C. & Winson, J. Influences of hippocampal place cell firing in the awake state on the activity of these cells during subsequent sleep episodes. *J Neurosci* **9**, 2907-2918 (1989). https://doi.org:10.1523/JNEUROSCI.09-08-02907.1989

Wilson, M. A. & McNaughton, B. L. Reactivation of hippocampal ensemble memories during sleep. *Science* **265**, 676-679 (1994). https://doi.org:10.1126/science.8036517

Louie, K. & Wilson, M. A. Temporally structured replay of awake hippocampal ensemble activity during rapid eye movement sleep. *Neuron* **29**, 145-156 (2001). https://doi.org:10.1016/s0896-6273(01)00186-6

Lee, A. K. & Wilson, M. A. Memory of sequential experience in the hippocampus during slow wave sleep. *Neuron* **36**, 1183-1194 (2002). https://doi.org:10.1016/s0896-6273(02)01096-6

Foster, D. J. & Wilson, M. A. Reverse replay of behavioural sequences in hippocampal place cells during the awake state. *Nature* **440**, 680-683 (2006). https://doi.org:10.1038/nature04587

Ji, D. & Wilson, M. A. Coordinated memory replay in the visual cortex and hippocampus during sleep. *Nat Neurosci* **10**, 100-107 (2007). https://doi.org:10.1038/nn1825

Diba, K. & Buzsaki, G. Forward and reverse hippocampal place-cell sequences during ripples. *Nat Neurosci* **10**, 1241-1242 (2007). https://doi.org:10.1038/nn1961

13   Carr, M. F., Jadhav, S. P. & Frank, L. M. Hippocampal replay in the awake state: a potential

substrate for memory consolidation and retrieval. *Nat Neurosci* **14**, 147-153 (2011).

https://doi.org:10.1038/nn.2732

14   Girardeau, G., Inema, I. & Buzsaki, G. Reactivations of emotional memory in the hippocampus-

amygdala system during sleep. *Nat Neurosci* **20**, 1634-1642 (2017).

https://doi.org:10.1038/nn.4637

15   Ghandour, K. *et al.* Orchestrated ensemble activities constitute a hippocampal memory engram.

Nat Commun **10**, 2637 (2019). https://doi.org:10.1038/s41467-019-10683-2

16   Gridchyn, I., Schoenenberger, P., O'Neill, J. & Csicsvari, J. Assembly-Specific Disruption of

Hippocampal Replay Leads to Selective Memory Deficit. *Neuron* **106**, 291-300 e296 (2020).

https://doi.org:10.1016/j.neuron.2020.01.021

17   Clawson, B. C. *et al.* Causal role for sleep-dependent reactivation of learning-activated sensory

ensembles for fear memory consolidation. *Nat Commun* **12**, 1200 (2021).

https://doi.org:10.1038/s41467-021-21471-2

18   Buzsaki, G. Hippocampal sharp wave-ripple: A cognitive biomarker for episodic memory and

planning. *Hippocampus* **25**, 1073-1188 (2015). https://doi.org:10.1002/hipo.22488

19   van de Ven, G. M., Trouche, S., McNamara, C. G., Allen, K. & Dupret, D. Hippocampal Offline

Reactivation Consolidates Recently Formed Cell Assembly Patterns during Sharp Wave-

Ripples. *Neuron* **92**, 968-974 (2016). https://doi.org:10.1016/j.neuron.2016.10.020

20   Colgin, L. L. Rhythms of the hippocampal network. *Nat Rev Neurosci* **17**, 239-249 (2016).

https://doi.org:10.1038/nrn.2016.21

21   Mau, W., Hasselmo, M. E. & Cai, D. J. The brain in motion: How ensemble fluidity drives

memory-updating and flexibility. *Elife* **9** (2020). https://doi.org:10.7554/eLife.63550

22   Chambers, K. C. Conditioned taste aversions. *World J Otorhinolaryngol Head Neck Surg* **4**, 92-

100 (2018). https://doi.org:10.1016/j.wjorl.2018.02.003

23   Cai, D. J., Mednick, S. A., Harrison, E. M., Kanady, J. C. & Mednick, S. C. REM, not incubation,

improves creativity by priming associative networks. *Proc Natl Acad Sci U S A* **106**, 10130-

10134 (2009). https://doi.org:10.1073/pnas.0900271106

24   Jezek, K. *et al.* Stress-induced out-of-context activation of memory. *PLoS Biol* **8**, e1000570

630   (2010). https://doi.org:10.1371/journal.pbio.1000570

631   25   Singh, D., Norman, K. A. & Schapiro, A. C. A model of autonomous interactions between

632   hippocampus and neocortex driving sleep-dependent memory consolidation. *Proc Natl Acad Sci*

633   *U S A* **119**, e2123432119 (2022). https://doi.org:10.1073/pnas.2123432119

634   26   Kurth-Nelson, Z. *et al.* Replay and compositional computation. *Neuron* (2023).

635   https://doi.org:10.1016/j.neuron.2022.12.028

636   27   Cai, D. J. *et al.* A shared neural ensemble links distinct contextual memories encoded close in

637   time. *Nature* **534**, 115-118 (2016). https://doi.org:10.1038/nature17955

638   28   Rashid, A. J. *et al.* Competition between engrams influences fear memory formation and recall.

639   *Science* **353**, 383-387 (2016). https://doi.org:10.1126/science.aaf0594

640   29   Phelps, E. A. & Sharot, T. How (and Why) Emotion Enhances the Subjective Sense of

641   Recollection. *Curr Dir Psychol Sci* **17**, 147-152 (2008). https://doi.org:10.1111/j.1467-

642   8721.2008.00565.x

643   30   LaBar, K. S. & Cabeza, R. Cognitive neuroscience of emotional memory. *Nat Rev Neurosci* **7**,

644   54-64 (2006). https://doi.org:10.1038/nrn1825

645   31   Dunsmoor, J. E., Murty, V. P., Davachi, L. & Phelps, E. A. Emotional learning selectively and

646   retroactively strengthens memories for related events. *Nature* **520**, 345-348 (2015).

647   https://doi.org:10.1038/nature14106

648   32   Diekelmann, S. & Born, J. The memory function of sleep. *Nat Rev Neurosci* **11**, 114-126 (2010).

649   https://doi.org:10.1038/nrn2762

650   33   Havekes, R. *et al.* Sleep deprivation causes memory deficits by negatively impacting neuronal

651   connectivity in hippocampal area CA1. *Elife* **5** (2016). https://doi.org:10.7554/eLife.13424

652   34   Cai, D. J., Shuman, T., Gorman, M. R., Sage, J. R. & Anagnostaras, S. G. Sleep selectively

653   enhances hippocampus-dependent memory in mice. *Behav Neurosci* **123**, 713-719 (2009).

654   https://doi.org:10.1037/a0016415

655   35   Clewett, D., Dunsmoor, J., Bachman, S. L., Phelps, E. A. & Davachi, L. Survival of the salient:

656   Aversive learning rescues otherwise forgettable memories via neural reactivation and post-

657    encoding hippocampal connectivity. *Neurobiol Learn Mem* **187**, 107572 (2022).

658    https://doi.org:10.1016/j.nlm.2021.107572

659 36  Lewis, P. A., Knoblich, G. & Poe, G. How Memory Replay in Sleep Boosts Creative Problem-

660    Solving. *Trends Cogn Sci* **22**, 491-503 (2018). https://doi.org:10.1016/j.tics.2018.03.009

661 37  Bittner, K. C., Milstein, A. D., Grienberger, C., Romani, S. & Magee, J. C. Behavioral time scale

662    synaptic plasticity underlies CA1 place fields. *Science* **357**, 1033-1036 (2017).

663    https://doi.org:10.1126/science.aan3846

664 38  Denny, C. A. *et al.* Hippocampal memory traces are differentially modulated by experience,

665    time, and adult neurogenesis. *Neuron* **83**, 189-201 (2014).

666    https://doi.org:10.1016/j.neuron.2014.05.018

667 39  Joo, H. R. & Frank, L. M. The hippocampal sharp wave-ripple in memory retrieval for immediate

668    use and consolidation. *Nat Rev Neurosci* **19**, 744-757 (2018). https://doi.org:10.1038/s41583-

669    018-0077-1

670 40  Magnus, C. J. *et al.* Chemical and genetic engineering of selective ion channel-ligand

671    interactions. *Science* **333**, 1292-1296 (2011). https://doi.org:10.1126/science.1206606

672 41  Ghandour, K. & Inokuchi, K. Memory reactivations during sleep. *Neurosci Res* (2022).

673    https://doi.org:10.1016/j.neures.2022.12.018

674 42  Malvache, A., Reichinnek, S., Villette, V., Haimerl, C. & Cossart, R. Awake hippocampal

675    reactivations project onto orthogonal neuronal assemblies. *Science* **353**, 1280-1283 (2016).

676    https://doi.org:10.1126/science.aaf3319

677 43  Villette, V., Malvache, A., Tressard, T., Dupuy, N. & Cossart, R. Internally Recurring

678    Hippocampal Sequences as a Population Template of Spatiotemporal Information. *Neuron* **88**,

679    357-366 (2015). https://doi.org:10.1016/j.neuron.2015.09.052

680 44  Buzsaki, G. Neural syntax: cell assemblies, synapsembles, and readers. *Neuron* **68**, 362-385

681    (2010). https://doi.org:10.1016/j.neuron.2010.09.023

682 45  Rajasethupathy, P. *et al.* Projections from neocortex mediate top-down control of memory

683    retrieval. *Nature* **526**, 653-659 (2015). https://doi.org:10.1038/nature15389

46   Gava, G. P. *et al.* Integrating new memories into the hippocampal network activity space. *Nat*
Neurosci **24**, 326-330 (2021). https://doi.org:10.1038/s41593-021-00804-w

47   Buzsaki, G. & Mizuseki, K. The log-dynamic brain: how skewed distributions affect network
operations. *Nat Rev Neurosci* **15**, 264-278 (2014). https://doi.org:10.1038/nrn3687

48   Luccioli, S., Ben-Jacob, E., Barzilai, A., Bonifazi, P. & Torcini, A. Clique of functional hubs
orchestrates population bursts in developmentally regulated neural networks. *PLoS Comput Biol*
10, e1003823 (2014). https://doi.org:10.1371/journal.pcbi.1003823

49   Rolls, E. T. The mechanisms for pattern completion and pattern separation in the hippocampus.
Front Syst Neurosci **7**, 74 (2013). https://doi.org:10.3389/fnsys.2013.00074

50   Zaki, Y. *et al.* Hippocampus and amygdala fear memory engrams re-emerge after contextual
fear relapse. *Neuropsychopharmacology* **47**, 1992-2001 (2022). https://doi.org:10.1038/s41386-
022-01407-0

51   Josselyn, S. A. & Tonegawa, S. Memory engrams: Recalling the past and imagining the future.
Science **367** (2020). https://doi.org:10.1126/science.aaw4325

52   Zhou, Y. *et al.* CREB regulates excitability and the allocation of memory to subsets of neurons
in the amygdala. *Nat Neurosci* **12**, 1438-1443 (2009). https://doi.org:10.1038/nn.2405

53   Bocchio, M., Nabavi, S. & Capogna, M. Synaptic Plasticity, Engrams, and Network Oscillations
in Amygdala Circuits for Storage and Retrieval of Emotional Memories. *Neuron* **94**, 731-743
(2017). https://doi.org:10.1016/j.neuron.2017.03.022

54   Ryan, T. J., Roy, D. S., Pignatelli, M., Arons, A. & Tonegawa, S. Memory. Engram cells retain
memory under retrograde amnesia. *Science* **348**, 1007-1013 (2015).
https://doi.org:10.1126/science.aaa5542

55   Choi, J. H. *et al.* Interregional synaptic maps among engram cells underlie memory formation.
Science **360**, 430-435 (2018). https://doi.org:10.1126/science.aas9204

56   Abdou, K. *et al.* Synapse-specific representation of the identity of overlapping memory engrams.
Science **360**, 1227-1231 (2018). https://doi.org:10.1126/science.aat3810

57   Han, J. H. *et al.* Selective erasure of a fear memory. *Science* **323**, 1492-1496 (2009).
https://doi.org:10.1126/science.1164139

58   Silva, A. J., Zhou, Y., Rogerson, T., Shobe, J. & Balaji, J. Molecular and cellular approaches to

memory allocation in neural circuits. *Science* **326**, 391-395 (2009).

https://doi.org:10.1126/science.1174519

59   Yokose, J. *et al.* Overlapping memory trace indispensable for linking, but not recalling, individual

memories. *Science* **355**, 398-403 (2017). https://doi.org:10.1126/science.aal2690

60   van Dijk, M. T. & Fenton, A. A. On How the Dentate Gyrus Contributes to Memory

Discrimination. *Neuron* **98**, 832-845 e835 (2018). https://doi.org:10.1016/j.neuron.2018.04.018

61   Lohnas, L. J. *et al.* Time-resolved neural reinstatement and pattern separation during memory

decisions in human hippocampus. *Proc Natl Acad Sci U S A* **115**, E7418-E7427 (2018).

https://doi.org:10.1073/pnas.1717088115

62   LeDoux, J. E. Emotion circuits in the brain. *Annu Rev Neurosci* **23**, 155-184 (2000).

https://doi.org:10.1146/annurev.neuro.23.1.155

63   Maren, S. Neurobiology of Pavlovian fear conditioning. *Annu Rev Neurosci* **24**, 897-931 (2001).

https://doi.org:10.1146/annurev.neuro.24.1.897

64   Moscovitch, A. & LoLordo, V. M. Role of safety in the Pavlovian backward fear conditioning

procedure. *J Comp Physiol Psychol* **66**, 673-678 (1968). https://doi.org:10.1037/h0026548

65   VM, K. N. & Stuber, G. D. The learning of prospective and retrospective cognitive maps within

neural circuits. *Neuron* **109**, 3552-3575 (2021). https://doi.org:10.1016/j.neuron.2021.09.034

66   Aly, M. H., Abdou, K., Okubo-Suzuki, R., Nomoto, M. & Inokuchi, K. Selective engram

coreactivation in idling brain inspires implicit learning. *Proc Natl Acad Sci U S A* **119**,

e2201578119 (2022). https://doi.org:10.1073/pnas.2201578119

67   Fernandez-Ruiz, A. *et al.* Long-duration hippocampal sharp wave ripples improve memory.

Science **364**, 1082-1086 (2019). https://doi.org:10.1126/science.aax0758

68   Karlsson, M. P. & Frank, L. M. Awake replay of remote experiences in the hippocampus. *Nat*

Neurosci **12**, 913-918 (2009). https://doi.org:10.1038/nn.2344

69   Jackson, J. C., Johnson, A. & Redish, A. D. Hippocampal sharp waves and reactivation during

awake states depend on repeated sequential experience. *J Neurosci* **26**, 12415-12426 (2006).

https://doi.org:10.1523/JNEUROSCI.4118-06.2006

70 Gillespie, A. K. *et al.* Hippocampal replay reflects specific past experiences rather than a plan

for subsequent choice. *Neuron* **109**, 3149-3163 e3146 (2021).

https://doi.org:10.1016/j.neuron.2021.07.029

71 Varga, A. W., Kang, M., Ramesh, P. V. & Klann, E. Effects of acute sleep deprivation on motor

and reversal learning in mice. *Neurobiol Learn Mem* **114**, 217-222 (2014).

https://doi.org:10.1016/j.nlm.2014.07.001

72 Denis, D. *et al.* The roles of item exposure and visualization success in the consolidation of

memories across wake and sleep. *Learn Mem* **27**, 451-456 (2020).

https://doi.org:10.1101/lm.051383.120

73 Wagner, U., Gais, S., Haider, H., Verleger, R. & Born, J. Sleep inspires insight. *Nature* **427**,

352-355 (2004). https://doi.org:10.1038/nature02223

74 Klinzing, J. G., Niethard, N. & Born, J. Mechanisms of systems memory consolidation during

sleep. *Nat Neurosci* **22**, 1598-1610 (2019). https://doi.org:10.1038/s41593-019-0467-3

75 Ressler, K. J. *et al.* Post-traumatic stress disorder: clinical and translational neuroscience from

cells to circuits. *Nat Rev Neurol* **18**, 273-288 (2022). https://doi.org:10.1038/s41582-022-00635-

8

76 Boschen, M. J., Neumann, D. L. & Waters, A. M. Relapse of successfully treated anxiety and

fear: theoretical issues and recommendations for clinical practice. *Aust N Z J Psychiatry* **43**, 89-

100 (2009). https://doi.org:10.1080/00048670802607154

Methods

Subjects

Adult C57BL/6J mice from Jackson Laboratories were used in all experiments. Mice arrived group-housed in cages of 4 mice/cage and were singly housed for the experiment. For behavioral experiments where mice did not undergo surgery, mice were ordered to arrive at 12 weeks of age and underwent behavioral testing 1-2 weeks from then. For experiments where mice underwent surgery, mice were ordered to arrive at 8-9 weeks of age and underwent behavioral testing about 4-6 weeks after the arrival date.

Viral constructs

For calcium imaging experiments, AAV1-Syn-GCaMP6f-WPRE-SV40 (titer: 2.8 x 10^13 GC/mL) was purchased from AddGene and was diluted 1:4 in sterile 1x PBS. Mice had 300nL of the diluted virus injected into the right hemisphere of dorsal CA1. For PSAM experiments, AAV5-Syn-PSAM4-GlyR-IRES-eGFP (2.4 x 10^13 GC/mL) was purchased from AddGene. Mice had the virus injected at stock titer bilaterally into dorsal and ventral hippocampus, 300nL per injection site.

Surgery

Mice were anesthetized with 1 to 2% isoflurane for surgical procedures and placed into a stereotaxic frame (David Kopf Instruments, Tujunga, CA). Eye ointment was applied to prevent desiccation, and mice were kept on a heated pad to prevent hypothermia. Surgery was performed with aseptic technique. After surgery, carprofen (5 mg/kg) was administered every day for the following three days, and ampicillin (20 mg/kg) was administered every day for the following 7 days. For calcium imaging experiments, dexamethasone (0.2 mg/kg) was also administered for the following 7 days.

For PSAM experiments, AAV5-Syn-PSAM4-GlyR-IRES-eGFP was injected at stock concentration. Mice had 300nL of the virus injected bilaterally into dorsal hippocampus (AP: -2mm, ML: +/-1.5mm, DV: -1.5mm) and 300nL injected bilaterally into ventral hippocampus (AP: -3mm, ML: +/-3.2mm, DV: -4mm), for a total of 4 injections and 1.2uL injected per mouse, using a glass pipette and Nanoject

injector. The pipette was slowly lowered to the injection site, the virus was injected at 2nL/sec, and then

the pipette remained for 5min before being removed to allow for virus diffusion. Mice had their incision

sutured following surgery and had betadine applied to the site to prevent infection.

For calcium imaging experiments, mice underwent two serial surgeries spaced one month apart, as

described before[1]. During the first surgery, a 1mm diameter craniotomy was made above the dorsal

hippocampus on the right hemisphere (centered at AP -2mm, ML +1.5mm from bregma). An anchor

screw was screwed into the skull on the contralateral hemisphere at approximately AP -1mm and ML -

2.5mm from bregma. 300nL of AAV1-Syn-GCaMP6f was injected into dorsal CA1 of the hippocampus

on the right hemisphere (AP -2mm, ML +1.5mm, DV -1.2mm). Virus was injected as described in

PSAM experiments above. After the pipette was removed, the mouse remained on the stereotaxic

frame for 20min to allow for complete diffusion of the virus. After the 20min of diffusion, the cortex

below the craniotomy was aspirated with a 25-gauge blunt syringe needle attached to a vacuum pump,

while constantly being irrigated with cortex buffer. When the striations of the corpus callosum were

visible, the 25-gauge needle was replaced with a 27-gauge needle for finer tuned aspiration. Once most

of corpus callosum was removed, bleeding was controlled using surgical foam (Surgifoam), and then a

1mm diameter x 4mm length GRIN lens (GRINTECH) was slowly lowered into the craniotomy. The lens

was fixed with cyanoacrylate, and then dental acrylic was applied to cement the implant in place and

cover the rest of the exposed skull. The top of the exposed lens was covered with Kwik-Sil (World

Precision Instruments) to protect it and the Kwik-Sil was covered with dental cement. Four weeks later,

mice were returned to attach the baseplate, visually guided by a Miniscope. The overlying dental

cement was drilled off and the Kwik-Sil was removed to reveal the top of the lens. The Miniscope with

an attached baseplate was lowered near the implanted lens and the field of view was monitored in real-

time on a computer. The Miniscope was rotated until a well-exposed field of view was observed, at

which point the baseplate was fixed to the implant with cyanoacrylate and dental cement. The mouse

did not receive post-operative drugs after this surgery since it was not invasive.

Behavioral procedures

Prior to all experiments, mice were handled for one minute each day for at least one week. On at least

four of those days, mice were transported to the testing room and handled there. On the rest of the

days, the mice were handled in the vivarium. In calcium imaging experiments, mice were handled and

habituated for 2 weeks instead of 1, during which they were habituated to having the Miniscope

attached and detached from its head. To become accustomed to the weight of the Miniscope, they

were placed in their homecage with the Miniscope attached for 5min per day for at least 5 days.

In Retrospective and Prospective memory-linking behavioral experiments, mice were exposed to the

Neutral context for 10 minutes to explore. During Aversive encoding, mice were placed in a novel

context and allowed to explore for 2 minutes. Then, mice received a 2-second foot shock of either

0.25mA (low shock) or 1.5mA (high shock). One minute after the first shock, they received a second

shock of the sample duration and amplitude, with a third shock following 1 minute after the second. 30

seconds after the third shock, the mice were removed and placed back in their homecage. On the

following three days, mice were tested in the previously experienced Aversive and Neutral contexts, as

well as a completely Novel context that they had not been exposed to prior. The features of the Neutral

and Novel contexts were counter-balanced and were made up of different olfactory, auditory, lighting,

and tactile cues. The Aversive context was always the same with distinct cues from the Neutral and

Novel contexts. In the PSAM experiment, mice were tested in either the Aversive, Neutral, or Novel

context. In the Prospective versus Retrospective memory-linking experiment, mice were tested in the

Aversive context first, and then half the mice were tested in the Neutral context and the other half in the

Novel context. In the Low vs High Shock experiments, mice were tested in the Aversive context first,

followed by testing in the Neutral and Novel context counter-balanced; half the mice received Neutral

recall and then Novel context exposure the next day, and the other half Novel context exposure and

then Neutral recall. All testing was done in Med Associates chambers. Behavioral data were analyzed

using the Med Associates software for measuring freezing. In experiments where mice were tethered

with a Miniscope, behavioral data were analyzed using our previously published open-source

behavioral tracking pipeline, ezTrack[2]. In the Prospective versus Retrospective memory-linking

timecourse experiments, the Aversive learning experience was distinct: mice explored for 2min, then

administered one 0.75mA, 2-second long foot shock and removed from the context 30sec following this

shock.

<u>Drug injections</u>

uPSEM-817 tartrate was made in a solution of 0.1mg/mL in saline and injected intraperitoneally at a

dose of 1mg/kg (10mL/kg injection volume). Saline was used as a vehicle. The first injection was done

as soon as the mice were brought back to the vivarium after Aversive encoding (~3min after the end of

Aversive encoding). The next 3 injections were done every 3 hours to cover a 12-hour timespan of

inhibition.

<u>Calcium imaging Miniscope recordings</u>

Open-source V4 Miniscopes (https://github.com/Aharoni-Lab/Miniscope-v4) were connected to a

coaxial cable which connected to a Miniscope data acquisition board (DAQ) 3.3. The DAQ connected to

a computer via a USB3.0. Data was collected via the Miniscope QT Software

(https://github.com/Aharoni-Lab/Miniscope-DAQ-QT-Software) at 30 frames per second. Miniscopes

and DAQ boards were all purchased from Open Ephys.

When performing calcium imaging with concurrent behavior in the Med Associates boxes, mice were

brought into the testing room from the vivarium, taken out of their homecage, and had the Miniscope

attached. They were placed back into their homecage for 1min. Then, they were removed from their

homecage and placed in the testing chamber. To record calcium and behavior, the Med Associates

software sent a continuous TTL pulse to record from the Miniscope while the behavior was concurrently

tracked via Med Associates cameras. After the session was complete, the mice were immediately

returned to their homecage, then the Miniscope was removed, and the mouse was returned to the

vivarium. One mouse was brought to the testing room at a time so that mice did not idly wait in the

testing room with partial recall cues from the room present.

Offline calcium imaging recordings were done in the mouse's homecage for the 1 hour following Neutral

encoding and following Aversive encoding. During these recordings, mice were placed back in their

homecage and the homecage was placed in a large rectangular and opaque storage bin to occlude

distal cues, with a webcam (Logitech C920e) overlying the homecage to track behavior during the

recording. Using the Miniscope QT Software with two devices connected (Miniscope and webcam),

calcium imaging and behavior were concurrently tracked. After the offline recording was complete, mice

were removed from their homecage, the Miniscope was removed, they were returned to their

homecage and returned to the vivarium immediately thereafter. The same procedure was undergone

for the experiment in Extended Figure 3.

<u>Miniscope data processing and data alignment</u>

To extract calcium transients from the calcium imaging data, we employed our previously published

open-source calcium imaging data processing pipeline, Minian[3]. Briefly, videos were pre-processed for

background fluorescence and sensor noise, and motion corrected. Then, putative cell bodies were

detected to feed into a constrained non-negative matrix factorization algorithm to decompose the 3-

dimensional video array into a 3-dimensional array representing the spatial footprint of each cell, as

well as a 2-dimensional matrix representing the calcium transients of each cell. The calcium transients

were then deconvolved to extract the estimated time of each calcium transient. These deconvolved

calcium activities were analyzed in these studies, after undergoing various transformations depending

on the specific analysis (see below). Cells recorded across sessions within a mouse were cross-

registered using a previously published open-source cross-registration algorithm, CellReg, using the

spatial correlations of nearby cells to determine whether highly correlated footprints close in space are

likely to be the same cell across sessions[4].

To align calcium imaging data with behavior, behavior recordings were first aligned to an idealized

template assuming a perfect sampling rate. This meant that if a recording session was 5min long, this

meant that there should be 300sec * 30frames/sec = 9000frames. All behavior recordings were within 4

frames of this perfect template. Calcium recordings recorded with a much more variable and dynamic

sampling rate. Then, for each behavior frame, the closest calcium imaging frame was aligned to that

frame, using the computer timestamp of that frame in milliseconds. No calcium imaging frame was re-

used more than twice.

General statistics and code/data availability

All analyses and statistics were done using custom-written Python and R scripts. Code detailing all the

analysis in this manuscript will be made available upon publication (https://github.com/denisecailab).

Calcium imaging data used in this manuscript will be made available using the Neurodata Without

Borders framework to seamlessly share data across institutions[5]. Statistical significance was assessed

with two-tailed paired and unpaired t-tests, as well as one-way, two-way, or three-way ANOVAs, linear

mixed effects models, or Chi-square test where appropriate. Significant effects or interaction were

followed with post-hoc testing with the use of orthogonal contrasts or with Benjamini-Hochberg

corrections for multiple comparisons. Significance levels were set to α=0.05. Significance for

comparisons: *p<=0.05; **p<0.01; ***p<0.001; ****p<0.0001. Sample sizes were chosen based on

previous similar studies. The investigators were not blinded to behavioral testing in calcium imaging

studies but were blinded to behavioral testing in all other experiments.

Ensemble reactivation analysis

To measure ensemble reactivation across the offline period, for each mouse, the matrix of neural

activity that was recorded during the offline session was z-scored along both axes (cells and time).

Cells were then broken up into ensembles based on whether they were previously observed to be

active. Previously active cells were defined based on whether they had a corresponding matched cell

via CellReg. On Offline1 after Neutral encoding, cells were either previously matched to an active cell

during Neutral encoding (Neutral ensemble) or had no previously matched cell (Remaining ensemble).

On Offline2, cells had a matched cell only with Neutral encoding and not Aversive encoding (Neutral

ensemble), a matched cell with Aversive encoding and not Neutral encoding (Aversive ensemble), a

matched cell on both Neutral encoding and Aversive encoding (Neutral ∩ Aversive ensemble), or no

matched cell (Remaining ensemble). For each ensemble, the activity of cells was averaged across

cells, and then averaged across time for each timebin.

Burst participation analysis

To measure population bursts, for each mouse, all cells that were recorded during that session were z-

scored along the time dimension, such that each cell was normalized to its own activity. By doing this,

no cell overly contributed to population bursts by having a very high amplitude event. Then, the mean

population activity across the whole population was computed across the session and that 1-

dimensional trace was z-scored. Time periods when the mean population activity reached above a

threshold of z=2 were considered burst events. During each of these burst events, each cell was

considered to have "participated" if its activity was above z=2 during the event. For each ensemble (as

defined in the previous section), the fraction of the ensemble that participated in each event was

computed, and then this was averaged across all events. The average participation of each ensemble

was compared across ensembles and across Low vs High Shock groups.

Ensemble co-participation analysis

To measure ensemble co-participation during bursts, first bursts were defined based on the z-scored

mean population activity of the whole population. Then, for each burst event, the z-scored mean

population activity was computed for the Neutral ensemble and for the Aversive ensemble (see

Ensemble reactivation analysis for ensemble definitions). For each population-level burst event, the

"participation" of the Neutral ensemble or Aversive ensemble was measured based on whether the

ensemble's mean population activity was above the z=2 threshold during the population level event.

The burst events where one ensemble participated without the other ensembles were considered

independent participations. The burst events where multiple ensembles simultaneously participated in

were considered co-participations. The fraction of burst events where each ensemble independently

and co- participated in were computed. Then, the same computation was made for all non-burst periods

to ask how frequently the ensembles burst independently and coincidentally outside of burst events.

<u>Time-lagged cross-correlation analysis</u>

To measure cross-correlations, first mean ensemble activities were computed for the Neutral ∩

Aversive, Neutral, and Aversive ensembles (see previous two sections). Then, each time series was

broken up into 120 sec bins. The Neutral ∩ Aversive ensemble was separately correlated with the

Neutral ensemble and the Aversive ensemble bin by bin. For each time bin, cross-correlations were

computed for lags up to a maximum of 5 frames (or ~160ms). The maximum correlation was taken for

each time bin, and the average correlation across time bins was computed. This led to, for each mouse,

an average correlation between the Neutral ∩ Aversive ensemble and the Neutral ensemble, and an

average correlation between the Neutral ∩ Aversive ensemble and the Aversive ensemble, across the

offline period.

<u>Encoding-to-Recall population vector correlation analysis</u>

To measure correlations between encoding and recall activity patterns, first for each mouse, only cells

that were active during both the encoding and recall session were included in the analysis and were

aligned across the two sessions. For the encoding session, the mean population activity across the

entire session was computed to produce one vector. Then, the recall session was broken up into 30-

second bins and the mean population activity vector was computed for each bin. The encoding vector

was correlated with each recall vector, as described before[6]. Finally, the correlations across all the

recall bins were averaged to produce one average correlation between encoding and recall, for each

mouse.

Methods References

Cai, D. J. *et al.* A shared neural ensemble links distinct contextual memories encoded close in time. *Nature* **534**, 115-118 (2016). https://doi.org:10.1038/nature17955

Pennington, Z. T. *et al.* ezTrack: An open-source video analysis pipeline for the investigation of animal behavior. *Sci Rep* **9**, 19979 (2019). https://doi.org:10.1038/s41598-019-56408-9

Dong, Z. *et al.* Minian, an open-source miniscope analysis pipeline. *Elife* **11** (2022). https://doi.org:10.7554/eLife.70661

Sheintuch, L. *et al.* Tracking the Same Neurons across Multiple Days in Ca(2+) Imaging Data. *Cell Rep* **21**, 1102-1115 (2017). https://doi.org:10.1016/j.celrep.2017.10.013

Rubel, O. *et al.* The Neurodata Without Borders ecosystem for neurophysiological data science. *Elife* **11** (2022). https://doi.org:10.7554/eLife.78362

Zaki, Y. *et al.* Hippocampus and amygdala fear memory engrams re-emerge after contextual fear relapse. *Neuropsychopharmacology* **47**, 1992-2001 (2022). https://doi.org:10.1038/s41386-022-01407-0

Acknowledgments
This work was supported by the DP2 MH122399, R01 MH120162, Brain Research Foundation Award, Klingenstein-Simons Fellowship, NARSAD Young Investigator Award, McKnight Memory and Cognitive Disorder Award, One Mind-Otsuka Rising Star Research Award, Hirschl/Weill-Caulier Award, Mount Sinai Distinguished Scholar Award, and Friedman Brain Institute Award, to DJC; the CURE Taking Flight Award, American Epilepsy Society Junior Investigator Award, R03 NS111493, R21 DA049568, R01NS116357, RF1AG072497 to TS; NIMH F31MH126543 to YZ; NIMH K99 MH131792 and BBRF Young Investigator Award to ZTP; NIMH R01 MH113071, NIA R01 AG013622, and Dr. Miriam and Sheldon G. Adelson Medical Research Foundation to AJS; F32NS116416 to ZCW. We would like to thank Brandon Wei, Mimi La-Vu, Christopher Lee for experimental support, and the members of the Cai and Shuman labs for their feedback throughout the duration of the project. We would like to thank Dr. Daniel Aharoni and Federico Sangiuliano Jimka for Miniscope-related support. We thank Dr. Margot Tirole, Dr. Claudia Clopath, Geoffroy Delamare, and Sima Rabinowitz for thoughtful discussions and input regarding analyses. We thank Dr. Patrick Davis for discussions throughout the project and for comments on the manuscript. We thank Stellate Communications for graphical design assistance. We thank William Janssen for microscopy support.

Author Contributions
DJC conceived the study. YZ, ZP, DMR, TF, TS, and DJC designed experiments. YZ, ZP, DMR, TF, AL, SL, and ZCW conducted behavioral experiments. YZ conducted calcium imaging experiments. YZ, DMR, TF, SL, and ZCW conducted chemogenetic experiments. YZ and DJC analyzed data. ZD and ZP contributed to development of data processing algorithms. YZ, ZP, DMR, TF, AL, ZD, SCS, HC, AJS, Mv, TS, AF, KR, and DJC contributed to interpretation of results. YZ and DJC wrote the manuscript. YZ, ZP, DMR, TF, AL, ZD, ZCW, SCS, HC, AJS, Mv, TS, AF, KR, and DJC edited the manuscript.

Competing Interests
The authors declare no competing interests.

Extended Figure 1

Extended Figure 1. Behavioral experiment controls.

A) Schematic to test the timecourse of prospective memory-linking (top). Mice underwent Aversive encoding and then either 5h, 1d, or 2d later they underwent Neutral encoding. The following day, mice were tested in the previously experienced Neutral context. Mice froze significantly more in the Neutral context when the Neutral context occurred within 5h of the Aversive context, compared to when it occurred one day or more after Aversive encoding (bottom). Main effect of timepoint ($F_{2,24} = 3.689, p = 0.04$) (*5h, N = 10 mice; 1d, N = 9 mice; 2d, N = 8 mice*). Post-hoc tests revealed a trend for higher freezing in the 5h timepoint compared to the 1d or 2d timepoints: 1d ($t_{16.38} = 2.137, p = 0.07$), 2d ($t_{13.45} = 2.38, p = 0.07$).

B) Schematic to test the timecourse of retrospective memory-linking (top). Mice underwent Neutral encoding, followed by Aversive encoding in a separate context 5h, 1d, or 2d later. The day following Aversive encoding, they were tested in the previously experienced Neutral context. Mice froze no differently in the Neutral context regardless of how long before Aversive encoding the Neutral context was experienced (bottom). No main effect of timepoint ($F_{2,27} = 0.73, p = 0.49$) (*5h, N = 10 mice; 1d, N = 10 mice; 2d, N = 10 mice*).

C) Schematic of low vs high shock retrospective memory-linking experiment (without calcium imaging as a replication). Mice underwent Neutral encoding followed by a low or high shock Aversive encoding two days later. In the subsequent 3 days, mice were tested in the Aversive context, and then Neutral and Novel contexts, counterbalanced.

D) Mice froze more in the Aversive context in High Shock vs Low Shock mice ($t_{14} = 5.04, p = 0.00018$) (*Low Shock, N = 8 mice; High Shock, N = 8 mice*).

E) High Shock mice exhibited higher freezing in Neutral vs Novel recall, while Low Shock mice did not. A priori post-hoc test: *High Shock ($t_7 = 2.65, p = 0.033$), Low Shock ($t_7 = 1.21, p = 0.133$) (Low Shock, N = 8 mice; High Shock N = 8 mice).*

Extended Figure 2

 Extended Figure 2. Offline hippocampal activity is necessary to drive retrospective memory-linking

A) Representative histological verification of viral expression in dorsal and ventral hippocampus. Blue represents DAPI and green represents AAV5-Syn-PSAM-GFP.

B) Schematic of the behavioral experiment disrupting hippocampal activity during the offline period. Mice were injected with AAV5-Syn-PSAM-GFP into dorsal and ventral hippocampus. Mice all had a Neutral experience and two days later a strong Aversive experience. Right after Aversive encoding, mice either had the hippocampus inactivated for 12hrs using the PSAM agonist, PSEM, or were given saline as a control. To do this, mice were injected four times, every three hours, to extend the manipulation across a 12-hour period. Two days later, mice were tested in the Neutral or a Novel context for freezing.

C) Control (saline-treated) mice displayed retrospective memory-linking (i.e., higher freezing during Neutral vs Novel recall), while mice that received hippocampal inhibition (PSEM-treated) no longer displayed retrospective memory-linking. Significant interaction between Experimental Group (PSEM vs Sal) and Context (Neutral vs Novel) ($F_{1,42} = 4.00, p = 0.05$) (*Saline Neutral, N = 12 mice; Saline Novel, N = 10 mice; PSEM Neutral, N = 12 mice; PSEM Novel, N = 12 mice*). Post-hoc tests demonstrate higher freezing in Neutral vs Novel contexts in the Sal group ($t_{19.84} = 2.57, p = 0.03$) and no difference in freezing in Neutral vs Novel contexts in the PSEM group ($t_{22} = 0.31, p = 0.76$).

D) Schematic of the behavioral experiment as above, but this time to test the effects of hippocampal inactivation on Aversive memory recall. Mice all underwent the Neutral and Aversive experiences as before, as well as PSEM or saline injections following Aversive encoding (as in Extended Figure 2B); however, two days following Aversive encoding, mice were tested in the Aversive context to test for an intact aversive memory.

E) Mice froze no differently in the Aversive context whether they had received hippocampal inhibition or not ($t_{13.9} = 0.32, p = 0.748$) (*Saline, N = 7 mice; PSEM, N = 9 mice*).

Extended Figure 3

Extended Figure 3. Neurons active during Aversive encoding are selectively reactivated offline and during Aversive recall.

A) Representative maximum intensity projection of the field-of-view of one example session (left). Spatial footprints of all recorded cells during the session, randomly color-coded (right).

B) Schematic of a single aversive experience. Mice had an Aversive experience followed by a 1hr offline session in the homecage. The next day, mice were tested in the Aversive context, followed by a test in a Novel context one day later. Calcium imaging in hippocampal CA1 was performed during all sessions.

C) Mice acquired within-session freezing during Aversive encoding (left); main effect of time ($F_{8,56} = 12.59, p = 3.87e-10, N = 8$ mice). And mice responded robustly to all three foot shocks, though their locomotion generally decreased across shocks, driven by increased freezing (right); main effect of shock number ($F_{2,14} = 7.45, p = 0.0154, N = 8$ mice) and main effect of PreShock vs Shock ($F_{1,7} = 581, p = 5.38e-8, N = 8$ mice), and no interaction.

D) Mice displayed a modest decrease in locomotion across the 1hr offline period ($R^2 = 0.064, p = 1.9e-8, N = 8$ mice).

E) Mice froze significantly more in the Aversive context than in a Novel context during recall ($t_7 = 165, p = 4e-6, N = 8$ mice).

F) Cells that were active during Aversive encoding and reactivated offline were significantly more likely to be reactivated during Aversive recall than cells active during Aversive encoding and not reactivated offline ($t_7 = 19.41, p = 2e-7, N = 8$ mice).

G) A larger fraction of cells active during Aversive recall than during Novel recall were previously active during Aversive encoding ($t_7 = 6.897, p = 0.0002, N = 8$ mice).

H) During the offline period, ~40% of the population was made up of cells previously active during Aversive encoding (top). This Aversive ensemble was much more highly active than the rest of the population during the offline period (bottom; A.U.) ($t_7 = 8.538, p = 0.00006, N = 8$ mice).

I) Each cell's activity was compared during locomotion vs during quiet rest (left; A.U.). A regression line was fit to the cells in the Aversive ensemble and in the Remaining ensemble separately, for each mouse. The Remaining ensemble showed greater activity during locomotion than during quiet rest (i.e., a less positive slope). The Aversive ensemble showed relatively greater activity during quiet rest than locomotion (i.e., a more positive slope) across mice (right) ($t_7 = 5.76, p = 0.047, N = 8$ mice).

J) Cells that had high levels of activity (A.U.) during Aversive encoding continued to have high levels of activity during the offline period (example mouse; left). There was a linear relationship between how active a cell was during Aversive encoding and how likely it was to be reactivated during the offline period (all mice; right) ($R^2 = 0.726, p = 1.25e-23, N = 8$ mice).

K) During the offline period, cells that would go on to become active during recall were more highly active than the Remaining ensemble during the offline period. The top represents the proportion of each ensemble (legend to its right). The cells that would become active during both Aversive and Novel recall were most highly active (A.U.). There was no difference in activity in the cells that would go on to be active in Aversive or Novel. Main effect of Ensemble ($F_{3,21} = 27.81, p = 1.65e-7, N = 8$ mice). Post-hoc tests: for Aversive vs Novel ($t_7 = 1.33, p = 0.22$), for Remaining vs Aversive ∩ Novel ($t_7 = 11.95, p = 0.000007$), for Remaining vs Aversive ($t_7 = 3.97, p = 0.005$), for Remaining vs Novel ($t_7 = 7.47, p = 0.0001$).

Extended Figure 4

Extended Figure 4. Low vs High Shock calcium imaging supplementary analyses.

A) Mice acquired within-session freezing during Aversive encoding. Mice that received high shocks (1.5mA) displayed more freezing than mice that received low shocks (0.25mA) (*Low Shock, N = 10 mice; High Shock, N = 12 mice*).

B) Mice responded robustly to each foot shock. High shock mice responded more strongly to each shock than low mice did (*Low Shock, N = 10 mice; High Shock, N = 12 mice*).

C) Relative to the first calcium imaging recording, mice showed comparable fractions of observed cells across the remaining sessions (*Low Shock, N = 8 mice; High Shock, N = 10 mice*).

D) Locomotion across the 1hr offline period after Neutral encoding (Offline1) and after Aversive encoding (Offline2) in Low and High Shock mice. Mice showed decreased locomotion across the offline period on both days. Low Shock mice did not locomote differently from High Shock mice during either offline period (*Low Shock, N = 10 mice; High Shock, N = 12 mice*).

E) In High Shock mice, Neutral recall cells were composed of more Neutral encoding cells being reactivated, compared to Novel recall cells. In Low Shock mice, Neutral recall cells and Novel recall cells were composed of similar fractions of Neutral encoding cells being reactivated. Significant interaction between Context (Neutral vs Novel) and Amplitude (Low vs High Shock) ($F_{1,12} = 6.81, p = 0.022$) (*Low Shock, N = 6 mice; High Shock, N = 8 mice*). Post-hoc tests, *Low Shock ($t_5 = 1.34, p = 0.24$), High Shock ($t_7 = 10.22, p = 0.000037$).*

F) In High Shock mice, Neutral recall cells were composed of more Aversive encoding cells being reactivated, compared to Novel recall cells. In Low Shock mice, Neutral recall cells and Novel recall cells were composed of similar fractions of Aversive encoding cells being reactivated. Significant interaction between Context (Neutral vs Novel) and Amplitude (Low vs High Shock) ($F_{1,12} = 4.75, p = 0.0499$) (*Low Shock, N = 6 mice; High Shock, N = 8 mice*). Post-hoc tests, *Low Shock ($t_5 = 0.59, p = 0.58$), High Shock ($t_7 = 5.46, p = 0.0019$).*

G) During Offline1, burst event frequency gradually decreased across the hour ($F_{11,143} = 4.43, p = 1.0e-5$). No difference across shock amplitudes ($F_{11,13} = 0.31, p = 0.587$) (*Low Shock, N = 7 mice; High Shock, N = 8 mice*). Significant interaction between Time and Amplitude ($F_{11,143} = 1.87, p = 0.047$). Follow-up repeated measures ANOVAs showed that both Low and High Shock groups showed a significant decrease in event rate across time (*Low Shock: $F_{11,66} = 4.13, p = 0.0001$; High Shock: ($F_{11,77} = 2.43, p = 0.01$).*

H) During Offline2, burst event frequency decreased across time ($F_{11,143} = 6.69, p = 0.000054$). No difference across shock amplitudes ($F_{1,13} = 0.0056, p = 0.94$) (*Low Shock, N = 7 mice; High Shock, N = 8 mice*).

I) Example process of identifying ensemble co-participations during bursts. Data in this panel are down-sampled from 30Hz to 1Hz for visualization purposes. On the left, the bottom matrix represents the neuronal activities for all neurons recorded across the offline period, color-coded by ensemble (see Ensembles legend). The top black trace represents the z-scored mean population activity across the hour. The yellow line represents a time slice of representative bursts (expanded on the right). In the middle, the whole population mean population activity is shown again, with the mean population activity of the Neutral, Neutral ∩ Aversive, and Aversive ensembles shown below. From these population activities, the time periods above threshold for the whole population were considered whole population bursts, and within those, we measured how frequently the other ensembles participated in these bursts. On the right, we zoom into two example whole population bursts in yellow. In the first one, at 629 sec into the recording, the Neutral ∩ Aversive and Aversive ensembles participated, and in the second one, at 655 sec, only the Aversive ensemble participated.

J) During Offline2, bursts as defined by each ensemble (rather than by whole population) decreased across the hour, with comparable frequencies across ensembles and amplitudes (*Low Shock, N = 7 mice; High Shock, N = 8 mice*).

K) Time-lagged cross correlations between the N∩A ensemble and the Neutral and Aversive ensembles during the offline period. Each of the three ensembles (N∩A, Neutral, and Aversive) were binned into 120 sec bins. Each time bin of N∩A ensemble activity was cross-correlated with the corresponding time bin of Neutral ensemble and Aversive ensemble activity. Cross-correlations were computed with a maximum time lag of 5 frames (or, ~160ms). For each mouse, the correlations were averaged across all time bins to get an average cross-correlation between the N∩A ensemble and Neutral ensemble (i.e., N∩A x N) and the N∩A ensemble by Aversive ensemble (i.e., N∩A x A). There was a significant interaction between Ensemble Combination and Low vs High Shock group

1189   ($F_{1,13} = 6.70, p = 0.02$) (*Low Shock, N = 7 mice; High Shock, N = 8 mice*). Post-hoc tests revealed that in High
1190   Shock mice, N∩A x N correlations were higher than N∩A x A correlations ($t_7 = 3.97, p = 0.01$) whereas they were
1191   no different in Low Shock mice ($t_6 = 0.83, p = 0.44$).

 Extended Figure 5

Burst Classification

A

B Burst Example   Non-Burst Example

Burst Frequency Compared to Shuffle

C **D**

Population Activity Skew Compared to Shuffle

E **F**

Lack of Periodicity of Bursts

G **H**

Decrease in Locomotion During Bursts

I **J**

Aversive Ensemble Participation During Bursts

K **L**

Extended Figure 5. Neurons active during Aversive encoding selectively participate in burst events offline.

A) Example of a burst event quantified in this figure. The top trace represents the z-scored mean population activity within one of the offline recordings. Three timepoints were chosen (overlaid in circles), the middle representing the peak of a burst event, and the timepoints to its left and right representing t-2sec and t+2sec from the peak, respectively. The bottom three matrices represent binarized spatial footprints depicting the spatial footprints of the cells sufficiently active to participate in a burst (z>2). The matrices represent the timepoints of the three datapoints above it, ordered by time.

B) Representative process of extracting ensemble participations (one mouse example). The left is an example burst period, with the rows in the heatmap representing the activity of the recorded cells during that session, binarized by z>2 and color-coded by whether they were previously active during Aversive encoding (Aversive ensemble, blue) or if they were not previously active (Remaining ensemble, grey). The black trace above represents the z-scored mean population activity during this period, demonstrating a brief burst in activity accompanied by participation by a significant fraction of neurons. On the right is an example non-burst period, where mean population activity remains below threshold.

C) Neuron activities were circularly shuffled 1000 times relative to one another and the mean population activity was re-computed each time. This shuffling method preserved the autocorrelations for each neuron while disrupting the co-firing relationships between neurons. The burst frequency was computed for each of these shuffles to produce a shuffled burst frequency distribution (gray histogram), to which the true burst frequency was compared (blue dotted line). This is an example mouse.

D) The mean burst frequency for the shuffled distribution was computed and compared to the true burst frequency for each mouse. True burst frequencies were greater than shuffled burst frequencies in every mouse ($t_7 = 6.159$, $p = 0.000463$, $N = 8$ mice), suggesting that during the offline period, hippocampal CA1 neurons fire in a more coordinated manner than would be expected from shuffled neuronal activities.

E) As in Extended Figure 5C, neuron activities were shuffled, and mean population was re-computed each time. From this population activity trace, the skew of the distribution was computed. If there were distinct periods where many neurons simultaneously fired, we hypothesized that the true distribution of mean population activity would be more skewed with a strong right tail demonstrating large and brief deflections, compared to shuffled neuronal activities. We computed the skew of each shuffled mean population activity, to produce a distribution (gray histogram), to which the true mean population's skew was compared (blue dotted line). This is an example mouse.

F) The mean skew for the shuffled distribution was computed and compared to the true skew of the mean population activity for each mouse. The true skew was greater than the shuffled skew in every mouse ($t_7 = 13.36$, $p = 0.000003$, $N = 8$ mice), supporting the idea that the mean population activity undergoes brief burst-like activations requiring the coordinated activity of groups of neurons.

G) Matrix of burst events for an example mouse, stacked along the y-axis and centered on time t=0 (top), and the average mean population activity around each burst event (bottom).

H) As in Extended Figure 5G but averaged across all mice. Each thin line represents one mouse, and the thick black line represents the mean across mice with the grey ribbon around it representing the standard error ($N = 8$ mice). There is no periodicity to when these burst events occur.

I) Locomotion of an example mouse during each burst event stacked along the y-axis (top), and the mean locomotion around burst events (bottom). Mice showed a robust and brief slowing down ~1sec before each burst event, before increasing locomotion back up ~2sec later.

J) As in Extended Figure 5I but averaged across all mice. Each thin line represents one mouse, and the thick black line represents the mean across mice with the grey ribbon around it representing the standard error ($N = 8$ mice). This demonstrates a robust and reliable decrease in locomotion around the onset of burst events.

K) The burst event frequency decreased across the hour ($F_{11,77} = 6.91$, $p = 5.66e-8$, $N = 8$ mice).

L) A larger fraction of the Aversive ensemble vs the Remaining ensemble participated in each burst event (left) ($t_7 = 3.68$, $p = 0.0079$, $N = 8$ mice).