1    **Structures of trehalose-6-phosphate synthase, Tps1, from the fungal pathogen**
2    ***Cryptococcus neoformans*: a target for novel antifungals**
3

4    Erica J. Washington[1], Ye Zhou[2], Allen L. Hsu[3], Matthew Petrovich[3], Mario J. Borgnia[3], Alberto
5    Bartesaghi[1,2], Richard G. Brennan[1]
6

1.  Department of Biochemistry, Duke University School of Medicine, Durham, North Carolina,
27710 USA
2.  Department of Computer Science, Duke University, Durham, North Carolina, 27708 USA
3.  Genome Integrity and Structural Biology Laboratory, National Institute of Environmental
Health Sciences, National Institutes of Health, Department of Health and Human Services,
Research Triangle Park, NC 27709 USA
13

14   [*]To whom correspondence should be addressed: richard.brennan@duke.edu

15
16
17
18

19   **Running Title**
20   Cryo-EM structures of CnTps1

22   **Keywords**
23   Tps1, Cryptococcus, cryo-electron microscopy, trehalose, fungal pathogens, Basidiomycetes

**Abstract**

Invasive fungal diseases are a major threat to human health, resulting in more than 1.5 million deaths worldwide each year. Yet the arsenal of antifungal therapeutics remains limited and is in dire need of novel drugs that target additional fungal-specific biosynthetic pathways. One such pathway involves the biosynthesis of trehalose. Trehalose is a non-reducing disaccharide composed of two molecules of glucose that is required for pathogenic fungi, including *Candida albicans* and *Cryptococcus neoformans*, to survive in their human hosts. Trehalose biosynthesis is a two-step process in fungal pathogens. Trehalose-6-phosphate synthase (Tps1) converts UDP-glucose and glucose-6-phosphate to trehalose-6-phosphate (T6P). Subsequently, trehalose-6-phosphate phosphatase (Tps2) converts T6P to trehalose. The trehalose biosynthesis pathway has been identified as a top candidate for novel antifungal development based on quality, occurrence, specificity, and assay development. However, there are currently no known antifungal agents that target this pathway. As initial steps to develop Tps1 from *Cryptococcus neoformans* (CnTps1) as a drug target, we report the structures of full-length *apo* CnTps1 and CnTps1 in complex with uridine diphosphate (UDP) and glucose-6-phosphate (G6P). Both CnTps1 structures are tetramers and display D2 (222) molecular symmetry. Comparison of these two structures reveals significant movement towards the catalytic pocket by the N-terminus upon ligand binding and identifies key residues required for substrate-binding, which are conserved amongst other Tps1 enzymes, as well as residues that stabilize the tetramer. Intriguingly, an intrinsically disordered domain (IDD), encompassing residues M209 to I300, which is conserved amongst Cryptococcal species and closely related Basidiomycetes, extends from each subunit of the tetramer into the "solvent" but is not visible in the density maps. Although, activity assays revealed that the highly conserved IDD is not required for catalysis *in vitro*, we hypothesize that the IDD is required for *C. neoformans* Tps1-dependent thermotolerance and osmotic stress survival. Characterization of the substrate specificity of CnTps1 revealed that UDP-galactose, an epimer of UDP-glucose, is a very poor substrate and inhibitor of the enzyme and highlights the exquisite substrate specificity of Tps1. *In toto*, these studies expand our knowledge of trehalose biosynthesis in *Cryptococcus* and highlight the potential of developing antifungal therapeutics that disrupt the synthesis of this disaccharide or the formation of a functional tetramer and the use of cryo-EM in the structural characterization of CnTps1-ligand/drug complexes.

## Introduction

Invasive fungal diseases (IFDs) caused by pathogenic fungi such as *Cryptococcus*, *Candida* and *Aspergillus* are a major threat to human health, resulting in more than one and a half million deaths worldwide each year (1, 2). These mortality rates are surprisingly high, given the fact that fungal infections are usually associated with superficial infections of skin and nails. However, IFDs result in increased mortality in immunocompromised populations, including HIV-infected patients and patients receiving immunosuppressive therapies (1-3). Additionally, the severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2; COVID-19) pandemic has resulted in a significant population of COVID-19 positive patients with increased susceptibility to IFDs, as has been demonstrated for pulmonary aspergillus and invasive candidiasis infections (4-10).

Unfortunately, the current arsenal of antifungal drugs, consisting of the polyenes, azoles and echinocandins, is insufficient to manage the mortality caused by IFDs due to significant off-target effects, the rapid emergence of resistance to antifungal therapeutics and the emergence of intrinsically drug resistant fungal pathogens, such as *Candida auris* and *Candida glabrata* (11-13). Therefore, in addition to the development of new formulations of current antifungal drugs, there is a critical need for the development of new classes of broad-spectrum antifungal drugs that are fast-acting and safe. The identification of targets that are not present in the human host is critical to the development of antifungal drugs with low toxicity. This is often difficult, as both fungi and humans are eukaryotes. However, pathways involved in the fungal stress response have been implicated recently as key targets, including the trehalose biosynthesis pathway (14-17).

Trehalose is a non-reducing disaccharide composed of two glucose molecules linked by an α,α-1,1-glycosidic bond. Fungal cells synthesize trehalose to protect their proteins and membranes from stresses encountered during the infection process (18-25). The canonical trehalose biosynthesis machinery in pathogenic fungi consists of trehalose-6-phosphate synthase, Tps1, a glucosyltransferase that converts uridine diphosphate glucose (UDPG) and glucose-6-phosphate (G6P) to trehalose-6-phosphate (T6P). Subsequently, trehalose-6-phosphate phosphatase (Tps2), removes the phosphate group to generate the final product of trehalose (Figure 1A). In *Candida albicans* and *Aspergillus fumigatus* there is an additional protein in the pathway, Tps3, with no known enzymatic function. The machinery to synthesize trehalose is found in plants, insects, fungi and even bacteria, but not in humans (26). For this reason, an antifungal therapeutic that targets the trehalose biosynthesis pathway could well result in a drug with minimal off-target effects, leading to greatly reduced toxicity in patients. Indeed, disruption of

the *TPS1* gene in *Cryptococcus neoformans* results in fungi that are avirulent in mice and rabbits

and, interestingly, disruption of *TPS2* was followed by the accumulation of the toxic intermediate

trehalose 6-phosphate, causing fungal cell death (27). Additionally, in *C. neoformans* trehalose

acts as a stress protectant and is required for growth at high temperatures (27). Similarly,

disruption of the *TPS1* and *TPS2* genes in *Cryptococcus gattii* results in decreased infectivity in

mice (28). Similar phenotypes indicating the importance of Tps1 and Tps2 to cell survival and

sometimes virulence are also seen in trehalose biosynthesis mutants in *C. albicans* (29-31) and

*A. fumigatus* (32, 33). Therefore, we hypothesize that targeting the trehalose biosynthesis

pathway will result in the development of a novel broader-spectrum antifungal drug, which would

be both highly effective and, importantly, safe for use in immunocompromised patients.

Pertinent to the development of an antifungal therapeutic which targets the trehalose

biosynthesis pathway is an intimate knowledge and understanding of the structures of enzymes

and proteins involved in this important pathway. Several structures of Tps1 and Tps2 from

nematodes, bacteria and pathogenic fungi have been determined using x-ray crystallography (34-

43). Subsequently, compounds that inhibit trehalose biosynthesis enzyme functions by targeting

their catalytic pockets have been investigated but not yet advanced for clinical use (39, 44-49).

As a key starting point to develop inhibitors of CnTps1, we describe structures of this 669-residue

(74 kDa per protomer) enzyme in both its *apo* (substrate-free) form and in complex with UDP and

G6P using single particle cryo-electron microscopy (cryo-EM). Unlike Tps1 from *Candida albicans*

(40), CnTps1 has three major insertions, 55 residues at the N-terminus, 63 residues at the C-

terminus and a 92-residue intrinsically disordered domain (IDD) inserted within the core of the

enzyme, all of unknown function. Residues located in the catalytic pocket and the IDD were

assayed for their contribution to the activity of this enzyme. Intriguingly, the IDD of CnTps1 is not

required for CnTps1 catalytic activity *in vitro*. Substrate specificity was investigated further and

revealed the exquisite ability of CnTps1 to discriminate at the level of a single interaction. Hence,

this work lays the groundwork for subsequent CnTps1 structure-based drug design.

## Results

### Trehalose biosynthesis protein CnTps1 self-associates in bacterial two-hybrid assays

The trehalose biosynthesis pathway is a two-step process. Tps1, trehalose-6-phosphate synthase, converts uridine diphosphate glucose (UDPG) and glucose-6-phosphate (G6P) into trehalose-6-phosphate (T6P). The phosphate group is subsequently removed from T6P by trehalose-6-phosphate phosphatase (Tps2) to yield trehalose (Figure 1A) (17). Previously the structures of Tps1/OtsA from *Candida albicans* (40), *Aspergillus fumigatus* (40), *E. coli (35, 37, 38, 43)*, *Streptomyces venezuelae* (34) and *Magnaporthe oryzae* (43) were determined using x-ray crystallography. Additionally, the structure of the catalytically inactive Tps1-like N-terminal domain of *C. albicans* Tps2 was determined (41).

Inspection of the Tps1 crystal structures from *Candida albicans* (40), *Aspergillus fumigatus* (40), *E. coli* (37, 38), *Streptomyces venezuelae* (34) and *Magnaporthe oryzae* (43) and the catalytically inactive Tps1-like N-terminal domain of *C. albicans* Tps2 (41) revealed that Tps1 most likely exists as a homo-tetramer. To determine if subunits of the *Cryptococcus neoformans* var. *grubii* strain H99 Tps1, hereafter referred to as CnTps1, self-associate, a bacterial two-hybrid system based on the LexA DNA-binding repressor was established (50). Expression of full-length LexA-CnTps1 in the bacterial two-hybrid system was not detected. However, LexA-CnTps1$_{56-669}$, in which the unstructured 55 N-terminal residues of CnTps1 were deleted, does self-associate in the bacterial two-hybrid assay, as evidenced by a reduction in β-galactosidase activity after protein expression was induced with 1 mM IPTG (Figure 1B). An anti-LexA Western blot shows the expression of LexA-CnTps1$_{56-669}$ at the appropriate molecular weight (Figure 1C). These data indicate that CnTps1 can self-associate and support the hypothesis that CnTps1 forms homo-oligomers.

### The structure of the *apo* (substrate-free) CnTps1 homo-tetramer

CnTps1 is approximately 200 residues larger than other Tps1 orthologues, due to the addition of a 55-residue N-terminal extension, a 63-residue C-terminal extension and an internal insertion from residues M209 to I300 (Supplemental Figure 1). Each extension is predicted to be structurally disordered. In order to determine the structure of CnTps1, the full-length (74 kDa, 669-residue per protomer) CnTps1 protein was expressed with an N-terminal hexahistidine tag and purified from *E. coli* via Ni$^{2+}$-NTA affinity chromatography (Supplemental Figure 2A). Following confirmation that 6xHis-CnTps1 was enzymatically active (Supplemental Figure 2B), the protein sample was subjected to single particle cryo-electron microscopy analysis. 2,520 micrographs were collected from grids prepared with 6xHis-CnTps1 (Supplemental Figure 3). No additional

ligands were added to the protein sample. 2D classification, performed on 907 micrographs with

resolution higher than 4.0 Å, revealed tetrameric CnTps1 particles (Supplemental Figure 3). 3D

classification and initial refinement steps were performed without imposing symmetry

(Supplemental Figure 3), resulting in a 5.3 Å resolution map. Additional refinement and polishing

steps, including adding D2 symmetry, resulted in an *apo* CnTps1 map at 3.3 Å global resolution,

as estimated by the "gold standard" Fourier shell correlation (FSC) = 0.143 criteria (Data Table 1

and Supplemental Figure 4A,B). Four protomers can be readily differentiated in the *apo* CnTps1

cryo-EM map (Supplemental Figure 4C). The final map shows features of well-resolved side

chains of many amino acid residues (Supplemental Figure 4D). The atomic model for *apo*

CnTps1, which was built based on the cryo-EM reconstruction, supports the conclusion that *apo*

CnTps1 forms a homo-tetramer (Figure 2).

The four CnTps1 protomers of the tetramer, designated as protomers 1 – 4, assemble into

the tetrameric structure with dimensions of 100.0 x 110.3 x 107.6 Å (Figure 2A-C). The four

protomers interact via two α-helical domains that are found in the central region of the homo-

tetramer (Figure 2A). The unbound substrate-binding pockets, are open and outward-facing, and

therefore accessible to ligands and substrates (Figure 2A,B). The "bottom-facing" view of the

homo-tetramer shows a staggered appearance of the top and bottom pair of protomers (Figure

2C).

Each CnTps1 protomer is comprised of N and C-terminal lobes, containing residues 1-

378 and 379-669, respectively. Both lobes are spanned by an α-helix, formed by residues 572-

603 (Figure 2D). Each domain contains a modified Rossman fold that is characteristic of the

retaining glycosyltransferase family (40, 51) (Figure 2D), and similar to the published CaTps1 and

CaTps1-like N-terminal domain of CaTps2 structures (Supplemental Figure 5) (34, 41, 43). A

common feature that is observed in these substrate-free structures is a lack of density in portions

of the N-terminus, suggestive of conformational flexibility in the absence of ligands. Indeed, the

CnTps1 cryo-EM map shows clear density for secondary structures in the C-terminus, whilst the

density was poor in several parts of the N-terminal domain, including that for residues 1-58, 68-

100, 127, 180, 210-301, 354-357 and 364-368 (Figure 2D). Not unexpectedly, the N-terminus

extension, the C-terminus and the internal insertion, residues 209-300, are not visible in this

structure suggesting multiple accessible conformations or intrinsic disorder.

**The structure of the CnTps1-UDP-G6P complex**

In order to identify residues in the catalytic pocket of CnTps1 as well as to visualize any

conformational changes necessary for substrate binding and catalysis, we determined the cryo-

EM structure of the *C. neoformans* H99 Tps1 in complex with the substrate, G6P, and a product,

UDP, to 3.1 Å global resolution (Figure 3A-D). 3,182 movies were collected from grids prepared

with 6xHis-CnTps1 and an excess of UDP and G6P (Supplemental Figure 6). The 2D

classification revealed tetrameric structures like those observed for *apo* CnTps1 (Supplemental

Figure 6). The cryo-EM reconstruction of 6xHis-CnTps1 bound to UDP and G6P also revealed a

tetrameric assembly. Final refinement was performed with D2 symmetry imposed (Supplemental

Figure 6). The addition of G6P and UDP resulted in a higher global resolution of the CnTps1

structure when compared to the *apo* structure (Data Table 2). The substrate-binding pocket is

more ordered (Supplemental Figure 7A,B). Four protomers can be detected in the CnTps1-UDP-

G6P cryo-EM map (Supplemental Figure 7C) and the final map shows features of well-resolved

side chains (Supplemental Figure 7D).

As observed in the *apo* structure, the N and C-termini and the internal insertion remain

unstructured in the presence of ligands. The structure of CnTps1 bound to UDP and G6P is similar

to that of the published CaTps1-UDP-G6P complex and the N-terminus of CnTps2 with root mean

square deviations (rmsds) of 1.2 Å for 450 corresponding $C_\alpha$ atoms and 1.5 Å for 468

corresponding $C_\alpha$ atoms, respectively (40, 41, 52). Views of the CnTps1-UDP-G6P tetramer show

the central binding region of UDP and G6P inside each protomer (Figure 3A-D). A view of each

protomer of the CnTps1-UDP-G6P complex reveals the substrate-binding site, with the UDP

bound to the C-terminal domain and the G6P adjacent to the N-terminal domain (Figure 3D). The

binding sites of the ligands are in excellent agreement with the published structure of CaTps1-

UDP-G6P and CaTps1-UDPG (40).

**Ligand-induced conformational changes in *C. neoformans* Tps1**

Significant movement and conformational changes of the N-terminal domain are observed

due to ligand binding to CnTps1 (Figure 4A,B). Structural alignments of the individual N-terminal

domain (residues 56 – 378) and C-terminal domain (residues 379 – 603) of the *apo* Tps1 protein

onto the corresponding domains of UDP-G6P bound CnTps1 reveal rmsds of 1.1 and 0.4 Å,

respectively. Superposition of the complete structure of *apo* CnTps1 onto CnTps1-UDP-G6P

reveals an rmsd of 1.4 Å for 433 corresponding $C_\alpha$ atoms. These rmsd values are a result of

ligand-induced conformational "closing" and stabilization of CnTps1, with the movement occurring

primarily in the N-terminus. Thus, *apo* CnTps1 protomers are in a more "open" conformation as

compared to the ligand-bound subunits. Systematic analysis of the domain movements using the

program DynDom (53, 54) shows that the large movements of the N-terminal G6P-binding lobe

centered around α-helix 2 (residues 112 to 125), which is located on the surface of the N-terminal

domain, result in a rotation of approximately 20° (Supplemental Figure 8). This rotation causes

an inward, approximately 7 Å, movement of α-helix 2 (Figure 4B). The linker that facilitates this

movement includes residues 113 through 145. This rotation results in a 29% closure of the N and

C-terminal lobes around the substrate-binding pocket. Interestingly, the local resolution proximal

to CnTps1 α-helix 2 is the lowest in the structure, which may indicate flexibility and a propensity

for movement, consistent with the results of the DynDom analysis. Since the CnTps1-UDP-G6P

complex contains both a product (UDP) and a substrate (G6P) as ligands, the completely closed

form of the protein is likely not visualized in this structure. Regardless, the movement of this

domain would facilitate the $S_N$-i catalytic mechanism proposed for Tps1 proteins in the GT-B fold

retaining glycosyltransferase family, including *C. albicans* Tps1 and *E. coli* OtsA (35, 40, 51, 55).

Indeed, superposition of UDP onto the UDPG from the CaTps1-UDPG complex (PDB ID 5HUT)

reveals that the G6P substrate bound to CnTps1 is proximal to and properly aligned with the

glucose of the UDPG to allow the ready formation of the 1,1 glycosidic bond, required for the

formation of T6P (Supplemental Figure 9).

**CnTps1 substrate-binding residues**

We were able to detect strong density for the bound ligand and substrate-binding residues

of CnTps1 (Figure 5A). The UDP molecule is bound to the C-terminal portion of each protomer

(Figure 5B). The orientation of the uracil base and CnTps1 is mediated by a hydrogen bond

between the exocyclic O4 and the V492 amide NH group. The O2 and O3 hydroxyl groups of the

ribose ring of UDP form hydrogen bonds with the side chain of E522. The orientation of the

phosphate groups of UDP in CnTps1 is mediated by several interactions. One hydrogen bond is

made between the α-phosphate group and the backbone amide group of L518. In CaTps1 (40)

there is a conserved Arg-Lys pair (residues R280 and K285) that interacts with the phosphate.

We were unable to detect side chain density for the equivalent CnTps1 Arg, indicating movement

of the substrate-binding pocket, potentially ready to release the product UDP. There is, however,

a hydrogen bond between K420 and the β-phosphate. Based on an overlay with the CaTps1-

UDPG structure, we predicted that the glucose moiety in the native substrate, UDPG, would be

stabilized by CnTps1 D514 and positioned to deprotonate G6P (Figure 5B and Supplemental

Figure 10A) (40). The G6P molecule is located adjacent to the N-terminal lobe in each protomer.

A key interaction stabilizing G6P is residue R453, which forms hydrogen bonds and ionic

interactions with the G6P phosphate group (Figure 5C).

CnTps1 substrate-binding residues K420, D514, L518, E522 and R453 are conserved

amongst Tps1 proteins from *C. neoformans*, *C. albicans*, *A. fumigatus*, *M. oryzae*, *S. venezuelae*

and *E. coli* (Supplemental Figure 1 and Supplemental Figure 10B). In *C. albicans* Tps1 these

residues have been demonstrated to be required for hyphal development and thermotolerance

(40). Site-directed mutagenesis was used to generate variants of CnTps1 in which potential

substrate-binding residues were mutated as follows: E522A, K420A, D514A and R453A. All

CnTps1 variants eluted similarly to wild-type CnTps1 on the size exclusion column, indicating that

they are properly folded (Supplemental Figure 11). Tps1 coupled enzyme assays revealed that

mutation of the residues that interact with UDP/UDPG resulted in severe loss-of-function (Figure

5D). Mutation of R453 to an alanine, which is the single identified G6P-interacting residue,

resulted in greatly reduced CnTps1 activity as well (Figure 5E).

**Interactions within the CnTps1 homo-tetramer**

The interactions between several side chains in the interfaces between CnTps1 protomers

explain the formation and stabilization of the CnTps1 homo-tetramer. The two key interfaces

responsible for formation of the CnTps1 homo-tetramer are referred to as interface #1 and

interface #2 (Supplemental Figure 12A). According to PDBePISA analysis (56), the buried surface

areas for each interface are similar, with 1062.9 Å buried in interface #1 and 986.2 Å buried in

interface #2. Interestingly, the calculated free energy (ΔG) of the two interfaces is -10.7 kcal/mol

and 0.1 kcal/mol, respectively, which suggests the former interface is a critical component of

oligomerization.

The key interactions in interface #1 are the hydrogen bonds between N467 and E468, as

well as R403 and E481 (Supplemental Figure 12B,C). Interface #2 also has hydrogen bonds that

are critical to the stability of the homo-tetramer, including hydrogen bonds between the E314

backbone oxygen and R513 side chain (Supplemental Figure 12D). Ionic interactions between

R513 and E314 and between R322 and E538 also contribute to the formation of interface #2.

None of the interface residues directly contact UDP or G6P, indicating that any effect mutation of

these residues might have on activity would be allosteric. Our cryo-EM results confirm that the

homo-tetrameric form of CnTps1 accurately represents the tetrameric structure in solution and is

not an artifact of the crystal lattice packing.

**CnTps1 contains a conserved intrinsically disordered domain**

As described above, CnTps1 contains three unstructured insertions (Supplemental Figure

13A). These regions have likely prevented the determination of its structure with x-ray

crystallography, as these regions are the main differences between CnTps1 and other crystallized

Tps1 orthologues, such as CaTps1. The most prominent unstructured region in CnTps1 is an

internal insertion of 92 residues (residues M209-I300), referred to hereafter as the Intrinsically

Disordered Domain or IDD (Supplemental Figure 13A). Interestingly, the CnTps1 IDD is found

only in the Tps1 proteins in fungi within the fungal division, Basidiomycota (Supplemental Figure

13B). More specifically, the CnTps1 IDD is highly conserved within Cryptococcal species and

closely related Basidiomycetes, with *C. neoformans*, *C. gattii* and *C. deneoformans* containing

IDDs with greater than 92% identity (Supplemental Figure 13A,B). The CnTps1 IDD is not present

in the Ascomycota division or the Mucoromycota phylum (Supplemental Figure 13B).

As anticipated, density for the CnTps1 IDD was not detected in the structures of *apo*

CnTps1 or CnTps1-UDP-G6P, consistent with the idea that the IDD is conformationally flexible.

However, based on the CnTps1 structures, we can determine that the IDD exits and returns to

the structured regions in adjacent parts of the N-terminus of CnTps1. To determine if the IDD is

required for CnTps1 activity *in vitro*, we designed and purified a chimeric protein in which the 92-

residue CnTps1 IDD was replaced with the six residues (GNKKKN), which *C. albicans* Tps1

utilizes to connect α-helix 5 to β-strand 4 (Supplemental Figure 14). We refer to this protein as

CnTps1 ΔIDD.

To determine the effect that the CnTps1 IDD has on CnTps1 function, we purified CnTps1

ΔIDD and confirmed that it was reduced in size compared to wild-type CnTps1, as determined by

size exclusion chromatography (Supplemental Figure 11). Notably, deletion of the CnTps1 IDD

did not affect formation of the tetramer (Supplemental Figure 11). Circular dichroism experiments

also confirmed that CnTps1 ΔIDD is properly folded (Supplemental Figure 15). Coupled enzyme

activity assays with CnTps1 ΔIDD were performed and revealed that the CnTps1 IDD is not

required for enzymatic activity of CnTps1 *in vitro* (Supplemental Figure 13C).

**Specificity of CnTps1 catalytic pocket**

To assess the substrate specificity of the CnTps1 catalytic pocket, we tested the catalytic

activity of CnTps1 in the presence of UDP-Galactose (UDP-Gal). The hydroxyl group on the 4th

carbon is oriented in opposite orientation in glucose and galactose. In the CaTps1-UDPG

structure, all of the extracyclic hydroxyl groups are easily detected, revealing that the O4 hydroxyl

hydrogen forms a hydrogen bond with the peptide backbone of CaTps1 N382 (40). Similarly, the

peptide backbone of CnTps1-UDP-G6P N517 would interact with the O4 hydroxyl of the glucose

moiety of UDPG (Supplemental Figure 16A). Modeling UDP-Galactose into the catalytic pocket

reveals the O4 hydroxyl points away from N517, preventing the formation of this critical hydrogen

bond (Supplemental Figure 16B). The loss of this interaction has a drastic effect on the activity of

CnTps1 with UDP-Gal as demonstrated by a reduction of approximately 96%, compared to

CnTps1 with UDPG as a substrate (Figure 6A), suggesting that the correct positioning of the O4

hydrogen mediated by its interaction with N517 is required for CnTps1 activity. This interaction

likely helps orient the entire UDPG molecule in the CnTps1 substrate-binding pocket for efficient

substrate-assisted catalysis.

To further explain the effect of UDP-Gal on CnTps1 activity, we performed a competition

assay. The combination of equal molar ratios of UDPG and UDP-Gal to CnTps1 resulted in a 12%

reduction in activity, compared to only adding UDPG to the assay (Figure 6B). Interestingly, the

addition of 10-fold excess UDP-Gal compared to UDPG resulted in a reduction of CnTps1 activity

by approximately 30% (Figure 6B), indicating that UDP-Gal is capable of binding CnTps1 but is a

weak binder and likely a weak competitive inhibitor, by binding to the substrate-binding pocket of

CnTps1.

**Discussion**

In order to develop the trehalose biosynthesis pathway as an antifungal drug target, we

must improve our understanding of the structure and function of trehalose biosynthesis proteins.

Here we report the use of cryo-EM to determine the structures of *apo* CnTps1 and CnTps1 bound

to UDP and G6P.

We confirmed that CnTps1 forms a tetramer in solution, indicating that previously reported

structures of Tps1 proteins determined by x-ray crystallography and exhibiting tetrameric

complexes are not simply an artifact of the crystal lattice packing. Our current hypothesis is that

trehalose biosynthesis proteins also form hetero-tetrameric complexes. Interactions amongst Tps

proteins (Tps1, Tps2, Tps3 and Tsl1) have been demonstrated in *S. cerevisiae* (57, 58). The

function of these complexes may be to sequester the highly cytotoxic trehalose-6-phosphate

(T6P) from the cytoplasm, to target trehalose biosynthesis to a proper subcellular localization (59)

or to allow for quick and efficient production of trehalose when fungal pathogens are exposed to

environmental stress. Future work shall include confirmation of the formation of heterocomplexes

both *in vitro* and *in vivo* and determination of the structure of the heterocomplex using cryo-

electron microscopy.

We have also reported here a conformational change of the individual protomers of

CnTps1 upon binding of UDP and G6P. Comparison of the *apo* CnTps1 and CnTps1-UDP-G6P

structures reveals an inward movement of the N-terminus of CnTps1 when bound to ligand. Most

of the movement occurs in α-helix 2 and surrounding residues. The closure of the ligand-bound

CnTps1 protomer is consistent with substrate-assisted catalysis, the proposed enzymatic

mechanism of Tps1 enzymes. Consistent with previous reports of the predicted catalytic activity

of fungal Tps1 proteins, we also propose that UDPG binds first, followed by G6P. The proximity

of the substrates in the substrate-binding pocket of CnTps1 is demonstrated in Supplemental

Figure 9. However, we now can see aspects of the catalytic mechanism occurring in the context

of the CnTps1 homo-tetramer for the first time.

The more ordered CnTps1-UDP-G6P substrate-binding pocket allows us to observe

specific interactions between the ligands and CnTps1 substrate-binding residues. The

identification of substrate-binding residues, K420, D514, E522 and R453, provides key insights

into the mechanism of the Tps1 glycosyltransferase as well as demonstrates the key roles of

these residues in substrate specificity (Figure 5A-C). Mutation of any of these residues is sufficient

to inhibit CnTps1 catalytic activity (Figure 5D,E). Additionally, our determination of substrate-

binding variants which lead to a loss of activity of CnTps1 provides the framework for further

investigation of a fully closed substrate-trapped cryo-EM structure of the CnTps1 homo-tetramer

in the future. This structure will provide fuller information regarding the CnTps1 catalytic
mechanism by allowing visualization of CnTps1 bound to the native substrates, UDPG and G6P.

The substrate-binding residues identified in the catalytic pocket of Tps1 are highly
conserved amongst fungal pathogens, including *C. albicans* (Supplemental Figure 1 and
Supplemental Figure 10B), supporting the proposition that a broader-spectrum antifungal
therapeutic can be developed by targeting the catalytic pocket of Tps1. Additionally, we have
been able to demonstrate that the CnTps1 substrate-binding domain is highly specific and
constrained as underscored by its inability to utilize UDP-Galactose as a substrate, which is an
epimer of UDPG (Figure 6 and Supplemental Figure 16).

Although the key interfaces of the CnTps1 homo-tetramer are not as highly resolved as
the substrate-binding pocket, specific residues that may play a role in the formation of the homo-
tetramers were identified (Supplemental Figure 12). It remains to be determined whether
tetramerization of CnTps1 is required for catalytic activity. We have evidence that disruption of *C.*
*albicans* Tps1 oligomers, achieved by the mutation of two residues, results in a significant
reduction of the ability of CaTps1 to generate T6P and also prevents strains of *C. albicans tps1*Δ
expressing monomeric Tps1 from proliferating in mice (data not published). Therefore, in addition
to targeting the substrate-binding pocket of CnTps1, the interfaces critical to the formation of the
CnTps1 tetramer may also be novel antifungal drug targets.

Finally, we have studied the importance of a 92-residue (residues M209-I300)
unstructured region in CnTps1, now called the intrinsically disordered domain (IDD)
(Supplemental Figures 1 and 13A). Interestingly, the IDD is well-conserved amongst Cryptococcal
species and closely related fungi in the Basidiomycota division (Supplemental Figure 13B). We
have also demonstrated, with the generation of a CnTps1 ΔIDD protein, that these 92 residues
are not required for the catalytic activity of CnTps1 *in vitro* (Supplemental Figure 13C). These
data are consistent with the residues neither interacting with CnTps1, either intra- or inter-
molecularly, nor being proximal to the catalytic region of CnTps1.

CnTps1 is required for the growth of *C. neoformans* at high growth temperatures, 37 °C
(27). In the future, we shall determine if the CnTps1 IDD is critical for the survival of *C. neoformans*
at high temperatures and in the presence of osmotic stress, both of which are encountered when
*C. neoformans* transitions from the environment to the human host. We posit, therefore, that the
main function of the CnTps1 IDD may involve its posttranslational modification, protein-protein
interactions and/or targeting CnTps1 to specific subcellular localizations in fungal cells. Future
work to determine the function of the CnTps1 IDD will aid in the development of the CnTps1 IDD
as a novel Basidiomycete-specific antifungal drug target.

In summary, we have solved the first cryo-EM structure of a Tps1 homo-tetramer, the

formation of which is potentially required for efficient trehalose biosynthesis and is therefore a

viable antifungal drug target (Supplemental Figure 17). We identified key residues that play a role

in tetramerization, as well as conserved substrate-binding residues. We have demonstrated that

the substrate-binding site of Tps1 is highly conserved and specific for UDPG. Finally, we identified

a novel 92-residue insertion in CnTps1 that may contribute to *C. neoformans* Tps1-mediated

tolerance of temperature and osmotic stress, both of which are encountered in the human host.

In conclusion, our work presented here will facilitate the development of novel antifungal drugs

which target multiple aspects of the trehalose biosynthesis pathway.

**Materials and Methods**

**Bacterial two-hybrid**

Protein interactions were analyzed by a bacterial two-hybrid analysis based on a set of vectors described by Daines and Silver (60). The genes for *C. neoformans Tps1* were cloned into the plasmid pSR658, which contains an N-terminal LexA DNA-binding domain. The resulting plasmid was cloned into *E. coli* strain SU101 to test for homo-oligomerization and expression of the protein fusions was induced with 1 mM IPTG. For this purpose, 5 mL of LB culture supplemented with antibiotics were inoculated and grown to 0.5 to 0.6 OD600 at 37 °C. The culture was mixed with permeabilization buffer (100 mM $Na_2HPO_4$, 20 mM KCl, 2 mM $MgSO_4$, 0.8 mg/mL CTAB, 0.4 mg/mL deoxycholic acid sodium salt, and 5.4 μL/mL of β-mercaptoethanol). Samples were incubated at 30 °C for 30 minutes and subsequently incubated with the prewarmed substrate solution (60 mM $Na_2HPO_4$, 40 mM $NaH_2PO_4$, 1 mg/mL ONPG, 2 μL/mL of β-mercaptoethanol). After 25 minutes, the stop solution (1 M $Na_2CO_3$) was added, and the absorbance of the culture supernatant was measured at 420 nM using a SpectraMax M5 plate reader (Molecular Devices).

**CnTps1 Protein expression and purification**

The full-length *TPS1* gene from *Cryptococcus neoformans* strain H99 was codon-optimized for expression in *E. coli* (Genscript) and subcloned using ligation-independent cloning into pMCSG7 (61). The construct was then transformed into *E. coli* OverExpress C41(DE3) chemically competent cells engineered for high protein expression. Supernatants from lysed cultures, induced with 0.5 mM IPTG, were loaded onto a nickel column (Ni-NTA, Qiagen) and washed in a buffer (50 mM Tris pH 8.0, 300 mM NaCl, 5 mM $MgCl_2$ and 5% glycerol) and eluted with increasing amounts of imidazole, which ranged from 40 mM to 400 mM imidazole, in this buffer. The fractions containing 6xHis-CnTps1 were then pooled, concentrated to 5 mL, and run over a S200 size exclusion column (HiLoad 26/600 Superdex 200pg, Cytiva) in pre-cooled buffer containing 20 mM Tris pH 8.0, 300 mM NaCl, 5% glycerol and 2 mM β-mercaptoethanol. Size exclusion fractions containing 6xHis-CnTps1 were pooled and concentrated to 1 mg/mL for downstream experimental applications such as cryo-electron microscopy and activity assays.

**CnTps1 Cryo-EM grid preparation**

6xHis-CnTps1 was purified as described above and concentrated in a buffer containing 20 mM Tris pH 8.0, 300 mM NaCl, 5% glycerol and 2 mM β-mercaptoethanol. For grids prepared

with *apo* CnTps1, 3 μL of 0.75 mg/mL 6xHis-CnTps1 was applied to glow-discharged carbon Quantifoil grids. After a 15 s incubation, the grids were blotted for 2 s to remove excess protein and rapidly plunged into liquid ethane (-182 °C) using a Leica EM GP2 (Leica Microsystems) operated at 95% humidity and 22 °C. For determination of the structure of the CnTps1-UDP-G6P complex, 0.5 mg/mL 6xHis-CnTps1 was incubated with 10 mM uridine di-phosphate (UDP, Sigma) and 10 mM glucose-6-phosphate (G6P, Sigma) for 18 hours at 4 °C. 3 μL of the CnTps1-UDP-G6P mixture was deposited onto glow-discharged UltrAuFoil grids. After a 15 s incubation, the grids were blotted for 2 s to remove excess protein and rapidly plunged into liquid ethane (-182 °C) using a Leica EM GP2 (Leica Microsystems) operated at 95% humidity and 22 °C. Grids were transferred to liquid nitrogen for storage until data collection.

**CnTps1 cryo-EM data collection**

After screening for high grid quality, using a Talos Arctica cryo-electron microscope (Thermo Fisher Scientific), a total of 2,520 micrographs from the cryo-EM grids containing *apo* CnTps1 were collected on a Titan Krios cryo-electron microscope (Thermo Fisher Scientific), at 300 kV equipped with a K3 detector (Gatan) at a nominal magnification of 105,000x and defocus values from -2.5 μm to -0.8 μm. The pixel size was 0.65 Å. The total dose was 62 e$^-$ Å$^{-2}$.

The cryo-EM grids containing CnTps1-UDP-G6P were also screened on a Talos Arctica cryo-electron microscope (Thermo Fisher Scientific). A total of 3,182 micrographs of CnTps1-UDP-G6P were collected on a Titan Krios cryo-electron microscope (Thermo Fisher Scientific), at 300 kV equipped with a K3 detector (Gatan) at a nominal magnification of 81,000x and defocus values from -2.5 μm to -0.8 μm. The pixel size was 1.08 Å. The total dose was 62 e$^-$ Å$^{-2}$.

**Cryo-EM data processing**

For *apo* CnTps1, 2,520 dose-fractionated image stacks were aligned using the drift correction routines implemented in RELION3.0 (62) and the contrast transfer function was determined on the motion corrected non-dose-weighted sum of frames using CTFFIND4.1 (63). CnTps1 particles were boxed out using template-free Auto-picking in RELION3.0. Two consecutive rounds of 2D classification were preformed to obtain a clean set of particles. The initial model was generated in RELION3.0, and 3 rounds of 3D classification were performed with resulting reconstructions showing non-uniform angular distributions. To improve the alignment, 907 micrographs were selected with the CTF max fitting resolution cutoff as 4 Å. A total of 456,020 4x binned clean particles were used for further processing. A new *ab initio* 3D reference was generated using 16,000 particles followed by 3D classification, and particles were classified into

5 classes. 207,081 particle images corresponding to best class were kept and subjected to 3D

refinement without symmetry resulting in a 5.3 Å resolution map. Particles were then re-centered

and re-extracted without binning and further refined using a shape mask to ~3.6 Å resolution.

After CTF refinement and Bayesian polishing in RELION3.0, the polished particles were imported

into cryoSPARC (64). Homogeneous refinement was performed without imposing symmetry

resulting in resolution of 3.6 Å.  After imposing D2 symmetry, the final global resolution of the

reconstruction improved to 3.3 Å. A summary of the data processing steps is shown in

Supplemental Figure 3.

For determination of the structure of CnTps1 bound to UDP-G6P, dose-fractionated

movies were aligned with MotionCor2 (65) and CTF estimation was performed using

CTFFIND4.1. Aligned micrographs were sorted and selected according to max fitting resolution

with cutoff as 4 Å with cryoSPARC Curate Exposures yielding 2,617 images. CnTps1 particles

were boxed out with template-free Blob picker. Approximately, 2.8 million particles were extracted

and subjected to 2D classification using a circular mask (180 Å radius) to focus the alignment on

one monomer.  A total of 2,079,376 clean particles were selected and subjected to *ab initio*

Reconstruction using 4 classes. A conformationally homogeneous class showed well resolved

features and accounted for ~850k particles which were kept and subjected to 3D refinement

(without imposing symmetry) resulting in a 3.1 Å resolution map. To improve the density of the

flexible loops around the substrate binding cave, 3D variability analysis was conducted using

cryoSPARC, and the particles were classified into 8 clusters. Further non-uniform refinement

using only particles assigned to cluster 1 resulted in the best map (3.5 Å resolution) that showed

well-resolved features on the flexible loop region. A summary of the data processing steps is

shown in Supplemental Figure 6.

**Cryo-EM model building**

Model building of the *apo* CnTps1 structure was initiated by using the CaTps2 N-terminal

domain crystal structure (PDB ID 5XDF) as the starting model. α-helix 2 had to be rebuilt. CnTps1

residues were mutated to match the correct sequence using COOT (66). Coordinates were then

fitted manually in Coot (66) followed by iterative refinement using Phenix (67) real space

refinement to improve the quality of the models. Following completion of the model of a single

protomer, multiple copies of the models were generated and docked into the map, using UCSF

Chimera 1.14, to form the homo-tetramer (68). The CnTps1-UDP-G6P model was generated

using the same workflow. The exception is that the initial model of the complex was the *apo*

CnTps1 model.

**Tps1 Activity Assay**

The catalytic activity of Tps1 was measured utilizing a continuous enzyme coupled assay

as previously reported (69). Briefly, 6xHis-CnTps1 protein was concentrated in a buffer containing

20 mM Tris pH 8.0, 300 mM NaCl, 5% glycerol and 2 mM β-mercaptoethanol. The assay was

carried out in a buffer containing 50 mM HEPES, pH 7.8, 100 mM KCl, 5 mM $MgCl_2$ and 2 mM

DTT. A final concentration of 3 μM, 1 mM and 1 mM were utilized for Tps1, uridine di-phosphate

glucose and glucose-6-phosphate, respectively. When reported, UDP-Galactose (UDP-Gal,

Sigma) was used in the assay. Activity assays were performed in clear, flat-bottomed 96-well

plates and the decrease in absorbance at 340 nm was recorded using a SpectraMax M5 plate

reader (Molecular Devices). The decrease in absorbance at 340 nm was analyzed for the 200 s

of the reaction, which corresponds to the initial rate of the reaction.

**Circular Dichroism spectroscopy**

Far-UV CD spectra of CnTps1 ΔIDD were recorded on an AVIV 435 CD

Spectrophotometer in a 1 mm sample cell. Measurements were taken from 200 to 260 nm with a

wavelength step of 1.0 nm and a 1 s averaging time. Each spectrum is the average of 3 scans.

CnTps1 ΔIDD was buffer exchanged into CD buffer (20 mM $NaH_2PO_4$ (pH 7.5), 300 mM NaF and

5% glycerol) and concentrated to a final concentration of 0.5 mg/mL.

**Accession Numbers**

Cryo-EM density maps of *apo* CnTps1 and CnTps1-UDP-G6P have been deposited in the Electron Microscopy Databank (EMDB) with accession codes EMD-29338 and EMD-29172, respectively. Atomic models of *apo* CnTps1 and CnTps1-UDP-G6P have been deposited in the RCSB Protein Data Bank (PDB) with accession codes PDB ID 8FO1 and PDB ID 8FHW, respectively.

**Acknowledgements**

This work was funded by grant 1P01AI104533 from the U.S. National Institutes of Health to R.G.B. and the NIH Tri-Institutional Molecular Mycology and Pathogenesis Training Grant T32AI052080 to E.W. Cryo-EM grid preparation and screening were carried out at the Genome Integrity and Structural Biology Laboratory at the National Institute of Environmental and Health Sciences in Research Triangle Park, NC. Cryo-EM data collection was performed at the Duke University Shared Materials Instrumentation Facility (SMIF), a member of the North Carolina Research Triangle Nanotechnology Network (RTNN), which is supported by the National Science Foundation as part of the National Nanotechnology Coordinated Infrastructure (NNCI) (Grant ECCS-2025064). We thank Dr. Mark Walters, Allen Hsu, and Megan Kopp for cryo-EM data collection support. We thank Dr. Brady Travis for cryo-EM data processing support. This study utilized the computational resources offered by Duke Research Computing.

**Author Contributions**

E.J.W. and R.G.B. designed the experiments and analyzed the biochemical data. E.J.W., Y.Z., A.B., and R.G.B. analyzed the structural data. E.J.W. generated purification constructs, purified proteins, determined their structures and performed biochemical characterizations. M.P., A.H., Y.Z., M.J.B. and A.B. provided cryo-EM consulting and experimental input. E.J.W., Y.Z. and R.G.B. wrote the manuscript with significant input from A.B. All authors have read and approved the manuscript.

**Conflict of Interest Statement**

The authors declare that they have no conflict of interest.

 **Figure Legends**

 **Figure 1. Trehalose-6-phosphate synthase (Tps1) from *Cryptococcus neoformans* self-**
 **associates. A)** Schematic of the canonical trehalose biosynthesis pathway in fungi. **B)** Bacterial
 two-hybrid results indicate a self-association of LexA-CnTps1$_{56-669}$ by a reduction in β-galactoside
 activity after induction of expression with 1 mM IPTG. Error bars represent standard error of
 triplicate biological replicates (N=3). Statistically significant differences were demonstrated with
 an unpaired Student's *t*-test (**** $P < 0.0001$). **C)** Anti-LexA Western blot confirms the expression
 of LexA-CnTps1$_{56-669}$.

 **Figure 2. Structure of the *apo Cryptococcus neoformans* Tps1 homo-tetramer.**
 **A)** Structure of the *apo C. neoformans* Tps1 homo-tetramer. The density map is shown in grey,
 overlayed with the model as a ribbon diagram. The protomers are colored and labelled in light
 blue, green, navy and magenta. The dimensions of the "front" view of the tetramer are labelled in
 Å. **B)** Structure of the tetramer viewed after a 90° rotation around the vertical axis. **C)** Structure of
 the tetramer viewed after a 90° rotation around the horizontal axis. **D)** Ribbon diagram of a single
 protomer of the *apo* CnTps1 cryo-EM structure with the N and C-termini labelled.

 **Figure 3. Structure of the *Cryptococcus neoformans* Tps1 homo-tetramer bound to UDP**
 **and G6P.**
 **A)** Structure of the *C. neoformans* Tps1 homo-tetramer bound to UDP and G6P. The density map
 is shown in grey, overlayed with the model as a ribbon diagram. The protomers are colored and
 labelled in light blue, green, navy and magenta. The dimensions of the "front" view of the tetramer
 are labelled in Å. Ligands UDP and G6P are shown with a space-filling representation. **B)**
 Structure of the tetramer bound to UDP and G6P viewed after a 90° rotation around the vertical
 axis. **C)** Structure of the tetramer bound to UDP and G6P viewed after a 90° rotation around the
 horizontal axis. **D)** Ribbon diagram of a single protomer of the CnTps1-UDP-G6P structure with
 the N and C-termini labelled and ligands in the substrate-binding pocket shown as atom-colored,
 space-filling molecules.

 **Figure 4. The binding of UDP and G6P induces a conformational change in CnTps1. A)**
 Overlay of the *apo* CnTps1 protomer (light blue) with the CnTps1 protomer bound to UDP and
 G6P (green). Density (grey mesh) is shown around ligands UDP and G6P in the substrate-binding
 pocket. Position of α-helix 2 is indicated with a black box. **B)** Overlay of the *apo* CnTps1 protomer

(light blue) with CnTps1-UDP-G6P (green) viewed after a 90° rotation around the horizontal axis.

Electron microscopy density is shown with UDP and G6P in the substrate-binding pocket. The

position of α-helix 2 is boxed, with the black line demonstrating the movement of approximately 7

Å.

**Figure 5. Substrate-binding residues of CnTps1. A)** Cryo-EM density for UDP and G6P, shown

as light grey mesh. UDP and G6P are shown as atom-colored sticks. **B)** View of the CnTps1

residues involved in binding UDP. UDP and residues of the C-terminal domain are shown as

atom-colored sticks. Hydrogen bonds are shown by dashes. Key atoms of the UDP molecule are

labelled. **C)** View of CnTps1 residue R453, which binds G6P. G6P and CnTps1 R453 are shown

as atom-colored sticks. Hydrogen bonds are shown by dashes. **D)** Relative activity of wild-type

CnTps1 and UDP-binding mutants. Error bars represent the standard error of three independent

measurements. **E)** Relative activity of wild-type CnTps1 and the G6P-binding mutant R453A.

Error bars represent the standard error of three independent measurements.

**Figure 6. The CnTps1 substrate-binding pocket is highly specific for UDPG. A)** Relative

activity of wild-type CnTps1 utilizing either 1 mM UDPG or 1 mM UDP-Gal or UDP as substrates.

Error bars represent the standard error of three independent measurements. **B)** Relative activity

of wild-type CnTps1 with 1 mM UDPG and increasing concentrations of UDP-Gal. Error bars

represent the standard error of three independent measurements. 1 mM G6P is present in all

experiments.

**References**

1. G. D. Brown *et al.*, Hidden killers: human fungal infections. *Science translational medicine* **4**, 165rv113 (2012).
2. F. Bongomin, S. Gago, R. O. Oladele, D. W. Denning, Global and Multi-National Prevalence of Fungal Diseases-Estimate Precision. *J Fungi (Basel)* **3** (2017).
3. Anonymous, Stop neglecting fungi. *Nat Microbiol* **2**, 17120 (2017).
4. A. Alanio, S. Delliere, S. Fodil, S. Bretagne, B. Megarbane, Prevalence of putative invasive pulmonary aspergillosis in critically ill patients with COVID-19. *Lancet Respir Med* **8**, e48-e49 (2020).
5. J. P. Gangneux, M. E. Bougnoux, E. Dannaoui, M. Cornet, J. R. Zahar, Invasive fungal diseases during COVID-19: We should be prepared. *J Mycol Med* **30**, 100971 (2020).
6. M. Hoenigl, Invasive Fungal Disease complicating COVID-19: when it rains it pours. *Clinical infectious diseases : an official publication of the Infectious Diseases Society of America* 10.1093/cid/ciaa1342 (2020).
7. P. Koehler *et al.*, COVID-19 associated pulmonary aspergillosis. *Mycoses* **63**, 528-534 (2020).
8. E. F. J. Meijer, A. S. M. Dofferhoff, O. Hoiting, J. B. Buil, J. F. Meis, Azole-Resistant COVID-19-Associated Pulmonary Aspergillosis in an Immunocompetent Host: A Case Report. *J Fungi (Basel)* **6** (2020).
9. D. Moser *et al.*, COVID-19 Impairs Immune Response to Candida albicans. *Front Immunol* **12**, 640644 (2021).
10. B. Posteraro *et al.*, Pan-Echinocandin-Resistant Candida glabrata Bloodstream Infection Complicating COVID-19: A Fatal Case Report. *J Fungi (Basel)* **6** (2020).
11. Y. Lee, E. Puumala, N. Robbins, L. E. Cowen, Antifungal Drug Resistance: Molecular Mechanisms in Candida albicans and Beyond. *Chem Rev* **121**, 3390-3411 (2021).
12. J. R. Perfect, The antifungal pipeline: a reality check. *Nature reviews. Drug discovery* **16**, 603-616 (2017).
13. J. R. Perfect, G. M. Cox, Drug resistance in Cryptococcus neoformans. *Drug resistance updates : reviews and commentaries in antimicrobial and anticancer chemotherapy* **2**, 259-269 (1999).
14. E. V. LeBlanc, E. J. Polvi, A. O. Veri, G. G. Prive, L. E. Cowen, Structure-guided approaches to targeting stress responses in human fungal pathogens. *The Journal of biological chemistry* **295**, 14458-14472 (2020).
15. J. R. Perfect, J. L. Tenor, Y. Miao, R. G. Brennan, Trehalose pathway as an antifungal target. *Virulence* 10.1080/21505594.2016.1195529, 1-7 (2016).
16. K. M. Pianalto, J. A. Alspaugh, New Horizons in Antifungal Therapy. *J Fungi (Basel)* **2** (2016).
17. A. Thammahong, S. Puttikamonkul, J. R. Perfect, R. G. Brennan, R. A. Cramer, Central Role of the Trehalose Biosynthesis Pathway in the Pathogenesis of Human Fungal Infections: Opportunities and Challenges for Therapeutic Development. *Microbiology and molecular biology reviews : MMBR* **81** (2017).
18. N. Benaroudj, D. H. Lee, A. L. Goldberg, Trehalose accumulation during cellular stress protects cells and cellular proteins from damage by oxygen radicals. *The Journal of biological chemistry* **276**, 24261-24267 (2001).
19. J. H. Crowe, L. M. Crowe, J. F. Carpenter, C. Aurell Wistrom, Stabilization of dry phospholipid bilayers and proteins by sugars. *The Biochemical journal* **242**, 1-10 (1987).
20. J. H. Crowe, L. M. Crowe, D. Chapman, Preservation of membranes in anhydrobiotic organisms: the role of trehalose. *Science* **223**, 701-703 (1984).

21.  J. H. Crowe, F. A. Hoekstra, L. M. Crowe, Anhydrobiosis. *Annu Rev Physiol* **54**, 579-599
(1992).
22.  O. Fernandez, L. Bethencourt, A. Quero, R. S. Sangwan, C. Clement, Trehalose and plant
stress responses: friend or foe? *Trends Plant Sci* **15**, 409-417 (2010).
23.  T. Hottiger, T. Boller, A. Wiemken, Rapid changes of heat and desiccation tolerance
correlated with changes of trehalose content in Saccharomyces cerevisiae cells subjected
to temperature shifts. *FEBS Lett* **220**, 113-115 (1987).
24.  T. Hottiger, C. De Virgilio, M. N. Hall, T. Boller, A. Wiemken, The role of trehalose
synthesis for the acquisition of thermotolerance in yeast. II. Physiological concentrations
of trehalose increase the thermal stability of proteins in vitro. *Eur J Biochem* **219**, 187-193
(1994).
25.  B. R. Steen *et al.*, Cryptococcus neoformans gene expression during experimental
cryptococcal meningitis. *Eukaryotic cell* **2**, 1336-1349 (2003).
26.  A. D. Elbein, Y. T. Pan, I. Pastuszak, D. Carroll, New insights on trehalose: a
multifunctional molecule. *Glycobiology* **13**, 17R-27R (2003).
27.  E. W. Petzold *et al.*, Characterization and regulation of the trehalose synthesis pathway
and its importance in the pathogenicity of Cryptococcus neoformans. *Infection and
immunity* **74**, 5877-5887 (2006).
28.  P. Ngamskulrungroj *et al.*, The trehalose synthesis pathway is an integral part of the
virulence composite for Cryptococcus gattii. *Infection and immunity* **77**, 4584-4596 (2009).
29.  M. Martinez-Esparza *et al.*, Role of trehalose-6P phosphatase (TPS2) in stress tolerance
and resistance to macrophage killing in Candida albicans. *Int J Med Microbiol* **299**, 453-
464 (2009).
30.  P. Van Dijck, L. De Rop, K. Szlufcik, E. Van Ael, J. M. Thevelein, Disruption of the Candida
albicans TPS2 gene encoding trehalose-6-phosphate phosphatase decreases infectivity
without affecting hypha formation. *Infection and immunity* **70**, 1772-1782 (2002).
31.  O. Zaragoza, M. A. Blazquez, C. Gancedo, Disruption of the Candida albicans TPS1 gene
encoding trehalose-6-phosphate synthase impairs formation of hyphae and decreases
infectivity. *Journal of bacteriology* **180**, 3809-3815 (1998).
32.  N. Al-Bader *et al.*, Role of trehalose biosynthesis in Aspergillus fumigatus development,
stress response, and virulence. *Infection and immunity* **78**, 3007-3018 (2010).
33.  S. Puttikamonkul *et al.*, Trehalose 6-phosphate phosphatase is required for cell wall
integrity and fungal virulence but not trehalose biosynthesis in the human fungal pathogen
Aspergillus fumigatus. *Molecular microbiology* **77**, 891-911 (2010).
34.  M. D. Asencion Diez *et al.*, The Production and Utilization of GDP-glucose in the
Biosynthesis of Trehalose 6-Phosphate by Streptomyces venezuelae. *The Journal of
biological chemistry* **292**, 945-954 (2017).
35.  J. C. Errey *et al.*, Mechanistic insight into enzymatic glycosyl transfer with retention of
configuration through analysis of glycomimetic inhibitors. *Angewandte Chemie* **49**, 1234-
1237 (2010).
36.  J. D. Farelli *et al.*, Structure of the trehalose-6-phosphate phosphatase from Brugia malayi
reveals key design principles for anthelmintic drugs. *PLoS pathogens* **10**, e1004245
(2014).
37.  R. P. Gibson, C. A. Tarling, S. Roberts, S. G. Withers, G. J. Davies, The donor subsite of
trehalose-6-phosphate synthase: binary complexes with UDP-glucose and UDP-2-deoxy-
2-fluoro-glucose at 2 A resolution. *The Journal of biological chemistry* **279**, 1950-1955
(2004).
38.  R. P. Gibson, J. P. Turkenburg, S. J. Charnock, R. Lloyd, G. J. Davies, Insights into
trehalose synthesis provided by the structure of the retaining glucosyltransferase OtsA.
*Chemistry & biology* **9**, 1337-1346 (2002).

39.    C. M. Harvey *et al.*, Structural Analysis of Binding Determinants of Salmonella
713            typhimurium Trehalose-6-phosphate Phosphatase Using Ground-State Complexes.
714            *Biochemistry* **59**, 3247-3257 (2020).

715    40.    Y. Miao *et al.*, Structural and In Vivo Studies on Trehalose-6-Phosphate Synthase from
716            Pathogenic Fungi Provide Insights into Its Catalytic Mechanism, Biological Necessity, and
717            Potential for Novel Antifungal Drug Design. *mBio* **8** (2017).

718    41.    Y. Miao *et al.*, Structures of trehalose-6-phosphate phosphatase from pathogenic fungi
719            reveal the mechanisms of substrate recognition and catalysis. *Proceedings of the National*
720            *Academy of Sciences of the United States of America* **113**, 7148-7153 (2016).

721    42.    S. Shan, H. Min, T. Liu, D. Jiang, Z. Rao, Structural insight into dephosphorylation by
722            trehalose 6-phosphate phosphatase (OtsB2) from Mycobacterium tuberculosis. *FASEB J*
723            **30**, 3989-3996 (2016).

724    43.    S. Wang *et al.*, Crystal structures of Magnaporthe oryzae trehalose-6-phosphate synthase
725            (MoTps1) suggest a model for catalytic process of Tps1. *The Biochemical journal* **476**,
726            3227-3240 (2019).

727    44.    M. Cross *et al.*, A suicide inhibitor of nematode trehalose-6-phosphate phosphatases. *Sci*
728            *Rep* **9**, 16165 (2019).

729    45.    S. Kapil, C. Petit, V. N. Drago, D. R. Ronning, S. J. Sucheck, Synthesis and in Vitro
730            Characterization of Trehalose-Based Inhibitors of Mycobacterial Trehalose 6-Phosphate
731            Phosphatases. *Chembiochem* **20**, 260-269 (2019).

732    46.    C. Kern *et al.*, Trehalose-6-phosphate synthase from the cat flea Ctenocephalides felis
733            and Drosophila melanogaster: gene identification, cloning, heterologous functional
734            expression and identification of inhibitors by high throughput screening. *Insect molecular*
735            *biology* **21**, 456-471 (2012).

736    47.    S. Klutts *et al.*, Purification, cloning, expression, and properties of mycobacterial trehalose-
737            phosphate phosphatase. *The Journal of biological chemistry* **278**, 2093-2100 (2003).

738    48.    C. Liu, D. Dunaway-Mariano, P. S. Mariano, Rational design of reversible inhibitors for
739            trehalose 6-phosphate phosphatases. *European journal of medicinal chemistry* **128**, 274-
740            286 (2017).

741    49.    Y. T. Pan, A. D. Elbein, Inhibition of the trehalose-P synthase of mycobacteria by various
742            antibiotics. *Archives of biochemistry and biophysics* **335**, 258-266 (1996).

743    50.    M. Dmitrova *et al.*, A new LexA-based genetic system for monitoring and analyzing protein
744            heterodimerization in Escherichia coli. *Mol Gen Genet* **257**, 205-212 (1998).

745    51.    L. L. Lairson, B. Henrissat, G. J. Davies, S. G. Withers, Glycosyltransferases: structures,
746            functions, and mechanisms. *Annu Rev Biochem* **77**, 521-555 (2008).

747    52.    L. Holm, P. Rosenstrom, Dali server: conservation mapping in 3D. *Nucleic Acids Res* **38**,
748            W545-549 (2010).

749    53.    R. A. Lee, M. Razaz, S. Hayward, The DynDom database of protein domain motions.
750            *Bioinformatics* **19**, 1290-1291 (2003).

751    54.    G. P. Poornam, A. Matsumoto, H. Ishida, S. Hayward, A method for the analysis of domain
752            movements in large biomolecular complexes. *Proteins* **76**, 201-212 (2009).

753    55.    S. S. Lee *et al.*, Mechanistic evidence for a front-side, SNi-type reaction in a retaining
754            glycosyltransferase. *Nature chemical biology* **7**, 631-638 (2011).

755    56.    E. Krissinel, K. Henrick, Inference of macromolecular assemblies from crystalline state.
756            *Journal of molecular biology* **372**, 774-797 (2007).

757    57.    W. Bell *et al.*, Composition and functional analysis of the Saccharomyces cerevisiae
758            trehalose synthase complex. *The Journal of biological chemistry* **273**, 33311-33319
759            (1998).

760    58.    A. Reinders *et al.*, Structural analysis of the subunits of the trehalose-6-phosphate
761            synthase/phosphatase complex in Saccharomyces cerevisiae and their function during
762            heat shock. *Molecular microbiology* **24**, 687-695 (1997).

763    59.    A. Thammahong, A. K. Caffrey-Card, S. Dhingra, J. J. Obar, R. A. Cramer, Aspergillus
764            fumigatus Trehalose-Regulatory Subunit Homolog Moonlights To Mediate Cell Wall
765            Homeostasis through Modulation of Chitin Synthase Activity. *mBio* **8** (2017).

766    60.    D. A. Daines, R. P. Silver, Evidence for multimerization of neu proteins involved in
767            polysialic acid synthesis in Escherichia coli K1 using improved LexA-based vectors.
768            *Journal of bacteriology* **182**, 5267-5270 (2000).

769    61.    W. H. Eschenfeldt, S. Lucy, C. S. Millard, A. Joachimiak, I. D. Mark, A family of LIC vectors
770            for high-throughput cloning and purification of proteins. *Methods in molecular biology* **498**,
771            105-115 (2009).

772    62.    S. H. Scheres, RELION: implementation of a Bayesian approach to cryo-EM structure
773            determination. *J Struct Biol* **180**, 519-530 (2012).

774    63.    A. Rohou, N. Grigorieff, CTFFIND4: Fast and accurate defocus estimation from electron
775            micrographs. *J Struct Biol* **192**, 216-221 (2015).

776    64.    A. Punjani, J. L. Rubinstein, D. J. Fleet, M. A. Brubaker, cryoSPARC: algorithms for rapid
777            unsupervised cryo-EM structure determination. *Nature methods* **14**, 290-296 (2017).

778    65.    S. Q. Zheng *et al.*, MotionCor2: anisotropic correction of beam-induced motion for
779            improved cryo-electron microscopy. *Nature methods* **14**, 331-332 (2017).

780    66.    P. Emsley, K. Cowtan, Coot: model-building tools for molecular graphics. *Acta
781            crystallographica. Section D, Biological crystallography* **60**, 2126-2132 (2004).

782    67.    P. D. Adams *et al.*, PHENIX: a comprehensive Python-based system for macromolecular
783            structure solution. *Acta crystallographica. Section D, Biological crystallography* **66**, 213-
784            221 (2010).

785    68.    E. F. Pettersen *et al.*, UCSF Chimera--a visualization system for exploratory research and
786            analysis. *J Comput Chem* **25**, 1605-1612 (2004).

787    69.    J. C. Errey *et al.*, Mechanistic Insight into Enzymatic Glycosyl Transfer with Retention of
788            Configuration through Analysis of Glycomimetic Inhibitors. *Angew. Chem.-Int. Edit.* **49**,
789            1234-1237 (2010).

790    70.    A. Idnurm, J. L. Reedy, J. C. Nussbaum, J. Heitman, Cryptococcus neoformans virulence
791            gene discovery through insertional mutagenesis. *Eukaryotic cell* **3**, 420-429 (2004).

792    71.    J. A. Fraser, R. L. Subaran, C. B. Nichols, J. Heitman, Recapitulation of the sexual cycle
793            of the primary fungal pathogen Cryptococcus neoformans var. gattii: implications for an
794            outbreak on Vancouver Island, Canada. *Eukaryotic cell* **2**, 1036-1045 (2003).

795    72.    D. L. Toffaletti, T. H. Rude, S. A. Johnston, D. T. Durack, J. R. Perfect, Gene transfer in
796            Cryptococcus neoformans by use of biolistic delivery of DNA. *Journal of bacteriology* **175**,
797            1405-1411 (1993).

 **Figure 1**

**Figure 2**

**Figure 3**

**Figure 4**

A

*apo* CnTps1
CnTps1-UDP-G6P

90°

B

**Figure 5**

**Figure 6**

A

B