1 **Structures of trehalose-6-phosphate synthase, Tps1, from the fungal pathogen**
2 ***Cryptococcus neoformans*: a target for novel antifungals**
3

4 Erica J. Washington[1*], Ye Zhou[2], Allen L. Hsu[3], Matthew Petrovich[3], Jennifer L. Tenor[4], Dena L.
5 Toffaletti[4], Ziqiang Guan[1], John R. Perfect[4], Mario J. Borgnia[3], Alberto Bartesaghi[1,2], Richard G.
6 Brennan[1*]
7

1. Department of Biochemistry, Duke University School of Medicine, Durham, North Carolina,
27710 USA
2. Department of Computer Science, Duke University, Durham, North Carolina, 27708 USA
3. Genome Integrity and Structural Biology Laboratory, National Institute of Environmental
Health Sciences, National Institutes of Health, Department of Health and Human Services,
Research Triangle Park, NC 27709 USA
4. Division of Infectious Diseases, Department of Medicine, Duke University School of
Medicine, Durham, North Carolina, 27710 USA
16

17 [*]To whom correspondence should be addressed: richard.brennan@duke.edu and
18 erica.washington1@duke.edu

19
20
21
22
23 **Running Title**
24 Cryo-EM structures of CnTps1
25
26 **Keywords**
27 Cryptococcus, cryo-electron microscopy, trehalose, fungal pathogens, Basidiomycetes

Abstract

Invasive fungal diseases are a major threat to human health, resulting in more than 1.5 million annual deaths worldwide. The arsenal of antifungal therapeutics remains limited and is in dire need of novel drugs that target additional biosynthetic pathways that are absent from humans. One such pathway involves the biosynthesis of trehalose. Trehalose is a disaccharide that is required for pathogenic fungi to survive in their human hosts. In the first step of trehalose biosynthesis, trehalose-6-phosphate synthase (Tps1) converts UDP-glucose and glucose-6-phosphate to trehalose-6-phosphate. Here, we report the structures of full-length *Cryptococcus neoformans* Tps1 (CnTps1) in unliganded form and in complex with uridine diphosphate and glucose-6-phosphate. Comparison of these two structures reveals significant movement towards the catalytic pocket by the N-terminus upon ligand binding and identifies residues required for substrate-binding, as well as residues that stabilize the tetramer. Intriguingly, an intrinsically disordered domain (IDD), which is conserved amongst Cryptococcal species and closely related Basidiomycetes, extends from each subunit of the tetramer into the "solvent" but is not visible in density maps. We determined that the IDD is not required for *C. neoformans* Tps1-dependent thermotolerance and osmotic stress survival. Studies with UDP-galactose highlight the exquisite substrate specificity of CnTps1. *In toto*, these studies expand our knowledge of trehalose biosynthesis in *Cryptococcus* and highlight the potential of developing antifungal therapeutics that disrupt the synthesis of this disaccharide or the formation of a functional tetramer and the use of cryo-EM in the structural characterization of CnTps1-ligand/drug complexes.

Significance Statement

Fungal infections are responsible for over a million deaths worldwide each year. Biosynthesis of
a disaccharide, trehalose, is required for multiple pathogenic fungi to transition from the
environment to the human host. Enzymes in the trehalose biosynthesis pathway are absent in
humans and, therefore, are potentially significant targets for novel antifungal therapeutics. One
enzyme in the trehalose biosynthesis is trehalose-6-phosphate synthase (Tps1). Here, we
describe the cryo-electron microscopy structures of the CnTps1 homo-tetramer in the
unliganded form and in complex with a substrate and a product. These structures and
subsequent biochemical analysis reveal key details of substrate-binding residues and substrate
specificity. These structures should facilitate structure-guided design of inhibitors against
CnTps1.

Introduction

Invasive fungal diseases (IFDs) caused by pathogenic fungi such as *Cryptococcus*, *Candida* and *Aspergillus* are a major threat to human health, resulting in more than one and a half million deaths worldwide each year (1, 2). These mortality rates are surprisingly high, given the fact that fungal infections are usually associated with superficial infections of skin and nails. However, IFDs result in increased mortality in immunocompromised populations, including HIV-infected patients and patients receiving immunosuppressive therapies (1-3). Additionally, the severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2; COVID-19) pandemic has resulted in a significant population of COVID-19 positive patients with increased susceptibility to IFDs, as has been demonstrated for pulmonary aspergillus and invasive candidiasis infections (4-10).

Unfortunately, the current arsenal of antifungal drugs, consisting of the polyenes, azoles and echinocandins, is insufficient to manage the mortality caused by IFDs due to significant off-target effects, the rapid emergence of resistance to antifungal therapeutics and the emergence of intrinsically drug resistant fungal pathogens, such as *Candida auris* and *Candida glabrata* (11-14) Therefore, in addition to the development of new formulations of current antifungal drugs, there is a critical need for the development of new classes of broad-spectrum antifungal drugs that are fast-acting and safe. The identification of targets that are not present in the human host is critical to the development of antifungal drugs with low toxicity. This is often difficult, as both fungi and humans are eukaryotes. However, pathways involved in the fungal stress response have been implicated recently as key targets, including the trehalose biosynthesis pathway (14-17).

Trehalose is a non-reducing disaccharide composed of two glucose molecules linked by an α,α-1,1-glycosidic bond. Fungal cells synthesize trehalose to protect their proteins and membranes from stresses encountered during the infection process (18-25). The canonical trehalose biosynthesis machinery in pathogenic fungi commences with trehalose-6-phosphate synthase, Tps1, a glucosyltransferase that converts uridine diphosphate glucose (UDPG) and glucose-6-phosphate (G6P) to trehalose-6-phosphate (T6P). Subsequently, trehalose-6-phosphate phosphatase (Tps2), removes the phosphate group to generate the final product of trehalose (Figure 1A). In *Candida albicans* and *Aspergillus fumigatus* there is an additional protein in the pathway, Tps3, with no known enzymatic function. The machinery to synthesize trehalose is found in plants, insects, fungi and bacteria, but not in humans (26). For this reason, an antifungal therapeutic that targets the trehalose biosynthesis pathway could well result in a drug with minimal off-target effects, leading to greatly reduced toxicity in patients. Indeed,

disruption of the *TPS1* gene in *Cryptococcus neoformans* results in fungi that are avirulent in
mice and rabbits and, interestingly, disruption of *TPS2* was followed by the accumulation of the
toxic intermediate trehalose 6-phosphate, causing fungal cell death (27). Additionally, in *C.*
neoformans trehalose acts as a stress protectant and is required for growth at high
temperatures (27). Similarly, disruption of the *TPS1* and *TPS2* genes in *Cryptococcus gattii*
results in decreased infectivity in mice (28). Similar phenotypes indicating the importance of
Tps1 and Tps2 to cell survival and sometimes virulence are also seen in trehalose biosynthesis
mutants in *C. albicans* (29-31) and *A. fumigatus* (32, 33). Therefore, we hypothesize that
targeting the trehalose biosynthesis pathway will result in the development of a novel broader-
spectrum antifungal drug, which would be both highly effective and, importantly, safe for use in
immunocompromised patients.

The trehalose biosynthesis pathway and its contribution to the response of fungi to the
increasingly warmer global temperatures is also important to understand. Climate change plays
a critical role in the emergence of fungal diseases by reducing the threshold between
environmental temperatures and mammalian basal temperatures (34-38). When fungi,
especially those possessing virulence factors, acquire thermotolerance they also acquire the
possibility of becoming more pathogenic. Importantly, the trehalose biosynthesis pathway in *C.*
neoformans plays a role in thermotolerance (39). CnTps1 is required for *C. neoformans* to grow
at higher temperatures (28). Similarly, the ability of trehalose to accumulate in *C. neoformans*
grown at 37 °C is dependent on CnTps1 (28). Therefore, understanding the function of CnTps1
and its structure should help us understand how environmental fungi evolve thermotolerance to
become pathogenic.

Pertinent to the development of an antifungal therapeutic which targets the trehalose
biosynthesis pathway is an intimate knowledge and understanding of the structures of enzymes
and proteins involved in this critical pathway. Several structures of Tps1 and Tps2 from
nematodes, bacteria and pathogenic fungi have been determined using x-ray crystallography
(40-49). Subsequently, compounds that inhibit trehalose biosynthesis enzymes functions by
targeting their catalytic pockets have been investigated but not yet advanced for clinical use (45,
50-55). As a key starting point to develop inhibitors of CnTps1, we describe structures of this
669-residue (74 kDa per protomer) enzyme in both its unliganded (substrate-free) form and in
complex with UDP and G6P using single particle cryo-electron microscopy (cryo-EM). Unlike
Tps1 from *Candida albicans* (46), CnTps1 has three major insertions, 55 residues at the N-
terminus, 63 residues at the C-terminus and a 92-residue intrinsically disordered domain (IDD)
inserted within the core of the enzyme, all of unknown function. Residues located in the catalytic

pocket and the IDD were assayed for their contribution to the activity of this enzyme. Substrate
specificity was investigated further and revealed the exquisite ability of CnTps1 to discriminate
at the level of a single interaction. Hence, this work lays the groundwork for subsequent CnTps1
structure-based drug design.

Results

Trehalose biosynthesis protein CnTps1 self-associates in bacterial two-hybrid assays

The trehalose biosynthesis pathway is a two-step process. Tps1, trehalose-6-phosphate synthase, converts uridine diphosphate glucose (UDPG) and glucose-6-phosphate (G6P) into trehalose-6-phosphate (T6P). The phosphate group is subsequently removed from T6P by trehalose-6-phosphate phosphatase (Tps2) to yield trehalose (Figure 1A) (17). Previously the structures of Tps1/OtsA from *Candida albicans* (46), *Aspergillus fumigatus* (46), *E. coli (41, 43, 44, 49)*, *Streptomyces venezuelae* (40) and *Magnaporthe oryzae* (49) were determined using x-ray crystallography. Additionally, the structure of the catalytically inactive Tps1-like N-terminal domain of *C. albicans* Tps2 was determined (47).

Inspection of the Tps1 crystal structures from *Candida albicans* (46), *Aspergillus fumigatus* (46), *E. coli* (43, 44), *Streptomyces venezuelae* (40) and *Magnaporthe oryzae* (49) and the catalytically inactive Tps1-like N-terminal domain of *C. albicans* Tps2 (47) revealed that Tps1 most likely exists as a homo-tetramer. To determine if subunits of the *Cryptococcus neoformans* var. *grubii* strain H99 Tps1, hereafter referred to as CnTps1, self-associate, a bacterial two-hybrid system based on the LexA DNA-binding repressor was established (56). Expression of full-length LexA-CnTps1 in the bacterial two-hybrid system was not detected. However, LexA-CnTps1$_{56-669}$, in which the unstructured 55 N-terminal residues of CnTps1 were deleted, does self-associate in the bacterial two-hybrid assay, as evidenced by a reduction in β-galactosidase activity after protein expression was induced with 1 mM IPTG (Figure 1B). An anti-LexA Western blot shows the expression of LexA-CnTps1$_{56-669}$ at the appropriate molecular weight (Figure 1C). These data indicate that CnTps1 can self-associate at physiological concentrations and support the hypothesis that CnTps1 forms homo-oligomers.

The structure of the unliganded (substrate-free) CnTps1 homo-tetramer

CnTps1 is approximately 200 residues larger than other Tps1 orthologues, due to the addition of a 55-residue N-terminal extension, a 63-residue C-terminal extension and an internal insertion from residues M209 to I300 (Supplemental Figure 1). Each extension is predicted to be structurally disordered. In order to determine the structure of CnTps1, the full-length (74 kDa, 669- residue per protomer) CnTps1 protein was expressed with an N-terminal hexahistidine tag and purified from *E. coli* via Ni^{2+}-NTA affinity chromatography (Supplemental Figure 2A). Following confirmation that 6xHis-CnTps1 was enzymatically active (Supplemental Figure 2B), the protein sample was subjected to single particle cryo-electron microscopy analysis. 2,520 micrographs were collected from grids prepared with 6xHis-CnTps1 (Supplemental Figure 3).

No additional ligands were added to the protein sample. 2D classification, performed on 907

micrographs with resolution higher than 4.0 Å, revealed tetrameric CnTps1 particles

(Supplemental Figure 3). 3D classification and initial refinement steps were performed without

imposing symmetry (Supplemental Figure 3), resulting in a 5.3 Å resolution map. Additional

refinement and polishing steps, including adding D2 symmetry, resulted in an unliganded

CnTps1 map at 3.3 Å global resolution, as estimated by the "gold standard" Fourier shell

correlation (FSC) = 0.143 criteria (Data Table 1 and Supplemental Figure 4A,B). Four protomers

can be readily observed in the unliganded CnTps1 cryo-EM map (Supplemental Figure 4C). The

final map shows features of well-resolved side chains of many amino acid residues

(Supplemental Figure 4D). The atomic model for unliganded CnTps1, which was built based on

the cryo-EM reconstruction, supports the conclusion that unliganded CnTps1 forms a homo-

tetramer (Figure 2).

The four CnTps1 protomers, designated as protomers 1 – 4, assemble into the

tetrameric structure with dimensions of 100.0 x 110.3 x 107.6 Å (Figure 2A-C). The four

protomers interact via two α-helical domains that are found in the central region of the homo-

tetramer (Figure 2A). The unbound substrate-binding pockets, are open and outward-facing,

and therefore accessible to ligands and substrates (Figure 2A,B). The "bottom-facing" view of

the homo-tetramer shows a staggered appearance of the top and bottom pair of protomers

(Figure 2C).

Each CnTps1 protomer is comprised of N and C-terminal lobes, containing residues 1-

378 and 379-669, respectively. Both lobes are spanned by an α-helix, formed by residues 572-

603 (Figure 2D). Each domain contains a modified Rossman fold that is characteristic of the

retaining glycosyltransferase family (46, 57) (Figure 2D) and similar to the published CaTps1

and CaTps1-like N-terminal domain of CaTps2 structures (Supplemental Figure 5) (40, 47, 49).

A common feature that is observed in these substrate-free structures is a lack of density in

portions of the N-terminus, suggestive of conformational flexibility in the absence of ligands.

Indeed, the CnTps1 cryo-EM map shows clear density for secondary structures in the C-

terminus, whilst the density was poor in several parts of the N-terminal domain, including that for

residues 1-58, 68-100, 127, 180, 210-301, 354-357 and 364-368 (Figure 2D). Not unexpectedly,

the N-terminus extension, the C-terminus and the internal insertion, residues 209-300, are not

visible in this structure suggesting multiple accessible conformations or intrinsic disorder.

The structure of the CnTps1-UDP-G6P complex

In order to identify residues in the catalytic pocket of CnTps1 as well as to visualize any
conformational changes necessary for substrate binding and catalysis, we determined the cryo-
EM structure of the *C. neoformans* H99 Tps1 in complex with the substrate, G6P, and a
product, UDP, to 3.1 Å global resolution (Figure 3A-D). 3,182 movies were collected from grids
prepared with 6xHis-CnTps1 and an excess of UDP and G6P (Supplemental Figure 6). The 2D
classification revealed a tetrameric structure like those observed for unliganded CnTps1
(Supplemental Figure 6). Final refinement was performed with D2 symmetry imposed
(Supplemental Figure 6). The addition of G6P and UDP resulted in a higher global resolution of
the CnTps1 structure when compared to the unliganded structure (Data Table 2). The substrate-
binding pocket is more ordered (Supplemental Figure 7A,B). Four protomers can be detected in
the CnTps1-UDP-G6P cryo-EM map (Supplemental Figure 7C) and the final map shows
features of well-resolved side chains (Supplemental Figure 7D).

As observed in the unliganded structure, the N and C-termini and the internal insertion
remain unstructured in the presence of ligands. The structure of CnTps1 bound to UDP and
G6P is similar to that of the published CaTps1-UDP-G6P complex and the N-terminus of
CnTps2 with root mean square deviations (rmsds) of 1.2 Å for 450 corresponding C_α atoms and
1.5 Å for 468 corresponding C_α atoms, respectively (46, 47, 58). Views of the CnTps1-UDP-G6P
tetramer show the central binding region of UDP and G6P inside each protomer (Figure 3A-D).
The CnTps1-UDP-G6P complex UDP bound to the C-terminal domain and the G6P adjacent to
the N-terminal domain (Figure 3D). The binding sites of the ligands are in excellent agreement
with the published structure of CaTps1-UDP-G6P and CaTps1-UDPG (46).

Ligand-induced conformational changes in *C. neoformans* Tps1

Significant movement and conformational changes of the core of the N-terminal domain
are observed due to ligand binding to CnTps1 (Figure 4A,B). Structural alignments of the
individual N-terminal domain (residues 56 – 378) and C-terminal domain (residues 379 – 603) of
the unliganded Tps1 protein onto the corresponding domains of UDP-G6P bound CnTps1
reveal rmsds of 1.1 and 0.4 Å, respectively. Superposition of the complete structure of
unliganded CnTps1 onto CnTps1-UDP-G6P reveals an rmsd of 1.4 Å for 433 corresponding C_α
atoms. These rmsd values are a result of ligand-induced conformational "closing" of CnTps1,
with the movement occurring primarily in the N-terminus. Thus, unliganded CnTps1 protomers
are in a more "open" conformation as compared to the ligand-bound subunits. Systematic
analysis of the domain movements using the program DynDom (59, 60) shows that the large
movements of the N-terminal G6P-binding lobe centered around α-helix 2 (residues 112 to 125),

which is located on the surface of the N-terminal domain, result in a rotation of approximately
20° (Supplemental Figure 8). This rotation causes an inward, approximately 7 Å movement of α-
helix 2 (Figure 4B). The linker that facilitates this movement includes residues 113 through 145.
This rotation results in a 29% closure of the N and C-terminal lobes around the substrate-
binding pocket. Interestingly, the local resolution proximal to CnTps1 α-helix 2 is the lowest in
the unliganded structure, which indicates flexibility and a propensity for movement, consistent
with the results of the DynDom analysis. Since the CnTps1-UDP-G6P complex contains both a
product (UDP) and a substrate (G6P) as ligands, the completely closed form of the protein is
likely not visualized in this structure. Regardless, the movement of this domain would facilitate
the S_N-i catalytic mechanism proposed for Tps1 proteins in the GT-B fold retaining
glycosyltransferase family, including *C. albicans* Tps1 and *E. coli* OtsA (41, 46, 57, 61). Indeed,
superposition of UDP onto the UDPG from the CaTps1-UDPG complex (PDB ID 5HUT) reveals
that the G6P substrate bound to CnTps1 is proximal to and properly aligned with the glucose of
the UDPG to allow the ready formation of the 1,1 glycosidic bond, required for the formation of
T6P (Supplemental Figure 9).
248
CnTps1 substrate-binding residues
We were able to detect strong density for the bound ligand and substrate-binding
residues of CnTps1 (Figure 5A). The UDP molecule is bound to the C-terminal portion of each
protomer (Figure 5B). The orientation of the uracil base and CnTps1 is mediated by a hydrogen
bond between the exocyclic O4 and the V492 amide NH group. The O2 and O3 hydroxyl groups
of the ribose ring of UDP form hydrogen bonds with the side chain of E522. The orientation of
the phosphate groups of UDP in CnTps1 is mediated by several interactions. One hydrogen
bond is made between the α-phosphate group and the backbone amide group of L518. In
CaTps1 (46) there is a conserved Arg-Lys pair (residues R280 and K285) that interacts with the
phosphates. We were unable to detect side chain density for the equivalent CnTps1 Arg,
indicating movement of the substrate-binding pocket, potentially ready to release the product
UDP. There is, however, a hydrogen bond between K420 and the β-phosphate. Based on an
overlay with the CaTps1-UDPG structure, we predicted that the glucose moiety in the native
substrate, UDPG, would be stabilized by CnTps1 residue D514 and positioned to deprotonate
G6P (Figure 5B and Supplemental Figure 10A) (46). The G6P molecule is located adjacent to
the N-terminal lobe in each protomer. A key interaction stabilizing G6P is residue R453, which
interacts with the G6P phosphate group (Figure 5C).

CnTps1 substrate-binding residues K420, D514, L518, E522 and R453 are conserved
amongst Tps1 proteins from *C. neoformans*, *C. albicans*, *A. fumigatus*, *M. oryzae*, *S.*
venezuelae and *E. coli* (Supplemental Figure 1 and Supplemental Figure 10B). In *C. albicans*
Tps1 these residues have been demonstrated to be required for hyphal development and
thermotolerance (46). Site-directed mutagenesis was used to generate variants of CnTps1 in
which potential substrate-binding residues were mutated as follows: E522A, K420A, D514A and
R453A. All CnTps1 variants eluted similarly to wild-type CnTps1 on the size exclusion column,
indicating that they are properly folded (Supplemental Figure 11). Tps1 coupled enzyme assays
revealed that mutation of the residues that interact with UDP/UDPG resulted in severe loss-of-
function (Figure 5D). Mutation of R453 to an alanine, which is the single identified G6P-
interacting residue, resulted in greatly reduced CnTps1 activity as well (Figure 5E).
277

Interactions within the CnTps1 homo-tetramer
The interactions between several side chains in the interfaces between CnTps1
protomers explain the formation and stabilization of the CnTps1 homo-tetramer. The two key
interfaces responsible for formation of the CnTps1 homo-tetramer are referred to as interface #1
and interface #2 (Supplemental Figure 12A). PDBePISA (62), revealed the buried surface areas
for each interface are similar with 1062.9 Å buried in interface #1 and 986.2 Å buried in interface
#2. The calculated free energy (ΔG) of the two interfaces is -10.7 kcal/mol and 0.1 kcal/mol,
respectively, which suggests the former interface is a critical component of oligomerization,
whilst the latter might allow the formation of a Tps1-Tps2 hetero-oligomer.
The key interactions in interface #1 are the hydrogen bonds between N467 and E468, as
well as R403 and E481 (Supplemental Figure 12B,C). Interface #2 also has hydrogen bonds
that help stabilize the homo-tetramer, including hydrogen bonds between the E314 backbone
oxygen and R513 side chain (Supplemental Figure 12D). Ionic interactions between R513 and
E314 and between R322 and E538 also contribute to the formation of interface #2. None of the
interface residues directly contact UDP or G6P, indicating that any effect mutation of these
residues might have on activity would be allosteric. Our cryo-EM results confirm that the homo-
tetrameric form of CnTps1 accurately represents the tetrameric structure in solution.
295

CnTps1 contains a conserved intrinsically disordered domain
CnTps1 contains three unstructured insertions (Supplemental Figure 13A). These
regions have prevented the determination of its structure by x-ray crystallography, as these
regions are the main differences between CnTps1 and other crystallized Tps1 orthologues. The

most prominent unstructured region in CnTps1 is an internal insertion of 92 residues (residues

M209-I300), referred to hereafter as the Intrinsically Disordered Domain or IDD (Supplemental

Figure 13A). Interestingly, the CnTps1 IDD is found only in the Tps1 proteins in fungi within the

fungal division, Basidiomycota (Supplemental Figure 13B). More specifically, the CnTps1 IDD is

highly conserved within Cryptococcal species and closely related Basidiomycetes, with *C.*

neoformans, *C. gattii* and *C. deneoformans* containing IDDs with greater than 92% identity

(Supplemental Figure 13A,B). The CnTps1 IDD is not present in the Ascomycota division or the

Mucoromycota phylum (Supplemental Figure 13B).

As anticipated, density for the CnTps1 IDD was not detected in the structures of

unliganded CnTps1 or CnTps1-UDP-G6P, consistent with the idea that the IDD is

conformationally flexible. However, based on the CnTps1 structures, we can determine that the

IDD exits and returns to the structured regions in adjacent parts of the N-terminus of CnTps1.

To determine if the IDD is required for CnTps1 activity *in vitro*, we designed and purified a

chimeric protein in which the 92-residue CnTps1 IDD was replaced with the six residues

(GNKKKN) of *C. albicans* Tps1 that connect α-helix 5 to β-strand 4 (Supplemental Figure 14).

We refer to this protein as CnTps1 ΔIDD.

To determine the effect that the CnTps1 IDD has on CnTps1 function, we purified

CnTps1 ΔIDD and confirmed that it was reduced in size compared to wild-type CnTps1, as

determined by size exclusion chromatography (Supplemental Figure 11). Notably, deletion of

the CnTps1 IDD did not affect formation of the tetramer (Supplemental Figure 11). Circular

dichroism experiments also confirmed that CnTps1 ΔIDD is properly folded (Supplemental

Figure 15). Coupled enzyme activity assays with CnTps1 ΔIDD were performed and revealed

that the CnTps1 IDD is not required for enzymatic activity of CnTps1 *in vitro* (Supplemental

Figure 13C). Indeed, wild-type CnTps1 and the CnTps1 ΔIDD variant have similar catalytic

efficiencies (Supplemental Table 1).

CnTps1 is required for the growth of *C. neoformans* at 37 °C (27). To determine if the

CnTps1 IDD is required for the CnTps1-mediated thermotolerance, strains of *C. neoformans*

H99 lacking the CnTps1 IDD were generated. Wild-type *C. neoformans*, the *tps1Δ* strain and

the *TPS1^{IDDΔ}* strain were able to grow similarly at 30 °C (Supplemental Figure 16). At 37 °C, the

C. neoformans TPS1^{IDDΔ} strain had wild-type levels of growth, compared to the temperature-

sensitive phenotype observed for *C. neoformans tps1Δ* (Supplemental Figure 16). Furthermore,

only the *C. neoformans tps1Δ* strain could not grow at 39 °C (Supplemental Figure 16). The

complementation strain of *C. neoformans TPS1^{IDDΔ}* strain expressing the wild-type *TPS1* grew

like wild-type *C. neoformans* at 30 °C, 37 °C and 39 °C (Supplemental Figure 16). The ability of

the strain lacking the CnTps1 IDD to grow at the higher temperatures indicates that the IDD is

not required for growth of *C. neoformans* at temperatures similar to the temperatures observed

in human hosts.

337   *C. neoformans tps1Δ* and *C. neoformans TPS1*[IDDΔ] were unable to grow at 37 °C or 39

°C on YPD plates containing 1 M sorbitol (Supplemental Figure 16). However, wild-type *C.*

neoformans, *C. neoformans TPS1*[IDDΔ] and the complementation strain grew in the presence of 1

M sorbitol at 30 °C, 37 °C, and 39 °C (Supplemental Figure 16). These data indicate that *C.*

neoformans tps1Δ is sensitive to osmotic stress, but the CnTps1-mediated tolerance of osmotic

stress does not require the IDD of CnTps1.

Specificity of CnTps1 catalytic pocket

345   To assess the substrate specificity of the CnTps1 catalytic pocket, we tested the

catalytic activity of CnTps1 in the presence of UDP-Galactose (UDP-Gal). The hydroxyl groups

on the 4[th] carbon are oriented differently in glucose and galactose. In the CaTps1-UDPG

structure, all of the extracyclic hydroxyl groups are easily detected, revealing that the O4

hydroxyl hydrogen forms a hydrogen bond with the peptide backbone of CaTps1 N382 (46).

Similarly, the peptide backbone of CnTps1-UDP-G6P residue N517 would interact with the O4

hydroxyl of the glucose moiety of UDPG (Supplemental Figure 17A). Modelling UDP-Galactose

into the catalytic pocket reveals the O4 hydroxyl points away from N517, preventing the

formation of this hydrogen bond (Supplemental Figure 17B). The loss of this single interaction

has a significant effect on the activity of CnTps1 with UDP-Gal as demonstrated by a reduction

of only 4% of this substrate and a reduction in catalytic efficiency, compared to CnTps1 with

UDPG as a substrate (~50% use of this substrate) (Figure 6A and Supplemental Table 1),

suggesting that a stereochemically proper positioning of the O4, mediated by its interaction with

N517, is a critical component of CnTps1 substrate-assisted catalysis.

359   To test the effect of UDP-Gal on CnTps1 activity further, we performed a competition

assay. The combination of equal molar ratios of UDPG and UDP-Gal to CnTps1 resulted in a

12% reduction in activity, compared to only adding UDPG to the assay (Figure 6B).

Interestingly, the addition of 10-fold excess UDP-Gal compared to UDPG resulted in a reduction

of CnTps1 activity by approximately 30% (Figure 6B), indicating that UDP-Gal is capable of

binding CnTps1 but is a significantly weaker binder and a poor competitive inhibitor. Further,

we confirmed that UDP-Gal is a poor substrate of CnTps1 using an orthogonal mass

spectrometry-based assay to detect the formation of a disaccharide-phosphate after incubation

of CnTps1 with either UDPG or UDP-Gal and the loss of these substrates over 4 minutes

(Figure 6C,D). Incubation of CnTps1 with UDPG resulted in the formation of more disaccharide-

phosphate (T6P) compared to UDP-Gal (Figure 6C). Consistent with these results, more UDPG

was expended in this assay compared to UDP-Gal (Figure 6D). In summary, these results

confirm the strong preference of CnTps1 for UDPG.

Discussion

In order to develop the trehalose biosynthesis pathway as an antifungal drug target, we must improve our understanding of the structure and function of trehalose biosynthesis proteins. Here we report the use of cryo-EM to determine the structures of unliganded CnTps1 and CnTps1 bound to UDP and G6P.

We confirmed that CnTps1 forms a tetramer in solution. Our current hypothesis is that trehalose biosynthesis proteins also form hetero-tetrameric complexes. Interactions amongst Tps proteins (Tps1, Tps2, Tps3 and Tsl1) have been demonstrated in *S. cerevisiae* (63, 64). The function of these complexes may be to sequester the highly cytotoxic trehalose 6-phosphate (T6P) from the cytoplasm, to target trehalose biosynthesis to a proper subcellular localization (65) or to allow for quick and efficient production of trehalose when fungal pathogens are exposed to environmental stress. Future work will include confirmation of the formation of heterocomplexes between CnTps1 and Tps2 both *in vitro* and *in vivo* and determination of the structure of the heterocomplex using cryo-electron microscopy.

We also described herein a conformational change of CnTps1 upon binding of UDP and G6P. Comparison of the unliganded CnTps1 and CnTps1-UDP-G6P structures reveals an inward movement of the N-terminus of CnTps1 when bound to ligand. Most of the movement occurs in α-helix 2 and surrounding residues. The closure of the ligand-bound CnTps1 protomer is consistent with substrate-assisted catalysis, the proposed enzymatic mechanism of Tps1 enzymes (46). Consistent with previous reports of the predicted catalytic activity of fungal Tps1 proteins, we also propose that UDPG binds first, followed by G6P. The proximity of the substrates in the substrate-binding pocket of CnTps1 is demonstrated in Supplemental Figure 9. However, we now can see aspects of the catalytic mechanism occurring in the context of the CnTps1 homo-tetramer.

The more ordered CnTps1-UDP-G6P substrate-binding pocket allows us to observe specific interactions between the ligands and CnTps1 substrate-binding residues. The identification of substrate-binding residues, K420, D514, E522 and R453, provides insights into the mechanism of the Tps1 glycosyltransferase as well as demonstrates the key roles of these residues in substrate specificity (Figure 5A-C). Mutation of any of these residues is sufficient to inhibit CnTps1 catalytic activity (Figure 5D,E). Additionally, our determination of substrate-binding variants which lead to a loss of activity of CnTps1 provides the framework for future investigation of a fully closed substrate-trapped cryo-EM structure of the CnTps1 homo-tetramer. This structure will provide fuller information regarding the CnTps1 catalytic mechanism by allowing visualization of CnTps1 bound to its substrates, UDPG and G6P.

The substrate-binding residues identified in the catalytic pocket of Tps1 are conserved
amongst fungal pathogens, including *C. albicans* (Supplemental Figure 1 and Supplemental
Figure 10B), supporting the proposition that a broader-spectrum antifungal therapeutic can be
developed by targeting the catalytic pocket of Tps1. Additionally, we have been able to
demonstrate that the CnTps1 substrate-binding domain is highly specific as underscored by its
significantly reduced ability to utilize UDP-Galactose as a substrate, an epimer of UDPG (Figure
6 and Supplemental Figure 17).

Although the key interfaces of the CnTps1 homo-tetramer are not as highly resolved as
the substrate-binding pocket, specific residues that may play a role in the formation of the
homo-tetramers were identified (Supplemental Figure 12). It remains to be determined whether
oligomerization of CnTps1 is required for catalytic activity. If so, in addition to targeting the
substrate-binding pocket of CnTps1, the interfaces critical to the formation of the CnTps1
tetramer may also be novel antifungal drug targets.

Finally, we have studied the importance of a 92-residue (residues M209-I300)
unstructured region in CnTps1, now called the intrinsically disordered domain (IDD)
(Supplemental Figures 1 and 13A). Intriguingly, the IDD is well-conserved amongst
Cryptococcal species and closely related fungi in the Basidiomycota division (Supplemental
Figure 13B). We have demonstrated, with the generation of a CnTps1 ΔIDD protein, that these
92 residues are not required for the catalytic activity of CnTps1 *in vitro* (Supplemental Figure
13C). These data are consistent with the residues not interacting with CnTps1, neither intra nor
intermolecularly, or being proximal to the catalytic region of CnTps1. Indeed, the IDD does not
affect the *in vitro* catalytic efficiency of CnTps1 significantly (Supplemental Table 1).

CnTps1 is required for the growth of *C. neoformans* at 37 °C (27). Surprisingly we
demonstrated that the CnTps1 IDD is not critical for the survival of *C. neoformans* at higher
temperatures and in the presence of osmotic stress, both of which are encountered when *C.*
neoformans transitions from the environment to the human host (Supplemental Figure 16). We
posit, therefore, that a function of this highly conserved IDD may involve its posttranslational
modification, ability to interact with other cellular proteins, targeting CnTps1 to specific
subcellular localizations in fungal cells or self-association into a condensate (66, 67). Future
work to determine the function of the CnTps1 IDD will aid in the development of the CnTps1 IDD
as a novel Basidiomycete-specific antifungal drug target.

CnTps1 has the GT-B fold of the retaining glycosyltransferase family which is
characterized largely by a large substrate-binding cleft between two lobes (68). GT-B fold,
retaining glycosyltransferases function in a metal-independent manner (68). Several structures

of proteins in the GT-B glycosyltransferase family including sucrose synthase from *N. europaea*

have been determined (69). While these enzymes have some characteristics that are similar,

there are also key differences that make them unique and the context in which they function is

also critical. Therefore, to capture the entire wealth of characteristics amongst

glycosyltransferases, it is important for new structures to be determined. For example, a

comparison of *N. europaea* sucrose synthase and CnTps1 reveals that the key lobes are

similar, with an rmsd of 1.4 A (Supplemental Figure 18). However, the sucrose synthase protein

has two extra domains (SSN-1, SSN-2) that are not found in CnTps1 whereas CnTps1 contains

the IDD, which is not found in sucrose synthase (69). Another surprising difference between

these two glycosyltransferases is the lack of the strict conservation of the triplet of conserved

catalytic residues amongst members of the GT-B retaining glycosyltransferase family (69).

Sucrose synthase and other glycosyltransferases contain a conserved Arg-Lys-Glu triad, in

CnTps1 the Glu is replaced by the smaller acidic amino acid residue Asp (69).

In summary, we have determined the first cryo-EM structures of a fungal Tps1 homo-

tetramer in its unliganded and ligand-bound conformation. In the future, we shall determine

whether a tetramer is potentially required for efficient trehalose biosynthesis and thus, disrupting

the oligomerization state of Tps1 might make a viable antifungal drug target (Supplemental

Figure 19). We identified key residues that play a role in tetramerization, as well as conserved

substrate-binding residues. We have demonstrated that the substrate-binding site of Tps1 is

conserved and specific for UDPG. Finally, we identified a 92-residue IDD in CnTps1 that does

not contribute to *C. neoformans* Tps1-mediated tolerance of temperature and osmotic stress,

both of which are encountered in the human host. However, the conservation of the CnTps1

IDD points strongly towards a necessary function in *Cryptococcus* species as well as

Basidiomycetes. In conclusion, our work presented here should facilitate the development of

novel antifungal drugs which target multiple aspects of the initial step in the trehalose

biosynthesis pathway.

Materials and Methods

Bacterial two-hybrid

Protein interactions were analyzed by a bacterial two-hybrid analysis based on a set of vectors described by Daines and Silver (70). The genes for *C. neoformans Tps1* were cloned into the plasmid pSR658, which contains an N-terminal LexA DNA-binding domain. The resulting plasmid was cloned into *E. coli* strain SU101 to test for homo-oligomerization and expression of the protein fusions was induced with 1 mM IPTG. For this purpose, 5 mL of LB culture supplemented with antibiotics were inoculated and grown to 0.5 to 0.6 OD600 at 37 °C. The culture was mixed with permeabilization buffer (100 mM Na_2HPO_4, 20 mM KCl, 2 mM $MgSO_4$, 0.8 mg/mL CTAB, 0.4 mg/mL deoxycholic acid sodium salt, and 5.4 µL/mL of β-mercaptoethanol). Samples were incubated at 30 °C for 30 minutes and subsequently incubated with the prewarmed substrate solution (60 mM Na_2HPO_4, 40 mM NaH_2PO_4, 1 mg/mL ONPG, 2 µL/mL of β-mercaptoethanol). After 25 minutes, the stop solution (1 M Na_2CO_3) was added, and the absorbance of the culture supernatant was measured at 420 nM using a SpectraMax M5 plate reader (Molecular Devices).

CnTps1 Protein expression and purification

The full-length *TPS1* gene from *Cryptococcus neoformans* strain H99 was codon-optimized for expression in *E. coli* (Genscript) and subcloned using ligation-independent cloning into pMCSG7 (71). The construct was transformed into *E. coli* OverExpress C41(DE3) chemically competent cells engineered for high protein expression. Supernatants from lysed cultures, induced with 0.5 mM IPTG, were loaded onto a nickel column (Ni-NTA, Qiagen) and washed in a buffer (50 mM Tris pH 8.0, 300 mM NaCl, 5 mM $MgCl_2$ and 5% glycerol) and eluted with increasing amounts of imidazole, which ranged from 40 mM to 400 mM imidazole, in this buffer. The fractions containing 6xHis-CnTps1 were then pooled, concentrated to 5 mL, and run over a S200 size exclusion column (HiLoad 26/600 Superdex 200pg, Cytiva) in pre-cooled buffer containing 20 mM Tris pH 8.0, 300 mM NaCl, 5% glycerol and 2 mM β-mercaptoethanol. Size exclusion fractions containing 6xHis-CnTps1 were pooled and concentrated to 1 mg/mL for downstream experimental applications such as cryo-electron microscopy and activity assays.

CnTps1 Cryo-EM grid preparation

6xHis-CnTps1 was purified as described above and concentrated in a buffer containing 20 mM Tris pH 8.0, 300 mM NaCl, 5% glycerol and 2 mM β-mercaptoethanol. For grids

prepared with unliganded CnTps1, 3 µL of 0.75 mg/mL 6xHis-CnTps1 was applied to glow-

discharged carbon Quantifoil grids. After a 15 s incubation, the grids were blotted for 2 s to

remove excess protein and rapidly plunged into liquid ethane (-182 °C) using a Leica EM GP2

(Leica Microsystems) operated at 95% humidity and 22 °C. For determination of the structure of

the CnTps1-UDP-G6P complex, 0.5 mg/mL 6xHis-CnTps1 was incubated with 10 mM uridine

diphosphate (UDP, Sigma) and 10 mM glucose-6-phosphate (G6P, Sigma) for 18 hours at 4 °C.

3 µL of the CnTps1-UDP-G6P mixture was deposited onto glow-discharged UltrAuFoil grids.

After a 15 s incubation, the grids were blotted for 2 s to remove excess protein and rapidly

plunged into liquid ethane (-182 °C) using a Leica EM GP2 (Leica Microsystems) operated at

95% humidity and 22 °C. Grids were transferred to liquid nitrogen for storage until data

collection.

CnTps1 cryo-EM data collection

After screening for high grid quality, using a Talos Arctica cryo-electron microscope

(Thermo Fisher Scientific), a total of 2,520 micrographs from the cryo-EM grids containing

unliganded CnTps1 were collected on a Titan Krios cryo-electron microscope (Thermo Fisher

Scientific), at 300 kV equipped with a K3 detector (Gatan) at a nominal magnification of

105,000x and defocus values from -2.5 µm to -0.8 µm. The pixel size was 0.65 Å. The total

dose was 62 $e^- \cdot Å^{-2}$.

The cryo-EM grids containing CnTps1-UDP-G6P were also screened on a Talos Arctica

cryo-electron microscope (Thermo Fisher Scientific). A total of 3,182 micrographs of CnTps1-

UDP-G6P were collected on a Titan Krios cryo-electron microscope (Thermo Fisher Scientific),

at 300 kV equipped with a K3 detector (Gatan) at a nominal magnification of 81,000x and

defocus values from -2.5 µm to -0.8 µm. The pixel size was 1.08 Å. The total dose was 62

$e^- \cdot Å^{-2}$.

Cryo-EM data processing

For unliganded CnTps1, 2,520 dose-fractionated image stacks were aligned using the

drift correction routines implemented in RELION3.0 (72) and the contrast transfer function was

determined on the motion corrected non-dose-weighted sum of frames using CTFFIND4.1 (73).

CnTps1 particles were boxed out using template-free Auto-picking in RELION3.0. Two

consecutive rounds of 2D classification were preformed to obtain a clean set of particles. The

initial model was generated in RELION3.0, and 3 rounds of 3D classification were performed

with resulting reconstructions showing non-uniform angular distributions. To improve the

alignment, 907 micrographs were selected with the CTF max fitting resolution cutoff as 4 Å. A

total of 456,020 4x binned clean particles were used for further processing. A new *ab initio* 3D

reference was generated using 16,000 particles followed by 3D classification, and particles were

classified into 5 classes. 207,081 particle images corresponding to best class were kept and

subjected to 3D refinement without symmetry resulting in a 5.3 Å resolution map. Particles were

then re-centered and re-extracted without binning and further refined using a shape mask to

~3.6 Å resolution. After CTF refinement and Bayesian polishing in RELION3.0, the polished

particles were imported into cryoSPARC (74). Homogeneous refinement was performed without

imposing symmetry resulting in resolution of 3.6 Å. After imposing D2 symmetry, the final global

resolution of the reconstruction improved to 3.3 Å. A summary of the data processing steps is

shown in Supplemental Figure 3.

For determination of the structure of CnTps1 bound to UDP-G6P, dose-fractionated

movies were aligned with MotionCor2 (75) and CTF estimation was performed using

CTFFIND4.1. Aligned micrographs were sorted and selected according to max fitting resolution

with a cutoff of 4 Å with cryoSPARC Curate Exposures yielding 2,617 images. CnTps1 particles

were boxed out with template-free Blob picker. Approximately, 2.8 million particles were

extracted and subjected to 2D classification using a circular mask (180 Å radius) to focus the

alignment on one protomer. A total of 2,079,376 clean particles were selected and subjected to

ab initio Reconstruction using 4 classes. A conformationally homogeneous class showed well

resolved features and accounted for ~850k particles which were kept and subjected to 3D

refinement (without imposing symmetry) resulting in a 3.1 Å resolution map. To improve the

density of the flexible loops around the substrate binding pocket, 3D variability analysis was

conducted using cryoSPARC, and the particles were classified into 8 clusters. Further non-

uniform refinement using only particles assigned to cluster 1 resulted in the best map (3.5 Å

resolution) that showed well-resolved features on the flexible loop region surrounding the

catalytic pocket. A summary of the data processing steps is shown in Supplemental Figure 6.

Cryo-EM model building

Model building of the unliganded CnTps1 structure was initiated by using the CaTps2 N-

terminal domain crystal structure (PDB ID 5XDF) as the starting model. α-helix 2 had to be

rebuilt. CnTps1 residues were mutated to match the correct sequence using COOT (76).

Coordinates were then fitted manually in Coot (76) followed by iterative refinement using Phenix

(77) real space refinement to improve the quality of the models. Following completion of the

model of a single protomer, multiple copies of the models were generated and docked into the

map, using UCSF Chimera 1.14, to form the homo-tetramer (78). The CnTps1-UDP-G6P model

was generated using the same workflow. The exception is that the initial model of the complex

was the unliganded CnTps1 model.

Tps1 Activity Assay

The catalytic activity of Tps1 was measured utilizing a continuous enzyme coupled

assay as previously reported (79). Briefly, 6xHis-CnTps1 protein was concentrated in a buffer

containing 20 mM Tris pH 8.0, 300 mM NaCl, 5% glycerol and 2 mM β-mercaptoethanol. The

assay was carried out in a buffer containing 50 mM HEPES, pH 7.8, 100 mM KCl, 5 mM $MgCl_2$

and 2 mM DTT. A final concentration of 3 μM, 1 mM and 1 mM were utilized for Tps1, uridine

diphosphate glucose and glucose-6-phosphate, respectively. When reported, UDP-Galactose

(UDP-Gal, Sigma) was used in the assay. Activity assays were performed in clear, flat-bottomed

96-well plates and the decrease in absorbance at 340 nm was recorded using a SpectraMax M5

plate reader (Molecular Devices). The decrease in absorbance at 340 nm was analyzed for the

200 s of the reaction, which corresponds to the initial rate of the reaction.

Circular Dichroism spectroscopy

Far-UV CD spectra of CnTps1 ΔIDD were recorded on an AVIV 435 CD

Spectrophotometer in a 1 mm sample cell. Measurements were taken from 200 to 260 nm

with a wavelength step of 1.0 nm and a 1 s averaging time. Each spectrum is the average of 3

scans. CnTps1 ΔIDD was buffer exchanged into CD buffer (20 mM NaH_2PO_4 (pH 7.5),

300 mM NaF and 5% glycerol) and concentrated to a final concentration of 0.5 mg/mL.

Generation of CnTps1 strains

Strains and primers used in this study are listed in Supplemental Table 2. The following

strains were constructed in this study: *tps1Δ*, *TPS1^{IDDΔ}*, and *TPS1^{IDDΔ}::TPS1.* To construct the

tps1Δ deletion mutant strain, three PCRs were prepared: the 5' flanking region (AD131/AD132),

the nourseothricin (NAT) drug selection cassette (AD133/AD134) amplified from plasmid, pIA3

(80), and the 3' flanking region (AD135/AD136). These PCR fragments were gel extracted,

purified using the Qiaquick gel extraction kit (Qiagen), and cloned into linearized pUC19 by In-

Fusion cloning (Takara Bio). To construct the *TPS1^{IDDΔ}* strain, four PCRs were prepared: the 5'

flank region (AD3710/AD3711), the neomycin (NEO) drug selection cassette amplified from

plasmid, pJAF1 (81) (AD3712/AD3713), 5' flank region closest to *TPS1* and the *TPS1* gene up

to the IDD (AD3714/AD3715), and part of the *TPS1* gene downstream of the IDD

(AD3716/AD3717). Primers AD3715 and AD3716 have the Ca IDD nucleotide sequence codon

optimized for *Cryptococcus* for replacement of the CnTPS1 IDD. These fragments were cloned

into linearized pUC19 using the Hi-Fi Assembly Cloning Kit (New England Biolabs). The *tps1*Δ

mutant was confirmed by PCR and the TPS1*^{IDDΔ}* strain was confirmed by PCRs and sequencing

to confirm the replacement of the CnTps1IDD with the CaTps1IDD. Lastly, a wild-type copy of

TPS1 was introduced into the *TPS1*^{IDDΔ} strain. Three PCRs were performed to amplify: the 5'

flanking region (AD3748/AD3749), the nourseothricin drug selection cassette (NAT)

(AD3750/AD3751), and additional 5' flank region and part of the *TPS1* gene (AD3572/AD3573).

These fragments were cloned into linearized pUC19 using the Hi-Fi Assembly Cloning Kit (New

England Biolabs). The plasmids carrying the desired inserts were transformed into strains by

biolistic transformation as previously reported (82). Strains were confirmed by PCR using

primers listed in Supplemental Table 2 and sequencing.

LC/MS/MS

Samples for LC/MS/MS were prepared by incubating by 2 μM CnTps1 with 1 mM G6P

and either 1 mM UDPG or 1 mM UDP-Gal for 4 minutes in a reaction buffer consisting of 20 mM

Tris pH 8.0, 300 mM NaCl, 5% glycerol and 2 mM β-mercaptoethanol. The reactions were

stopped by the addition of methanol. Samples were centrifuged at 10,000 rpm at room

temperature for 2 minutes. Subsequently, the supernatant was collected and submitted for

analysis.

Reverse-phase liquid chromatography-electrospray ionization/tandem mass

spectrometry (LC-ESI/MS/MS) was performed using a Shimadzu LC system (comprising a

solvent degasser, two LC-10A pumps and a SCL-10A system controller) coupled to a high-

resolution TripleTOF5600 mass spectrometer (AB Sciex, Framingham, MA). LC was operated

at a flow rate of 200 μl/min with a linear gradient as follows: 100% of mobile phase A was held

isocratically for 2 min and then linearly increased to 100% mobile phase B over 5 min and held

at 100% B for 2 min. Mobile phase A was a mixture of water/acetonitrile (98/2, v/v) containing

0.1% acetic acid. Mobile phase B was a mixture of water/acetonitrile (10/90, v/v) containing

0.1% acetic acid. A Zorbax SB-C8 reversed-phase column (5 μm, 2.1 x 50 mm) was obtained

from Agilent (Palo Alto, CA). The LC eluent was introduced into the ESI source of the mass

spectrometer. Instrument settings for negative ion ESI/MS and MS/MS analysis of lipid species

were as follows: Ion spray voltage (IS) = -4500 V; Curtain gas (CUR) = 20 psi; Ion source gas 1

(GS1) = 20 psi; De-clustering potential (DP) = -55 V; Focusing potential (FP) = -150 V. Data

acquisition and analysis were performed using the Analyst TF1.5 software (AB Sciex,
Framingham, MA).

Data, Materials, and Software Availability

Cryo-EM density maps of unliganded CnTps1 and CnTps1-UDP-G6P have been deposited in the Electron Microscopy Databank (EMDB) (https://www.ebi.ac.uk/emdb) with accession codes EMD-29338 and EMD-29172, respectively (83, 84). Atomic models of unliganded CnTps1 and CnTps1-UDP-G6P have been deposited in the RCSB Protein Data Bank (PDB) (https://www.rcsb.org) with accession codes PDB ID 8FO1 and PDB ID 8FHW, respectively (85, 86). The enzymatic data in this study have been deposited in the Standards for Reporting Enzymology Data (STRENDA) database with the SRN HSMHV7 and VZVQFC. All software used in this study is publicly available. The paper does not report any original code. All other data are included in the manuscript and/or supplemental data.

Acknowledgements

This work was funded by grant 1P01AI104533 from the U.S. National Institutes of Health to R.G.B. and the NIH Tri-Institutional Molecular Mycology and Pathogenesis Training Grant T32AI052080 to E.W. Cryo-EM grid preparation and screening were carried out at the Genome Integrity and Structural Biology Laboratory at the National Institute of Environmental and Health Sciences in Research Triangle Park, NC. Cryo-EM data collection was performed at the Duke University Shared Materials Instrumentation Facility (SMIF), a member of the North Carolina Research Triangle Nanotechnology Network (RTNN), which is supported by the National Science Foundation as part of the National Nanotechnology Coordinated Infrastructure (NNCI) (Grant ECCS-2025064). We thank Dr. Mark Walters, Allen Hsu, and Megan Kopp for cryo-EM data collection support. We thank Dr. Brady Travis for cryo-EM data processing support. This study utilized the computational resources offered by Duke Research Computing. The open access option is selected.

Author Contributions

E.J.W. and R.G.B. designed the experiments and analyzed the biochemical data. E.J.W., Y.Z., A.B., and R.G.B. analyzed the structural data. E.J.W. generated purification constructs, purified proteins, determined their structures and performed biochemical characterizations. J.L.T., D.L.T., and J.R.P. designed and performed cellular experiments. Y.G. performed mass spectrometry experiments. M.P., A.H., Y.Z., M.J.B. and A.B. provided cryo-EM consulting and experimental input. E.J.W., Y.Z. and R.G.B. wrote the manuscript with significant input from A.B. All authors have read and approved the manuscript.

671 **Competing Interest**

672 The authors declare that they have no competing interests.

 Figure Legends

Figure 1. Trehalose-6-phosphate synthase (Tps1) from *Cryptococcus neoformans* self-
associates. A) Schematic of the canonical trehalose biosynthesis pathway in fungi. **B)** Bacterial
two-hybrid results indicate a self-association of LexA-CnTps1$_{56-669}$ by a reduction in β-
galactoside activity after induction of expression with 1 mM IPTG. Error bars represent standard
error of triplicate biological replicates (N=3). Statistically significant differences were
demonstrated with an unpaired Student's *t*-test (**** $P < 0.0001$). **C)** Anti-LexA Western blot
confirms the expression of LexA-CnTps1$_{56-669}$.

Figure 2. Structure of the unliganded *Cryptococcus neoformans* Tps1 homo-tetramer.
A) Structure of the unliganded *C. neoformans* Tps1 homo-tetramer. The density map is shown
in grey, overlayed with the model as a ribbon diagram. The protomers are colored and labelled
in light blue, green, navy and magenta. The dimensions of the "front" view of the tetramer are
labelled in Å. **B)** Structure of the tetramer viewed after a 90° rotation around the vertical axis. **C)**
Structure of the tetramer viewed after a 90° rotation around the horizontal axis. **D)** Ribbon
diagram of a single protomer of the unliganded CnTps1 cryo-EM structure with the N and C-
termini labelled. **E)** Density for unliganded CnTps1 is shown in grey.

Figure 3. Structure of the *Cryptococcus neoformans* Tps1 homo-tetramer bound to UDP
and G6P.
A) Structure of the *C. neoformans* Tps1 homo-tetramer bound to UDP and G6P. The density
map is shown in grey, overlayed with the model as a ribbon diagram. The protomers are colored
and labelled in light blue, green, navy and magenta. The dimensions of the "front" view of the
tetramer are labelled in Å. Ligands UDP and G6P are shown with a space-filling representation.
B) Structure of the tetramer bound to UDP and G6P viewed after a 90° rotation around the
vertical axis. **C)** Structure of the tetramer bound to UDP and G6P viewed after a 90° rotation
around the horizontal axis. **D)** Ribbon diagram of a single protomer of the CnTps1-UDP-G6P
structure with the N and C-termini labelled and ligands in the substrate-binding pocket shown as
atom-colored, space-filling molecules. **E)** Density for CnTps1-UDP-G6P is shown in grey.

Figure 4. The binding of UDP and G6P induces a conformational change in CnTps1. A)
Overlay of the unliganded CnTps1 protomer (light blue) with the CnTps1 protomer bound to
UDP and G6P (green). Density (grey mesh) is shown around ligands UDP and G6P in the

catalytic pocket. Position of α-helix 2 is indicated with a black box. **B)** Overlay of the unliganded

CnTps1 protomer (light blue) with CnTps1-UDP-G6P (green) viewed after a 90° rotation around

the horizontal axis. Electron microscopy density is shown for UDP and G6P in the substrate-

binding pocket. The position of α-helix 2 is boxed, with the black line demonstrating the

movement of approximately 7 Å.

Figure 5. Substrate-binding residues of CnTps1. A) Cryo-EM density for UDP and G6P,

shown as light grey mesh. UDP and G6P are shown as atom-colored sticks. **B)** View of the

CnTps1 residues involved in binding UDP. UDP and residues of the C-terminal domain are

shown as atom-colored sticks. Hydrogen bonds are shown by dashes. Key atoms of the UDP

molecule are labelled. **C)** View of CnTps1 residue R453, which binds G6P. G6P and CnTps1

R453 are shown as atom-colored sticks. Hydrogen bonds are shown by dashes. **D)** Relative

activity of wild-type CnTps1 and UDP-binding mutants. Error bars represent the standard error

of three independent measurements. **E)** Relative activity of wild-type CnTps1 and the G6P-

binding mutant R453A. Error bars represent the standard error of three independent

measurements.

Figure 6. The CnTps1 substrate-binding pocket is highly specific for UDPG. A) Relative

activity of wild-type CnTps1 utilizing either 1 mM UDPG or 1 mM UDP-Gal or UDP as

substrates. Error bars represent the standard error of three independent measurements. **B)**

Relative activity of wild-type CnTps1 with 1 mM UDPG and increasing concentrations of UDP-

Gal. Error bars represent the standard error of three independent measurements. 1 mM G6P is

present in all experiments.   **C)** Shown are the extracted ion chromatograms of [M-H]$^-$ at

m/z 421.08 for the formation of a disaccharide-phosphate (D-P). **D)** Shown are the extracted

ion chromatograms of [M-H]$^-$ at *m/z* 565.05 for UDPG and UDP-Gal at zero and four minutes

post catalysis.

Table Legends

Data Table 1. Cryo-EM data collection and refinement statistics for unliganded CnTps1.

Data collection and refinement statistics regarding the cryo-EM structure of unliganded CnTps1.

FSC, Fourier shell correlation.

Data Table 2. Cryo-EM data collection and refinement statistics for CnTps1-UDP-G6P.

Data collection and refinement statistics regarding the cryo-EM structure of CnTps1 in complex

with UDP and G6P. FSC, Fourier shell correlation.

744
745

References

1. G. D. Brown *et al.*, Hidden killers: human fungal infections. *Science translational medicine* **4**, 165rv113 (2012).
2. F. Bongomin, S. Gago, R. O. Oladele, D. W. Denning, Global and Multi-National Prevalence of Fungal Diseases-Estimate Precision. *J Fungi (Basel)* **3** (2017).
3. Anonymous, Stop neglecting fungi. *Nat Microbiol* **2**, 17120 (2017).
4. A. Alanio, S. Delliere, S. Fodil, S. Bretagne, B. Megarbane, Prevalence of putative invasive pulmonary aspergillosis in critically ill patients with COVID-19. *Lancet Respir Med* **8**, e48-e49 (2020).
5. J. P. Gangneux, M. E. Bougnoux, E. Dannaoui, M. Cornet, J. R. Zahar, Invasive fungal diseases during COVID-19: We should be prepared. *J Mycol Med* **30**, 100971 (2020).
6. M. Hoenigl, Invasive Fungal Disease complicating COVID-19: when it rains it pours. *Clinical infectious diseases : an official publication of the Infectious Diseases Society of America* 10.1093/cid/ciaa1342 (2020).
7. P. Koehler *et al.*, COVID-19 associated pulmonary aspergillosis. *Mycoses* **63**, 528-534 (2020).
8. E. F. J. Meijer, A. S. M. Dofferhoff, O. Hoiting, J. B. Buil, J. F. Meis, Azole-Resistant COVID-19-Associated Pulmonary Aspergillosis in an Immunocompetent Host: A Case Report. *J Fungi (Basel)* **6** (2020).
9. D. Moser *et al.*, COVID-19 Impairs Immune Response to Candida albicans. *Front Immunol* **12**, 640644 (2021).
10. B. Posteraro *et al.*, Pan-Echinocandin-Resistant Candida glabrata Bloodstream Infection Complicating COVID-19: A Fatal Case Report. *J Fungi (Basel)* **6** (2020).
11. S. E. Jacobs *et al.*, Candida auris Pan-Drug-Resistant to Four Classes of Antifungal Agents. *Antimicrobial agents and chemotherapy* **66**, e0005322 (2022).
12. Y. Lee, E. Puumala, N. Robbins, L. E. Cowen, Antifungal Drug Resistance: Molecular Mechanisms in Candida albicans and Beyond. *Chem Rev* **121**, 3390-3411 (2021).
13. J. R. Perfect, G. M. Cox, Drug resistance in Cryptococcus neoformans. *Drug resistance updates : reviews and commentaries in antimicrobial and anticancer chemotherapy* **2**, 259-269 (1999).
14. J. R. Perfect, J. L. Tenor, Y. Miao, R. G. Brennan, Trehalose pathway as an antifungal target. *Virulence* 10.1080/21505594.2016.1195529, 1-7 (2016).
15. E. V. LeBlanc, E. J. Polvi, A. O. Veri, G. G. Prive, L. E. Cowen, Structure-guided approaches to targeting stress responses in human fungal pathogens. *The Journal of biological chemistry* **295**, 14458-14472 (2020).
16. K. M. Pianalto, J. A. Alspaugh, New Horizons in Antifungal Therapy. *J Fungi (Basel)* **2** (2016).
17. A. Thammahong, S. Puttikamonkul, J. R. Perfect, R. G. Brennan, R. A. Cramer, Central Role of the Trehalose Biosynthesis Pathway in the Pathogenesis of Human Fungal Infections: Opportunities and Challenges for Therapeutic Development. *Microbiology and molecular biology reviews : MMBR* **81** (2017).
18. N. Benaroudj, D. H. Lee, A. L. Goldberg, Trehalose accumulation during cellular stress protects cells and cellular proteins from damage by oxygen radicals. *The Journal of biological chemistry* **276**, 24261-24267 (2001).
19. J. H. Crowe, L. M. Crowe, J. F. Carpenter, C. Aurell Wistrom, Stabilization of dry phospholipid bilayers and proteins by sugars. *The Biochemical journal* **242**, 1-10 (1987).
20. J. H. Crowe, L. M. Crowe, D. Chapman, Preservation of membranes in anhydrobiotic organisms: the role of trehalose. *Science* **223**, 701-703 (1984).

795   21.   J. H. Crowe, F. A. Hoekstra, L. M. Crowe, Anhydrobiosis. *Annu Rev Physiol* **54**, 579-599
796   (1992).
797   22.   O. Fernandez, L. Bethencourt, A. Quero, R. S. Sangwan, C. Clement, Trehalose and
798   plant stress responses: friend or foe? *Trends Plant Sci* **15**, 409-417 (2010).
799   23.   T. Hottiger, T. Boller, A. Wiemken, Rapid changes of heat and desiccation tolerance
800   correlated with changes of trehalose content in Saccharomyces cerevisiae cells
801   subjected to temperature shifts. *FEBS Lett* **220**, 113-115 (1987).
802   24.   T. Hottiger, C. De Virgilio, M. N. Hall, T. Boller, A. Wiemken, The role of trehalose
803   synthesis for the acquisition of thermotolerance in yeast. II. Physiological concentrations
804   of trehalose increase the thermal stability of proteins in vitro. *Eur J Biochem* **219**, 187-
805   193 (1994).
806   25.   B. R. Steen *et al.*, Cryptococcus neoformans gene expression during experimental
807   cryptococcal meningitis. *Eukaryotic cell* **2**, 1336-1349 (2003).
808   26.   A. D. Elbein, Y. T. Pan, I. Pastuszak, D. Carroll, New insights on trehalose: a
809   multifunctional molecule. *Glycobiology* **13**, 17R-27R (2003).
810   27.   E. W. Petzold *et al.*, Characterization and regulation of the trehalose synthesis pathway
811   and its importance in the pathogenicity of Cryptococcus neoformans. *Infection and
812   immunity* **74**, 5877-5887 (2006).
813   28.   P. Ngamskulrungroj *et al.*, The trehalose synthesis pathway is an integral part of the
814   virulence composite for Cryptococcus gattii. *Infection and immunity* **77**, 4584-4596
815   (2009).
816   29.   M. Martinez-Esparza *et al.*, Role of trehalose-6P phosphatase (TPS2) in stress tolerance
817   and resistance to macrophage killing in Candida albicans. *Int J Med Microbiol* **299**, 453-
818   464 (2009).
819   30.   P. Van Dijck, L. De Rop, K. Szlufcik, E. Van Ael, J. M. Thevelein, Disruption of the
820   Candida albicans TPS2 gene encoding trehalose-6-phosphate phosphatase decreases
821   infectivity without affecting hypha formation. *Infection and immunity* **70**, 1772-1782
822   (2002).
823   31.   O. Zaragoza, M. A. Blazquez, C. Gancedo, Disruption of the Candida albicans TPS1
824   gene encoding trehalose-6-phosphate synthase impairs formation of hyphae and
825   decreases infectivity. *Journal of bacteriology* **180**, 3809-3815 (1998).
826   32.   N. Al-Bader *et al.*, Role of trehalose biosynthesis in Aspergillus fumigatus development,
827   stress response, and virulence. *Infection and immunity* **78**, 3007-3018 (2010).
828   33.   S. Puttikamonkul *et al.*, Trehalose 6-phosphate phosphatase is required for cell wall
829   integrity and fungal virulence but not trehalose biosynthesis in the human fungal
830   pathogen Aspergillus fumigatus. *Molecular microbiology* **77**, 891-911 (2010).
831   34.   A. Bergman, A. Casadevall, Mammalian endothermy optimally restricts fungi and
832   metabolic costs. *mBio* **1** (2010).
833   35.   A. Casadevall, Global warming could drive the emergence of new fungal pathogens. *Nat
834   Microbiol* **8**, 2217-2219 (2023).
835   36.   M. A. Garcia-Solache, A. Casadevall, Global warming will bring new fungal diseases for
836   mammals. *mBio* **1** (2010).
837   37.   N. E. Nnadi, D. A. Carter, Climate change and the emergence of fungal pathogens.
838   *PLoS pathogens* **17**, e1009503 (2021).
839   38.   V. A. Robert, A. Casadevall, Vertebrate endothermy restricts most fungi as potential
840   pathogens. *The Journal of infectious diseases* **200**, 1623-1626 (2009).
841   39.   C. Gancedo, C. L. Flores, The importance of a functional trehalose biosynthetic pathway
842   for the life of yeasts and fungi. *FEMS Yeast Res* **4**, 351-359 (2004).
843   40.   M. D. Asencion Diez *et al.*, The Production and Utilization of GDP-glucose in the
844   Biosynthesis of Trehalose 6-Phosphate by Streptomyces venezuelae. *The Journal of
845   biological chemistry* **292**, 945-954 (2017).

41. J. C. Errey *et al.*, Mechanistic insight into enzymatic glycosyl transfer with retention of
configuration through analysis of glycomimetic inhibitors. *Angewandte Chemie* **49**, 1234-
1237 (2010).
42. J. D. Farelli *et al.*, Structure of the trehalose-6-phosphate phosphatase from Brugia
malayi reveals key design principles for anthelmintic drugs. *PLoS pathogens* **10**,
e1004245 (2014).
43. R. P. Gibson, C. A. Tarling, S. Roberts, S. G. Withers, G. J. Davies, The donor subsite of
trehalose-6-phosphate synthase: binary complexes with UDP-glucose and UDP-2-
deoxy-2-fluoro-glucose at 2 A resolution. *The Journal of biological chemistry* **279**, 1950-
1955 (2004).
44. R. P. Gibson, J. P. Turkenburg, S. J. Charnock, R. Lloyd, G. J. Davies, Insights into
trehalose synthesis provided by the structure of the retaining glucosyltransferase OtsA.
Chemistry & biology **9**, 1337-1346 (2002).
45. C. M. Harvey *et al.*, Structural Analysis of Binding Determinants of Salmonella
typhimurium Trehalose-6-phosphate Phosphatase Using Ground-State Complexes.
Biochemistry **59**, 3247-3257 (2020).
46. Y. Miao *et al.*, Structural and In Vivo Studies on Trehalose-6-Phosphate Synthase from
Pathogenic Fungi Provide Insights into Its Catalytic Mechanism, Biological Necessity,
and Potential for Novel Antifungal Drug Design. *mBio* **8** (2017).
47. Y. Miao *et al.*, Structures of trehalose-6-phosphate phosphatase from pathogenic fungi
reveal the mechanisms of substrate recognition and catalysis. *Proceedings of the*
National Academy of Sciences of the United States of America **113**, 7148-7153 (2016).
48. S. Shan, H. Min, T. Liu, D. Jiang, Z. Rao, Structural insight into dephosphorylation by
trehalose 6-phosphate phosphatase (OtsB2) from Mycobacterium tuberculosis. *FASEB J*
30, 3989-3996 (2016).
49. S. Wang *et al.*, Crystal structures of Magnaporthe oryzae trehalose-6-phosphate
synthase (MoTps1) suggest a model for catalytic process of Tps1. *The Biochemical*
journal **476**, 3227-3240 (2019).
50. M. Cross *et al.*, A suicide inhibitor of nematode trehalose-6-phosphate phosphatases.
Sci Rep **9**, 16165 (2019).
51. S. Kapil, C. Petit, V. N. Drago, D. R. Ronning, S. J. Sucheck, Synthesis and in Vitro
Characterization of Trehalose-Based Inhibitors of Mycobacterial Trehalose 6-Phosphate
Phosphatases. *Chembiochem* **20**, 260-269 (2019).
52. C. Kern *et al.*, Trehalose-6-phosphate synthase from the cat flea Ctenocephalides felis
and Drosophila melanogaster: gene identification, cloning, heterologous functional
expression and identification of inhibitors by high throughput screening. *Insect molecular*
biology **21**, 456-471 (2012).
53. S. Klutts *et al.*, Purification, cloning, expression, and properties of mycobacterial
trehalose-phosphate phosphatase. *The Journal of biological chemistry* **278**, 2093-2100
(2003).
54. C. Liu, D. Dunaway-Mariano, P. S. Mariano, Rational design of reversible inhibitors for
trehalose 6-phosphate phosphatases. *European journal of medicinal chemistry* **128**,
274-286 (2017).
55. Y. T. Pan, A. D. Elbein, Inhibition of the trehalose-P synthase of mycobacteria by various
antibiotics. *Archives of biochemistry and biophysics* **335**, 258-266 (1996).
56. M. Dmitrova *et al.*, A new LexA-based genetic system for monitoring and analyzing
protein heterodimerization in Escherichia coli. *Mol Gen Genet* **257**, 205-212 (1998).
57. L. L. Lairson, B. Henrissat, G. J. Davies, S. G. Withers, Glycosyltransferases: structures,
functions, and mechanisms. *Annu Rev Biochem* **77**, 521-555 (2008).
58. L. Holm, P. Rosenstrom, Dali server: conservation mapping in 3D. *Nucleic Acids Res* **38**,
W545-549 (2010).

59. R. A. Lee, M. Razaz, S. Hayward, The DynDom database of protein domain motions. *Bioinformatics* **19**, 1290-1291 (2003).

60. G. P. Poornam, A. Matsumoto, H. Ishida, S. Hayward, A method for the analysis of domain movements in large biomolecular complexes. *Proteins* **76**, 201-212 (2009).

61. S. S. Lee *et al.*, Mechanistic evidence for a front-side, SNi-type reaction in a retaining glycosyltransferase. *Nature chemical biology* **7**, 631-638 (2011).

62. E. Krissinel, K. Henrick, Inference of macromolecular assemblies from crystalline state. *Journal of molecular biology* **372**, 774-797 (2007).

63. W. Bell *et al.*, Composition and functional analysis of the Saccharomyces cerevisiae trehalose synthase complex. *The Journal of biological chemistry* **273**, 33311-33319 (1998).

64. A. Reinders *et al.*, Structural analysis of the subunits of the trehalose-6-phosphate synthase/phosphatase complex in Saccharomyces cerevisiae and their function during heat shock. *Molecular microbiology* **24**, 687-695 (1997).

65. A. Thammahong, A. K. Caffrey-Card, S. Dhingra, J. J. Obar, R. A. Cramer, Aspergillus fumigatus Trehalose-Regulatory Subunit Homolog Moonlights To Mediate Cell Wall Homeostasis through Modulation of Chitin Synthase Activity. *mBio* **8** (2017).

66. S. Boeynaems *et al.*, Protein Phase Separation: A New Phase in Cell Biology. *Trends Cell Biol* **28**, 420-435 (2018).

67. Z. G. Qian, S. C. Huang, X. X. Xia, Synthetic protein condensates for cellular and metabolic engineering. *Nature chemical biology* **18**, 1330-1340 (2022).

68. K. W. Moremen, R. S. Haltiwanger, Emerging structural insights into glycosyltransferase-mediated synthesis of glycans. *Nature chemical biology* **15**, 853-864 (2019).

69. R. Wu *et al.*, The Crystal Structure of Nitrosomonas europaea Sucrose Synthase Reveals Critical Conformational Changes and Insights into Sucrose Metabolism in Prokaryotes. *Journal of bacteriology* **197**, 2734-2746 (2015).

70. D. A. Daines, R. P. Silver, Evidence for multimerization of neu proteins involved in polysialic acid synthesis in Escherichia coli K1 using improved LexA-based vectors. *Journal of bacteriology* **182**, 5267-5270 (2000).

71. W. H. Eschenfeldt, S. Lucy, C. S. Millard, A. Joachimiak, I. D. Mark, A family of LIC vectors for high-throughput cloning and purification of proteins. *Methods in molecular biology* **498**, 105-115 (2009).

72. S. H. Scheres, RELION: implementation of a Bayesian approach to cryo-EM structure determination. *J Struct Biol* **180**, 519-530 (2012).

73. A. Rohou, N. Grigorieff, CTFFIND4: Fast and accurate defocus estimation from electron micrographs. *J Struct Biol* **192**, 216-221 (2015).

74. A. Punjani, J. L. Rubinstein, D. J. Fleet, M. A. Brubaker, cryoSPARC: algorithms for rapid unsupervised cryo-EM structure determination. *Nature methods* **14**, 290-296 (2017).

75. S. Q. Zheng *et al.*, MotionCor2: anisotropic correction of beam-induced motion for improved cryo-electron microscopy. *Nature methods* **14**, 331-332 (2017).

76. P. Emsley, K. Cowtan, Coot: model-building tools for molecular graphics. *Acta crystallographica. Section D, Biological crystallography* **60**, 2126-2132 (2004).

77. P. D. Adams *et al.*, PHENIX: a comprehensive Python-based system for macromolecular structure solution. *Acta crystallographica. Section D, Biological crystallography* **66**, 213-221 (2010).

78. E. F. Pettersen *et al.*, UCSF Chimera--a visualization system for exploratory research and analysis. *J Comput Chem* **25**, 1605-1612 (2004).

79. J. C. Errey *et al.*, Mechanistic Insight into Enzymatic Glycosyl Transfer with Retention of Configuration through Analysis of Glycomimetic Inhibitors. *Angew. Chem.-Int. Edit.* **49**, 1234-1237 (2010).

80.   A. Idnurm, J. L. Reedy, J. C. Nussbaum, J. Heitman, Cryptococcus neoformans
949   virulence gene discovery through insertional mutagenesis. *Eukaryotic cell* **3**, 420-429
950   (2004).
951   81.   J. A. Fraser, R. L. Subaran, C. B. Nichols, J. Heitman, Recapitulation of the sexual cycle
952   of the primary fungal pathogen Cryptococcus neoformans var. gattii: implications for an
953   outbreak on Vancouver Island, Canada. *Eukaryotic cell* **2**, 1036-1045 (2003).
954   82.   D. L. Toffaletti, T. H. Rude, S. A. Johnston, D. T. Durack, J. R. Perfect, Gene transfer in
955   Cryptococcus neoformans by use of biolistic delivery of DNA. *Journal of bacteriology*
956   **175**, 1405-1411 (1993).
957   83.   B. R. Washington EJ, Cryo-EM structure of *Cryptococcus neoformans* trehalose-6-
958   phosphate synthase homotetramer in complex with uridine diphosphate and glucose-6-
959   phosphate. https://www.emdataresource.org/EMD-29172 (2023).
960   84.   B. R. Washington EJ, Cryo-EM structure of *Cryptococcus neoformans* trehalose-6-
961   phosphate synthase homotetramer in apo form. https://www.emdataresource.org/EMD-
962   29338 (2024).
963   85.   E. J. Washington, Brennan, R.G., Cryo-EM structure of *Cryptococcus neoformans*
964   trehalose-6-phosphate synthase homotetramer in apo form.
965   https://www.rcsb.org/structure/8fo1 (2024).
966   86.   E. J. Washington, Brennan, R.G., Cryo-EM structure of *Cryptococcus neoformans*
967   trehalose-6-phosphate synthase homotetramer in complex with uridine diphosphate and
968   glucose-6-phosphate. https://www.rcsb.org/structure/8fhw (2024).
969
970

971 **Tables**

973 **Data Table 1. Cryo-EM data collection and refinement statistics for unliganded CnTps1.**

EM data collection statistics

Protein	CnTps1
EMDB	EMD-29338
Microscope	FEI Titan Krios
Voltage (kV)	300
Detector	Gatan K3
Magnification (nominal)	105,000
Pixel size (Å/pix)	0.65
Frames per exposure	60
Exposure (e^- $Å^{-2}$)	62
Defocus range	-2.5 – -0.8 um
Micrographs collected	2,520
Particles extracted/final	456,020/207,081
Symmetry imposed	D2
Map sharpening B-factor	-145
Unmasked resolution at 0.143 FSC (Å)	3.8
Masked resolution at 0.143 FSC (Å)	3.3

Model refinement and validation statistics

PDB	8FO1
Composition	
Atoms	6178 (Hydrogens: 3069)
Residues	403
Ligands	0
RMSD bonds (Å)	0.007
RMSD angles (°)	0.591
Ramachandran	
Favored (%)	92.51
Allowed (%)	7.49
Outliers (%)	0.00
Rotamer outliers (%)	0.00
Clash score	4.99
C-beta outliers (%)	0.00
CaBLAM outliers (%)	0.81
CC (mask)	0.85
MolProbity score	1.73

Data Table 2. Cryo-EM data collection and refinement statistics for CnTps1-UDP-G6P.

EM data collection statistics

Protein	CnTps1-UDP-G6P
EMDB	EMD-29172
Microscope	FEI Titan Krios
Voltage (kV)	300
Detector	Gatan K3
Magnification (nominal)	81,000
Pixel size (Å/pix)	1.08
Frames per exposure	60
Exposure (e⁻ Å⁻²)	62
Defocus range	-2.5 – -0.8 um
Micrographs collected	3,182
Particles extracted/final	2,003,550/854,608
Symmetry imposed	D2
Map sharpening B-factor	-159
Unmasked resolution at 0.143 FSC (Å)	3.8
Masked resolution at 0.143 FSC (Å)	3.1

Model refinement and validation statistics

PDB	8FHW
Composition	
Atoms	6639 (Hydrogens: 3220)
Residues	433
Ligands	UDP:1
	G6P: 1
RMSD bonds (Å)	0.005
RMSD angles (°)	0.538
Ramachandran	
Favored (%)	96.20
Allowed (%)	3.80
Outliers (%)	0.00
Rotamer outliers (%)	0.00
Clash score	3.98
C-beta outliers (%)	0.00
CaBLAM outliers (%)	0.49
CC (mask)	0.82
MolProbity score	1.44

A

UDP-Glucose

Glucose-6-phosphate

Trehalose-6-phosphate

Trehalose

B

C

EV   CnTps1₅₆₋₆₀₉

****

EV   CnTps1₅₆₋₆₀₉
+   -   +   -   1 mM IPTG

A

apo CnTps1
CnTps1-UDP-G6P

α2

90°

B

~ 7 Å

α2