A mathematical framework for the emergence of winners and losers in cell competition

Thomas F. Pak[a], Joe M. Pitt-Francis[b], Ruth E. Baker[a]

[a]Mathematical Institute, University of Oxford, Andrew Wiles Building, Radcliffe Observatory Quarter, Woodstock Road, Oxford, OX2 6GG, UK
[b]Department of Computer Science, University of Oxford, Wolfson Building, Parks Road, Oxford, OX1 3QD, UK

Abstract

Cell competition is a process in multicellular organisms where cells interact with their neighbours to determine a "winner" or "loser" status. The loser cells are eliminated through programmed cell death, leaving only the winner cells to populate the tissue. Cell competition is context-dependent; a given cell type can win or lose depending on the cell type it is competing against. Hence, winner/loser status is an *emergent* property. A key question in cell competition is: how do cells acquire their winner/loser status? In this paper, we propose a mathematical framework for studying the emergence of winner/loser status based on a set of quantitative criteria that distinguishes competitive from non-competitive outcomes. We apply this framework in a cell-based modelling context, to both highlight the crucial role of active cell death in cell competition and identify the factors that drive cell competition.

Keywords: cell-based model, vertex-based model, programmed cell death, epithelial tissue

1. Introduction

Cell competition is a process that occurs in multicellular organisms where cells composing genetically heterotypic tissues interact to determine their relative fitness and acquire a winner or loser status [1–6]. The loser cells are then eliminated through programmed cell death, leaving only winner cells to populate the tissue. Cell competition is context-dependent: the competing cell types are both viable in homotypic conditions,

and acquire a winner/loser status only when exposed to each other in the same tissue.
The main function of cell competition is to improve the overall fitness of the tissue by
removing suboptimal cells. For example, during development of the *Drosophila* wing,
cell competition serves as a homeostatic mechanism that stabilises tissue growth and
ensures consistent wing shape [7]. It can also play a role in tumour suppression by elim-
inating cells with proto-oncogenic mutations [8]. However, this is not the case for all
proto-oncogenic mutations: overexpression of *Myc* results in mutants that outcompete
wild-type cells in a process known as super-competition [9]. This allows precancerous
cells to expand within a tissue at the expense of healthy cells, without producing de-
tectable morphological abnormalities. Cell competition can therefore also contribute to
the early stages of tumour development.

The underlying mechanisms of cell competition are not yet fully understood. While
progress has been made in identifying the drivers of cell competition and the path-
ways downstream of winner/loser identification, the intra- and intercellular processes
by which cells determine winner/loser status are still unclear. Mathematical modelling,
particularly cell-based modelling, has the potential to provide insight into the mecha-
nisms of cell competition. Cell-based models allow researchers to define the behaviours
of individual cells and study their effects at the population level. Because cell compe-
tition is a process that unfolds at the population level but is mediated by interactions
at the cellular level, cell-based models are potentially an effective tool for exploring the
most pertinent questions in cell competition. However, current cell-based models of
cell competition assume *a priori* winner/loser identities [10–13]. Although such models
can simulate processes occurring downstream of winner/loser identification, they do not
address *how* cells become winners or losers in the first place. In this paper, we propose
a mathematical framework to address precisely this question.

1.1. Emergence of winner/loser status

33 In our framework, we do not assume that certain cells are winners or losers *a pri-*
34 *ori.* Instead, we consider cell-based models with two cell types that vary only in their

Figure 1: Illustration of the cell competition criteria. The two cell types A and B fulfil the cell competition criteria if (i) cell type A is homotypically viable, (ii) cell type B is homotypically viable, and (iii) cell type A is heterotypically viable and cell type B is heterotypically nonviable *or*, conversely, cell type A is heterotypically nonviable and cell type B is heterotypically viable.

parameters and investigate the conditions that lead to competitive outcomes. Because this approach involves detecting rather than asserting winners and losers, we need a stringent definition of what a "competitive outcome" entails. We consider two defining features of cell competition: both of the competing cell types are viable when grown in homotypic conditions, and the loser cells are completely eliminated in heterotypic conditions. Therefore, to identify competitive outcomes between two competing cell types in a cell-based model, we evaluate their viability in both homotypic and heterotypic conditions. An interaction between two cell types is classified as competitive if both cell types are viable in a homotypic environment and only one cell type remains viable in a heterotypic environment. These are the **cell competition criteria**, illustrated in Figure 1.

We use these cell competition criteria to identify parameter regimes that are associ-

ated with cell competition. Our approach has two important advantages over modelling frameworks that hardcode winner/loser identities. Firstly, it allows us to determine whether a given cell-based model is capable of displaying cell competition. Secondly, characterising the parameter regimes that lead to competitive outcomes helps us identify and analyse the factors that drive cell competition. Finally, we note that our framework respects the context-dependent nature of cell competition; winner/loser status is treated as an emergent property that exists only in the relationship between two cell types and is not inherent to any particular cell type.

1.2. Outline

In this paper, we apply the framework for emergent winner/loser status to two different models. The first model is discussed in Section 2, where we investigate whether differences in mechanical parameters in a vertex-based model constitute a sufficient mechanism for cell competition. We perform a large parameter sweep to search for competitive outcomes, but we do not find any evidence for competitive outcomes, suggesting that an active mechanism of cell death is necessary for cell competition. Motivated by these results, we introduce a modelling framework in Section 3 that simulates the intercellular exchange of death signals and the intracellular initiation of apoptosis. We use this modelling framework in Section 4 to construct the second model, where cells interact by emitting death signals. To investigate the potential for competitive outcomes in this model, we use a combination of analytical reasoning and computational simulations. Not only are we able to produce competitive outcomes, but we also identify additional biologically relevant competition regimes. Finally, in Section 5, we provide a discussion of the main results, and propose a conceptual model of cell competition based on two key cellular properties: tolerance to, and emission of, death signals. We examine the experimental evidence in support of this model and suggest novel cell competition experiments inspired by it.

2. Cell competition via differing biomechanical properties

Mechanical cell competition is a special case of cell competition, observed specifically in epithelia, that is mediated through mechanical interactions [14]. The losers in this interaction are more sensitive to cell compression than the winners and initiate apoptosis in response to cell crowding [15, 16]. In addition, we note that epithelial tissues shed live cells in response to cell crowding under homotypic conditions [17, 18]. In this case, cells undergo a "passive" form of cell death because they are extruded from the tissue as a result of mechanical interactions, and only die after being removed from the tissue. In this section, we investigate the question: are differences in biomechanical properties, combined with passive cell extrusion, sufficient to engender cell competition? A suitable cell-based framework for simulating the mechanical interactions in epithelial tissues is vertex-based modelling, since it has been shown to reproduce the dynamics of epithelial tissues in a variety of developmental processes [19]. The overall strategy of this section, then, is to construct a heterotypic vertex-based model that allows for the independent variation of mechanical parameters between two cell types, and to test whether this variation is sufficient to give rise to competitive outcomes.

2.1. Vertex-based model

In vertex-based modelling, the epithelial tissue is represented by a polygonal mesh where each polygon corresponds to an epithelial cell, and the dynamics of the tissue is based on the motion of the mesh vertices. In particular, the equation of motion for vertex i with position $\mathbf{r}_i$, experiencing the total force $\mathbf{F}_i$, has the form [20]

$$\mu \frac{\mathrm{d}\mathbf{r}_i}{\mathrm{d}t} = \mathbf{F}_i \,, \tag{1}$$

where μ is the friction coefficient. The force acting on vertex i is given by

$$\mathbf{F}_i = \nabla_i E \,, \tag{2}$$

where ∇_i is the gradient of an energy function E with respect to the spatial coordinates of vertex i. We use the energy function presented in [21], which describes three major

biomechanical properties: cell elasticity, cell contractility, and cell–cell adhesion;

$$E = \sum_\alpha \frac{K_\alpha}{2} \left(S_\alpha - S_\alpha^0\right)^2 + \sum_\alpha \frac{\Gamma_\alpha}{2} L_\alpha^2 + \sum_{\langle i,j \rangle} \Lambda_{ij} \ell_{ij} \,. \tag{3}$$

The first term represents cell elasticity, i.e. the cell's resistance against deformation. The parameters K_α and S_α^0 are the elasticity constant and the target cell area of cell α, respectively, while S_α is the cell area of cell α. The second term models cell contractility, with Γ_α and L_α corresponding to the contractility constant and the cell perimeter of cell α, respectively. The final term represents cell–cell adhesion, which is implemented as a line tension acting on cell–cell interfaces. For each edge $\langle i,j \rangle$ connecting the vertices i and j, this line tension is the product of the line tension constant, Λ_{ij}, and the edge length, ℓ_{ij}.

In addition to vertex dynamics, the vertex-based model also evolves through mesh rearrangements that allow cells to exchange neighbours, to proliferate, and to be extruded from the tissue. During cell division, a new edge is formed that bisects the mother cell and results in two daughter cells. Cell extrusion, on the other hand, is achieved by the "T2 swap", which removes cells when their cell area falls below a certain threshold. There are many technical details involved with mesh rearrangements, and we refer the reader to [22] for further details.

Motivated by experiments with *in vitro* cell cultures [23], we assume a two-phase cell cycle model. The first phase corresponds to the G1 phase, and we lump together the S, G2 and M phases in the second phase. For brevity, we refer to the second phase as the G2 phase. For cell α, the duration of G1 phase is exponentially distributed with mean $t_{\text{G1},\alpha}$. The G2 phase lasts for the fixed duration $t_{\text{G2},\alpha}$. At the end of the G2 phase, cell division occurs as described above.

We divide the cell population into two non-overlapping sets that correspond to two distinct cell types, A and B. The mechanical and cell cycle constants for each cell are

determined by its cell type. In particular, the elasticity constant K_αp is given as

$$K_\alpha = \begin{cases} K_A & \text{for } \alpha \in A \\ K_B & \text{for } \alpha \in B \end{cases}, \tag{4}$$

and the target cell area S_α^0, contractility constant Γ_α, and cell cycle constants $t_{\text{G1},\alpha}$ and $t_{\text{G2},\alpha}$ are determined analogously. Since the line tension parameter is dependent on the edge type, rather than the cell type, we need to specify values for every pairing of cell types. In addition, we need to account for edges at the boundary of the tissue, which border a cell on one side and empty space on the other. Denoting the two cells sharing the edge $\langle i, j \rangle$ as α and β, we write

$$\Lambda_{ij} = \begin{cases} \Lambda_{AA} & \text{for } \alpha, \beta \in A \\ \Lambda_{BB} & \text{for } \alpha, \beta \in B \\ \Lambda_{AB} & \text{for } \alpha \in A, \beta \in B \\ \Lambda_A & \text{for } \alpha \in A, \beta \in \emptyset \\ \Lambda_B & \text{for } \alpha \in B, \beta \in \emptyset \end{cases}, \tag{5}$$

where $\beta \in \emptyset$ signifies that $\langle i, j \rangle$ is a boundary edge. Furthermore, we impose that each cell division results in cells that are of the same type as the mother cell, i.e. a cell of type A divides into two daughter cells of type A.

We implemented the heterotypic vertex-based model within Chaste, an open-source simulation package for computational physiology and biology [24] that includes a range of cell-based models [25]. We refer the reader to the following GitHub repository for the code of the heterotypic vertex-based model: `https://github.com/ThomasPak/cell-competition`.

2.2. Methods

After constructing the heterotypic vertex-based model, the next step is to determine whether it is capable of generating competitive outcomes. We performed a systematic parameter sweep of the heterotypic vertex-based model, and evaluated the outputs

against the cell competition criteria defined in Section 1.1. In particular, we varied the parameters of the model as indicated in Table S1, giving rise to 3^7 unique parameter sets. For every parameter set, we conducted two heterotypic simulations: one where the cell types are initially randomly distributed in the tissue, and one where the cell types are initially spatially segregated. In both cases, the population is initially composed of 18 A-type cells and 18 B-type cells. In addition, for each heterotypic simulation we conducted a homotypic simulation initially containing 36 A-type cells.

We characterised the proliferative behaviour of cells using the **survival frequency**. For each homotypic simulation, we computed the homotypic survival frequency, denoted by $\hat{\lambda}$, as

$$\hat{\lambda} = \frac{\# \text{ divisions}}{\# \text{ divisions} + \# \text{ deaths}} \,. \tag{6}$$

For each heterotypic simulation, we computed the heterotypic survival frequencies as

$$\hat{\xi}_{A|B} = \frac{\# \text{ A divisions}}{\# \text{ A divisions} + \# \text{ A deaths}} \,, \tag{7}$$

$$\hat{\xi}_{B|A} = \frac{\# \text{ B divisions}}{\# \text{ B divisions} + \# \text{ B deaths}} \,, \tag{8}$$

for cell types A and B, respectively.

2.3. Results

Since the survival frequency indicates whether a cell type has increased or declined in number over the course of a simulation, we can use this quantity to evaluate the viability of a cell type. Concretely, we consider a cell type in a given simulation as viable if its survival frequency is greater than or equal to a half, since that signifies that the population size has grown or stayed the same, respectively, and as nonviable otherwise.

We found that cell type B is consistently viable in all heterotypic and homotypic simulations. Considering the viability of cell type A in each heterotypic simulation and its homotypic counterpart, we observed occurrences of all possible combinations of homotypic and heterotypic viability:

	$\hat{\lambda}_A < 1/2$	$\hat{\lambda}_A \geq 1/2$
$\hat{\xi}_{A|B} < 1/2$	84	7
$\hat{\xi}_{A|B} \geq 1/2$	58	4 225

Table 1: Count of simulation pairs for every combination of homotypic and heterotypic viability of cell type A. Note that simulation pairs in category (iii) satisfy the cell competition criteria.

(i) $\hat{\lambda}_A < 1/2$, $\hat{\xi}_{A|B} < 1/2$: cell type A is nonviable in homotypic and heterotypic conditions;

(ii) $\hat{\lambda}_A < 1/2$, $\hat{\xi}_{A|B} \geq 1/2$: cell type A is nonviable in homotypic conditions and viable in heterotypic conditions;

(iii) $\hat{\lambda}_A \geq 1/2$, $\hat{\xi}_{A|B} < 1/2$: cell type A is viable in homotypic conditions and nonviable in heterotypic conditions;

(iv) $\hat{\lambda}_A \geq 1/2$, $\hat{\xi}_{A|B} \geq 1/2$: cell type A is viable in homotypic and heterotypic conditions.

We assigned each pair of heterotypic and homotypic simulations into these four cases based on the observed homotypic and heterotypic survival frequencies, and list the results in Table 1. For the overwhelming majority of parameter sets, cell type A was viable both in homotypic and heterotypic conditions. Only seven out of 4 374 parameter sets, approximately 0.16%, satisfy the cell competition criteria encoded in category (iii).

To determine whether those seven parameter sets represent true cases of cell competition, and not just a statistical fluke, we conducted an additional simulation suite targeting those seven parameter sets in particular. Specifically, we ran 20 pairs of simulations (one heterotypic simulation with random initial conditions and one homotypic simulation) for each targeted parameter set. We evaluated the viability of cell type A as before and counted the number of simulation pairs that satisfy the cell competition criteria. One of the parameter sets yielded three simulation pairs out of 20 that satisfied the cell competition criteria, while the other parameter sets yielded even fewer matches.

This strongly suggests that the competitive outcomes we encountered in the parameter sweep were statistical flukes rather than genuine examples of competitive behaviour. Indeed, a statistical analysis of the results (Section S1 of the supplementary material) shows that there is insufficient evidence in support of competitive behaviour.

2.4. Discussion

In this section, we constructed a heterotypic vertex-based model to investigate whether differences in mechanical properties are sufficient to give rise to cell competition. We performed a large, systematic parameter sweep with this model, and found that we could not reliably reproduce competitive behaviour for any of the simulated parameter sets. We conclude that, for the parameter space investigated, simply varying the mechanical parameters in the heterotypic vertex-based model is not sufficient for cell competition. This agrees with experiments suggesting that cell competition generally depends on an active mechanism of cell death, such as apoptosis [26], and that mechanical cell competition is no exception in this respect [15, 16]. These results do not mean that mechanical interactions cannot play a role in cell competition. They do strongly suggest, however, that passive cell death alone is insufficient for cell competition and that mechanical interactions must be paired with an active mechanism of cell death to generate competitive outcomes.

3. Cell competition via exchange of death signals

The results of the previous section suggest that cell competition requires an **active** and **non-autonomous** mechanism of cell death. This observation is also supported experimentally [7, 8, 26]. Therefore, the aim in this section is to develop a modelling framework for cell competition implementing such an active and non-autonomous mechanism for cell death. The core idea is that cells exchange "death signals" with their neighbours and that these signals are accumulated by the cell into an abstract quantity called the "death clock". When the death clock reaches a threshold value, apoptosis is triggered. We do not attach the death signal to a concrete biological mechanism because

there are multiple competing hypotheses regarding the mode of intercellular communication taking place in cell competition, and because the mode of communication likely depends on the type of cell competition under consideration.

We first discuss the underlying biological assumptions and modelling choices in Section 3.1, before introducing the death clock framework in Section 3.2. In Section 3.3, we derive the survival probability of cells, which allows us to analyse the death clock framework from a theoretical perspective. In Section 3.4, we discuss the computational cell-based models implementing the death clock framework that we use to validate theoretical predictions.

3.1. Assumptions

A series of studies involving mathematical modelling and experiments have revealed the importance of threshold mechanisms in the initiation of apoptosis [27–30]. For instance, it was shown that death ligand-induced apoptosis requires a threshold proportion of ligand to receptor numbers to be reached [29, 30]. Given this precedent, we propose a model in which competition-induced apoptosis is triggered by the accumulation of death signals reaching a threshold value.

Further, it has been established in the literature that apoptosis and the cell cycle are closely coupled [31–34]. Notably, the regulatory protein *Myc* is known to affect both cell cycle progression and apoptosis [35–37]. On the one hand, *Myc* is necessary for the transition of G1 to S phase, and it induces cell cycle progression in quiescent cells [35, 37]. On the other hand, *Myc* has been associated with increased rates of cell death [36]. Coupled with the fact that differential *Myc* expression results in cell competition [9], we hypothesise that apoptosis, competition, and the cell cycle are interrelated. Concretely, we assume that the cell is only susceptible to competition-induced apoptosis in G1 phase, and that from S phase onwards, the cell is committed to division. As in the vertex-based model introduced in Section 2.1, we assume a two-phase cell cycle model, where we treat the duration of the G1 phase as a random variable and lump together the S, G2 and M phases into the G2 phase, which lasts for

 a fixed duration.

 3.2. Death clock framework

The death clock framework consists of two coupled cellular processes: the cell cycle and the death clock, where the death clock governs the initiation of apoptosis in response to death signals. We consider the cell cycle to be an **autonomous** process, meaning that it is not affected by other cells. On the other hand, the death clock is a **non-autonomous** process because it is driven by extracellular signals produced by other cells. Together, these processes determine whether and when the cell divides or initiates apoptosis.

At division, we sample a stochastic G1 duration, denoted as t^*, from the **G1 duration distribution** $\mathcal{C}$, i.e.

$$t^* \sim \mathcal{C} \, , \tag{9}$$

where $\mathcal{C}$ is subject to the constraints that (i) $t^* \in [0, \infty)$ and (ii) $E(t^*) = t_{\mathrm{G1}}$, and we term t_{G1} the **autonomous G1 duration**. If apoptosis is not triggered by the death clock, the cell spends a duration t^* in G1 phase and then transitions into G2 phase. After spending a fixed duration, t_{G2}, in G2 phase, the cell divides and the process repeats for each of the daughter cells.

We model the accumulation of death signals using an ordinary differential equation (ODE) model in which the **death clock**, denoted by $\tau(t)$, evolves according to the ODE

$$\frac{\mathrm{d}\tau}{\mathrm{d}t} = f(t) \, , \tag{10}$$

where $f(t) \geq 0$ is the **death signal** experienced by the cell. At birth, the death clock of a cell is initialised to zero, i.e. $\tau(t = 0) = 0$. The apoptosis rule is then

$$\text{Cell is in G1 phase} \quad \textbf{and} \quad \tau(t) \text{ reaches } T_\dagger$$

$$\Downarrow$$

$$\text{initiate apoptosis} \, ,$$

Figure 2: Death clock flowchart. The "Sample" step corresponds to Equation (9), and the "Integrate" step corresponds to Equation (10). The condition in the decision block is the survival condition, corresponding to Equation (11).

where $T_\dagger$ is the **death threshold**. We define the **survival condition** as

$$\tau(t^*) < T_\dagger. \tag{11}$$

We note that there are two potential sources of uncertainty in the death clock framework: variability in G1 duration and in the death signal. The former originates from the cell cycle, the latter from intercellular interactions, and both contribute to the decision of the cell to initiate apoptosis. Our framework can thus be regarded as a minimalist model of autonomous and non-autonomous processes interacting to govern competition-induced apoptosis. The death clock framework is summarised by the flowchart in Figure 2.

3.3. Survival probability

In order to predict the viability of a cell population, we must determine the probability of cells surviving. This problem is intractable when considering the uncertainty in the death signal and in the cell cycle simultaneously. To make analytic progress, we fix the death signal and consider only variance in the cell cycle, which lets us derive an expression for the "survival probability". We define this survival probability, which we denote by θ, as the probability that the survival condition (Equation (11)) is satisfied,

273 i.e.

$$\theta \equiv P(\tau(t^*) < T_\dagger) \,. \tag{12}$$

274 Furthermore, assuming that $f(t)$ is a non-negative integrable function, we define

$$F(t) \equiv \int_0^t f(t') \, \mathrm{d}t' \,, \tag{13}$$

275 such that the value of the death clock at time t^* is $F(t^*)$. This lets us write the survival
276 condition as $F(t^*) < T_\dagger$. We define the pseudoinverse function of $F(t)$ as

$$F^{-1}(\tau) \equiv \min\{t \in [0, \infty) : F(t) = \tau\} \,, \tag{14}$$

277 so that we can reformulate the survival condition as

$$t^* < F^{-1}(T_\dagger) \,. \tag{15}$$

278 Substituting this into Equation (12), and denoting the cumulative distribution function
279 for the distribution of t^* as $\Psi(t)$, we obtain

$$\theta = P\left(t^* < F^{-1}(T_\dagger)\right) = \Psi\left(F^{-1}(T_\dagger)\right) \,. \tag{16}$$

As a special case, consider the constant death signal $f(t) = c$, where $c > 0$ is a positive
constant. We then have $F(t) = ct \Rightarrow F^{-1}(\tau) = \tau/c \Rightarrow \theta = \Psi(T_\dagger/c)$.

3.4. Cell-based death clock models

So far, we have described the processes leading to competition-induced apoptosis
from the perspective of a single cell. The death clock framework can be embedded in any
cell-based model that (i) provides cells with an extracellular environment from which
to derive a death signal and (ii) includes a cellular operation for initiating apoptosis.
In this paper, we implement the death clock mechanism in two particular cell-based
models: the vertex-based model (Section 2.1), and a well-mixed model. In the vertex-
based model, a cell interacts only with cells in its local neighbourhood. In the well-mixed
model, on the other hand, each cell interacts with every other cell on an equal basis. In

Symbol	Description
$\tau_\alpha(t)$	Death clock
t_α^*	Sampled G1 duration
t_α^0	Birth time
$\mathcal{C}_\alpha$	G1 duration distribution
$t_{\mathrm{G2},\alpha}$	G2 duration
$f_\alpha(\cdot)$	Death signal function
$T_{\dagger,\alpha}$	Death threshold

Table 2: Summary of cell vector elements.

Section 4, we use these models in a complementary manner. Here, we present a high-level outline of the well-mixed and vertex-based models. For a detailed discussion of their numerical implementations, see Sections S2 and S3 in the supplementary material. We provide the code for both models in the following GitHub repository: `https://github.com/ThomasPak/cell-competition`.

3.4.1. Well-mixed model

For each cell α, we represent its state with the cell vector $\mathbf{y}_\alpha(t)$, where $\alpha = 1, \ldots, N(t)$, and $N(t)$ is the number of cells at time t. We write the cell vector as

$$\mathbf{y}_\alpha(t) \equiv \left[\begin{array}{ccccccc} \tau_\alpha(t) & t_\alpha^* & t_\alpha^0 & \mathcal{C}_\alpha & t_{\mathrm{G2},\alpha} & f_\alpha(\cdot) & T_{\dagger,\alpha} \end{array} \right], \tag{17}$$

and summarise its contents in Table 2. The state of the system, denoted $S(t)$, is then

$$S(t) \equiv \left\{ \mathbf{y}_1(t), \mathbf{y}_2(t), \ldots, \mathbf{y}_{N(t)}(t) \right\}. \tag{18}$$

We evolve the death clock for each cell α as

$$\frac{\mathrm{d}\tau_\alpha}{\mathrm{d}t} = f_\alpha(\mathbf{x}_\alpha(t)), \tag{19}$$

where $f_\alpha(\cdot)$ is the death signal function and $\mathbf{x}_\alpha(t)$ is the "input vector" representing the extracellular environment. Since the cell population is well-mixed, this environment is composed of every cell except itself, i.e. $\mathbf{x}_\alpha(t) = \left[\mathbf{y}_1(t), \ldots, \mathbf{y}_{\alpha-1}(t), \mathbf{y}_{\alpha+1}(t), \ldots, \mathbf{y}_{N(t)}(t) \right]$.

In addition, we define two discrete operations: cell division and cell death. When a cell's age reaches its total cell cycle duration, the division operation is triggered which constructs two daughter cells; one in a new cell vector and the other reusing the mother cell vector. When a cell's death clock reaches the death threshold in G1 phase, the cell is removed from the population. See Section S2 in the supplementary material for further implementation details.

3.4.2. Vertex-based model

In the vertex-based model, the death clock for each cell is similarly evolved using Equation (19). However, the input vector $\mathbf{x}_\alpha(t)$ is constrained to contain only information about the local extracellular environment of cell α, for instance the states of its direct neighbours. Apoptosis is implemented in the vertex-based model by shrinking the target cell area, S_α^0, to zero, which causes the cell to contract until it is extruded from the tissue. We implemented the death clock model by augmenting the basic vertex-based model, introduced in Section 2, with the death clock framework. See Section S3 in the supplementary material for further implementation details.

4. The G2 death signal model

We now explore a particular form of the death signal function, $f_\alpha(\cdot)$, which we term the G2 death signal. In the G2 death signal model, cells emit death signals to their neighbours while they are in G2 phase. This choice is motivated by the observation that cell competition often manifests as patches of proliferating cells inducing apoptosis in neighbouring cells to make room for themselves. In the death clock framework, cells in G2 phase are committed to division, so we chose to associate the death signal with the decision to proliferate. Moreover, there is experimental evidence suggesting that there is a link between cell cycle progression and death signals [38, 39]. Concretely, the G2 death signal is defined as

$$f(t) = cg(t)\,, \tag{20}$$

where $g(t)$ is the proportion of neighbouring cells in G2 phase, i.e.

$$g(t) = \begin{cases} \dfrac{\text{\# neighbours in G2}}{\text{\# neighbours}} & \text{if \# neighbours} > 0 \\ 0 & \text{otherwise} \end{cases}, \qquad (21)$$

and c is a positive constant.

We first examine the behaviour of the G2 death signal model in homotypic populations in Section 4.1, before moving on to heterotypic populations in Section 4.2. The goal is characterise the parameter regimes for which the population (or subpopulations in a heterotypic population) are viable or nonviable. In Section 4.3, we combine the results for homotypic and heterotypic populations to identify biologically relevant competition regimes. In particular, we apply the cell competition criteria to show that the G2 death signal model produces competitive outcomes. Moreover, additional competition regimes beyond the classical ones already posed in the literature are identified.

4.1. Homotypic populations

The fact that cells only emit death signals in G2 phase leads to an important trade-off; cells in G1 phase are vulnerable to death signals and do not generate death signals, whereas cells in G2 phase are impervious to death signals but do generate death signals. This raises the question: what is the impact of changing the proportion of the cell cycle that is spent in G1 or G2 phase on the survival probability, given a fixed total cell cycle duration? In order to investigate this question, we denote the total cell cycle duration as t_{G}, and define β as the fraction of the cell cycle that is spent, on average, in G1 phase, so that

$$t_{\mathrm{G1}} = \beta t_{\mathrm{G}}, \quad t_{\mathrm{G2}} = (1 - \beta)t_{\mathrm{G}}. \qquad (22)$$

In order to derive the homotypic survival probability, we first obtain an expression for the G2 death signal under homotypic conditions. Despite the fact that the model is stochastic, it turns out that the death signal is not only relatively stable, but even predictable. In particular, we observe that the system is **ergodic**, in the sense that the average proportion of cells in G2 phase relative to the population well approximates

the average proportion of the cell cycle spent in G2 phase. More precisely, we state that the system is ergodic if, on average,

$$\frac{\#\text{ cells in G2}}{\#\text{ cells}} \approx \frac{\text{G2 duration}}{\text{cell cycle duration}}\,. \tag{23}$$

Furthermore, if the system is well-mixed, then we can approximate $g(t)$ as

$$g(t) \approx \frac{\#\text{ cells in G2}}{\#\text{ cells}}\,. \tag{24}$$

Combining Equations (23) and (24), we have

$$g(t) \approx \frac{\text{G2 duration}}{\text{cell cycle duration}} = 1 - \beta\,, \tag{25}$$

so that the death signal is

$$f(t) = cg(t) \approx c(1 - \beta)\,. \tag{26}$$

We can derive the **homotypic survival probability**, denoted λ, as

$$\lambda = \Psi\left(\frac{T_\dagger}{c(1-\beta)}\right)\,. \tag{27}$$

For the exponential cell cycle model in particular, we obtain

$$\lambda = 1 - \exp\left(-\frac{T_\dagger}{ct_{\mathrm{G}}\beta(1-\beta)}\right)\,. \tag{28}$$

In order to simplify the notation, we introduce the dimensionless parameter η,

$$\eta \equiv \frac{T_\dagger}{ct_{\mathrm{G}}}\,, \tag{29}$$

which can be interpreted as a normalised death threshold. We thus write the homotypic survival probability as a function of two dimensionless parameters:

$$\lambda(\beta, \eta) = 1 - \exp\left(-\frac{\eta}{\beta(1-\beta)}\right)\,, \tag{30}$$

and validate this expression via simulation in Section S4 of the supplementary material.

Figure 3: Homotypic proliferation regimes. (a) Diagram of homotypic proliferation regimes. The homotypic viability curve is given by Equation (31). Red: Viable Regime. Grey: Nonviable Regime. (b) Estimated homotypic survival frequency, $\overline{\lambda}$, defined in Equation (32), for the well-mixed and vertex-based models. The homotypic viability curve is plotted using a black line.

4.1.1. Homotypic proliferation regimes

Based on the homotypic survival probability λ, we distinguish between two proliferation regimes for homotypic populations:

Nonviable Regime $\left\{\lambda \leq \frac{1}{2}\right\}$. Cells are equally or more likely to die than to proliferate, hence the population declines. We say that cell types in this regime are **nonviable**.

Viable Regime $\left\{\lambda > \frac{1}{2}\right\}$. Cells are more likely to proliferate than to die, hence the population grows. We say that cell types in this regime are **viable**.

We define the **homotypic viability curve** as the curve satisfying $\lambda = 1/2$. This curve separates the Nonviable Regime from the Viable Regime. For the exponential cell cycle model, the homotypic viability curve is given by

$$\eta = \ln(2)\beta(1 - \beta).$$ (31)

This analysis therefore predicts that a population is viable if $\eta > \ln(2)/4$, and for $\eta \leq \ln(2)/4$ it is viable for extreme values of β and nonviable otherwise (Figure 3(a)).

4.1.2. Computational validation of homotypic proliferation regimes

We use computational simulation to determine whether the viability of homotypic populations *in silico* matches the homotypic proliferation regimes as predicted by the

homotypic viability curve. Further details are provided in Section S5 of the supplementary material.

For each simulation k, we computed the homotypic survival frequency, denoted by $\hat{\lambda}_k$, using Equation (6). For every unique parameter set, we averaged the homotypic survival frequency as

$$\overline{\lambda} = \frac{1}{N_{\text{sim}}} \sum_{k=1}^{N_{\text{sim}}} \hat{\lambda}_k \,, \tag{32}$$

where N_{sim} is the number of simulations for the given parameter set.

We expect that nonviable populations tend to have a survival frequency below a half, i.e. $\hat{\lambda}_k < 1/2$, and vice versa for viable populations. Figure 3(a) predicts that cell types below the homotypic viability curve are nonviable and cell types above the curve are viable. To verify these predictions, we visualise $\overline{\lambda}$ in Figure 3(b) for both the well-mixed and vertex-based models.

The left-hand plot in Figure 3(b) shows that the observed border between nonviable and viable regimes closely matches predictions for the well-mixed model. We see that for small η values, the survival frequency is asymmetrical with respect to β, with higher survival frequencies for $\beta < 1/2$ than $\beta > 1/2$. The reason for this discrepancy is discussed in Section S4.5. In short, for low η values, the rate of apoptosis is so high that the limiting factor is the number of cells susceptible to apoptosis, rather than the survival probability. For small β values, cells spend less time in G1 phase and are therefore susceptible for a shorter amount of time.

The right-hand plot in Figure 3(b) also shows good agreement between theory and simulations for the vertex-based model, although the border is less finely resolved than in the well-mixed case. We also observe the same asymmetry for small η values as seen in the well-mixed model.

4.2. Heterotypic populations

Similarly to Section 2, we split the cell population into two cell types, denoted A and B, to create a heterotypic population. Each cell type has its own cell cycle model, $\Psi_A(t)$ and $\Psi_B(t)$, death signal function, $f_A(t)$ and $f_B(t)$, and death threshold, $T_{\dagger,A}$ and $T_{\dagger,B}$.

407 We assume that the cell cycle models and death signal functions are the same in both cell
408 types, except in their parameters. With $\Psi(\cdot)$ as the common cell cycle model, the cell
409 cycle models are thus parameterised as $\Psi_A(t) = \Psi(t\,; t_{\mathrm{G1},A}), \Psi_B(t) = \Psi(t\,; t_{\mathrm{G1},B})$. Sim-
410 ilarly, the death signal functions are parameterised as $f_A(t) = c_A g(t), f_B(t) = c_B g(t)$,
411 with $g(t)$ as defined in Equation (21).

We demonstrated for homotypic populations that the proportion of the cell cycle spent in G1 phase, β, is an important nondimensional parameter in determining the survival probability. In analogy with Equation (22), we define β_A and β_B for heterotypic populations such that:

$$
\begin{aligned}
t_{\mathrm{G1},A} &= \beta_A t_{\mathrm{G},A}, t_{\mathrm{G2},A} = (1 - \beta_A) t_{\mathrm{G},A}\,; \\
t_{\mathrm{G1},B} &= \beta_B t_{\mathrm{G},B}, t_{\mathrm{G2},B} = (1 - \beta_B) t_{\mathrm{G},B}\,.
\end{aligned}
\tag{33}
$$

412 In the rest of this section, we first derive an expression for the heterotypic survival
413 probability. Then, we define and analyse the heterotypic survival difference, as well as
414 the homotypic survival difference. Using these quantities, we classify different types of
415 competitive interactions in Section 4.2.1. Finally, we derive the heterotypic proliferation
416 regimes in Section 4.2.2, and validate them with simulations in Section 4.2.3.

We generalise the ergodic approximation to obtain expressions for the heterotypic survival probabilities of cell types A and B. First, we assume that the ergodic property holds for both cell types separately. For cell type A, we have

$$
\frac{\#\text{ A cells in G2}}{\#\text{ A cells}} \approx \frac{\text{G2 duration of A cells}}{\text{cell cycle duration of A cells}}
$$
$$
= 1 - \beta_A\,,
\tag{34}
$$

417 and an analogous expression can be derived for cell type B. We denote the number of
418 A-type and B-type cells with $n_A(t)$ and $n_B(t)$, respectively, so that we can write the
419 fraction of cells in G2 phase for the whole population as

$$
\frac{\#\text{ cells in G2}}{\#\text{ cells}} = \frac{\#\text{ A cells in G2} + \#\text{ B cells in G2}}{n_A(t) + n_B(t)}\,.
\tag{35}
$$

420 We substitute Equation (34) and its analogue for cell type B to obtain

$$
\frac{\#\text{ cells in G2}}{\#\text{ cells}} \approx \frac{n_A(t)(1 - \beta_A) + n_B(t)(1 - \beta_B)}{n_A(t) + n_B(t)}\,.
\tag{36}
$$

To simplify notation, we define the weighted average

$$\langle 1 - \beta \rangle(t) \equiv \frac{n_A(t)(1 - \beta_A) + n_B(t)(1 - \beta_B)}{n_A(t) + n_B(t)} . \tag{37}$$

Assuming that the population is well-mixed, i.e. that Equation (24) holds, we can approximate $g(t)$ as

$$g(t) \approx \frac{\# \text{ cells in G2}}{\# \text{ cells}} \approx \langle 1 - \beta \rangle(t) . \tag{38}$$

For cell type A, the death signal is thus approximated as $f_A(t) = c_A g(t) \approx c_A \langle 1 - \beta \rangle(t)$. Note that the quantity $\langle 1 - \beta \rangle(t)$ is not constant with respect to time because it depends on $n_A(t)$ and $n_B(t)$. This is unlike the homotypic case, where the death signal is approximated by the constant quantity $1 - \beta$. Therefore, even with the ergodic approximation we cannot derive an exact heterotypic survival probability. Nonetheless, we can define the *instantaneous* heterotypic survival probability at time t as the survival probability of a cell *assuming* a constant death signal of magnitude $f_A(t)$, i.e.

$$\xi_{A|B}(t) = \Psi_A \left(\frac{T_{\dagger,A}}{c_A \langle 1 - \beta \rangle(t)} \right) , \tag{39}$$

where we use the symbol $\xi_{A|B}(t)$ to denote the instantaneous survival probability at time t for cell type A in a heterotypic population with cell type B. Similarly, for cell type B, we have

$$\xi_{B|A}(t) = \Psi_B \left(\frac{T_{\dagger,B}}{c_B \langle 1 - \beta \rangle(t)} \right) . \tag{40}$$

In order to derive the instantaneous heterotypic survival probability for the exponential cell cycle model in particular, we first define the dimensionless parameters

$$\eta_A \equiv \frac{T_{\dagger,A}}{c_A t_{G,A}} , \quad \eta_B \equiv \frac{T_{\dagger,B}}{c_B t_{G,B}} , \tag{41}$$

in analogy with Equation (29). We can then derive that the instantaneous heterotypic survival probabilities are

$$\xi_{A|B}(t) = 1 - \exp \left(-\frac{\eta_A}{\beta_A \langle 1 - \beta \rangle(t)} \right) , \tag{42}$$

$$\xi_{B|A}(t) = 1 - \exp \left(-\frac{\eta_B}{\beta_B \langle 1 - \beta \rangle(t)} \right) , \tag{43}$$

for cell types A and B, respectively. For brevity, we omit the word "instantaneous" going forward and use the symbols $\langle 1 - \beta \rangle$ and $\xi_{A|B}$ instead of $\langle 1 - \beta \rangle(t)$ and $\xi_{A|B}(t)$, except when we wish to emphasise their time dependence. Furthermore, in the rest of the paper we will assume an exponential cell cycle model, unless stated otherwise.

Comparing the expressions for the heterotypic survival probability and the homotypic survival probability (Equation (30)), we see that they are almost identical, except that the weighted average $\langle 1 - \beta \rangle$ is used instead of $1 - \beta$. We note that if $n_B = 0$, then $\langle 1 - \beta \rangle = 1 - \beta_A$ and vice versa for $n_A = 0$. In other words, when either cell type A or B is absent, we recover the homotypic survival probability of the other cell type.

Even though the instantaneous heterotypic survival probabilities $\xi_{A|B}(t)$ and $\xi_{B|A}(t)$ evolve with time, we now show that the sign of their difference is invariant with respect to system state, and therefore only depends on model parameters. This enables us to partition the parameter space based on which cell type has a higher survival probability.

We define the **heterotypic survival difference** between cell types A and B as

$$\Delta_{A|B}^{\neq} \equiv \xi_{A|B} - \xi_{B|A}. \tag{44}$$

The sign of the heterotypic survival difference tells us which cell type is at a proliferative advantage. If $\Delta_{A|B}^{\neq} > 0$, then we say that A-type cells are **winner cells** and B-type cells are **loser cells**, and vice versa for $\Delta_{A|B}^{\neq} < 0$. Moreover, if $\Delta_{A|B}^{\neq} = 0$, we say that the cell types are in **coexistence**, since neither cell type has an advantage over the other.

We define winners and losers here in a weak sense; if the population were to reproduce indefinitely, the winner cells would come to dominate the heterotypic population. It is not specified whether the loser population is eliminated. The classical definition of winners and losers, however, is based on the stronger condition of loser elimination. In Section 4.3, we will refine our terminology and differentiate winners and losers into more precise categories, which include classical winners and losers.

We also note that this definition of winners and losers relies on the assumption that $t_{G,A} = t_{G,B}$, such that differences in survival probability alone determine relative

proliferative success. In the general case, however, differences in the total cell cycle duration can also affect the dynamics of heterotypic populations. For instance, a cell type with a lower survival probability may become more abundant than the competing cell type by dividing more rapidly. However, for the sake of simplicity we do not consider such cases in this paper, and instead characterise population dynamics solely in terms of survival probabilities.

To obtain an expression for the sign of $\Delta_{A|B}^{\neq}$, we substitute Equations (42) and (43) into Equation (44) and rearrange to give

$$\Delta_{A|B}^{\neq} = \exp\left(-\frac{\eta_B}{\beta_B \langle 1 - \beta \rangle}\right) - \exp\left(-\frac{\eta_A}{\beta_A \langle 1 - \beta \rangle}\right) . \tag{45}$$

Since $\exp(\cdot)$ is a monotonically increasing function, we have $\mathrm{sgn}(\exp(x) - \exp(y)) = \mathrm{sgn}(x - y)$. Applying the sign function thus yields

$$\mathrm{sgn}\left(\Delta_{A|B}^{\neq}\right) = \mathrm{sgn}\left(\frac{\eta_A}{\beta_A \langle 1 - \beta \rangle} - \frac{\eta_B}{\beta_B \langle 1 - \beta \rangle}\right)$$
$$= \mathrm{sgn}\left(\frac{\eta_A}{\beta_A} - \frac{\eta_B}{\beta_B}\right) . \tag{46}$$

To interpret Equation (46), we note that η and β both affect a cell's sensitivity to the death signal. Increasing η corresponds to a higher death threshold, and thus a lower sensitivity, and decreasing β shortens the time spent in G1 phase, during which a cell is vulnerable to competition-induced apoptosis. This suggests that we can interpret η/β as a cell's tolerance to death signals. Therefore, Equation (46) states that the relative tolerance to death signals determines winner/loser status, with the most tolerant cell type becoming the winner.

Since the sign of $\Delta_{A|B}^{\neq}$ depends only on model parameters, we can partition the parameter space into two regions for which $\Delta_{A|B}^{\neq} > 0$ and $\Delta_{A|B}^{\neq} < 0$, respectively. We define the **coexistence curve** for fixed β_B and η_B as the curve in (β_A, η_A)–space that satisfies $\Delta_{A|B}^{\neq} = 0$. From Equation (46), we derive that the coexistence curve is given by

$$\frac{\eta_A}{\beta_A} - \frac{\eta_B}{\beta_B} = 0 . \tag{47}$$

We validate this result using simulations of the well-mixed and vertex-based models in Section S6 of the supplementary material.

A nonzero heterotypic survival difference by itself does not prove that a competitive interaction, or indeed any interaction, is taking place. After all, two non-interacting cell types with different intrinsic survival probabilities would also result in a nonzero heterotypic survival difference. Hence, now we define a metric that quantifies the change in survival probability as a result of heterotypic interactions. In particular, we define the **homotypic survival difference** as

$$\Delta_{\overline{A|B}}^{=} \equiv \xi_{A|B} - \lambda_A \,, \quad \Delta_{\overline{B|A}}^{=} \equiv \xi_{B|A} - \lambda_B \,, \tag{48}$$

for cell types A and B, respectively. The homotypic survival difference compares the fitness of a cell type in a heterotypic environment to its fitness in a homotypic environment.

The sign of the homotypic survival difference indicates whether a cell type is more or less fit as a result of the heterotypic interaction, compared to homotypic conditions. If $\Delta_{\overline{A|B}}^{=} > 0$, then we say that cell type A is more fit when competing with cell type B, and vice versa for $\Delta_{\overline{A|B}}^{=} < 0$. A positive homotypic survival difference indicates that the cell type benefits from the interaction, but it does not mean that the interaction is mutualistic, since in that case both cell types would need to benefit from the interaction (i.e. $\Delta_{\overline{A|B}}^{=}, \Delta_{\overline{B|A}}^{=} > 0$). We show below that such an interaction is impossible in the G2 death signal model. Finally, if $\Delta_{\overline{A|B}}^{=} = 0$, then we say that cell type A is in **neutral competition** with cell type B, since the presence of cell type B does not produce a net change in the fitness of cell type A.

Focusing our derivation on the homotypic survival difference of cell type A, we apply the sign function to give

$$\mathrm{sgn}\left(\Delta_{\overline{A|B}}^{=}\right) = \mathrm{sgn}\left(\frac{1}{\langle 1 - \beta \rangle} - \frac{1}{1 - \beta_A}\right) \,. \tag{49}$$

We expand $\langle 1 - \beta \rangle(t)$ to give

$$\frac{1}{\langle 1 - \beta \rangle(t)} - \frac{1}{1 - \beta_A} =$$

$$\frac{n_B(t)(\beta_B - \beta_A)}{[n_A(t)(1 - \beta_A) + n_B(t)(1 - \beta_B)]\,(1 - \beta_B)}. \quad (50)$$

506 The denominator on the right-hand side is strictly positive, so we only need to consider

507 the sign of the numerator. Equation (50) indicates that the sign of $\Delta_{A|B}^=$ is dependent

508 on the system state. In the degenerate case of $n_B(t) = 0$, we are reduced to a homotypic

509 population composed solely of A-type cells, and thus $\Delta_{A|B}^= = 0$. However, if we limit

510 our scope to the heterotypic case, i.e. $n_A(t), n_B(t) > 0$, we can rewrite Equation (49) as

$$\text{sgn}\left(\Delta_{A|B}^=\right) = \text{sgn}(\beta_B - \beta_A). \quad (51)$$

511 For cell type B, we derive an analogous expression:

$$\text{sgn}\left(\Delta_{B|A}^=\right) = \text{sgn}(\beta_A - \beta_B). \quad (52)$$

512 Comparing Equations (51) and (52), we derive the following identity:

$$\text{sgn}\left(\Delta_{A|B}^=\right) = -\text{sgn}\left(\Delta_{B|A}^=\right). \quad (53)$$

In other words, the homotypic survival differences of two competing cell types have

opposite signs. Hence, one cell type's loss is another cell type's gain, and a mutualistic

relationship is impossible.

For the heterotypic survival difference, we factored out the death signal, $\langle 1 - \beta \rangle$, to

find an expression for the sign of $\Delta_{A|B}^{\neq}$ and found that winner/loser status is determined

by the difference in *tolerance* to death signals. Here, in contrast, we factored out the

tolerance to death signals, η/β, to find that the sign of the homotypic survival difference

depends on the difference in β. In the ergodic approximation, a larger value of $1 - \beta$

corresponds to a greater death signal. This is because the amount of time spent in G2

phase, during which cells emit death signals, is proportional to $1 - \beta$, which suggests

that we can interpret $1 - \beta$ as the cell's emission rate of death signals. Rewriting

Equation (51) as

$$\text{sgn}\left(\Delta_{A|B}^=\right) = \text{sgn}\left((1 - \beta_A) - (1 - \beta_B)\right), \quad (54)$$

$\Delta^{\neq}_{A\|B}$ \\ $\Delta^{=}_{A\|B}$	+	0	−
+	A direct winner B direct loser	A neutral winner B neutral loser	A indirect winner B indirect loser
0	Coexistence	Neutral coexistence	Coexistence
−	A indirect loser B indirect winner	A neutral loser B neutral winner	A direct loser B direct winner

Table 3: Classification of competitive interactions based on the heterotypic survival difference, $\Delta^{\neq}_{A|B}$, and the homotypic survival difference, $\Delta^{=}_{A|B}$.

shows that the sign of the homotypic survival difference is determined by the difference in *emission* of death signals. In particular, cell type A fares better in heterotypic conditions if cell type B has a lower emission of death signal than cell type A, and vice versa.

Equations (51) and (52) show that the signs of $\Delta^{=}_{A|B}$ and $\Delta^{=}_{B|A}$ are independent of the system state, except in the degenerate homotypic cases $n_A(t) = 0$ and $n_B(t) = 0$. We can therefore partition the parameter space into two regions: one where $\Delta^{=}_{A|B} > 0 \wedge \Delta^{=}_{B|A} < 0$, and one where $\Delta^{=}_{A|B} < 0 \wedge \Delta^{=}_{B|A} > 0$. We define the **neutral competition curve** as the curve in (β_A, η_A)–space that satisfies $\Delta^{=}_{A|B} = 0$ for fixed values of β_B and η_B. From Equation (51), we derive that the neutral competition curve is given by

$$\beta_B - \beta_A = 0 \,. \tag{55}$$

We validate this result using simulations of the well-mixed and vertex-based models in Section S7 of the supplementary material.

 4.2.1. Classification of competitive interactions

In this section, we combine the results for the heterotypic and homotypic survival

differences to classify different types of competitive interactions. There are nine different

types of competitive interactions (Table 3). We can group these outcomes by considering

their symmetry. Given that $\Delta^{\neq}_{A|B} = -\Delta^{\neq}_{B|A}$, we can group the nine types of competitive

interactions into five distinct categories:

Neutral coexistence $\left\{\Delta^{\neq}_{A|B} = 0, \ \Delta^{=}_{\overline{A}|B} = 0\right\}$. This is the degenerate case where nei-

ther cell type has a relative survival advantage, and both cell types have the same

survival probability as in homotypic conditions. The competitive interaction is

neutral because there is no effect on either cell type's absolute fitness, and the

cell types coexist because they have the same fitness.

Coexistence $\left\{\Delta^{\neq}_{A|B} = 0, \ \Delta^{=}_{\overline{A}|B} \neq 0\right\}$. The cells experience a change in absolute fit-

ness compared to the homotypic environment, but there is no relative survival

advantage for either cell type. Therefore, neither cell type dominates.

Neutral competition $\left\{\Delta^{\neq}_{A|B} \neq 0, \ \Delta^{=}_{\overline{A}|B} = 0\right\}$. The nonzero heterotypic survival dif-

ference means that there is a difference in relative fitness. Thus, winners and

losers emerge, with the winner cell type dominating the population. However,

neither cell type experiences a difference in absolute fitness compared to homo-

typic conditions.

Indirect competition $\left\{\Delta^{\neq}_{A|B} \neq 0, \ \text{sgn}\left(\Delta^{\neq}_{A|B}\right) \neq \text{sgn}\left(\Delta^{=}_{\overline{A}|B}\right) \neq 0\right\}$. As in neutral com-

petition, winners and losers emerge from the competitive interaction. The sign of

the homotypic survival difference is nonzero and opposite to the sign of the het-

erotypic survival difference, which means that the losers experience an increase in

absolute fitness compared to homotypic conditions, and the winners experience a

decrease.

Direct competition $\left\{\Delta^{\neq}_{A|B} \neq 0, \ \text{sgn}\left(\Delta^{\neq}_{A|B}\right) = \text{sgn}\left(\Delta^{=}_{\overline{A}|B}\right)\right\}$. Similar to the other

types of competition, the population splits into winner and loser cells. In contrast

Figure 4: Diagram situating the different types of competitive interactions in (β_A, η_A)–space, given fixed values for β_B and η_B. The full and dashed lines correspond to the coexistence and neutral competition curves, respectively. The green dot corresponds to the neutral coexistence point.

to indirect competition, however, the homotypic survival difference has the same sign as the heterotypic survival difference, meaning that the winners are fitter than in the homotypic environment, and the losers less fit.

All types of competition involve one cell type (the winners) becoming more abundant than the other cell type (the losers). The distinction between them is based on the change in fitness experienced by the winners and losers compared to homotypic conditions. In neutral competition, there is no change in fitness for either the winners or losers. In indirect competition, the winners become less fit and the losers more fit, potentially leading to a scenario where a previously nonviable loser cell type is "rescued" by the interaction with the winner cell type and becomes viable. In direct competition, the winners become more fit and the losers less fit, potentially leading to a previously viable loser cell type becoming nonviable as a result of the interaction. We therefore expect any competitive outcomes to fall under this category.

As discussed previously, we can partition cross sections of parameter space using the coexistence curve and the neutral competition curve. In Figure 4, we plot these curves in (β_A, η_A)–space for fixed values of β_B and η_B. The curves translate to straight

₅₈₀ lines, on which we find the coexistence and neutral competition regimes. Furthermore,

₅₈₁ we find the **neutral coexistence point** at their intersection, $\beta_A = \beta_B, \eta_A = \eta_B$,

₅₈₂ which corresponds to the degenerate case where the competing cell types have identical

₅₈₃ parameters. Finally, we see that the curves divide the cross section into four sectors,

₅₈₄ with the top left and bottom right sectors corresponding to direct competition, and the

₅₈₅ top right and bottom left sectors corresponding to indirect competition.

₅₈₆ *4.2.2. Heterotypic proliferation regimes*

₅₈₇ Earlier, we defined winners and losers in a weak sense based on which cell type

₅₈₈ becomes more abundant. Although this is an important precondition for cell compe-

₅₈₉ tition, the definition of winners and losers in the cell competition criteria is based on

₅₉₀ the viability of the competing cell types. In this section, we investigate the viability of

₅₉₁ winners and losers.

₅₉₂ Regardless of the type of competitive interaction, winners become the dominant

₅₉₃ species in the population over time by definition. Therefore, we expect that the

₅₉₄ population-weighted average death signal, $\langle 1 - \beta \rangle(t)$, approaches the intrinsic death

₅₉₅ signal of the winner cell type. Assuming for now that cell type A is the winner, i.e.

₅₉₆ $\Delta_{A|B}^{\neq} > 0$, we have

$$\langle 1 - \beta \rangle(t) \to 1 - \beta_A \quad \text{as} \quad t \to \infty \,. \tag{56}$$

Hence, when considering the long-term behaviour of the population, we can substitute
$1 - \beta_A$ for $\langle 1 - \beta \rangle$ into the heterotypic survival probability for cell types A and B to
obtain the **asymptotic survival probabilities**:

$$\xi_{A|B}(t \to \infty) = 1 - \exp\left(-\frac{\eta_A}{\beta_A(1 - \beta_A)} \right) \,, \tag{57}$$

$$\xi_{B|A}(t \to \infty) = 1 - \exp\left(-\frac{\eta_B}{\beta_B(1 - \beta_A)} \right) \,. \tag{58}$$

₅₉₇ Comparing Equation (57) with Equation (30), we find that the asymptotic survival

₅₉₈ probability of cell type A is equal to its homotypic survival probability, λ_A. The het-

₅₉₉ erotypic viability of winners is thus determined by their homotypic viability. We denote

the right-hand side of Equation (58) as

$$\xi_{B|A}^{\infty} \equiv 1 - \exp\left(-\frac{\eta_B}{\beta_B(1-\beta_A)}\right), \tag{59}$$

so that we can write the asymptotic survival probabilities more succinctly as

$$\xi_{A|B}(t \to \infty) = \lambda_A, \tag{60}$$

$$\xi_{B|A}(t \to \infty) = \xi_{B|A}^{\infty}. \tag{61}$$

Conversely, if cell type B is the winner, i.e. $\Delta_{A|B}^{\neq} < 0$, we have

$$\xi_{B|A}(t \to \infty) = \lambda_B, \tag{62}$$

$$\xi_{A|B}(t \to \infty) = \xi_{A|B}^{\infty}, \tag{63}$$

where $\xi_{A|B}^{\infty}$ is defined analogously to Equation (59).

We use the asymptotic survival probability to characterise the viability of competing cell types in a heterotypic population. Assuming that cell type A is the winner, we distinguish between the following outcomes:

Case $\left\{\lambda_A \leq \frac{1}{2}\right\}$. If the winner cells are not viable, then the losers are also not, since they have, by definition, a lower survival probability than the winners. Hence, both winners and losers go extinct.

Case $\left\{\lambda_A > \frac{1}{2}\right\}$. The winner cells are homotypically viable and therefore remain viable. Whether or not the losers are viable depends on $\xi_{B|A}^{\infty}$.

Subcase $\left\{\xi_{B|A}^{\infty} \leq \frac{1}{2}\right\}$. The loser cells are heterotypically nonviable and are eliminated from the tissue.

Subcase $\left\{\xi_{B|A}^{\infty} > \frac{1}{2}\right\}$. The losers are heterotypically viable and persist in the tissue.

We thus have three distinct proliferation regimes for $\Delta_{A|B}^{\neq} > 0$. Three analogous proliferation regimes exist for $\Delta_{A|B}^{\neq} < 0$, for a total of six proliferation regimes overall. We

cannot visualise four-dimensional $(\beta_A, \eta_A, \beta_B, \eta_B)$–space directly, so we first provide an outline of the proliferation regimes, and then sketch them in cross sections for particular values of β_B and η_B.

Firstly, the **coexistence hypersurface** $\Delta^{\neq}_{A|B} = 0$ divides the parameter space into two subspaces, $\Delta^{\neq}_{A|B} > 0$ and $\Delta^{\neq}_{A|B} < 0$, where cell types A and B are the respective winners. Secondly, for $\Delta^{\neq}_{A|B} > 0$, we have two regions where $\lambda_A > 1/2$ and $\lambda_A < 1/2$, respectively. The boundary is given by the **A winner viability hypersurface** $\lambda_A = 1/2$. The region in which the winner is viable, i.e. $\lambda_A > 1/2$, is further split into two parts, based on whether the loser is viable ($\xi^{\infty}_{B|A} > 1/2$) or nonviable ($\xi^{\infty}_{B|A} < 1/2$), by the **B loser viability hypersurface** $\xi^{\infty}_{B|A} = 1/2$. We divide the subspace $\Delta^{\neq}_{A|B} < 0$, where cell type B is the winner, in an analogous manner. Hence, in total we have five hypersurfaces that delineate the heterotypic proliferation regimes: the coexistence hypersurface, two winner viability hypersurfaces and two loser viability hypersurfaces.

We visualise the heterotypic proliferation regimes using cross sections for particular values of β_B and η_B in (β_A, η_A)–space. In these cross sections, the hypersurfaces become the following curves:

$$\text{Coexistence curve:} \quad \Delta^{\neq}_{A|B} = 0 \Leftrightarrow \eta_A = \frac{\eta_B}{\beta_B}\beta_A \,, \tag{64}$$

$$\text{A winner curve:} \quad \lambda_A = \frac{1}{2} \Leftrightarrow \eta_A = \ln(2)\beta_A(1 - \beta_A) \,, \tag{65}$$

$$\text{B loser curve:} \quad \xi^{\infty}_{B|A} = \frac{1}{2} \Leftrightarrow \beta_A = 1 - \frac{\eta_B}{\ln(2)\beta_B} \,, \tag{66}$$

$$\text{A loser curve:} \quad \xi^{\infty}_{A|B} = \frac{1}{2} \Leftrightarrow \eta_A = \ln(2)(1 - \beta_B)\beta_A \,. \tag{67}$$

The B winner viability hypersurface does not map onto a curve in (β_A, η_A)–space because it depends only on β_B and η_B. We therefore consider the cases $\lambda_B < 1/2$ and $\lambda_B > 1/2$ in separate cross sections.

If $\eta_B/\beta_B > \ln(2)$, then Equation (66) does not have a solution for positive β_A, so the B loser viability curve is not present in cross sections for which this is the case. We therefore consider this case in a separate cross section. It can be easily verified

Figure 5: Heterotypic proliferation regimes: diagram and well-mixed results. (a) Diagrams for Cross Sections I, II, and III, situating the different heterotypic proliferation regimes. The green dot corresponds to the neutral coexistence point. Grey: cell types A and B are nonviable. Green: cell type A is nonviable, cell type B is viable. Orange: cell type A is viable, cell type B is nonviable. Red: cell types A and B are viable. (b) Estimated heterotypic survival frequency of cell types A and B using the well-mixed model. The top row displays the estimated heterotypic survival frequency of cell type A, $\overline{\xi_{A|B}}$, defined in Equation (68). The bottom row displays the estimated heterotypic survival frequency of cell type B, $\overline{\xi_{B|A}}$, also defined in Equation (68). All curves are the same as in (a).

that $\eta_B/\beta_B > \ln(2)$ implies $\lambda_B > 1/2$, so we only need to consider three distinct cross sections (Figure 5(a)):

Cross Section I $\{\beta_B = 0.2, \eta_B = 0.2\}$. This cross section satisfies $\eta_B/\beta_B > \ln(2)$. We see three distinct regimes. Above the coexistence curve, both cell types are viable, with cell type A as the winner. Between the coexistence curve and the A loser viability curve, cell type B is the winner and both cell types are viable. Below the A loser viability curve, only cell type B is viable. We note that there are no values of β_A and η_A for which cell type B is nonviable. Therefore, regardless of the competing cell type, cell type B is always viable.

Cross Section II $\{\beta_B = 0.8, \eta_B = 0.2\}$. This cross section satisfies $\eta_B/\beta_B < \ln(2)$ and $\lambda_B > 1/2$. We identify five distinct regimes. Below the coexistence curve, we see the same two regimes as in Cross Section I. The wedge-shaped region between the coexistence curve and the A loser viability curve is particularly interesting because it partly overlaps with the area under the homotypic viability curve of cell type A. The A-type cells in this region are nonviable under homotypic conditions, but are viable when interacting with cell type B and are therefore "rescued" by the competitive interaction. This is also present in Cross Section I, but it is more visible here. We note that this region is contained within the indirect competition sector because only an indirect competitive interaction can increase the fitness of loser cells.

We see three regimes above the coexistence curve. Below the A winner viability curve, the winning A-type cells are nonviable, which renders both cell types nonviable. Above this curve, the winner A-type cells are viable. In this subspace, the survival of cell type B depends on β_A. To the left of the B loser viability curve, the death signal emitted by cell type A is sufficiently high to eliminate cell type B, whereas, on the other side, the death signal is not strong enough to eliminate cell type B, so cell type B survives.

Cross Section III $\{\beta_B = 0.4, \eta_B = 0.1\}$. This cross section satisfies $\eta_B/\beta_B < \ln(2)$ and $\lambda_B < 1/2$. Below the coexistence curve, where cell type B is the winner, both cell types

are nonviable because cell type B is homotypically nonviable. Above the coexistence curve, we find the same regimes as in Cross Section II. Since cell type B is homotypically nonviable in this cross section, we note that the top right triangular region, where cell type B is heterotypically viable, corresponds to nonviable loser rescue, and thus is analogous to the wedge-shaped areas discussed in Cross Section II. Again, this area is fully contained within the indirect competition sector.

4.2.3. Computational validation of heterotypic proliferation regimes

In this section, we validate the predicted heterotypic proliferation regimes by conducting simulations of the well-mixed and vertex-based models. For the vertex-based model, we conducted simulations with both segregated and random initial conditions. Further details are provided in Section S8 of the supplementary material.

To estimate the survival frequency for a particular parameter set, we averaged the heterotypic survival frequencies across repeated simulations as

$$\overline{\xi_{A|B}} = \frac{1}{N_{\text{sim}}} \sum_{k=1}^{N_{\text{sim}}} \hat{\xi}_{A|B,k} \,, \quad \overline{\xi_{B|A}} = \frac{1}{N_{\text{sim}}} \sum_{k=1}^{N_{\text{sim}}} \hat{\xi}_{B|A,k} \,. \tag{68}$$

The results for the well-mixed model are given in Figure 5(b). The top and bottom rows show the survival frequency for cell types A and B, respectively. When compared to Figure 5(a), we see an excellent agreement between the simulations and predictions.

The results for the vertex-based model with random and segregated initial conditions are provided in Figures 6(a) and 6(b), respectively. In Figure 6(a), we can see roughly similar proliferation regimes as in the well-mixed case, but the contours do not align perfectly with the predicted curves. Moreover, in Cross Section II, for cell type A, we expect to see a sequence of red–blue–red–blue regions from top left to bottom right, but instead we see a gradual transition from red to blue. In addition, for high β_A, we see red regions for cell type A that extend below their predicted limits in all cross sections.

In Figure 6(b), we see significant deviations from the predicted proliferation regimes. When comparing the plots for cell type A with the results for the homotypic proliferation regimes in Figure 3(b), we see that A-type cells essentially behave as if they were in a

Figure 6: Heterotypic proliferation regimes: vertex-based results. (a–b) Estimated heterotypic survival frequency of cell types A and B using the vertex-based model with random and segregated initial conditions. See Figure 5(b) for legend. (a) Random initial conditions. (b) Segregated initial conditions.

Figure 7: Diagrams of competition regimes for Cross Sections I and II. The green dot corresponds to the neutral coexistence point. The labels W and L are used to refer to the winner and loser cell types, respectively, i.e. W = A, L = B for $\Delta_{A|B}^{\neq} > 0$ and W = B, L = A for $\Delta_{A|B}^{\neq} < 0$. The symbol $\xi_{L|W}^{\infty}$ refers to the asymptotic survival probability of the loser cell type. Linear hatch: homotypic viability. Cross hatch: cell competition. Red: complete cell competition. Orange: incomplete cell competition. Green: indirect cell competition. See also Table 4 for the legend.

homotypic environment. Similarly, the heterotypic viability of cell type B corresponds to its viability in homotypic conditions, regardless of the parameters of cell type A. These results suggest that the segregated cell types behave like homotypic populations.

4.3. Classification of competition regimes

Now that we have characterised the proliferation regimes of homotypic and heterotypic populations, and classified different types of competitive interactions, we combine these classifications to identify different competition regimes in the G2 death signal model. The first condition of the cell competition criteria is that both cell types are homotypically viable, i.e. $\lambda_A, \lambda_B > 1/2$. In order to satisfy $\lambda_A > 1/2$, we only consider the parameter space above the homotypic viability curve, as shown in Figure 7. To satisfy the viability condition for cell type B, we only consider cross sections that satisfy $\lambda_B > 1/2$. In particular, Cross Section III does not satisfy this condition, so we only plot Cross Sections I and II in Figure 7. We define the **homotypic viability regime**

Regime	λ_W, λ_L	$\Delta^{\neq}_{W|L}$	$\xi^{\infty}_{L|W}$	$\Delta^{=}_{\bar{L}|W}$	Legend
Homotypic viability	$> 1/2$	-	-	-	╲╲╲
Coexistence	$> 1/2$	$= 0$	-	-	▬
Competition	$> 1/2$	> 0	-	-	-
Loser elimination	$> 1/2$	> 0	$\leq 1/2$	$\underline{< 0}$	-
Loser survival	$> 1/2$	> 0	$> 1/2$	-	-
Cell competition	$> 1/2$	> 0	-	< 0	⊠⊠
Complete cell competition	$> 1/2$	> 0	$< 1/2$	$\underline{\leq 0}$	▮
Critical cell competition	$> 1/2$	> 0	$= 1/2$	$\underline{\leq 0}$	▬
Incomplete cell competition	$> 1/2$	> 0	$> 1/2$	< 0	▮
Neutral competition	$> 1/2$	> 0	$\underline{\geq 1/2}$	$= 0$	▬ ▬
Indirect competition	$> 1/2$	> 0	$\underline{\geq 1/2}$	> 0	▮

Table 4: Classification of competition regimes. The competition regime (bolded) can be subdivided in two ways: loser elimination and loser survival regimes (top section), or cell competition, neutral competition, and indirect competition regimes (bottom section). The underlined conditions are implied by the other conditions on the same row. The legend column maps the regimes onto areas and curves plotted in Figure 7.

 as

$$\lambda_A, \lambda_B > \frac{1}{2}\,. \tag{69}$$

The second condition is that only one cell type remains viable when the cell types compete. This implies a nonzero heterotypic survival difference, i.e. $\Delta^{\neq}_{A|B} \neq 0$, splitting the homotypic viability regime into the **coexistence regime**

$$\lambda_A, \lambda_B > \frac{1}{2} \quad \wedge \quad \Delta^{\neq}_{A|B} = 0\,, \tag{70}$$

and the **competition regime**

$$\lambda_A, \lambda_B > \frac{1}{2} \quad \wedge \quad \Delta^{\neq}_{A|B} \neq 0\,. \tag{71}$$

The competition regime is further subdivided according to which cell type is the winner. The G2 death signal model is symmetric with respect to swapping cell type labels, so the choice of winner or loser is arbitrary. Therefore, for ease of notation, we henceforth label the winner cell type with W and the loser cell type with L, such that $\Delta^{\neq}_{W|L} > 0$ is true by construction.

As we saw in Section 4.2.2, the viability of the winner cell type is determined by its homotypic viability, which is guaranteed by Equation (69). Therefore, we only need to impose further that the loser cell type is heterotypically nonviable, i.e. $\xi^{\infty}_{L|W} \leq 1/2$. We define the **loser elimination regime** as

$$\lambda_W, \lambda_L > \frac{1}{2} \quad \wedge \quad \Delta^{\neq}_{W|L} > 0 \quad \wedge \quad \xi^{\infty}_{L|W} \leq \frac{1}{2}\,, \tag{72}$$

and the **loser survival regime** as

$$\lambda_W, \lambda_L > \frac{1}{2} \quad \wedge \quad \Delta^{\neq}_{W|L} > 0 \quad \wedge \quad \xi^{\infty}_{L|W} > \frac{1}{2}\,. \tag{73}$$

The loser elimination regime satisfies the cell competition criteria and is non-empty. We therefore conclude that the G2 death signal model is capable of producing competitive outcomes.

We can further refine the competition regimes by considering, in addition, the type of competitive interaction. Figure 7 shows that the neutral competition curve, defined by

$\Delta_{\overline{L}|W}^{=} = 0$, runs through the loser survival regime. We define the **neutral competition**
regime as

$$\lambda_W, \lambda_L > \frac{1}{2} \quad \wedge \quad \Delta_{W|L}^{\neq} > 0 \quad \wedge \quad \Delta_{\overline{L}|W}^{=} = 0 \,. \tag{74}$$

The neutral competition curve separates the loser survival regime into two subregimes
where $\Delta_{\overline{L}|W}^{=} < 0$ and $\Delta_{\overline{L}|W}^{=} > 0$, respectively. In the case of $\Delta_{\overline{L}|W}^{=} < 0$, the fitness of
losers is reduced by the winners, but not enough to cause loser elimination. We define
this as the **incomplete cell competition regime**

$$\lambda_W, \lambda_L > \frac{1}{2} \,\wedge\, \Delta_{W|L}^{\neq} > 0 \,\wedge\, \xi_{L|W}^{\infty} > \frac{1}{2} \,\wedge\, \Delta_{\overline{L}|W}^{=} < 0 \,. \tag{75}$$

In addition, we can partition the loser elimination regime into the **complete cell**
competition regime

$$\lambda_W, \lambda_L > \frac{1}{2} \quad \wedge \quad \Delta_{W|L}^{\neq} > 0 \quad \wedge \quad \xi_{L|W}^{\infty} < \frac{1}{2} \,, \tag{76}$$

and the **critical cell competition regime**

$$\lambda_W, \lambda_L > \frac{1}{2} \quad \wedge \quad \Delta_{W|L}^{\neq} > 0 \quad \wedge \quad \xi_{L|W}^{\infty} = \frac{1}{2} \,, \tag{77}$$

which is the threshold regime between complete and incomplete cell competition. The
common feature of complete, critical, and incomplete cell competition is that the win-
ners negatively impact the losers. We group these regimes under the **cell competition**
regime

$$\lambda_W, \lambda_L > \frac{1}{2} \quad \wedge \quad \Delta_{W|L}^{\neq} > 0 \quad \wedge \quad \Delta_{\overline{L}|W}^{=} < 0 \,. \tag{78}$$

Finally, on the other side of the neutral competition curve we have $\Delta_{\overline{L}|W}^{=} > 0$, where
loser cells have a higher fitness than in homotypic conditions. We denote this as the
indirect competition regime

$$\lambda_W, \lambda_L > \frac{1}{2} \quad \wedge \quad \Delta_{W|L}^{\neq} > 0 \quad \wedge \quad \Delta_{\overline{L}|W}^{=} > 0 \,. \tag{79}$$

We plot the competition regimes in Figure 7 for Cross Sections I and II, and summarise
them in Table 4.

The competition regimes let us discriminate between different types of winners and losers. We define *complete winners*, *critical winners*, *incomplete winners*, *neutral winners*, and *indirect winners* as the winner cell types in the respective competition regimes, and define different types of losers analogously. In this terminology, complete and critical winners and losers correspond to the classical definition of winners and losers in cell competition.

5. Discussion

We stated in the introduction that there are two important advantages in treating winner/loser status as an emergent property rather than hardcoded identities: (i) we can test whether a given cell-based model is capable of producing competitive outcomes; and (ii), if so, analyse the conditions that give rise to competitive outcomes in that model. We demonstrated the first capability in Section 2 by showing that differences in mechanical properties alone (i.e. without a mechanism for active cell death) are insufficient to generate competitive outcomes in a vertex-based model of an epithelial tissue, which agrees with experimental observations that cell competition depends on the initiation of cell death in loser cells [26]. This result motivated our decision to propose a modelling framework for cell competition with an active mechanism of cell death in Section 3. We then introduced the G2 death signal model in Section 4 and thoroughly studied its behaviour for homotypic and heterotypic populations, identifying the conditions and parameters that lead to competitive outcomes. In the rest of this section, we will interpret and discuss these results, propose specific ideas for novel cell competition experiments, and outline potential future research directions.

Spatial mixing is required for cell competition. In Section 4.2.3, we observed that the occurrence of competitive outcomes in the vertex-based model depends on the initial spatial patterning of cell types. When the cell types are distributed randomly, we observe competitive outcomes, but when they are segregated, we do not. In fact, the behaviour of the segregated cell types is virtually identical to that of isolated homotypic

₇₆₇ populations. This result agrees with experimental observations that spatial mixing is
₇₆₈ required for cell competition [40], and has been replicated in other cell-based models of
₇₆₉ cell competition [13].

₇₇₀ Our derivation of heterotypic proliferation regimes is based on the assumption that
₇₇₁ the population is well-mixed, which is only true *locally* at heterotypic clone boundaries
₇₇₂ in the vertex-based model, where cells sample the death signal of both cell types. Within
₇₇₃ clones, however, cells interact only with cells of the same type, so they behave more
₇₇₄ similar to a homotypic population. The degree of competition therefore depends on the
₇₇₅ amount of heterotypic contact between cell types, which is determined by the level of
₇₇₆ spatial mixing.

₇₇₇ *Tolerance and emission.* When deriving the heterotypic survival difference, we found
₇₇₈ that the relative abundance of cell types in a tissue is determined by their tolerance to
₇₇₉ death signals (i.e. η/β). Furthermore, when deriving the homotypic survival difference,
₇₈₀ we showed that the difference in death signal emission (i.e. $1-\beta$) between two competing
₇₈₁ cell types determines the impact of the heterotypic interaction compared to homotypic
₇₈₂ conditions. Also, in Section 4.2.2, we demonstrated that loser elimination depends
₇₈₃ on the relationship between the tolerance of the loser and the emission of the winner.
₇₈₄ From these observations, we infer that tolerance to, and emission of, death signals are
₇₈₅ the fundamental cell properties driving cell competition in the G2 death signal model.
₇₈₆ Here, we present a transformation of parameters that explicitly describes the behaviour
₇₈₇ of the model in terms of tolerance and emission. We also show that the transformed
₇₈₈ parameters allow us to describe the competition regimes using intuitive and elegant
₇₈₉ expressions.

₇₉₀ We define the **tolerance** and **emission** of cell type X, respectively denoted $\tilde{\eta}_X$ and
₇₉₁ d_X, as follows:

$$\tilde{\eta}_X \equiv \frac{\eta_X}{\ln(2)\beta_X}, \quad d_X \equiv 1 - \beta_X. \tag{80}$$

₇₉₂ We can formulate the **homotypic viability** condition, $1/2 < \lambda_X$, using $\tilde{\eta}_X$ and d_X by

substituting the homotypic survival probability (Equation (30)) and rearranging:

$$\frac{1}{2} < \lambda_X \quad \Leftrightarrow \quad 1 - \beta_X < \frac{\eta_X}{\ln(2)\beta_X} \quad \Leftrightarrow \quad d_X < \tilde{\eta}_X \,. \tag{81}$$

The last inequality reads as the condition that cells must have a higher tolerance than emission to be homotypically viable. The biological interpretation is that cells must be capable of tolerating the death signal that they themselves emit in order to survive on their own. The **loser elimination** condition, $\xi_{L|W}^{\infty} < 1/2$, can also be expressed using tolerance and emission. Denoting the winner and loser cell types using the labels W and L, respectively, we substitute the asymptotic survival probability of the loser (Equation (59)) to obtain

$$\xi_{L|W}^{\infty} < \frac{1}{2} \quad \Leftrightarrow \quad \frac{\eta_L}{\ln(2)\beta_L} < 1 - \beta_W \quad \Leftrightarrow \quad \tilde{\eta}_L < d_W \,. \tag{82}$$

This means that the winner cells must emit death signals at a rate that the loser cells cannot tolerate in order to eliminate the loser cell type from the tissue.

To satisfy the cell competition criteria, we require that both cell types are homotypically viable, i.e. $d_W < \tilde{\eta}_W$ and $d_L < \tilde{\eta}_L$, and that the loser is eliminated, i.e. $\tilde{\eta}_L < d_W$. Combining these expressions, we can summarise the conditions on model parameters that satisfy the cell competition criteria in a single statement:

$$d_L < \tilde{\eta}_L < d_W < \tilde{\eta}_W \,, \tag{83}$$

which can be read as

loser emission < loser tolerance <

winner emission < winner tolerance . (84)

This corresponds to the **complete cell competition** regime that we defined in Section 4.3. In a similar manner, we can express all competition regimes in terms of tolerance and emission:

Cell competition: $d_L < \tilde{\eta}_L < \tilde{\eta}_W \wedge d_L < d_W$.

Figure 8: Diagram of competition regimes using the transformed parameters $\tilde{\eta}_X$ and d_X, defined in Equation (80). The green dot corresponds to the neutral coexistence point. The same conventions apply as in Figure 7. See Table 4 for the legend.

Complete cell competition: $d_L < \tilde{\eta}_L < d_W < \tilde{\eta}_W$.

Critical cell competition: $d_L < \tilde{\eta}_L = d_W < \tilde{\eta}_W$.

Incomplete cell competition: $d_L < d_W < \tilde{\eta}_L < \tilde{\eta}_W$.

Neutral competition: $d_L = d_W < \tilde{\eta}_L < \tilde{\eta}_W$.

Indirect competition: $d_W < d_L < \tilde{\eta}_L < \tilde{\eta}_W$.

These relationships can be verified visually in Figure 8, which shows the competition regimes in transformed parameter space.

The tolerance–emission model of cell competition. Based on Equation (84), we make the following biological prediction: **cell competition requires that winner cells have a higher tolerance to death signals *and* a higher rate of death signal emission than loser cells**. The implicit assumption in this statement is that cells emit and tolerate some form of death signal, which can be contact-based, ligand-based, mechanical stress-based, etc. Intuitively, regardless of the type of death signal, winners must expose losers to a sufficiently high level of death signal to eliminate them, while still being able to withstand it themselves.

Importantly, this model implies that mutations resulting in cell competition, such as *Minutes* and *Myc*, are pleiotropic because they simultaneously alter the tolerance to, and emission of, death signals. As a corollary, mutations which affect only one or neither, do *not* engender cell competition. This potentially explains why some mutations related to proliferation rates result in cell competition, and others do not [7]. In this view, the inhibition of apoptosis can be seen as a mutation that results in an infinite tolerance, without affecting emission. Indeed, it has been shown in experiments that inhibiting apoptosis prevents cell competition [7, 26].

This observation raises the question: do mutations exist that increase the emission of death signals, without affecting tolerance? If so, they would be challenging to culture, since such mutants would not tolerate their own death signal and thus be homotypically nonviable. However, the tolerance–emission model suggests that such a mutation would be viable if it were paired with apoptosis inhibition. Our model therefore predicts that a hypothetical emission-enhancing mutation combined with apoptosis inhibition would result in a new species of super-competitors.

Experimental support   Experimental evidence from *Myc*-based cell competition supports the tolerance–emission hypothesis. In [41], the authors demonstrated that the ligand Spätzle is necessary for the elimination of loser cells in the *Drosophila* wing disc, forming what the authors term a "killing signal". They also observed that Spätzle is produced in wild-type conditions at a rate that is tolerated by the wild-type cells, and that the production of Spätzle is upregulated in *Myc* super-competitors without inducing cell death in *Myc* mutants. *Myc* mutants therefore emit more death signal than wild-type cells, while simultaneously being less sensitive to it.

In this experiment, the death signal takes on the form of a diffusible death ligand. While the principles of the tolerance and emission should still apply in this case, there is one important difference; namely that death ligands can diffuse away from the site of heterotypic contact. This could potentially explain why we observe loser cell death at a distance in *Myc*-based cell competition [7], but not in *Minutes*-based cell competition.

According to our model, death ligand secretion is upregulated in mutant winner cells in the former case, and downregulated in mutant loser cells in the latter case.

We also find support for the tolerance–emission model in mechanical cell competition, specifically in cultures of Madine–Darby canine kidney (MDCK) cells [10]. The authors discovered that cell proliferation is in part modulated by the composition of cell types in the cellular neighbourhood. In particular, winner cells are more prolific when they are specifically surrounded by loser cells. This agrees with our observations that winner cells benefit from proximity to loser cells because the losers emit a lower level of death signal.

Experimental validation   To validate the tolerance–emission hypothesis, we must extrapolate the model predictions to experimental conditions that have not yet been tested. We predicted in Section 4.2.2 that homotypically nonviable loser cells can be rescued through indirect competition. This occurs when a winner cell type has a lower emission rate than the loser cell type, creating an environment in which losers can proliferate even if they are not viable on their own. The challenge in reproducing this result experimentally, however, is that we first need to identify an intrinsically nonviable mutant cell type. We therefore propose an alternative experiment that replicates this phenomenon with known cell types.

In particular, consider a triple co-culture where cell type A outcompetes cell type B and cell type B outcompetes cell type C. Cell types B and C are both eliminated in a background of cell type A, which mimics the intrinsic nonviability of cell types B and C. The tolerance–emission model predicts that the emission of death signals by cell type C is tolerated by cell type B. Therefore, if we inhibit apoptosis in cell type C, we expect to see: (i) C-type clones forming in an A-type background; and (ii) the survival of B-type cells exclusively inside the C-type clones. This outcome would be analogous to the rescue of a homotypically nonviable loser by indirect competition, with cell types B and C as the indirect losers and winners, respectively.

The function of cell competition. The prevalent hypothesis is that cell competition is a mechanism for maintaining tissue health by eliminating unfit cells. However, what is meant by "fitness" in this context is not clear [42]. The classical definition of fitness is based on *reproductive success* and early experiments linked this to cell competition, with winner cells having higher intrinsic proliferation rates than losers [43, 44]. However, not all mutations that increase proliferation rates result in cell competition [7]. In cell competition, fitness is perhaps more accurately defined as a measure of *competitive success*, which can determined by pairwise contests between cell types. In the tolerance–emission model, competitive success is a combination of tolerance and emission, and lacks a causal relationship with proliferation rates.

Competitive fitness is therefore not the same as reproductive fitness, but then why are they often linked in practice? We speculate that differential proliferation rates are not the mechanism of cell competition, but the *target* of cell competition. Cell competition evolved to optimise reproductive fitness, but uses competitive fitness as an imperfect means to communicate it. In other words, competitive fitness serves as a proxy for reproductive fitness, and it evolved in a trade-off with other factors such as the costs of cell competition.

Furthermore, we expect that the target of cell competition depends on the function of the host tissue. In the *Drosophila* wing disc, the tissue expands from 50 to 50 000 cells in the span of four days, so the aim of cell competition here is to optimise for reproductive fitness. MDCK cells, on the other hand, were derived from kidney tubules, where their function is to form a mechanically resilient barrier. In this case, cell competition is linked to mechanical cell compression, which we hypothesise acts as a proxy for the cell's ability to contribute to the structural integrity of the tissue.

Future work. The framework presented in this paper can be applied to any cell-based model to study hypothetical mechanisms of cell competition. Moreover, the cell competition criteria are sufficiently abstract that they can potentially be translated to models of cell competition that are not cell-based, such as Lotka–Volterra models [45].

We have emphasised that the death clock framework is agnostic with respect to the death signal, and that it can be used to represent different kinds of cell competition mechanisms. Of particular interest are diffusible ligands and mechanical compression as death signals. Research shows that cell competition in the *Drosophila* wing disc involves the use of diffusible death ligands [41, 46, 47]. A death clock model based on the secretion (i.e. emission) and recognition of death ligands is therefore an obvious next step toward a more biologically accurate representation of the cell competition process. Section 2 suggests that differences in mechanical properties alone do not generate competitive outcomes in a heterotypic vertex-based model. However, they may still play a role in cell competition when paired with an active mechanism for cell death. Research shows that mechanical compression triggers apoptosis in loser cells during mechanical cell competition [16]. Therefore, in this context, cell compression could be a suitable death signal. Further research is needed to examine models that incorporate diffusible ligands and mechanical compression as death signals.

6. Acknowledgements

The authors would like to acknowledge the use of the University of Oxford Advanced Research Computing (ARC) facility [48] and GNU Parallel [49] in carrying out this work. This work was supported by the Biotechnology and Biological Sciences Research Council (BBSRC) [BB/M011224/1].

7. Declaration of interests

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

References

[1] L. A. Johnston, Competitive interactions between cells: death, growth, and geography, Science 324 (5935) (2009) 1679–1682. `doi:10.1126/science.1163862`.

[2] S. Vivarelli, L. Wagstaff, E. Piddini, Cell wars: regulation of cell survival and proliferation by cell competition, Essays In Biochemistry 53 (2012) 69–82. `doi: 10.1042/bse0530069`.

[3] R. Levayer, E. Moreno, Mechanisms of cell competition: themes and variations, Journal of Cell Biology 200 (6) (2013) 689–698. `doi:10.1083/jcb.201301051`.

[4] J. P. Vincent, A. G. Fletcher, L. A. Baena-Lopez, Mechanisms and mechanics of cell competition in epithelia, Nature Reviews Molecular Cell Biology 14 (9) (2013) 581–591. `doi:10.1038/nrm3639`.

[5] M. Amoyel, E. A. Bach, Cell competition: how to eliminate your neighbours, Development 141 (5) (2014) 988–1000. `doi:10.1242/dev.079129`.

[6] S. Bowling, K. Lawlor, T. A. Rodríguez, Cell competition: the winners and losers of fitness selection, Development 146 (2019) dev167486. `doi:10.1242/dev.167486`.

[7] C. De La Cova, M. Abril, P. Bellosta, P. Gallant, L. A. Johnston, Drosophila myc regulates organ size by inducing cell competition, Cell 117 (1) (2004) 107–116. `doi:10.1016/S0092-8674(04)00214-4`.

[8] M. Norman, K. A. Wisniewska, K. Lawrenson, P. Garcia-Miranda, M. Tada, M. Kajita, H. Mano, S. Ishikawa, M. Ikegawa, T. Shimada, Y. Fujita, Loss of Scribble causes cell competition in mammalian cells, Journal of Cell Science 125 (1) (2012) 59–66. `doi:10.1242/JCS.085803`.

[9] E. Moreno, K. Basler, dMyc transforms cells into super-competitors, Cell 117 (1) (2004) 117–129. `doi:10.1016/S0092-8674(04)00262-4`.

[10] A. Bove, D. Gradeci, Y. Fujita, S. Banerjee, G. Charras, A. R. Lowe, Local cellular neighborhood controls proliferation in cell competition, Molecular Biology of the Cell 28 (23) (2017). `doi:10.1091/mbc.e17-06-0368`.

[11] S. W. Lee, Y. Morishita, Possible roles of mechanical cell elimination intrinsic to growing tissues from the perspective of tissue growth efficiency and homeostasis, PLoS Computational Biology 13 (7) (2017). `doi:10.1371/journal.pcbi.1005651`.

[12] A. Tsuboi, S. Ohsawa, D. Umetsu, Y. Sando, E. Kuranaga, T. Igaki, K. Fujimoto, Competition for space is controlled by apoptosis-induced change of local epithelial topology, Current Biology 28 (13) (2018). `doi:10.1016/j.cub.2018.05.029`.

[13] D. Gradeci, A. Bove, G. Vallardi, A. R. Lowe, S. Banerjee, G. Charras, Cell-scale biophysical determinants of cell competition in epithelia, eLife 10 (2021) e61011. `doi:10.7554/eLife.61011`.

[14] C. Brás-Pereira, E. Moreno, Mechanical cell competition, Current Opinion in Cell Biology 51 (2018) 15–21. `doi:10.1016/j.ceb.2017.10.003`.

[15] R. Levayer, C. Dupont, E. Moreno, Tissue crowding induces caspase-dependent competition for space, Current Biology 26 (5) (2016) 670–677. `doi:10.1016/j.cub.2015.12.072`.

[16] L. Wagstaff, M. Goschorska, K. Kozyrska, G. Duclos, I. Kucinski, A. Chessel, L. Hampton-O'Neil, C. R. Bradshaw, G. E. Allen, E. L. Rawlins, P. Silberzan, R. E. Salas, E. Piddini, Mechanical cell competition kills cells via induction of lethal p53 levels, Nature Communications 7 (2016). `doi:10.1038/ncomms11373`.

[17] E. Marinari, A. Mehonic, S. Curran, J. Gale, T. Duke, B. Baum, Live-cell delamination counterbalances epithelial growth to limit tissue overcrowding, Nature 484 (7395) (2012) 542–545. `doi:10.1038/nature10984`.

[18] G. T. Eisenhoffer, P. D. Loftus, M. Yoshigi, H. Otsuna, C. B. Chien, P. A. Morcos, J. Rosenblatt, Crowding induces live cell extrusion to maintain homeostatic cell numbers in epithelia, Nature 484 (7395) (2012) 546–549. `doi:10.1038/nature10999`.

[19] A. G. Fletcher, M. Osterfield, R. E. Baker, S. Y. Shvartsman, Vertex models of epithelial morphogenesis, Biophysical Journal 106 (11) (2014) 2291–2304. `doi:10.1016/j.bpj.2013.11.4498`.

[20] E. M. Purcell, Life at low Reynolds number, American Journal of Physics 45 (1) (1977) 3–11. `doi:10.1119/1.10903`.

[21] R. Farhadifar, J. C. Röper, B. Aigouy, S. Eaton, F. Jülicher, The influence of cell mechanics, cell-cell interactions, and proliferation on epithelial packing, Current Biology 17 (24) (2007) 2095–2104. `doi:10.1016/j.cub.2007.11.049`.

[22] A. G. Fletcher, J. M. Osborne, P. K. Maini, D. J. Gavaghan, Implementing vertex dynamics models of cell populations in biology within a consistent computational framework, Progress in Biophysics and Molecular Biology 113 (2) (2013) 299–326. `doi:10.1016/J.PBIOMOLBIO.2013.09.003`.

[23] J. A. Smith, L. Martin, Do cells cycle?, Proceedings of the National Academy of Sciences of the United States of America 70 (4) (1973) 1263–1267. `doi:10.1073/pnas.70.4.1263`.

[24] G. R. Mirams, C. J. Arthurs, M. O. Bernabeu, R. Bordas, J. Cooper, A. Corrias, Y. Davit, S.-J. Dunn, A. G. Fletcher, D. G. Harvey, M. E. Marsh, J. M. Osborne, P. Pathmanathan, J. Pitt-Francis, J. Southern, N. Zemzemi, D. J. Gavaghan, Chaste: an open source C++ library for computational physiology and biology, PLoS Computational Biology 9 (3) (2013) e1002970. `doi:10.1371/journal.pcbi.1002970`.

[25] J. M. Osborne, A. G. Fletcher, J. M. Pitt-Francis, P. K. Maini, D. J. Gavaghan, Comparing individual-based approaches to modelling the self-organization of multicellular tissues, PLoS Computational Biology 13 (2) (2017) e1005387. `doi:10.1371/journal.pcbi.1005387`.

[26] E. Moreno, K. Basler, G. Morata, Cells compete for decapentaplegic survival factor to prevent apoptosis in Drosophila wing development, Nature 416 (6882) (2002) 755–759. `doi:10.1038/416755a`.

[27] S. Legewie, N. Blüthgen, H. Herzel, Mathematical modeling identifies inhibitors of apoptosis as mediators of positive feedback and bistability, PLoS Computational Biology 2 (9) (2006) e120. `doi:10.1371/journal.pcbi.0020120`.

[28] M. Rehm, H. J. Huber, H. Dussmann, J. H. M. Prehn, Systems analysis of effector caspase activation and its control by X-linked inhibitor of apoptosis protein, The EMBO Journal 25 (18) (2006) 4338–4349. `doi:10.1038/sj.emboj.7601295`.

[29] M. Bentele, I. Lavrik, M. Ulrich, S. Stößer, D. W. Heermann, H. Kalthoff, P. H. Krammer, R. Eils, Mathematical modeling reveals threshold mechanism in CD95-induced apoptosis, Journal of Cell Biology 166 (6) (2004) 839–851. `doi:10.1083/jcb.200404158`.

[30] I. N. Lavrik, A. Golks, D. Riess, M. Bentele, R. Eils, P. H. Krammer, Analysis of CD95 threshold signaling: triggering of CD95 (FAS/APO-1) at low concentrations primarily results in survival signaling, Journal of Biological Chemistry 282 (18) (2007) 13664–13671. `doi:10.1074/jbc.M700434200`.

[31] K. L. King, J. A. Cidlowski, Cell cycle and apoptosis: common pathways to life and death, Journal of Cellular Biochemistry 58 (2) (1995) 175–180. `doi:10.1002/jcb.240580206`.

[32] K. L. King, J. A. Cidlowski, Cell cycle regulation and apoptosis, Annual Review of Physiology 60 (1) (1998) 601–617. `doi:10.1146/annurev.physiol.60.1.601`.

[33] B. Pucci, M. Kasten, A. Giordano, Cell cycle and apoptosis, Neoplasia 2 (4) (2000) 291–299. doi:10.1038/sj.neo.7900101.

[34] K. Vermeulen, Z. N. Berneman, D. R. Van Bockstaele, Cell cycle and apoptosis, Cell Proliferation 36 (3) (2003) 165–175. doi:10.1046/j.1365-2184.2003.00267.x.

[35] L. M. Facchini, L. Z. Penn, The molecular role of Myc in growth and transformation: recent discoveries lead to new insights, The FASEB Journal 12 (9) (1998) 633–651. doi:10.1096/fasebj.12.9.633.

[36] S. B. McMahon, MYC and the control of apoptosis, Cold Spring Harbor Perspectives in Medicine 4 (7) (2014) a014407. doi:10.1101/cshperspect.a014407.

[37] G. Bretones, M. D. Delgado, J. León, Myc and cell cycle control, Biochimica et Biophysica Acta - Gene Regulatory Mechanisms 1849 (5) (2015) 506–516. doi:10.1016/j.bbagrm.2014.03.013.

[38] N. Ninov, D. A. Chiarelli, E. Martín-Blanco, Extrinsic and intrinsic mechanisms directing epithelial cell sheet replacement during Drosophila metamorphosis, Development 134 (2) (2007) 367–379. doi:10.1242/dev.02728.

[39] Y. Nakajima, E. Kuranaga, K. Sugimura, A. Miyawaki, M. Miura, Nonautonomous apoptosis is triggered by local cell cycle progression during epithelial replacement in Drosophila, Molecular and Cellular Biology 31 (12) (2011) 2499–2512. doi:10.1128/mcb.01046-10.

[40] R. Levayer, B. Hauert, E. Moreno, Cell mixing induced by myc is required for competitive tissue invasion and destruction, Nature 524 (7566) (2015) 476–480. doi:10.1038/nature14684.

[41] L. Alpar, C. Bergantiños, L. A. Johnston, Spatially restricted regulation of Spätzle/Toll signaling during cell competition, Developmental Cell 46 (6) (2018) 1–14. doi:10.1016/j.devcel.2018.08.001.

[42] K. Maheden, The field of cell competition comes of age: semantics and technological synergy, Frontiers in Cell and Developmental Biology 10 (2022) 891569. `doi:10.3389/fcell.2022.891569`.

[43] G. Morata, P. Ripoll, Minutes: mutants of Drosophila autonomously affecting cell division rate, Developmental Biology 42 (2) (1975) 211–221. `doi:10.1016/0012-1606(75)90330-9`.

[44] P. Simpson, G. Morata, Differential mitotic rates and patterns of growth in compartments in the Drosophila wing, Developmental Biology 85 (2) (1981) 299–308. `doi:10.1016/0012-1606(81)90261-X`.

[45] S. Nishikawa, A. Takamatsu, S. Ohsawa, T. Igaki, Mathematical model for cell competition: predator-prey interactions at the interface between two groups of cells in monolayer tissue, Journal of Theoretical Biology 404 (2016). `doi:10.1016/j.jtbi.2016.05.031`.

[46] S. N. Meyer, M. Amoyel, C. Bergantiños, C. De La Cova, C. Schertel, K. Basler, L. A. Johnston, An ancient defense system eliminates unfit cells from developing tissues during cell competition, Science 346 (6214) (2014). `doi:10.1126/science.1258236`.

[47] N. Senoo-Matsuda, L. A. Johnston, Soluble factors mediate competitive and cooperative interactions between cells expressing different levels of Drosophila Myc, Proceedings of the National Academy of Sciences of the United States of America 104 (47) (2007) 18543–18548. `doi:10.1073/pnas.0709021104`.

[48] A. Richards, University of Oxford Advanced Research Computing (2015). `doi:10.5281/zenodo.22558`.

[49] O. Tange, GNU Parallel: the command-line power tool, ;login: The USENIX Magazine 36 (1) (2011) 42–47. `doi:10.5281/zenodo.16303`.