The RNA-DNA world and the emergence of DNA-encoded heritable traits

Suvam Roy[1] and Supratim Sengupta[1,*]

[1]Department of Physical Sciences, Indian Institute of Science Education and Research Kolkata,
Mohanpur-741246, India
[*]Address for correspondence: supratim.sen@iiserkol.ac.in

March 14, 2023

Abstract

The RNA world hypothesis confers a central role to RNA molecules in information encoding and catalysis. Even though evidence in support of this hypothesis has accumulated from both experiments and computational modelling, the transition from an RNA world to a world where heritable genetic information is encoded in DNA remains an open question. Recent experiments show that both RNA and DNA templates can extend complementary primers using free RNA/DNA nucleotides, either non-enzymatically or in the presence of a replicase ribozyme. Guided by these experiments, we analyse protocellular evolution with an expanded set of reaction pathways made possible through the presence of DNA nucleotides. We allow for the possibility of auto-catalytic synthesis of replicase ribozymes from DNA templates encoding the replicase, after the chance emergence of a replicase through non-enzymatic reactions. By encapsulating these reactions inside three different types of protocellular compartments, each subject to distinct modes of selection, we show how protocells containing replicase-encoded DNA in low copy numbers and replicases in high copy numbers can dominate the population. Our work suggests a plausible pathway for the transition from an RNA world to a mixed RNA-DNA world characterized by Darwinian evolution, where DNA sequences encode heritable phenotypes.

Keywords— Origin of Life, RNA World, Ribozyme, Heritable Traits

1 Introduction

The RNA world hypothesis states that a living system composed entirely of RNA may have emerged prior to the origin of the current DNA-protein world. The ability of RNA to store information [1, 2] and catalyse reactions [3–5] suggest that they may indeed have played a key role in the emergence of a self-sustaining chemical system capable of undergoing Darwinian evolution. However, the plausibility of such an RNA world relies on explaining how self-replicating protocells encapsulating RNA molecules first emerged from a primordial soup containing basic chemical building blocks. Several experiments have shown that both ribonucleotides [6–8] and deoxy-ribonucleotides [9, 10] can be spontaneously created in prebiotic environments. Non-enzymatic processes of concatenation and template-directed primer extension, albeit with activated RNA nucleotides [11], along with environmental cycling [12, 13] give rise to long RNA polymers that can fold into complex secondary structures [14]. It seems plausible that the first ribozymes emerged by chance through such processes, as a result of extensive sampling of the sequence space through error-prone replication [15] in a prebiotic environment. Recent work has provided evidence of phenotype bias, indicating that certain secondary structures which appear most frequently while sampling sequences of fixed length happen to be the ones that are found in nature [16]. Remarkably, those structures can be generated by sampling of a relatively small region of the sequence space, implying that

the chance of emergence of a functional bio-molecule may not be too small. However, the random creation of a few functional bio-molecules is not sufficient to guarantee their proliferation especially in a spatially open environment dominated by selfish parasitic sequences. It is therefore imperative for template-directed replication processes to be encapsulated in compartments since such segregation can help in preventing parasites from overwhelming the system [17]. If distinct ribozymes are created with small probabilities inside a protocell during replication by the rolling circle mechanism [18–22], they can act to not only synergistically enhance each others' formation but also increase the likelihood of creation of other functionally distinct ribozymes. This provides a pathway towards increasing functional complexity of the protocell and can allow for the proliferation of protocells containing a functionally diverse set of ribozymes [23]. However, such an evolution is not Darwinian in nature and the ribozymes generated cannot be considered as heritable traits since they appear by chance during the error-prone replication process and are also prone to degradation. How then is it possible to create a self-sustained chemical system that does not rely on random ribozyme production for its survival? A possible resolution to this problem in the RNA world was proposed by Takeuchi *et al.* [24] who showed that genome-like strands can appear inside vesicles due to symmetry-breaking between two complementary strands of a self-replicating ribozyme. One strand would then act as a genome and its complementary strand as an enzyme. Since there is very little free-energy difference

between a template and its complement, it seems unlikely that the reverse complement of a template strand would fold into a complex secondary structure that is characteristic of a ribozyme. Therefore, although ribozymes can possibly emerge in an RNA world [14], RNA alone might not have been sufficient for the emergence of heredity.

We believe the answer to this question lies in the appearance of information-encoding DNA templates that act like primitive pseudo-genes. A collection of such pseudo-genes, each encoding a different functional phenotype, would then constitute a primitive albeit fragmented genome. However, unlike ribonucleotides [25–33], free DNA nucleotides can not spontaneously polymerize into long DNA strands. Therefore the transition to a DNA world must have been facilitated by RNA. Indeed, experiments on mixed RNA-DNA template-primer systems have shown that RNA is capable of extending DNA primers both non-enzymatically [34] and under the action of a polymerase ribozyme [35]. Therefore, non-enzymatic and enzymatic reverse transcription of RNA strands might have been the primary mechanism of DNA strand creation in a primordial world.

In this work, we show how the presence of DNA nucleotides increases the space of possible reactions and opens up the possibility of creating single-stranded DNA sequences that can act as information-encoding templates. Even though the non-enzymatic template-directed replication process produces mostly useless sequences (parasites), we assume that bio-molecules like replicase ribozymes can be created with small probabilities. We further assume that the replication of RNA templates using DNA nucleotides can create, with a small probability, the DNA sequence encoding the replicase ribozyme. Those DNA sequences can act as templates, whose accurate, auto-catalytic replication with RNA nucleotides can generate new replicases. Such templates can speed up the replication process inside a protocell which in-turn can ensure rapid growth in the number of strands and enhanced likelihood of creation of new replicases and templates. A population of protocells can eventually evolve to increase the number of protocells containing both replicase-encoding templates and replicases thereby marking the onset of Darwinian evolution characterized by encoding of a heritable trait like the replicase ribozyme in DNA sequences. The nature of the protocellular compartment determines how likely it is for such protocells to dominate the population under reasonable conditions. Proliferation of such protocells are observed for coacervate droplets, water-in-oil droplets as well as vesicular compartments, but coacervate droplets which preferentially select for replicases [36], and water-in-oil droplets, allow for the dominance of such protocells under diverse conditions that are even robust to decrease in the error threshold for accurate replication. Intriguingly, in our model, evolution leads to the spontaneous emergence of protocells that contain the replicase-encoding DNA template in low copy numbers and replicases in high copy numbers. Our work shows how the creation of DNA-encoded heritable phenotypes through RNA-templated replication can lead to protocells capable of undergoing Darwinian evolution.

2 Methods

The reactions inside each protocell depends on the nature of the components present in it. The presence of DNA nucleotides allows for the non-enzymatic replication of RNA templates using deoxy-ribonucleotides [34, 37] leading to the creation of

DNA strands. The number of strands inside a protocell initially grow through non-enzymatic template-directed replication based on the reactions described below (see section 1 of the Supplementary information for details about estimation of the non-enzymatic replication rates). Enzymatic reactions are activated once a replicase emerges as a result of a chance non-enzymatic replication event. Table-1 gives both the non-enzymatic and enzymatic replication rates for the various reactions. The evolution of a population of such protocells occurs due to protocellular growth followed by division and selection in a manner that depends on the nature of the encapsulating compartment.

2.1 Reactions

Initially, we consider protocells containing RNA templates T_r only. Non-enzymatic replication of T_r by RNA nucleotides (with a rate $K_{rr}^n = 0.00255 T_r \ (h^{-1})$) will mostly create RNA parasites (P) (which are non-functional sequences that are neither templates nor ribozymes). However, following the argument given in the Introduction, we assume that replicase (R) ribozymes can be created with a small probability (p_R) during the error-prone, non-enzymatic replication process (see reaction (1)). The chance creation of a replicase eventually leads to the emergence of a more complex reaction network as indicated by the blue arrows in Fig.1. We also assume unlimited monomer supply by the environment to keep the rates independent of monomer concentrations. P/R on the RHS of reactions 1-9 indicates that both both parasites (P) and replicase (R) are produced albeit with different probabilities that are indicated in brackets. $T_d/T_{dP}/T_{dR}$ on the LHS of reactions 1-9 indicate that any one of them is used as a template and the corresponding template will also be present on the RHS of the reactions.

$$T_r \xrightarrow{K_{rr}^n} T_r + P(1 - p_R)/R(p_R) \qquad (1)$$

Table 1: Non-enzymatic and enzymatic replication rates for a template of length L=200 nt inside a protocell of radius 57 nm along with the corresponding percentage errors ($\varepsilon = 100 \times er/L$). $r = RNA$, $d = DNA$, $T = template$, $R = replicase$. K_{rd}^n, K_{rd}^e represent non-enzymatic and enzymatic replication rates from RNA to DNA etc.

Reaction	Rate (h^{-1})	Error (%)	Reference
K_{rr}^n	$0.00255 T$	$\sim 36\%$	[15, 38]
K_{rd}^n	$0.001646 T$	High	[34]
K_{dr}^n	$0.000823 T$	High	[34]
K_{dd}^n	$0.00255 T$	$\sim 32\%$	[15, 37]
K_{rr}^e	$0.46447 R$	$\sim 8\%$	[35, 39]
K_{rd}^e	$0.226 R$	$\sim 3\%$	[35]
K_{dr}^e	$0.1434 R$	$\sim 3\%$	[40]
K_{dd}^e	0	NA	[35]

Non-enzymatic replication of T_r by DNA nucleotides will create DNA templates, the majority of which will be non-functional since they will encode RNA parasites (T_{dP}) but a small fraction will encode (T_{dR}) replicase ribozymes.

$$T_r \xrightarrow{K_{rd}^n} T_r + T_{dP}(1 - p_R)/T_{dR}(p_R) \qquad (2)$$

Figure 1: Pictorial representation of the reaction network inside a protocell. Red (Blue) arrows indicate non-enzymatic (enzymatic) reactions. Ch-1, Ch-2 and Ch-3 denote 3 different channels for non-enzymatic replication of the 3 different types of DNA templates. The numbers 1-9 correspond to the reaction numbers specified below.

Non-enzymatic replication of DNA templates T_d (replica of T_r that can get created by enzymatic replication described later on), T_{dP} and T_{dR} by RNA nucleotides will create parasites and replicase (rate $K_{dr}^n = 0.000823\,(T_d + T_{dP} + T_{dR})\;(h^{-1})$). Similarly non-enzymatic replication of DNA templates T_d, T_{dP} and T_{dR} by DNA nucleotides can create T_{dP} and T_{dR} (rate $K_{dd}^n = 0.00255\,(T_d + T_{dP} + T_{dR})\;(h^{-1})$).

$$T_d/T_{dP}/T_{dR} \xrightarrow{K_{dr}^n} T_d/T_{dP}/T_{dR} + P(1-p_R)/R(p_R) \quad (3)$$

$$T_d/T_{dP}/T_{dR} \xrightarrow{K_{dd}^n} T_d/T_{dP}/T_{dR} + T_{dP}(1-p_R)/T_{dR}(p_R) \quad (4)$$

Initially, the processes inside a protocell are driven only by non-enzymatic reactions. However, upon the chance emergence of replicase ribozymes, enzymatic replication of T_r by RNA nucleotides (see reaction 5) can create new copies of the template with relatively higher accuracy. Nevertheless, parasites will continue to be created, since even enzymatic replication in a primordial world is error-prone (Table-1) in the absence of proof-reading mechanisms. We define a function $f(\varepsilon) = 1/(1 + exp(\varepsilon - e_T))$ where e_T is the error-threshold to quantify the likelihood of accurate replication under the action of a replicase ribozyme. This depends on the percentage error (ε) during such enzymatic replications (see Table-1). Fig-S1 shows the variation of this probability of accurate replication for these 3 types of enzymatic replication processes with different values of the error threshold e_T. Enzymatic replication of T_r by DNA nucleotides will similarly create (see reaction 6) a DNA version T_d of the template T_r or DNA template T_{dP} encoding a parasitic sequence, with different likelihoods modulated by the function $f(\varepsilon)$.

$$T_r + R \xrightarrow{K_{rr}^e} T_r + R + P(1-f(\varepsilon_{rr}))/T_r(f(\varepsilon_{rr})) \quad (5)$$

$$T_r + R \xrightarrow{K_{rd}^e} T_r + R + T_d(f(\varepsilon_{rd}))/T_{dP}(1-f(\varepsilon_{rd})) \quad (6)$$

The three types of DNA templates in our model allow for three different enzymatic replications of DNA templates (with a rate $K_{dr}^e = 0.1434\,R\;(h^{-1})$). Enzymatic replication of T_d by RNA nucleotides will create primarily parasites but can also produce a RNA template T_r. In the case of T_{dP} enzymatic replication by RNA nucleotides will create only parasites. Finally, autocatalytic replication of T_{dR} by RNA nucleotides can recreate a replicase depending on the error-threshold.

$$T_d + R \xrightarrow{K_{dr}^e} T_d + R + T_r(f(\varepsilon_{dr}))/P(1-f(\varepsilon_{dr})) \quad (7)$$

$$T_{dP} + R \xrightarrow{K_{dr}^e} T_{dP} + R + P \quad (8)$$

$$T_{dR} + R \xrightarrow{K_{dr}^e} T_{dR} + R + R(f(\varepsilon_{dr}))/P(1-f(\varepsilon_{dr})) \quad (9)$$

Reaction 9 marks the onset of one way information flow from the DNA template sequence to the replicase it encodes and can be considered to be a manifestation of a primitive central dogma. Fig.1 provides a pictorial representation of the nine types of replication reactions that can occur inside a protocell. Additionally, we also consider degradation of both RNA and DNA molecules. As DNA is more stable than RNA, the degradation rate of DNA (h_d) is taken to be lower than RNA (h_r). We use $h_r = 0.0008\;h^{-1}$, $h_d = 0.00008\;h^{-1}$ for all simulations.

$$T_r/P/R \xrightarrow{h_r} \Phi$$
$$T_d/T_{dP}/T_{dR} \xrightarrow{h_d} \Phi \quad (10)$$

The differential equations determining the time evolution of the abundances of the six different types of molecules, as a result of reactions (1)-(10), are given in the *Supplementary information*.

2.2 Protocellular compartments and population evolution

We considered N=400 protocells each initially containing 10 RNA templates T_r. To account for variation between different protocells, we assumed that the templates inside different protocells have different replicase creation probabilities, with the probability taken randomly from the range ($0.5\,p_R$, $1.5\,p_R$). The

Figure 2: Stochastic simulation of a population of water-in-oil droplets containing strands. Heatmaps for **A:** average fraction of replicase ribozymes per droplet; **B:** fraction of droplets containing both replicase (R) and replicase encoding DNA template (T_{dR}); **C:** average reaction propensity of reaction 9 per droplet; with different non-enzymatic ribozyme creation probabilities and error thresholds of enzymatic replications. The heatmaps are generated by taking both ensemble average and time average of the quantities at equilibrium.

two key parameters in our model are the creation probability (p_R) of R/T_{dR} via error-prone, non-enzymatic replication and the error-threshold (e_T) in case of enzymatic reactions. We varied these 2 parameters and carried out stochastic simulations of the evolving population. The dynamics of the evolving population can be best understood by tracking the average fraction of replicase per protocell, fraction of protocells containing both replicase R, the corresponding DNA strand encoding it (T_{dR}) and the relative propensity of reaction 9 (see *Supplementary information* for details.)

The three different types of protocellular compartments in our model are distinguished by the distinct selection mechanisms they undergo during the evolution of the protocellular population.

In the first scenario, our choice of protocellular compartments made out of water-in-oil droplets was inspired by recent RNA host-parasite experiments [41–45]. In the experiments, the protocellular compartments are subjected to periodic washout-mixing cycles where a fraction of droplets are randomly removed from the population followed by supply of empty compartments which are then mixed with existing compartments leading to random redistribution of components from the filled to empty compartments. We model this system by considering pairwise competition between protocells that is initiated whenever the number of strands inside all droplets lie in the range ($V/5$, V). During such competition, each droplet competes with another randomly chosen droplet and replaces it by dividing into 2 daughter protocells with components of the surviving protocells being randomly distributed between the two daughter droplets.

Another alternative compartment we consider are coacervate droplets formed by liquid-liquid phase separation [46–49]. Such coacervate droplets can undergo growth and division under environmental cycling [50, 51]. Template-directed reactions were shown to occur inside such droplets [52] which can preferentially sequester and replicate long and potentially functional RNA molecules like ribozymes. Further, coacervate droplets are shown to up concentrate ribozymes [36] and ribozymes show enhanced activity inside those droplets [46, 53]. Such preference shown by coacervate droplets for containing functional RNA molecules comes from the observation that coacervates prefer molecules which exhibit electrostatic interactions [54] as such interactions inhibit their Ostwald ripening and increase their stability [55]. Therefore in our coacervate droplet model we consider a competition similar to the water-in-oil droplet model

but with selection dependent not just on the number of strands but on the fraction of replicase ribozymes inside the droplet. This amounts to defining a fitness function f_i for each droplet involved in pairwise competition with $f_i \propto \frac{R^i}{T_r^i + P^i + T_d^i + T_{dP}^i + T_{dR}^i}$. The droplet selected on the basis of its fitness divides into 2 daughter droplets, one of which replaces the droplet eliminated during pairwise competition.

Finally, vesicular compartments with lipid bilayer membranes were also considered. The volume of the vesicle is determined by the number of strands inside it and vesicle division occurs when the number of strands reach an upper bound set to $V = 1000$. During the division process, the components of the vesicle are randomly distributed among the two daughter vesicles while an existing vesicle is eliminated with a probability proportional to the difference in the number of strands between itself and the dividing vesicle. This results in the preferential elimination of smaller vesicles [56] while keeping the population size fixed.

These three different types of competition between protocells leads to distinct conditions for the dominance of protocells containing a replicase ribozyme whose synthesis is brought about primarily through template-directed, replicase-catalysed replication of a DNA sequence (T_{dR}) encoding the enzyme.

3   Results

The presence of DNA nucleotides inside a single protocell can enhance the process of replicase creation and ensure sustained growth of strands. When enzymatic reactions are activated due to the chance emergence of a replicase, the presence of even a single DNA template ensures rapid growth of both DNA and RNA strands that further facilitate replicase production (see section 4 of Supplementary Information for details). In the subsections below, we describe how the encapsulation of strands in three different types of protocellular compartments affect the outcome of competition between protocells in an evolving population.

3.1   Water-in-Oil droplet model

Fig-2 shows the heatmaps at equilibrium for the average fraction of replicase per droplet (panel A), fraction of droplets containing both replicase R & the DNA template (T_{dR}) encoding

Figure 3: Stochastic simulation of a population of water-in-oil droplets ($p_R = 0.002$, $e_T = 4$). Time evolution of the **A:** Average number of different types of strands per droplet; **B:** Percentage of droplets containing different types of strands; **C:** Average reaction propensities of the 9 types of reactions per droplet. Panels **A** and **B** have the same legends.

it (panel B); and the relative propensity of reaction 9 (panel C) which provides a measure of the efficacy of replicase creation from its encoded DNA template. A very low non-enzymatic probability of replicase creation ($p_R < 0.002$) is not favourable for ensuring the presence of both R and T_{dR} in majority of droplets (see lower part of Fig-2(B)). A low value of p_R leads to low copy numbers of both R and T_{dR} through non-enzymatic processes which primarily lead to the creation of parasites (P) or parasite encoding DNA (T_{dP}). The enzymatic creation of replicases through reaction 9 is suppressed since that pathway depends on the easy availability of both R and T_{dR}. Even when an increase in the error threshold increases the probability of creation of T_r and T_d through reactions 5-7, it does not have an impact because of the suppression in creation of R and T_{dR} ensures that the reactant concentrations for 5-7 remain quite low. Even if the probability of non-enzymatic creation of replicases is relatively high (upper left region of Fig-2), low enzymatic replication fidelity due to very low error thresholds make parasite creation through enzymatic reaction channels 5-9 more likely, leading to lower average replicase fraction per droplet (see upper left region of Fig-2(A)).

When the non-enzymatic replicase creation probability (p_R) as well as the error threshold (e_T) are high, replicases are produced with a high likelihood both non-enzymatically and enzymatically since the replicase phenotype is more robust to replication errors. However, the resultant replicases are predominantly used to catalyze reactions 5-7 as evident from the higher reaction propensities of these reactions in the upper right region of Fig-S4(B). Replicases available for catalyzing their own formation are relatively fewer and reaction 9 is sub-dominant (see upper right region of Fig-2(C)). Moreover, since T_r and T_d are created with higher fidelity in this region of parameter space. the relative fraction of R is comparatively low as seen in Fig-2(A). In contrast, reaction 9 is dominant for moderately low non-enzymatic replicase creation probability and moderate to high error-threshold (see centre-right region of Fig-2(C)) leading to a comparatively larger average fraction of replicases per protocell. Fig-3 shows the time evolution of the average number of different strands (Fig-3(A)) for a point lying in this region. The percentage of droplets containing both R and T_{dR} (DNA template encoding R) is $\sim 48\%$ (Fig-3(B)) and the average propensity of reaction 9, responsible for replicase creation using a DNA template encoding the replicase, is large (Fig-3(C)). Even though parasites continue to be formed, primarily through reaction 8, efficient creation of replicases both non-enzymatically and enzymatically is sufficient to ensure that close to a majority of

protocells contain both R and T_{dR}. The continuous regeneration of both R and T_{dR} can be sustained in the population even if the non-enzymatic reactions are eventually switched off after the onset of enzymatic replication of R (see Fig-S5). Intriguingly, we also find symmetry breaking between the number of T_{dR} and R with the average number of former and latter being ~ 1 per droplet and ~ 100 per droplet respectively (see Fig-3(A)). The crucial role played by reaction 9 in sustaining replicase formation and aiding the proliferation of droplets containing both R and T_{dR} is evident from Fig-S6 which shows the outcome of switching off reaction 9. Doing so prevents the asymmetry in copy numbers from emerging, drastically reduces the average number of *both* R and T_{dR} to below one per droplet and the percentage of droplets containing both R and T_{dR} to $< 20\%$ of the population. This indicates that replicase creation through non-enzymatic processes alone cannot drive the evolution of the protocellular population towards increasing functional complexity.

3.2 Coacervate droplet model

In the case of coacervate droplets, selection pressure acts in favour of droplets containing more ribozymes. The region of parameter space where the fraction of droplets containing both R & T_{dR} exceeds $\sim 50\%$ also overlaps with the region where the reaction propensity of 9 is large (Fig-4) and is characterized by moderately low values ($\geq \mathscr{O}(10^{-3})$) of non-enzymatic creation probability of a replicase. T_{dR} creation inside coacervate droplets can be amplified by harnessing the products (T_d and T_{dP}) of the replicase catalysed reaction 6 and utilizing them to drive the non-enzymatic reactions 3-4 occurring through reaction channels Ch-1 and Ch-3. Hence, replicase creation through reaction 9 leads to a positive feedback that enhances the propensity of non-enzymatic reactions and induces the creation of even more replicases. Droplets which exploit such positive feedback loops resulting from creation of R & T_{dR} will be preferentially selected (see Fig-S8(B) and text below it in the Supplementary information) over droplets containing only parasites and can flourish even after non-enzymatic reactions are turned off (Fig-S9). The symmetry breaking between R & T_{dR} is also observed in this region (Fig-S8(A)), with the asymmetry between DNA templates and replicases vanishing in the absence of reaction 9 (see Fig-S10).

Figure 4: Stochastic simulation of a population of coacervate droplets containing strands. Heatmaps for **A:** average fraction of replicase ribozymes per droplet; **B:** fraction of droplets containing both replicase (R) and replicase encoding DNA template (T_{dR}); **C:** average reaction propensity of reaction 9 per droplet; with different non-enzymatic ribozyme creation probabilities and error thresholds of enzymatic replications. The heatmaps are generated by taking both ensemble average and time average of the quantities at equilibrium.

3.3 Vesicle model

For such compartments, selection of a protocell depends only on the number but not on the type of strands in it, unlike the previous case. Since replicase formation is favoured both non-enzymatically (through reactions 1 and 3) and enzymatically (through reaction 9) for high p_R and high e_T respectively, the fraction of replicase per protocell increases with the non-enzymatic replicase creation probability (p_R) and error threshold (e_T). Surprisingly, the parameter region of highest replicase fraction per vesicle is distinct from the region with the highest propensity of reaction 9, which occurs for low e_T (see Fig-S11). For high e_T, reactions 5-8 collectively dominate over reaction 9 and the non-enzymatic reactions (compare Fig-S11 and S12). Moreover, they use up replicases as catalysts producing parasites, or templates that feed the non-enzymatic reactions (see Fig-1); thereby reducing the propensity of reaction 9 which depends on the presence of both R and TdR. For low error-thresholds, the fidelity of replicase creation through reaction 9 is reduced thereby creating a negative feedback loop that reduces the propensity of enzymatic reactions 5-8 and leading to the decrease in number of both RNA and DNA templates. In this regime, the contribution of non-enzymatic reactions to the creation of replicase-encoding DNA template can be significant at high p_R (see Fig-S13 in the supplementary information) and when those reactions are switched off, the average number of T_{dR} per vesicle drops below 1 and the percentage of vesicles containing T_{dR} drops below 50% (see Fig-S14 in the supplementary information). Even though reaction 9 is sub-dominant compared to other enzymatic reactions (see Fig-S13(C)), it is essential for ensuring that the average number of replicases inside a vesicle exceeds the number of DNA templates encoding the replicase (see Fig-S15 and section 7 of supplementary information).

4 Discussion

Evolution of a protocellular population, aided by the coupling between growth in the number of internal components and division can occur through physical processes only [23, 57]. However, the onset of Darwinian evolution requires heritable and selectively advantageous phenotypes encoded in sequences to be accurately replicated so that they can spread through the population. In this work, we show how the transition to an epoch characterized by Darwinian evolution can be facilitated by the presence of DNA nucleotides. The RNA-templated creation of DNA sequences using DNA nucleotides lead to new reaction pathways that increase the likelihood of replicase formation through both non-enzymatic and enzymatic channels. Once enzymatic pathways are established, they can help in sustainably creating replicases by increasing both the speed and accuracy of replicase formation. Such enzymatic replicase creation induces symmetry breaking in the number of replicases and replicase-encoding DNA templates, increasing the copy number of the former relative to the latter.

Changing the nature of the protocellular compartment affects the mode of selection and consequently the ability of protocells containing replicases to spread through the population. It is most desirable to get sustained replication and proliferation of a protocell containing both the encoded DNA template (T_{dR}) and corresponding replicase (R) for low error thresholds and low values of non-enzymatic replicase creation probability. These conditions are satisfied for coacervate and water-in-oil droplet models and we therefore conclude that those compartments will be more effective in ensuring the dominance of protocells containing R and T_{dR} in high and low copy numbers respectively.

Even though several aspects of our model can be subject to experimental validation, we believe the most compelling test would be to demonstrate the efficacy of auto-catalytic replicase creation through reaction 9 inside a protocell. Although ribozyme catalysed replication of sequences longer than 100 nucleotides has not yet been experimentally observed, a possible alternative approach might involve ribozyme catalysed replication of different fragments of a complete DNA template encoding the ribozyme. The fragments generated can then self-assemble to form the whole ribozyme [58–60]. The versatility of the replicase thus created and its role in catalysing several other reactions, especially in the presence of both RNA and DNA nucleotides, can create a self-sustained chemical system inside a protocell. Although we focused on replicase creation only in this work, other functional ribozymes can also be sustainably produced in a similar fashion. We envisage a scenario where multiple functional ribozymes each encoded by their specific DNA templates gradually emerge, providing an increasing selective advantage to the protocell encapsulating them and consequently facilitating their proliferation in the population.

The epoch of evolution discussed here predates the Darwinian transition and is also likely to have been of a communal nature [61] characterized by widespread horizontal transfer of

sequence elements across protocells. It would be interesting to explore how such horizontal transfer could have driven the evolution towards increasing complexity characterized by the presence of functionally diverse components inside such progenotes. We hope that our work, which establishes a proof of concept for the emergence and proliferation of encoded heritable phenotypes, will motivate the design of novel *in vitro* evolution experiments that will eventually help in further unravelling the mystery of the origin of life.

5 Data accessibility

All codes used to generate the results in the manuscript and in the electronic supplementary material are available at the following Github link.

6 Acknowledgments

S.R. is supported by an INSPIRE graduate fellowship given by SERB, India. S.S. acknowledges partial support through a MATRICS grant (MTR/2020/000446 given by SERB, India), that was utilized for purchasing computing equipment.

References

[1] S. Brenner, F. Jacob, and M. Meselson. An unstable intermediate carrying information from genes to ribosomes for protein synthesis. *Nature*, 190(4776):576–581, may 1961.

[2] Francois Gros, H. Hiatt, Walter Gilbert, C. G. Kurland, R. W. Risebrough, and J. D. Watson. Unstable ribonucleic acid revealed by pulse labelling of escherichia coli. *Nature*, 190(4776):581–585, may 1961.

[3] Kelly Kruger, Paula J. Grabowski, Arthur J. Zaug, Julie Sands, Daniel E. Gottschling, and Thomas R. Cech. Self-splicing RNA: Autoexcision and autocyclization of the ribosomal RNA intervening sequence of tetrahymena. *Cell*, 31(1):147–157, nov 1982.

[4] B. C. Stark, R. Kole, E. J. Bowman, and S. Altman. Ribonuclease p: an enzyme with an essential RNA component. *Proceedings of the National Academy of Sciences*, 75(8):3717–3721, aug 1978.

[5] Cecilia Guerrier-Takada, Katheleen Gardiner, Terry Marsh, Norman Pace, and Sidney Altman. The RNA moiety of ribonuclease p is the catalytic subunit of the enzyme. *Cell*, 35(3):849–857, dec 1983.

[6] Matthew W. Powner, Béatrice Gerland, and John D. Sutherland. Synthesis of activated pyrimidine ribonucleotides in prebiotically plausible conditions. *Nature*, 459(7244):239–242, may 2009.

[7] Brian J. Cafferty, David M. Fialho, Jaheda Khanam, Ramanarayanan Krishnamurthy, and Nicholas V. Hud. Spontaneous formation and base pairing of plausible prebiotic nucleotides in water. *Nature Communications*, 7:11328, apr 2016.

[8] Sidney Becker, Jonas Feldmann, Stefan Wiedemann, Hidenori Okamura, Christina Schneider, Katharina Iwan, Antony Crisp, Martin Rossa, Tynchtyk Amatov, and Thomas Carell. Unified prebiotically plausible synthesis of pyrimidine and purine RNA ribonucleotides. *Science*, 366(6461):76–82, oct 2019.

[9] Jianfeng Xu, Václav Chmela, Nicholas J. Green, David A. Russell, Mikołaj J. Janicki, Robert W. Góra, Rafał Szabla, Andrew D. Bond, and John D. Sutherland. Selective prebiotic formation of RNA pyrimidine and DNA purine nucleosides. *Nature*, 582(7810):60–66, June 2020.

[10] Jianfeng Xu, Nicholas J. Green, David A. Russell, Ziwei Liu, and John D. Sutherland. Prebiotic photochemical coproduction of purine ribo- and deoxyribonucleosides. *Journal of the American Chemical Society*, 143(36):14482–14486, September 2021.

[11] K. Adamala and J. W. Szostak. Nonenzymatic template-directed RNA synthesis inside model protocells. *Science*, 342(6162):1098–1100, nov 2013.

[12] Bruce Damer and David Deamer. The hot spring hypothesis for an origin of life. *Astrobiology*, 20(4):429–452, apr 2020.

[13] Suvam Roy and Supratim Sengupta. The effect of environment on the evolution and proliferation of protocells of increasing complexity. *Life*, 12(8):1227, aug 2022.

[14] Suvam Roy, Niraja V. Bapat, Julien Derr, Sudha Rajamani, and Supratim Sengupta. Emergence of ribozyme and tRNA-like structures from mineral-rich muddy pools on prebiotic earth. *Journal of Theoretical Biology*, 506:110446, dec 2020.

[15] Kevin Leu, Eric Kervio, Benedikt Obermayer, Rebecca M. Turk-MacLeod, Caterina Yuan, Jesus-Mario Luevano, Eric Chen, Ulrich Gerland, Clemens Richert, and Irene A. Chen. Cascade of reduced speed and accuracy after errors in enzyme-free copying of nucleic acid sequences. *Journal of the American Chemical Society*, 135(1):354–366, dec 2012.

[16] Kamaludin Dingle, Fatme Ghaddar, Petr Šulc, and Ard A Louis. Phenotype bias determines how natural RNA structures occupy the morphospace of all possible shapes. *Molecular Biology and Evolution*, 39(1), sep 2021.

[17] Vismay Shah, Jonathan de Bouter, Pauli Quinn, Andrew Tupper, and Paul G. Higgs. Survival of RNA replicators is much easier in protocells than in surface-based, spatial systems. *Life*, 9(3):65, aug 2019.

[18] Rika Kusumoto-Matsuo, Tadahito Kanda, and Iwao Kukimoto. Rolling circle replication of human papillomavirus type 16 DNA in epithelial cell extracts. *Genes to Cells*, 16(1):23–33, nov 2010.

[19] José-Antonio Daròs, Santiago F Elena, and Ricardo Flores. Viroids: an ariadne's thread into the RNA labyrinth. *EMBO reports*, 7(6):593–598, jun 2006.

[20] Ricardo Flores, María-Eugenia Gas, Diego Molina-Serrano, María-Ángeles Nohales, Alberto Carbonell, Selma Gago, Marcos De la Peña, and José-Antonio Daròs. Viroid replication: Rolling-circles, enzymes and ribozymes. *Viruses*, 1(2):317–334, sep 2009.

[21] Andrew S. Tupper and Paul G. Higgs. Rolling-circle and strand-displacement mechanisms for non-enzymatic RNA replication at the time of the origin of life. *Journal of Theoretical Biology*, 527:110822, oct 2021.

[22] Felipe Rivera-Madrinan, Katherine Di Iorio, and Paul G. Higgs. Rolling circles as a means of encoding genes in the RNA world. *Life*, 12(9):1373, September 2022.

[23] Suvam Roy and Supratim Sengupta. Evolution towards increasing complexity through functional diversification in a protocell model of the RNA world. *Proceedings of the Royal Society B: Biological Sciences*, 288(1963), nov 2021.

[24] Nobuto Takeuchi, Paulien Hogeweg, and Kunihiko Kaneko. The origin of a primordial genome through spontaneous symmetry breaking. *Nature Communications*, 8(1), aug 2017.

[25] Sudha Rajamani, Alexander Vlassov, Seico Benner, Amy Coombs, Felix Olasagasti, and David Deamer. Lipid-assisted synthesis of RNA-like polymers from mononucleotides. *Origins of Life and Evolution of Biospheres*, 38(1):57–74, nov 2007.

[26] Veronica DeGuzman, Wenonah Vercoutere, Hossein Shenasa, and David Deamer. Generation of oligonucleotides under hydrothermal conditions by non-enzymatic polymerization. *Journal of Molecular Evolution*, 78(5):251–262, may 2014.

[27] Laura Toppozini, Hannah Dies, David W. Deamer, and Maikel C. Rheinstädter. Adenosine monophosphate forms ordered arrays in multilamellar lipid matrices: Insights into assembly of nucleic acid for primitive life. *PLoS ONE*, 8(5):e62810, may 2013.

[28] Jay G. Forsythe, Sheng-Sheng Yu, Irena Mamajanov, Martha A. Grover, Ramanarayanan Krishnamurthy, Facundo M. Fernández, and Nicholas V. Hud. Ester-mediated amide bond formation driven by wet-dry cycles: A possible path to polypeptides on the prebiotic earth. *Angewandte Chemie International Edition*, 54(34):9871–9875, jul 2015.

[29] Irena Mamajanov, Patrick J. MacDonald, Jingya Ying, Daniel M. Duncanson, Garrett R. Dowdy, Chelsea A. Walker, Aaron E. Engelhart, Facundo M. Fernández, Martha A. Grover, Nicholas V. Hud, and F. Joseph Schork. Ester formation and hydrolysis during wet–dry cycles: Generation of far-from-equilibrium polymers in a model prebiotic reaction. *Macromolecules*, 47(4):1334–1343, feb 2014.

[30] Chaitanya Mungi and Sudha Rajamani. Characterization of RNA-like oligomers from lipid-assisted nonenzymatic synthesis: Implications for origin of informational molecules on early earth. *Life*, 5(1):65–84, jan 2015.

[31] Wenhua Huang and James P. Ferris. One-step, regioselective synthesis of up to 50-mers of RNA oligomers by montmorillonite catalysis. *Journal of the American Chemical Society*, 128(27):8914–8919, jul 2006.

[32] Prakash C. Joshi, Michael F. Aldersley, John W. Delano, and James P. Ferris. Mechanism of montmorillonite catalysis in the formation of RNA oligomers. *Journal of the American Chemical Society*, 131(37):13369–13374, sep 2009.

[33] Tue Hassenkam, Bruce Damer, Gabriel Mednick, and David Deamer. AFM images of viroid-sized rings that self-assemble from mononucleotides through wet–dry cycling: Implications for the origin of life. *Life*, 10(12):321, nov 2020.

[34] Kevin Leu, Benedikt Obermayer, Sudha Rajamani, Ulrich Gerland, and Irene A. Chen. The prebiotic evolutionary advantage of transferring genetic information from RNA to DNA. *Nucleic Acids Research*, 39(18):8135–8147, June 2011.

[35] Biswajit Samanta and Gerald F Joyce. A reverse transcriptase ribozyme. *eLife*, 6, sep 2017.

[36] Carsten Donau, Fabian Späth, Marilyne Sosson, Brigitte A. K. Kriebisch, Fabian Schnitter, Marta Tena-Solsona, Hyun-Seo Kang, Elia Salibi, Michael Sattler, Hannes Mutschler, and Job Boekhoven. Active coacervate droplets as a model for membraneless organelles and protocells. *Nature Communications*, 11(1), oct 2020.

[37] E. Kervio, B. Claasen, U. E. Steiner, and C. Richert. The strength of the template effect attracting nucleotides to naked DNA. *Nucleic Acids Research*, 42(11):7409–7420, may 2014.

[38] Niraja V. Bapat and Sudha Rajamani. Effect of co-solutes on template-directed nonenzymatic replication of nucleic acids. *Journal of Molecular Evolution*, 81(3-4):72–80, oct 2015.

[39] David P. Horning and Gerald F. Joyce. Amplification of RNA by an RNA polymerase ribozyme. *Proceedings of the National Academy of Sciences*, 113(35):9786–9791, aug 2016.

[40] David P. Horning, Saikat Bala, John C. Chaput, and Gerald F. Joyce. RNA-catalyzed polymerization of deoxyribose, threose, and arabinose nucleic acids. *ACS Synthetic Biology*, 8(5):955–961, May 2019.

[41] Norikazu Ichihashi, Kimihito Usui, Yasuaki Kazuta, Takeshi Sunami, Tomoaki Matsuura, and Tetsuya Yomo. Darwinian evolution in a translation-coupled RNA replication system within a cell-like compartment. *Nature Communications*, 4(1), October 2013.

[42] Yohsuke Bansho, Taro Furubayashi, Norikazu Ichihashi, and Tetsuya Yomo. Host–parasite oscillation dynamics and evolution in a compartmentalized RNA replication system. *Proceedings of the National Academy of Sciences*, 113(15):4045–4050, March 2016.

[43] Tomoaki Yoshiyama, Tetsuo Ichii, Tetsuya Yomo, and Norikazu Ichihashi. Automated in vitro evolution of a translation-coupled RNA replication system in a droplet flow reactor. *Scientific Reports*, 8(1), August 2018.

[44] Taro Furubayashi, Kensuke Ueda, Yohsuke Bansho, Daisuke Motooka, Shota Nakamura, Ryo Mizuuchi, and Norikazu Ichihashi. Emergence and diversification of a host-parasite RNA ecosystem through darwinian evolution. *eLife*, 9, July 2020.

[45] Ryo Mizuuchi, Taro Furubayashi, and Norikazu Ichihashi. Evolutionary transition from a single RNA replicator to a multiple replicator network. *Nature Communications*, 13(1), March 2022.

[46] Björn Drobot, Juan M. Iglesias-Artola, Kristian Le Vay, Viktoria Mayr, Mrityunjoy Kar, Moritz Kreysing, Hannes Mutschler, and T-Y Dora Tang. Compartmentalised RNA catalysis in membrane-free coacervate protocells. *Nature Communications*, 9(1), September 2018.

[47] Christine D. Keating. Aqueous phase separation as a possible route to compartmentalization of biological molecules. *Accounts of Chemical Research*, 45(12):2114–2124, February 2012.

[48] Raghav R. Poudyal, Fatma Pir Cakmak, Christine D. Keating, and Philip C. Bevilacqua. Physical principles and extant biology reveal roles for RNA-containing membraneless compartments in origins of life chemistry. *Biochemistry*, 57(17):2509–2519, March 2018.

[49] Shogo Koga, David S. Williams, Adam W. Perriman, and Stephen Mann. Peptide–nucleotide microdroplets as a step towards a membrane-free protocell model. *Nature Chemistry*, 3(9):720–724, August 2011.

[50] Hadi M. Fares, Alexander E. Marras, Jeffrey M. Ting, Matthew V. Tirrell, and Christine D. Keating. Impact of wet-dry cycling on the phase behavior and compartmentalization properties of complex coacervates. *Nature Communications*, 11(1), October 2020.

[51] David Zwicker, Rabea Seyboldt, Christoph A. Weber, Anthony A. Hyman, and Frank Jülicher. Growth and division of active droplets provides a model for protocells. *Nature Physics*, 13(4):408–413, December 2016.

[52] Raghav R. Poudyal, Rebecca M. Guth-Metzler, Andrew J. Veenis, Erica A. Frankel, Christine D. Keating, and Philip C. Bevilacqua. Template-directed RNA polymerization and enhanced ribozyme catalysis inside membraneless compartments formed by coacervates. *Nature Communications*, 10(1), jan 2019.

[53] Raghav R. Poudyal, Christine D. Keating, and Philip C. Bevilacqua. Polyanion-assisted ribozyme catalysis inside complex coacervates. *ACS Chemical Biology*, 14(6):1243–1248, June 2019.

[54] N. Amy Yewdall, Alain A.M. André, Tiemei Lu, and Evan Spruijt. Coacervates as models of membraneless organelles. *Current Opinion in Colloid & Interface Science*, 52:101416, April 2021.

[55] Karina K. Nakashima, Merlijn H. I. van Haren, Alain A. M. André, Irina Robu, and Evan Spruijt. Active coacervate droplets are protocells that grow and resist ostwald ripening. *Nature Communications*, 12(1), June 2021.

[56] I. A. Chen. The emergence of competition between model protocells. *Science*, 305(5689):1474–1476, sep 2004.

[57] I. A. Chen, R. W. Roberts, and J. W. Szostak. The emergence of competition between model protocells. *Science*, 305(5689):1474–1476, sep 2004.

[58] Lijun Zhou, Derek K. O'Flaherty, and Jack W. Szostak. Assembly of a ribozyme ligase from short oligomers by nonenzymatic ligation. *Journal of the American Chemical Society*, 142(37):15961–15965, August 2020.

[59] Eric J. Hayden and Niles Lehman. Self-assembly of a group i intron from inactive oligonucleotide fragments. *Chemistry & Biology*, 13(8):909–918, August 2006.

[60] Nilesh Vaidya, Michael L. Manapat, Irene A. Chen, Ramon Xulvi-Brunet, Eric J. Hayden, and Niles Lehman. Spontaneous network formation among cooperative RNA replicators. *Nature*, 491(7422):72–77, October 2012.

[61] Carl Woese. The universal ancestor. *Proceedings of the National Academy of Sciences*, 95(12):6854–6859, June 1998.