Persistence of SARS-CoV-2 and its surrogate, bacteriophage Phi6, on surfaces and in water

Ana K. Pitol [1*], Samiksha Venkatesan [1], Grant Hughes [1]

[1]Departments of Vector Biology and Tropical Disease Biology, Centre for Neglected Tropical Diseases,

Liverpool School of Tropical Medicine, Liverpool, L3 5QA, UK

*Corresponding author: ana.pitolgarcia@lstmed.ac.uk

ABSTRACT

The COVID-19 pandemic has motivated research on the persistence of infectious SARS-CoV-2 in environmental reservoirs such as surfaces and water. Viral persistence data has been collected for SARS-CoV-2 and its surrogates, including bacteriophage Phi6. Despite its wide use, no side-by-side comparisons between Phi6 and SARS-CoV-2 exist. Here, we quantified the persistence of SARS-CoV-2 and Phi6 on surfaces (plastic and metal) and in water and evaluated the influence that the deposition solution has on viral persistence by using four commonly used deposition solutions: two culture media (DMEM and Tryptone Soya Broth (TSB)), Phosphate Buffered Saline (PBS), and human saliva. Phi6 remained infectious in water significantly longer than SARS-CoV-2, having a half-life of 27 hours as compared with 15 hours for SARS-CoV-2. The persistence of viruses on surfaces was significantly influenced by the virus used and the deposition solution, but not by the surface material. Phi6 remained infectious significantly longer than SARS-CoV-2 when the inoculation solution was culture media (DMEM, TSB) and saliva. Using culture media and saliva led to half-lives between 9 hours and 2 weeks for Phi6, as compared to 0.5 to 2 hours for SARS-CoV-2. Using PBS as a deposition solution led to half-lives shorter than 4 hours for both viruses on all surfaces. Our results showed that, although it has been frequently used as a surrogate for coronaviruses, bacteriophage Phi6 is not an adequate surrogate for studies quantifying SARS-CoV-2 persistence, as it over-estimates infectiousness. Additionally, our findings reveal the need of using adequate deposition solutions when evaluating viral persistence on surfaces.

KEYWORDS

SARS-CoV-2, bacteriophage Phi6, surrogate, survival, surface, fomite, water

INTRODUCTION

SARS-CoV-2 is primarily transmitted through respiratory droplets and aerosols[1]. Nevertheless, the RNA of SARS-CoV-2 has been extensively detected on environmental reservoirs, with concentrations as high as 10^5 genome copies (gc) per swab on surfaces, and $>10^5$ gc mL^{-1} in wastewater samples[2–7]. The extensive contamination of environmental reservoirs with the SARS-CoV-2 RNA has been the source of concern throughout the COVID-19 pandemic[8]. However, viral RNA concentration is higher than that of infectious viral particles on a given sample[9,10]. Hence, using data on RNA contamination to estimate the risks associated with people interacting with contaminated environments could lead to overestimating the risks. Therefore, it is important to understand the persistence of infectious SARS-CoV-2 to estimate the magnitude of the risks associated with people interacting with contaminated environments.

Since the beginning of the pandemic, there have been a number of studies quantifying the persistence and inactivation of SARS-CoV-2 on surfaces and in liquids under different environmental conditions such as temperature and humidity[9–16]. However, there is a lack of consistency in the experimental designs of these studies, which could lead to biased outcomes. For example, in studies of SARS-CoV-2 persistence on surfaces, a wide range of deposition solutions has been used to inoculate the virus, these range from culture media and Bovine Serum Albumin (BSA) containing media[16,17] to bodily fluids such as human saliva and mucus[10,18]. The solutions used for virus inoculation have very different characteristics which can influence viral survival[19]. Additionally, working with SARS-CoV-2 requires handling the virus in high containment facilities, which presents many challenges and limits the type of experiments that can be performed. For example, it is not advisable to do experimental work using SARS-CoV-2 with human volunteers, such as testing SARS-CoV-2 persistence on human hands or quantifying the transfer of the virus between surfaces and hands.

Therefore, efforts have been made to use surrogates, instead of SARS-CoV-2, to understand the mechanisms of its survival, inactivation, and transfer [20–24]. Bacteriophages are frequently used as surrogates of pathogenic viruses because they are safe, inexpensive, and do not require contaminant facilities [25–27]. Bacteriophage Phi6 is one of the few bacteriophages that has a lipid envelope [28]. Therefore, it has been used as a surrogate for enveloped viruses such as SARS and MERS coronaviruses, influenza virus, and Ebola [20–22,27,29–32]. To this end, bacteriophage Phi6 has been used as a surrogate for SARS-CoV-2 in studies evaluating virus persistence on surfaces [22,24,32], virus persistence in water and wastewater [33], virus inactivation [20,21,30], virus transfer from surfaces to hands[34] and virus recovery from fingertips [23]. While these studies provide important information on viral persistence and transfer, there is a gap in our knowledge related to the adequacy of using bacteriophage Phi6 as a proxy for SARS-CoV-2. As such, there is an urgent need to evaluate the suitability of using bacteriophage Phi6 as a surrogate for SARS-CoV-2 in persistence studies.

Here we evaluated the persistence of SARS-CoV-2 and bacteriophage Phi6 in water. Additionally, we assessed the persistence of SARS-CoV-2 and Phi6 on surfaces of two different non-porous materials: plastic (PVC) and metal (stainless steel). Since the deposition solution used to inoculate the virus on the surface could have an influence on virus persistence, we evaluated four commonly used deposition solutions: Phosphate Buffered Saline (PBS), Tryptic Soy Broth (TSB), cell culture media (DMEM), and human saliva. Finally, we used the data to assess the suitability of using bacteriophage Phi6 as a surrogate for SARS-CoV-2 in studies of virus persistence in water and on surfaces.

METHODS

Bacteriophage Phi6 production and enumeration

Bacteriophage Phi6 (DSM 21518) and its host *Pseudomonas syringae* (DSM 21482) were obtained from the DSMZ (Deutsche Sammlung von Mikroorganismen und Zellkulturen). To culture the bacteriophage Phi6, we used a protocol adapted from Pitol *et al*. 2017 [35]. Briefly, 100 mL of Tryptic Soy Broth (TSB; Millipore) containing log-phase *P. syringae* was inoculated with 100 µL of a 10^8 Plaque Forming Units (PFU) mL^{-1} stock of bacteriophage Phi6 and incubated overnight. The following day, the media was centrifuged for 15 min at 5000 rpm and the supernatant was filtered using a 0.45 µm filter unit. Aliquots of the supernatant with a final concentration of 1.5 x10^{11} PFU mL^{-1} were stored at 4°C for subsequent assays.

To enumerate Phi6 we used the standard Double Agar Layer assay[35]. Briefly, soft TSB agar (0.5% Agar) was inoculated with 100 µL of an overnight culture of *P. syringae* and 100 µL of the sample containing Phi6. Samples were mixed and poured on hard agar (1.5%. Agar) TSB plates and incubated at 25°C overnight. Negative controls (100 µL of an overnight culture of *P. syringae* with no bacteriophage) were included in each experiment. All dilutions were quantified in duplicates, and all experiments were performed in triplicates.

SARS-CoV-2 production and enumeration

SARS-CoV-2, Delta variant, passage 4 (SARS-CoV-2/human/GBR/Liv_273/2021, OK392641)[36] was amplified and quantified using Vero E6 cells (African green monkey kidney cells, Public Health England). Vero cells were maintained at 37°C and 5% CO_2 in Dulbecco's Minimal Essential Medium (DMEM; Corning) supplemented with 10 % Foetal Bovine Serum (FBS; Sigma-Aldrich) and 0.05 mg mL^{-1} of gentamicin (Gibco). To amplify SARS-CoV-2, a flask T-150 of confluent Vero E6 was inoculated with 20 µL of a 10^6 PFU mL^{-1} stock of SARS-CoV-2 Delta variant, passage 4, and incubated for 72h. Subsequently,

the media was recovered and centrifuged for 15 min at 5000 rpm to remove the remaining cells and cell debris. Finally, the stock solution, SARS-CoV-2 passage 5 at a concentration of 10^7 PFU mL^{-1}, was concentrated using the Amicon Ultra Centrifugal Filter (100kDa; Merk Millipore Amicon) to a final concentration of ~10^8 PFU mL^{-1}.

Standard plaque assay was performed to quantify infectious virus as previously described [37]. Briefly, samples were serially diluted and inoculated in a confluent monolayer of Vero E6 cells. One hour after infection, an agarose media overlay (2% Agarose in DMEM supplemented with 2% FBS) was applied to the cell monolayer and incubated at 37°C and 5% CO_2 for 72h. Subsequently, the cells were fixed with Formalin 10% (VWR International) and stained with Crystal Violet (Sigma-Aldrich) and plaques were counted. All experiments were conducted in a containment level 3 laboratory by personnel trained in the relevant code of practices and standard operating procedures.

Surfaces and deposition solutions

To study the persistence of SARS-CoV-2 and Phi6 on surfaces, each virus was suspended in four different deposition solutions to a final concentration of ~10^6 PFU mL^{-1}. The solutions used were: Phosphate Buffered Saline (PBS; Gibco), Tryptic Soy Broth (TSB; Millipore), DMEM supplemented with 2 % FBS, and human saliva. Saliva was collected at Unilever Research Port Sunlight as described elsewhere [36,38]. Briefly, healthy donors provided a stimulated daytime saliva sample for which they were given a piece of gum to chew (Wrigley's Turbulence). Subjects were given a maximum of five sterile 30 ml containers in which they were asked to provide a 20–25 ml sample of saliva per container. Subjects were requested to leave 30 min after eating or drinking before providing a saliva sample. Saliva samples were stored overnight at -80°C prior sterilization using to gamma irradiation (Systagenix, UK, Cobolt 60 turntable, dose rate 1.2 kGy/h, minimum dose 32.1 kGy). After sterilization, saliva was stored at 4°C until used.

To study virus persistence on surfaces, we selected two non-porous materials: Plastic (PVC plastic vinyl), and metal (stainless steel). Circular coupons of stainless steel and plastic were disinfected by soaking in 70% ethanol (VWR International) for 30 minutes. Subsequently, the coupons were thoroughly rinsed with deionized water before allowing them to dry inside a class II biological safety cabinet for one hour.

Persistence of SARS-CoV-2 and Phi6 on surfaces

The plastic and steel coupons were inoculated by pipetting a 1 µL droplet of deposition solution (PBS, TSB, DMEM, saliva) containing ~10^6 PFU mL^{-1} of virus (SARS-CoV-2 or Phi6) in the centre of the coupon. The inoculated coupons were incubated at 25°C in a container partially opened to allow evaporation of the droplets. Coupons were sampled at different time points by pipetting up and down 15 times using 50 µL cell culture media (DMEM supplemented with 2% FBS) and samples were stored at -80°C before being quantified together at the end of the experiment. Positive controls, consisting of 50 µL DMEM inoculated with 1 µL of virus (SARS-CoV-2 or Phi6) suspended in each of the four viral matrices (PBS, TSB, DMEM, Saliva), were run alongside each treatment. Experiments were repeated three times.

Persistence of SARS-CoV-2 and Phi6 in water

A stock of 10^8 PFU mL^{-1} of virus (SARS-CoV-2 or Phi6) was serially diluted in mineral water (Volvic Natural Mineral Water) to a final concentration of 10^6 PFU mL^{-1} and aliquoted in samples of 50 µL. The samples were incubated at 25°C and recovered at different time points. Subsequently, they were diluted in 200 µL of culture media (DMEM supplemented by 2% FBS) to a final volume of 250 µL, and stored at -80°C, before being quantified. Positive controls, consisting of 200 µL DMEM inoculated with 50 µL of virus (SARS-CoV-2 or Phi6) suspended in water, were collected alongside each treatment. Experiments were repeated three times.

Data analysis

All data analysis was performed using R statistical software (Version 4.1.0). Multiple regression analysis was used to assess the difference in the number of viruses on the water as a function of time and virus (Phi6, SARS-CoV-2). In the experiments which determined virus persistence on surfaces, multiple regression analysis was used to evaluate the survival of viruses as a function of virus, deposition solution (PBS, TSB, DMEM, saliva), surface material (plastic, steel), and time.

Linear regression was used to calculate the first-order decay constant, k (h^{-1}), as the slope of ln (N/N_0) versus time in hours (Equation 1), where N is the number of viruses at time = t, and N_0 is the number of viruses at the beginning of the experiment (t=0). Data with concentrations below the limit of detection (LOD) was excluded from the regression analysis. The decay constant, k, was used to estimate the half-life of the virus, t_{50} (h^{-1}), and the time required for 90% of the reduction in the number of viruses, t_{90} (h^{-1}). The t_{50} and the t_{90} were estimated using equations 2 and 3 respectively.

(Equation 1)
$$\ln \frac{N}{N_0} = -kt$$

(Equation 2)
$$t_{50} = \frac{\ln 2}{-k}$$

(Equation 3)
$$t_{90} = \frac{\ln 10}{-k}$$

RESULTS AND DISCUSSION

SARS-CoV-2 and Phi6 survival in water

A multiple linear regression (MLR) was calculated to predict the concentration of viruses in the water as a function of time and virus type ($F_{(2,59)}$ = 29.73, $p<0.001$), $R^2=0.49$). The persistence of the viruses in water was significantly influenced by the virus used (MLR, $p<0.001$). The half-life of bacteriophage Phi6 was 27 hours as compared with 15 hours for SARS-CoV-2 (Table 2, Figure 2). Therefore, bacteriophage Phi6 significantly overestimated the persistence of SARS-CoV-2 in the water. The decay rates reported here for both viruses, Phi6 and SARS-CoV-2, fall within the range of data reported elsewhere[33,39]. The time taken to achieve a 90% reduction in the titter (t_{90}) for SARS-CoV-2 in mineral water was 50 hours. Our findings were comparable to other studies using autoclaved or filtered river water and tap water at similar temperatures (20-25°C), t_{90}= 48-79.2 hours [14,15,40]. Additionally, we obtained a t_{90} of 89 hours for Phi6, which is consistent with previous research that shows a range of t_{90} between 74 and 179 hours for autoclaved or filtered river water and tap water at similar temperatures [27,33]. The variation observed in virus survival between different studies can be explained by several factors including the chemical composition of the water used in the experiments (e.g., oxygen level, pH, presence of organic matter) and environmental variables such as light exposure and temperature [41]. Therefore, a quantitative comparison between different studies is difficult, which highlights the importance of performing a side-by-side comparison between the viruses when evaluating the suitability of using viral surrogates.

Table 2. Linear Regression Models for Persistence of SARS-CoV-2 and Bacteriophage Phi6 on Water

Virus	Liquid Matrix	t_{90} (h)	t_{50} (h)	k, SE k (h^{-1})
SARS-CoV-2	Mineral water	50.44	15.19	-0.046, 0.004
Phi6	Mineral water	88.93	26.77	-0.026, 0.002

t_{90} : Time required for 90% of the reduction in the number of viruses (hours)
t_{50}: Half-life or time required for 50% of the reduction in the number of viruses (hours)
k: First order decay constant (hours $^{-1}$)

Figure 2. Persistence of SARS-CoV-2 and bacteriophage Phi6 on mineral water. Number of virus per sample are shown in red circles for Phi6 and blue triangles for SARS-CoV-2. The red and blue continuous lines show the fitted linear model for Phi6 and SARS-CoV-2 respectively, with grey areas showing the 95% CI for each regression line. The black dotted line shows the limit of detection for the assays.

Persistence of SARS-CoV-2 and Phi6 on surfaces

MLR was used to predict the concentration of viruses as a function of time, virus type, deposition solution, and surface material ($F_{(6, 376)} = 12.01$, $p<0.001$). Virus type, deposition solution, and time were significant predictors of virus concentration (MLR, virus type $p<0.001$, deposition solution $p<0.001$, time $p<0.001$). Conversely, the influence of surface material (metal vs plastic) on virus persistence was not statistically significant ($p=0.57$). The most marked observation to emerge from the data comparing the survival of SARS-CoV-2 with that of bacteriophage Phi6 on surfaces was that Phi6 survives significantly longer than SARS-CoV-2 (Figure 1, Table 1). The half-life of Phi6 was 403 hours (~2 weeks) on plastic and 250 hours (~10 days) on metal when the deposition solution was TSB. Conversely, inoculating the surfaces with SARS-CoV-2 suspended in TSB resulted in half-lives of 1.3 and 1.9 hours for metal and plastic respectively. Similar patterns of higher persistence of Phi6 as compared with SARS-CoV-2 were obtained when using saliva and DMEM as deposition solutions (Table 1). The half-life of Phi6 on plastic

was 1-3 days when the deposition solution was saliva, as compared to 0.8-1.2 hours for SARS-CoV-2.

Using DMEM as deposition solution, the half-life of Phi6 on plastic and metal were 12.3 and 9.3 hours

respectively, as compared with halve-lives of 1.3 and 0.5 hours for SARS-CoV-2. Interestingly, when

viruses were suspended in PBS, their half-lives were short, regardless of the virus used. Using PBS as a

deposition solution led to Phi6 half-lives of 1.7 hours and 1.5 hours in plastic and metal respectively, as

compared to 4 and 2 hours for SARS-CoV-2.

Figure 1. Virus survival on surfaces as a function of surface material (plastic vs metal), virus type (SARS-CoV-2 vs Phi6) and deposition solution (DMEM, PBS, TSB, and saliva). Data shows the mean and the standard deviation of triplicates. Red dotted line shows the limit of detection of the assays.

Table 1. Summary of the Mean Linear Regression Model Parameters for the Persistence of SARS-CoV-2 and Phi6 on Surfaces.

Virus	Deposition solution	Surface material	T_{90} (h)	T_{50} (h)	k, SE k (h^{-1})
SARS-CoV-2	DMEM	plastic	4.24	1.28	-0.543, 0.125 *
SARS-CoV-2	TSB	plastic	6.35	1.91	-0.363, 0.080
SARS-CoV-2	PBS	plastic	4.27	1.29	-0.539, 0.042
SARS-CoV-2	Saliva	plastic	4.13	1.24	-0.557, 0.099
SARS-CoV-2	DMEM	metal	1.63	0.49	-1.414, 0.086 *
SARS-CoV-2	TSB	metal	4.32	1.30	-0.533, 0.063
SARS-CoV-2	PBS	metal	2.00	0.60	-1.150, 0.223*
SARS-CoV-2	Saliva	metal	2.68	0.81	-0.859, 0.138
Phi6	DMEM	plastic	40.81	12.28	-0.056, 0.031
Phi6	TSB	plastic	1338.77	403.01	-0.002, 0.001
Phi6	PBS	plastic	1.66	0.50	-1.389, 0.000*
Phi6	Saliva	plastic	268.96	80.96	-0.009, 0.002
Phi6	DMEM	metal	31.03	9.34	-0.074, 0.009
Phi6	TSB	metal	829.92	249.83	-0.003, 0.001
Phi6	PBS	metal	1.51	0.46	-1.520, 0.086
Phi6	Saliva	metal	93.92	28.27	-0.025, 0.003

*Only 2 time points were considered in the regression, as the virus was inactivated within 6 hours.

In addition to the higher persistence of bacteriophage Phi6 over SARS-CoV-2, our data indicate that the deposition solution had a significant influence on virus survival (Table 1, Figure 1). This finding is consistent with previous research that suggest that the survival of viruses, including SARS-CoV-2, is highly influenced by the liquid matrix used to suspend the virus[19,42]. Pastorino *et al.* showed that higher protein content in the deposition solution increases viral persistence on surfaces[19]. In the present study we used four deposition solutions: TSB (20 grams of protein per litter, g L^{-1}), human saliva (0.7 - 2.5 g L^{-1}, based on data by Lin *et al* [43]), DMEM supplemented with 2% FBS (~1.2 g L^{-1}), and PBS (0 g L^{-1}). Our data showed that

bacteriophage Phi6 survives the longest in TSB, followed by saliva, DMEM, and lastly PBS, which correlates with protein concentration of the deposition solution. Other components within deposition solutions that have been shown to influence the stability of the viruses is the pH [42,44], salt concentration [42,45], and the presence of polysaccharides [46] in the solution.

Our data on SARS-CoV-2 survival on surfaces are consistent with multiple research studies that showed that the half-life of SARS-CoV-2 at 20-27°C was between 1.5 and 9 hours in plastic surfaces [10,13,16,47], and between 3.4 and 7.8 hours in stainless steel [9,13,16]. Nevertheless, other studies have shown a longer survival rate for SARS-CoV-2 on surfaces at similar temperatures, with half-lives in the order of days rather than hours [17,48]. Differences between the studies include using different deposition solutions, inoculum volumes, light exposure during the experiment, and drying times. Since factors such as temperature, light exposure, and experimental variables such as the composition of the deposition solution used to inoculate the surfaces play a crucial role in the survival of viruses, it is desirable for future research to carefully select these factors when evaluating the survival of emerging pathogens in the environment.

We acknowledge our study has limitations, which are inherent when working with a category 3 pathogen such as SARS-CoV-2. For example, due to biosafety requirements, samples were placed in a container inside an incubator with no light or active air movement, which are factors that have been shown to influence virus survival[41,49]. Therefore, the decay of the viruses presented here may not adequately reflect the decay that viruses have in scenarios with sunlight exposure and air movement. Additionally, experiments were performed at ambient humidity, which fluctuated between 25 and 50%. Temperature and humidity have shown to have an influence on the persistence of SARS-CoV-2 on the surfaces [13,18] . For example, one study showed that the half-life of SARS-CoV-2 in stainless steel ranges from 3 to 70 hours, depending on the environmental temperature and humidity [13]. Therefore, adequate control of both temperature and humidity is advisable to when evaluating the persistence of pathogens.

Despite the limitations of this study, our findings highlight the importance of evaluating the suitability of using viral surrogates by performing side-by-side comparisons with the pathogen of interest to control all variables that could potentially influence the outcome. Our results showed that, although it has been frequently used as a surrogate for coronaviruses, bacteriophage Phi6 is not an adequate surrogate for studies quantifying SARS-CoV-2 persistence, as it over-estimates infectiousness. Additionally, our findings reveal the need of using adequate deposition solutions when evaluating viral persistence on surfaces. Future research on the persistence of pathogens should place careful consideration on the methodology used, selecting a deposition solution, inoculation method, and environmental parameters that adequately mimic real-life scenarios.

ACKNOWLEDGMENT

Carole Philpotts (Unilever) is thanked for her assistance in supply of saliva. This work was supported by UKRI grants (20197 and 85336) awarded to GLH. GLH was further supported by the BBSRC (BB/T001240/1, BB/V011278/1, and BB/W018446/1), the UKRI (85336), the EPSRC (V043811/1), a Royal Society Wolfson Fellowship (RSWF\R1\180013), the NIHR (NIHR2000907) and the Bill and Melinda Gates Foundation (INV-048598). This work was additionally supported by a Pandemic Institute grant (TPIMPX01) awarded to AKP. The Pandemic Institute is formed of seven founding partners: The University of Liverpool, Liverpool School of Tropical Medicine, Liverpool John Moores University, Liverpool City Council, Liverpool City Region Combined Authority, Liverpool University Hospital Foundation Trust, and Knowledge Quarter Liverpool. The views expressed are those of the author(s) and not necessarily those of the Pandemic Institute.

REFERENCES

1. National Center for Immunization and Respiratory Diseases (NCIRD), Division of Viral Diseases. Scientific Brief: SARS-CoV-2 Transmission. *CDC COVID-19 Science Briefs*. Published online May 7, 2021. Accessed November 18, 2022. https://www.ncbi.nlm.nih.gov/books/NBK570442/

2. Harvey AP, Fuhrmeister ER, Cantrell ME, et al. Longitudinal Monitoring of SARS-CoV-2 RNA on High-Touch Surfaces in a Community Setting. *Environ Sci Technol Lett*. 2020;8(2):168-175. doi:10.1021/acs.estlett.0c00875

3. Amoah ID, Pillay L, Deepnarian N, et al. Detection of SARS-CoV-2 RNA on contact surfaces within shared sanitation facilities. *Int J Hyg Environ Health*. 2021;236:113807. doi:10.1016/J.IJHEH.2021.113807

4. Abrahão JS, Sacchetto L, Rezende IM, et al. Detection of SARS-CoV-2 RNA on public surfaces in a densely populated urban area of Brazil: A potential tool for monitoring the circulation of infected patients. *Science of The Total Environment*. 2021;766:142645. doi:10.1016/J.SCITOTENV.2020.142645

5. Gonçalves J, da Silva PG, Reis L, et al. Surface contamination with SARS-CoV-2: A systematic review. *Science of The Total Environment*. 2021;798:149231. doi:10.1016/J.SCITOTENV.2021.149231

6. Moore G, Rickard H, Stevenson D, et al. Detection of SARS-CoV-2 within the healthcare environment: a multi-centre study conducted during the first wave of the COVID-19 outbreak in England. *Journal of Hospital Infection*. 2021;108:189-196. doi:10.1016/J.JHIN.2020.11.024

7. Peccia J, Zulli A, Brackney DE, et al. Measurement of SARS-CoV-2 RNA in wastewater tracks community infection dynamics. *Nature Biotechnology 2020 38:10*. 2020;38(10):1164-1167. doi:10.1038/s41587-020-0684-z

8. Geng Y, Wang Y. Stability and transmissibility of SARS-CoV-2 in the environment. *J Med Virol*. 2023;95(1):e28103. doi:10.1002/JMV.28103

9.   Bonil L, Lingas G, Coupeau D, et al. Survival of SARS-CoV-2 on Non-Porous Materials in an Experimental Setting Representative of Fomites. *Coatings 2021, Vol 11, Page 371*. 2021;11(4):371. doi:10.3390/COATINGS11040371

10.   Matson MJ, Yinda CK, Seifert SN, et al. Effect of Environmental Conditions on SARS-CoV-2 Stability in Human Nasal Mucus and Sputum. *Emerg Infect Dis*. 2020;26(9):2276. doi:10.3201/EID2609.202267

11.   Raiteux J, Eschlimann M, Marangon A, et al. Inactivation of SARS-CoV-2 by Simulated Sunlight on Contaminated Surfaces. *Microbiol Spectr*. 2021;9(1). doi:10.1128/SPECTRUM.00333-21

12.   Harbourt DE, Haddow AD, Piper AE, et al. Modeling the stability of severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) on skin, currency, and clothing. *PLoS Negl Trop Dis*. 2020;14(11):e0008831. doi:10.1371/JOURNAL.PNTD.0008831

13.   Kwon T, Gaudreault NN, Richt JA, Vilibic-Cavlek T, Savic V. Environmental Stability of SARS-CoV-2 on Different Types of Surfaces under Indoor and Seasonal Climate Conditions. *Pathogens 2021, Vol 10, Page 227*. 2021;10(2):227. doi:10.3390/PATHOGENS10020227

14.   de Oliveira LC, Torres-Franco AF, Lopes BC, et al. Viability of SARS-CoV-2 in river water and wastewater at different temperatures and solids content. *Water Res*. 2021;195:117002. doi:10.1016/J.WATRES.2021.117002

15.   Sala-Comorera L, Reynolds LJ, Martin NA, O'Sullivan JJ, Meijer WG, Fletcher NF. Decay of infectious SARS-CoV-2 and surrogates in aquatic environments. *Water Res*. 2021;201:117090. doi:10.1016/J.WATRES.2021.117090

16.   van Doremalen N, Bushmaker T, Morris DH, et al. Aerosol and Surface Stability of SARS-CoV-2 as Compared with SARS-CoV-1. *N Engl J Med*. Published online 2020:1-3. doi:10.1056/NEJMc2004973

17.   Riddell S, Goldie S, Hill A, Eagles D, Drew TW. The effect of temperature on persistence of SARS-CoV-2 on common surfaces. *Virol J*. 2020;17(1):1-7. doi:10.1186/S12985-020-01418-7/FIGURES/2

18. Biryukov J, Boydston JA, Dunning RA, et al. Increasing Temperature and Relative Humidity Accelerates Inactivation of SARS-CoV-2 on Surfaces. *mSphere*. 2020;5(4). doi:10.1128/mSphere.00441-20

19. Pastorino B, Touret F, Gilles M, de Lamballerie X, Charrel RN. Prolonged Infectivity of SARS-CoV-2 in Fomites. *Emerg Infect Dis*. 2020;26(9):2256. doi:10.3201/EID2609.201788

20. String GM, White MR, Gute DM, Mühlberger E, Lantagne DS. Selection of a SARS-CoV-2 Surrogate for Use in Surface Disinfection Efficacy Studies with Chlorine and Antimicrobial Surfaces. *Environ Sci Technol Lett*. 2021;8:995-1001. doi:doi.org/10.1021/acs.estlett.1c00593

21. Gomes M, Bartolomeu M, Vieira C, et al. Photoinactivation of Phage Phi6 as a SARS-CoV-2 Model in Wastewater: Evidence of Efficacy and Safety. *Microorganisms 2022, Vol 10, Page 659*. 2022;10(3):659. doi:doi.org/10.3390/microorganisms10030659

22. Baker CA, Gutierrez A, Gibson KE. Factors Impacting Persistence of Phi6 Bacteriophage, an Enveloped Virus Surrogate, on Fomite Surfaces. *Appl Environ Microbiol*. 2022;88(7). doi:10.1128/AEM.02552-21/SUPPL_FILE/AEM.02552-21-S0001.PDF

23. Baker CA, Gibson KE. Phi 6 recovery from inoculated fingerpads based on elution buffer and methodology. *J Virol Methods*. 2022;299:114307. doi:10.1016/J.JVIROMET.2021.114307

24. Fedorenko A, Grinberg M, Orevi T, Kashtan N. Survival of the enveloped bacteriophage Phi6 (a surrogate for SARS-CoV-2) in evaporated saliva microdroplets deposited on glass surfaces. *Scientific Reports 2020 10:1*. 2020;10(1):1-10. doi:10.1038/s41598-020-79625-z

25. Gallandat K, Lantagne D. Selection of a Biosafety Level 1 (BSL-1) surrogate to evaluate surface disinfection efficacy in Ebola outbreaks: Comparison of four bacteriophages. *PLoS One*. 2017;12(5):e0177943. doi:10.1371/JOURNAL.PONE.0177943

26. Aranha-Creado H, Brandwein H. Application of Bacteriophages as Surrogates for Mammalian Viruses: A Case for Use in Filter Validation Based on Precedents and Current Practices in Medical and Environmental Virology. *PDA J Pharm Sci Technol*. 1999;53(2).

27. Aquino De Carvalho N, Stachler EN, Cimabue N, Bibby K. Evaluation of Phi6 Persistence and Suitability as an Enveloped Virus Surrogate. *Environ Sci Technol*. 2017;51(15):8692-8700. doi:10.1021/ACS.EST.7B01296/SUPPL_FILE/ES7B01296_SI_001.PDF

28. Vidaver AK, Koski RK, Etten JL van. Bacteriophage φ6: a Lipid-Containing Virus of Pseudomonas phaseolicola. *J Virol*. 1973;11(5):799-805. doi:10.1128/JVI.11.5.799-805.1973

29. Ye Y, Ellenberg RM, Graham KE, Wigginton KR. Survivability, Partitioning, and Recovery of Enveloped Viruses in Untreated Municipal Wastewater. *Environ Sci Technol*. 2016;50(10):5077-5085. doi:10.1021/ACS.EST.6B00876/ASSET/IMAGES/LARGE/ES-2016-00876F_0005.JPEG

30. Kim K, Jothikumar N, Sen A, Murphy JL, Chellam S. Removal and inactivation of an enveloped virus surrogate by iron conventional coagulation and electrocoagulation. *Environ Sci Technol*. 2021;55(4):2674-2683. doi:10.1021/ACS.EST.0C07697/SUPPL_FILE/ES0C07697_SI_001.PDF

31. Prussin AJ, Schwake DO, Lin K, Gallagher DL, Buttling L, Marr LC. Survival of the enveloped virus Phi6 in droplets as a function of relative humidity, absolute humidity, and temperature. *Appl Environ Microbiol*. 2018;84(12). doi:10.1128/AEM.00551-18/ASSET/88D4C88A-F78E-43D8-A182-4B5D2920F76C/ASSETS/GRAPHIC/ZAM0121885520005.JPEG

32. Whitworth C, Mu Y, Houston H, et al. Persistence of bacteriophage phi 6 on porous and nonporous surfaces and the potential for its use as an ebola virus or coronavirus surrogate. *Appl Environ Microbiol*. 2020;86(17). doi:10.1128/AEM.01482-20/ASSET/47BA9BA8-F156-4290-8C1F-B48A89070141/ASSETS/GRAPHIC/AEM.01482-20-F0002.JPEG

33. Silverman AI, Boehm AB. Systematic Review and Meta-Analysis of the Persistence and Disinfection of Human Coronaviruses and Their Viral Surrogates in Water and Wastewater. *Environ Sci Technol Lett*. 2020;7(8):544-553. doi:10.1021/ACS.ESTLETT.0C00313/ASSET/IMAGES/LARGE/EZ0C00313_0004.JPEG

34. Anderson CE, Boehm AB. Transfer Rate of Enveloped and Nonenveloped Viruses between Fingerpads and Surfaces. *Appl Environ Microbiol*. 2021;87(22). doi:10.1128/AEM.01215-21/SUPPL_FILE/AEM.01215-21-S0001.PDF

35.   Pitol AK, Bischel HN, Kohn T, Julian TR. Virus Transfer at the Skin-Liquid Interface. *Environ Sci Technol*. 2017;51(24):14417-14425. doi:10.1021/acs.est.7b04949

36.   Anderson ER, Patterson EI, Richards S, et al. CPC-containing oral rinses inactivate SARS-CoV-2 variants and are active in the presence of human saliva. *J Med Microbiol*. 2022;71(2):1508. doi:10.1099/JMM.0.001508

37.   Anderson ER, Hughes GL, Patterson EI. Inactivation of SARS-CoV-2 on surfaces and in solution with Virusend (TX-10), a novel disinfectant. *Access Microbiol*. 2021;3(4). doi:10.1099/ACMI.0.000228

38.   Dawes C, Tsang RWL, Suelzle T. The effects of gum chewing, four oral hygiene procedures, and two saliva collection techniques, on the output of bacteria into human whole saliva. *Arch Oral Biol*. 2001;46(7):625-632. doi:10.1016/S0003-9969(01)00017-6

39.   Mahlknecht J. Presence and persistence of SARS-CoV-2 in aquatic environments: A mini-review. *Curr Opin Environ Sci Health*. 2022;29:100385. doi:10.1016/J.COESH.2022.100385

40.   Bivins A, Greaves J, Fischer R, et al. Persistence of SARS-CoV-2 in Water and Wastewater. *Environ Sci Technol Lett*. 2020;7(12):937-942. doi:10.1021/acs.estlett.0c00730

41.   Pinon A, Vialette M. Survival of Viruses in Water. *Intervirology*. 2018;61(5):214-222. doi:10.1159/000484899

42.   Lin K, Schulte CR, Marr LC. Survival of MS2 and Φ6 viruses in droplets as a function of relative humidity, pH, and salt, protein, and surfactant concentrations. *PLoS One*. 2020;15(12):e0243505. doi:10.1371/JOURNAL.PONE.0243505

43.   Lin LY, Chang CC. [Determination of protein concentration in human saliva]. *Gaoxiong Yi Xue Ke Xue Za Zhi*. 1989;5(7):389-397. Accessed November 20, 2022. https://europepmc.org/article/med/2810445

44.   Luo B, Schaub A, Glas I, et al. Acidity of expiratory aerosols controls the infectivity of airborne influenza virus and SARS-CoV-2. *medRxiv*. Published online June 21, 2022:2022.03.14.22272134. doi:10.1101/2022.03.14.22272134

45.   Lin K, Marr LC. Humidity-Dependent Decay of Viruses, but Not Bacteria, in Aerosols and Droplets Follows Disinfection Kinetics. *Environ Sci Technol*. 2020;54(2):1024-1032. doi:10.1021/ACS.EST.9B04959/SUPPL_FILE/ES9B04959_SI_001.PDF

46.   Malenovská H. The influence of stabilizers and rates of freezing on preserving of structurally different animal viruses during lyophilization and subsequent storage. *J Appl Microbiol*. 2014;117(6):1810-1819. doi:10.1111/JAM.12654

47.   Johnson GT, Loehle C, Zhou SS, et al. Evaluation of the Survivability of SARS-CoV-2 on Cardboard and Plastic Surfaces and the Transferability of Virus from Surface to Skin. *Occupational Diseases and Environmental Medicine*. 2021;9(2):63-73. doi:10.4236/ODEM.2021.92006

48.   Liu Y, Li T, Deng Y, et al. Stability of SARS-CoV-2 on environmental surfaces and in human excreta. *J Hosp Infect*. 2021;107:105. doi:10.1016/J.JHIN.2020.10.021

49.   Marzoli F, Bortolami A, Pezzuto A, et al. A systematic review of human coronaviruses survival on environmental surfaces. *Science of The Total Environment*. 2021;778:146191. doi:10.1016/J.SCITOTENV.2021.146191