N-WASP-dependent branched actin polymerization attenuates B-cell receptor signaling by increasing the molecular density of receptor clusters

Anshuman Bhanja[1][†], Margaret K. Seeley-Fallen[1], Michelle Lazzaro[1][^], Arpita Upadhyaya[2,3,4], and Wenxia Song[1]*

[1]Department of Cell Biology and Molecular Genetics, University of Maryland, College Park, MD 20742, USA

[2]Biophysics Program, University of Maryland, College Park, MD, 20742, USA.

[3]Department of Physics, University of Maryland, College Park, MD, 20742, USA. arpitau@umd.edu.

[4]Institute for Physical Science and Technology, University of Maryland, College Park, MD, 20742, USA.

[†]Current address: Department of Pharmacology, University of California, San Diego, CA, 92034, USA

[^]Current address: AstraZeneca, Gaithersburg, MD 20878, United States

[*]Authors for correspondence: wenxsong@umd.edu and arpitau@umd.edu

Abstract

Antigen-induced B-cell receptor (BCR) signaling is critical for initiating and regulating B-cell activation. The actin cytoskeleton plays essential roles in BCR signaling. Upon encountering cell-surface antigens, actin-driven B-cell spreading amplifies signaling, while B-cell contraction following spreading leads to signal attenuation. However, the mechanism by which actin dynamics switch BCR signaling from amplification to attenuation is unknown. Here, we show that Arp2/3-mediated branched actin polymerization is required for B-cell contraction. Contracting B-cells generate centripetally moving actin foci from lamellipodial F-actin networks in the B-cell plasma membrane region contacting antigen-presenting surfaces. Actin polymerization driven by N-WASP, but not WASP, initiates these actin foci and facilitates non-muscle myosin II recruitment to the contact zone, creating actomyosin ring-like structures. Furthermore, B-cell contraction increases BCR molecular density in individual clusters, leading to decreased BCR phosphorylation. Increased BCR molecular density reduced levels of the stimulatory kinase Syk, the inhibitory phosphatase SHIP-1, and their phosphorylated forms in individual BCR clusters. These results suggest that N-WASP-activated Arp2/3, coordinating with myosin, generates centripetally moving foci and contractile actomyosin ring-like structures from lamellipodial networks, enabling contraction. B-cell contraction attenuates BCR signaling by pushing out both stimulatory kinases and inhibitory phosphatases from BCR clusters, providing novel insights into actin-facilitated signal attenuation.

Introduction

B-cell-mediated antibody responses are essential for eliminating invading pathogens. B-cell receptors (BCRs) expressed on the B-cell surface detect the presence of cognate antigens. The binding of antigen to the BCR leads to the activation of signaling cascades (Reth and Wienands 1997; Dal Porto *et al.* 2004; Kwak *et al.* 2019), which induce transcriptional programs that prepare B-cells for proliferation and differentiation (Kurosaki *et al.* 2010; Shlomchik *et al.* 2019; Wang *et al.* 2020). BCR signaling also induces rapid antigen internalization, processing, and presentation for T-cell recognition, which provides the second signal required for B-cell clonal expansion and differentiation to high-affinity antibody-secreting cells and memory B-cells (Song *et al.* 1995; Gitlin *et al.* 2014). BCR signaling is tightly regulated by various external and internal factors to activate antibody responses that are specific and also qualitatively and quantitatively matched to the encountered antigen.

Antigen-induced receptor reorganization activates BCRs at the B-cell surface. Antigen binding leads to BCR clustering at lipid rafts, which enables raft-resident kinases, such as the Src kinases Lyn and Fyn, to phosphorylate immunoreceptor tyrosine-based activation motifs (ITAMs) in the cytoplasmic domains of the BCR signaling subunit CD79a/b heterodimer (Reth 1994; Pierce 2002; Sohn *et al.* 2008). Doubly phosphorylated ITAMs recruit and activate spleen tyrosine kinase (Syk), which in turn activates multiple downstream signaling pathways, including phospholipase Cγ2 (PLCγ2), Bruton's tyrosine kinase (Btk), Ras-GTPase, and phosphatidylinositol-3 kinase (PI3K), initiating signaling cascades (Kurosaki 2000; Dal Porto *et al.* 2004; Tanaka and Baba 2020). Signaling activation also induces the recruitment and activation of inhibitory phosphatases, such as SH2-containing tyrosine phosphatase-1 (SHP-1) and phosphatidylinositol-5 phosphatase-1 (SHIP-1), to BCR signaling complexes (Brauweiler *et al.* 2000; Gross *et al.* 2009; Franks and Cambier 2018). SHIP-1 and SHP-1 negatively regulate

BCR signaling by inactivating the plasma membrane docking lipids for stimulatory kinases, such

as Btk and Akt (Aman *et al.* 1998; Bolland *et al.* 1998) and dephosphorylating BCR and its

downstream signaling molecules (Mizuno *et al.* 2000; Adachi *et al.* 2001), respectively. The

interplay between the stimulatory kinases and the inhibitory phosphatases controls the balance

of antibody responses against pathogens and self. Deficiencies in stimulatory kinases by

mutations, such as Btk, cause X-linked agammaglobulinemia (XLA) (Kinnon *et al.* 1993), while

deficiencies in the inhibitory phosphatases SHP-1 or SHIP-1 result in autoimmune diseases

(Pao *et al.* 2007; Leung *et al.* 2013). However, the mechanisms by which BCR signaling

attenuation is initiated and regulated remain elusive.

B-cells encounter both soluble and membrane-associated antigens *in vivo*. Membrane-

associated antigens include antigens on antigen-presenting cells, like follicular dendritic cells in

B-cell follicles or germinal centers, and pathogenic cells, like bacteria, parasites, and cancer

cells (Batista and Harwood 2009; Depoil *et al.* 2009; Gonzalez *et al.* 2009; Cyster 2010). The

binding of multi-valent soluble antigen and membrane-associated antigen with any valency

induces dynamic reorganization of surface BCRs into microclusters, triggering BCR signaling.

Subsequently, surface BCRs continue moving to the plasma membrane region contacting

antigen-presenting surfaces or to one B-cell pole in the case of soluble antigen, which leads to

the growth and merger of BCR microclusters and the formation of immunological synapses

(Carrasco *et al.* 2004; Harwood and Batista 2010) or supra-molecular activation complexes

(Unanue and Karnovsky 1973; Schreiner and Unanue 1977; Tolar *et al.* 2009b; Harwood and

Batista 2010).

Multiple mechanisms have been proposed for signaling initiation by antigen-induced surface

BCR reorganization. The binding of surface BCRs to membrane-associated antigen has been

shown to induce a conformational change in the BCR extracellular domain, exposing a proximal

membrane region of membrane IgM that promotes receptor clustering (Tolar *et al.* 2009a; Shen

et al. 2019). Antigen-binding also changes the conformation of the BCR cytoplasmic domains

from a closed to an open form, facilitating the recruitment of signaling molecules (Tolar *et al.*

2005). Alternatively, antigen-BCR interaction can increase the molecular spacing of BCRs in

clusters, facilitating BCR interaction with signaling molecules (Yang and Reth 2010; Kläsener *et*

al. 2014). The inhibitory phosphatases SHP-1 and SHIP-1 are also recruited to BCR clusters

(Adachi *et al.* 2001; Seeley-Fallen *et al.* 2014). However, the mechanisms underlying

phosphatase recruitment and its relationship with BCR signaling activation complexes are

unknown.

The actin cytoskeleton is essential for antigen-induced BCR reorganization on the B-cell surface

in response to both soluble and membrane-associated antigen (Harwood and Batista 2011; Liu

et al. 2013b; Song *et al.* 2013; Hoogeboom and Tolar 2016). Early signaling of the BCR triggers

transient actin depolymerization via Rap GTPase and its downstream target cofilin (Freeman *et*

al. 2011), which disassembles the existing cortical actin network that confines the lateral

movement of surface BCRs (Treanor *et al.* 2010). Following this transient depolymerization,

BCR signaling activates rapid actin polymerization through the actin nucleation promoting

factors Wiskott-Aldrich Syndrome Protein (WASP) and neuronal-WASP (N-WASP), modulating

BCR mobility and leading to F-actin accumulation at BCR-antigen interaction sites (Liu *et al.*

2013a; Rey-Suarez *et al.* 2020). Actin polymerization and treadmilling in B-cells stimulated by

soluble antigen drive surface BCRs to cluster and move to one pole of the B-cell, forming a BCR

cap (Schreiner and Unanue 1977; Liu *et al.* 2012a). Upon interacting with membrane-associated

antigen, B-cells organize actin into two dynamic structures. One treadmills outwards, driving B-

cell membrane spreading and expanding the B-cell membrane region contacting the antigen-

presenting surface (contact zone), which enables more BCRs to engage antigen (Bolger-Munro

et al. 2019). The other creates retrograde flow towards the center of the B-cell contact zone,

driving the centripetal movement and growth of BCR microclusters (Liu *et al.* 2012b). Together,

these processes amplify BCR signaling (Batista *et al.* 2010; Harwood and Batista 2011). The

extent of B-cell spreading and the amount of BCR-antigen complexes gathered in the contact

zone depend on BCR binding affinity and the density of antigen on the membrane (Fleire *et al.*

2006). Subsequent to spreading, B-cells contract, reducing the contact area, which drives BCR

microclusters to merge into central clusters to form immunological synapses (Fleire *et al.* 2006;

Liu *et al.* 2013a).

The actin cytoskeleton closely interplays with BCR signaling. We have previously shown that

mouse primary B-cells with Btk deficiency (Liu *et al.* 2011) or double knockouts of the actin

nucleation promoting factors WASP and N-WASP (Liu *et al.* 2013a) fail to spread on antigen-

presenting surfaces and establish stable interactions with membrane-associated antigen,

drastically reducing BCR signaling. Btk can activate WASP and N-WASP through Vav, the

guanine nucleotide exchange factor of Cdc42 and Rac and phosphatidylinositol-5 kinase that

generates phosphatidylinositol-4,5-biphosphate (PI4,5P$_2$) (Sharma *et al.* 2009; Padrick and

Rosen 2010). Surprisingly, B-cell-specific knockout of N-WASP (cNKO) but not WASP germline

knockout (WKO) enhances B-cell spreading, delays B-cell contraction, and inhibits the

centralization of BCR clusters at the contact zone (Liu *et al.* 2013a). Consistent with these

findings, Bolger-Munro et al. have shown that knockdown of the actin nucleation factor Arp2/3

downstream of WASP and N-WASP disrupts the dynamic actin reorganization induced by

membrane-associated antigen required for BCR microcluster growth and merger (Bolger-Munro

et al. 2019). Using B-cell-specific and germline knockouts, we showed that in addition to N-

WASP, the actin motor non-muscle myosin IIA (NMIIA) (Seeley-Fallen *et al.* 2022) and the

actin-binding adaptor protein Abp1 (Seeley-Fallen *et al.* 2014) are required for B-cell contraction

and BCR central cluster formation. Finally, Wang et al. recently showed that in the presence of

the adhesion molecule ICAM on the antigen-presenting surface, B-cells form contractile

actomyosin arcs, driving centripetal movement of BCR clusters in the B-cell contact zone (Wang

et al. 2022). Together, these findings indicate a critical role of different actin networks in B-cell

contraction and BCR signaling.

Our previous studies demonstrate that B-cell contraction is critical for BCR signaling

attenuation. Delay or inhibition of B-cell contraction due to deficiencies in actin regulators, N-

WASP, Abp1, or NMIIA, prolongs and/or increases BCR signaling by enhancing the activation of

stimulatory kinases and suppressing the activation of inhibitory phosphatases, which elevates

the production of autoantibody levels in mice (Liu *et al.* 2013a; Seeley-Fallen *et al.* 2014;

Seeley-Fallen *et al.* 2022). N-WASP and NMIIA are not known to interact with kinases and

phosphatases directly. The growth, merger, and centralization of antigen-BCR complexes at the

B-cell contact zone during the B-cell contraction phase are associated with reduced BCR

signaling capability (Liu *et al.* 2013a). These findings suggest that B-cell contraction is one of

the mechanisms for inducing BCR signaling attenuation. However, how the actin cytoskeleton

facilitates the transition of the spreading to contraction phase and how B-cell contraction

switches BCR signaling from amplification to attenuation is unknown.

This study explores the mechanisms by which the actin cytoskeleton remodels from spreading

lamellipodia to contractile structures and by which B-cell contraction suppresses BCR signaling.

Our results show that the contractile actomyosin structures responsible for B-cell contraction

originate from branched actin at spreading lamellipodia. N-WASP/Arp2/3- mediated actin

polymerization prolongs the lifetime of the lamellipodial actin structures, enables NMIIA

recruitment, and drives their movement to the center of the B-cell contact zone. B-cell

contraction increases the molecular density within individual BCR-antigen clusters, which

promotes the disassociation of both the stimulatory kinase Syk and the inhibitory phosphatases

SHIP-1 from BCR clusters, leading to signal attenuation. Our study reveals a new mechanism

underlying BCR signal downregulation.

Results

Arp2/3, activated by N-WASP but not WASP, is required for B-cell contraction

Arp2/3-generated branched F-actin is known to drive lamellipodial expansion for B-cell

spreading (Bolger-Munro *et al.* 2019). However, whether branched F-actin is involved in the

subsequent contraction phase is unknown. To address this, we perturbed the polymerization of

branched F-actin using the Arp2/3 inhibitor CK-666 (50 µM) while using its inactive derivative

CK-689 as a control. Pre-warmed splenic B-cells from WT C57BL/6 mice were incubated with

monobiotinylated Fab' fragment of anti-mouse IgM+G attached to planar lipid bilayers (Fab'-

PLB) by biotin-streptavidin interaction and imaged live at 37°C using interference reflection

microscopy (IRM). The contact area of B-cells treated with CK-689 rapidly increased upon

contacting Fab'-PLB and reached a maximum at ≥0.5 min after the initial contact (*Figure 1A,*

Figure 1-figure supplement 1, and Figure 1-Video 1A). Following maximal spreading, most B-

cells reduced the area of their contact zone, indicating contraction (*Figure 1A, Figure 1-figure*

supplement 1, and Figure 1-Video 1A). We classified a B-cell as contracting if its contact zone

area reduced by ≥5% for at least 10 sec after reaching a maximum value. Based on the average

timing for B-cell maximal spreading, we treated B-cells with CK-666 at the beginning of the

incubation (Time 0), before B-cell spreading initiation, or at 2 min, when all B-cells had already

spread. The effectiveness of CK-666 was detected by reduced Arp2/3 staining in the contact

zone (*Figure 1-figure supplement 2*). As expected, CK-666 treatment at 0 but not 2 min reduced

the kinetics of B-cell spreading (*Figure 1-figure supplement* 3). Importantly, CK-666 treatment at

0 and 2 min both significantly reduced the percentage of B-cells undergoing contraction (*Figure*

1A-C and Figure 1-Video 1A). The inhibitory effect of CK-666 on B-cell contraction, particularly

CK-666 treatment after B-cell spreading, suggests that Arp2/3-mediated branched actin

polymerization is required for B-cell contraction.

WASP and N-WASP are actin nucleation-promoting factors upstream of Arp2/3 that are

expressed in B-cells (Padrick and Rosen 2010). To determine if either or both were responsible

for activating Arp2/3 for B-cell contraction, we utilized the N-WASP inhibitor wiskostatin (Wisko),

and splenic B-cells from B-cell-specific N-WASP knockout mice (cNKO) and germline WASP

knockout (WKO) mice (Westerberg *et al.* 2012; Liu *et al.* 2013a). Wisko has been shown to

inhibit N-WASP activation while enhancing WASP activation in B-cells (*Figure 1-figure*

supplement 4) (Liu *et al.* 2013a). We found that both Wisko (10 µM) (*Figure 1D and E and*

Figure 1-Video 1B) and cNKO (*Figure 1F and G and Figure 1-Video 1C*), but not WKO (*Figure*

1H and I and Figure 1-Video 1D), significantly reduced the percentage of B-cells undergoing

contraction, compared to the vehicle, flox, or WT controls. These results suggest that N-WASP-

but not WASP-activated Arp2/3 mediates branched actin polymerization for B-cell contraction.

Arp2/3, downstream of N-WASP, generates inner F-actin foci, driving B-cell contraction

To understand how Arp2/3 drives B-cells to transition from spreading to contraction, we

identified F-actin structures associated with contracting B-cells that were sensitive to CK-666

treatment. We visualized F-actin by phalloidin staining and compared F-actin organization in the

contact zone of B-cells at the spreading (2 min) and contraction (4 min) phases using TIRF.

While B-cells in both spreading and contraction phases exhibited phalloidin staining outlining the

contact zone (*Fig. 2A, green arrows*), only B-cells in the contracting phase showed interior

phalloidin patches brighter than the phalloidin staining at the periphery of the contact zone

(*Figure 2A, purple arrows*). These F-actin patches were organized into a ring-like structure and

resided ~1 µm behind the spreading front, surrounding an F-actin-poor center (*Figure 2A, purple*

arrows). Here, we refer these F-actin patches as inner F-actin foci. We identified inner F-actin

foci based on whether their peak fluorescence intensity (FI) was ≥2 fold of the mean

fluorescence intensity (MFI) of phalloidin in the no-foci area, had diameters of ≥250 nm, and

were located 1 μm away from the edge of the contact zone. We found that such inner F-actin

foci were detected in >60% of B-cells in the contracting phase (4 min) but only in <20% of B-

cells in the spreading phase (2 min) (*Figure 2B*). CK-666 treatment at 0 min, which inhibited B-

cell contraction, significantly reduced the percentage of B-cells showing inner F-actin foci at 4

min but not at 2 min (*Figure 2B*). CK-666 treatment did not affect the phalloidin staining outlining

the contact zone (*Figure 2A*). Similarly, the percentage of cNKO B-cells showing inner F-actin

foci was drastically reduced at 4 min but not at 2 min (*Figure 2C*). These results suggest that the

formation of these inner F-actin foci is associated with B-cell contraction.

We further quantified the number of inner F-actin foci in individual B-cells from mice expressing

the LifeAct-GFP transgene (Riedl *et al.* 2010) (*Figure 2D, E, and G*), which allowed us to

monitor F-actin reorganization using live-cell imaging (*Figure 2-Video 1*), or using phalloidin

staining in flox control and cNKO B-cells (*Figure 2F*). Consistent with the results with phalloidin

staining, CK-666 treatment at time 0 significantly reduced the number of inner F-actin foci in the

contact zone (*Figure 2D and Figure 1-Video 1A and B*). When we followed the same B-cells

before and after CK-666 treatment at 2 min, many of the inner F-actin foci formed before the

treatment (1 min 50 sec) disappeared after the treatment (2 min 20 sec), significantly reducing

the number of inner F-actin foci (*Figure 2E*). Similar to the CK-666 treatment, cNKO cells

significantly reduced the number of inner F-actin foci in the contact zone, compared to flox

controls (*Figure 2F*). In contrast, WKO, which does not affect B-cell contraction, did not

significantly change the number of inner F-actin foci, compared to WT control B-cells (*Figure 2G*

and Figure 2-Video 1C and D). Thus, N-WASP- but not WASP-activated Arp2/3 drives the

formation and the maintenance of contraction-associated inner F-actin foci.

Inner F-actin foci are derived from lamellipodial actin networks supporting the spreading membrane

We next examined the formation of contraction-associated inner F-actin foci utilizing live-cell TIRF imaging of B-cells from mice expressing LifeAct-GFP. We generated kymographs along eight lines from the center of each contact zone using time-lapse images of LifeAct-GFP (*Figure 3A and B and Figure 3-figure supplement 1*). Analysis of these kymographs showed that most F-actin foci were first detected closely behind lamellipodial F-actin networks. Following maximal spreading, these F-actin foci moved centripetally while increasing in intensity, becoming inner F-actin foci, in cells transitioning from spreading to contraction (*Figure 3B, top panel, and Figure 3-figure supplement 1*). However, such centripetally moving F-actin foci were not detected in B-cells that did not undergo contraction (*Figure 3B, bottom panel*). We quantified the percentage of the eight kymographs from each cell that exhibited lamellipodia-derived inner F-actin foci and found, on average, that six out of eight kymographs from contracting cells showed lamellipodia-derived inner F-actin foci, compared to only one or two kymographs from non-contracting cells (*Figure 3C*). To examine the temporal relationship between the generation of lamellipodia-derived inner F-actin foci and contraction, we plotted the percentage of kymographs with lamellipodia-derived inner F-actin foci over time with the spreading to contracting transition time set as 0 (*Figure 3D*). We found that the percentage of kymographs showing lamellipodia-derived inner F-actin foci peaked at almost the same time when the spreading transitioned to contraction (*Figure 3D*), suggesting a close temporal relationship between the two events. Furthermore, the N-WASP inhibitor Wisko, but not WKO, significantly inhibited the formation of lamellipodia-derived inner F-actin foci (*Figure 3E*). Thus, inner F-actin foci originate from branched actin driven lamellipodia and form simultaneously with the transition of B-cell spreading to contraction.

N-WASP-activated Arp2/3 generates inner F-actin foci by sustaining the lifetime and the centripetal movement of lamellipodial F-actin

We examined the mechanism by which N-WASP and Arp2/3 generate inner F-actin foci by measuring their relative lifetime and mobility using kymographs generated from TIRF time-lapse images of B-cells expressing LifeAct-GFP. Inner F-actin foci were identified as described above, and their tracks were manually determined (*Figure 4A, black dashed lines*). The time window in which an inner F-actin patch could be detected in a kymograph was measured as the relative lifetime, as actin foci could move away from the kymograph line or the TIRF evanescent field (*Figure 4A, right panels*). The distance each F-actin focus moved during its lifetime was used to calculate its speed (*Figure 4A, right panels*). Compared to CK-689-treated B-cells, CK-666 treatment at time 0 significantly reduced the relative lifetime and the centripetal speed of inner F-actin foci (*Figure 4B-D*). After B-cells were treated with CK-666 at 2 min (the average time for B-cells to reach the maximal spreading), the relative lifetime and the centripetal speed of inner F-actin foci were significantly lower than before the treatment in the same cell (*Figure 4E-G, line-linked dots*). Similarly, the N-WASP inhibitor Wisko significantly reduced the relative lifetime and the centripetal speed of inner F-actin foci (*Figure 4H-J*); however, WKO had no significant effect (*Figure 4K-M*). These results show that N-WASP and Arp2/3 mediated branched actin polymerization prolongs the lifetime of lamellipodia-derived F-actin foci and drives them to move inward in the B-cell contact zone.

N-WASP coordinates with NMII to generate inner actin foci and NMII ring-like structures

As non-muscle myosin II (NMII) is required for B-cell contraction (Seeley-Fallen *et al.* 2022), we examined the relationship between the formation of inner F-actin foci and the recruitment and reorganization of NMII in the contact zone using TIRF imaging of B-cells from mice expressing a GFP-NMIIA transgene. Upon interacting with Fab'-PLB, the GFP-NMIIA MFI in the contact zone of untreated WT B-cells increased rapidly in the first minute and slowly afterward (*Figure 5A and*

B and Figure 5-Video 1). Wiskostatin treatment significantly reduced the GFP-NMIIA MFI in the

contact zone (*Figure 5B*) and its initial rate of increase (as determined by the slope of the GFP-

NMIIA MFI increase at 0~30 sec time window) (*Figure 5C*). Kymographs generated from time-

lapse TIRF images of B-cells from mice expressing GFP-NMIIA and LifeAct-RFP showed that

recruited NMIIA accumulated between lamellipodia and inner F-actin foci when the foci moved

centripetally away from lamellipodia (*Figure 5D, white arrow, and Figure 5-Video 1*). Recruited

NMIIA reorganized with inner F-actin foci to form a ring-like structure in the contact zone (*Figure*

5A, D, and E, and Figure 5-Video 1). The percentage of B-cells with NMIIA ring-like structures,

visualized by immunostaining, increased over time as more B-cells underwent contraction

(*Figure 5E and F*). Compared to flox controls, the percentage of cNKO B-cells with NMIIA ring-

like structures was significantly decreased (*Figure 5E and F*). Wiskostatin treatment also

reduced NMIIA recruitment and ring-like structure formation (*Figure 5-Video 1*). Surprisingly, the

percentage of WKO B-cells with NMIIA ring-like structure was higher than that of flox control B-

cells (*Figure 5E and F*). Thus, N-WASP and Arp2/3 mediated branched actin polymerization

promotes the recruitment and the reorganization of NMII ring-like structures.

We next tested whether recruited NMII contributed to the formation of inner actin foci by

inhibiting its motor activity using blebbistatin, which is known to impede B-cell contraction

(Seeley-Fallen *et al.* 2022). Inner F-actin foci in the B-cell contact zone were identified by

phalloidin staining, as described in Figure 2. Treatment with blebbistatin significantly reduced

the percentages of WT splenic B cells exhibiting inner F-actin foci ring-like structures (*Figure 5G*

and H). This result suggests that while NMII is recruited with the help of N-WASP and Arp2/3

mediated actin polymerization, its motor activity is critical for the maturation of F-actin structures

associated with B-cell contraction.

B-cell contraction increases the BCR molecular density in individual clusters

To understand how B-cell contraction promotes BCR signaling attenuation, we examined the

impact of B-cell contraction on the properties of BCR clusters. We first measured the MFI of

AF546-Fab' attached to PLB gathered by B-cells into the contact zone as an indication of the

overall BCR molecular density. The clustering of AF546-Fab' on PLB by B-cell binding reflects

surface BCR clustering, as B-cell binding to transferrin (Tf)-tethered PLB does not cause

surface BCRs to cluster and be phosphorylated (*Figure 6-figure supplement 1*) (Liu *et al.* 2011;

Liu *et al.* 2013a). The MFI of AF546-Fab' in the B-cell contact zone increased over time.

Treatment with CK-666, Wisko, or cNKO all reduced the AF546-Fab' MFI, particularly during the

time window of B-cell contraction in controls (*Figure 6A-G, purple rectangles, and Figure 6-*

Video 1). Notably, the rates of increase in AF546-Fab' MFI, calculated from the slopes of

AF546-Fab' MFI versus time plots in individual cells, were significantly higher during B-cell

contraction than before B-cell contraction in control cells and conditions (*Figure 6D-K*).

Significantly, inhibiting B-cell contraction by CK-666 treatment at 0 (*Figure 6A, D, and H, and*

Figure 6-Video 1A and D) or 2 min (*Figure 6A, E, and I*), Wisko (*Figure 6B, F, and J, and Figure*

6-Video 1B and E), and cNKO (*Figure 6C, G, and K, and Figure 6-Video 1C and F*) reduced the

increases in AF546-Fab' accumulation rates. We further examined the peak FI of AF546-Fab' in

individual microclusters as a measure of the BCR molecular density in individual clusters.

AF546-Fab' clusters were identified based on their diameters ≥250 nm, peak FI ≥1.1 fold

outside the B-cell contact zone, and trackable for ≥20 sec (*Figure 6L and Figure 6-figure*

supplement 2). AF546-Fab' microclusters could not be identified during the early stage of B-cell

spreading. Time-lapse imaging by TIRF enabled us to measure the rate of increase in AF546-

Fab' peak FI in individual clusters (*Figure 6M and Figure 6-figure supplement 2*). Consistent

with our observation of AF546-Fab' MFI increase in the contact zone, the peak FI of individual

clusters increased at a faster rate during contraction than after contraction (when the contact

area no longer decreased) (*Figure 6N*). Furthermore, CK-666 (*Figure 6O*), Wisko (*Figure 6P*),

and cNKO (*Figure 6Q*) all significantly reduced the rate of increase in AF546-Fab' peak FI in

individual clusters. These results show that B-cell contraction significantly increases the

molecular density of BCRs in BCR clusters.

Increased BCR molecular density by B-cell contraction reduces BCR phosphorylation

levels in individual microclusters

To examine how BCR molecular density influenced BCR signaling capability, we

immunostained B-cells interacting with AF546-Fab'-PLB for 1, 3, 5, and 7 min, for

phosphorylated CD79a (pCD79a, Y182) and performed IRM and TIRF imaging (*Figure 7A and*

H). We analyzed equal numbers of AF546-Fab' clusters selected randomly, in the contact zone

of B-cells interacting with Fab'-PLB for 1, 3, 5, and 7 min, using a gradient threshold of their

AF546-Fab' MFI, 1.1 to 4.1 fold of the background, and with diameter ≥250 nm (*Figure 6-figure*

supplement 2). We determined the MFI of pCD79a and AF546-Fab' in individual clusters and

plotted the MFI ratio of pCD79a relative to AF546-Fab' (reflecting the relative level of BCR

phosphorylation) versus the AF546-Fab' peak FI (reflecting BCR molecular density) in individual

clusters (*Figure 7B and C*). The dot plots show that individual AF546-Fab' clusters with relatively

low peak FI displayed increasing MFI ratios of pCD79a to AF546-Fab' as the AF546-Fab' peak

FI rose in flox control B-cells (*Figure 7B*). Past a certain level of AF546-Fab' peak FI, the MFI

ratios of pCD79a to AF546-Fab' decreased as the AF546-Fab' peak FI in individual clusters

further increased (*Figure 7A and B*). Inhibition of B-cell contraction by cNKO reduced the

AF546-Fab' peak FI in individual clusters, maintaining it within a relatively low range, where the

MFI ratios of pCD79a to AF546-Fab' increased with the AF546-Fab' peak FI (*Figure 7C and D*).

Additionally, the average pCD79a to Fab' MFI ratios were much higher in cNKO B-cells than flox

control B-cells, when comparing AF546-Fab' clusters with the same range of peak FI (*Figure*

7B-D). During the contraction stage (5 and 7 min), flox control B-cells exhibited increased

AF546-Fab' MFI (*Figure 7E*) but decreased pCD79a MFI (*Figure 7F*) and the MFI ratios of

pCD79a relative to AF54-Fab' (*Figure 7G*) in individual clusters, compared to the spreading

stage (1 and 3 min). In contrast, non-contracting cNKO B-cells only slightly increased AF546-

Fab' MFI (*Figure 7E*) but significantly increased pCD79a MFI (*Figure 7F*) and the pCD79a to

AF546-Fab' MFI ratios (*Figure 7G*) in individual clusters at the contraction stage compared to

the spreading stage. Consequently, individual clusters in non-contracting cNKO B-cells had

significantly lower AF546-Fab' MFI but significantly higher pCD79a MFI and pCD79a to AF546-

Fab' FIRs than contracting flox control B-cells at 5 and 7 min but not at 1 and 3 min (*Figure 7E-*

G). Similarly, inhibiting B-cell contraction by treatment with the Arp2/3 inhibitor CK-666 at 2 min

post-stimulation reduced AF546-Fab' MFI (*Figure 7I*) but increased pCD79a MFI (*Figure 7J*)

and the MFI ratios of pCD79a relative to AF546-Fab' (*Figure 7K*) in individual clusters. These

data suggest that increases in BCR molecular density in BCR clusters during B-cell contraction

inhibit BCR phosphorylation.

Increased BCR molecular density by B-cell contraction promotes the disassociation of

the stimulatory kinase Syk from BCR microclusters

Increased BCR molecular density may promote signaling attenuation by inducing the

disassociation and/or dephosphorylation of stimulatory kinases from and at BCR clusters. To

test this hypothesis, we analyzed the relative amounts of Syk, a major stimulatory kinase in the

BCR signaling pathway, and its phosphorylated form pSyk (Y519/520) in individual BCR

clusters in relation to the molecular density of BCRs. Splenic B-cells were incubated with Fab'-

PLB for 3 and 7 min (when most cells were at the spreading and contraction phase,

respectively), fixed, permeabilized, stained for Syk (*Figure 8A*) or pSyk (*Figure 8E*), and imaged

using TIRF. We measured MFI ratios of Syk relative to AF546-Fab' in individual BCR clusters to

reflect the relative amount of Syk associated with individual BCR clusters and analyzed their

relationship with AF546-Fab' peak FI (reflecting the molecular density within BCR clusters).

AF546-Fab' clusters were detected and analyzed as described above. In flox control B-cells, the

highest fractions of BCR clusters had an AF546-Fab' peak FI at the150~200 range (*Figure 8A,*

*top panels, and Figure 8B, brown line and symbol*s). The average MFI ratios of Syk to Fab' in

individual clusters increased at a low peak FI range (<140) and did not significantly decrease

until Fab' peak FI reached a relatively high range (>280) (*Figure 8B, black line and symbol*). B-

cells from cNKO mice exhibited a reduction in the Fab' peak FI of clusters (*Figure 8A, bottom*

panels, and Figure 8C and D, brown line and symbol) but an increase in the average Syk to

Fab' MFI ratios in clusters in a wide range of Fab' peak FI, when compared to flox control B-

cells (*Figure 8C and D, black lines and symbols*). However, the average Syk to Fab' MFI ratios

of cNKO B-cells decreased in clusters with Fab' peak FI $\geq$220 and reduced to levels similar to

those in flox control B-cells in clusters with Fab' peak FI $\geq$240 (*Figure 8C and D, black lines and*

symbols). These data suggest that increases in the molecular density of BCR clusters, reflected

by Fab' peak FI, induce disassociation of Syk from BCR clusters in both flox control and cNKO

B-cells. Our results also show that BCR clusters in cNKO B-cells have significantly higher levels

of Syk association than in flox control B-cells, even though they have similar Fab' peak

intensities.

We next analyzed the relationship of the pSyk level in individual clusters with BCR molecular

density using the method described above. We found that the MFI ratios of pSyk relative to

AF546-Fab' gradually decreased with increases in Fab' peak FI in both flox control and cNKO

B-cells (*Figure 8E-G, black lines and symbols*), even though we did not observe an increase in

pSyk to Fab' MFI ratio at the low Fab' peak FI range. Similar to the Syk to Fab' MFI ratio, the

average pSyk to Fab' MFI ratios of individual clusters in cNKO B-cells were much higher than

those in flox control B-cells, except for those at the high Fab' peak FI range (*Figure 8E-H*). To

confirm this result, we analyzed equal numbers of pSyk clusters in the same cells, based on the

criteria of $\geq$1.3 fold increase in the peak FI compared to the background outside the contact

zone with a diameter of $\geq$250 nm. Similar to clusters identified by AF546-Fab', the average MFI

ratios of pSyk to Fab' in these pSyk clusters decreased with increase in their Fab' peak FI in

both flox control and cNKO B-cells (*Figure 8I-K, black lines and symbols*). Again, the average

pSyk to Fab' MFI ratios of pSyk clusters were much higher in cNKO than flox control B-cells, but

were reduced to similar levels in clusters with relatively high Fab' peak FI (*Figure 8K, black line*

and symbols). Similar reductions of total Syk and pSyk with increasing molecular density of

BCR clusters suggest that the disassociation of Syk from BCR clusters, caused by cell

contraction-induced increases in molecular density, contributes to BCR signaling attenuation.

Increased BCR molecular density by B-cell contraction promotes disassociation of the

inhibitory phosphatase SHIP-1 from BCR microclusters

The inhibitory phosphatase SHIP-1 is essential for B-cell signaling attenuation (Brauweiler *et al.*

2000; Liu *et al.* 2011), suggesting that increases in the molecular density of BCR clusters by B-

cell contraction may promote SHIP-1 recruitment. We used the methods described above to

address this hypothesis, staining cells for total SHIP-1 (*Figure 9A*) and phosphorylated SHIP-1

(pSHIP-1 Y1020) (*Figure 9E*). We found that the average SHIP-1 to Fab' MFI ratios in both flox

control and cNKO B-cells decreased with Fab' peak FI at similar rates (*Figure 9A-D, black lines*

and symbols), even though inhibition of contraction by cNKO reduced the Fab' peak FI of BCR

clusters (*Figure 9A-D, brown lines and symbols*). Notably, the reduction in the SHIP-1 to Fab'

MFI ratios with increasing Fab' peak FI occurred at the lowest detectable Fab' peak FI (*Figure*

9B-D, brown lines and symbols), when the Syk to Fab' MFI ratios increased and were sustained

(*Figure 8B-D*). Furthermore, this reciprocal relationship between the SHIP-1 to Fab' MFI ratio

and Fab' peak FI continued over the entire Fab' peak FI range. It also remained the same in

both flox control and cNKO B-cells (*Figure 9B-D*). These results suggest that SHIP-1

disassociates from BCR clusters as their molecular density increases, and that the SHIP-1

disassociation is more sensitive to the molecular density of BCR clusters than Syk

disassociation.

Similar to the relationship of total SHIP-1 with BCR molecular density, the average MFI ratios of

pSHIP-1 relative to AF546-Fab' in individual Fab' clusters decreased with increases in Fab'

peak FI at similar rates in flox control and cNKO B-cells (*Figure 9E-H, black lines and symbols*),

even though inhibition of contraction by cNKO reduced Fab' peak FI of BCR clusters (*Figure 9E-*

H, brown lines and symbols). The average pSHIP-1 to Fab' MFI ratios in individual pSHIP-1

clusters, detected and analyzed in the same way as pSyk clusters, showed the same decrease

with increases in their Fab' peak FI in both flox control and cNKO B-cells (*Figure 9I-K, black*

lines and symbols). Notably, the average pSHIP-1 to Fab' MFI ratios in individual Fab' or

pSHIP-1 clusters in flox control and cNKO B-cells were at similar levels at the same Fab' peak

FI ranges (*Figure 9H and K, black lines and symbols*). These results indicate that contraction-

induced molecular density increases within individual BCR clusters do not induce preferential

recruitment of SHIP-1; instead, it promotes the disassociation of SHIP-1 from BCR clusters.

Discussion

When binding membrane-associated antigen, naive follicular B-cells undergo actin-mediated

spreading followed by a contraction, which amplifies BCR signaling and promotes

immunological synapse formation (Fleire *et al.* 2006; Harwood and Batista 2011). We previously

showed that B-cell contraction after spreading on antigen-presenting surfaces promotes BCR

signaling attenuation (Liu *et al.* 2013a; Seeley-Fallen *et al.* 2022). However, how the actin

cytoskeleton reorganizes as B-cells transition from spreading to contraction and how B-cell

contraction downregulates BCR signaling have been elusive. Here, we demonstrate that inner

F-actin foci formed at the contact zone distal to the lamellipodial F-actin network promote B-cells

to switch from spreading to contraction. These inner foci are derived from the lamellipodial F-

actin network that mediates spreading, are generated by N-WASP- but not WASP-activated

Arp2/3-mediated branched actin polymerization, and facilitate NMII recruitment. N-WASP-

activated actin polymerization coordinates with NMII to form actomyosin ring-like structures,

enabling B-cell contraction. B-cell contraction increases BCR molecular density in existing

clusters, which promotes the disassociation of both the stimulatory kinase Syk and the inhibitory

phosphatase SHIP-1, leading to signaling attenuation.

One significant finding of this study is that Arp2/3-mediated polymerization of branched actin is

required for B-cell contraction. Arp2/3-mediated branched actin polymerization is known to drive

B-cell spreading and to create actin centripetal flow at the contact zone between B-cells and

antigen-presenting surface (Bolger-Munro *et al.* 2019). The actin structure that supports B- and

T-cell spreading to form the immunological synapse with antigen-presenting cells is similar to

lamellipodial F-actin networks found in adherent cells (Bunnell *et al.* 2001; Koestler *et al.* 2008;

Bolger-Munro *et al.* 2019). Lamellipodial F-actin networks consist primarily of branched actin

filaments polymerizing against the plasma membrane interspersed with bundled actin filaments

(Krause and Gautreau 2014; Skau and Waterman 2015). Contractile actin structures, such as

stress fibers, are typically generated from bundled actin filaments, as observed in adherent and

migrating cells (Levayer and Lecuit 2012; Tojkander *et al.* 2015; Hammer *et al.* 2019). Here, we

show that inner F-actin foci, generated by Arp2/3-mediated actin polymerization, transition B-

cells from spreading to contraction. Furthermore, these inner F-actin foci are directly derived

from lamellipodial F-actin networks. Based on the observed dynamics, we infer that instead of

polymerizing against the plasma membrane, Arp2/3 appears to nucleate actin polymerization in

the opposite direction, sustaining actin foci and their movement away from the lamellipodia. The

natural retraction of lamellipodia may contribute to B-cell contraction, but the requirement of

inner actin foci argues against it as a primary mechanism for B-cell contraction. This study does

not exclude the involvement of bundled actin filaments. Based on the requirement of formin-

activated bundled actin filaments for lamellipodial F-actin networks and B-cell spreading (Wang

et al. 2022), we can speculate that bundled actin filaments may play a role in the formation and

movement of these inner F-actin foci, as well as in NMII recruitment.

Our work provides new insights into distinct functions of the actin nucleation-promoting factors

WASP and N-WASP in controlling cell morphology and signaling. Immune cells, including B-

cells, express both hematopoietic-specific WASP and the ubiquitous homolog of WASP, N-

WASP. These two share high sequence homology, activation mechanism, and Arp2/3 activation

function (Padrick and Rosen 2010). We previously identified distinct functions of these two

factors unique to B-cells. While both are required for B-cell spreading, N-WASP plays a unique

role in B-cell contraction. WASP and N-WASP double knockout B-cells fail to spread on antigen-

presenting surfaces, but B-cell-specific N-WASP knockout enhances B-cell spreading and

delays B-cell contraction (Liu *et al.* 2013a). WASP and N-WASP appear to have a competitive

relationship in B-cells, suppressing each other's activation (Liu *et al.* 2013a). WASP is activated

first during B-cell spreading and primarily localized at the periphery of the B-cell contact zone.

N-WASP is activated later during B-cell contraction and scattered across the B-cell contact zone

(Liu *et al.* 2013a). Here, we reveal the exact role of N-WASP in B-cell contraction – to generate

inner F-actin foci from lamellipodial F-actin networks by activating Arp2/3-mediated actin

polymerization. N-WASP- but not WASP-activated actin polymerization prolongs the relative

lifetime of F-actin foci and facilitates their inward motion toward the center of the contact zone.

The delayed activation time and the location in the interior of the contact zone likely give N-

WASP a unique opportunity to generate inner F-actin foci. As cNKO only delays and reduces

but does not block B-cell contraction and inner F-actin foci formation, other actin factors may be

involved, such as the WASP-family verprolin-homologous protein (WAVE) (Rotty *et al.* 2013).

Our recently published data show that NMII is required for B-cell contraction (Seeley-Fallen *et*

al. 2022). Activated NMII is recruited to the B-cell contact zone in a SHIP-1-dependent manner.

NMII levels reach a plateau at the beginning of B-cell contraction, and NMII forms a peripheral

ring surrounding the contact zone during B-cell contraction. Here, we further show that in

addition to SHIP-1, N-WASP but not WASP is also involved in NMII recruitment and NMII ring-

like structure formation, probably by initiating the generation of inner F-actin foci. Collectively,

these findings suggest that SHIP-1 coordinates with N-WASP to recruit NMII. Our previous

finding that SHIP-1 promotes B-cell contraction by facilitating N-WASP activation (Liu *et al.*

2011; Liu *et al.* 2013a) supports this notion. However, it is surprising that the recruitment of NMII

activated by BCR signaling to the contact zone is associated with inhibitory signaling molecules.

BCR downstream signaling, including Ca^{2+} fluxes and Rho-family GTPase activation, likely

activates NMII motor activity (Vicente-Manzanares *et al.* 2009). The activation switches NMII

from the incompetent folded conformation to the competent extended conformation, enabling

NMII molecules to bind to F-actin and assemble into contractible bipolar filaments and stacks

(Matsumura 2005). A denser F-actin organization has been shown to increase NMII filament

stacking (Fenix and Burnette 2018). Here, we showed that the inner F-actin foci generated by

N-WASP-activated Arp2/3 are more stable and denser and thus likely promote NMII binding and

stacking more efficiently than F-actin generated by WASP-activated Arp2/3. Indeed, our

kymograph analysis revealed similar time windows and spatial locations for NMII accumulation

and for generating inner F-actin foci from lamellipodial F-actin, supporting our hypothesis. While

NMII recruitment is facilitated by inner F-actin foci, the motor activity of NMII is required for the

formation of the ring-like structures of inner F-actin foci in return, likely by driving their

centripetal movement, thereby providing mechanical feedback for actin reorganization.

However, the structural organization of inner F-actin foci and recruited NMII and their

coordinated dynamics require further analysis with higher-resolution microscopy techniques.

Wang et al. (Wang *et al.* 2022) recently showed that B-cells generate actomyosin arcs when

interacting with membrane-associated antigen in the presence of adhesion molecules

expressed on professional antigen-presenting cells. In contrast to T-cells, B-cells can also

respond to membrane-associated antigen without the help of adhesion molecules, when

engaging antigen on pathogenic cells, like bacteria, parasites, and cancer cells. While the

relationship between actomyosin arcs and inner F-actin foci remains to be explored, the

coordinated formation of inner F-actin foci and NMII ring-like structures revealed in our study

suggests that these structures are likely the precursors of actomyosin arcs. These structures

can potentially be enhanced and matured by adhesion molecule interactions between B-cells

and antigen-presenting membranes.

Our findings suggest an increase in the molecular density within BCR clusters or B-cell

synapses as one of the mechanisms by which B-cell contraction promotes BCR signaling

attenuation. During contraction, the molecular density of individual BCR clusters, measured by

their MFI and peak FI of BCR clustered Fab', increased faster than during the spreading and

post-contraction phases. Surprisingly, increases in molecular density induced disassociation of

both the stimulatory kinase Syk and the inhibitory phosphatase SHIP-1, contradicting the

existing dogma of sequential association of stimulatory and inhibitory signaling molecules with

the BCR (Franks and Cambier 2018). Interestingly, the disassociation of SHIP-1 is much more

sensitive to BCR molecular density than Syk disassociation. Syk does not disassociate until the

molecular density of BCR clusters reaches the top 10% range, while SHIP-1 disassociates from

BCR clusters with increases over the entire detectable range of molecular densities. When B-

cell contraction is inhibited, the molecular density of BCR clusters is reduced to a range with

limited Syk disassociation and normal SHIP-1 disassociation, probably resulting in high Syk to

SHIP-1 molecular ratios in BCR clusters than contracting B-cells, increasing BCR signaling

levels. The consistency of the data from AF546-Fab' and pSyk or pSHIP-1 clusters supports this

notion. However, the resolution of our widefield TIRF imaging limited our ability to examine

nascent BCR clusters that actively recruit signaling molecules. Available antibodies did not allow

us to simultaneously stain Syk and SHIP-1 or their phosphorylated forms and determine their

relative levels in the same cluster. The timing and dynamics of Syk and SHIP-1 recruitment to

individual BCR clusters and the direct relationship between Syk and SHIP-1 association and

disassociation with BCR clusters and their phosphorylation remain to be further investigated.

How the molecular density within BCR clusters promotes disassociation of signaling molecules

is unknown. We speculate that molecular crowding may displace signaling molecules out of

BCR clusters. Supporting this possibility, disassociation of the 145 kDa SHIP-1 is likely to be

more sensitive to changes in the molecular density of BCR clusters, compared to disassociation

of the 72 kDa Syk. However, these hypotheses need to be further examined. An interesting

finding of this study is that the pCD79a and pSyk levels were higher in individual BCR clusters

of cNKO B-cells than those of flox control B-cells, even when BCR clusters had similar

molecular densities. These data suggest additional mechanisms for B-cell contraction to

promote BCR signaling attenuation. In addition to increasing molecular density within BCR

clusters, the contractile forces generated by actomyosin rings on the B-cell membrane and BCR

clusters may cause changes to the conformations of both BCRs and associated signaling

molecules and their lateral interactions in the membrane. These changes may favor the

disassociation rather than the association of signaling molecules with BCR clusters. Recently

solved molecular assembly structures of both human and mouse BCRs (Dong *et al.* 2022; Su *et*

al. 2022) support conformational and lateral molecular interaction changes as possible

mechanisms for BCR signaling regulation. Similar to the differential impact of the molecular

density on the disassociation of Syk and SHIP-1 from BCR clusters, B-cell contraction may

differentially affect the interaction of surface BCRs with cytoplasmic and membrane-anchored

signaling molecules, such as the lipidated and lipid raft-resident Src kinase Lyn that is

responsible for phosphorylating both the immunoreceptor tyrosine-based activation motif (ITAM)

of CD79a and the immunoreceptor tyrosine-based inhibitory motif (ITIM) that SHIP-1 binds to

(Franks and Cambier 2018).

Following antigen-induced BCR signaling at immunological synapses, B-cells internalize BCR-

captured antigen for processing and presentation. Actomyosin is required for BCR-mediated

endocytosis of antigen, particularly surface- and membrane-associated antigen (Natkanski *et al.*

2013; Hoogeboom *et al.* 2018; Maeda *et al.* 2021). NMII-mediated traction forces pull BCR-

bound antigen off presenting surfaces for endocytosis in an affinity-dependent manner. We

have previously shown that when antigens are tightly attached to a surface, high-affinity binding

of the BCR to antigens tears the B-cell membrane in an NMII-dependent manner, which triggers

lysosome exocytosis and lysosomal enzyme-mediated cleavages of antigen from the associated

surface, allowing antigen endocytosis (Maeda *et al.* 2021). While actomyosin plays essential

roles in both B-cell contraction and the subsequent antigen endocytosis, whether the

actomyosin structure responsible for B-cell contraction also mediates BCR endocytosis remains

an interesting and open question. Our early finding that N-WASP is required for BCR

endocytosis (Liu *et al.* 2013a) supports this notion. BCR endocytosis can reduce B-cell surface

signaling by removing the BCR from the cell surface, transitioning the BCR into an intracellular

signaling state (Chaturvedi *et al.* 2008).

The results presented here have revealed novel insights into the mechanisms underlying actin-

facilitated signaling attenuation of the BCR. Taking the previous and current data of our and

other labs together, we propose a new working model for such actin-mediated signaling

downregulation (*Figure 10*). Upon encountering membrane-associated antigen, such as antigen

presented by follicular dendritic cells and on the surfaces of pathogens, mature follicular B-cells

undergo rapid spreading, primarily driven by WASP-mediated branched actin polymerization,

which maximizes B-cell contact with antigen-presenting surfaces and BCR-antigen

engagement, amplifying signaling. Following maximal spreading, N-WASP distal to lamellipodial

networks activates Arp2/3-mediated branched actin polymerization, which initiates the formation

of inner F-actin foci from lamellipodia towards the center of the B-cell contact zone (*Figure 10*).

NMII is then preferentially recruited to these relatively stable inner foci, which promotes the

centripetal movement of inner F-actin foci and the maturation of ring-like actomyosin structures,

enabling B-cell contraction. B-cell contraction pushes the BCR microclusters formed during B-

cell spreading to the center of the contact zone, increasing their molecular density. Increased

molecular density promotes the disassociation of signaling molecules from BCR clusters,

probably due to crowding and conformational changes, leading to signal downregulation (*Figure*

10). Inhibitory signaling molecules likely activate the actin reorganization that drives B-cell

contraction by promoting the activation and recruitment of NMII to the B-cell contact zone (Liu *et*

al. 2013a; Seeley-Fallen *et al.* 2022) by hitherto unknown mechanisms. The actomyosin

structures responsible for B-cell contraction may further drive B-cells to internalize engaged

antigen, further downregulating signaling at the B-cell surface and initiating antigen processing

and presentation. Thus, actin reorganization downstream of inhibitory phosphatases reinforces

signaling attenuation by driving B-cell contraction.

Materials and Methods

Key Reagents Table

Reagent type	Designation	Source	Identifiers	Additional Information
B-cells from C57BL/6 mice	WT	Jackson Laboratories	000664	
B-cells from WASP$^{-/-}$ mice	WKO	Jackson Laboratories	019458	
B-cells from CD19$^{Cre/+}$N-WASP$^{flox/flox}$ mice	cNKO	Lisa Westerberg laboratory		
B-cells from N-WASP$^{flox/flox}$ mice	Flox control	Lisa Westerberg laboratory		

B-cells from LifeAct-GFP mice	LifeAct-GFP	Roberto Weigert laboratory		
B-cells from LifeAct-RFP mice	LifeAct-RFP	Klaus Ley laboratory		
B-cells from GFP-NMIIA mice	GFP-NMIIA	Robert Adelstein laboratory		
Reagent	1,2-dioleoyl-sn-glycero-3-phosphocholine	Avanti Polar Lipids	850375 P	5 mM
Reagent	1,2-dioleoyl-sn-glycero-3-phosphoethanolamine-cap-biotin	Avanti Polar Lipids	870273 C	50 μM
Reagent	Streptavidin	Jackson Immuno Research	016-000-084	1 μg/ml
Antibody	F(ab')$_2$ fragment of goat IgG anti-mouse Ig(G+M) (polyclonal)	Jackson Immuno Research	115-006-068	-
Antibody	Cy3-Fab fragment of goat anti-mouse IgG+M (polyclonal)	Jackson Immuno Research	115-167-020	2.5 μg per 1 x 10^6 cells
Chemical	2-Mercaptoethylamine HCL	Thermo Fisher Scientific	20408	50 mM
Reagent	EZ-Link^{TM} Maleimide-PEG$_2$-biotin	Thermo Fisher Scientific	A39261	20 mM per mM of protein
Commercial Kit	Alexa Fluor 546 antibody labeling kit	Thermo Fisher Scientific	A20183	-
Ligand	Biotinylated holo-transferrin	Jackson Immuno Research	015-060-050	-
Antibody	Rat IgG$_{2b}$ anti-mouse CD90.2 (Thy1.2) (monoclonal)	Biolegend	105351	1 μl per 2x10^6 cells
Inhibitor	CK-689	Millipore Sigma	182517-25MG	50 μM
Inhibitor	CK-666	Millipore Sigma	SML0006-5MG	50 μM
Inhibitor	Wiskostatin	Millipore Sigma	W2270-5MG	10 μM
Inhibitor	Blebbistatin	Cayman Chemicals	13013	50 μM
Antibody	Rabbit IgG anti-mouse Arp2 antibody (polyclonal)	Abcam	ab47654	1:100
Antibody	Rabbit IgG anti-mouse pWASP antibody (polyclonal)	Thermo Fisher Scientific	PA5-105572	1:100
Antibody	Rabbit IgG anti-mouse pN-WASP antibody (polyclonal)	Thermo Fisher Scientific	PA5-105307	1:100
Antibody	AF488-goat IgG anti-rabbit IgG antibody (polyclonal)	Thermo Fisher Scientific	A-11034	1:200
Antibody	AF546-goat IgG anti-rabbit IgG antibody (polyclonal)	Thermo Fisher Scientific	A-11035	1:200
Reagent	Acti-stain-488 Phalloidin	Cytoskeleton	PHDG1-A	200 nM
Reagent	Acti-stain-555 Phalloidin	Cytoskeleton	PHDH1-A	200 nM

	Rabbit IgG anti-mouse NMIIA antibody (polyclonal)	Abcam	ab75590	1:100
Antibody				
Antibody	Rabbit IgG anti-mouse pCD79a (Y182) (monoclonal)	Cell Signaling Technology	14732S	1:100
Antibody	Rabbit IgG anti-mouse pSyk (Y519/520) (monoclonal)	Cell Signaling Technology	2710S	1:100
Antibody	Rabbit IgG anti-mouse pSHIP-1 (Y1020) (polyclonal)	Cell Signaling Technology	3941S	1:100
Antibody	Rabbit IgG anti-mouse Syk (polyclonal)	Thermo Fisher Scientific	PA5-17812	1:100
Antibody	Rabbit IgG anti-mouse SHIP-1 (polyclonal)	Thermo Fisher Scientific	PA5-115894	1:100

Mice and B-cell isolation

Wild-type (WT) C57BL/6 mice and WASP knockout (WKO) mice on the C57BL/6 background

were purchased from Jackson Laboratories. C57BL/6 mice expressing LifeAct-GFP transgene

(Riedl *et al.* 2010) were kindly provided by Dr. Roberto Weigert at National Cancer Institute,

USA. C57BL/6 mice expressing the LifeAct-RFP transgene were kindly provided by Dr. Klaus

Ley at La Jolla Institute for Allergy & Immunology. C57BL/6 mice expressing the GFP-non-

muscle myosin IIA (NMIIA) transgene were kindly provided by Dr. Robert Adelstein at the

National Heart, Lung, and Blood Institute, USA. WKO mice were bred with LifeAct-GFP mice to

obtain LifeAct-GFP-expressing WKO mice. GFP-NMIIA mice were crossed with LifeAct-RFP

mice to generate mice expressing both transgenes. N-WASP$^{flox/flox}$ on a 129Sv background

(Cotta-de-Almeida *et al.* 2007) were kindly provided by Dr. Lisa Westerberg at Karolinska

Institute, Sweden. B-cell-specific N-WASP knockout mice (cNKO, CD19$^{Cre/+}$N-WASP$^{flox/flox}$) and

floxed littermate controls (N-WASP$^{flox/flox}$) were obtained by breeding N-WASP$^{flox/flox}$ mice with

CD19$^{Cre/+}$ mice on a C57BL/6 background.

Primary B-cells were isolated from the spleens of 6 to 18 week-old male or female mice, using a

previously published protocol (Sharma *et al.* 2009). Briefly, mononuclear cells were isolated by

Ficoll density-gradient centrifugation (Millipore Sigma), and T-cells were eliminated by

complement-mediated cytolysis with anti-mouse CD90.2 mAb (BD Biosciences) and guinea pig

complement (Innovative Research Inc). Monocytes and dendritic cells were eliminated by

panning at 37°C and 5% CO_2. Isolated B-cells were kept on ice in DMEM (Lonza) supplemented

with 0.6% BSA (Thermo Fisher Scientific). All work involving mice was approved by the

Institutional Animal Care and Usage Committee of the University of Maryland.

Pseudo-antigen-coated planar lipid bilayers

Planar lipid bilayers (PLB) were prepared using a previously described method (Dustin *et al.*

2007; Liu *et al.* 2012a). Briefly, liposomes were generated from a mixture of 5 mM (total

concentration) 1,2-dioleoyl-sn-glycero-3-phosphocholine and 1,2-dioleoyl-sn-glycero-3-

phosphoethanolamine-cap-biotin (Avanti Polar Lipids) at a 100:1 molar ratio by sonication.

Glass coverslips, cleaned overnight with Piranha solution (KMG chemicals), were attached to 8-

well chambers (Lab-Tek) and incubated with liposomes (0.05 mM) in PBS for 20 min at room

temperature and washed with PBS. The chambers were then incubated with 1 µg/ml

streptavidin (Jackson ImmunoResearch Laboratories) for 10 min, washed with PBS, and then

incubated with 10 µg/ml mono-biotinylated Fab' fragment of goat anti-mouse Ig(G+M) (pseudo-

antigen) (Fab'-PLB) for 10 min, followed by PBS wash. For transferrin-coated PLB (Tf-PLB), 16

16 µg/ml biotinylated holoTF was used. To visualize Fab' clustering, a mixture of 0.5 µg/ml

Alexa Fluor (AF) 546-labeled and 9.5 µg/ml unlabeled mono-biotinylated Fab' fragment of goat

anti-mouse Ig(G+M) was used. The lateral mobility of AF546-Fab' on the PLB was tested using

fluorescence recovery after photobleaching (FRAP) to ensure ≥85% AF546 fluorescence

recovery within 1 min after photobleaching using a Zeiss LSM 710 equipped with a 60X oil-

immersion objective.

Mono-biotinylated Fab' fragments were generated as previously described (Liu *et al.* 2011).

Briefly, the disulfide bond linking the two Fab fragments of F(ab')$_2$ goat anti-mouse Ig(G+M) was

reduced using 2-mercaptoethylamine HCL (Thermo Fisher Scientific) and biotinylated by using

maleimide-PEG$_2$-biotin (Thermo Fisher Scientific). Mono-biotinylated Fab' fragments of goat

anti-mouse Ig(G+M) were labeled with AF546 using an Alexa Fluor 546 antibody labeling kit

(Thermo Fisher Scientific). The molar ratio of AF546 to Fab' in the AF546-labeled Fab' was ~2

determined by a Nanodrop spectrophotometer (Nanodrop Technologies).

Total internal fluorescence microscopy

To visualize molecules proximal to interacting sites between B-cells and Fab'- or Tf-PLB, we

utilized total internal reflection fluorescence microscopy (TIRF) and interference reflection

microscopy (IRM). Images were acquired using a Nikon TIRF system on an inverted microscope

(Nikon TE2000-PFS, Nikon Instruments Inc.) equipped with a 60X, NA 1.49 Apochromat TIRF

objective (Nikon), a Coolsnap HQ2 CCD camera (Roper Scientific), and two solid-state lasers of

wavelength 491 and 561 nm. IRM, AF488, and AF546 images were acquired sequentially.

The plasma membrane area of B-cells contacting PLB was determined using IRM images and

custom MATLAB codes. Whether a B-cell contracted or not was determined using the area

versus time plots, wherein if a B-cell's contact zone area reduced by $\geq$5% for at least 10 sec

after reaching a maximum value, it was classified as contracting. To image intracellular

molecules, B-cells were incubated with Fab'-PLBs for varying lengths of time in PBS at 37°C,

fixed with 4% paraformaldehyde, permeabilized with 0.05% saponin, and stained for various

molecules. For live-cell imaging, B-cells were pre-warmed to 37°C and imaged as soon as B-

cells were dropped into coverslip chambers coated with Fab'-PLB containing PBS in a humidity

chamber at 37°C, at 2 sec per frame, up to 7 min. All images from multiple independent

experiments were analyzed using NIH ImageJ and custom MatLab scripts. Acquired

fluorescence intensity (FI) data were normalized to the one with the lowest FI.

Inhibitors

CK-666 (50 µM, Millipore Sigma) was used to perturb Arp2/3 activity (Nolen *et al.* 2009), and its

non-functional derivative CK-689 (50 µM, Millipore Sigma) as a control. CK-666 or CK-689 was

added at either 0 min, the start of incubation with Fab'-PLB, or at 2 min when most B-cells

reached maximum spreading. Notably, B-cells take approximately 1 to 1.5 min to land on PLB.

The time when B-cells were added to Fab'-PLB is referred to as 0 min, and the time when B-

cells landed on PLB as the start of spreading. The effectiveness of CK-666 was determined by

its inhibitory effects on the recruitment of Arp2/3, stained by an anti-Arp2 antibody (Abcam), to

the B-cell contact zone using TIRF (*Figure 1-figure supplement 2*). Wiskostatin (Wisko, 10 µM,

Millipore Sigma) was used to perturb N-WASP activity in B-cells (Peterson *et al.* 2004), with

DMSO used as a vehicle control. Splenic B-cells were pretreated with Wisko for 10 min at 37°C

before and during incubation with Fab'-PLB. The effectiveness of Wisko was determined by its

inhibitory effects on the level of phosphorylated N-WASP in the B-cell contact zone using

immunostaining and TIRF (*Figure 1-figure supplement 4A and C)*. Possible effects of Wisko on

WASP activation were evaluated by measuring the mean fluorescence intensity (MFI) of

phosphorylated WASP in the contact zone of B-cells treated with or without Wisko for 10 min

(*Figure 1-figure supplement 4B and D*). Blebbistatin (Bleb, 50 µM, Cayman Chemicals) was

used to inhibit the NMII motor activity (Seeley-Fallen *et al.* 2022). Splenic B-cells were

pretreated with Bleb for 20 min before and during incubation with Fab'-PLB.

Analysis of the actin cytoskeleton

F-actin was visualized by phalloidin staining in fixed cells and by LifeAct-GFP or LifeAct-RFP

expressed by primary B-cells from transgenic mice in live cells. Inner F-actin foci, visualized by

both phalloidin staining and LifeAct-GFP, were identified using NIH ImageJ based on the

following three criteria: 1) ≥250 nm in diameter, 2) peak fluorescence intensity (FI) ≥2 fold higher

than the FI of a nearby area containing no foci, and 3) ≥1 µm away from the outer edge of the B-

cell contact zone. The horizontal or vertical length of a focus was measured, and the lesser of

the two values was used as its diameter. To analyze the spatiotemporal relationship between

inner F-actin foci and the lamellipodial F-actin network, we generated 8 radially and equally

spaced kymographs from each cell using time-lapse images and MATLAB. Each kymograph

was either classified as contracting or not contracting based on the movement of the leading

edge of the B-cell contact zone. The percentage of 8 kymographs from each cell exhibiting inner

F-actin foci that could be traced back to lamellipodial F-actin when the B-cell switched from

spreading to contraction was determined.

The relative speeds and lifetimes of F-actin foci were determined using three kymographs from

each B-cell (positioned to track as many inner F-actin foci as possible) and F-actin foci

emerging during a 60-sec window right after B-cell maximal spreading. Only those remaining

visible in individual kymographs for at least 4 sec were analyzed. The time duration of each F-

actin focus detected in a kymograph was used to determine the relative lifetime. The distances

individual F-actin foci moved during their lifetimes were used to determine the relative speed.

The term 'relative' has been used because the disappearance of an F-actin focus could be due

to its movement away from the region used for the kymograph or vertically from the TIRF

evanescent field of excitation.

To determine the rate of myosin recruitment, GFP-NMIIA MFI in the B-cell contact zone was

plotted over time, and the slope for the initial segment of the MFI versus the time plot was

determined using linear regression. The percentage of B-cells with the NMIIA ring was

determined by visual inspection.

Analysis of BCR-Fab' clusters

BCR clusters were identified by clustering of AF546-Fab' on PLBs using custom code by

MATLAB. We utilized the median FI of the Fab'-PLB within the same area outside but near the

B-cell contact zone as the background. The median, but not mean FI value, was used for

background FI calculations to minimize fluctuations due to debris. To ensure that individual

clusters with varying FI were detected as distinct objects, we used 16 graded thresholds from

1.1 to 4.1 fold of the background (0.2 fold apart) to acquire 16 sets of binary masks for each

frame of time-lapse images from each cell. When clusters were detected by multiple thresholds

at the same location, only the one identified by the highest threshold was retained. Objects that

were smaller than 250 nm in diameter or could not be tracked for at least 20 sec were

eliminated. This allowed us to detect the position of BCR-Fab' clusters and track them over time

until they merged with other clusters. The horizontal or vertical length of a focus was measured,

and the lesser of the two values was used as its diameter. We chose the peak FI as a metric for

the extent of Fab' clustering by the BCR, because it does not rely on the area occupied by each

cluster. The rate of peak FI increase was determined by linear regression of peak FI versus time

curves for each cluster for a given window of time.

BCR signaling

Splenic B-cells were incubated with AF546-Fab'-PLB at 37°C and fixed at 1, 3, 5, 7, and 9 min.

After fixation, cells were stained for phosphorylated CD79a (pCD79a, Y182) (Cell Signaling

Technology), Syk (pSyk, Y519/520) (Cell Signaling Technology), SHIP-1 (pSHIP-1, Y1020) (Cell

Signaling Technology), or total Syk or SHIP-1 proteins (Thermo Fisher Scientific). We identified

BCR-Fab' clusters as described. We determined the peak FI of AF546-Fab' and the MFI of

AF546-Fab', pCD79a, pSyk, pSHIP-1, Syk, or SHIP-1 within each cluster. The ratio of pCD79a

MFI to AF546-Fab' MFI in the same cluster was used to estimate the relative phosphorylation

level of BCRs in that cluster. This MFI ratio was plotted against AF546-Fab' peak FI to analyze

the relationship between BCR phosphorylation and BCR density of individual clusters. We

calculated the MFI ratio of Syk to Fab' or SHIP-1 to Fab' in individual clusters to estimate the

relative recruitment level of Syk or SHIP-1 to BCR clusters. We plotted these MFI ratios against

Fab' peak FI to determine the relationship between Syk and SHIP-1 recruitment levels and BCR

density in individual clusters. We calculated the MFI ratio of pSyk to Fab' or pSHIP-1 to Fab' to

estimate the amount of pSyk or pSHIP-1 relative to BCR in individual clusters. We plotted these

MFI ratios against Fab' peak FI to determine the relationship between Syk or SHIP-1

phosphorylation and BCR density in individual clusters. We identified pSyk or pSHIP-1 puncta

using the criteria: $\geq$1.3 fold of the background staining outside the B-cell contact zone and $\geq$250

nm in diameter. We calculated the MFI ratio of pSyk to Fab' or pSHIP-1 to Fab' in individual

puncta to estimate the amount of pSyk or pSHIP-1 relative to BCR in individual puncta. We

plotted these MFI ratios against Fab' peak FI to determine the relationship between Syk or

SHIP-1 phosphorylation and BCR density in individual puncta. We determined the fraction of

clusters with graded increases in Fab' peak FI (every 20 FI units) out of the total to analyze the

distribution of clusters with different Fab' peak FI. We also determine the average MFI ratios of

pCD79a, Syk, pSyk, SHIP-1, and pSHIP-1 relative to Fab' in each graded cluster population.

We further divided AF546-Fab' clusters into three populations based on their peak AF546-Fab'

FI, relatively low (<190), medium (190-280), and high (>280, detected only in contracted cells)

and compared the MFI ratios of pCD79a, pSyk, pSHIP-1, Syk, and SHIP-1 relatively to AF546-

Fab' among the three populations using statistical analysis described below. The data were

generated from 3 independent experiments with 20~23 (60~69 total) cells and 125~140

individual clusters (1500~1680 total) per condition, per time point, and per experiment.

Statistical analysis

Statistical analysis was performed using the Mann-Whitney U non-parametric test for unpaired

groups having different sample sizes or the Student's *t*-test for paired groups with the same

sample size. To compare curves, Kolmogorov-Smirnov test was used. Statistical analyses were

performed in Microsoft Excel, GraphPad Prism, and MATLAB. All data are presented as mean $\pm$

SEM (standard error of the mean). When testing multiple hypotheses, *p*--values acquired using

t-tests were corrected using the Benjamini-Hochberg/Yekutieli method for false discovery rate

control.

MATLAB scripts

All MATLAB scripts used for this study are available as supplemental materials.

Acknowledgements

This work was supported by NIH grants GM064625 (NWA and WS), GM145313 (AU), and

AI122205 (AU and WS). We would like to acknowledge Amy Beaven and Kenneth Class for

technical support from the Confocal Microcopy and Flow Cytometry cores. We would like to

thank Dr. Norma W. Andrews (University of Maryland, College Park) for her careful reading and

critical comments on the manuscript.

References

Adachi, T., Wienands, J., Wakabayashi, C., Yakura, H., Reth, M. and Tsubata, T. (2001) 'SHP-1
requires inhibitory co-receptors to down-modulate B cell antigen receptor-mediated
phosphorylation of cellular substrates', *J Biol Chem*, 276(28), 26648-55, available:
http://dx.doi.org/10.1074/jbc.M100997200.

Aman, M.J., Lamkin, T.D., Okada, H., Kurosaki, T. and Ravichandran, K.S. (1998) 'The inositol
phosphatase SHIP inhibits Akt/PKB activation in B cells', *J Biol Chem*, 273(51), 33922-8,
available: http://dx.doi.org/10.1074/jbc.273.51.33922.

Batista, F.D. and Harwood, N.E. (2009) 'The who, how and where of antigen presentation to B
cells', *Nat Rev Immunol*, 9(1), 15-27, available: http://dx.doi.org/10.1038/nri2454.

Batista, F.D., Treanor, B. and Harwood, N.E. (2010) 'Visualizing a role for the actin cytoskeleton
in the regulation of B-cell activation', *Immunol Rev*, 237(1), 191-204, available:
http://dx.doi.org/10.1111/j.1600-065X.2010.00943.x.

Bolger-Munro, M., Choi, K., Scurll, J.M., Abraham, L., Chappell, R.S., Sheen, D., Dang-Lawson,
M., Wu, X., Priatel, J.J., Coombs, D., Hammer, J.A. and Gold, M.R. (2019) 'Arp2/3
complex-driven spatial patterning of the BCR enhances immune synapse formation,

BCR signaling and B cell activation', *Elife*, 8, available:
http://dx.doi.org/10.7554/eLife.44574.

Bolland, S., Pearse, R.N., Kurosaki, T. and Ravetch, J.V. (1998) 'SHIP modulates immune
receptor responses by regulating membrane association of Btk', *Immunity*, 8(4), 509-16,
available: http://dx.doi.org/10.1016/s1074-7613(00)80555-5.

Brauweiler, A.M., Tamir, I. and Cambier, J.C. (2000) 'Bilevel control of B-cell activation by the
inositol 5-phosphatase SHIP', *Immunol Rev*, 176, 69-74.

Bunnell, S.C., Kapoor, V., Trible, R.P., Zhang, W. and Samelson, L.E. (2001) 'Dynamic actin
polymerization drives T cell receptor-induced spreading: a role for the signal
transduction adaptor LAT', *Immunity*, 14(3), 315-29, available:
http://dx.doi.org/10.1016/s1074-7613(01)00112-1.

Carrasco, Y.R., Fleire, S.J., Cameron, T., Dustin, M.L. and Batista, F.D. (2004) 'LFA-1/ICAM-1
interaction lowers the threshold of B cell activation by facilitating B cell adhesion and
synapse formation', *Immunity*, 20(5), 589-99.

Chaturvedi, A., Dorward, D. and Pierce, S.K. (2008) 'The B cell receptor governs the subcellular
location of Toll-like receptor 9 leading to hyperresponses to DNA-containing antigens',
Immunity, 28(6), 799-809, available: http://dx.doi.org/10.1016/j.immuni.2008.03.019.

Cotta-de-Almeida, V., Westerberg, L., Maillard, M.H., Onaldi, D., Wachtel, H., Meelu, P., Chung,
U.I., Xavier, R., Alt, F.W. and Snapper, S.B. (2007) 'Wiskott Aldrich syndrome protein
(WASP) and N-WASP are critical for T cell development', *Proc Natl Acad Sci U S A*,
104(39), 15424-9, available: http://dx.doi.org/10.1073/pnas.0706881104.

Cyster, J.G. (2010) 'B cell follicles and antigen encounters of the third kind', *Nat Immunol*,
11(11), 989-96, available: http://dx.doi.org/10.1038/ni.1946.

Dal Porto, J.M., Gauld, S.B., Merrell, K.T., Mills, D., Pugh-Bernard, A.E. and Cambier, J. (2004)
'B cell antigen receptor signaling 101', *Mol Immunol*, 41(6-7), 599-613, available:
http://dx.doi.org/10.1016/j.molimm.2004.04.008.

Depoil, D., Weber, M., Treanor, B., Fleire, S.J., Carrasco, Y.R., Harwood, N.E. and Batista, F.D.
(2009) 'Early events of B cell activation by antigen', *Sci Signal*, 2(63), pt1, available:
http://dx.doi.org/10.1126/scisignal.263pt1.

Dong, Y., Pi, X., Bartels-Burgahn, F., Saltukoglu, D., Liang, Z., Yang, J., Alt, F.W., Reth, M. and
Wu, H. (2022) 'Structural principles of B cell antigen receptor assembly', *Nature*,
612(7938), 156-161, available: http://dx.doi.org/10.1038/s41586-022-05412-7.

Dustin, M.L., Starr, T., Varma, R. and Thomas, V.K. (2007) 'Supported planar bilayers for study
of the immunological synapse', *Curr Protoc Immunol*, Chapter 18, Unit 18.13, available:
http://dx.doi.org/10.1002/0471142735.im1813s76.

Fenix, A.M. and Burnette, D.T. (2018) 'Assembly of myosin II filament arrays: Network
Contraction versus Expansion', *Cytoskeleton (Hoboken)*, 75(12), 545-549, available:
http://dx.doi.org/10.1002/cm.21487.

Fleire, S.J., Goldman, J.P., Carrasco, Y.R., Weber, M., Bray, D. and Batista, F.D. (2006) 'B cell
ligand discrimination through a spreading and contraction response', *Science*,
312(5774), 738-41, available: http://dx.doi.org/10.1126/science.1123940.

Franks, S.E. and Cambier, J.C. (2018) 'Putting on the Brakes: Regulatory Kinases and
Phosphatases Maintaining B Cell Anergy', *Front Immunol*, 9, 665, available:
http://dx.doi.org/10.3389/fimmu.2018.00665.

Freeman, S.A., Lei, V., Dang-Lawson, M., Mizuno, K., Roskelley, C.D. and Gold, M.R. (2011)
'Cofilin-mediated F-actin severing is regulated by the Rap GTPase and controls the
cytoskeletal dynamics that drive lymphocyte spreading and BCR microcluster formation',
J Immunol, 187(11), 5887-900, available: http://dx.doi.org/10.4049/jimmunol.1102233.

Gitlin, A.D., Shulman, Z. and Nussenzweig, M.C. (2014) 'Clonal selection in the germinal centre
by regulated proliferation and hypermutation', *Nature*, 509(7502), 637-40, available:
http://dx.doi.org/10.1038/nature13300.

Gonzalez, S.F., Pitcher, L.A., Mempel, T., Schuerpf, F. and Carroll, M.C. (2009) 'B cell
acquisition of antigen in vivo', *Curr Opin Immunol*, 21(3), 251-7, available:
http://dx.doi.org/10.1016/j.coi.2009.05.013.

Gross, A.J., Lyandres, J.R., Panigrahi, A.K., Prak, E.T. and DeFranco, A.L. (2009)
'Developmental acquisition of the Lyn-CD22-SHP-1 inhibitory pathway promotes B cell
tolerance', *J Immunol*, 182(9), 5382-92, available:
http://dx.doi.org/10.4049/jimmunol.0803941.

Hammer, J.A., Wang, J.C., Saeed, M. and Pedrosa, A.T. (2019) 'Origin, Organization,
Dynamics, and Function of Actin and Actomyosin Networks at the T Cell Immunological
Synapse', *Annu Rev Immunol*, 37, 201-224, available: http://dx.doi.org/10.1146/annurev-
immunol-042718-041341.

Harwood, N.E. and Batista, F.D. (2010) 'Early events in B cell activation', *Annu Rev Immunol*,
28, 185-210, available: http://dx.doi.org/10.1146/annurev-immunol-030409-101216.

Harwood, N.E. and Batista, F.D. (2011) 'The cytoskeleton coordinates the early events of B-cell activation', *Cold Spring Harb Perspect Biol*, 3(2), available: http://dx.doi.org/10.1101/cshperspect.a002360.

Hoogeboom, R., Natkanski, E.M., Nowosad, C.R., Malinova, D., Menon, R.P., Casal, A. and Tolar, P. (2018) 'Myosin IIa Promotes Antibody Responses by Regulating B Cell Activation, Acquisition of Antigen, and Proliferation', *Cell Rep*, 23(8), 2342-2353, available: http://dx.doi.org/10.1016/j.celrep.2018.04.087.

Hoogeboom, R. and Tolar, P. (2016) 'Molecular Mechanisms of B Cell Antigen Gathering and Endocytosis', *Curr Top Microbiol Immunol*, 393, 45-63, available: http://dx.doi.org/10.1007/82_2015_476.

Kinnon, C., Hinshelwood, S., Levinsky, R.J. and Lovering, R.C. (1993) 'X-linked agammaglobulinemia--gene cloning and future prospects', *Immunol Today*, 14(11), 554-8, available: http://dx.doi.org/10.1016/0167-5699(93)90187-P.

Kläsener, K., Maity, P.C., Hobeika, E., Yang, J. and Reth, M. (2014) 'B cell activation involves nanoscale receptor reorganizations and inside-out signaling by Syk', *Elife*, 3, e02069.

Koestler, S.A., Auinger, S., Vinzenz, M., Rottner, K. and Small, J.V. (2008) 'Differentially oriented populations of actin filaments generated in lamellipodia collaborate in pushing and pausing at the cell front', *Nat Cell Biol*, 10(3), 306-13, available: http://dx.doi.org/10.1038/ncb1692.

Krause, M. and Gautreau, A. (2014) 'Steering cell migration: lamellipodium dynamics and the regulation of directional persistence', *Nat Rev Mol Cell Biol*, 15(9), 577-90, available: http://dx.doi.org/10.1038/nrm3861.

Kurosaki, T. (2000) 'Functional dissection of BCR signaling pathways', *Curr Opin Immunol*, 12(3), 276-81.

Kurosaki, T., Shinohara, H. and Baba, Y. (2010) 'B cell signaling and fate decision', *Annu Rev Immunol*, 28, 21-55, available: http://dx.doi.org/10.1146/annurev.immunol.021908.132541.

Kwak, K., Akkaya, M. and Pierce, S.K. (2019) 'B cell signaling in context', *Nat Immunol*, 20(8), 963-969, available: http://dx.doi.org/10.1038/s41590-019-0427-9.

Leung, W.H., Tarasenko, T., Biesova, Z., Kole, H., Walsh, E.R. and Bolland, S. (2013) 'Aberrant antibody affinity selection in SHIP-deficient B cells', *Eur J Immunol*, 43(2), 371-81, available: http://dx.doi.org/10.1002/eji.201242809.

987   Levayer, R. and Lecuit, T. (2012) 'Biomechanical regulation of contractility: spatial control and
988   dynamics', *Trends Cell Biol*, 22(2), 61-81, available:
989   http://dx.doi.org/10.1016/j.tcb.2011.10.001.

990
991   Liu, C., Bai, X., Wu, J., Sharma, S., Upadhyaya, A., Dahlberg, C.I., Westerberg, L.S., Snapper,
992   S.B., Zhao, X. and Song, W. (2013a) 'N-WASP is essential for the negative regulation of
993   B cell receptor signaling', *PLoS Biol*, 11(11), e1001704, available:
994   http://dx.doi.org/10.1371/journal.pbio.1001704.

995
996   Liu, C., Fallen, M.K., Miller, H., Upadhyaya, A. and Song, W. (2013b) 'The actin cytoskeleton
997   coordinates the signal transduction and antigen processing functions of the B cell
998   antigen receptor', *Front Biol (Beijing)*, 8(5), 475-485, available:
999   http://dx.doi.org/10.1007/s11515-013-1272-0.

1000
1001   Liu, C., Miller, H., Hui, K.L., Grooman, B., Bolland, S., Upadhyaya, A. and Song, W. (2011) 'A
1002   balance of Bruton's tyrosine kinase and SHIP activation regulates B cell receptor cluster
1003   formation by controlling actin remodeling', *J Immunol*, 187(1), 230-9, available:
1004   http://dx.doi.org/10.4049/jimmunol.1100157.

1005
1006   Liu, C., Miller, H., Orlowski, G., Hang, H., Upadhyaya, A. and Song, W. (2012a) 'Actin
1007   reorganization is required for the formation of polarized B cell receptor signalosomes in
1008   response to both soluble and membrane-associated antigens', *J Immunol*, 188(7), 3237-
1009   46, available: http://dx.doi.org/10.4049/jimmunol.1103065.

1010
1011   Liu, C., Miller, H., Sharma, S., Beaven, A., Upadhyaya, A. and Song, W. (2012b) 'Analyzing
1012   actin dynamics during the activation of the B cell receptor in live B cells', *Biochem
1013   Biophys Res Commun*, 427(1), 202-6, available:
1014   http://dx.doi.org/10.1016/j.bbrc.2012.09.046.

1015
1016   Maeda, F.Y., van Haaren, J.J., Langley, D.B., Christ, D., Andrews, N.W. and Song, W. (2021)
1017   'Surface-associated antigen induces permeabilization of primary mouse B-cells and
1018   lysosome exocytosis facilitating antigen uptake and presentation to T-cells', *Elife*, 10,
1019   available: http://dx.doi.org/10.7554/eLife.66984.

1020
1021   Matsumura, F. (2005) 'Regulation of myosin II during cytokinesis in higher eukaryotes', *Trends
1022   Cell Biol*, 15(7), 371-7, available: http://dx.doi.org/10.1016/j.tcb.2005.05.004.

1023
1024   Mizuno, K., Tagawa, Y., Mitomo, K., Arimura, Y., Hatano, N., Katagiri, T., Ogimoto, M. and
1025   Yakura, H. (2000) 'Src homology region 2 (SH2) domain-containing phosphatase-1
1026   dephosphorylates B cell linker protein/SH2 domain leukocyte protein of 65 kDa and
1027   selectively regulates c-Jun NH2-terminal kinase activation in B cells', *J Immunol*, 165(3),
1028   1344-51, available: http://dx.doi.org/10.4049/jimmunol.165.3.1344.

1029

1030 Natkanski, E., Lee, W.Y., Mistry, B., Casal, A., Molloy, J.E. and Tolar, P. (2013) 'B cells use
1031   mechanical energy to discriminate antigen affinities', *Science*, 340(6140), 1587-90,
1032   available: http://dx.doi.org/10.1126/science.1237572.

1033
1034 Nolen, B.J., Tomasevic, N., Russell, A., Pierce, D.W., Jia, Z., McCormick, C.D., Hartman, J.,
1035   Sakowicz, R. and Pollard, T.D. (2009) 'Characterization of two classes of small molecule
1036   inhibitors of Arp2/3 complex', *Nature*, 460(7258), 1031-4, available:
1037   http://dx.doi.org/10.1038/nature08231.

1038
1039 Padrick, S.B. and Rosen, M.K. (2010) 'Physical mechanisms of signal integration by WASP
1040   family proteins', *Annu Rev Biochem*, 79, 707-35, available:
1041   http://dx.doi.org/10.1146/annurev.biochem.77.060407.135452.

1042
1043 Pao, L.I., Lam, K.P., Henderson, J.M., Kutok, J.L., Alimzhanov, M., Nitschke, L., Thomas, M.L.,
1044   Neel, B.G. and Rajewsky, K. (2007) 'B cell-specific deletion of protein-tyrosine
1045   phosphatase Shp1 promotes B-1a cell development and causes systemic autoimmunity',
1046   *Immunity*, 27(1), 35-48, available: http://dx.doi.org/10.1016/j.immuni.2007.04.016.

1047
1048 Peterson, J.R., Bickford, L.C., Morgan, D., Kim, A.S., Ouerfelli, O., Kirschner, M.W. and Rosen,
1049   M.K. (2004) 'Chemical inhibition of N-WASP by stabilization of a native autoinhibited
1050   conformation', *Nat Struct Mol Biol*, 11(8), 747-55, available:
1051   http://dx.doi.org/10.1038/nsmb796.

1052
1053 Pierce, S.K. (2002) 'Lipid rafts and B-cell activation', *Nat Rev Immunol*, 2(2), 96-105, available:
1054   http://dx.doi.org/10.1038/nri726.

1055
1056 Reth, M. (1994) 'B cell antigen receptors', *Curr Opin Immunol*, 6(1), 3-8.

1057
1058 Reth, M. and Wienands, J. (1997) 'Initiation and processing of signals from the B cell antigen
1059   receptor', *Annu Rev Immunol*, 15, 453-79, available:
1060   http://dx.doi.org/10.1146/annurev.immunol.15.1.453.

1061
1062 Rey-Suarez, I., Wheatley, B.A., Koo, P., Bhanja, A., Shu, Z., Mochrie, S., Song, W., Shroff, H.
1063   and Upadhyaya, A. (2020) 'WASP family proteins regulate the mobility of the B cell
1064   receptor during signaling activation', *Nat Commun*, 11(1), 439, available:
1065   http://dx.doi.org/10.1038/s41467-020-14335-8.

1066
1067 Riedl, J., Flynn, K.C., Raducanu, A., Gärtner, F., Beck, G., Bösl, M., Bradke, F., Massberg, S.,
1068   Aszodi, A., Sixt, M. and Wedlich-Söldner, R. (2010) 'Lifeact mice for studying F-actin
1069   dynamics', *Nat Methods*, 7(3), 168-9, available: http://dx.doi.org/10.1038/nmeth0310-
1070   168.

1071

1072 Rotty, J.D., Wu, C. and Bear, J.E. (2013) 'New insights into the regulation and cellular functions
1073   of the ARP2/3 complex', *Nat Rev Mol Cell Biol*, 14(1), 7-12, available:
1074   http://dx.doi.org/10.1038/nrm3492.

1075
1076 Schreiner, G.F. and Unanue, E.R. (1977) 'Capping and the lymphocyte: models for membrane
1077   reorganization', *J Immunol*, 119(5), 1549-51.

1078
1079 Seeley-Fallen, M.K., Lazzaro, M., Liu, C., Li, Q.Z., Upadhyaya, A. and Song, W. (2022) 'Non-
1080   Muscle Myosin II Is Essential for the Negative Regulation of B-Cell Receptor Signaling
1081   and B-Cell Activation', *Front Immunol*, 13, 842605, available:
1082   http://dx.doi.org/10.3389/fimmu.2022.842605.

1083
1084 Seeley-Fallen, M.K., Liu, L.J., Shapiro, M.R., Onabajo, O.O., Palaniyandi, S., Zhu, X., Tan, T.H.,
1085   Upadhyaya, A. and Song, W. (2014) 'Actin-binding protein 1 links B-cell antigen
1086   receptors to negative signaling pathways', *Proc Natl Acad Sci U S A*, 111(27), 9881-6,
1087   available: http://dx.doi.org/10.1073/pnas.1321971111.

1088
1089 Sharma, S., Orlowski, G. and Song, W. (2009) 'Btk regulates B cell receptor-mediated antigen
1090   processing and presentation by controlling actin cytoskeleton dynamics in B cells', *J
1091   Immunol*, 182(1), 329-39.

1092
1093 Shen, Z., Liu, S., Li, X., Wan, Z., Mao, Y., Chen, C. and Liu, W. (2019) 'Conformational change
1094   within the extracellular domain of B cell receptor in B cell activation upon antigen
1095   binding', *Elife*, 8, available: http://dx.doi.org/10.7554/eLife.42271.

1096
1097 Shlomchik, M.J., Luo, W. and Weisel, F. (2019) 'Linking signaling and selection in the germinal
1098   center', *Immunol Rev*, 288(1), 49-63, available: http://dx.doi.org/10.1111/imr.12744.

1099
1100 Skau, C.T. and Waterman, C.M. (2015) 'Specification of Architecture and Function of Actin
1101   Structures by Actin Nucleation Factors', *Annu Rev Biophys*, 44, 285-310, available:
1102   http://dx.doi.org/10.1146/annurev-biophys-060414-034308.

1103
1104 Sohn, H.W., Tolar, P. and Pierce, S.K. (2008) 'Membrane heterogeneities in the formation of B
1105   cell receptor-Lyn kinase microclusters and the immune synapse', *J Cell Biol*, 182(2),
1106   367-79, available: http://dx.doi.org/10.1083/jcb.200802007.

1107
1108 Song, W., Cho, H., Cheng, P. and Pierce, S.K. (1995) 'Entry of B cell antigen receptor and
1109   antigen into class II peptide-loading compartment is independent of receptor cross-
1110   linking', *J Immunol*, 155(9), 4255-63.

1111
1112 Song, W., Liu, C., Seeley-Fallen, M.K., Miller, H., Ketchum, C. and Upadhyaya, A. (2013) 'Actin-
1113   mediated feedback loops in B-cell receptor signaling', *Immunol Rev*, 256(1), 177-89,
1114   available: http://dx.doi.org/10.1111/imr.12113.

1115

1116 Su, Q., Chen, M., Shi, Y., Zhang, X., Huang, G., Huang, B., Liu, D. and Liu, Z. (2022) 'Cryo-EM
1117   structure of the human IgM B cell receptor', *Science*, 377(6608), 875-880, available:
1118   http://dx.doi.org/10.1126/science.abo3923.

1119
1120 Tanaka, S. and Baba, Y. (2020) 'B Cell Receptor Signaling', *Adv Exp Med Biol*, 1254, 23-36,
1121   available: http://dx.doi.org/10.1007/978-981-15-3532-1_2.

1122
1123 Tojkander, S., Gateva, G., Husain, A., Krishnan, R. and Lappalainen, P. (2015) 'Generation of
1124   contractile actomyosin bundles depends on mechanosensitive actin filament assembly
1125   and disassembly', *Elife*, 4, e06126, available: http://dx.doi.org/10.7554/eLife.06126.

1126
1127 Tolar, P., Hanna, J., Krueger, P.D. and Pierce, S.K. (2009a) 'The constant region of the
1128   membrane immunoglobulin mediates B cell-receptor clustering and signaling in
1129   response to membrane antigens', *Immunity*, 30(1), 44-55, available:
1130   http://dx.doi.org/10.1016/j.immuni.2008.11.007.

1131
1132 Tolar, P., Sohn, H.W., Liu, W. and Pierce, S.K. (2009b) 'The molecular assembly and
1133   organization of signaling active B-cell receptor oligomers', *Immunol Rev*, 232(1), 34-41,
1134   available: http://dx.doi.org/10.1111/j.1600-065X.2009.00833.x.

1135
1136 Tolar, P., Sohn, H.W. and Pierce, S.K. (2005) 'The initiation of antigen-induced B cell antigen
1137   receptor signaling viewed in living cells by fluorescence resonance energy transfer', *Nat
1138   Immunol*, 6(11), 1168-76, available: http://dx.doi.org/10.1038/ni1262.

1139
1140 Treanor, B., Depoil, D., Gonzalez-Granja, A., Barral, P., Weber, M., Dushek, O., Bruckbauer, A.
1141   and Batista, F.D. (2010) 'The membrane skeleton controls diffusion dynamics and
1142   signaling through the B cell receptor', *Immunity*, 32(2), 187-99, available:
1143   http://dx.doi.org/10.1016/j.immuni.2009.12.005.

1144
1145 Unanue, E.R. and Karnovsky, M.J. (1973) 'Redistribution and fate of Ig complexes on surface of
1146   B lymphocytes: functional implications and mechanisms', *Transplant Rev*, 14, 184-210,
1147   available: http://dx.doi.org/10.1111/j.1600-065x.1973.tb00107.x.

1148
1149 Vicente-Manzanares, M., Ma, X., Adelstein, R.S. and Horwitz, A.R. (2009) 'Non-muscle myosin
1150   II takes centre stage in cell adhesion and migration', *Nat Rev Mol Cell Biol*, 10(11), 778-
1151   90, available: http://dx.doi.org/10.1038/nrm2786.

1152
1153 Wang, H., Morse, H.C. and Bolland, S. (2020) 'Transcriptional Control of Mature B Cell Fates',
1154   *Trends Immunol*, 41(7), 601-613, available: http://dx.doi.org/10.1016/j.it.2020.04.011.

1155
1156 Wang, J.C., Yim, Y.I., Wu, X., Jaumouille, V., Cameron, A., Waterman, C.M., Kehrl, J.H. and
1157   Hammer, J.A. (2022) 'A B-cell actomyosin arc network couples integrin co-stimulation to
1158   mechanical force-dependent immune synapse formation', *Elife*, 11, available:
1159   http://dx.doi.org/10.7554/eLife.72805.

1160

1161 Westerberg, L.S., Dahlberg, C., Baptista, M., Moran, C.J., Detre, C., Keszei, M., Eston, M.A.,
1162   Alt, F.W., Terhorst, C., Notarangelo, L.D. and Snapper, S.B. (2012) 'Wiskott-Aldrich
1163   syndrome protein (WASP) and N-WASP are critical for peripheral B-cell development
1164   and function', *Blood*, 119(17), 3966-74, available: http://dx.doi.org/10.1182/blood-2010-
1165   09-308197.

1166

1167 Yang, J. and Reth, M. (2010) 'Oligomeric organization of the B-cell antigen receptor on resting
1168   cells', *Nature*, 467(7314), 465-9, available: http://dx.doi.org/10.1038/nature09357.

1169

 Figure legend

 Figure 1. Arp2/3, activated by N-WASP but not WASP, is required for B-cell contraction.

Splenic B-cells were incubated with planar lipid bilayers coated with monobiotinylated Fab'

fragment of goat anti-mouse IgG+M (Fab'-PLB) in the absence and presence of various

inhibitors and imaged live at 37°C by interference reflection microscopy (IRM). The B-cell

plasma membrane area contacting Fab'-PLB (B-cell contact zone) was measured using IRM

images and custom MATLAB scripts. (**A**) Representative IRM images of splenic B-cells from

C57BL/6 mice treated with CK-689 or CK-666 (50 μM) before (0 min) and after maximal

spreading (2 min). (**B**) Representative plots of the B-cell contact area versus time from one

contracting cell and one non-contracting cell. (**C**) Percentages (±SEM) of B-cells that underwent

contraction after treatment with CK-666 or CK-689. A B-cell was classified as contracting if its

contact zone area was reduced by ≥5% for at least 10 sec after reaching a maximum value. (**D**)

Representative IRM images of splenic B-cells from C57BL/6 mice treated with DMSO or

Wiskostatin (Wisko, 10 μM) 10 min before and during incubation with Fab'-PLB. (**E**)

Percentages (±SEM) of B-cells that underwent contraction after treatment with Wisko or DMSO.

(**F**) Representative IRM images of splenic B-cells from flox control and B-cell-specific N-WASP

knockout (cNKO) mice. (**G**) Percentages (±SEM) of cNKO or flox control B-cells that underwent

contraction. (**H**) Representative IRM images of splenic B-cells from WT or WASP knockout mice

(WKO). (**I**) Percentages (±SEM) of WKO or WT B-cells that underwent contraction. Data points

in **C**, **E**, **G**, and **I** represent three independent experiments, ~25 cells per condition per

experiment, with each color representing one experiment. Scale bar, 2 μm. *p <0.05,

***p<0.001, by paired student's t-test.

Figure supplement 1. B-cells spread and contract on Fab'-coated-planar lipid bilayers.

Figure supplement 2. CK-666 significantly decreases Arp2/3 recruitment to the B-cell contact

zone.

Figure supplement 3. CK-666 treatment before but not after maximal B-cell spreading

decreases the spreading kinetics.

Figure supplement 4. Wiskostatin treatment inhibits N-WASP activation while enhancing

WASP activation in B-cells.

Figure 1–Video 1. Effects of CK-666, Wiskostatin, conditional N-WASP knockout, and WASP

knockout on B-cell contraction.

1201

Figure 2. Arp2/3, downstream of N-WASP, generates inner F-actin foci, driving B-cell

contraction. (**A-C**) WT splenic B-cells were treated with CK-689 or CK-666 (50 µM) during

incubation with Fab'-PLB (**A and B**), and flox control and cNKO B-cells were incubated with

Fab'-PLB at 37°C **(C)**. Cells were fixed at 2 and 4 min, permeabilized, stained for F-actin with

phalloidin, and analyzed using TIRF. Shown are representative TIRF images of phalloidin

staining in the contact zone of CK-689 and CK-666-treated B-cells (**A,** left panels) and

fluorescence intensity (FI) profiles of phalloidin staining along a line crossing cells (**A,** right

panels). Green arrows indicate lamellipodial F-actin, purple arrows indicate inner F-actin foci on

the line, and blue arrows indicate all inner actin foci forming a ring-like structure. Percentages of

cells (per image) (±SEM) with inner F-actin foci forming ring-like distribution among CK-689-

versus CK-666-treated cells (**B**) and flox control versus cNKO B-cells (**C**) before (2 min) and

after (4 min) contraction were determined by visual inspection of phalloidin FI line-profiles

across the B-cell contact zones. Data points in **B** and **C** represent 3 independent experiments,

with each color representing one experiment, 5 images per condition per experiment, and ~15

cells per image. (**D-G**) Inner F-actin foci were identified by their diameter ($\geq$250 nm), peak FI ($\geq$2

fold of no foci area, and location (1 µm away from the outer edge) and quantified as the number

per cell using TIRF images. (**D**) Shown are representative images of splenic B-cells from

LifeAct-GFP-expressing mice treated with CK-689 or CK-666 from 0 min during incubation with

Fab'-PLB at 37°C (top) and the average number (±SEM) of inner LifeAct-GFP foci per cell

(bottom) at 2 min. (**E**) LifeAct-GFP B-cells were treated with CK-666 at 2 min. Shown are

representative TIRF images of LifeAct-GFP in the contact zone of B-cells (top) and the average

number (±SEM) of inner LifeAct-GFP foci (bottom) in the same cell 10s before and 20s after

CK-666 treatment. Arrows indicate disappeared actin foci after CK666 treatment. (**F**) Shown are

representative TIRF images of phalloidin-staining in the contact zone of flox control and cNKO

B-cells after incubating with Fab'-PLB for 2 min (top) and the average number (±SEM) of inner

F-actin foci per cell (bottom). (**G**) Shown are representative TIRF images of WT and WKO B-

cells expressing LifeAct-GFP incubated with Fab'-PLB for 2 min and the average number

(±SEM) of inner F-actin foci per cell (bottom). Data points represent individual cells from three

independent experiments with 10 (**D, F, and G**) or 6 (**E**) cells per condition per experiment.

Scale bar, 2 µm. ** p <0.01, *** p<0.001, by non-parametric student's t-test.

Figure 2-Video 1. Effects of CK-666 and WKO on F-actin foci formation.

1233

Figure 3. Inner F-actin foci are originated from lamellipodia behind the spreading

membrane. Mouse splenic B-cells from LifeAct-GFP transgenic mice were treated with DMSO

or Wisko (10 µM), imaged live using TIRF during incubation with Fab'-PLB at 37°C, and

analyzed using kymographs generated by NIH ImageJ. (**A**) One frame of TIRF time-lapse

images of LifeAct-GFP in the contact zone of B-cells treated with DMSO or Wisko and a WKO

B-cell. Lines indicate eight kymographs that were randomly generated from each cell. (**B**)

Representative kymographs were generated from TIRF time-lapse images of LifeAct-GFP at the

red line in (A). Top panel, a contracting cell. Arrows indicate the start of contraction with inner F-

actin foci originating from lamellipodia. Bottom panel, a non-contracting cell. Lamellipodia-

derived inner F-actin foci were identified by their LifeAct-GFP FI ≥2 fold of their nearby region,

migrating out of the lamellipodial F-actin toward the center of the contact zone, and trackable for

>8 sec. (**C**) Percentages (±SEM) of kymographs showing inner F-actin foci originating from

lamellipodia per cell that did and did not undergo contraction. Data were generated from 3

independent experiments with ~10 cells per condition per experiment. (**D**) A histogram of inner

F-actin foci emerging (expressed as percentages of the total events, blue line) over time relative

to the time of B-cell contraction (defined as 0 sec, indicated by a purple dash line and arrow).

Data were generated from 5 independent experiments with ~9 cells per condition per

experiment. (**E**) Percentage (±SEM) of inner F-actin foci originated from lamellipodia observed

in 8 randomly positioned kymographs of each DMSO- or Wisko-treated WT or untreated WKO

B-cell. Data were generated from 3 independent experiments with ~10 cells per condition per

experiment. Scale bars, 2 µm. *** $p<0.001$, by non-parametric student's *t*-test.

Figure supplement 1. Emerging of Inner F-actin foci from lamellipodia.

1256

Figure 4. N-WASP-activated Arp2/3 sustains the lifetime and the centripetal movement of

inner F-actin foci. LifeAct-GFP-expressing B-cells were incubated with or without various

inhibitors and imaged live by TIRF during interaction with Fab'-PLB at 37°C. Three kymographs

were generated for each cell using time-lapse images and positioned to track as many inner

actin foci as possible. (**A**) A representative kymograph from TIRF time-lapse images of a

DMSO-treated B-cell (Left panels). Inner F-actin foci were identified as described in Figure 3,

and those that emerged during the 60 sec window after maximal spreading (white rectangles in

B) and can be tracked for ≥ 4 sec (dashed lines) in individual kymographs were analyzed.

Relative lifetimes of inner F-actin foci were measured using the duration each focus could be

detected in a kymograph. The relative distances traveled by the foci were measured using the

displacement of each focus in a kymograph. Relative speed was calculated for each inner F-

actin focus by dividing its relative distance by its relative lifetime (right panels). (**B-D**) B-cells

were treated with CK-689 or CK-666 (50 µM) from the beginning of the incubation with Fab'-PLB

(0 min). Shown are representative kymographs (**B**), relative lifetimes (**C**), and relative speed (**D**)

of inner F-actin foci in CK-689- versus CK-666-treated B-cells. (**E-G**) B-cells were treated with

CK-666 at maximal spreading (2 min). Shown are a representative kymograph of a CK-666-

treated cell (**E**), relative lifetimes (**F**), and relative speeds (**G**) of inner F-actin foci in 30-sec

windows before the inhibition and 10 sec after the inhibition in the same cells (linked by blue

lines). (**H-J**) B-cells were treated with DMSO or Wisko (10 µM) 10 min before and during

interaction with Fab'-PLB. Shown is a representative kymograph of a Wisko-treated B-cell (**H**),

relative lifetime (**I**), and relative speed (**J**) of inner F-actin foci in DMSO versus Wisko-treated B-

cells. (**K-M**) LifeAct-GFP-expressing WT and WKO B-cells were incubated with Fab'-PLB.

Shown are a representative kymograph of a WKO B-cell (**K**), relative lifetime (**L**), and relative

speed (**M**) of WT versus WKO B-cells. Data points represent the averaged values from inner F-

actin foci in individual cells, with 3 kymographs per cell and ~12 cells per condition per

experiment from 3 independent experiments. Scale bar, 2 µm. * p <0.05, ** p <0.01, ***

p<0.001, by non-parametric and paired student's t-test.

1284

Figure 5. N-WASP facilitates NMII recruitment and ring-like structure formation, and NMII

promotes inner F-actin foci ring maturation. (**A-C**) B-cells from mice expressing the GFP

fusion of non-muscle myosin IIA (GFP-NMIIA) transgene were treated with DMSO or Wisko (10

µM) 10 min before and during incubation with Fab'-PLB. The B-cell contact zones were imaged

live using TIRF. Shown are representative TIRF images of DMSO- and Wisko-treated B-cells at

30 sec (during spreading) and 2 min 30 sec (after maximal spreading) post landing (**A,** Scale

bars, 2 µm), the averaged GFP-NMIIA MFI (±SEM) (**B**), and the initial rates of increasing

(±SEM) of GFP-NMIIA in the contact zone (the slope of the initial GFP-NMIIA MFI versus time

curves of individual cells) (**C**). Data points represent individual cells from 3 independent

experiments with ~6 cells per condition per experiment. * p <0.05, ***p<0.001, by Kolmogorov-

Smirnov test (**B**) or non-parametric student's t-test (**C**). (**D**) Primary B-cells from mice

expressing both GFP-NMIIA and LifeAct-RFP transgenes were incubated with Fab'-PLB at 37°C

and imaged live by TIRF. Shown are a representative TIRF image of a cell and a kymograph

generated from time-lapse TIRF images at the yellow line. The purple arrow indicates the

starting point of contraction, the white arrow GFP-NMIIA recruitment proximal to the spreading

membrane, and the yellow arrow an F-actin (LifeAct-RFP) focus originating at the lamellipodia

and moving away from the spreading membrane. (**E, F**) Primary B-cells from flox control, WKO,

and cNKO mice were incubated with Fab'-PLB for indicated times. Cells were fixed,

permeabilized, stained for NMII light chain, and imaged by IRM and TIRF. Shown are

representative IRM and TIRF images (**E**) and percentages (±SD) of B-cells with the NMII ring-

like structure in individual images (**F**), identified by visual inspection. The data were generated

from 3 independent experiments with 5 images per condition per experiment. Scale bars, 2 µm.

* p <0.05, ** p <0.01, *** p<0.001, by non-parametric student's t-test. (**G**, **H**) WT splenic B-cells

were treated with DMSO or Blebbistatin (Bleb, 50 µM) 20 min before and during incubation with

Fab'-PLB at 37°C. Cells were fixed at 2 and 4 min, permeabilized, stained with phalloidin, and

imaged by TIRF. Shown are representative TIRF images of the B-cell contact zone (**G**) and

percentages of cells (±SEM) with inner F-actin foci forming ring-like distribution (**H**), determined

as described in Figure 2. Data were generated from 3 independent experiments with ~50 cells

per condition per experiment with different color dots representing individual experiments. Scale

bar, 2 µm. * p <0.05, ** p<0.01 by paired student's t-test.

Figure 5-Video 1. Wiskostatin treatment inhibits NMII ring-like structure formation.

1316

Figure 6. B-cell contraction increases the molecular density within BCR clusters. Primary

B-cells from WT mice were treated with CK-689 or CK-666 from the beginning of the incubation

with AF546-Fab'-PLB (0 min) or at the maximal B-cell spreading (2 min) (**A, D, E, H, I, N, O**).

WT B-cells were treated with DMSO or Wisko (10 µM) 10 min before and during the incubation

with AF546-Fab'-PLB (**B, F, J, P**). B-cells from flox control and cNKO mice were incubated with

AF546-Fab'-PLB (**C, G, K, Q**). The B-cell contact zones were imaged live by TIRF. (**A-C**)

Representative time-lapse images at 30 sec (during B-cell spreading) and 2 min 30 sec (after

maximal spreading) after cell landed. Scale bars, 2 µm. (**D-G**) The MFI of AF546-Fab' in the

contact zone was plotted over time. Purple rectangles indicate the contraction phase based on

the changes in the average B-cell contact area over time. (**H-K**) Rates of AF546-Fab' MFI

increases in the B-cell contact zone before and during contraction were determined by the slope

of AF546-Fab' MFI versus time plots. Data points represent individual cells and were generated

from 3 independent experiments with 6-12 cells per experiment. * p <0.05, ** p <0.01, ***

p<0.001, by Kolmogorov-Smirnov test (D-G) or paired student's t-test (H-K). (**L**) A

representative frame from a time-lapse of a CK-689-treated B-cell (left) shows AF546-Fab'

clusters 30 sec after contraction began, and enlarged time-lapse images (right) show a single

AF546-Fab' cluster over a 40-sec time window after contraction began. AF546 clusters were

identified using the criteria: ≥250 nm in diameter, ≥1.1 fold of FI outside the B-cell contact zone,

and trackable for ≥ 20 sec. Scale bars, 2 μm. (**M**) The peak FI of an AF546-Fab' cluster was

measured over time, and the increasing rate of AF546-Fab' peak FI of this cluster was

determined by the slope of the plot. (**N-Q**) The rates (±SEM) of increase in AF546-Fab' peak FI

in individual clusters were compared between during and after contraction (**N**), between B-cells

treated with CK-689 and CK-666 from 0 min (**O**), between DMSO- and Wisko-treated B-cells

(**P**), and between flox control and cNKO B-cells (**Q**) after B-cells reached maximal spreading.

Data points represent individual cells, the averaged slopes of clusters detected in one B-cell,

from 3 independent experiments with 6~12 cells per condition per experiment. *p <0.05, ** p

<0.01, ***p<0.001, by non-parametric student's t-test.

Figure supplement 1. Fab'-PLB, but not Tf-PLB, induces BCR clustering and phosphorylation.

Figure supplement 2. Tracking and analyzing AF546-Fab' clusters in the B-cell contact zone.

Figure 6-Video 1. Inhibition of B-cell contraction reduces the molecular density within BCR

clusters.

1348

Figure 7. Increased molecular density in BCR clusters leads to reductions in BCR

phosphorylation. (**A-G**) Flox control and cNKO B-cells incubated with AF546-Fab'-PLB were

fixed at 1, 3, 5, and 7 min, permeabilized, stained for phosphorylated CD79a (pCD79a,

Tyr182), and imaged using TIRF and IRM. (**A**) Representative IRM and TIRF images of a flox

control versus a cNKO B-cell at 7 min. Scale bars, 2 μm. (**B-D**) Ratios of pCD79a MFI relative to

AF546-Fab' MFI were plotted against AF546-Fab' peak FI in individual AF546-Fab' clusters in

the contact zone of flox control (**B**), cNKO B-cells (**C**), or flox control and cNKO B-cells overlay

(**D**). AF546-Fab' clusters were identified as described in Figure 6 and Figure 6-figure

supplement 2. Blue dots represent individual AF546-Fab' clusters with an equal number of

clusters from the 4 time points. The black line and diamond symbols represent the average

ratios of pCD79a MFI to Fab' MFI in individual BCR-Fab' clusters within the indicated Fab' peak

FI range. The brown line and square symbols represent the fraction of the AF546-Fab' clusters

out of the total, within the indicated Fab' peak FI range. Clusters were divided into three

populations based on their peak AF546-Fab' FI, relatively low (<190), medium (190-280), and

high (>280, detected only in contracted cells), and the pCD79a to AF546-Fab' MFI ratios of the

three populations were compared (**B** and **C**). Data were generated from 3 independent

experiments with ~20 cells and ≥125 clusters per condition per experiment. * p <0.05, ***

p<0.001, by non-parametric student's t-test. The p-values in **D** were corrected using the

Benjamini-Hochberg/Yekutieli method for false discovery rate control. (**E-G**) The MFI (±SEM) of

AF546-Fab' (**E**) and pCD79a (**F**) and the MFI ratio (±SEM) of pCD79a relative to AF546-Fab'

(**G**) in individual AF546-Fab' clusters at indicated times were compared between flox control and

cNKO B-cells and between different times. (**H-K**) WT B-cells treated with CK-689 or CK-666

after 2 min-incubation with AF546-Fab'-PLB. (**H**) Representative IRM and TIRF images of a CK-

689- versus a CK-666-treated B-cell at 7 min. Scale bars, 2 μm. (**I-K**) The MFI (±SEM) of

AF546-Fab' (**I**) and pCD79a (**J**) and the MFI ratio (±SEM) of pCD79a relative to AF546-Fab' (**K**)

in individual AF546-Fab' clusters were compared between CK-689- and CK-666-treated B-cells

after 7 min stimulation. Data points represent individual clusters. Horizontal solid lines in the

violin plots represent the mean, while the dotted lines represent the quartiles of the distribution.

Data were generated from 3 independent experiments with ~20 cells per condition per

experiment. * p <0.05, *** p<0.001, by non-parametric student's t-test.

1379

Figure 8. Effects of BCR-Fab' density on the association of Syk with BCR-Fab' clusters

and its phosphorylation. Primary B-cells from flox control and cNKO mice were incubated with

AF546-Fab'-PLB at 37°C, fixed at 3 or 7 min, stained for total Syk and phosphorylated Syk

(pSyk Y519/520), and imaged by IRM and TIRF. (**A**) Representative IRM and TIRF images of

Syk staining in a flox control versus a cNKO B-cell at 7 min (Scale bars, 2 μm). (**B-D**) Ratios of

Syk MFI relative to AF546-Fab' MFI were plotted against AF546-Fab' peak IF in individual

AF546-Fab' clusters in the contact zone of Flox control (**B**) and cNKO B-cells (**C**) and their

overlay (**D**). AF546-Fab' clusters were identified as described in Figure 6 and Figure 6-figure

supplement 2 from an equal number of cells after 3- and 7-min stimulation. Blue dots represent

individual AF546-Fab' clusters with an equal number of clusters from each time point. The black

line and diamond symbols represent the average ratios of Syk MFI to Fab' MFI in individual

AF546-Fab' clusters at indicated Fab' peak FI ranges. The brown line and square symbols

represent the fraction of the AF546-Fab' clusters out of the total at indicated Fab' peak FI

ranges. (**E**) Representative IRM and TIRF images of pSyk staining in a flox control versus a

cNKO B-cell at 7 min (Scale bars, 2 μm). (**F-H**) Ratios of pSyk MFI relative to AF546-Fab' MFI

were plotted against AF546-Fab' peak IF in individual AF546-Fab' clusters in the contact zone

of flox control (**F**) and cNKO B-cells (**G**) and their overlay (**H**). Blue dots represent individual

AF546-Fab' clusters with an equal number of clusters from each time point. The black line and

diamond symbols represent the average ratios of pSyk MFI to Fab' MFI in individual AF546-Fab'

clusters at indicated Fab' peak FI ranges. The brown line and square symbols represent the

fraction of the AF546-Fab' clusters out of the total at indicated Fab' peak FI ranges. (**I-K**) MFI

ratios of pSyk relative to AF546-Fab' were plotted against AF546-Fab' peak IF in individual pSyk

puncta in the contact zone of Flox control (**I**) and cNKO B-cells (**J**) and their overlay (**K**). pSyk

puncta were identified using the criteria: FI ≥1.3 fold of the background outside the B-cell

contact zone and diameter ≥250 nm. Blue dots represent individual pSyk puncta with an equal

number of clusters from each time point. The black line and diamond symbols represent the

average ratios of pSyk MFI to Fab' MFI in individual pSyk puncta at indicated Fab' peak FI

ranges. The brown line and square symbols represent the fraction of the pSyk puncta out of the

total at indicated Fab' peak FI ranges. Clusters were divided into three populations based on

their peak AF546-Fab' FI, relatively low (<190), medium (190-280), and high (>280, detected

only in contracted cells), and the Syk (**B** and **C**) or pSyk (**F**, **G**, **I**, and **J**) to AF546-Fab' MFI

ratios of the three populations were compared. Data were generated from 3 independent

experiments with ~23 cells and ≥125 clusters per condition per experiment. * p <0.05, ** p

<0.01, *** p<0.001, by non-parametric student's *t*-test, between AF546-Fab' cluster group with

different Fab' peak FI ranges. The *p*-values in **D, H,** and **K** were corrected using the Benjamini-

Hochberg/Yekutieli method for false discovery rate control.

1416

Figure 9. The effects of BCR-Fab' density on the association of SHIP-1 with BCR-Fab'

clusters and its phosphorylation. Primary B-cells from flox control and cNKO mice were

incubated with AF546-Fab'-PLB at 37°C, fixed at 3 or 7 min, stained for total SHIP-1 and

phosphorylated SHIP-1 (pSHIP-1 Tyr1020), and imaged by TIRF. (**A**) Representative IRM and

TIRF images of SHIP-1 staining in a flox control versus a cNKO B-cell at 7 min (Scale bars, 2

μm). (**B-D**) Ratios of SHIP-1 MFI relative to AF546-Fab' MFI were plotted against AF546-Fab'

peak IF in individual AF546-Fab' clusters in the contact zone of Flox control (**B**) and cNKO B-

cells (**C**). An overlay of flox control and cNKO plots is also shown (**D**). AF546-Fab' clusters were

identified as described in Figure 6 and Figure 6-figure supplement 2 from an equal number of

cells after 3 and 7 min stimulation. Blue dots represent individual AF546-Fab' clusters with an

equal number of clusters from each time point. The black line and diamond symbols represent

the average ratios of SHIP-1 MFI to Fab' MFI in individual AF546-Fab' clusters at indicated Fab'

peak FI ranges. The brown line and square symbols represent the fraction of the AF546-Fab'

clusters out of the total at indicated Fab' peak FI ranges. (**E**) Representative IRM and TIRF

images of pSHIP-1 staining in a flox control versus a cNKO B-cell at 7 min (Scale bars, 2 μm).

(**F-H**) Ratios of pSHIP-1 MFI relative to AF546-Fab' MFI were plotted against AF546-Fab' peak

IF in individual AF546-Fab' clusters in the contact zone of flox control (**F**) and cNKO B-cells (**G**)

and their overlay (**H**). Blue dots represent individual AF546-Fab' clusters with an equal number

of clusters from each time point. The black line and diamond symbols represent the average

ratios of pSHIP-1 MFI to Fab' MFI in individual AF546-Fab' clusters at indicated Fab' peak FI

ranges. The brown line and square symbols represent the fraction of the AF546-Fab' clusters

out of the total at indicated Fab' peak FI ranges. (**I-K**) MFI ratios of pSHIP-1 relative to AF546-

Fab' were plotted against AF546-Fab' peak IF in individual pSHIP-1 puncta in the contact zone

of Flox control (**I**) and cNKO B-cells (**J**) and their overlay (**K**). pSHIP-1 puncta were identified

using the criteria: FI ≥1.5 fold of the background outside the B-cell contact zone and diameter

≥250 nm. Blue dots represent individual pSHIP-1 puncta with an equal number of clusters from

each time point. The black line and diamond symbols represent the average ratios of pSHIP-1

MFI to Fab' MFI in individual pSHIP-1 puncta at indicated Fab' peak FI ranges. The brown line

and square symbols represent the fraction of the total pSHIP-1 puncta at indicated Fab' peak FI

ranges. Clusters were divided into three populations based on their peak AF546-Fab' FI,

relatively low (<200), medium (200-300), and high (>300, detected only in contracted cells), and

the SHIP-1 (**B** and **C**) or pSHIP-1 (**F**, **G**, **I**, and **J**) to AF546-Fab' MFI ratios of the three

populations were compared. Data were generated from 3 independent experiments with ~23

cells and ≥125 clusters per condition per experiment. * p <0.05, ** p <0.01, *** p<0.001, by non-

parametric student's *t*-test, between AF546-Fab' cluster group with different Fab' peak FI

ranges. The *p*-values in **D, H,** and **K** were corrected using the Benjamini-Hochberg/Yekutieli

method for false discovery rate control.

1454

Figure 10. A working model for cell traction-mediated BCR signaling attenuation. When

encountering membrane-associated antigen, mature follicular B-cells undergo rapid spreading,

primarily driven by WASP-mediated branched actin polymerization, which maximizes B-cell

contact with antigen-presenting surface and BCR-antigen engagement, amplifying signaling.

Upon reaching maximal spreading, N-WASP distal to lamellipodial networks activates Arp2/3-

mediated branched actin polymerization, which initiates the generation of inner F-actin foci from

lamellipodia towards the center of the B-cell contact zone. NMII is then preferentially recruited to

these relatively stable inner foci, which in turn promotes the centripetal movement of inner F-

actin foci and the maturation of ring-like actomyosin structures, enabling B-cell contraction. B-

cell contraction pushes the BCR microclusters formed during B-cell spreading to the center of

the contact zone, increasing their molecular density. Increased molecular density promotes the

disassociation of signaling molecules from BCR clusters, probably due to crowding and

conformational changes, leading to signaling attenuation.

1468

Figure 1

1469

Figure 2

Figure 3

A

DMSO

Wisko

WKO

LifeAct

Contraction begins

Not contracting

2 µm

1 µm

0   1   2

0   1   2

Time (min)

0   1   2

B
% Lamellipodia-derived inner F-actin foci / cell

***

Contraction: +   −

C
% Lamellipodia-derived F-actin foci

Contraction begins

Time (s)

D
% Lamellipodia-derived inner F-actin foci / cell

***

DMSO   Wisko   WKO

1471

Figure 4

Figure 5

A

00:30   02:30

DMSO

Wisko

NMII

B

NMII MFI in contact zone

*** / *

DMSO
Wisko

Time (s)

C

NMII recruitment rate (units/s)

***

DMSO   Wisko

D

Contraction begins

LifeAct NMII

Time (s)

NMII   LifeAct

E

| Flox | WKO | cNKO |
| IRM NMII | IRM NMII | IRM NMII |

1 min

3 min

5 min

7 min

F

WKO cNKO Flox

% Cells with myosin ring

*   ***   **   **

1 min   3 min   5 min   7 min

G

DMSO   Bleb

2 min

4 min

F-actin
(Phalloidin)

H

% Cells with inner F-actin foci

DMSO Bleb

**   *

2 min   4 min

1473

Figure 6

Figure 8

1476

Figure 9

Figure 10

1479 **Supplementary materials for this manuscript include the following:**

1480 7 figures

1481 4 videos

Figure 1-S1

Time (min:sec)

Figure 1-figure supplement 1. B-cells spread and contract on Fab'-coated-planar lipid bilayers. Splenic B-cells were pre-warmed to 37°C and incubated with planar lipid bilayers coated with monobiotinylated Fab' fragment of goat anti-mouse IgG+M (Fab'-PLB) and imaged live at 37°C by interference reflection microscopy (IRM). Shown are individual frames from a time-lapse IRM image of one B-cell. The plasma membrane area contacting with Fab'-PLB (B-cell contact zone) visualized by IRM increased between 0-1 min after landing, indicating spreading, and decreased after 1 min 30 sec, indicating contraction. Scale bar, 2 μm.

Figure 1-S2

Figure 1-figure supplement 2. CK-666 significantly decreases Arp2/3 recruitment to the B-cell contact zone. WT splenic B-cells were treated with CK-689 or CK-666 (50 μM) before (0 min) and after maximal spreading (2 min) during incubation with Fab'-PLB at 37°C. Cells were fixed at 3 and 7 min, permeabilized, stained for Arp2, and imaged using IRM and TIRF. Shown are representative images (**A**) and the MFI of Arp2 in the contact zone at 3 min and 7 min compared between B-cell treated with CK-689 (black dots), CK-666 from 0 min (red dots), and CK-666 from 2 min (purple dots) (**B**). Data points represent individual cells from 3 independent experiments with ~20 cells per condition per experiment. Scale bar, 2 μm. *** $p<0.001$, by non-parametric student's *t*-test.

Figure 1-S3

Figure 1-figure supplement 3. CK-666 treatment before but not after maximal B-cell spreading decreased the spreading kinetics. WT splenic B-cells were treated with CK-689 or CK-666 (50 µM) before (0 min) and after maximal spreading (2 min) during incubation with Fab'-PLB and imaged live at 37°C by IRM. The area occupied by the B-cell contact zone was measured using IRM images and custom codes made in MATLAB. The mean spreading rate of each cell during its early spreading phase was quantified using the contact area versus the time curve of that cell by linear regression. The averaged spreading rates (±SEM) were generated from 3 independent experiments with ~15 cells per condition per experiment. * $p > 0.05$, ** $p < 0.01$, by non-parametric student's *t*-test.

Figure 1-S4

Figure 1-figure supplement 4. Wiskostatin treatment inhibits N-WASP activation while enhancing WASP activation in B-cells. WT splenic B-cells were pre-treated with Wiskostatin (Wisko, 10 µM) or DMSO (control) for 10 min at 37°C before and during incubation with Fab'-PLB. Cells were fixed at 3 and 7 min, permeabilized, stained for phosphorylated N-WASP (pN-WASP) or WASP (pWASP), and imaged using IRM and TIRF. Shown are representative IRM and TIRF images of pN-WASP (**A**) and pWASP (**B**) at the B-cell contact zone and the MFI (±SEM) of pN-WASP (**C**) and pWASP (**D**) in the B-cell contact zone, comparing between DMSO and Wisko-treated B-cells. Data points represent individual cells from 3 independent experiments with ~25 cells per condition per experiment. Scale bar, 2 µm. ** $p<0.01$, *** $p<0.001$, by non-parametric student's *t*-test.

Figure 3-S1

1521

Figure 3-figure supplement 1. Emerging of Inner F-actin foci from lamellipodia. Splenic B-

cells from LifeAct-GFP transgenic mice were treated with DMSO, imaged live using TIRF and

IRM during incubation with Fab'-PLB at 37°C, and analyzed using kymographs that were

randomly generated from each cell. Shown are six examples of the kymographs used for

analysis. Arrows indicate the emergence of inner F-actin foci near the lamellipodia.

Lamellipodia-derived inner F-actin foci were identified by their LifeAct-GFP FI ≥2 fold of their

nearby region, inside location in the contact zone, migrating away the lamellipodial F-actin, and

trackable for ≥8 sec.

1530

Figure 6-S1

Figure 6-figure supplement 1. Fab'-PLB, but not Tf-PLB, induces BCR clustering and phosphorylation. WT splenic B-cells were pre-labeled with Cy3-Fab fragment of goat anti-mouse IgM+G at a concentration of 2.5 μg per 10^6 cells at 4°C for 30 min, followed by incubation with Fab'-PLBs or Tf-PLBs for 5 min at 37°C. Cells were fixed, permeabilized, stained for pCD79a, and imaged using IRM and TIRF. Shown are representative IRM and TIRF images from three independent experiments. Scale bar, 2 μm.

Figure 6-S2

A Time elapsed since contraction began (s):
0   20   40   60   80   100   120   140

IRM

B
Cluster 1
Cluster 2
Cluster 3

Fab'

C
Cluster 1
Cluster 2
Cluster 3

D Cluster tracks over 140s since contraction began

E Fab' Peak Intensity vs Time

Fab' Peak Intensity

Time elapsed since contraction began (s)

Cluster 1   Cluster 2   Cluster 3

F

Fab' Peak Intensity

Cluster 1   Cluster 2   Cluster 3

Fab'

1538

Figure 6-figure supplement 2. Tracking and analyzing AF546-Fab' clusters in the B-cell

contact zone. WT splenic B-cells were incubated with AF546-Fab'-PLB at 37°C and imaged

live using IRM and TIRF. Shown are individual frames from time-lapse images of IRM (**A**) and

TIRF (**B**), showing AF546-Fab' clusters within the contact zone of one DMSO-treated (vehicle

control for Wisko) B-cell for 140 sec since the beginning of contraction. Fab' FI is shown as heat

maps using NIH ImageJ. The boundary of the contact zone, detected using IRM images by a

custom MATLAB script, is shown in yellow dashed lines. Arrows point to three representative

clusters among the other clusters detected in the contact zone using custom MATLAB codes.

Cluster detection masks for the three representative clusters are shown (**C**). Moving tracks for

the three AF546-Fab' clusters are shown alongside the initial (black dashed lines) and final state

(gray dashed lines) of the contact zone (**D**). Tracks were generated by following the peak of

AF546 FI in each cluster as it moved. AF546-Fab' peak FI versus time curves for the three

representative clusters are plotted over the duration that each cluster could be detected (**E**).

Surface plots (2.5-D plots) of AF546-Fab' FI show a zoomed-in region consisting of each of the

three AF546-Fab' clusters (**F**). Colors in (**B**) and (**F**) are scaled to AF546-Fab' FI values. Scale

bars, 1 µm.

Video Legends

Figure 1-Video 1. Effects of CK-666, Wiskostatin, conditional N-WASP knockout, and WASP knockout on B-cell contraction. Splenic B-cells were incubated with planar lipid bilayers coated with monobiotinylated Fab' fragment of goat anti-mouse IgG+M (Fab'-PLB) in the absence and presence of various inhibitors and imaged live at 37°C at one frame per 2 seconds by interference reflection microscopy (IRM). Shown are representative IRM time-lapse images of WT B-cells treated with CK-689 or CK-666 (50 µM) before (0 min) and after maximal spreading (2 min) (**A**), WT B-cells treated with DMSO or Wiskostatin (Wisko, 10 µM) 10 min before and during incubation with Fab'-PLB (**B**), B-cells from flox control and B-cell-specific N-WASP knockout (cNKO) mice (**C**), and B-cells from WT or WASP knockout mice (WKO) (**D**). The frame in which the contact zone first appears was considered time 0. The videos are sped up by 20x compared to real time. Scale bar, 2 µm.

Figure 2-Video 1. Effects of CK-666 and WKO on F-actin foci formation. Splenic B-cells from LifeAct-GFP transgenic mice were incubated with Fab'-PLB in the absence and presence of CK-689 or CK-666 and imaged live at 37°C at one frame per 2 seconds by IRM and total internal reflection fluorescence microscopy (TIRF). Shown are representative IRM (**A** and **C**) and TIRF (**B** and **D**) time-lapse images of WT B-cells treated with CK-689 or CK-666 (50 µM) before and after maximal spreading (**A** and **B**) and WT and WKO B-cells expressing LifeAct-GFP (**C** and **D**). The video is sped up by 20x compared to real time. Scale bar, 2 µm.

Figure 5-Video 1. Wiskostatin treatment inhibits NMII ring-like structure formation. B-cells from mice expressing GFP fusion of non-muscle myosin IIA (GFP-NMIIA) and LifeAct-RFP transgenes were treated with DMSO or Wisko (10 µM) 10 min before and during incubation with Fab'-PLB. The B-cell contact zones were imaged live at one frame per 2 seconds using IRM

1580 and TIRF. Shown are representative IRM and TIRF time-lapse images. The video is sped up by

1581 20x compared to real time. Scale bar, 2 µm.

1582

Figure 6-Video 1. Inhibition of B-cell contraction reduces the molecular density within

BCR clusters. Splenic B-cells were incubated with AF546-Fab'-PLB in the absence and

presence of various inhibitors and imaged live at 37^{o}C at one frame per 2 seconds by IRM and

TIRF. Shown are representative IRM **(A, C,** and **E)** and TIRF **(B, D,** and **F)** time-lapse images of

WT B-cells treated with CK-689 or CK-666 (50 µM) before and after maximal spreading (**A** and

B), WT B-cells treated with DMSO or Wisko (10 µM) 10 min before and during incubation with

Fab'-PLB (**C** and **D**), and flox control or cNKO B-cells (**E** and **F**). TIRF images (**B, D,** and **F**) are

shown as AF546-Fab' FI maps. The video is sped up by 20x compared to real time. Scale bar, 2

µm.