# Reciprocal discoidin domain receptor signaling strengthens integrin adhesion to connect adjacent tissues

[#]Kieop Park[1], [#]Ranjay Jayadev[1], [#]Sara G. Payne[1,2], Isabel W. Kenny-Ganzert[1], Qiuyi Chi[1], Daniel S. Costa[1], William Ramos-Lewis[1], Siddharthan Balachandar Thendral[1], and [*]David R. Sherwood[1]

[1] Department of Biology, Duke University, Box 90338, Durham, NC 27708, USA

[2] Department of Cell Biology, Duke University Medical Center, Durham, NC 27708, USA

[#]Shared first author
[*]Correspondence: david.sherwood@duke.edu

Running title: DDR-2 signaling connects tissues

Key words: Discoidin domain receptor, type IV collagen, integrin, endocytosis, Ras, basement membrane, tissue connection

## Abstract

Separate tissues connect through adjoining basement membranes to carry out molecular

barrier, exchange, and organ support functions. Cell adhesion at these connections must be

robust and balanced to withstand independent tissue movement. Yet, how cells achieve

synchronized adhesion to connect tissues is unknown. Here, we have investigated this question

using the *C. elegans* utse-seam tissue connection that supports the uterus during egg-laying.

Through genetics, quantitative fluorescence, and cell specific molecular disruption, we show

that type IV collagen, which fastens the linkage, also activates the collagen receptor discoidin

domain receptor 2 (DDR-2) in both the utse and seam. RNAi depletion, genome editing, and

photobleaching experiments revealed that DDR-2 signals through LET-60/Ras to coordinately

strengthen an integrin adhesion in the utse and seam that stabilizes their connection. These

results uncover a synchronizing mechanism for robust adhesion during tissue connection,

where collagen both affixes the linkage and signals to both tissues to bolster their adhesion.

**Introduction**

Basement membrane (BM) is a planar, cell-associated extracellular matrix (ECM) that underlies or surrounds most tissues. BM provides tissues with structural support, barrier functions, and signaling platforms (Morrissey and Sherwood 2015, Jayadev and Sherwood 2017, Pozzi, Yurchenco et al. 2017). The two core BM matrix components are the heterotrimeric proteins laminin and type IV collagen. Laminin is comprised of an α, β, and γ subunit. Once secreted, laminin associates with cell surface receptors and self-assembles into a sheet-like network. Type IV collagen is a large protein comprised of two α1-like and one α2-like chains, and these heterotrimers cross-link to form a grid, which is thought to be connected to the laminin network through bridging molecules (Hohenester and Yurchenco 2013). Type IV collagen provides BM with tensile strength, which allows BM to structurally support tissues and resist mechanical forces (Fidler, Boudko et al. 2018). Type IV collagen is a ligand for several vertebrate integrin receptors and the collagen-specific receptor tyrosine kinase discoidin domain receptor 1 (DDR1) (Khoshnoodi, Pedchenko et al. 2008, Brown, Cummings et al. 2017). Integrin and DDR1 interactions with type IV collagen mediate a wide variety cell-matrix functions, including cell migration and cell adhesion (Khoshnoodi, Pedchenko et al. 2008, Castro-Sanchez, Soto-Guzman et al. 2010, Borza and Pozzi 2014, Xiao, Jiang et al. 2015). DDR receptors and integrin also functionally interact within cells (Leitinger 2014); however, the roles of these interactions in native tissue settings remains poorly defined.

At tissue interfaces juxtaposed BMs usually slide along each other and maintain distinct tissue boundaries (Sherwood and Sternberg 2003, Brown 2011). At specific sites, however, the BMs of neighboring tissues link to stabilize tissue interactions and build complex organs (Keeley and Sherwood 2019). Examples include the kidney, where podocytes and endothelial BMs connect to form the glomerular BM blood filtration unit (Abrahamson 1985); the brain, where the astrocyte end feet and endothelial BMs join to build the blood-brain barrier (Sixt, Engelhardt et

al. 2001); and the somite, where somite and epidermal BMs link to maintain somite-epidermal

association during development (Feitosa, Zhang et al. 2012). Approximately 20 different BM-BM

linkages between neighboring tissues have been documented (Gao, Mruk et al. 2017, Keeley

and Sherwood 2019, Welcker, Stein et al. 2021). Disruption of these BM-BM tissue connections

appear to underlie human diseases, such as in Alport's syndrome, where defects in kidney and

hearing are associated with tissue linkage disruption (Merchant, Burgess et al. 2004, Naylor,

Morais et al. 2021). Despite the prevalence and importance of BM-BM tissue connections, the

challenge of studying tissue interactions *in vivo* in vivo has hampered our understanding of the

mechanisms establishing and maintaining these linkages.

The simple tissues, visual transparency, and experimental tractability of *C. elegans* offers a

powerful model to study BM-BM tissue interactions. Further, most *C. elegans* BM components

and receptors are endogenously tagged with genetically encoded fluorophores to visualize

localization, levels, and dynamics (Keeley, Hastie et al. 2020, Jayadev, Morais et al. 2022). A

stable BM-BM tissue connection between the large, multinucleated uterine utse cell and

epidermal seam cells stabilizes the uterus during egg laying (Vogel and Hedgecock 2001,

Morrissey, Keeley et al. 2014, Gianakas, Keeley et al. 2023). Disruption of the utse-seam BM-

BM tissue linkage results in prolapse of the uterus (**Rup**tured phenotype) out of the animal

during egg-laying. Formation of the utse-seam attachment is initiated by the deposition of the

matrix proteins fibulin and hemicentin during the last larval stage of development. Hemicentin

promotes the recruitment of type IV collagen, which accumulates at high levels at the BM-BM

tissue connection and strengthens the adhesion, allowing it to resist the strong mechanical

forces of egg-laying (Gianakas, Keeley et al. 2023). The integrin, $\alpha$INA-1, one of two *C. elegans*

alpha integrins, is also expressed in the utse and is required to prevent uterine-prolapse during

egg-laying. However, its regulation and function at the utse-seam connection is unclear

(Morrissey, Keeley et al. 2014, Gianakas, Keeley et al. 2023). Mechanisms that link juxtaposed

BMs appear to be shared, as hemicentin-1 promotes BM-BM tissue connection in the zebrafish

fin fold, hemicentin-2 and fibulin-1 mediate zebrafish somite-epidermis BM-BM association, and

hemicentin-1 stabilizes BM-BM linkage at myotendinous junctions in mice (Carney, Feitosa et

al. 2010, Feitosa, Zhang et al. 2012, Welcker, Stein et al. 2021). In addition, type IV collagen

bridges the podocyte BM and endothelial BM to form the glomerular BM in mice and humans

(Suleiman, Zhang et al. 2013, Naylor, Morais et al. 2021). While significant insights are being

gained on matrix composition at BM-BM tissue connections, it remains unknown how tissues

coordinate and strengthen their adhesive activity at linkage sites. Cells at tissue connections

must resist the forces of tissue shifting arising from growth, muscle contraction, and blood flow,

and it is unclear how tissues synchronize and bolster adhesions at tissue linkage sites.

Here, we discover that the *C. elegans* discoidin domain receptor DDR-2, an ortholog to the two

vertebrate collagen binding DDR tyrosine kinase receptors (RTK), coordinates and strengthens

an integrin mediated cell adhesion to support the utse-seam BM-BM tissue linkage during its

formation. Using genetics, endogenous tagging, and tissue-specific molecular perturbations, we

show that DDR-2 is expressed and functions in the utse and seam cells to mediate utse-seam

attachment between the mid and late L4 larval stages. We reveal that assembly of type IV

collagen linking the juxtaposed BMs triggers the internalization of DDR-2 in both the utse and

seam cells into endocytic vesicles, a signaling compartment for DDR/RTK receptors. In the *ddr-*

*2* knockout mutant, gaps in utse-seam attachment occurred at this time. Through tissue-specific

perturbation, quantitative fluorescence analysis, and fluorescence recovery after photobleaching

(FRAP), we provide evidence that DDR-2 signals through LET-60/Ras to increase the levels

and stability of the α-integrin INA-1 at the utse-seam linkage. Together, these results reveal a

mechanism for synchronizing a robust adhesion during tissue linkage, where type IV collagen,

which bridges and fastens the BM-BM tissue linkage, also serves as a signal to coordinate and

bolster cell-matrix adhesion in cells on both sides of the tissue connection.

## Results

### The utse and seam cells associate during the early-L4 stage

The utse is an H-shaped syncytial uterine cell that underlies the *C. elegans* uterus and opens
along the central crossbar region during egg-laying to allow embryo passage (Ghosh and
Sternberg 2014). The utse is flanked laterally on both sides of the animal by a string of
epidermal seam cells that run the length of the body. The utse cell and seam cells are each
encased in a BM, which become linked by a specialized matrix whose known functional
components include hemicentin, fibulin, and type IV collagen (***Figure 1A***) (Gianakas, Keeley et
al. 2023). The BM-BM tissue connection supports the uterus during egg-laying and disruption of
the linkage results in uterine prolapse (***Figure 1B and C***) (Vogel and Hedgecock 2001). The
utse and seam cells are in contact by the mid L4 larval stage (Gianakas, Keeley et al. 2023), but
when the initial contact is established is not known.

The utse is formed from the fusion of one specialized uterine cell, the anchor cell, with eight
adjacent uterine pi cell progeny, forming the syncytial nine-nuclei utse during the early L4 stage
(Newman, White et al. 1996, Ghosh and Sternberg 2014). To determine when the utse first
contacts the seam cells, we used an anchor cell-specific mCherry fluorophore marker (*cdh-3p::mCherry::PLCδ$^{PH}$*; ***Figure 1D***, arrow) (Hagedorn, Yashiro et al. 2009) that diffuses into the
uterine pi cell progeny during fusion and simultaneously viewed a GFP seam cell marker
(*scmp::GFP::CAAX*) (Chapman, Li et al. 2008). We found that the utse contacts the seam cells
during the early L4 stage as the utse syncytium forms (***Figure 1D***, arrowheads; n=5/5 cases
observed). Shortly after utse-seam contact, the BM-BM linkage is initiated at the mid L4 stage
by the assembly of low levels of hemicentin, which is secreted by the utse and fibulin-1, which
hemicentin recruits from the extracellular fluid. Fibulin-1 and hemicentin protect the utse-seam
cell connection from mechanical forces arising from body movement and uterine and vulval

muscle contractions prior to egg-laying (Gianakas et al. 2023). Hemicentin also promotes

recruitment of type IV collagen from the extracellular fluid, which allows the utse-seam BM-BM

connection to resist the strong mechanical forces from egg-laying (Gianakas et al. 2023)

(***Figure 1D***). Together, these results show that the utse and seam cells are in contact prior to

deposition of matrix components that strengthen the utse-seam BM-BM tissue connection.

**Loss of discoidin domain receptor-2 (DDR-2) causes uterine prolapse**

As type IV collagen is critical to the BM-BM connection, we were next interested in

understanding how receptors for type IV collagen regulate the utse-seam BM-BM linkage. The

two main receptors for type IV collagen are integrin and the discoidin domain receptor (DDR)

(Fidler, Boudko et al. 2018). *C. elegans* harbor two integrin heterodimers composed of different

α chains with a shared β chain: αINA-1/ βPAT-3 and αPAT-2/ βPAT-3 (Clay and Sherwood

2015). The integrin INA-1 is expressed in the utse and its loss leads to uterine prolapse,

suggesting a possible role in utse-seam attachment (Morrissey, Keeley et al. 2014, Gianakas,

Keeley et al. 2023). Whether DDRs function at the utse-seam connection, however, is unknown.

*C. elegans* have two DDR genes, *ddr-1* and *ddr-2,* which are paralogs and orthologs to the two

mammalian DDRs (Vogel, Abdulhussein et al. 2006)*.* Like mammalian counterparts, the *C.*

*elegans* DDRs are comprised of discoidin and discoidin-like domains, followed by a

transmembrane domain and a kinase domain (***Figure 2A***). To determine if *C. elegans* DDRs

play a role in utse-seam BM-BM linkage, we examined several *ddr-1* and *ddr-2* mutants (***Figure***

***2A***). We first screened for the Rup phenotype caused by uterine prolapse. Animals harboring

the *ddr-1(ok874)* deletion allele, which eliminates the intracellular portion of DDR-1, and animals

with the *ddr-1(tm382)* allele containing an early stop codon truncating the majority of the

predicted protein (Unsoeld, Park et al. 2013), did not show a Rup phenotype (***Figure 2A and***

**B**). In contrast, animals homozygous for the *ddr-2(tm797)* allele, which introduces an early stop

codon and is thought to be a null or strong loss-of-function (Unsoeld, Park et al. 2013), exhibited

a significant Rup/uterine prolapse defect. Animals carrying the *ddr-2(ok574)* allele, which

deletes a portion of the intracellular kinase domain (Unsoeld, Park et al. 2013), also trended

towards displaying a significant Rup phenotype (**Figure 2A and B**). We further generated a full-

length *ddr-2* deletion allele, *ddr-2(qy64)*, and confirmed that complete loss of *ddr-2* led to

penetrant uterine prolapse (**Figure 2A and B**). Taken together, these results indicate that DDR-

2 protects against uterine prolapse and may play a role in mediating utse-seam BM-BM

attachment.

**DDR-2 maintains utse-seam attachment during formation of the tissue connection**

To determine if the Rup phenotype observed in *ddr-2* mutants is a result of perturbations in the

utse-seam BM-BM tissue connection, we examined utse-seam association from the late L3

through young adult stages in animals harboring the *ddr-2(qy64)* deletion allele (**Figure 2C**). We

visualized utse-seam association using the utse and seam cell membrane markers *cdh-*

*3p::mCherry::PLCδ$^{PH}$* and *scmp::GFP::CAAX,* respectively. Animals homozygous for *ddr-*

*2(qy64)* had normal utse-seam cell association at the early L4 stage; however, beginning at the

mid L4 stage, gaps in the utse-seam attachment were detected in about 30% of animals. The

frequency of gaps increased to ~40% of animals by the mid-to-late L4 and persisted at a similar

penetrance to young adulthood (**Figure 2C and D**). These observations suggest that DDR-2

functions around the mid L4 to late L4 stages to promote utse-seam attachment. Next, we

wanted to determine if the utse-seam detachment observed in *ddr-2(qy64)* mutants led to

uterine prolapse. Wild-type and *ddr-2(qy64)* animals were assessed at the L4 larval stage for

utse-seam attachment defects and tracked to the 72-hour adult stage, where they were

examined for the Rup phenotype. All wild-type L4 larvae had intact utse-seam attachment and

did not rupture by 72-hour adulthood (n=8/8 animals). However, among nine *ddr-2(qy64)* mutant

animals examined, four exhibited utse-seam detachments at the L4 stage, all of which

underwent uterine prolapse (Rup) by adulthood. The remaining five *ddr-2* mutant animals, which

had an intact utse-seam linkage, did not display the Rup phenotype. Taken together, these

results suggest that DDR-2 promotes utse-seam attachment during early stages of tissue

linkage.

**DDR-2 is found within endocytic vesicles in the utse and seam cells during initial stages**

**of tissue attachment**

As utse-seam detachment in *ddr-2* mutant animals was first observed at the mid L4 larval stage,

we next sought to determine if DDR-2 was expressed in the utse, seam, or both cell types at

this time. Between the mid L4 and late L4 stage, the utse-seam tissue connection forms into a

tongue-and-groove association between the utse (tongue) and seam (groove) (***Figure 2D and***

***Figure 2—video 1***). To resolve whether DDR2 is localized within the seam, utse, or both (Vogel

and Hedgecock 2001), we used either a utse (*cdh-3p::mCherry::PLCδ$^{PH}$*) or seam cell marker

(*scmp::2xmKate2::PLCδ$^{PH}$*), and imaged endogenously tagged DDR-2 protein (DDR-2::mNG)

(Keeley, Hastie et al. 2020) in lateral confocal sections from the L4 to young adult stages

(***Figure 3A***). We first examined the seam cells contacting the utse (***Figure 3B***). At the early L4

stage DDR-2::mNG was faintly visible at the surface of the seam cells and found in a few

intracellular punctae (***Figure 3B,*** n=10/10 animals). By the mid L4 stage, DDR-2 was

predominantly found in punctae in the region of attachment with the utse. Levels of punctae

increased until the late L4 stage and then declined sharply by the young adult (***Figure 3B***).

Notably, at the late L4, when DDR-2 punctate localization was at its highest, we could not detect

any DDR-2 signal at the cell surface (***Figure 3B,*** n = 9/9). We further examined DDR-2::mNG in

seam cells that were not contacting the utse and found that DDR-2 levels were uniformly low

from the early L4 through the young adult stages (**Figure 3—figure supplement 1A**). We next

investigated DDR-2 localization in the utse contacting the seam. Like the seam cells at the

tissue connection site, DDR-2 in the utse was punctate and peaked in levels between the mid

and late L4 stages and then declined rapidly (**Figure 3C**). We conclude that DDR-2 is present at

high levels and localized within punctate structures in the seam and utse cells at the time when

utse-seam defects are observed in *ddr-2* mutant animals.

Receptor tyrosine kinases (RTKs), including DDRs, are activated upon ligand binding and then

rapidly internalized into endocytic vesicles: first the early endosome (marked by RAB-5),

followed by subsequent sorting for either lysosome-mediated degradation via late endosomes

(marked by RAB-7), or recycled back to the cell surface in recycling endosomes (marked by

RAB-11) (Goh and Sorkin 2013). In vertebrates, the exposure of cells to collagen results in the

internalization of DDR1b into Rab5a positive early endosomes (Mihai, Chotani et al. 2009).

Phosphorylation of DDR1 does not occur until the onset of endocytosis and recycling,

suggesting that receptor activation occurs in endocytic vesicles (Mihai, Chotani et al. 2009, Fu,

Valiathan et al. 2013). As *C. elegans* DDR-2 appeared to be functioning during the mid L4 to

late L4 stages, when levels of DDR-2 were highest and present in vesicular structures, we

sought to determine whether DDR-2 was internalized within endocytic vesicles. We examined

localization of DDR-2::mNG punctae in relation to utse-expressed markers of early endosomes

(*cdh-3p::mCherry::rab-5*), late endosomes (*cdh-3p::mCherry::rab-7*), and recycling endosomes

(*cdh-3p::mCherry::rab-11*) (Sato, Norris et al. 2014). DDR-2 was co-localized with all three

classes of endosomes in the utse, with the highest overlap with RAB-5 early endosomes

(**Figure 3—figure supplement 1B**). In addition, we examined an early endosome marker

expressed in seam cells (*scmp::mKate2::rab-5*) and observed DDR-2::mNG punctae co-

localized with RAB-5 vesicles (**Figure 3—figure supplement 1C**). The presence of DDR-2 in

endocytic vesicles within the utse and seam suggested that DDR-2 signaling may be required

for utse-seam attachment. Consistent with this, animals harboring the *ddr-2(ok574)* mutant

allele, where a portion of the intracellular kinase domain is deleted (**Figure 2A**), exhibited

significant utse-seam detachments at the late L4 stage (**Figure 4A**). Taken together, these

observations suggest that DDR-2 functions in the utse and seam cells to promote attachment at

the tissue linkage site between the mid and late L4 stages, where it may signal within endocytic

vesicles.

**Type IV collagen promotes DDR-2 vesiculation**

DDR signaling is triggered by collagen binding, and vertebrate DDRs bind multiple collagens

with different binding affinities (Leitinger 2014). Genetic interaction studies in *C. elegans*

revealed that transmembrane collagen COL-99, which is most similar to transmembrane

collagen type XIII, XXIII, and XXV in vertebrates (Tu, Huhtala et al. 2015), may act as a ligand

for *C. elegans* DDR-1 and DDR-2 during axon guidance and that type IV collagen might act as a

ligand for DDR-2 during axon regeneration (Hisamoto, Nagamori et al. 2016, Taylor, Unsoeld et

al. 2018). *C. elegans* BMs also harbor type XVIII collagen (CLE-1) (Keeley, Hastie et al. 2020).

Thus, we next investigated whether any of these cell and BM-associated collagens could

function as a ligand for DDR-2 at the utse-seam BM-BM tissue connection site.

Type IV collagen enriches at the utse-seam BM-BM linkage during its maturation and helps the

connected tissues resist the mechanical forces of egg-laying. In contrast, type XVIII collagen is

not present enriched at the linkage (Gianakas, Keeley et al. 2023). Whether COL-99 localizes to

the utse-seam tissue linkage is unknown. We thus first examined an endogenously tagged

COL-99 (mNG::COL-99) (Jayadev, Morais et al. 2022) and compared its localization and levels

to type IV collagen ($\alpha$1-type IV collagen::mNG (EMB-9::mNG)) and type XVIII collagen (CLE-

1::mNG) (Keeley, Hastie et al. 2020) at the utse-seam connection in late L4 animals. COL-99

was not detected while EMB-9 was present at nearly 8-fold higher levels than CLE-1,

suggesting that type IV collagen might be the ligand for DDR-2 (***Figure 4B***). Furthermore, the

enrichment of type IV collagen levels at the utse-seam connection site occurs rapidly from the

mid L4 to late L4 stage (Gianakas, Keeley et al. 2023), correlating with the buildup of DDR-2 in

endosomes in the utse and seam cells.

As exposure to collagen triggers vesiculation of vertebrate DDR1b into endocytic vesicles, we

wanted to determine whether type IV collagen controls vesiculation of DDR-2 at the utse-seam

BM-BM connection. RNAi-mediated depletion of type IV collagen from the L1 through the late

L4 larval stages resulted in a sharp decrease in vesicular DDR-2::mNG in the utse and seam

cells at the linkage site, accompanied by enrichment of DDR-2 at cell surfaces (***Figure 4C***;

***Figure 4—figure supplement 1A***). In contrast, reduction of COL-99 and type XVIII collagen did

not significantly affect DDR-2 vesiculation (***Figure 4C***; ***Figure 4—figure supplement 1B and***

***C***). Depletion of hemicentin, which plays a crucial role in recruiting type IV collagen to the BM-

BM linking matrix (Gianakas, Keeley et al. 2023), also resulted in a significant reduction in DDR-

2 vesiculation (***Figure 4D***; ***Figure 4—figure supplement 1D***). In addition, we did not detect

DDR-2 at the cell surface, suggesting that hemicentin has a role in recruiting DDR-2 to the site

of utse-seam attachment. Together, these findings indicate that type IV collagen assembly at

the site of utse-seam BM-BM tissue linkage triggers DDR-2 vesiculation and may act as a ligand

for DDR-2 activation to promote utse-seam attachment.

**DDR-2 functions in the utse and the seam to promote utse-seam attachment**

Since DDR-2 localizes to both the utse and seam cells, we next investigated where DDR-2

functions to promote utse-seam BM-BM attachment. We generated genome edited lines with

single-copy insertions of dominant negative (DN) *ddr-2* (Bernadskaya, Brahmbhatt et al. 2019)

driven by the *cdh-3* promoter for utse expression or the *scm* promoter for seam expression.

Using the utse-specific membrane marker (*nas-22p::2xmKate2::PLCδ^PH*) (Park, Pan et al. 2010)

and a seam cell membrane marker (*wrt-2p::GFP::PLCδ^PH*) (Wildwater, Sander et al. 2011), we

found significant utse-seam detachments in both utse-expressed and seam-expressed *DN ddr-2*

animals at the late L4 stage (***Figure 4F and G***). Consistent with the timing and vesicular pattern

of DDR-2 localization, these results indicate that DDR-2 functions in both the utse and seam

between the mid and late L4 stages to promote utse-seam attachment.

**Loss of DDR-2 increases type IV collagen at the utse-seam BM-BM adhesion site**

The matrix components fibulin-1 and hemicentin play early roles in mediating utse-seam

attachment and hemicentin promotes recruitment of type IV collagen and further enrichment of

fibulin, which strengthens the BM-BM attachment (Gianakas et al. 2023). We thus next asked

whether DDR-2 promotes utse-seam linkage by regulating the assembly of fibulin, hemicentin,

or type IV collagen at the connecting matrix. Knockdown of *ddr-2* by RNAi did not alter fibulin

levels at the late L4 stage (***Figure 5—figure supplement 1A and B***). However, we observed a

modest ~20% reduction in hemicentin upon depletion of DDR-2 (***Figure 5—figure supplement***

***1C***). As loss of hemicentin results in utse-seam detachment at the late L4 stage (Gianakas,

Keeley et al. 2023), we tested whether a ~20% reduction of hemicentin could cause these

defects by initiating RNAi at the early L4, just before the utse forms (***Methods***). We found that

utse-seam attachments were intact in all animals at the late L4 stage upon ~20% reduction of

hemicentin (n=13/13 and 12/12 control and *him-4* RNAi-treated animals examined respectively;

***Figure 5—figure supplement 1D***), suggesting that the utse-seam detachment defect in *ddr-2*

mutant animals is not caused by a modest reduction in hemicentin levels. Interestingly, there

was an ~40% increase in type IV collagen levels at the utse-seam connection after RNAi

mediated reduction of *ddr-2*, indicating a possible feedback mechanism between DDR-2

signaling and collagen assembly (***Figure 5—figure supplement 1E***). In sum, these results

indicate that knockdown of *ddr-2* does not decrease the functional levels of key matrix

components at the site of utse-seam BM-BM connection, suggesting that DDR-2 does not

promote utse-seam attachment by regulating the assembly of the BM-BM linking matrix.

**DDR-2 promotes integrin function to mediate utse-seam tissue attachment**

Studies have indicated that vertebrate DDRs are not strong adhesive receptors (Xu, Bihan et al.

2012). Consistent with this, we detected weak DDR-2::mNG fluorescence signals at the cell

membranes of the utse and seam. However, we observed strong DDR-2 signals in a vesicular

pattern in both tissues. We thus assessed several molecular activities regulated by DDR

signaling that might mediate utse-seam attachment. Vertebrate DDRs promote matrix

metalloproteinase (MMP) expression and localization (Leitinger 2014). The *C. elegans* genome

harbors six MMP genes, named zinc metalloproteinase 1-6 (*zmp-1-6*) (Altincicek, Fischer et al.

2010). We examined four available reporters of ZMP localization (ZMP-1::GFP, ZMP-2::GFP,

ZMP-3::GFP, and ZMP-4::GFP) (Kelley, Chi et al. 2019). Only ZMP-4 was detected at the utse-

seam connection and its localization was not altered by knockdown of *ddr-2* (***Figure 5—figure***

***supplement 1F***), suggesting that DDR-2 does not promote utse-seam linkage through

regulation of MMPs.

DDRs also modulate integrin activity (Leitinger 2014). Cell culture studies have shown that

vertebrate DDRs can enhance integrin adhesion and integrin cell surface levels in multiple cell

types (Xu, Bihan et al. 2012, Staudinger, Spano et al. 2013, Bayer, Grither et al. 2019). *C.*

*elegans* harbor only two integrin receptors, made up of either the α subunit INA-1 or PAT-2

dimerized to the sole β subunit PAT-3 (Clay and Sherwood 2015). We previously found that

depletion of INA-1 results in a Rup phenotype, but the specific effect on utse-seam BM-BM

tissue attachment was not determined (Morrissey, Keeley et al. 2014). Thus, we next asked

whether DDR-2 promotes utse-seam attachment through regulation of integrin. First, we

analyzed the localization and levels of endogenously tagged INA-1 (αINA-1::mNG) and PAT-2

(αPAT-2::2xmNG) (Jayadev, Chi et al. 2019) and found that both integrin α subunits were

expressed in the utse and seam tissues and localized to the utse-seam connection region from

the mid-L4 stage through young adulthood (**Figure 5A and B; Figure 5—figure supplement**

**2A and B**). Their levels mirrored that of DDR-2, with a peak at the late L4 stage, followed by a

drop off in the young adult stage (**Figure 5A and B)**. However, INA-1 levels were at least sixfold

greater than PAT-2 (note, each PAT-2 is linked to two mNG molecules, see **Methods**),

suggesting it may have a dominant role in adhesion (**Figure 5A and B**). Loss of *ddr-2* resulted

in a significant reduction in levels of both INA-1 and PAT-2 at the utse-seam BM-BM attachment

through the L4 larval stage (**Figure 5B**). Taken together these data suggest DDR-2 could

function to promote integrin adhesion within each tissue at the utse-seam connection.

We focused on a possible role for DDR-2 in regulating INA-1 at the utse-seam linkage, as INA-1

was present at higher levels compared to PAT-2 (**Figure 5A and B**) and was more enriched at

cell surfaces (**Figure 5—figure supplement 2C**). Furthermore, PAT-2 anchors muscle

attachments (Moerman and Williams 2006, Gieseler, Qadota et al. 2017) (**Figure 5A**) and its

loss leads to paralysis, which eliminates mechanical stress on the tissue connection and

prevents defects in the utse-seam attachment (Gianakas, Keeley et al. 2023). To determine if

DDR-2 promotes integrin activity, we reduced INA-1 levels at the late L4 stage to 30% by RNAi

and assessed whether loss of *ddr-2* enhanced the Rup and utse-seam splitting defects (**Figure**

**5—figure supplement 2D and E)**. Loss of *ddr-2* strongly enhanced the uterine prolapse defect

caused by RNAi-mediated knockdown of *ina-1* (**Figure 5C; Figure 5—figure supplement 2D**).

In addition, loss of *ddr-2* enhanced the frequency of utse-seam detachments at the late L4 stage

upon depletion of INA-1 (**Figure 5D**). We note that utse-seam detachments were not detected

upon RNAi against *ina-1* alone, likely due to incomplete knockdown at this timepoint (**Figure**

**5—figure supplement 2D and E**). Together, these observations are consistent with a possible

role for DDR-2 in regulating integrin function to promote utse-seam connection.

To determine where integrin functions to promote utse-seam attachment, we used a previously

characterized dominant negative strategy, where expression of the β integrin PAT-3 lacking the

extracellular domain (referred to as *DN integrin*; **Figure 5E**) inhibits endogenous integrin

function (Martin-Bermudo and Brown 1999, Lee, Cram et al. 2001, Hagedorn, Yashiro et al.

2009). We used the utse-specific $zmp\text{-}1^{mK50\text{-}51}$ promoter (**Figure 5—figure supplement 2F**)

(Hagedorn, Yashiro et al. 2009) and the seam-specific *wrt-2* promoter to disrupt integrin activity

in the respective tissues through the time of utse-seam tissue linkage (late L3 to young adult

stages, **Figure 1D**). Expression of *DN integrin* in either the utse or the seam cells resulted in

utse-seam detachment at the late L4 (**Figure 5F and G**). Taken together, these findings indicate

that DDR-2 may promote utse-seam BM-BM adhesion in part by regulating integrin levels and

function.

**DDR-2 promotes vinculin localization and integrin stability at the utse-seam linkage**

As DDR-2 promotes integrin localization to the utse-seam BM-BM linkage and integrin activity

promotes utse-seam attachment, we further explored how DDR-2 may regulate integrin. Talin

and vinculin are core components of the integrin adhesion complex that support integrin

activation (Chastney, Conway et al. 2021). We thus examined endogenously tagged talin

(GFP::TLN-1) (Walser, Umbricht et al. 2017) and vinculin (DEB-1::mNG) localization and

detected both proteins at the utse-seam attachment site. Levels of each protein ramped up to

the late L4 and then decreased by the young adult stage, mirroring DDR-2 (**Figure 6A and B**).

However, loss of *ddr-2* only decreased DEB-1 levels (**Figure 6B**). Confirming a role in utse-

seam connection, depletion of DEB-1 by RNAi resulted in utse-seam detachment at the late L4

stage (*Figure 6C; Figure 6—figure supplement 1*). These data show DDR-2 also regulates

vinculin levels at the utse-seam linkage and that vinculin is required for utse-seam attachment.

Integrin adhesions form stable complexes that anchor cells to ECM (Chastney, Conway et al.

2021). To determine whether DDR-2 regulates integrin complex stability at the site of utse-seam

BM-BM tissue connection, we analyzed the dynamics of INA-1 by performing Fluorescence

Recovery After Photobleaching (FRAP) experiments. We focused on INA-1::mNG at the late L4

stage when DDR-2 peaks in levels. Strikingly, loss of *ddr-2* nearly doubled the rate of recovery

of INA-1::mNG fluorescence signal in the bleached region 10 minutes post-bleaching as

compared to wild-type animals (*Figure 6D*). These observations indicate that DDR-2 stabilizes

the INA-1 integrin adhesion complex at the utse-seam adhesion site.

**Ras acts in the DDR-2 pathway and promotes integrin adhesion at the utse-seam linkage**

Vertebrate DDR2 activates multiple signaling networks, including the PI3K/Akt and Ras/Raf/Erk

cascades (Payne and Huang 2014, Chen, Kong et al. 2021). Notably, Ras regulates integrin

activity in several mammalian cell types (Zhang, Vuori et al. 1996, Conklin, Ada-Nguema et al.

2010, Sandri, Caccavari et al. 2012, Lilja, Zacharchenko et al. 2017). We thus hypothesized that

DDR-2 could promote integrin stabilization through Ras signaling. We first strongly depleted

LET-60 (see *Methods*) the *C. elegans* Ras ortholog, and observed penetrant utse-seam

attachment defects, which were not worsened in *ddr-2* null mutant animals (*Figure 7A*). This

suggests that DDR-2 and LET-60/Ras function in the same pathway to mediate the utse-seam

connection. Next, we analyzed INA-1::mNG levels and stability after LET-60 reduction. RNAi

against *let-60* resulted in significantly lower INA-1 levels at the utse-seam BM-BM connection

site (*Figure 7B*, pre-bleach panels and *Figure 7—figure supplement 1*). Further, like loss of

*ddr-2*, FRAP analysis indicated that depletion of LET-60/Ras reduced the stability of INA-1 at

the utse-seam linkage site (*i.e.,* there was more rapid recovery of INA-1 after photobleaching

(**Figure 7B**). We also generated animals harboring a gain-of-function mutation in Ras (LET-

60$^{G13E}$, *let-60(qy203)*) (Singh and Han 1995). Strikingly, utse-seam detachments were observed

in *let-60* gain-of-function animals (**Figure 7C**) and FRAP experiments revealed that INA-1 had

an approximately two-fold greater stability at the utse-seam BM-BM connection site (**Figure**

**7D**). Collectively, these findings suggest that Ras acts downstream of DDR-2 to set the

appropriate levels and stability of INA-1 mediated adhesion during the formation the utse-seam

BM-BM tissue connection.

**Discussion**

To form organs such as the brain, kidney, and lung, separate tissues connect through adjoining

basement membrane (BM) to mediate complex functions, such as molecular barrier, blood

filtration, and gas exchange, respectively (Keeley and Sherwood 2019). These linkages resist

the independent movement of each tissue and must be strong and balanced. How cells

coordinate and strengthen adhesion at BM-BM linkage sites, however, is unknown. Using the *C.*

*elegans* utse-seam BM-BM tissue connection, we show that type IV collagen, which is the

molecular glue that fastens the BM-BM linkage, also activates the *C. elegans* collagen receptor

DDR-2 in the utse and seam tissues. DDR-2 activity synchronizes and bolsters an integrin

mediated adhesion to stabilize the utse-seam BM-BM tissue connection during its formation

(**Figure 8**).

The linkage between the multi-nucleated utse cell and epithelial seam cells allows the utse to

maintain the uterus within the animal during powerful egg-laying muscle contractions (Vogel and

Hedgecock 2001). We found that contact between the BM encased utse and seam cells occurs

at the time the utse cell is formed during the early L4 larval stage. This association happens just

prior to assembly of type IV collagen at the BM-BM linkage, a matrix component that allows the

linkage to resist the mechanical forces of egg-laying (Gianakas, Keeley et al. 2023). Even

before egg-laying, however, the utse-seam linkage resists the forces of body wall and uterine

muscle contractions, as defects in matrix assembly linking the BMs results in utse-seam

detachment during the mid L4 stage (Gianakas, Keeley et al. 2023). Through analysis of genetic

mutations in the *C. elegans* receptor tyrosine kinase (RTK) DDR-2, an ortholog to the two

vertebrate DDR receptors (DDR1 & DDR2) (Unsoeld, Park et al. 2013), we discovered that loss

of *ddr-2* results in utse-seam detachment beginning at the mid L4 stage. Analysis of the BM-BM

connecting matrix that links the BMs, which is composed of hemicentin, fibulin-1, and type IV

collagen, revealed it was still assembled at functional levels. This suggests that the utse and

seam detachment did not arise from a reduction in deposition of BM-BM linking matrix, but

rather a defect in tissue adhesion to the BM-BM linkage site. Our results also indicated that

DDR-2 functions in both the utse and seam to promote adhesion, as tissue specific expression

of a dominant negative DDR-2 in either tissue resulted in utse-seam detachment. Loss of DDR1

in mice disrupts the BM-BM connection between the podocytes and endothelial cells in the

kidney glomeruli and leads to defects in blood filtration (Gross, Beirowski et al. 2004),

suggesting a possible shared function of DDR receptors in mediating proper linkage at sites of

tissue connection.

In vitro studies using DDR-specific collagen binding peptides have revealed that DDRs only

modestly contribute directly to cell adhesion (Xu, Bihan et al. 2012). Instead, DDRs are thought

to initiate downstream signaling cascades that promote cell adhesion (Borza and Pozzi 2014).

Consistent with this notion, we observed that DDR-2::mNG was not detectable at utse and

seam cell surfaces when the *ddr-2* mutant attachment defect occurred between the mid to late

L4 stages. Instead, DDR-2 was present in early, late, and recycling endosomes of both utse and

seam cells. Work in a mouse osteoblast and a human kidney cell line has shown that DDR1

shifts from localization at the cell membrane to aggregation and internalization into early

endosomes within minutes of exposure to collagen (Mihai, Chotani et al. 2009). Examination of

the timing of DDR activation has suggested that full tyrosine receptor activation occurs within

the endocytic vesicles (Mihai, Chotani et al. 2009, Fu, Valiathan et al. 2013)—a signaling hub of

other RTKs (Murphy, Padilla et al. 2009, Villasenor, Kalaidzidis et al. 2016). We also found that

internalization of DDR-2 at the utse-seam connection correlated with the assembly of type IV

collagen at the BM-BM linkage and was dependent on type IV collagen deposition. As DDR-2 is

activated by ligand-induced clustering of the receptor (Juskaite, Corcoran et al. 2017, Corcoran,

Juskaite et al. 2019), it suggests that the BM-BM linking type IV collagen network, which is

specifically assembled at highly levels at the BM-BM connection, clusters and activates DDR-2

in the utse and seam cells to coordinate adhesion at the BM-BM linkage.

Vertebrate DDRs can promote integrin mediated cell-matrix adhesion in a variety of cell types.

For example, overexpression of DDR1 enhances integrin matrix adhesion in human embryonic

kidney cells and in human and mouse fibroblast (Xu, Bihan et al. 2012, Staudinger, Spano et al.

2013, Borza, Bolas et al. 2022), and high levels of DDR2 enhances integrin activation in cancer

associated fibroblasts in mice (Bayer, Grither et al. 2019). Several mechanisms for DDR-

mediated enhancement of integrin adhesion have been proposed, including increased integrin

surface levels and Rap1-mediated talin recruitment (Staudinger, Spano et al. 2013, Bayer,

Grither et al. 2019). Loss of *C. elegans rap-1* has not been reported to have a Rup/uterine

prolapse phenotype (Frische, Pellis-van Berkel et al. 2007) and we found that DDR-2 did not

promote talin recruitment to the utse-seam connection. DDR-2 did, however, increase the levels

of the two *C. elegans* α integrins—INA-1 and PAT-2—at the utse-seam linkage (Clay and

Sherwood 2015), as well as the integrin activating protein DEB-1/vinculin (Bays and DeMali

2017). Supporting a functional role for DDR-2 in promoting INA-1 mediated utse-seam

attachment, *ddr-2* null mutants strongly enhanced the utse-seam detachment and uterine

prolapse defect (Rup phenotype) of animals with reduced INA-1 function. Further, like DDR-2,

we found that INA-1 was expressed in the utse and seam cells and through tissue-specific

expression of a dominant negative integrin, we determined that integrin activity is required in

both tissues to promote adhesion. To understand how DDR-2 affects the strength of INA-1

mediated utse-seam adhesion, we performed photobleaching experiments on endogenously

tagged INA-1::mNG, and discovered that DDR-2 stabilizes INA-1-mediated adhesion. Through

genetic interaction studies, RNAi, and expression of constitutively active Ras (*C. elegans* LET-

60), we provide evidence that DDR-2 signals through Ras to promote INA-1 stability. Ras

GTPases are a key downstream mediator of DDR signaling (Chen, Kong et al. 2021). Ras has

been implicated in activating integrins via a poorly understood mechanism that may involve

integrin receptor trafficking (Zhang, Vuori et al. 1996, Conklin, Ada-Nguema et al. 2010, Sandri,

Caccavari et al. 2012). Interestingly, expression of the constitutively active form of Ras further

increased INA-1 stability at the utse-seam linkage and led to a highly penetrant utse-seam

detachment defect. Notably, our photobleaching experiments showed that there was a wide

range of integrin stability at the utse-seam attachment site in wild-type animals, likely reflecting

low to high adhesion strength. However, Ras activation narrowed integrin to within a high

stability range (i.e. high adhesion strength), while *ddr-2* loss as well as Ras reduction shifted

integrin towards a narrow, low stability range (low adhesion strength). Together, our

observations suggest that DDR-2 signaling through Ras facilitates an increased range of

integrin adhesion at the utse-seam connection, which is required to maintain the utse-seam

linkage.

The utse-seam adhesion at the BM-BM tissue linkage is built in the L4 larval stage and then

maintained in the adult to support egg-laying (Gianakas, Keeley et al. 2023). Our evidence

indicates that DDR-2 activity is highest and perhaps only required between the mid and late L4

larval stages (~four-hour period), as the utse-seam detachment defect peaked by the mid-to-

late L4 stage and did not worsen at the late L4 and young adult stages. This timing aligns with a

spike in DDR-2 levels in the utse and seam cells and is also when DDR-2 most strongly

promoted INA-1 integrin and vinculin levels at the utse-seam linkage. It is possible that DDR-2 is

specifically required between the mid and late L4 stages to facilitate a wide range of integrin

adhesion. This could adjust the cell-matrix adhesion strength as the BM-BM connecting matrix

assembles, strengthens, and changes composition during this time window, shifting from a

hemicentin and fibulin-rich to a type IV collagen-dominated matrix (Gianakas, Keeley et al.

2023). Alternatively, a range of integrin adhesion strength may be required to maintain the utse-

seam linkage, while allowing the utse and seam to undergo morphogenetic changes to form the

tongue (utse) and groove (seam) morphology that secures the tissue linkage. Collectively, our

findings support the idea that although the ligands for DDRs are stable collagen molecules,

DDRs can function as dynamic sensors of the external environment, allowing cells and tissues

to coordinate their activities with changes to the extracellular surroundings.

## Materials and Methods

### Key resources table

| Reagent type (species) or resource | Designation | Source or reference | Identifiers | Additional information |
|---|---|---|---|---|
| Strain, strain background (*C. elegans*) | NK2617 | (Gianakas, Keeley et al. 2023) | | *qyIs23 [cdh-3p::mCh::PH]* II; *lqIs80 [scmp::GFP::CAAX]* IV |
| Strain, strain background (*C. elegans*) | N2 | Caenorhabditis Genetics Center (CGC) | | Wild type (ancestral) |
| Strain, strain background (*C. elegans*) | RB970 | | | *ddr-1(ok874)* X |
| Strain, strain background (*C. elegans*) | VH1387 | (Unsoeld, Park et al. 2013) | | *ddr-1(tm382)* X |
| Strain, strain background (*C. elegans*) | RB788 | | | *ddr-2(ok574)* X |
| Strain, strain background (*C. elegans*) | VH1383 | (Unsoeld, Park et al. 2013) | | *ddr-2(tm797)* X |

| Strain, strain background (*C. elegans*) | NK2511 | This study | | *ddr-2(qy64)* X |
|---|---|---|---|---|
| Strain, strain background (*C. elegans*) | NK2655 | This study | | *qyIs23 [cdh-3p::mCh::PH]* II*; lqIs80 [scmp::GFP::CAAX]* IV; *ddr-2(qy64)* X |
| Strain, strain background (*C. elegans*) | NK2640 | This study | | *qy91 [nas-22p::2xmKate2::PH]* II; *lqIs80 [scmp::GFP::CAAX]* IV |
| Strain, strain background (*C. elegans*) | NK2682 | This study | | *qy91 [nas-22p::2xmKate2::PH]* II; *lqIs80 [scmp::GFP::CAAX]* IV; *ddr-2(qy64)* X |
| Strain, strain background (*C. elegans*) | | This study | | *qy44 [ddr-2::mNG]* X; *qyEx605 [scmp::2xmKate2:PH]* |

| Strain, strain background (*C. elegans*) | NK2620 | This study | | *qyIs23 [cdh-3p::mCh::PH]* II*; qy44 [ddr-2::mNG]* X |
| --- | --- | --- | --- | --- |
| Strain, strain background (*C. elegans*) | NK2763 | This study | | *qyIs256 [cdh-3p::mCherry::rab-5] ?; qy44 [ddr-2::mNG]* X |
| Strain, strain background (*C. elegans*) | NK2766 | This study | | *qyIs252 [cdh-3p::mCherry::rab-7] ?; qy44 [ddr-2::mNG]* X |
| Strain, strain background (*C. elegans*) | NK2769 | This study | | *qyIs205 [cdh-3p::mCherry::rab-11] ?; qy44 [ddr-2::mNG]* X |
| Strain, strain background (*C. elegans*) | NK2901 | This study | | *qy188 [scmp::mKate2::rab-5]* I*; qy44 [ddr-2::mNG]* X |
| Strain, strain background (*C. elegans*) | NK2926 | This study | | *qy91 [nas-22p::2xmKate2::PH]* II; *lqIs80 [scmp::GFP::CAAX]* IV; *ddr-2(ok574)* X |

| Strain, strain background (*C. elegans*) | NK2705 | (Jayadev, Morais et al. 2022) | | *qy118 [col-99::mNG (internal tag)]* IV |
|---|---|---|---|---|
| Strain, strain background (*C. elegans*) | NK2322 | (Keeley, Hastie et al. 2020) | | *qy22 [cle-1::mNG]* I |
| Strain, strain background (*C. elegans*) | NK2326 | (Keeley, Hastie et al. 2020) | | *qy24 [emb-9::mNG (internal tag)]* III |
| Strain, strain background (*C. elegans*) | NK2830 | This study | | *qy166 [cdh-3p::dominant negative ddr-2::mNG]* I; *qy91 [nas-22p::2xmKate2::PH]* II; *heIs63 [wrt-2p::GFP::PH + wrt-2p::GFP::H2B + lin-48p:mCherry]* V |
| Strain, strain background (*C. elegans*) | NK2848 | This study | | *qy175 [scmp::dominant negative ddr-2::mKate2]* I; *qy91 [nas-22p::2xmKate2::PH]* |

| | | | | II; *heIs63 [wrt-2p::GFP::PH + wrt-2p::GFP::H2B + lin-48p:mCherry]* V |
|---|---|---|---|---|
| Strain, strain background (*C. elegans*) | NK2585 | (Jayadev, Morais et al. 2022) | | *qy83 [emb-9::mRuby2 (internal tag)]* III |
| Strain, strain background (*C. elegans*) | NK2422 | (Keeley, Hastie et al. 2020) | | *qy33 [him-4::mNG]* X |
| Strain, strain background (*C. elegans*) | NK2324 | (Jayadev, Chi et al. 2019) | | *qy23 [ina-1::mNG]* III |
| Strain, strain background (*C. elegans*) | NK2479 | (Jayadev, Chi et al. 2019) | | *qy49 [pat-2::2xmNG]* III |
| Strain, strain background (*C. elegans*) | NK2825 | This study | | *qy23 [ina-1::mNG]* III; *ddr-2(qy64)* X |
| Strain, strain background (*C. elegans*) | NK2858 | This study | | *qy49 [pat-2::2xmNG]* III; *ddr-2(qy64)* X |
| Strain, strain background (*C. elegans*) | NK2804 | This study | | *qyIs23 [cdh-3p::mCh::PH]* II; *heIs63 [wrt-* |

| Strain, strain background (*C. elegans*) | | | | 2p::GFP::PH + wrt-2p::GFP::H2B + lin-48p:mCherry] V |
|---|---|---|---|---|
| Strain, strain background (*C. elegans*) | NK2824 | This study | | *qyIs23 [cdh-3p::mCh::PH]* II; *qyIs15 [zmp-1$^{mk50-51}$p::dominant negative integrin b-pat-3]* IV;*heIs63 [wrt-2p::GFP::PH + wrt-2p::GFP::H2B + lin-48p:mCherry]* V |
| Strain, strain background (*C. elegans*) | NK2934 | This study | | *qy91 [nas-22p::2xmKate2::PH]* II; *unc-119(ed4)* III; *lqIs80 [scmp::GFP::CAAX]* IV; *qyEx604 [wrt-2p::dominant negative integrin b-pat-3 + unc-119(+)]* |
| Strain, strain background (*C. elegans*) | NK2579 | (Keeley, Hastie et al. 2020) | | *qy62 [mNG::fbl-1] IV* |

| | | | | |
|---|---|---|---|---|
| Strain, strain background (*C. elegans*) | NK932 | (Kelley, Chi et al. 2019) | | *qyIs190 [zmp-4p::zmp-4::GFP]* |
| Strain, strain background (*C. elegans*) | NK268 | (Hagedorn, Yashiro et al. 2009) | | *qyIs17 [zmp-1$^{mk50-51}$p mCherry]* |
| Strain, strain background (*C. elegans*) | AH3437 | (Walser, Umbricht et al. 2017) | | *zh117 [GFP::tln-1]* I |
| Strain, strain background (*C. elegans*) | NK2478 | This study | | *qy48 [deb-1::mNG]* IV |
| Strain, strain background (*C. elegans*) | NK2854 | This study | | *zh117 [GFP::tln-1]* I; *ddr-2(qy64)* X |
| Strain, strain background (*C. elegans*) | NK2860 | This study | | *qy48 [deb-1::mNG]* IV; *ddr-2(qy64)* X |
| Strain, strain background (*C. elegans*) | NK2944 | This study | | *qy91 [nas-22p::2xmKate2::PH*] II; *lqIs80 [scmp::GFP::CAAX]* IV; *let-60(qy203)* IV |

| Strain, strain background (*C. elegans*) | NK2957 | This study | | *qy23 [ina-1::mNG]* III; *let-60(qy203)* IV |
|---|---|---|---|---|
| Strain, strain background (*E. coli*) | *emb-9* RNAi | (Kamath and Ahringer 2003) | | Clone from Ahringer library (L4440 vector backbone) |
| Strain, strain background (*E. coli*) | *col-99* RNAi | (Rual, Ceron et al. 2004) | | Clone from Vidal library (L4440 vector backbone) |
| Strain, strain background (*E. coli*) | *cle-1* RNAi | (Rual, Ceron et al. 2004) | | Clone from Vidal library (L4440 vector backbone) |
| Strain, strain background (*E. coli*) | *ina-1* RNAi | (Rual, Ceron et al. 2004) | | Clone from Vidal library (L4440 vector backbone) |
| Strain, strain background (*E. coli*) | *let-60* RNAi | (Rual, Ceron et al. 2004) | | Clone from Vidal library (L4440 vector backbone) |
| Strain, strain background (*E. coli*) | *deb-1* RNAi | (Rual, Ceron et al. 2004) | | Clone from Vidal library (L4440 vector backbone) |
| Strain, strain background (*E. coli*) | *him-4* RNAi | (Gianakas, Keeley et al. 2023) | | T444t vector backbone |

| Strain, strain background (*E. coli*) | *ddr-2* RNAi | This study | | T444t vector backbone |
|---|---|---|---|---|
| Sequence-based reagent | *ddr-2* knockout (*ddr-2(qy64)*) sgRNA 1 | This study | | ATCCTGACATAGA TGAGCGT |
| Sequence-based reagent | *ddr-2* knockout (*ddr-2(qy64)*) sgRNA 2 | This study | | GTCATTGGTGCAC ACTTCTC |
| Sequence-based reagent | *ddr-2* knockout (*ddr-2(qy64)*) sgRNA 3 | This study | | AAGTGTGCACCAA TGACTGG |
| Sequence-based reagent | *deb-1::mNG* (*deb-1(qy48)*) sgRNA 1 | This study | | AGTTGGACCACAT TGGCTTT |
| Sequence-based reagent | *deb-1::mNG* (*deb-1(qy48)*) sgRNA 2 | This study | | ATTTAGAAGTTGG ACCACAT |
| Sequence-based reagent | *let-60* gain-of-function (*let-60(qy203)*) sgRNA 1 | This study | | CTTGTGGTAGTTG GAGATGG |
| Sequence-based reagent | Primer *ddr-2(qy64)* homology arm forward | This study | | TTTTCAGAGTCTC CGACGCTCATCTA |

| | | | | |
|---|---|---|---|---|
| Sequence-based reagent | Primer *ddr-2(qy64)* homology arm reverse | This study | | TAAATATTATTCTG AGAATATA |
| Sequence-based reagent | Primer *ddr-2(qy64)* genotyping forward | This study | | TGGTAATTGATGA GAGGGTG |
| Sequence-based reagent | Primer *ddr-2(qy64)* genotyping reverse | This study | | TGTCGTTTCGACA CCGGCAA 1.8 kb band |
| | Primer *let-60(qy203)* homology arm forward | This study | | ATGACGGAGTACA AGCTTGTGGTAGT TGGAGATGGAGA AGT |
| | Primer *let-60(qy203)* homology arm reverse | This study | | TACCCTTTTCTGA AAAAAGACGC |
| Sequence-based reagent | Primer *nas-22* promoter forward | This study | | TCAAAGCGTCAAG CTTTACG |

| Sequence-based reagent | Primer *nas-22* promoter reverse | This study | | TTTTCTGTCTACAA GGACTGTGC |
|---|---|---|---|---|
| Sequence-based reagent | Primer *scm* promoter forward | (Chapman, Li et al. 2008) | | AACCTCCACCAGA TGGTTGGCG |
| Sequence-based reagent | Primer *scm* promoter reverse | (Chapman, Li et al. 2008) | | CCCGGGGATCCG TCCACTCT |
| Sequence-based reagent | Primer *cdh-3* promoter forward | This study | | CTAGAGCATGATG TCCTTAC |
| Sequence-based reagent | Primer *cdh-3* promoter reverse | This study | | CAAAACGGACCG ACCGTCCC |
| Sequence-based reagent | Primer *dominant negative ddr-2* forward | This study | | ATGAAGTTGCTGC TGTATCT |
| Sequence-based reagent | Primer *dominant negative ddr-2* reverse | This study | | TCTGCTCACGCAA ATCAACT |

| Sequence-based reagent | Primer *wrt-2* promoter forward | This study | | TCAGAACTCTAAT ACTTACT |
|---|---|---|---|---|
| Sequence-based reagent | Primer *wrt-2* promoter reverse | This study | | CCGAGAAACAATT GGCAGGT |
| Sequence-based reagent | Primer *dominant negative integrin b pat-3* forward | (Hagedorn, Yashiro et al. 2009) | | TCTAGAGGATCCC GGGGAT |
| Sequence-based reagent | Primer *dominant negative integrin b pat-3* reverse | (Hagedorn, Yashiro et al. 2009) | | ATTTAGTTGGCTT TTCCAGC |
| Sequence-based reagent | Primer *ddr-2* RNAi forward | This study | | ATGAAGTTGCTGC TGTATCT |
| Sequence-based reagent | Primer *ddr-2* RNAi reverse | This study | | ATGAATATGAGGA GAAGTGTGC |
| Recombinant DNA reagent | Plasmid: pCFJ352 | Addgene | RRID: Addgene_3 0539 | sgRNA targeting MosSCI insertion |

| | | | | site on Chr I, from Erik Jorgensen |
|---|---|---|---|---|
| Recombinant DNA reagent | Plasmid: pDD122 | (Dickinson, Ward et al. 2013) | RRID: Addgene_4 7550 | sgRNA targeting MosSCI insertion site on Chr II |
| | Plasmid: pAP087 | (Pani and Goldstein 2018) | | starter SEC repair template plasmid for single copy insertion at the ttTi5605 site on chromosome II |
| Recombinant DNA reagent | Plasmid: pAP088 | (Pani and Goldstein 2018) | | starter SEC repair template plasmid for single copy insertion at the ttTi4348 site on chromosome I |
| Chemical compound, drug | Ampicillin | Sigma-Aldrich | #A0166 | |
| Chemical compound, drug | Isopropyl β-D-1-thiogalactopyra noside | Sigma-Aldrich | #I6758 | |
| Chemical compound, drug | Hygromycin B | Sigma-Aldrich | #H3274 | |
| Chemical compound, drug | Sodium azide | Sigma-Aldrich | #S2002 | |

| Chemical compound, drug | Levamisole | Sigma-Aldrich | #L9756 | |
|---|---|---|---|---|
| Software, algorithm | µManager v.1.4.23 | | | |
| Software, algorithm | Zen Black | | | |
| Software, algorithm | Fiji/ImageJ | | | |
| Software, algorithm | GraphPad Prism v9 | | | |

***C. elegans* culture and strains**

*C. elegans* strains used in this study are listed in the ***Key resources table***. Worms were reared

on nematode growth medium plates seeded with OP50 *Escherichia coli* at 16°C, 18°C, or 20°C

according to standard procedures (Stiernagle 2006). We used *C. elegans* vulval development to

accurately stage animals through the morphogenesis of the utse-seam connection (late L3 to

young adult stage) (Mok, Sternberg et al. 2015).

**Generation of genome-edited strains**

To generate the genome-edited mNeonGreen (mNG) knock-in allele for *deb-1* (*qy48*), we used

CRISPR/Cas9 genome editing with a self-excising cassette (SEC) for drug selection

(hygromycin treatment) as described previously (Dickinson, Ward et al. 2013, Keeley, Hastie et

al. 2020). An 18 amino acid flexible linker attached to the mNG fluorophore was inserted in

frame and directly upstream of the stop codon. sgRNA sequences directing Cas9 cleavage near

the C-terminus are provided in the ***Key resources table***.

The *ddr-2* knockout allele *ddr-2(qy64)*, an 8615 bp deletion corresponding to the coding

sequence of DDR-2, was also generated with the SEC CRISPR method detailed above with one

modification—we generated a new starter repair plasmid lacking the fluorescent tag. Primers

used to amplify homology arms, sgRNA sequences to direct Cas9 cleavage, and genotyping

primers used to verify genome-edited knockout animals are provided in the **Key resources**

**table**.

For the LET-60$^{G13E}$ gain-of function mutation, *let-60(qy203)*, we used the same strategy as the

*ddr-2* knockout allele. The point mutation was incorporated into the primer for homology arm

amplification. Note that this mutation is identical to the *let-60(n1046)* allele.

**Generation of transgenic strains**

For the utse marker *qy91 [nas-22p::2xmKate2::pH]*, we generated a single-copy transgene

inserted into the ttTi5605 transposon insertion on chromosome II (Frokjaer-Jensen, Davis et al.

2012) using the SEC CRISPR method detailed above. A 2 kb *nas-22* promoter, two tandem

mKate2 fluorophores attached to a PH domain, and an *unc-54* 3' UTR fragment were ligated in

order via Gibson assembly into the pAP087 starter repair plasmid containing homology arms for

the ttTi5605 site. The sgRNA directing Cas9 cleavage near this region is contained in the

pDD122 vector.

To express mKate2-tagged RAB-5 in the seam, we generated a single-copy transgene inserted

into the ttTi4348 transposon insertion site on chromosome I (Frokjaer-Jensen, Davis et al. 2012)

using SEC CRISPR. Briefly, the scm promoter, mKate2 fluorophore, rab-5 genomic sequence,

and *unc-54* 3' UTR were Gibson assembled in order into the pAP088 starter repair plasmid

containing homology arms for the ttTi4348 site. The sgRNA directing Cas9 cleavage near this

region is contained in the pCFJ352 vector.

For tissue-specific expression of dominant negative (DN) DDR-2 in the utse or seam, we also

generated single-copy transgenes inserted into the ttTi4348 site. For utse expression, the

following fragments were Gibson assembled in order into pAP088: a 1.5kb *cdh-3* promoter

fragment, *DN ddr-2*, mNG fluorophore, and *unc-54* 3' UTR. Expression in transgenic animals

was verified by assessing mNG fluorescence. For seam expression, the following fragments

were Gibson assembled in order into pAP088: the *scm* promoter fragment, *DN ddr-2*, and

mKate2 fluorophore. Expression in transgenic animals was verified by assessing mKate2

fluorescence.

The seam cell marker *qyEx605 [scmp::2xmKate2::PH]* was built by Gibson assembling a 1.3 kb

*scm* promoter fragment, a fragment with two tandem mKate2 fluorophores attached to the PH

domain, and *unc-54* 3' UTR fragment in order. The construct was expressed as

extrachromosomal arrays as described previously (Yochem and Herman 2003).

To build the construct for *DN integrin* expressed in the seam, we first amplified the *DN integrin*

fragment from animals harboring the *qyIs15* allele (see **Key resources table**). The *DN integrin*

fragment is composed of a *pes-10* enhancer element, the *ost-1* signal peptide, the *pat-3* β-tail

fragment, and *unc-54* 3' UTR as described previously (Lee Cram et al. 2001). The *DN integrin*

fragment was fused to a 1.3 kb *wrt-2* promoter by PCR. The construct was expressed

extrachromosomally.

**RNAi**

All RNAi constructs except for *ddr-2* RNAi were obtained from the Ahringer and Vidal RNAi

libraries (Kamath and Ahringer 2003, Rual, Ceron et al. 2004) or generated previously

(Gianakas, Keeley et al. 2023). For the *ddr-2* clone, a ~2.4 kb cDNA fragment corresponding to

the longest *ddr-2* transcript was amplified by PCR and inserted into the T444t vector as

described previously (Gianakas et al 2023). RNAi experiments were performed using the

feeding method (Timmons, Court et al. 2001) according to previously detailed protocols

(Jayadev, Chi et al. 2019, Gianakas, Keeley et al. 2023). For experiments assessing the utse-

seam connection, RNAi was initiated in synchronized L1 larvae and animals were examined at

the late L4 stage (~ 44 h treatment). For examination of uterine prolapse, RNAi was performed

from the L1 through adulthood (~ 120 h treatment). For depletion of hemicentin initiated at the

early L4 stage, synchronized L1 worm were grown on control RNAi for ~ 38 h to the early L4,

and then transferred onto *him-4* RNAi to the late L4 (~5-6 h treatment). We verified knockdown

efficiencies for all RNAi experiments except for *let-60* RNAi by including a control with the

relevant fluorescent tagged target protein, achieving between ~75 to 100% reduction. We

verified *let-60* RNAi knockdown efficiency by plate level assessment of brood size, which was

strongly reduced compared to control animals, as previously reported (Ceron, Rual et al. 2007).

**Microscopy and image processing**

All fluorescence images (except for photobleaching experiments) were acquired at 20°C on

Zeiss Axio Imager A1 microscopes controlled by the µManager software v.1.4.23 or v.2.0.1

(Edelstein, Amodaj et al. 2010), equipped with a Hamamatsu ImagEM electron multiplying

charge-coupled device camera or an Orca-Fusion scientific complementary metal oxide

semiconductor camera, Zeiss 40x and 100x Plan Apochromat (1.4 numerical aperture) oil

immersion objectives, Yokogawa CSU-10 spinning disc confocal scan heads, and 488 nm, 505

nm, and 561 nm laser lines. Worms were mounted on 5% noble agar pads containing 0.01M

sodium azide for imaging.

For photobleaching experiments, images were acquired at 20°C on a Zeiss 880 single-point

scanning confocal attached to a Zeiss Axio Observer Z1 microscope, with a Marzhauser linearly

encoded stage, a 40x Plan Neofluar (1.3 numerical aperture) oil immersion objective, and a 488

nm laser line. Worms were anesthetized by soaking in 5 mM levamisole in M9 buffer for 15 min

and then transferred to 4% noble agar pads. Coverslips were then placed and sealed on top

with valap (equal weight Vaseline, lanolin, and paraffin) and flooded with 5 mM levamisole.

Coverslip sealing kept the animals hydrated during the experimental timeframe, ensuring

accurate imaging of endogenous protein dynamics (detailed in (Kelley, Wang et al. 2017)). A

rectangle region within one arm of the BM-BM connection BLINK was photobleached using 30

iterations of simultaneous 405 nm and 488 nm excitation at 100% laser power for a total

bleaching time of 1.5 seconds. Worms were imaged prior and immediately following bleaching,

and then again 10 min later.

Brightfield images of worms were acquired at 20°C on a Zeiss Axio Zoom V16 stereo

fluorescence microscope controlled by Zen 3.2 software, equipped with an Axiocam digital

camera and a 3x objective. Worms were imaged without immobilization.

To image the utse and seam cells ventrally, we manually oriented animals in the ventral

orientation (Kelley, Wang et al. 2017) and acquired z-stacks at 0.37 μm intervals, capturing the

entire H-shaped structure of the utse and the seam cells flanking both sides of the utse at 100x

magnification. Ventral views of the utse and seam shown in figures are maximum intensity

projections of these z-stacks. 3D isosurface renderings of the utse and seam cells shown in

*Figure 2D* were generated with Imaris 7.4 software (Bitplane). We acquired lateral z-stacks at

0.37 μm intervals to fully capture the superficial half of the utse-seam attachment.

For lateral imaging of DDR-2::mNG together with the utse, seam cells, or endosome markers

expressed in the utse or seam, we acquired z-stacks at 0.37 μm intervals to capture as much of

each tissue that was in focus superficially at 40x magnification. Images shown in figures are

single slices where DDR-2 was sharply in focus within the tissue of interest. The same

parameters were used for imaging INA-1::mNG and PAT-2::mNG in the utse and seam. Images

in **Figure 5A** correspond to the z-slices where INA-1 and PAT-2 were most in focus at the BM-

BM connection region. In **Figure 5—figure supplement 2A and B**, images shown are sum

projections of the seam or utse regions.

To image DDR-2::mNG specifically at the utse-seam attachment site, we acquired lateral z-

stacks at 0.37 μm intervals capturing both the surface of the utse and the seam cells. We used

a utse cell marker as a reference to accurately capture this region (see also schematic in **Figure**

**3A**). Images shown in **Figure 4** are sum projections of these z-stacks.

For lateral imaging of respective tagged proteins at the BM-BM connection region, we acquired

single slices (40x magnification) at the middle focal plane where the BM-BM connection signal

was sharply in focus. All images were process in Fiji 2.0 (Schindelin, Arganda-Carreras et al.

2012).

**Image analysis and quantification**

To assess utse-seam detachment, we used maximum intensity projections of the utse and

seam viewed ventrally as detailed above. We defined utse-seam detachment as any gap in the

utse-seam attachment of at least 4 square pixels in area. We preferred the ventral view over 3D

rendering for this metric as gaps were variable in size and location (typically occurring in one or

two arms of the utse on opposite sides). The ventral projections allowed us to reliably examine

all four arms of the utse and whether they were in contact with the seam from animal to animal.

The visual representation of a gap in a ventral maximum projection is an underestimate of the

true size of the detachment, as the tongue-in-groove association of the utse and seam can only

be seen in a 3D surface rendering of the utse and seam viewed from the side. We thus used a

binary scoring system (animals with or without utse-seam detachments) instead of quantifying

the size of the detached region. We note that ventral imaging of the utse and seam sometimes

results in small utse-seam detachments as we manually orient the animals on the slide. To

control for this, we acquired large datasets (at least 40 animals examined) for every control

strain we used to visualize the utse-seam attachment with. We found that small detachments

related to ventral imaging occurred in ~8-12% of animals, depending on the genetic

background. All datasets shown in figures have been normalized to account for this.

All quantifications of mean fluorescence intensity were done on raw images in Fiji 2.0. We drew

~5-pixel long and 2-pixel wide linescans to obtain raw values of mean fluorescence intensity.

For measurement within the BM-BM connection region, we positioned the linescan within the

arm that was most in focus. Other regions where linescans were performed are indicated in

figures. Background intensity values were obtained with similar linescans in adjacent regions

with no visible fluorescence signal. Note that we used two tandem mNG fluorophores to

visualize PAT-2 at the utse-seam attachment site (PAT-2::2xmNG) as it was present at very low

levels. We imaged PAT-2::mNG and PAT-2::2xmNG in this region at the same exposure and

found that the latter was approximately two times brighter (PAT-2::2xmNG mean fluorescence

intensity 3941 ± 847 A.U., PAT-2::mNG 2037 ± 465 A.U., n = 10 animals examined each).

Taken together with quantifications in *Figure 5B*, INA-1 is thus likely present at six-fold higher

levels than PAT-2 at the utse-seam attachment site. For additional quantification related to

FRAP, see below.

For FRAP analysis, we measured mean fluorescence intensity within the bleached region. In

addition, we performed an equivalent linescan within the unbleached arm at the BM-BM

connection to calculate a bleach correction factor to account for general photobleaching during

image acquisition across the duration of the experiment, as previously described (Gianakas,

Keeley et al. 2023). Briefly, the background-corrected fluorescence intensity measurement on

the side of the BM-BM connection where FRAP was not performed at the 10 min post-bleach

timepoint was divided by the respective value at the pre-bleach timepoint to obtain the bleach

correction factor. Fluorescence intensity measurements at the pre-bleach and post-bleach

timepoints on the side of the BM-BM connection where FRAP was performed were then

multiplied by the respective bleach correction factor to normalize these values to the 10 min

post-bleach fluorescence intensity measurement in the same region.

To count the number of DDR-2 punctae at the site of utse-seam attachment, we used sum

projections of DDR-2::mNG capturing the surface of the utse and seam (described earlier). We

restricted measurements to the utse-seam connection corresponding to one utse arm.

Brightness and contrast was then adjusted to manually count individual punctae, as DDR-2

punctae showed a large variance in size and fluorescence intensity. Note that in the type IV

collagen depletion condition, we only counted discrete puncta and did not include the large cell

surface accumulation of DDR-2 in the analysis.

To determine co-localization of DDR-2 with markers of endocytic vesicles, we also used a

manual approach due to the large variability of the size and fluorescence intensity of punctae.

We selected a single slice where the utse or the seam was sharply in focus, and restricted

measurements to a single utse arm or a region of the seam corresponding to a single utse arm.

For each animal, the number, and positions of DDR-2 punctae were noted and compared to the

respective endosome marker (RAB-5, RAB-7, or RAB-11). If a DDR-2 puncta physically

overlapped with or was in contact with an endosome punctum, it was classified as one that co-

localized with the endosomal marker. We then calculated the percentage overlap of DDR-2

punctae with the vesicle marker for each animal (corresponding to each data point within

boxplots in *Figure 3—figure supplement 1B and C*).

**Scoring of uterine prolapse**

Synchronized L1 larvae were plated (~20 animals per plate) and after 24 h, the exact number of

worms on each plate was recorded. Plates were then visually screened for ruptured worms

(uterine prolapse) every 24 h, up to 120 h post L1 plating. Ruptured animals were scored and

removed from plates to avoid double counting and after 120 h, the percentage of animals with

uterine prolapse was calculated. At least 50 animals were screened for every experiment.

**Illustrations of utse-seam tissue connection**

For schematics in Figure 1A and Figure 8, model protein sizing was scaled according to

previously generated models (Keeley, Hastie et al. 2020). We sized hemicentin, fibulin, type IV

collagen, integrin, and DDR-2. We also approximated the span of the basement membranes of

the utse and seam cells (~50 nm each), as well as the BM-BM connecting matrix (~200 nm),

based on electron microscopy of the utse-seam attachment (Vogel and Hedgecock 2001).

**Statistical analysis**

Statistical analysis was performed in GraphPad Prism 9. Normality of datasets was assessed

using the D'Agostino-Pearson normality test. We used parametric tests for datasets that

followed a Gaussian distribution and non-parametric tests for those that did not. For

comparisons of means between two populations, we used either an unpaired two-tailed

Student's $t$ test or a Mann-Whitney $U$ test. For comparisons of means between three or more

populations, we performed either a one-way ANOVA followed by post hoc Dunnett's test or

Kruskal-Wallis $H$ test with post hoc Dunn's test. For comparisons between two categorical

variables, we used the Fisher's exact test. All graphs were prepared in GraphPad Prism. Figure

legends indicate sample sizes, statistical tests used, and $P$ values.

**Data availability**

All data generated in this study are included in the manuscript and supporting files.

**Acknowledgements**

We would like to thank C. Borza, A. Pozzi, B. Hoffman, and D. Reiner for helpful discussions, C.

Gianakas for comments on the manuscript, and M. Boxem and H. Hutter for strains. Some

strains were provided by the CGC, which is funded by NIH Office of Research Infrastructure

Programs (P40 OD010440). S.G.P. was supported by graduate research fellowship NICHD F31

HD97901. This work was supported by R35GM118049, R21OD028766, and R21OD032430 to

D.R.S.

**Additional Information**

**Funding**

| Funder | Grant reference number | Author |
|---|---|---|
| National Institutes of Health | R35GM118049 | David R. Sherwood |
| National Institutes of Health | R21OD028766 | David R. Sherwood |
| National Institutes of Health | R21OD032430 | David R. Sherwood |
| National Institute of Child Health and Human Development | F31 HD97901 | Sara G. Payne |

**Author Contributions**

Kieop Park, conceptualization, data curation, formal analysis, investigation, visualization, methodology, writing-original draft; Ranjay Jayadev, conceptualization, data curation, formal analysis, investigation, visualization, methodology, writing-original draft; Sara G. Payne, Conceptualization, data curation, formal analysis, investigation, visualization, methodology, writing-original draft, funding acquisition; Isabel W. Kenny-Ganzert, investigation, methodology; Qiuyi Chi, investigation, methodology; Daniel S. Costa, methodology; William Ramos-Lewis, investigation, methodology; Siddharthan Balachandar Thendral, visualization; David R. Sherwood, supervision, writing-original draft, funding acquisition, project administration.

**Competing Interests**

No competing interests declared for all authors

**Author ORCID**

Kieop Park  https://orcid.org/0000-0002-6380-4730

Ranjay Jayadev  https://orcid.org/0000-0003-0465-0337

Sara G. Payne  https://orcid.org/0000-0002-3572-9112

Isabel Kenny-Ganzert  https://orcid.org/0000-0001-6306-4472

Qiuyi Chi  https://orcid.org/0000-0003-1297-4163

Daniel S. Costa   https://orcid.org/0000-0003-2179-8800

Will Ramos-Lewis  https://orcid.org/0000-0002-4627-4571

Siddharthan Balachandar Thendral https://orcid.org/0000-0002-4674-7033

David R Sherwood  https://orcid.org/0000-0002-4448-6917

## References

Abrahamson, D. R. (1985). "Origin of the glomerular basement membrane visualized after in vivo labeling of laminin in newborn rat kidneys." <u>J Cell Biol</u> **100**(6): 1988-2000.

Altincicek, B., M. Fischer, M. Fischer, K. Luersen, M. Boll, U. Wenzel and A. Vilcinskas (2010). "Role of matrix metalloproteinase ZMP-2 in pathogen resistance and development in Caenorhabditis elegans." <u>Dev Comp Immunol</u> **34**(11): 1160-1169.

Bayer, S. V., W. R. Grither, A. Brenot, P. Y. Hwang, C. E. Barcus, M. Ernst, P. Pence, C. Walter, A. Pathak and G. D. Longmore (2019). "DDR2 controls breast tumor stiffness and metastasis by regulating integrin mediated mechanotransduction in CAFs." <u>Elife</u> **8**.

Bays, J. L. and K. A. DeMali (2017). "Vinculin in cell-cell and cell-matrix adhesions." <u>Cell Mol Life Sci</u> **74**(16): 2999-3009.

Bernadskaya, Y. Y., S. Brahmbhatt, S. E. Gline, W. Wang and L. Christiaen (2019). "Discoidin-domain receptor coordinates cell-matrix adhesion and collective polarity in migratory cardiopharyngeal progenitors." <u>Nat Commun</u> **10**(1): 57.

Borza, C. M., G. Bolas, X. Zhang, M. B. Browning Monroe, M. Z. Zhang, J. Meiler, M. J. Skwark, R. C. Harris, L. A. Lapierre, J. R. Goldenring, M. Hook, J. Rivera, K. L. Brown, B. Leitinger, M. J. Tyska, M. Moser, R. T. Bottcher, R. Zent and A. Pozzi (2022). "The Collagen Receptor Discoidin Domain Receptor 1b Enhances Integrin beta1-Mediated Cell Migration by Interacting With Talin and Promoting Rac1 Activation." <u>Front Cell Dev Biol</u> **10**: 836797.

Borza, C. M. and A. Pozzi (2014). "Discoidin domain receptors in disease." <u>Matrix Biol</u> **34**: 185-192.

Brown, K. L., C. F. Cummings, R. M. Vanacore and B. G. Hudson (2017). "Building collagen IV smart scaffolds on the outside of cells." <u>Protein Sci</u> **26**(11): 2151-2161.

Brown, N. H. (2011). "Extracellular matrix in development: insights from mechanisms conserved between invertebrates and vertebrates." <u>Cold Spring Harb Perspect Biol</u> **3**(12).

Carney, T. J., N. M. Feitosa, C. Sonntag, K. Slanchev, J. Kluger, D. Kiyozumi, J. M. Gebauer, J. Coffin Talbot, C. B. Kimmel, K. Sekiguchi, R. Wagener, H. Schwarz, P. W. Ingham and M. Hammerschmidt (2010). "Genetic analysis of fin development in zebrafish identifies furin and hemicentin1 as potential novel fraser syndrome disease genes." <u>PLoS Genet</u> **6**(4): e1000907.

Castro-Sanchez, L., A. Soto-Guzman, N. Navarro-Tito, R. Martinez-Orozco and E. P. Salazar (2010). "Native type IV collagen induces cell migration through a CD9 and DDR1-dependent pathway in MDA-MB-231 breast cancer cells." <u>Eur J Cell Biol</u> **89**(11): 843-852.

Ceron, J., J. F. Rual, A. Chandra, D. Dupuy, M. Vidal and S. van den Heuvel (2007). "Large-scale RNAi screens identify novel genes that interact with the C. elegans retinoblastoma pathway as well as splicing-related components with synMuv B activity." <u>BMC Dev Biol</u> **7**: 30.

Chapman, J. O., H. Li and E. A. Lundquist (2008). "The MIG-15 NIK kinase acts cell-autonomously in neuroblast polarization and migration in C. elegans." <u>Dev Biol</u> **324**(2): 245-257.

Chastney, M. R., J. R. W. Conway and J. Ivaska (2021). "Integrin adhesion complexes." <u>Curr Biol</u> **31**(10): R536-R542.

Chen, L., X. Kong, Y. Fang, S. Paunikar, X. Wang, J. A. L. Brown, E. Bourke, X. Li and J. Wang (2021). "Recent Advances in the Role of Discoidin Domain Receptor Tyrosine Kinase 1 and Discoidin Domain Receptor Tyrosine Kinase 2 in Breast and Ovarian Cancer." <u>Front Cell Dev Biol</u> **9**: 747314.

Clay, M. R. and D. R. Sherwood (2015). "Basement Membranes in the Worm: A Dynamic Scaffolding that Instructs Cellular Behaviors and Shapes Tissues." <u>Curr Top Membr</u> **76**: 337-371.

Conklin, M. W., A. Ada-Nguema, M. Parsons, K. M. Riching and P. J. Keely (2010). "R-Ras regulates beta1-integrin trafficking via effects on membrane ruffling and endocytosis." <u>BMC Cell Biol</u> **11**: 14.

Corcoran, D. S., V. Juskaite, Y. Xu, F. Gorlitz, Y. Alexandrov, C. Dunsby, P. M. W. French and B. Leitinger (2019). "DDR1 autophosphorylation is a result of aggregation into dense clusters." Sci Rep **9**(1): 17104.

Dickinson, D. J., J. D. Ward, D. J. Reiner and B. Goldstein (2013). "Engineering the Caenorhabditis elegans genome using Cas9-triggered homologous recombination." Nat Methods **10**(10): 1028-1034.

Edelstein, A., N. Amodaj, K. Hoover, R. Vale and N. Stuurman (2010). "Computer control of microscopes using microManager." Curr Protoc Mol Biol **Chapter 14**: Unit14 20.

Feitosa, N. M., J. Zhang, T. J. Carney, M. Metzger, V. Korzh, W. Bloch and M. Hammerschmidt (2012). "Hemicentin 2 and Fibulin 1 are required for epidermal-dermal junction formation and fin mesenchymal cell migration during zebrafish development." Dev Biol **369**(2): 235-248.

Fidler, A. L., S. P. Boudko, A. Rokas and B. G. Hudson (2018). "The triple helix of collagens - an ancient protein structure that enabled animal multicellularity and tissue evolution." J Cell Sci **131**(7).

Frische, E. W., W. Pellis-van Berkel, G. van Haaften, E. Cuppen, R. H. Plasterk, M. Tijsterman, J. L. Bos and F. J. Zwartkruis (2007). "RAP-1 and the RAL-1/exocyst pathway coordinate hypodermal cell organization in Caenorhabditis elegans." EMBO J **26**(24): 5083-5092.

Frokjaer-Jensen, C., M. W. Davis, M. Ailion and E. M. Jorgensen (2012). "Improved Mos1-mediated transgenesis in C. elegans." Nat Methods **9**(2): 117-118.

Fu, H. L., R. R. Valiathan, R. Arkwright, A. Sohail, C. Mihai, M. Kumarasiri, K. V. Mahasenan, S. Mobashery, P. Huang, G. Agarwal and R. Fridman (2013). "Discoidin domain receptors: unique receptor tyrosine kinases in collagen-mediated signaling." J Biol Chem **288**(11): 7430-7437.

Gao, Y., D. Mruk, H. Chen, W. Y. Lui, W. M. Lee and C. Y. Cheng (2017). "Regulation of the blood-testis barrier by a local axis in the testis: role of laminin alpha2 in the basement membrane." FASEB J **31**(2): 584-597.

Ghosh, S. and P. W. Sternberg (2014). "Spatial and molecular cues for cell outgrowth during C. elegans uterine development." Dev Biol **396**(1): 121-135.

Gianakas, C. A., D. P. Keeley, W. Ramos-Lewis, K. Park, R. Jayadev, I. W. Kenny, Q. Chi and D. R. Sherwood (2023). "Hemicentin-mediated type IV collagen assembly strengthens juxtaposed basement membrane linkage." J Cell Biol **222**(1).

Gieseler, K., H. Qadota and G. M. Benian (2017). "Development, structure, and maintenance of C. elegans body wall muscle." WormBook **2017**: 1-59.

Goh, L. K. and A. Sorkin (2013). "Endocytosis of receptor tyrosine kinases." Cold Spring Harb Perspect Biol **5**(5): a017459.

Gross, O., B. Beirowski, S. J. Harvey, C. McFadden, D. Chen, S. Tam, P. S. Thorner, N. Smyth, K. Addicks, W. Bloch, Y. Ninomiya, Y. Sado, M. Weber and W. F. Vogel (2004). "DDR1-deficient mice show localized subepithelial GBM thickening with focal loss of slit diaphragms and proteinuria." Kidney Int **66**(1): 102-111.

Hagedorn, E. J., H. Yashiro, J. W. Ziel, S. Ihara, Z. Wang and D. R. Sherwood (2009). "Integrin acts upstream of netrin signaling to regulate formation of the anchor cell's invasive membrane in C. elegans." Dev Cell **17**(2): 187-198.

Hisamoto, N., Y. Nagamori, T. Shimizu, S. I. Pastuhov and K. Matsumoto (2016). "The C. elegans Discoidin Domain Receptor DDR-2 Modulates the Met-like RTK-JNK Signaling Pathway in Axon Regeneration." PLoS Genet **12**(12): e1006475.

Hohenester, E. and P. D. Yurchenco (2013). "Laminins in basement membrane assembly." Cell Adh Migr **7**(1): 56-63.

Jayadev, R., Q. Chi, D. P. Keeley, E. L. Hastie, L. C. Kelley and D. R. Sherwood (2019). "alpha-Integrins dictate distinct modes of type IV collagen recruitment to basement membranes." J Cell Biol **218**(9): 3098-3116.

Jayadev, R., M. Morais, J. M. Ellingford, S. Srinivasan, R. W. Naylor, C. Lawless, A. S. Li, J. F. Ingham, E. Hastie, Q. Chi, M. Fresquet, N. M. Koudis, H. B. Thomas, R. T. O'Keefe, E. Williams,

A. Adamson, H. M. Stuart, S. Banka, D. Smedley, C. Genomics England Research, D. R. Sherwood and R. Lennon (2022). "A basement membrane discovery pipeline uncovers network complexity, regulators, and human disease associations." Sci Adv **8**(20): eabn2265.

Jayadev, R. and D. R. Sherwood (2017). "Basement membranes." Curr Biol **27**(6): R207-R211.

Juskaite, V., D. S. Corcoran and B. Leitinger (2017). "Collagen induces activation of DDR1 through lateral dimer association and phosphorylation between dimers." Elife **6**.

Kamath, R. S. and J. Ahringer (2003). "Genome-wide RNAi screening in Caenorhabditis elegans." Methods **30**(4): 313-321.

Keeley, D. P., E. Hastie, R. Jayadev, L. C. Kelley, Q. Chi, S. G. Payne, J. L. Jeger, B. D. Hoffman and D. R. Sherwood (2020). "Comprehensive Endogenous Tagging of Basement Membrane Components Reveals Dynamic Movement within the Matrix Scaffolding." Dev Cell **54**(1): 60-74 e67.

Keeley, D. P. and D. R. Sherwood (2019). "Tissue linkage through adjoining basement membranes: The long and the short term of it." Matrix Biol **75-76**: 58-71.

Kelley, L. C., Q. Chi, R. Caceres, E. Hastie, A. J. Schindler, Y. Jiang, D. Q. Matus, J. Plastino and D. R. Sherwood (2019). "Adaptive F-Actin Polymerization and Localized ATP Production Drive Basement Membrane Invasion in the Absence of MMPs." Dev Cell **48**(3): 313-328 e318.

Kelley, L. C., Z. Wang, E. J. Hagedorn, L. Wang, W. Shen, S. Lei, S. A. Johnson and D. R. Sherwood (2017). "Live-cell confocal microscopy and quantitative 4D image analysis of anchor-cell invasion through the basement membrane in Caenorhabditis elegans." Nat Protoc **12**(10): 2081-2096.

Khoshnoodi, J., V. Pedchenko and B. G. Hudson (2008). "Mammalian collagen IV." Microsc Res Tech **71**(5): 357-370.

Lee, M., E. J. Cram, B. Shen and J. E. Schwarzbauer (2001). "Roles for beta(pat-3) integrins in development and function of Caenorhabditis elegans muscles and gonads." J Biol Chem **276**(39): 36404-36410.

Leitinger, B. (2014). "Discoidin domain receptor functions in physiological and pathological conditions." Int Rev Cell Mol Biol **310**: 39-87.

Lilja, J., T. Zacharchenko, M. Georgiadou, G. Jacquemet, N. De Franceschi, E. Peuhu, H. Hamidi, J. Pouwels, V. Martens, F. H. Nia, M. Beifuss, T. Boeckers, H. J. Kreienkamp, I. L. Barsukov and J. Ivaska (2017). "SHANK proteins limit integrin activation by directly interacting with Rap1 and R-Ras." Nat Cell Biol **19**(4): 292-305.

Martin-Bermudo, M. D. and N. H. Brown (1999). "Uncoupling integrin adhesion and signaling: the betaPS cytoplasmic domain is sufficient to regulate gene expression in the Drosophila embryo." Genes Dev **13**(6): 729-739.

Merchant, S. N., B. J. Burgess, J. C. Adams, C. E. Kashtan, M. C. Gregory, P. A. Santi, R. Colvin, B. Collins and J. B. Nadol, Jr. (2004). "Temporal bone histopathology in alport syndrome." Laryngoscope **114**(9): 1609-1618.

Mihai, C., M. Chotani, T. S. Elton and G. Agarwal (2009). "Mapping of DDR1 distribution and oligomerization on the cell surface by FRET microscopy." J Mol Biol **385**(2): 432-445.

Moerman, D. G. and B. D. Williams (2006). "Sarcomere assembly in C. elegans muscle." WormBook: 1-16.

Mok, D. Z., P. W. Sternberg and T. Inoue (2015). "Morphologically defined sub-stages of C. elegans vulval development in the fourth larval stage." BMC Dev Biol **15**: 26.

Morrissey, M. A., D. P. Keeley, E. J. Hagedorn, S. T. H. McClatchey, Q. Chi, D. H. Hall and D. R. Sherwood (2014). "B-LINK: a hemicentin, plakin, and integrin-dependent adhesion system that links tissues by connecting adjacent basement membranes." Dev Cell **31**(3): 319-331.

Morrissey, M. A. and D. R. Sherwood (2015). "An active role for basement membrane assembly and modification in tissue sculpting." J Cell Sci **128**(9): 1661-1668.

Murphy, J. E., B. E. Padilla, B. Hasdemir, G. S. Cottrell and N. W. Bunnett (2009). "Endosomes: a legitimate platform for the signaling train." Proc Natl Acad Sci U S A **106**(42): 17615-17622.

Naylor, R. W., M. Morais and R. Lennon (2021). "Complexities of the glomerular basement membrane." <u>Nat Rev Nephrol</u> **17**(2): 112-127.

Newman, A. P., J. G. White and P. W. Sternberg (1996). "Morphogenesis of the C. elegans hermaphrodite uterus." <u>Development</u> **122**(11): 3617-3626.

Pani, A. M. and B. Goldstein (2018). "Direct visualization of a native Wnt in vivo reveals that a long-range Wnt gradient forms by extracellular dispersal." <u>Elife</u> **7**.

Park, J. O., J. Pan, F. Mohrlen, M. O. Schupp, R. Johnsen, D. L. Baillie, R. Zapf, D. G. Moerman and H. Hutter (2010). "Characterization of the astacin family of metalloproteases in C. elegans." <u>BMC Dev Biol</u> **10**: 14.

Payne, L. S. and P. H. Huang (2014). "Discoidin domain receptor 2 signaling networks and therapy in lung cancer." <u>J Thorac Oncol</u> **9**(6): 900-904.

Pozzi, A., P. D. Yurchenco and R. V. Iozzo (2017). "The nature and biology of basement membranes." <u>Matrix Biol</u> **57-58**: 1-11.

Rual, J. F., J. Ceron, J. Koreth, T. Hao, A. S. Nicot, T. Hirozane-Kishikawa, J. Vandenhaute, S. H. Orkin, D. E. Hill, S. van den Heuvel and M. Vidal (2004). "Toward improving Caenorhabditis elegans phenome mapping with an ORFeome-based RNAi library." <u>Genome Res</u> **14**(10B): 2162-2168.

Sandri, C., F. Caccavari, D. Valdembri, C. Camillo, S. Veltel, M. Santambrogio, L. Lanzetti, F. Bussolino, J. Ivaska and G. Serini (2012). "The R-Ras/RIN2/Rab5 complex controls endothelial cell adhesion and morphogenesis via active integrin endocytosis and Rac signaling." <u>Cell Res</u> **22**(10): 1479-1501.

Sato, K., A. Norris, M. Sato and B. D. Grant (2014). "C. elegans as a model for membrane traffic." <u>WormBook</u>: 1-47.

Schindelin, J., I. Arganda-Carreras, E. Frise, V. Kaynig, M. Longair, T. Pietzsch, S. Preibisch, C. Rueden, S. Saalfeld, B. Schmid, J. Y. Tinevez, D. J. White, V. Hartenstein, K. Eliceiri, P. Tomancak and A. Cardona (2012). "Fiji: an open-source platform for biological-image analysis." <u>Nat Methods</u> **9**(7): 676-682.

Sherwood, D. R. and P. W. Sternberg (2003). "Anchor cell invasion into the vulval epithelium in C. elegans." <u>Dev Cell</u> **5**(1): 21-31.

Singh, N. and M. Han (1995). "sur-2, a novel gene, functions late in the let-60 ras-mediated signaling pathway during Caenorhabditis elegans vulval induction." <u>Genes Dev</u> **9**(18): 2251-2265.

Sixt, M., B. Engelhardt, F. Pausch, R. Hallmann, O. Wendler and L. M. Sorokin (2001). "Endothelial cell laminin isoforms, laminins 8 and 10, play decisive roles in T cell recruitment across the blood-brain barrier in experimental autoimmune encephalomyelitis." <u>J Cell Biol</u> **153**(5): 933-946.

Staudinger, L. A., S. J. Spano, W. Lee, N. Coelho, D. Rajshankar, M. P. Bendeck, T. Moriarty and C. A. McCulloch (2013). "Interactions between the discoidin domain receptor 1 and beta1 integrin regulate attachment to collagen." <u>Biol Open</u> **2**(11): 1148-1159.

Stiernagle, T. (2006). "Maintenance of C. elegans." <u>WormBook</u>: 1-11.

Suleiman, H., L. Zhang, R. Roth, J. E. Heuser, J. H. Miner, A. S. Shaw and A. Dani (2013). "Nanoscale protein architecture of the kidney glomerular basement membrane." <u>Elife</u> **2**: e01149.

Taylor, J., T. Unsoeld and H. Hutter (2018). "The transmembrane collagen COL-99 guides longitudinally extending axons in C. elegans." <u>Mol Cell Neurosci</u> **89**: 9-19.

Timmons, L., D. L. Court and A. Fire (2001). "Ingestion of bacterially expressed dsRNAs can produce specific and potent genetic interference in Caenorhabditis elegans." <u>Gene</u> **263**(1-2): 103-112.

Tu, H., P. Huhtala, H. M. Lee, J. C. Adams and T. Pihlajaniemi (2015). "Membrane-associated collagens with interrupted triple-helices (MACITs): evolution from a bilaterian common ancestor and functional conservation in C. elegans." <u>BMC Evol Biol</u> **15**: 281.

Unsoeld, T., J. O. Park and H. Hutter (2013). "Discoidin domain receptors guide axons along
longitudinal tracts in C. elegans." Dev Biol **374**(1): 142-152.
Villasenor, R., Y. Kalaidzidis and M. Zerial (2016). "Signal processing by the endosomal
system." Curr Opin Cell Biol **39**: 53-60.
Vogel, B. E. and E. M. Hedgecock (2001). "Hemicentin, a conserved extracellular member of
the immunoglobulin superfamily, organizes epithelial and other cell attachments into oriented
line-shaped junctions." Development **128**(6): 883-894.
Vogel, W. F., R. Abdulhussein and C. E. Ford (2006). "Sensing extracellular matrix: an update
on discoidin domain receptor function." Cell Signal **18**(8): 1108-1116.
Walser, M., C. A. Umbricht, E. Frohli, P. Nanni and A. Hajnal (2017). "beta-Integrin de-
phosphorylation by the Density-Enhanced Phosphatase DEP-1 attenuates EGFR signaling in C.
elegans." PLoS Genet **13**(1): e1006592.
Welcker, D., C. Stein, N. M. Feitosa, J. Armistead, J. L. Zhang, S. Lutke, A. Kleinridders, J. C.
Bruning, S. A. Eming, G. Sengle, A. Niehoff, W. Bloch and M. Hammerschmidt (2021).
"Hemicentin-1 is an essential extracellular matrix component of the dermal-epidermal and
myotendinous junctions." Sci Rep **11**(1): 17926.
Wildwater, M., N. Sander, G. de Vreede and S. van den Heuvel (2011). "Cell shape and Wnt
signaling redundantly control the division axis of C. elegans epithelial stem cells." Development
**138**(20): 4375-4385.
Xiao, Q., Y. Jiang, Q. Liu, J. Yue, C. Liu, X. Zhao, Y. Qiao, H. Ji, J. Chen and G. Ge (2015).
"Minor Type IV Collagen alpha5 Chain Promotes Cancer Progression through Discoidin Domain
Receptor-1." PLoS Genet **11**(5): e1005249.
Xu, H., D. Bihan, F. Chang, P. H. Huang, R. W. Farndale and B. Leitinger (2012). "Discoidin
domain receptors promote alpha1beta1- and alpha2beta1-integrin mediated cell adhesion to
collagen by enhancing integrin activation." PLoS One **7**(12): e52209.
Yochem, J. and R. K. Herman (2003). "Investigating C. elegans development through mosaic
analysis." Development **130**(20): 4761-4768.
Zhang, Z., K. Vuori, H. Wang, J. C. Reed and E. Ruoslahti (1996). "Integrin activation by R-ras."
Cell **85**(1): 61-69.
1062

# Figure 1

**Morphogenesis of the *C. elegans* utse-seam basement membrane-basement membrane tissue connection.**
(**A**) Left and center: Schematic depicting the utse-seam connection. The utse (a syncytial uterine cell) and the seam epidermal cells are both encased in basement membranes (BMs), which are linked by a BM-BM connecting matrix. Right: ventral fluorescence z-projection showing the utse and seam cells visualized by the markers *cdh-3p::mCherry::PLC$^{δPH}$* and *scmp::GFP::CAAX*, respectively. (**B**) Schematic illustrating the role of the utse-seam connection in supporting the uterus during egg-laying muscle contractions. Disruption of the connection results in uterine prolapse. (**C**) Lateral brightfield images of adult worms on control RNAi treatment or RNAi against a key component of the BM-BM linkage, α1 type IV collagen/*emb-9* (RNAi fed from the L1 onward); note the uterine prolapse after loss of collagen. (**D**) A schematic summarizing the development of the utse-seam connection. Fluorescence images shown are ventral z-projections of the utse and the seam at the respective developmental stages from the late L3 to young adult (hours post-hatch at 20°C). Arrow indicates the uterine anchor cell (AC) that fuses with adjacent uterine cells to form the utse. During syncytium formation, the utse makes contact with the seam cells (arrowheads). Scale bars, 20 μm.

**Figure 2** with 1 supplement

**DDR-2 promotes utse-seam attachment and protects against uterine prolapse.**

(**A**) Domain structure of *C. elegans* discoidin domain receptors (DDRs) and exon-intron gene structures of *ddr-1* and *ddr-2*. Exons are color coded according to respective domains. Gene regions that are deleted in respective mutant alleles (italicized text) are demarcated by solid black lines. (**B**) Frequency of uterine prolapse in *ddr-1* and *ddr-2* mutant animals compared to wild type. Data are shown as mean prolapse percentage ± SD, derived from three independent trials (n = 50 animals screened per trial). α1 type IV collagen/*emb-9* RNAi was used as a positive control. ****$P \leq 0.0001$, **$P \leq 0.01$, *$P \leq 0.05$; one-way ANOVA with post hoc Dunnett's test. (**C**) Ventral fluorescence z-projections of the utse *(cdh-3p::mCherry::PLC $^{\delta PH}$)* and seam *(scmp::GFP::CAAX)* cells in wild-type and *ddr-2* knockout *(ddr-2(qy64))* animals from the late L3 to young adult developmental stages. Arrows indicate the AC prior to utse formation and arrowheads denote early contacts between the utse and seam. Dotted lines with asterisks indicate regions of utse-seam detachment in *ddr-2(qy64)* animals. (**D**) Left: 3D isosurface renderings of the utse and seam in representative late L4 wild-type and *ddr-2(qy64)* animals (see also ***Figure 2—video 1***). Right: Quantification of the percentage of animals with utse-seam detachment. Wild type, early L4 to young adult: n = 0/13, n = 0/14, n = 0/22, n = 0/21, n = 0/18 animals with detachments respectively; *ddr-2(qy64)*, early L4 to young adult: n = 1/13, n = 4/14, n = 9/22, n = 7/18, n = 7/15 animals with detachments respectively. **$P \leq 0.01$, n.s. (not significant), $P > 0.05$; Fisher's exact test. Scale bars, 20 $\mu$m.

**Source data 1.** Source data for ***Figure 2***. Raw data of uterine prolapse frequency used to generate ***Figure 2B*** in Microsoft Excel format.

**Figure 2—video 1.** Animation of 3D isosurface rendering of the utse and seam in a representative late L4 wild-type animal; related to ***Figure 2D***.

## **Figure 3** with 1 supplement

**DDR-2 levels peak in the utse and seam during the formation of the BM-BM connection.**
(**A**) Schematic illustrating lateral z-sectioning to determine slices that capture the seam region in contact with the utse and vice-versa. (**B**) Left: Lateral fluorescence images of DDR-2::mNG visualized with the seam marker *scmp::2xmKate2::PLC $^{\delta PH}$* from the early L4 to young adult stages. Dotted yellow regions indicate DDR-2 signal in the seam contacting the utse. White arrowhead denotes DDR-2 signal at the cell surface. Right: Quantification of mean DDR-2::mNG fluorescence intensity in the dotted regions (n ≥ 9 for each developmental stage). ****$P$ ≤ 0.0001, one-way ANOVA with post hoc Dunnett's test. (**C**) Left: Lateral fluorescence images of DDR-2::mNG visualized with the utse marker *cdh-3p::mCherry::PLC $^{\delta PH}$* from the early L4 to young adult stages. Dotted yellow regions indicate DDR-2 signal in the utse contacting the seam. Right: Quantification of mean DDR-2::mNG fluorescence intensity in dotted yellow regions (n = 10 all stages). **$P$ ≤ 0.01, one-way ANOVA with post hoc Dunnett's test. Scale bars, 20 $\mu$m. Box edges in boxplots represent the 25th and 75th percentiles, the line in the box denotes the median value, and whiskers mark the minimum and maximum values.

**Source data 1.** Source data for *Figure 3*. Raw data of DDR-2::mNG fluorescence intensity quantification in the seam and utse used to generate boxplots in *Figure 3B and C* in Microsoft Excel format. Note that boxplot in *Figure 3—figure supplement 1A* was also generated from these data.
**Source data 2.** Source data for *Figure 3—figure supplement 1*. Raw data of DDR-2::mNG and mCherry-tagged vesicle marker co-localization analysis used to generate boxplots in *Figure 3—figure supplement 1B and C* in Microsoft Excel format.

**Figure supplement 1.** DDR-2 is found within endocytic vesicles; related to *Figure 3B and C*.

# Figure 3—figure supplement 1

**DDR-2 is found within endocytic vesicles.**
Related to ***Figure 3***. (**A**) Lateral fluorescence images of DDR-2::mNG in the seam cells from the early L4 to young adult stages. Note that these images are the same animals as in ***Figure 3B*** with a larger cropped field of view to include regions of the seam not in contact with the utse. Quantification of mean fluorescence intensity in the blue and yellow boxed regions is shown on the right (n ≥ 9 all stages). Boxplots for yellow boxed regions are reproduced from ***Figure 3B***. n.s. (not significant), $P > 0.05$; Kruskal-Wallis *H* test. ****$P ≤ 0.0001$, one-way ANOVA with post hoc Dunnett's test. (**B**) Lateral fluorescence images of DDR-2::mNG within a single arm of the utse tissue at the mid L4 stage, visualized together with mCherry-tagged RAB-5, RAB-7, or RAB-11. Boxed regions are magnified on the right. Yellow arrowheads denote DDR-2 punctae in contact with the respective vesicle marker, while blue arrowheads correspond to DDR-2 puncta that not in contact with vesicle markers. The boxplot shows quantification of percentage overlap of DDR-2 punctae with vesicle markers (n = 11 animals per marker). (**C**) Left: Lateral fluorescence images of DDR-2::mNG and mKate2::RAB-5 in the seam region contacting the utse. Arrowheads denote DDR-2 punctae in contact with (yellow) or not in contact with (blue) mCherry::RAB-5 punctae. Right: Percentage overlap of DDR-2 punctae with RAB-5 (n = 15). Scale bars, 20 $\mu$m. Box edges in boxplots represent the 25th and 75th percentiles, the line in the box denotes the median value, and whiskers mark the minimum and maximum values.

# Figure 4 with 1 supplement

**DDR-2 endocytosis is triggered by type IV collagen and DDR-2 is required in both the utse and seam to promote tissue connection.**
(**A**) Top: Ventral fluorescence z-projections of the utse *(nas-22p::2xmKate2::PLC $^{\delta PH}$)* and seam *(scmp::GFP::CAAX)* cells in late L4 wild-type and *ddr-2* kinase domain truncation mutant *(ddr-2(ok574))* animals. Dotted lines with asterisks indicate regions of utse-seam detachment. Bottom: Quantification of the percentage of animals with utse-seam detachment. Wild type, n = 0/26; *ddr-2(ok574)*, n = 5/21 animals with detachments respectively. **$P \leq 0.01$, Fisher's exact test.
(**B**) Top: Lateral fluorescence images of COL-99 (mNG::COL-99), type XVIII collagen (CLE-1::mNG), and α1 type IV collagen (EMB-9::mNG) at the utse-seam connection (dotted yellow regions) of late L4 animals. All images were acquired at the same exposure. Bottom: Quantification of mean fluorescence intensity (n ≥ 10 for each fluorescent protein). ****$P \leq 0.0001$, unpaired two-tailed Student's *t* test. (**C**) Top: Lateral fluorescence z-projections of DDR-2::mNG at the utse-seam connection in control or *emb-9* RNAi-treated late L4 animals. Note that the field of view contains a single utse arm contacting the seam tissue. Arrowheads indicate cell surface accumulation of DDR-2 upon collagen IV depletion. Bottom: Quantification of the average number of discrete DDR-2 punctae at the utse-seam connection on control, *col-99*, *cle-1*, or *emb-9* RNAi treatments (n ≥ 13 animals for all treatments). ****$P \leq 0.0001$, n.s. (not significant), $P > 0.05$; Kruskal-Wallis *H* test with post hoc Dunn's test. (**D**) Top: Lateral fluorescence z-projections of DDR-2::mNG at the utse-seam attachment in control or hemicentin/*him-4* RNAi-treated late L4 animals. Bottom: Quantification of the average number of discrete DDR-2 punctae at the utse-seam attachment on control or *him-4* RNAi-treated animals (n ≥ 13 all treatments). ****$P \leq 0.0001$, Mann-Whitney *U* test. (**E**) Schematic of dominant negative (DN) *ddr-2*. (**F**) Ventral fluorescence z-projections of the utse *(nas-22p::2xmKate2::PLC $^{\delta PH}$)* and seam *(wrt-2p::GFP::PLC $^{\delta PH}$)* tissues in late L4 stage wild-type animals and animals expressing *DN ddr-2* in the utse (driven by *cdh-3* promoter) or the seam (driven by *scm* promoter). Dotted lines with asterisks indicate regions of utse-seam detachment. (**G**) Quantification of the percentage of animals with utse-seam detachment. Wild type, n = 0/23; *DN ddr-2(utse)*, n = 5/23; *DN ddr-2(seam)*, n = 7/25 animals with detachments respectively. **$P \leq 0.01$ 0.01, *$P \leq 0.05$; Fisher's exact test. Scale bars, 20 μm. Box edges in boxplots represent the 25th and 75th percentiles, the line in the box denotes the median value, and whiskers mark the minimum and maximum values.

**Source data 1.** Source data for *Figure 4*. Raw data of fluorescence intensity quantification of tagged collagen ligands at the utse-seam connection used to generate boxplot in *Figure 4B*, and raw data of DDR-2::mNG puncta counts at the utse-seam connection upon collagen ligand or hemicentin depletion used to generate boxplots in *Figure 4C and D* in Microsoft Excel format.
**Source data 2.** Source data for *Figure 4—figure supplement 1*. Raw data of quantification of knockdown efficiencies of tagged collagen ligands and hemicentin at the utse-seam connection used to generate boxplots in *Figure 4—figure supplement 1* in Microsoft Excel format.

**Figure supplement 1.** Collagen and hemicentin knockdown efficiencies; related to *Figure 4C and D*.

# Figure 4—figure supplement 1

**Collagen and hemicentin knockdown efficiencies.**

Related to **Figure 4**. (**A**) Lateral fluorescence images of EMB-9::mRuby2 (type IV collagen) at the utse-seam BM-BM connection in late L4 control or *emb-9* RNAi-treated animals. Mean fluorescence intensity quantified on the right (n ≥ 8 all treatments). (**B**) Lateral fluorescence images of mNG::COL-99 (COL-99) in the body wall muscle tissue in late L4 control or *col-99* RNAi-treated animals. Mean fluorescence intensity quantified on the right (n ≥ 13 all treatments). COL-99 signal in the muscle was analyzed as COL-99 was not detected at the utse-seam connection. (**C**) Lateral fluorescence images of CLE-1::mNG (type XVIII collagen) at the utse-seam BM-BM attachment in late L4 control or *cle-1* RNAi-treated animals. Mean fluorescence intensity quantified on the right (n ≥ 16 all treatments). (**D**) Lateral fluorescence images of HIM-4::mNG (hemicentin) at the utse-seam BM-BM connection in late L4 control or *him-4* RNAi-treated animals. Mean fluorescence intensity quantified on the right (n = 10 all treatments). ****$P \leq 0.0001$, unpaired two-tailed Student's *t* test (EMB-9, CLE-1, and HIM-4) or Mann-Whitney *U* test (COL-99). Scale bars, 20 μm. Box edges in boxplots represent the 25th and 75th percentiles, the line in the box denotes the median value, and whiskers mark the minimum and maximum values.

# Figure 5 with 2 supplements

**DDR-2 regulates integrin levels at the utse-seam connection and integrin functions in the utse and seam to mediate tissue attachment.**
(**A**) Lateral fluorescence z-projections of integrin α subunits INA-1::mNG and PAT-2::2XmNG at the utse-seam connection (yellow arrowheads) from the mid L4 to young adult stages. Blue asterisks and arrowheads denote INA-1 signal in the developing vulva and PAT-2 at muscle attachment sites respectively. Fluorescence intensity at the utse-seam connection measured in yellow boxes. (**B**) Quantification of mean fluorescence intensity in wild-type and *ddr-2* knockout *(ddr-2(qy64))* animals (n ≥ 9 all conditions). Error bars represent SD. ****$P \leq 0.0001$, ***$P \leq 0.001$, **$P \leq 0.01$, *$P \leq 0.05$, n.s. (not significant), $P > 0.05$; unpaired two-tailed Student's *t* test. (**C**) Frequency of uterine prolapse in wild-type and *ddr-2(qy64)* animals treated with control or *ina-1* RNAi. Control n = 1/70, *ddr-2(qy64)* n = 4/60, *ina-1* RNAi n = 22/66, and *ddr-2(qy64)*; *ina-1* RNAi n = 31/60 animals with uterine prolapse respectively. ****$P \leq 0.0001$, *$P \leq 0.05$; Fisher's exact test.
(**D**) Frequency of utse-seam detachments in late L4 wild-type and *ddr-2(qy64)* animals treated with control or *ina-1* RNAi. Control n = 0/21, *ina-1* RNAi n = 0/25, *ddr-2(qy64)* n = 2/21, and *ddr-2(qy64)*; *ina-1* RNAi n = 9/21 animals with detachments respectively. ***$P \leq 0.001$, *$P \leq 0.05$; Fisher's exact test. (**E**) Schematic of dominant negative (DN) β integrin *pat-3*. (**F**) Ventral fluorescence z-projections of the utse *(cdh-3p::mCherry::PLC^{δPH}* or *nas-22p::2xmKate2::PLC^{δPH})* and seam *(wrt-2p::GFP::PLC^{δPH}* or *scmp::GFP::CAAX)* cells in late L4 wild-type animals and animals expressing *DN integrin* in the utse *(zmp-1^{mk50-51}* promoter) or the seam *(wrt-2* promoter). Dotted lines with asterisks indicate regions of utse-seam detachment. (**G**) Percentage of animals with utse-seam detachment. Wild type, n = 0/19; *DN integrin (utse)*, n = 4/18; *DN integrin (seam)*, n = 4/16 animals with detachments respectively. *$P \leq 0.05$, Fisher's exact test. Scale bars, 20 μm.

**Source data 1.** Source data for **Figure 5**. Raw data of INA-1/PAT-2 fluorescence intensity quantification used to generate **Figure 5B** in Microsoft Excel format.
**Source data 2.** Source data for **Figure 5—figure supplement 1**. Raw data of fluorescence intensity quantification of matrix components and ZMP-4 at the utse-seam BM-BM linkage used to generate boxplots in **Figure 5—figure supplement 1** in Microsoft Excel format.
**Source data 3.** Source data for **Figure 5—figure supplement 2**. Raw data of fluorescence intensity quantification of *ina-1* knockdown used to generate plots in **Figure 5—figure supplement 2D and E** in Microsoft Excel format.

**Figure supplement 1.** *ddr-2* loss does not reduce functional levels of fibulin, hemicentin, type IV collagen, and matrix metalloproteinase ZMP-4 at the utse-seam connection; related to **Figure 5**.
**Figure supplement 2.** The integrin α subunits INA-1 and PAT-2 are localized within both the utse and seam cells; related to **Figure 5**.

# Figure 5—figure supplement 1

**ddr-2 loss does not reduce functional levels of fibulin, hemicentin, type IV collagen, and matrix metalloproteinase ZMP-4 at the utse-seam connection.** Related to *Figure 5*. (**A**) Lateral fluorescence images of DDR-2::mNG at the utse-seam BM-BM connection in late L4 control or *ddr-2* RNAi-treated animals. Mean fluorescence intensity quantified on the right (n ≥ 60 all treatments). ****$P ≤ 0.0001$, Mann-Whitney *U* test. (**B**) Lateral fluorescence images of mNG::FBL-1 (fibulin) at the utse-seam connection in late L4 control or *ddr-2* RNAi-treated animals. Mean fluorescence intensity quantified on the right (n ≥ 30 all treatments). n.s. (not significant), $P > 0.05$, Mann-Whitney *U* test. (**C**) Lateral fluorescence images of HIM-4::mNG (hemicentin) at the utse-seam tissue linkage in late L4 control or *ddr-2* RNAi-treated animals. Mean fluorescence intensity quantified on the right (n ≥ 25 all treatments). **$P ≤ 0.01$, unpaired two-tailed Student's *t* test. (**D**) Lateral fluorescence images of HIM-4::mNG (hemicentin) at the utse-seam connection in late L4 control or *him-4* RNAi-treated animals. RNAi was initiated at the early L4 stage. Mean fluorescence intensity quantified on the right (n ≥ 14 all treatments). **$P ≤ 0.01$, unpaired two-tailed Student's *t* test.
(**E**) Lateral fluorescence images of EMB-9::mRuby2 (type IV collagen) at the utse-seam attachment site in late L4 control or *ddr-2* RNAi-treated animals. Mean fluorescence intensity quantified on the right (n ≥ 30 all treatments). ****$P ≤ 0.0001$, unpaired two-tailed Student's *t* test. (**F**) Lateral fluorescence images of matrix metalloproteinase ZMP-4::GFP at the utse-seam tissue connection in late L4 control or *ddr-2* RNAi-treated animals. Mean fluorescence intensity quantified on the right (n = 10 all treatments). n.s. (not significant), $P > 0.05$, Mann-Whitney *U* test. Scale bars, 20 µm. Box edges in boxplots represent the 25th and 75th percentiles, the line in the box denotes the median value, and whiskers mark the minimum and maximum values.

# Figure 5—figure supplement 2

**The integrin α subunits INA-1 and PAT-2 are localized within both the utse and seam cells.**
Related to *Figure 5*. (**A**) Lateral fluorescence z-projections of the integrin α subunit INA-1::mNG between the mid L4 and young adult stages. Localization in the seam and utse are indicated by solid and dotted boxed regions respectively (n ≥ 10 animals examined all stages). (**B**) Lateral fluorescence z-projections of the integrin α subunit PAT-2::2XmNG between the mid L4 and young adult stages. Localization in the seam and utse are indicated by solid and dotted boxed regions respectively (n ≥ 7 animals examined all stages). (**C**) Lateral fluorescence images of INA-1::mNG and PAT-2::2XmNG at the z-plane of the utse surface in late L4 animals (n ≥ 10 animals examined each). Arrow indicates strong INA-1 signal at the utse surface. (**D**) Lateral fluorescence images of INA-1::mNG at the utse-seam connection in late L4 control or *ina-1* RNAi-treated animals. Mean fluorescence intensity quantified on the right (n ≥ 9 all treatments). ****$P ≤ 0.0001$, unpaired two-tailed Student's *t* test. (**E**) Quantification of residual INA-1::mNG fluorescence intensity at the utse-seam interface at the indicated developmental stages following *ina-1* knockdown initiated at the L1 stage. Data are represented as percentages of mean INA-1::mNG at the respective stages in control animals (n ≥ 7 animals examined all stages). Error bars denote SD. (**F**) Lateral fluorescence z-projections of mCherry driven by the *zmp-1^{mK50-51}* promoter in the utse region in late L3 and young adult animals (n = 5 animals examined at each stage).

**Figure 6** with 1 supplement

**DDR-2 upregulates vinculin levels and stabilizes integrin at the utse-seam connection.**
(**A**) Lateral fluorescence z-projections of the integrin activators talin (GFP::TLN-1) and vinculin (DEB-1::mNG) at the utse-seam attachment region (bounded by yellow arrowheads) from the mid L4 to young adult stages. Vinculin was not detected at the utse-seam attachment region at the mid L4 stage (dotted yellow box). Blue arrowheads indicate fluorescence signal at muscle attachment sites. Fluorescence intensity at the utse-seam attachment region was measured in the solid yellow boxes. (**B**) Quantification of TLN-1 and DEB-1 mean fluorescence intensity in wild-type and *ddr-2* knockout *(ddr-2(qy64))* animals (n ≥ 9 all conditions). Error bars represent SD. ****$P \leq 0.0001$, **$P \leq 0.01$, *$P \leq 0.05$, n.s. (not significant), $P > 0.05$; unpaired two-tailed Student's *t* test. (**C**) Frequency of utse-seam detachments observed in control or *deb-1* RNAi-treated late L4 animals. Control n = 0/21, *ina-1* RNAi n = 5/17 animals with detachments respectively. *$P \leq 0.05$, Fisher's exact test. (**D**) Left: Lateral fluorescence images of INA-1::mNG at the utse-seam connection before photobleaching, immediately after photobleaching, and 10 min post-photobleaching in late L4 wild-type and *ddr-2(qy64)* animals. Box indicates bleached region. Right: Quantification of mean INA-1::mNG fluorescence intensity in the bleached region 10 min post-photobleaching. Wild type, n = 17; *ddr-2(qy64)*, n = 12. **$P \leq 0.01$, unpaired two-tailed Student's *t* test. Scale bars, 20 $\mu$m. Box edges in boxplots represent the 25th and 75th percentiles, the line in the box denotes the median value, and whiskers mark the minimum and maximum values.

**Source data 1.** Source data for *Figure 6*. Raw data of fluorescence intensity quantification of TLN-1 and DEB-1 as well as INA-1 recovery after photobleaching used to generate plots in *Figure 6B and D* in Microsoft Excel format.
**Source data 2.** Source data for *Figure 6—figure supplement 1*. Raw data of fluorescence intensity quantification of *deb-1* knockdown at the utse-seam connection used to generate boxplot in *Figure 6—figure supplement 1* in Microsoft Excel format.

**Figure supplement 1.** *deb-1* knockdown efficiency; related to *Figure 6C*.

# Figure 6—figure supplement 1

**_deb-1_ knockdown efficiency.**
Related to **_Figure 6_**. Lateral fluorescence images of DEB-1::mNG (vinculin) at the utse-seam connection in late L4 control or _deb-1_ RNAi-treated animals. Mean fluorescence intensity quantified on the right (n = 12 all treatments). ****$P ≤ 0.0001$, unpaired two-tailed Student's _t_ test. Scale bars, 20 $\mu$m. Box edges in boxplots represent the 25th and 75th percentiles, the line in the box denotes the median value, and whiskers mark the minimum and maximum values.

# Figure 7 with 1 supplement

**Ras/LET-60 acts in the same pathway as DDR-2 and controls integrin stability at the utse-seam tissue connection.**

(**A**) Left: Ventral fluorescence z-projections of the utse *(nas-22p::2xmKate2::PLC $^{\delta PH}$)* and seam *(scmp::GFP::CAAX)* tissues in wild-type and *ddr-2* knockout *(ddr-2(qy64))* animals on Ras/*let-60* RNAi treatment. Dotted lines with asterisks indicate regions of utse-seam detachment. Right: Percentage of animals with utse-seam detachment. Control n = 0/21, *let-60* RNAi n = 11/18, and *let-60* RNAi; *ddr-2(qy64)*, n = 10/18 animals with detachments respectively. n.s. (not significant), *P* > 0.05; Fisher's exact test. (B) Left: Lateral fluorescence images of INA-1::mNG at the utse-seam connection before photobleaching, immediately after photobleaching, and 10 min post-photobleaching in late L4 control or *let-60* RNAi-treated animals. Box indicates bleached region. Right: Quantification of mean INA-1::mNG fluorescence intensity in the bleached region 10 min post-photobleaching. Control n = 16, *let-60* RNAi n = 25. *P ≤ 0.05, unpaired two-tailed Student's *t* test. (C) Left: Ventral fluorescence z-projections of the utse and seam tissues in late L4 wild-type and *let-60* gain-of-function mutant *(let-60(qy203))* animals. Dotted lines with asterisks indicate regions of utse-seam detachment. Right: Quantification of utse-seam detachment frequency. Wild type, n = 0/26; *let-60(qy203)*, n = 11/11 animals with detachments respectively. ****P ≤ 0.0001, Fisher's exact test. (D) Left: Lateral fluorescence images of INA-1::mNG at the utse-seam attachment region before photobleaching, immediately after photobleaching, and 10 min post-photobleaching in late L4 wild-type and *let-60(qy203)* animals. Box indicates bleached region. Right: Quantification of mean INA-1::mNG fluorescence intensity in the bleached region 10 min post-photobleaching (n = 17 each genotype). *P ≤ 0.05, unpaired two-tailed Student's *t* test. Scale bars, 20 μm. Box edges in boxplots represent the 25th and 75th percentiles, the line in the box denotes the median value, and whiskers mark the minimum and maximum values.

**Source data 1.** Source data for **Figure 7**. Raw data of fluorescence intensity quantification of INA-1::mNG for photobleaching experiments used to generate boxplots in **Figure 7B and D** in Microsoft Excel format. Note that source data for **Figure 7—figure supplement 1** is found within **Source data 1**.

**Figure supplement 1.** Ras/*let-60* knockdown reduces INA-1::mNG levels at the utse-seam connection; related to **Figure 7B**.

# Figure 7—figure supplement 1

**Ras/*let-60* knockdown reduces INA-1::mNG levels at the utse-seam connection.**
Related to **Figure 7**. Quantification of mean INA-1::mNG fluorescence intensity at the utse-seam connection prior to photobleaching in control or Ras/*let-60* RNAi-treated late L4 animals (see representative images in **Figure 7B**, pre-bleach panel. Control n = 16; *let-60* RNAi n = 25. ****$P \leq$ 0.0001, unpaired two-tailed Student's *t* test. Scale bars, 20 *µ*m. Box edges in boxplots represent the 25th and 75th percentiles, the line in the box denotes the median value, and whiskers mark the minimum and maximum values.

# Figure 8

**Model of DDR-2 function at the utse-seam tissue connection.**
(**A**) At the early L4 larval stage, DDR-2 is predominantly found at the surfaces of utse and seam cells. (**B**) Between the mid and late L4 stages, type IV collagen assembles in the BM-BM connecting matrix to link the utse and seam tissues. Collagen addition and binding to DDR-2 triggers DDR-2 endocytosis and activation. (**C**) By the late L4, endocytic localization of DDR-2 reaches peak levels. Ras/LET-60 acts in the same pathway and may function downstream of DDR-2 to stabilize integrin adhesion. Thus, type IV collagen has a dual role in fastening the tissue connection and in signaling through DDR-2 to bolster cell adhesion at the linkage site.