# Modelling the viral dynamics of the SARS-CoV-2 Delta and Omicron variants in different cell types

Clare P. McCormack[1]*, Ada W. C. Yan[2]*, Jonathan C. Brown[2], Ksenia Sukhova[2], Thomas P. Peacock[2], Wendy S. Barclay[2], Ilaria Dorigatti[1]

[1]MRC Centre for Global Infectious Disease Analysis, School of Public Health, Imperial College London, London, United Kingdom

[2]Department of Infectious Disease, Imperial College London, London, United Kingdom

*contributed equally, joint corresponding authors

## Abstract

We use viral kinetic models fitted to viral load data from *in vitro* studies to explain why the SARS-CoV-2 Omicron variant replicates faster than the Delta variant in nasal cells, but slower than Delta in lung cells, which could explain Omicron's higher transmission potential and lower severity. We find that in both nasal and lung cells, viral infectivity is higher for Omicron but the virus production rate is higher for Delta. However, the differences are unequal between cell types, and ultimately leads to the basic reproduction number and growth rate being higher for Omicron in nasal cells, and higher for Delta in lung cells. In nasal cells, Omicron alone can enter via a TMPRSS2-independent pathway, but it is primarily increased efficiency of TMPRSS2-dependent entry which accounts for Omicron's increased activity. This work paves the way for using within-host mathematical models to understand the transmission potential and severity of future variants.

## Introduction

Since its designation as a variant of concern (VOC) by the World Health Organization on 26[th] November 2021[1], the Omicron (B.1.1.529\BA.1) variant of severe acute respiratory coronavirus 2 (SARS-CoV-2) has rapidly displaced the Delta (B.1.617.2) variant to become the dominant SARS-CoV-2 variant globally[2–4]. Analyses have demonstrated that Omicron can partially evade the immunity generated through previous infection and vaccination[5–8], thereby leading to reduced vaccine effectiveness against symptomatic disease[9].

Despite this reduction in vaccine effectiveness against symptomatic disease, the risk of severe outcomes (including hospital admission and death) following infection with Omicron is substantially lower than following infection with Delta, for both vaccinated and unvaccinated individuals[10,11]. While much remains to be understood about the mechanisms underpinning this observation, a reduction in the capacity of Omicron relative to Delta to replicate in lung cells has been suggested as a possible explanation[12].  On the other hand, Omicron's increased ability to replicate in nasal cells has been suggested as an explanation for Omicron's transmission advantage over Delta observed from epidemiological data[13–15].

It has been hypothesised that these differences in viral replication capacity in different cells can be attributed to how Omicron and Delta utilise different pathways to enter host cells[12,16–18]. All SARS-CoV-2 viruses can enter cells which express both ACE2 and TMPRSS2 proteins, via fusion of the viral and cell membranes or via early endosomes[19]. However, Omicron is also able to efficiently enter ACE2+ cells via fusion from the endosome after endocytosis, without the involvement of TMPRSS2[16,20]. This could expand the range of cells that Omicron can infect.  A complication is by entering cells via the endosome, SARS-CoV-2 may be inhibited by endosomal restriction factors such as interferon-induced IFITM proteins[21,22] or NCOA7[23]. Understanding how virological properties shape the viral dynamics of SARS-CoV-2 in different cell types can enhance our understanding of the mechanisms underlying the observed differences in transmissibility between the Omicron and Delta variants. The different entry pathways and their inhibition by endosomal restriction factors are shown in Figure 1.

**Figure 1. Illustration of TMPRSS2-dependent and TMPRSS2-independent entry pathways and their inhibition by endosomal restriction factors**. In ACE2+ TMPRSS2+ cells (right), virus can enter in a TMPRSS2-dependent way via membrane fusion either at the cell surface or the early endosome; alternatively, virus can enter in a TMPRSS2-independent way through the late endosome. In ACE2+ TMPRSS2- cells (left), only the TMPRSS2-independent pathway is available. Both pathways can be inhibited by endosomal restriction factors. Created with BioRender.com.

Mechanistic mathematical models, calibrated against virological data, are a powerful tool for exploring viral dynamics as they provide a framework to quantify key characteristics of different variants and build an understanding of the drivers of inter-individual variation in response to infection. For example, previous within-host modelling studies of SARS-CoV-2 have enabled an understanding of the effect of covariates such as age, sex, and disease severity on viral load[24], the degree of heterogeneity between individuals[25], and the potential effect of antivirals and masking on viral load[26–30]. Models fitted to data from vaccine trials have suggested correlates of protection[31], while more theoretical immunological models have also been developed with the ultimate aim of understanding the interplay between the immune response and disease severity[32–34]. Studies linking the within- and between-host scales have improved the understanding of the relationship between Ct values and infectiousness[25,35], suggested characteristics of optimal testing regimes[36] and provided explanations for population-level epidemiological observations, e.g. the apparent reduction in viral load observed during the declining phase of an epidemic[37].

The studies mentioned above have all fitted their models to data from *in vivo* studies. While dynamics in humans are of ultimate interest, modelling viral kinetics and the immune response *in vivo* is

extremely complex. Moreover, studies in humans are often limited, with little or no information on
the timing and amount of the viral inoculum, sparse longitudinal observations, and often no available
measurements before the start of symptom onset[25,35]. *In vitro* studies offer a simplification of *in vivo*
dynamics, where both the inoculum and method of inoculation is controlled, and where some
components of the immune response – such as the adaptive immune response – are eliminated. Viral
load can be measured from the start of infection, which is important for estimating the growth rate
and basic reproduction number of the virus, which in turn can be used to assess the potential for load
variants to outcompete existing variants.

Here, using viral kinetics models calibrated against viral replication data generated through
experimental infection studies in primary human nasal epithelial cells (hNECs) and immortalised Calu-
3 lung cells, we characterise and compare the dynamics of Omicron and Delta viruses in each cell type
by quantifying key properties such as the basic reproduction number and growth rate, and by
exploring how these properties vary for different entry pathways, and in the presence and absence of
functional endosomal restriction factors.

## Methods

### Experimental Design/Data

Full details of the viral kinetics experiments conducted are provided in Peacock et al.[20] . In brief, Calu-
3 and primary human nasal epithelial cells (hNECs) were inoculated with Omicron BA.1 and
Delta/B.1.617.2 isolates at an MOI of 0.001 (Calu-3) or 0.05 (hNECs), and incubated for 1 hr at 37°C.
The inoculum was then removed and, in the case of Calu-3 cells, replaced with 1 mL serum-free
DMEM. For Calu-3 cells, 100 µL of supernatant was collected for titration at 18, 24, 48 and 72 hours
post-infection. For hNECs, at the same timepoints post-infection, the supernatant was collected by
adding 200 µL of serum-free DMEM to the apical surface, incubating for 10 minutes, and removing for
collection.

Additional experiments were performed to investigate the effect of Camostat mesylate (henceforth
referred to as Camostat) and Amphotericin B on viral kinetics. Camostat is a serine protease inhibitor
and thus inhibits TMPRSS2, while Amphotericin B inhibits the restriction of viral endosomal entry by
endosomal restriction factors such as IFITM proteins[38–40]. Cells were pre-treated both basolaterally
and apically with either 50 µM of Camostat, 1µM of Amphotericin B or no drug for 2 hours prior to
infection, and this concentration of drug was maintained in the basolateral media throughout the
experiment. Infections and collection of timepoints were performed as above.

All experiments were performed in triplicate. Infectious viral titres were quantified by plaque assay

and viral genome numbers were quantified by E gene RT-qPCR.

**Mathematical Models**

*Model 1*

To gain an initial understanding of the differences between the observed infection dynamics of the

Omicron and Delta variants across each cell type, we first developed a simple model of the viral

dynamics which assumed a single virus entry pathway for both variants. A simple schematic of the

model is provided in Figure 2. Model equations are in the Supplementary Methods.

Target cells $T$ become infected at a rate $\beta$ (target cell infection rate) per infectious virion ($V_{Inf}$). The

target cell infection rate is defined as the proportion of cells infected per day, per unit inoculum of 1

pfu/mL. Following an eclipse phase of mean duration of $1/\tau$ days, infectious cells (I) produce infectious

and non-infectious virus. Infectious virus ($V_{Inf}$), as measured by plaque-assay, and total (infectious and

non-infectious) virus ($V_{RNA}$), as measured by RT-qPCR, are produced at rates $\omega_{Inf}$ and $\omega_{RNA}$

respectively. (While the RT-qPCR assay could detect viral RNA released from lysed cells as well as from

extracellular viral particles, the model assumes that all detected viral RNA originates from viral

particles.) The duration of the eclipse phase $1/\tau$ reflects the speed of a single viral replication cycle,

whereas the rate of viral production $\omega_{Inf}$ also reflects replication capacity. Infectious cells have a

mean lifespan of $1/\delta$ days, and infectious and non-infectious virus is assumed to decay at rates $\kappa_{Inf}$

and $\kappa_{RNA}$ respectively, with additional loss due to entry into cells. Cells which express the ACE2

receptor, which is required for SARS-CoV-2 virus entry, are considered target cells.

**Figure 2: Compartmental diagram of model 1**. Target cells $T$ become infected at a rate $\beta$ per infectious virion ($V_{Inf}$). Following
an eclipse phase of mean duration of $1/\tau$ days, infectious cells (I) produce infectious virus as measured by plaque-assay ($V_{Inf}$)
and total virus as measured by qPCR ($V_{RNA}$) at rates $\omega_{Inf}$ and $\omega_{RNA}$ respectively. Infected cells have a mean lifespan of $1/\delta$
days, and infectious and total virus are assumed to decay at rates $\kappa_{Inf}$ and $\kappa_{RNA}$ respectively.

*Model 2*

The experimental data showed that while Delta can only enter ACE2$^+$ cells by utilizing the TMPRSS2

protein, Omicron can enter these cells both through TMPRSS2- dependent and TMPRSS2-independent

pathways. Thus, in Model 2 we modify Model 1 to account for the two possible cell entry pathways,

where TMPRSS2-independent pathways can be used to infect all ACE2$^+$ cells, and TMPRSS2-dependent

pathways can only infect ACE2$^+$ TMPRSS2$^+$ cells. We assume that all ACE2$^+$ cells become infected

through TMPRSS2-independent pathways at a rate $\beta_E$ per infectious virion ($V_{Inf}$) and that ACE2$^+$

TMPRSS2$^+$ target cells ($T_+$) also become infected through TMPRSS2-dependent pathways at a rate $\beta_T$

per infectious virion. We assume that endosomal restriction factors decrease the infectivity through

the TMPRSS2-independent pathways by a factor $f_E$, and through TMPRSS2-dependent pathways by a

factor $f_T$. Cells infected through TMPRSS2-independent and TMPRSS2-dependent pathways ($L_E$ and $L_T$

respectively) undergo eclipse phases of mean duration $1/\tau_E$ and $1/\tau_T$ days respectively. Following

the eclipse phase, infectious cells (I) behave as per Model 1. The flow diagram of Model 2 is described

in Figure 3.

**Figure 3: Compartmental diagram for Model 2**. ACE2$^+$ TMPRSS2$^-$ target cells (T.) become infected through TMPRSS2-

independent pathways at a rate $\beta_E$ per infectious virion (V$_{Inf}$). ACE2$^+$ TMPRSS2$^+$ target cells (T$_+$) become infected through

TMPRSS2-independent pathways at a rate $\beta_E$ per infectious virion, and through TMPRSS2 -dependent pathways at a rate

$\beta_T$ per infectious virion. Endosomal restriction factors decrease the infectivity through TMPRSS2-independent pathways by

a factor $f_E$, and through TMPRSS2-dependent pathways by a factor $f_T$. Cells infected through TMPRSS2-independent and

TMPRSS2-dependent pathways ($L_E$ and $L_T$ respectively) undergo eclipse phases of mean duration $1/\tau_E$ and $1/\tau_T$ days

respectively. Following the eclipse phase, infectious cells (I) behave as per Model 1.

Model 2 is fitted to hNEC and Calu-3 data without drug, in the presence of Camostat, in the presence

of Amphotericin B, and in the presence of both drugs. For each virus strain/cell type combination, all

data (with and without drugs) is fitted simultaneously. Camostat is assumed to completely inhibit

TMPRSS2-dependent pathways (we set $\beta_T = 0$); although Camostat also inhibits other serine

proteases, but as TMPRSS2 is the main serine protease involved in SARS-CoV-2 virus entry, and

because the activation of S2' in TMPRSS2-independent pathways likely occurs by cathepsin proteases

rather than serine proteases, for simplicity we assume that Camostat does not affect TMPRSS2-

independent pathways. Amphotericin B is assumed to completely remove endosomal restriction (we

set $f_E = f_T = 0$).

*Inferential framework*

We calibrated the models to the observed data in a Bayesian inferential framework using Markov

Chain Monte Carlo (MCMC) methods. In the results, we reported the median and 95% credible interval

(CrI) of the parameter estimates. Full details of the model fitting algorithms are provided in the

Supplementary Methods.

**Results**

**Data Description**

**Figure 4: Summary of Data.** (A) Infectious viral load (pfu/mL) quantified by plaque assay and (B) RNA copy number/mL
quantified by RT-qPCR for Omicron (blue triangles) and Delta (red dots) for different cell types (columns) and drug treatments
(rows). The dotted line shows the limit of detection; markers on the lines represent measurements below the limit of
detection. Cells were infected on day 0.

Figure 4 summarises the study data. In control wells with no drug, temporal trends in viral growth
varied both between strains and cell type. In hNECs, Omicron grew more rapidly than Delta, with viral
load peaking by 1 day post infection (p.i.) for Omicron, compared with 2 days p.i. for Delta. However,
despite this initial growth advantage, viral load was higher for Delta compared to Omicron at 3 days
p.i.. In Calu-3 cells, the peak in viral load was observed 2-3 days p.i. for both Omicron and Delta. In
contrast with hNECs where the strain corresponding to the highest viral load was dependent on
measurement time, infectious and non-infectious viral load was higher for Delta than Omicron across
all time points for Calu-3 cells.

In the presence of the drug Camostat which inhibits serine proteases, and thus TMPRSS2-dependent
pathways, replication of Delta was severely inhibited in hNECs with no detectable increase in virus
throughout the course of the experiment. In contrast, Omicron successfully replicated in hNECs in the
presence of Camostat, albeit at a slower rate than in untreated wells. However, similar levels of viral

load were obtained by 2-3 days p.i. in both the presence and absence of Camostat. For both strains,

the presence of Amphotericin B had little impact on viral load dynamics in hNECs. In Calu-3 cells,

infectious virus was not detected at any point for either Omicron or Delta in the presence of Camostat.

In the presence of the drug Amphotericin B which is described as specifically inhibiting endosomal

restriction[22,39], Omicron infectious viral titres were consistently higher than in control wells, until 3

days p.i. when titres became similar. Delta geometric mean infectious viral titres were similar with

and without Amphotericin B.

Experiments with both drugs combined were only performed for Calu-3 cells. For both virus strains,

the infectious viral load was below the limit of detection in the presence of Camostat only. Adding

Amphotericin B raised the infectious viral load to above the limit of detection for all replicates for

Omicron, but only for some replicates for Delta (Figure 4).

**Viral Fitness**

**Model 1**

**Figure 5: Model 1 fits to data from hNECs without drugs.** Dots show the data, black lines show the maximum likelihood
fit, shaded areas show the 95% credible interval (CrI) and dotted lines show the limit of detection.

**Figure 6: Model 1 fits to data from Calu-3 cells without drugs**. Dots show the data, black lines show the maximum likelihood fit, shaded areas show the 95% credible interval (CrI) and dotted lines show the limit of detection.

For both Omicron and Delta, the simple model (Model 1) captured the viral dynamics observed in control wells of hNECs and Calu-3 cells (Figures 5, 6). For hNECs, we estimated a higher target cell infection rate $\beta$ for Omicron relative to Delta (2.13e-05 (95% CrI: 1.54e-05, 3.07e-05) cell (pfu/mL)$^{-1}$ day$^{-1}$ for Omicron, 8.89e-08 (95% CrI: 5.91e-08, 1.28e-07) cell (pfu/mL)$^{-1}$ day$^{-1}$ for Delta), but a lower infectious virus production rate $\omega_{Inf}$ for Omicron relative to Delta (4.07e+03 (95% CrI: 3.13e+03, 5.50e+03) pfu/mL cell$^{-1}$ day$^{-1}$ for Omicron, 3.42e+05 (95% CrI: 2.41e+05, 5.04e+04) pfu/mL cell$^{-1}$ day$^{-1}$ for Delta). All estimated parameters are presented in Supplementary Table 2. These results suggest an approximately 1.5-fold difference in the growth rate between these strains in hNECs, with a growth rate of 11.61 day$^{-1}$ (95% CrI: 10.43,12.99) for Omicron and 6.87 day$^{-1}$ (95% CrI: 6.33,7.50) for Delta. Furthermore, we estimated a higher basic reproduction number ($R_0$) for Omicron (65.04 (95% CrI: 51.94, 82.71)) compared with Delta (23.13 (95% CrI: 19.91, 27.23)), thereby suggesting an overall increase in viral fitness of Omicron relative to Delta in hNECs.

For Calu-3 cells, we estimated a higher target cell infection rate for Omicron than Delta (9.23e-07 (95% CrI: 6.17e-07, 1.36e-06) cell (pfu/mL)$^{-1}$ day$^{-1}$ for Omicron, 1.42e-07 (95% CrI: 9.80e-08, 2.05e-07) cell

(pfu/mL)$^{-1}$ day$^{-1}$ for Delta) and lower infectious virus production rate for Omicron than Delta (4.13e+02

(95% CrI: 2.85e+02,6.07e+02) cell (pfu/mL)$^{-1}$ day$^{-1}$ for Omicron, 3.73e+03 (95% CrI:

2.67e+03,5.32e+03) cell (pfu/mL)$^{-1}$ day$^{-1}$ for Delta). In contrast with hNECs, our results indicate an

overall increase in viral fitness of Delta relative to Omicron in Calu-3 cells as we estimated a higher

basic reproduction number of Delta (36.67 (95% CrI: 33.69, 39.81)) compared to Omicron (24.23 (95%

CrI: 22.14, 26.70)) in this cell type. In addition, we estimated a larger growth rate of Delta relative to

Omicron in Calu-3 cells, with a growth rate of 8.78 day$^{-1}$ (95% CrI: 8.40, 9.15) for Delta compared with

7.25 day$^{-1}$ (95% CrI: 6.86, 7.60) for Omicron (for details see Supplementary Table 2).

For both cell types, the target cell infection rate ($\beta$), which is positively correlated with the probability

of a virion successfully infecting a cell, was higher for Omicron than Delta. The infectious virus

production rate ($\omega_{Inf}$), which is directly proportional to the burst size (the total number of infectious

virions produced by one infected cell), was higher for Delta than Omicron.  However, the magnitude

of each of these differences meant that the basic reproduction number and growth rate, which are

both functions of the target cell infection rate and the infectious virus production rate (see

Supplementary Methods), were higher for Omicron in hNECs, but higher for Delta in Calu-3 cells

(Supplementary Table 2).

**Model 2**

**Figure 7. Model 2 fits to data from hNECs with and without drugs.**  Dots show the data, black lines show the maximum
likelihood fit, shaded areas show the 95% credible interval (CrI) and dotted lines show the limit of detection.  Markers on
the lines represent measurements below the limit of detection.

**Figure 8. Model 2 fits to data from Calu-3 cells with and without drugs**. Dots show the data, black lines show the
maximum likelihood fit, shaded areas show the 95% credible interval (CrI) and dotted lines show the limit of detection.
Markers on the lines represent measurements below the limit of detection.

When fitting Model 2 to the data, we found the same qualitative result that Omicron's $R_0$ is higher

than Delta's in hNECs, but lower in Calu-3 cells (Figures 7 and 8). In hNECs, $R_0$ for Omicron was

estimated to be 106.59 (95% CrI: 85.92, 134.72), while for Delta it was estimated to be 21.61 (95% CrI:

18.97, 25.30). In Calu-3 cells, $R_0$ for Omicron was estimated to be 16.53 (95% CrI: 15.15, 17.99), and

for Delta was estimated to be 27.87 (95% CrI: 25.60, 30.34). $R_0$ estimates were comparable between

Models 1 and 2, except for Omicron in hNECs, where the $R_0$ estimates obtained with Model 2 were

higher. This discrepancy occurred because Model 2 predicts that Omicron can use both TMPRSS2-

dependent and TMPRSS2-independent pathways in hNECs (see next section), whereas for all other

strain and cell combinations only TMPRSS2-dependent pathways are active. When only TMPRSS2-

dependent pathways are active, then in the absence of drug, Models 1 and Model 2 are the same. All

parameter estimates are presented in Supplementary Table 1.

**Role of Cell Entry Pathways**

As Camostat inhibits serine proteases including TMPRSS2, Camostat-sensitive viruses must be able to

use TMPRSS2-independent pathways for cell entry; thus, the fit of Model 2 to the data in the presence

of Camostat provides information on the role of cell entry pathways. We found that in hNECs, Omicron

was able to utilise both TMPRSS2-dependent and TMPRSS2-independent pathways. The median

estimates of $R_{0T}$ and $R_{0E}$ were 92.13 (95% CrI: 72.95, 118.27) and 15.44 (95% CrI 12.86, 18.52)

respectively (Supplementary Table 3).  On the other hand, in hNECs, Delta was unable to effectively

use TMPRSS2-independent pathways, as $R_{0E} < 1$ (median 0.15, 95% CrI: (0.07, 0.30)). In Calu-3s,

neither virus was able to use TMPRRS2-independent pathways effectively.  $R_{0E}$ was estimated to be

slightly greater than 1 for both Omicron (median 1.34 (95% CrI: 1.14, 1.55)) and Delta (median 1.71

(95% CrI: 1.47, 1.98)) but these small values of $R_0$ are insufficient for the viral load to exceed the limit

of detection during the time course of the experiment (Figure 8 Camostat panels).  Note that for

Omicron in hNECs, the estimate for $R_{0T}$ is comparable to the overall $R_0$ for Model 1.  This is because

the increase in viral load due to TMPRSS2-dependent pathways is much faster than through TMPRSS2-

independent pathway, so the TMPRSS2-dependent pathway is the main contributor to the initial

exponential increase, from which Model 1 estimates $R_0$.

It has been suggested that Omicron's use of TMPRRS2-independent pathways explains its fitness

advantage over Delta[12,16,17]. However, even the TMPRSS2-specific basic reproduction number is higher

for Omicron than Delta in hNECs, with a median $R_{0T}$ equal to 92.13 (95% CrI: 72.95, 118.27) for

Omicron compared to 21.46 (95% CrI:18.89, 25.05) for Delta.  In fact, $R_{0T}$ alone for Omicron is greater

than the overall $R_0$ for Delta.  Therefore, it is primarily Omicron's more efficient use of TMPRSS2-

dependent pathways, rather than its utilisation of TMPRSS2-independent pathways, which gives it a

fitness advantage over Delta in hNECs.

**Role of endosomal restriction**

As Amphotericin B inhibits restrictions to virus entry imposed by endosomal restriction factors, the fit

of Model 2 to the Amphotericin B data provides information on the role of endosomal restriction

factors in shaping the viral dynamics observed.  The parameters $f_E$ and $f_T$ capture the degree of

inhibition of TMPRSS2-independent and TMPRSS2-dependent pathways by endosomal restriction

factors respectively, and Amphotericin B is assumed to set these parameters to 0.  Comparing across

cell types, inhibition by endosomal restriction factors was higher in Calu-3 cells than hNECs across all

strains and pathways (higher values of $f_E$ and $f_T$ for Calu-3 cells compared to hNEC cells in

Supplementary Table 4), with the exception of TMPRSS2-independent pathways for Delta, where $f_E$

could not be estimated precisely. Inhibition was similar for both Omicron and Delta in hNECs, but

Omicron was more inhibited by endosomal restriction factors than Delta in Calu-3 cells for both

pathways (higher values of $f_E$ and $f_T$ for Omicron than Delta in Calu-3 cells in Supplementary Table

4). Comparing across pathways, $f_E$ and $f_T$ were similar for each virus strain in hNECs, but in Calu-3

cells, $f_E$ was higher than $f_T$ for both virus strains, suggesting more inhibition of TMPRSS2-independent

pathways than of TMPRSS2-dependent pathways.

Given that TMPRSS2-independent pathways are susceptible to inhibition by endosomal restriction

factors, and that previous variants were not able to use endosomal restriction pathways, one may

hypothesise that Omicron has overcome endosomal restriction in hNECs, and that TMPRSS2-

independent pathways would also be accessible by other virus-cell combinations if endosomal

restriction were lifted.  However, we predict that even if endosomal restriction is lifted for the three

other virus-cell combinations (as per the Amphotericin B experiments) $R_{0EX}$ remains low.  This

suggests that other factors contribute to Omicron's increased use of TMPRSS2-independent pathways

in hNECs. We predict that in the absence of endosomal restriction factors, Omicron would still have a

higher overall basic reproduction number and growth rate than Delta in hNECs, but these quantities

would be similar between variants in Calu-3 cells ($R_{0X}$ in Supplementary Table 3 and $r_X$ in

Supplementary Table 6).

**Changing the assumed eclipse phase duration**

A caveat of the above results is that we have fixed the length of the eclipse phase.  The eclipse phase

in this model reflects the speed of viral replication with the cell.  To date, studies which have measured

the duration of the eclipse phase in the SARS-CoV-2 viral life cycle have used viruses which enter target

cells through TMPRSS2-dependent pathways only[41,42]. Thus, the duration of the eclipse phase for

TMPRSS2-independent pathways is unclear and our knowledge of this component of the viral life cycle

is limited. To address this, we conducted a sensitivity analysis around the duration of the eclipse phase.

In Model 1, $\tau$ was set to 6, 4, or 2 day$^{-1}$.  In Model 2, $\tau_T$ was set to 6, 4 or 2 day$^{-1}$, and $\tau_E$ was set to 4,

2 or 1 day$^{-1}$.  We found that for Model 1, regardless of the duration of the eclipse phase assumed – as

long as it was the same between strains – Omicron had a higher $R_0$ than Delta in hNECs, and Delta

had a higher $R_0$ than Omicron in Calu-3 cells (Supplementary Table 4).  Also, all estimated $R_0$ values

decreased as the duration of the eclipse phase decreased, as expected theoretically.  This is because

if, at the start of infection, cells start producing virus sooner due to a short eclipse phase, then each

individual infected cell needs to lead to fewer secondary infections to reproduce the dynamics

observed.  Similarly, for Model 2, we found that regardless of the values of $\tau_E$ and $\tau_T$, as long as these

were assumed to be the same between strains, Omicron had a higher $R_0$ than Delta in hNECs, and

Delta had a higher $R_0$ than Omicron in Calu-3 cells; only Omicron in hNECs used TMPRSS2-independent

pathways efficiently; and endosomal restriction only had a significant effect in Calu-3 cells

(Supplementary Table 5).  However, allowing the duration of the eclipse phase to be different between

strains, no longer allows us to draw conclusions about whether $R_0$ is greater for Omicron or Delta

(Supplementary Tables 3 and 5). On the other hand, estimates of the growth rate $r$ only varied slightly

with the assumed duration of the eclipse phase for both models (Supplementary Tables 4 and 6).

## Discussion

By fitting a simple within-host model to viral kinetics data for Omicron and Delta in hNECs and Calu-3

cells we found that Omicron has a fitness advantage over Delta in hNECs, but Delta has a fitness

advantage over Omicron in Calu-3 cells, as measured by both the growth rate and basic reproduction

number of the virus. These findings are consistent with previously published studies showing faster

replication of Omicron compared with Delta in human nasal epithelial cells[14,16], and faster replication

of Delta compared with Omicron in the ex vivo explant cultures of human lungs[12] and Calu-3 cells[43].

We estimated that Omicron had a higher rate of infection $\beta$ in both cell types, which could be linked

to increased ACE2 binding affinity, likely due to mutations in the spike protein. Evidence is mixed as

to whether Omicron has a higher ACE2 binding affinity than Delta. Cameroni et al.[44] found that

Omicron has a higher human ACE2 binding affinity than Delta, but Mannar et al.[45] found that ACE2

binding affinity is similar for Omicron and Delta, though higher than the ancestral strain. On the other

hand, Zhang et al.[46] found that the Omicron spike protein required a higher level of ACE2 than Delta

for efficient membrane fusion. We also estimated that Omicron had a lower rate of infectious virus

production $\omega_{Inf}$ than Delta for both cell types. Changes to the rate of viral production could be due

to mutations in either spike or non-spike proteins. The basic reproduction number is positively

correlated with both the rate of infection $\beta$ and the rate of infectious virus production $\omega_{Inf}$ in our

models. In hNECs, the increase in the rate of infection $\beta$ for Omicron was greater than the decrease

in infectious virus production $\omega_{Inf}$, resulting in a larger basic reproduction number and growth rate

for Omicron compared to Delta; the converse was true in Calu-3 cells.

Previous studies have proposed that TMPRSS2-independent pathways which are only available to

Omicron are responsible for its faster growth in hNECs. Our study found that Omicron can use

TMPRSS2-independent pathways in hNECs, consistent with the previous studies. However, the

growth rate and basic reproduction number for TMPRSS2-independent pathways was low and

insufficient to explain the overall fitness advantage for Omicron over Delta in hNECs. On the other

hand, the growth rate and basic reproduction number for TMPRSS2-dependent pathways was also

higher for Omicron than Delta in hNECs and was sufficient to explain the overall fitness advantage.

Evidence for the continued importance of TMPRSS2-dependent pathways for Omicron was provided

by a study by Metzdorf et al.[47] which found that the growth of Omicron in the nose and lung was

attenuated in TMPRSS2 knockout mice. In Calu-3 cells, neither Omicron nor Delta can use TMPRSS2-

independent pathways, thereby providing further insight into how these viruses use different

pathways to enter target cells.

We estimated the degree of inhibition in viral growth imposed by endosomal restriction factors by

fitting our model to data where the cells were treated with Amphotericin B, as Amphotericin B inhibits

the action of endosomal restriction factors.  The main hypotheses tested were i) whether Omicron

was able to evade endosomal restriction in hNECs; ii) whether this evasion enabled Omicron to use

TMPRSS2-independent pathways; and iii) whether this was responsible for Omicron's growth

advantage.  We found that the fitness of Delta in hNECs without endosomal restriction, as measured

by the basic reproduction number $R_{0X}$ and the growth rate $r_X$, was still lower than that of Omicron in

hNECs with endosomal restriction. Hence, reduced endosomal restriction alone do not explain

Omicron's fitness advantage in hNECs.  Although we were able to quantify the degree of inhibition by

endosomal restriction factors for both pathways for Omicron in hNECs, we were unable to do so for

Delta, as doing so requires either the infectious viral load to be above the limit of detection in the

presence of Camostat (as was the case for Omicron in hNECs), or requires data on viral growth in the

presence of both drugs (as was the case in Calu-3 cells).  Hence, we were unable to make a direct

comparison at a pathway-specific level between Omicron and Delta in hNECs.  In Calu-3 cells, we found

that Omicron was more inhibited by endosomal restriction factors than Delta, for either TMPRSS2-

dependent or TMPRSS2-independent pathways.

The model has several limitations. First, we fixed the infected cell death rate, the rate at which viruses

lose infectivity, the RNA degradation rate and the duration of eclipse phases, according to values from

the literature.  It is currently unclear whether the duration of the eclipse phase differs between

variants. To test this, we conducted a sensitivity analysis where the model was fitted assuming

different lengths of the eclipse phase.  This sensitivity analysis showed that if eclipse phase lengths

were assumed to be the same between Omicron and Delta, while the estimated values of $R_0$ changed

with the assumed length of the eclipse phase, the trends and relationships in $R_0$ and the growth rate

estimates $r$ between Omicron and Delta (for example, that Omicron has a higher $R_0$ and growth rate

estimate $r$ than Delta in hNECs) remained unchanged. Allowing the eclipse phase to be different

between strains, no longer allows us to draw conclusions about whether $R_0$ is greater for Omicron or

Delta (Supplementary Tables 3 and 5), but the differences in $r$ are robust.  This result reflects the

finding that if the eclipse phase is allowed to vary, the growth rate $r$ is identifiable but $R_0$ is not.  Future

studies could measure the duration of the eclipse phase for each cell type and virus strain using single-

cycle growth kinetics experiments.

We also did not test the effect of changing the assumed values of the infected cell or virus decay rates.

Studies have shown that Omicron survives for longer on surfaces than the ancestral strain[48] and

Delta[49], so it is plausible that extracellular virus may also differ in stability in our experiments.

Moreover, fixing the parameters using values from previous studies may underestimate the
uncertainty in $R_0$ and r.

Another limitation of the model is that it assumes 100% effectiveness for both drugs.  Assuming 100%
effectiveness of Camostat means interpreting all remaining viral growth as due to TMPRSS2-
independent pathways, so if the drug is less than 100% effective, the contribution of TMPRSS2-
independent pathways will be overestimated.  Assuming 100% effectiveness of Amphotericin B means
interpreting viral growth in the presence of Amphotericin B as the viral growth in the complete
absence of endosomal restriction.  If the drug were less than 100% effective, the viral load in the
complete absence of endosomal restriction could be higher than that suggested by the data, so the
effect of endosomal restriction would be underestimated.

Last, the structure of the model is such that the initial growth rate is independent of the inoculum
size. It is plausible that differences in growth rate between hNECs and Calu-3 cells observed in the
experiments were partly due to the differing inoculum sizes used, as a higher inoculum could, for
example, trigger endosomal restriction factors more quickly – an effect which was not modelled.  A
higher multiplicity of infection (MOI) was used for hNECs as a lower MOI did not always successfully
infect these primary cells; repeating the Calu-3 experiments at the same higher MOI would strengthen
confidence in our comparison between cell types.

The models developed in this study can inform future studies using more complex models.  Models
capturing more immune components and how the virus spreads within the host are useful for
understanding contributors to disease severity and the effects of antivirals[32–34], but because they have
many parameters, it is often difficult to determine their values by fitting them to data.  This study can
help parameterize these models.

The models developed in this study can be applied to different SARS-CoV-2 virus strains to help
enhance our understanding of transmission potential of new strains as the virus continues to evolve.
While between-strain differences in the speed of viral replication is typically assessed by comparing
viral titre measurements at individual timepoints, adopting a mathematical modelling approach using
longitudinal timepoints enables us to estimate key characteristics of different strains such as the
growth rate and viral fitness, thus providing a deeper insight into the factors shaping observed
differences in viral dynamics at the individual and population level.

## References

1.  WHO. Classification of Omicron (B.1.1.529): SARS-CoV-2 Variant of Concern. *Who* 1 https://www.who.int/news/item/26-11-2021-classification-of-omicron-(b.1.1.529)-sars-cov-2-variant-of-concern (2021).

2.  GISAID. GISAID - hCov19 Variants. *Gisaid* https://gisaid.org/hcov19-variants/ (2022).

3.  WHO. Tracking SARS-CoV-2 variants. *Who* https://www.who.int/en/activities/tracking-SARS-Co Preprint at (2021).

4.  Viana, R. *et al.* Rapid epidemic expansion of the SARS-CoV-2 Omicron variant in southern Africa. *Nature* **603**, 679–686 (2022).

5.  Cele, S. *et al.* Omicron extensively but incompletely escapes Pfizer BNT162b2 neutralization. *Nature* **602**, 654–656 (2022).

6.  Planas, D. *et al.* Considerable escape of SARS-CoV-2 Omicron to antibody neutralization. *Nature* **602**, 671–675 (2022).

7.  Cao, Y. *et al.* Omicron escapes the majority of existing SARS-CoV-2 neutralizing antibodies. *Nature* **602**, 657–663 (2022).

8.  Dejnirattisai, W. *et al.* SARS-CoV-2 Omicron-B.1.1.529 leads to widespread escape from neutralizing antibody responses. *Cell* **185**, 467-484.e15 (2022).

9.  Andrews, N. *et al.* Covid-19 Vaccine Effectiveness against the Omicron (B.1.1.529) Variant. *New England Journal of Medicine* **386**, 1532–1546 (2022).

10. Nyberg, T. *et al.* Comparative analysis of the risks of hospitalisation and death associated with SARS-CoV-2 omicron (B.1.1.529) and delta (B.1.617.2) variants in England: a cohort study. *The Lancet* **399**, 1303–1312 (2022).

11. Wolter, N. *et al.* Early assessment of the clinical severity of the SARS-CoV-2 omicron variant in South Africa: a data linkage study. *The Lancet* **399**, 437–446 (2022).

12. Hui, K. P. Y. Y. *et al.* SARS-CoV-2 Omicron variant replication in human bronchus and lung ex vivo. **603**, (2022).

13. Willett, B. J. *et al.* SARS-CoV-2 Omicron is an immune escape variant with an altered cell entry pathway. *Nature Microbiology 2022 7:8* **7**, 1161–1179 (2022).

14. Barut, G. T. *et al.* The spike gene is a major determinant for the SARS-CoV-2 Omicron-BA.1 phenotype. *Nat Commun* **13**, 1–14 (2022).

15. Lyngse, F. P. *et al.* Household transmission of the SARS-CoV-2 Omicron variant in Denmark. *Nat Commun* **13**, 5573 (2022).

16. Willett, B. J. *et al.* SARS-CoV-2 Omicron is an immune escape variant with an altered cell entry pathway. *Nat Microbiol* (2022) doi:10.1038/s41564-022-01143-7.

17. Meng, B. *et al.* Altered TMPRSS2 usage by SARS-CoV-2 Omicron impacts infectivity and fusogenicity. *Nature 2022 603:7902* **603**, 706–714 (2022).

18. Suzuki, R. *et al.* Attenuated fusogenicity and pathogenicity of SARS-CoV-2 Omicron variant. *Nature* **603**, 700–705 (2022).

19.  Hoffmann, M. *et al.* SARS-CoV-2 Cell Entry Depends on ACE2 and TMPRSS2 and Is Blocked by
a Clinically Proven Protease Inhibitor. *Cell* **181**, 271-280.e8 (2020).

20.  Peacock, T. P. *et al.* The altered entry pathway and antigenic distance of the SARS-CoV-2
Omicron variant map to separate domains of spike protein. *bioRxiv* 2021.12.31.474653 (2022)
doi:10.1101/2021.12.31.474653.

21.  Winstone, H. *et al.* The Polybasic Cleavage Site in SARS-CoV-2 Spike Modulates Viral
Sensitivity to Type I Interferon and IFITM2. *J Virol* **95**, e02422-20 (2021).

22.  Peacock, T. P. *et al.* The furin cleavage site in the SARS-CoV-2 spike protein is required for
transmission in ferrets. *Nat Microbiol* **6**, 899–909 (2021).

23.  Khan, H. *et al.* TMPRSS2 promotes SARS-CoV-2 evasion from NCOA7-mediated restriction.
*PLoS Pathog* **17**, e1009820 (2021).

24.  Challenger, J. D. *et al.* Modelling upper respiratory viral load dynamics of SARS-CoV-2. *BMC*
*Med* **20**, 25 (2022).

25.  Ke, R., Zitzmann, C., Ho, D. D., Ribeiro, R. M. & Perelson, A. S. In vivo kinetics of SARS-CoV-2
infection and its relationship with a person's infectiousness. *Proc Natl Acad Sci U S A* **118**,
e2111477118 (2021).

26.  Gonçalves, A. *et al.* Timing of Antiviral Treatment Initiation is Critical to Reduce SARS-CoV-2
Viral Load. *CPT Pharmacometrics Syst Pharmacol* **9**, 509–514 (2020).

27.  Gonçalves, A. *et al.* SARS-CoV-2 viral dynamics in non-human primates. *PLoS Comput Biol* **17**,
(2021).

28.  Perelson, A. S. & Ke, R. Mechanistic Modeling of SARS-CoV-2 and Other Infectious Diseases
and the Effects of Therapeutics. *Clin Pharmacol Ther* **109**, 829–840 (2021).

29.  Koelle, K. *et al.* Masks Do No More Than Prevent Transmission: Theory and Data Undermine
the Variolation Hypothesis. Preprint at https://doi.org/10.1101/2022.06.28.22277028 (2022).

30.  Ito, K., Piantham, C. & Nishiura, H. Relative instantaneous reproduction number of Omicron
SARS-CoV-2 variant with respect to the Delta variant in Denmark. *J Med Virol* **94**, 2265–2268
(2022).

31.  Alexandre, M. *et al.* Modelling the response to vaccine in non-human primates to define
SARS-CoV-2 mechanistic correlates of protection. *Elife* **11**, e75427 (2022).

32.  Jenner, A. L. *et al.* COVID-19 virtual patient cohort suggests immune mechanisms driving
disease outcomes. *PLoS Pathog* **17**, e1009753 (2021).

33.  Rowlatt, C. F. *et al.* Modelling the within-host spread of SARS-CoV-2 infection, and the
subsequent immune response, using a hybrid, multiscale, individual-based model. Part I:
Macrophages. *bioRxiv* (2022) doi:10.1101/2022.05.06.490883.

34.  Getz, M. *et al.* Iterative community-driven development of a SARS-CoV-2 tissue simulator.
*bioRxiv* (2021) doi:10.1101/2020.04.02.019075.

35.  Marc, A. *et al.* Quantifying the relationship between SARS-CoV-2 viral load and infectiousness.
*Elife* **10**, e69302 (2021).

36.   Larremore, D. B. *et al.* Test sensitivity is secondary to frequency and turnaround time for
COVID-19 screening. *Sci Adv* **7**, eabd5393 (2021).

37.   Hay, J. A. *et al.* Estimating epidemiologic dynamics from cross-sectional viral load
distributions. *Science (1979)* **373**, eabh0635 (2021).

38.   Brass, A. L. *et al.* The IFITM Proteins Mediate Cellular Resistance to Influenza A H1N1 Virus,
West Nile Virus, and Dengue Virus. *Cell* **139**, 1243–1254 (2009).

39.   Lin, T.-Y. *et al.* Amphotericin B Increases Influenza A Virus Infection by Preventing IFITM3-
Mediated Restriction. *Cell Rep* **5**, 895–908 (2013).

40.   Qian, J. *et al.* Primate lentiviruses are differentially inhibited by interferon-induced
transmembrane proteins. *Virology* **474**, 10–18 (2015).

41.   Brahim Belhaouari, D. *et al.* The Strengths of Scanning Electron Microscopy in Deciphering
SARS-CoV-2 Infectious Cycle. *Front Microbiol* **11**, 2014 (2020).

42.   Hou, Y. J. *et al.* SARS-CoV-2 D614G variant exhibits efficient replication ex vivo and
transmission in vivo. *Science (1979)* **370**, 1464–1468 (2020).

43.   Shuai, H. *et al.* Attenuated replication and pathogenicity of SARS-CoV-2 B.1.1.529 Omicron.
*Nature* **603**, 693–699 (2022).

44.   Cameroni, E. *et al.* Broadly neutralizing antibodies overcome SARS-CoV-2 Omicron antigenic
shift. *Nature* **602**, 664–670 (2022).

45.   Mannar, D. *et al.* SARS-CoV-2 Omicron variant: Antibody evasion and cryo-EM structure of
spike protein–ACE2 complex. *Science (1979)* **375**, 760–764 (2022).

46.   Zhang, J. *et al.* Structural and functional impact by SARS-CoV-2 Omicron spike mutations. *Cell
Rep* **39**, 110729 (2022).

47.   Metzdorf, K. *et al.* TMPRSS2 Is Essential for SARS-CoV-2 Beta and Omicron Infection. *Viruses*
**15**, 271 (2023).

48.   Chin, A. W. H., Lai, A. M. Y., Peiris, M. & Man Poon, L. L. Increased Stability of SARS-CoV-2
Omicron Variant over Ancestral Strain. *Emerging Infectious Disease journal* **28**, 1515 (2022).

49.   Hirose, R. *et al.* Differences in environmental stability among SARS-CoV-2 variants of concern:
Both Omicron BA.1 and BA.2 have higher stability. *Clinical Microbiology and Infection* (2022)
doi:https://doi.org/10.1016/j.cmi.2022.05.020.

# Acknowledgements

CMC and ID acknowledge funding from the MRC Centre for Global Infectious Disease Analysis

(reference MR/R015600/1), jointly funded by the UK Medical Research Council (MRC) and the UK

Foreign, Commonwealth & Development Office (FCDO), under the MRC/FCDO Concordat agreement

and is also part of the EDCTP2 programme supported by the European Union. AWCY acknowledges

funding from an Imperial College Research Fellowship. ID acknowledges funding by the Wellcome

Trust and Royal Society (grant number 213494/Z/18/Z). This study was conducted as part of G2P-UK

National Virology consortium funded by MRC/UKRI (MR/W005611/1). For the purpose of open access,

the authors have applied a CC BY public copyright licence to any Author Accepted Manuscript version

arising from this submission.

## Author Contributions

580     CMC, AWCY, WSB and ID conceived the study; JCB and KS generated the *in-vitro* data; CMC, AWCY and

581     ID conceived the mathematical models; CMC performed the analysis for Model 1; AWCY performed

582     the analysis for Model 2; CMC, AWCY, JCB, TPP, WSB, and ID contributed to the interpretation of the

583     results; CMC and AWCY wrote the first draft of the manuscript; all authors reviewed and approved the

584     final version of the manuscript.

## Competing Interests

586     The authors declare no competing financial interests.

## Data Availability

588     All data is available at https://github.com/ada-w-yan/deltaomicronmodelling

## Code Availability

590     All code is available at https://github.com/ada-w-yan/deltaomicronmodelling

591