Human assembloids reveal the consequences of *CACNA1G* gene variants in the thalamocortical pathway

Ji-il Kim[1,2,5], Yuki Miura[1,2,5], Min-Yin Li[1,2], Omer Revah[1,2,3], Sridhar Selvaraj[4], Fikri Birey[1,2]
Xiangling Meng[1,2], Mayuri Vijay Thete[1,2], Sergey D. Pavlov[1,2], Jimena Andersen[1,2]
Anca M. Pașca[4], Matthew H. Porteus[4], John R. Huguenard[3] and Sergiu P. Pașca[1,2*]

[1] Department of Psychiatry and Behavioral Sciences, Stanford University, Stanford, CA 94305, USA
[2] Stanford Brain Organogenesis, Wu Tsai Neuroscience Institute, Stanford, CA 94305, USA
[3] Department of Neurology and Neurological Sciences, Stanford University, Stanford, CA 94305, USA
[4] Department of Pediatrics, Stanford University, Stanford, CA 94305, USA
[5] These authors contributed equally
[*] Correspondence to spasca@stanford.edu (S.P.P.)

ABSTRACT

Abnormalities in crosstalk between the thalamus and the cerebral cortex are thought to lead to severe neuropsychiatric disorders, such as epilepsy and psychotic disorders. Pathogenic variants in the *CACNA1G* gene, which encodes the α1G subunit of the thalamus-enriched T-type voltage-gated calcium channel Ca$_V$3.1, are associated with absence seizures, intellectual disability, and schizophrenia, but the cellular and circuit level consequences of these genetic variants in humans remain unknown. Here, we developed an *in vitro* human assembloid model of the thalamocortical pathway to systematically dissect the contribution of genetic variants in T-type calcium channels. We discovered that a *CACNA1G* variant (M1531V) associated with seizures led to hypersynchronous activity in the thalamus and in cortical neurons in thalamo-cortical assembloids. In contrast, *CACNA1G* loss, which has been associated with risk of schizophrenia, resulted in abnormal thalamocortical connectivity that was related to both increased spontaneous thalamic activity and aberrant thalamic axonal projections. Taken together, these results illustrate the utility of organoid and assembloid systems for interrogating human genetic disease risk variants at both cellular and circuit level.

Introduction

Communication between the thalamus and the cerebral cortex via reciprocal thalamocortical projections is essential for processing sensory information and for modulating cognitive functions in the mammalian nervous system (*1-3*). Anatomical or functional dysconnectivity in the thalamocortical circuit is often associated with psychiatric disorders, including schizophrenia and autism spectrum disorder (*4, 5*). Moreover, hypersynchronous thalamocortical activity is a core feature of absence seizures in epilepsy (*6, 7*). The thalamus-enriched voltage-dependent calcium channel Ca$_V$3.1, encoded by the *CACNA1G* gene, is a key mediator of activity in this circuit (*8, 9*). Importantly, pathogenic variants in *CACNA1G* are strongly associated with disease; putative loss-of-function variants have been linked to schizophrenia and intellectual disability (*10, 11*), while putative gain-of-function variants have been linked to epilepsy (*12-14*). However, the cell-specific consequences of different human *CACNA1G* gene variants are unknown. Further, how these cellular defects influence activity in the thalamocortical pathway is a mystery. Addressing this has been particularly challenging in the context of evolutionary changes in thalamocortical projections in primates, in which, for instance, thalamocortical projections innervate the dorsal forebrain at earlier stages than in rodents (*15*).

We previously introduced an assembloid approach to model network-level interactions of the developing nervous system by functionally integrating regionalized neural organoids (*16, 17*), and applied these models to study disease

mechanisms in neurodevelopmental diseases (*18-21*). Despite efforts to derive thalamus-like neural cells from mouse embryonic stem cells (*22*) and human induced pluripotent stem (hiPS) cells (*23*), an experimental model of human thalamocortical connectivity has not yet been developed *in vitro*.

Here, we developed a platform that combines assembloids, high-resolution live cell imaging, optogenetics, and extracellular recordings to investigate early development of the human thalamocortical pathway. Using hiPS cells, we generated human 3D regionalized neural organoids resembling the diencephalon, the developmental precursor to thalamic glutamatergic neurons. The neurons in human diencephalic organoids (hDiO) transcriptionally and electrophysiologically recapitulated features of thalamic glutamatergic neurons. When assembled with human cortical organoids (hCO), these neurons sent thalamocortical projections to the cortical neurons. Using optogenetics and simultaneous extracellular recording or calcium imaging, we demonstrated the formation of the functional thalamocortical pathway in these assembloids. To study *CACNA1G*-related channelopathies, we generated hiPS cell lines with gain- and loss-of-function variants in *CACNA1G*. Probing thalamo-cortical assembloids revealed spontaneous hypersynchronous neuronal activity associated with the gain-of-function *CACNA1G* variant; in contrast, a *CACNA1G* loss-of-function variant resulted in asynchronous hyperactivity following abnormal thalamocortical connectivity.

Results

Generation of 3D human diencephalic organoids (hDiO) from hiPS cells

To generate neural organoids that include thalamic glutamatergic neurons, we implemented guided differentiation approach to generate regionalized neural organoids from hiPS cells, as we have done previously for other nervous system domains (*18*, *19*, *24*, *25*). We applied small molecules that modulate the SHH (SAG), WNT (CHIR), and BMP (BMP7) signaling to differentiate cells to diencephalic-like fates (**Fig. 1A**, **Supplementary Fig. 1A**). The neural organoids treated with a combination of 1 μM CHIR, 100 nM SAG and 30 ng/mL of BMP7 showed a higher expression of the diencephalic markers *TCF7L2* and *OTX2* and a lower expression of the rostral forebrain marker *FOXG1* when compared to human cortical organoids (hCO) (**Supplementary Fig. 1B**). We refer to these regionalized organoids with the broader term human diencephalic organoids (hDiO) (*26*). To examine cellular diversity more comprehensively in these organoids, we performed droplet-based single-cell RNA sequencing (scRNAseq) at day 100 of differentiation (**Fig. 1A**, **Supplementary Fig. 1C**). Uniform Manifold

Figure 1. Generation of 3D human diencephalic organoids (hDiO). (**A**) Schematic describing the generation of 3D human diencephalic organoids (hDiO) from hiPS cells and scRNAseq analysis. (**B**) UMAP visualization of single cell RNA expression in hDiO at day 100 of *in vitro* differentiation (*n* = 15,653 cells from 3 hiPS cell lines). (**C**) VoxHunt spatial brain mapping of hDiO onto the BrainSpan dataset of the human developing brain (PCW15–24). (**D**) VoxHunt spatial brain mapping of hDiO onto the Allen Brain Institute E15.5 mouse brain data. (**E**) UMAP of integrated hDiO and human primary samples from PCW16–20. (**F**) UMAP visualization of thalamic glutamatergic markers *TCF7L2*, *SLC17A6* and *NTNG1*, and (**G**) percentage distribution of cell clusters in each sample. (**H**) CUBIC-cleared human primary samples immunostained with TCF7L2 (green) and RedDot2 nuclear marker (magenta). Scale bar: 1000 μm. (**I**) Immunostaining for TCF7L2 (green), and Hoechst (blue) in hCO and hDiO at day 80, and quantification of TCF7L2+ cells (*n* = 20 for hCO and *n* = 19 for hDiO from 3 differentiation experiments of 4 hiPS cell lines; Mann-Whitney test, ****$P < 0.0001$). Scale bar: 100 μm. (**J**) Immunostaining for GFP (green) and VGLUT2 (magenta) in hDiO at day 80, and quantification of VGLUT2+/EYFP+ cells (*n* = 18 from 2 differentiation experiments of 3 hiPS cell lines). Scale bar: 100 μm. (**K**) Whole-cell recording of hDiO neurons (flattened on an insert) and representative traces showing hyperpolarization-induced rebound spikes in AAV-hSYN1:: EYFP+ neurons. Cells were held at –50 mV for rebound spike recordings and were held at –60 mV for depolarization-induced spikes. Examples shown for –100 pA and –50 pA current induced rebound spikes. Rebound spikes were blocked by bath application of TTA-P2 (3 μM). Abbreviations: VFC; ventrolateral frontal cortex, V1C; primary visual cortex, STR; striatum, STC; Posterior superior temporal cortex, S1C; primary somatosensory cortex, OFC; orbital frontal cortex, MFC; medial frontal cortex, MD; mediodorsal nucleus of thalamus, M1C; primary motor cortex, ITC; inferior temporal cortex, IPC; posterior inferior parietal cortex, HIP; hippocampus, DFC; dorsolateral prefrontal cortex, CBC; cerebellar cortex, AMY; amygdala, A1C; primary auditory temporal cortex.

Approximation and Projection (UMAP) showed the presence of a large cluster of glutamatergic neurons that express thalamic glutamatergic neuronal markers *TCF7L2* and *SLC17A6*. This analysis also revealed clusters of GABAergic neurons (*SLC32A1*), neuronal progenitors (*GADD45G*), choroid plexus-like cells (*TTR*), astrocytes (*AQP4*), *OLIG2*-expressing cells, proliferating cells (*MKI67*), and a negligible number of cells expressing the rostral forebrain marker *FOXG1* or the dopaminergic neuron marker *TH* (**Fig. 1B**, **Supplementary Fig. 1D**). Unbiased mapping of scRNAseq data onto midgestational human samples using VoxHunt demonstrated hDiO approximates post-conceptional week (PCW) 15–24 (*27*, *28*) with high similarity to the human mediodorsal nucleus of the thalamus (MD) (**Fig. 1C**, **Supplementary Fig. 1E**). In addition, VoxHunt analysis using mouse *in situ* gene expression data indicated high scaled correlation of hDiO with the diencephalon of embryonic day 15.5 mouse (**Fig. 1D**). We also confirmed the presence of cells in the hDiO expressing early developing thalamic regional markers (*29*) for caudal thalamus (cTh) (*GBX2*, *CD47*), rostral thalamus (rTH) (*SOX14*, *GATA3*), prethalamus (PTh) (*GAD2*, *DLX5*) and epithalamus (Eth) (*TAC1*, *POU4F*), with neglectable expression of zona limitans intrathalamica (ZLI) (*PITX2*, *FOXA2*) and pretectum (rPT) (*1C1QL2*, *BHLHE23*) markers (**Supplementary Fig. 1F**). To further validate the transcriptional similarity between cells *in vitro* and their *in vivo* counterpart, we compared scRNAseq data from hDiO and the developing human thalamus (**Fig. 1E–G**, **Supplementary Fig. 1G**). We obtained a PCW20 diencephalic region, performed scRNAseq analysis, and combined this with the publicly available PCW16–20 human thalamus scRNAseq data (*30*). UMAP visualization demonstrated that thalamic glutamatergic neurons (*SLC17A6*), GABAergic neurons (*GAD1*), astrocytes (*GFAP*), glial progenitors (*SOX9*), oligodendrocytes (*SOX10*) and cycling cells (*MKI67*) are present in both hDiO and primary tissues. In contrast, endothelial cells (*FLT1*), microglia (*CSF1R*), mesenchymal cells (*COL1A1*), pericytes (*HIGD1*), blood cells (*HBA1*), and melanocytes (*DCT*) were only detected in primary tissue, while choroid plexus-like cells (*TTR*) were only found in hDiO (**Fig. 1E**, **Supplementary Fig. 1G**). Thalamic glutamatergic neurons (*TCF7L2$^+$/SLC17A6$^+$/NTNG1$^+$*) overlapped across datasets (**Fig. 1F**) and represented approximately up to 45% of cells in hDiO and primary tissues over time (**Fig. 1G**). To further examine expression of thalamic glutamatergic neuronal markers at protein levels in hDiO, we performed immunostaining for TCF7L2, which was highly expressed in human diencephalon tissue at PCW19 (**Fig. 1H**). We found expression of TCF7L2 in approximately 20% of cells in hDiO, but not in hCO (**Fig. 1I**). Moreover, approximately 80% of AAV-hSYN1::EYFP virally labeled neurons in hDiO were VGLUT2$^+$ glutamatergic neurons (**Fig. 1J**). Lastly, we characterized the electrophysiological properties of hDiO neurons. Patch-clamp recordings from AAV-hSYN1::EYFP labeled hDiO neurons showed spontaneous action potentials (**Supplementary Fig. 2A**), as well as characteristic post-

hyperpolarization rebound spiking (*31-33*) (**Fig. 1K**, **Supplementary Fig. 2B**) and fast T-type channel-dependent calcium influx as measured by barium current recordings (**Supplementary Fig. 2C, D**), which could be blocked by the T-type calcium channel blocker TTA-P2 (*34*, *35*). The current-voltage (*I-V*) curve (measured by barium currents) showed a typical low-voltage activation profile, with activation voltage threshold around −50 mV and a peak current at −30 mV. T-type currents also showed high reliability for repetitive trials (**Supplementary Fig. 2C**). In summary, neurons in hDiO transcriptionally and electrophysiologically resemble thalamic glutamatergic neurons.

Projections and functional connectivity in thalamo-cortical assembloids

To model the formation of thalamocortical connectivity *in vitro*, we used an assembloid approach as we have previously described for other regionalized neural organoids (*19*, *36*) (**Fig. 2A**). We virally labeled the hCO with hSYN1:: mCherry and the hDiO with hSYN1::EYFP and then assembled them by placing the two regionalized organoids in close contact in an Eppendorf tube. After 72 hours, the two organoids fused to generate a thalamo-cortical assembloid (**Fig. 2B**). To quantify the structural integration between the two organoids, we measured the fluorescence area on each side of the assembloid. We found a progressive increase in reciprocal, bilateral projections between hCO and hDiO over 30 days of assembly (**Fig. 2C, D**). This is consistent with early embryonic stage formation of thalamocortical and corticothalamic projections in primate embryos (*15*) and in contrast with the formation of unilateral projections in cortico-striatal assembloids (*19*). To verify synaptic connections between hDiO projection neurons and hCO neurons, we applied anterograde transsynaptic tracing (*37*) by expressing AAV1-hSYN1::Cre in hDiO and AAV-EF1α::DIO-tdTomato in hCO (**Fig. 2E**). At 1 month after assembly, we found EYFP$^+$/tdTomato$^+$ cells on the hCO side of the thalamo-cortical assembloids, suggestive of thalamocortical connectivity (**Fig. 2F**). We next used retrograde tracing by infecting hCO with rabies-ΔG-Cre-EGFP and AAV1-EF1α::rabies glycoprotein (G) and hDiO with AAV-EF1α::DIO-tdTomato (*38*, *39*) before generating thalamo-cortical assembloids (**Fig. 2G**). After assembly, we observed EGFP$^+$/tdTomato$^+$ double-positive cells on the hDiO side of thalamo-cortical assembloids (**Fig. 2H**). These experiments suggest transsynaptic, anterograde and retrograde transmission in assembloids through thalamocortical connections.

To examine whether thalamo-cortical assembloids form functional connections, we combined optogenetic stimulation with simultaneous extracellular recording in thalamo-cortical assembloids using a custom extracellular recording system (**Fig. 2I**, **Supplementary Fig. 3A**). Acute recordings from individual hDiO using Cambridge NeuroTech™ ultrathin and high-density electrode arrays (**Supplementary Fig. 3B**) revealed spontaneous neuronal activity (**Supplementary Fig. 3C**). Optogenetic stimulation of hDiO

Figure 2. Generation of human assembloids to study thalamocortical connectivity. (**A**) Schematic illustrating the thalamocortical pathway in assembloid model. (**B**) Representative image of a 3D thalamo-cortical assembloid; green, EYFP+ neurons in hDiO; magenta, mCherry+ neurons in hCO. (**C, D**) Quantification of EYFP and mCherry projections in assembloids. (*n* = 10–16 assembloids from 4 differentiation experiments of 3 hiPS cell lines; Kruskal-Wallis test, *P* = 0.0273, **P = 0.0028, ****P < 0.0001). (**E**) Schematic of anterograde viral tracing in thalamo-cortical assembloids using AAV1. (**F**) Representative image showing the presence of EYFP+/tdTomato+ cells in the a-hCO side of thalamo-cortical assembloids. (**G**) Schematic illustrating retrograde viral tracing in thalamo-cortical assembloids using rabies virus. (**H**) Representative image indicating EGFP+/tdTomato+ cells in a-hCO. (**I**) Schematic illustrating extracellular recording coupled with optogenetic stimulation of thalamo-cortical assembloids. Image showing insertion of each electrode into a-hCO and a-hDiO in thalamo-cortical assembloids (bottom). (**J**) Raster plot showing optically-evoked responses in thalamo-cortical assembloids (top) and randomly-selected time points (bottom). (**K**) Quantification of firing rates in a-hCO in assembloids before and after 590 nm light application (*n* = 10 units from 3 differentiation experiments of 2 hiPS cell lines; paired t-test, *P = 0.0102). (**L**) Schematic illustrating calcium imaging coupled with optogenetics in thalamo-cortical assembloids. (**M**) Representative images and (**N**) calcium imaging trace of cells in a-hCO following light stimulation of a-hDiO. (**O**) Median amplitude of ΔF/F signals of optically-evoked responses on the a-hCO side of thalamo-cortical assembloids (*n* = 176 cells from 2 differentiation experiments of 2 hiPS cell lines; Wilcoxon test, *P = 0.039). Abbreviations: Daf; days after fusion of assembloids.

expressing hSYN1::ChrimsonR–tdTomato (*40*) elicited increased firing of isolated single units (**Supplementary Fig. 3D, E**), which was blocked by sodium channel blocker tetrodotoxin (TTX) (**Supplementary Fig. 3E, F**). To probe functional connectivity in thalamo-cortical assembloids, we optogenetically stimulated hDiO of the assembloid (a-hDiO) while recording extracellular signals from the hCO side of the assembloid (a-hCO) (**Fig. 2I–K**). We found increased firing rates of single units in a-hCO following light illumination (**Fig. 2J, K**). In contrast, light illumination did not influence firing rates in a-hCO when the opsin was not expressed in a-hDiO (**Supplementary Fig. 3G–I**),

demonstrating that optically-evoked responses in the thalamo-cortical assembloids are not due to light-related artifacts. We also delivered AAV-hSYN1::GCaMP6f to a-hCO and AAV-hSYN1::ChrimsonR-tdTomato to a-hDiO, and then applied 590 nm light and measured calcium activity in a-hCO neurons after thalamic stimulation (**Fig. 2L**). Consistent with extracellular recording experiments, we detected significantly increased light-dependent calcium responses in a-hCO neurons in thalamo-cortical assembloids (**Fig. 2M–O**). Altogether, these results support the presence of anatomical and functional connectivity in hiPS cell-derived thalamo-cortical assembloids.

Figure 3. Neuronal activity in hDiO carrying pathogenic variants in the *CACNA1G* gene. (**A**) Schematic illustrating the generation of *CACNA1G* M1531V and KO hiPS cells using CRISPR/Cas9 genome editing. (**B**) Schematic illustrating the position of the M1531V variant in Ca$_V$3.1. (**C**) Schematic illustrating the strategy to introduce the *CACNA1G* M1531V point mutation. (**D**) Schematic illustrating the strategy to introduce homozygous *CACNA1G* KO via exon 1 deletion. (**E**) *CACNA1G* gene expression quantified in hDiO from control, *CACNA1G* KO and *CACNA1G* M1531V hiPS cell lines at day 100 (n = 3 differentiation experiments; one-way *ANOVA*, $_F$(2, 9) = 47.31, $P < 0.0001$). (**F**) Representative whole-cell recording traces of hDiO neurons labeled with AAVDJ-hSYN1::EYFP highlighting the hyperpolarization-induced rebound spikes in control, but not in *CANCA1G* KO hDiO neurons. (**G**) Quantification of the number of maximal rebound spikes (n = 12 for control and n = 7 neurons for *CACNA1G* KO; Mann-Whitney test, ***P = 0.0002). (**H**) Schematic illustrating the experimental design for probing spontaneous activity via extracellular recording from control, *CACNA1G* M1531V, and *CACNA1G* KO hDiOs. (**I**) Representative traces of spontaneous extracellular recording from control (left), *CACNA1G* M1531V (middle) and *CACNA1G* KO (right) hDiO. Representative long-term traces (upper) and short-term traces (middle) highlighting the spontaneous hypersynchronous activity in *CACNA1G* M1531V hDiO. Vertical scale bar for upper panel: 1000 µV, Horizontal scale bar for upper panel: 10 seconds, Vertical scale bar for middle panel: 100 µV, Horizontal scale bar for middle panel: 1 second. Lower panel depicts the percentage of electrode channels detecting activity. (**J**) Quantification of hypersynchronous activity in control, *CACNA1G* M1531V, and *CACNA1G* KO hDiOs (control, n = 11 organoids; M1531V, n = 14 organoids; KO, n = 11 organoids; Kruskal-Wallis test, *P = 0.0163, **P = 0.0037). (**K**) Quantification of hypersynchronous activity from *CACNA1G* M1531V hDiO before and after 3 µM TTA-P2 application (n = 9 organoids. Wilcoxon test, *P = 0.0312). (**L**) Firing rates of sorted single units from control, *CACNA1G* M1531V, and *CACNA1G* KO hDiOs (control, n = 56 units from 9 hDiOs; M1531V, n = 78 units from 12 hDiOs; KO, n = 107 units from 11 hDiOs; Kruskal- Wallis test, *P = 0.0336, ***P = 0.0004, ****P < 0.0001). (**M**) Representative spectral power analysis (upper) and band-filtered local field potential (LFP) (lower) in *CACNA1G* M1531V hDiO (same time window as in the middle panel in (**I**)).

Cellular defects in thalamic neurons carrying disease variants in the *CACNA1G* gene

The low-threshold voltage-dependent calcium channel Ca$_V$3.1, encoded by the *CACNA1G* gene, is highly expressed in thalamic neurons in primary human fetal tissue, as well as in hDiO (**Supplementary Fig. 4A, B**). Both presumed gain- and loss-of-function variants in *CACNA1G* lead to severe channelopathies (*10-14*), but the consequences of these

variants on human thalamic neurons and on their cortical neuronal targets remain unknown.

We generated hiPS cell lines with gain- and loss-of-function mutations in the *CACNA1G* gene (**Fig. 3A**). To derive a homozygous gain-of-function line, we used AAV6/Cas9 RNP genome-editing methods in hiPS cells (*41*) resulting in a substitution of methionine 1531 to valine in domain III of *CACNA1G* (*CACNA1G* M1531V; **Fig. 3B, C**,

Supplementary Fig. 4C–E). *De novo* heterozygous *CACNA1G* M1531V variant was reported in patients with early onset seizures and epileptic encephalopathy (*12, 13*). To generate homozygous *CACNA1G* knockout (*CACNA1G⁻/⁻*) hiPS cell lines, we used dual guide RNAs (gRNAs) targeting the exon 1 region of *CACNA1G* (**Fig. 3D**). We found that gene editing with a combination of 2 gRNAs efficiently caused 178 or 179 bp deletions in exon 1 (**Supplementary Fig. 4F–I**). We used 2 knockout hiPS cell lines for subsequent assays. The engineered hiPS cell lines maintained their pluripotency (**Supplementary Fig. 4J**) and could be differentiated into regionalized organoids (**Supplementary Fig. 4K**). We also confirmed a reduction of *CACNA1G* gene expression in *CACNA1G* KO hDiO by RT-qPCR (**Fig. 3E**).

Loss of *CACNA1G* resulted in significantly reduced rebound spiking (**Fig. 3F, G**). Moreover, T-type currents were significantly reduced in patched neurons in *CACNA1G* KO hDiO (**Supplementary Fig. 5A**).

To probe spontaneous neuronal activity in hDiO carrying *CACNA1G* variants, we performed extracellular recordings of control, *CACNA1G* M1531V, and *CACNA1G* KO hDiO (**Fig. 3H**). In M1531V hDiO, we detected striking hypersynchronous events across all recording electrodes as well as an increase in firing in between events (**Fig. 3I, J** and **Supplementary Fig. 5B**). Spectral power analysis showed increased power in both the high-frequency spike band and the local field potential (LFP) during hypersynchronous

Figure 4. Altered neuronal activity and connectivity in thalamo-cortical assembloids with *CACNA1G* pathogenic variants. (**A**) Schematic illustrating extracellular recording from control, *CACNA1G* M1531V and *CACNA1G* KO hCO-hDiO assembloids. (**B**) Representative traces of spontaneous activity in control (left), *CACNA1G* M1531V (middle), and *CACNA1G* KO (right) hCO-hDiO. Vertical scale bar: 100 µV, Horizontal scale bar: 500 ms. Lower panel depicts the percentage of channels with activity. (**C**) Quantification of hypersynchronous activity in a-hCO (upper) and a-hDiO (lower) derived from control, *CACNA1G* M1531V, and *CACNA1G* KO hCO–hDiO (control, $n = 9$ assembloids; M1531V, $n = 11$ assembloids; KO, $n = 5$ assembloids; Kruskal–Wallis test, n.s., not significant, $*P = 0.0242$, $**P = 0.0090$). (**D**) Percentage of a-hCO hypersynchronous events co-occurring a-hDiO (upper) and the percentage of a-hDiO hypersynchronous events co-occurring with a-hCO (lower) ($n = 6$ assembloids that showed hypersynchronous activity at least once). (**E**) Firing rates of sorted single units from control, *CACNA1G* M1531V, and *CACNA1G* KO a-hCOs (upper; control, $n = 17$ units from 7 assembloids; M1531V, $n = 24$ units from 7 assembloids; KO, $n = 32$ units from 5 assembloids; Kruskal-Wallis test, n.s., not significant, $*P = 0.0196$), and a-hDiOs (lower; control, $n = 25$ units from 7 assembloids; M1531V, $n = 48$ units from 10 assembloids; KO, $n = 35$ units from 6 assembloids; Kruskal- Wallis test, n.s., not significant, $*P = 0.0494$). (**F**) Schematic illustrating quantification of thalamocortical projections in hCO-hDiO assembloids. (**G**) Representative images showing thalamocortical projections in hCO-hDiO (left) and their quantification (control, $n = 18$ assembloids; *CACNA1G* M1531V, $n = 8$ assembloids; *CACNA1G* KO, $n = 14$ assembloids; one-way *ANOVA* with Fisher's LSD test; $_F(2, 37) = 4.796$, n.s., not significant, $*P = 0.0142$). (**H**) Representative images of time lapse growth cone dynamics (left) and quantification (control, $n = 82$ growth cones from 9 assembloids; *CACNA1G* KO, $n = 57$ growth cones from 6 assembloids; unpaired *t*-test, $**P = 0.0056$). Scale bar: 10 µm. (**I**) Schematic illustrating optogenetic stimulation and extracellular recordings in hCO-hDiO. (**J**) Representative extracellular recording traces from control and *CACNA1G* KO assembloids highlighting the response of a-hCO by a-hDiO optogenetic stimulation (top) and z-scored power of spike band frequency (bottom). Vertical scale bar: 100 µV, horizontal scale bar: 100 ms. (**K**) Quantification of z-scored power of spike band frequency in a-hCO during light stimulation (control, $n = 9$ assembloids; *CACNA1G* KO, $n = 10$ organoids; Mann-Whitney test, $*P = 0.0435$). (**L**) Firing rate heatmap of all sorted units from control and *CACNA1G* KO a-hCO. (**M**) Percentage of responding units of a-hCO after optogenetic stimulation of the counterpart a-hDiO (control, $n = 17$ units from 8 assembloids; *CACNA1G* KO, $n = 79$ units from 9 assembloids; Chi-square test, $*P = 0.0121$).

events (**Fig. 3M**). Of note, this synchronized neuronal activity was spontaneous, did not require drug or electrical stimulation, and was reminiscent of characteristic patterns of thalamic activity described in absence seizures (*42*). The T-type calcium channel blocker TTA-P2 blocked these spontaneous hypersynchronous activity events in *CACNA1G* M1531V hDiO (**Fig. 3K**), demonstrating that they are mediated by the T-type calcium channel. In contrast, in *CACNA1G* KO hDiO neurons, we found no hypersynchronous events, but we detected increased asynchronous spontaneous firing rates (**Fig. 3L**). These results suggest that different pathogenic variants in *CACNA1G* result in different functional defects in thalamic neurons: seizure-related variants elicited hypersynchronous neuronal activity with high spontaneous firing rates, while homozygous loss of the gene resulted in increased spontaneous firing rates without any hypersynchronous activity.

Altered connectivity in thalamo-cortical assembloids carrying *CACNA1G* pathogenic variants

To compare the circuit-level consequences of the thalamic defects identified in the *CACNA1G* variants, we generated thalamo-cortical assembloids and recorded spontaneous neuronal activity (**Fig. 4A**). Interestingly, we found that the hypersynchronous events in *CACNA1G* M1531V hDiO (**Fig. 3I, J**) were also present on the cortical side of thalamo-cortical assembloids (**Fig. 4B, C**, **Supplementary Fig. 5C–E**). Importantly, most of these events co-occurred in both regions of the assembloids (**Fig. 4D, Supplementary Fig. 5D, E**), further illustrating the formation of functional thalamocortical connections and suggesting transmission of these large, synchronous electrical events. Surprisingly, the increased spontaneous firing of thalamic neurons in *CACNA1G* KO and M1531V hDiO was also present on the cortical side of assembloids in *CACNA1G* KO, but not in M1531V (**Fig. 4E**).

This prompted us to investigate if additional defects, such as abnormal projections, might contribute to this functional defect. Therefore, we next probed anatomic and functional thalamocortical connectivity in assembloids derived from *CACNA1G* variant lines. Interestingly, confocal live imaging of thalamocortical projections in the assembloids (**Fig. 4F**), revealed an increase of projections in *CACNA1G* KO assembloids, but not in *CACNA1G* M1531V assembloids (**Fig. 4G**). To verify if this was related to abnormal axonal growth, we performed live imaging of growth cone dynamics in *CACNA1G* KO assembloids. We found that *CACNA1G* KO growth cones moved longer distances over time in lived-imaged assembloids (**Fig. 4H**). To examine whether increased thalamocortical projections in *CACNA1G* KO assembloids contribute to changes in functional connectivity, we recorded neuronal responses in a-hCO following optogenetic stimulation of ChrimsonR-tdTomato expressing a-hDiO (**Fig. 4I**). We detected an increase in neuronal activity in *CACNA1G* KO a-hCO. Power in the high-frequency band (spike power) was higher in *CACNA1G* KO a-hCO, but not in a-hDiO, following light stimulation (**Fig. 4J, K**; **Supplementary Fig. 5F**). In addition, the proportion of sorted single units that responded to light was also higher in *CACNA1G* KO than control (**Fig. 4L, M**). Altogether, experiments in thalamo-cortical assembloids illustrate a range of cellular- and circuit-level defects associated with *CACNA1G* channelopathies: a gain-of-function variant caused hypersynchronous activity across the thalamocortical pathway, while loss of *CACNA1G* resulted in abnormal projections and activity of thalamic neurons in the thalamocortical pathway.

Discussion

To elucidate the pathophysiology of human developmental channelopathies and to develop effective therapeutics, it is necessary to investigate the degree of ion channel dysfunction (i.e., gain- versus loss-of-function), cell-type specific expression and susceptibility, as well as circuit level consequences and compensation. In this study, we probed the cellular- and circuit-level consequences of *CACNA1G* pathogenic variants associated with different clinical presentations in a new human model of thalamocortical connectivity.

In-frame deletion and protein-truncating variants in *CACNA1G* have been associated with intellectual disability and schizophrenia (*10, 11*). We found that deletion of *CACNA1G* resulted in accelerated growth cone movement of thalamic neurons and increased projections in thalamo-cortical assembloids. Surprisingly, loss of this thalamus-enriched calcium channel also resulted in higher hDiO activity. Taken together, these defects suggest abnormal thalamocortical connectivity, which is consistent with functional imaging reports (*43-46*). In contrast, the M1531V point mutation associated with neonatal-onset epilepsy resulted in hypersynchronous, TTA-P2-sensitive events in thalamic neurons, which were simultaneously present in cortical neurons in assembloids. This is aligned with pathophysiological models of absence seizures that arise from T-type calcium channel-dependent burst firing of reciprocally connected neurons in the thalamus and neocortex (*7*).

There are several advantages to our human cellular model. It enables a multi-level assessment of genetic disease variants, including cell type-specific effects in regionalized organoids to inter-regional interactions in assembloids and the consequences (and potential compensation) in thalamocortical networks. As we have shown before, assembloid models are modular, allowing disentanglement of cell autonomous and non-autonomous effects. Our experimental system also allows simultaneous probing of interconnected neuronal populations.

Moving forward, the current assembloids system can be further improved. There are key GABAergic populations in both cortical circuits and in the thalamus (i.e., reticular neurons). Future studies could incorporate these cells through additional assembloid components. The thalamocortical circuit is part of a larger loop circuit, thus integration with striatal cells and other GABAergic cells of the midbrain could enable analysis of emergent features of these circuits. The development of thalamocortical circuitry is characterized by waves of activity that spread from the thalamus even in the absence of sensory input, which is thought to contribute to maturation and refinement of connections (*47, 48*). Larger field imaging coupled with integration of sensory input through either assembly with sensory cells or following transplantation may enable the study of wave-like phenomena and how these shape circuit formation (*49*). Overall, this *in vitro* platform has the potential to reveal cell and network level defects associated with genetic and environmental perturbations at early, previously inaccessible but critical stages of thalamocortical crosstalk in human development and, ultimately, to be used for testing of therapeutics.

References

1. S. M. Sherman, R. W. Guillery, The role of the thalamus in the flow of information to the cortex. Philos Trans R Soc Lond B Biol Sci 357, 1695-1708 (2002).
2. R. V. Rikhye, A. Gilra, M. M. Halassa, Thalamic regulation of switching between cortical representations enables cognitive flexibility. Nat Neurosci 21, 1753-1763 (2018).
3. D. S. Roy et al., Anterior thalamic circuits crucial for working memory. Proceedings of the National Academy of Sciences of the United States of America 119, e2118712119 (2022).
4. R. Ayub et al., Thalamocortical connectivity is associated with autism symptoms in high-functioning adults with autism and typically developing adults. Transl Psychiatry 11, 93 (2021).
5. D. S. Roy et al., Anterior thalamic dysfunction underlies cognitive deficits in a subset of neuropsychiatric disease models. Neuron 109, 2590-2603 e2513 (2021).
6. M. M. Huntsman, D. M. Porcello, G. E. Homanics, T. M. DeLorey, J. R. Huguenard, Reciprocal inhibitory connections and network synchrony in the mammalian thalamus. Science 283, 541-543 (1999).
7. P. M. Fogerson, J. R. Huguenard, Tapping the Brakes: Cellular and Synaptic Mechanisms that Regulate Thalamic Oscillations. Neuron 92, 687-704 (2016).
8. E. Carbone, H. Lux, A low voltage-activated, fully inactivating Ca channel in vertebrate sensory neurones. Nature 310, 501-502 (1984).
9. E. M. Talley et al., Differential distribution of three members of a gene family encoding low voltage-activated (T-type) calcium channels. Journal of Neuroscience 19, 1895- 1911 (1999).
10. A. M. Alazami et al., Accelerating novel candidate gene discovery in neurogenetic disorders via whole-exome sequencing of prescreened multiplex consanguineous families. Cell reports 10, 148-161 (2015).
11. T. Singh et al., Rare coding variants in ten genes confer substantial risk for schizophrenia. Nature 604, 509-516 (2022).
12. J. Chemin et al., De novo mutation screening in childhood-onset cerebellar atrophy identifies gain-of-function mutations in the CACNA1G calcium channel gene. Brain 141, 1998-2013 (2018).
13. M. Kunii et al., De novo CACNA1G variants in developmental delay and early-onset epileptic encephalopathies. J Neurol Sci 416, 117047 (2020).
14. P. Lory, S. Nicole, A. Monteil, Neuronal Cav3 channelopathies: recent progress and perspectives. Pflugers Archiv : European journal of physiology 472, 831-844 (2020).
15. A. Alzu'bi, J. Homman-Ludiye, J. A. Bourne, G. J. Clowry, Thalamocortical Afferents Innervate the Cortical Subplate much Earlier in Development in Primate than in Rodent. Cereb Cortex 29, 1706-1718 (2019).
16. S. P. Paşca, Assembling human brain organoids. Science 363, 126-127 (2019).
17. S. Kanton, S. P. Pasca, Human assembloids. Development 149, (2022).
18. F. Birey et al., Assembly of functionally integrated human forebrain spheroids. Nature 545, 54-59 (2017).
19. Y. Miura et al., Generation of human striatal organoids and corticostriatal assembloids from human pluripotent stem cells. Nat Biotechnol 38, 1421-1430 (2020).
20. F. Birey et al., Dissecting the molecular basis of human interneuron migration in forebrain assembloids from Timothy syndrome. Cell stem cell 29, 248-264 e247 (2022).
21. X. Meng et al., CRISPR screens in 3D assembloids reveal disease genes associated with human interneuron development. bioRxiv, 2022.2009.2006.506845 (2022).
22. A. Shiraishi, K. Muguruma, Y. Sasai, Generation of thalamic neurons from mouse embryonic stem cells. Development 144, 1211-1220 (2017).
23. Y. Xiang et al., hESC-Derived Thalamic Organoids Form Reciprocal Projections When Fused with Cortical Organoids. Cell stem cell 24, 487-497 e487 (2019).
24. A. M. Pasca et al., Functional cortical neurons and astrocytes from human pluripotent stem cells in 3D culture. Nat Methods 12, 671-678 (2015).
25. J. Andersen et al., Generation of Functional Human 3D Cortico-Motor Assembloids. Cell 183, 1913-1929 e1926 (2020).
26. S. P. Paşca et al., A nomenclature consensus for nervous system organoids and assembloids. Nature 609, 907-910 (2022).
27. H. J. Kang et al., Spatio-temporal transcriptome of the human brain. Nature 478, 483- 489 (2011).
28. J. S. Fleck et al., Resolving organoid brain region identities by mapping single-cell genomic data to reference atlases. Cell stem cell 28, 1148-1159 e1148 (2021).

29. Q. Guo, J. Y. Li, Defining developmental diversification of diencephalon neurons through single cell gene expression profiling. Development 146, dev174284 (2019).

30. A. Bhaduri et al., An atlas of cortical arealization identifies dynamic molecular signatures. Nature 598, 200-204 (2021).

31. R. Llinas, Y. Yarom, Electrophysiology of mammalian inferior olivary neurones in vitro. Different types of voltage‐dependent ionic conductances. The Journal of physiology 315, 549-567 (1981).

32. W. Guido, S.-M. Lu, S. M. Sherman, Relative contributions of burst and tonic responses to the receptive field properties of lateral geniculate neurons in the cat. Journal of neurophysiology 68, 2199-2211 (1992).

33. J. Huguenard, Low-threshold calcium currents in central nervous system neurons. Annual review of physiology 58, 329-348 (1996).

34. F. M. Dreyfus et al., Selective T-type calcium channel block in thalamic neurons reveals channel redundancy and physiological impact of I(T)window. The Journal of neuroscience : the official journal of the Society for Neuroscience 30, 99-109 (2010).

35. W. Choe et al., TTA-P2 is a potent and selective blocker of T-type calcium channels in rat sensory neurons and a novel antinociceptive agent. Mol Pharmacol 80, 900-910 (2011).

36. Y. Miura et al., Engineering brain assembloids to interrogate human neural circuits. Nature protocols 17, 15-35 (2022).

37. B. Zingg et al., AAV-mediated anterograde transsynaptic tagging: mapping corticocollicular input-defined neural pathways for defense behaviors. Neuron 93, 33- 47 (2017).

38. R. Etessami et al., Spread and pathogenic characteristics of a G-deficient rabies virus recombinant: an in vitro and in vivo study. Journal of General Virology 81, 2147-2153 (2000).

39. I. R. Wickersham et al., Monosynaptic restriction of transsynaptic tracing from single, genetically targeted neurons. Neuron 53, 639-647 (2007).

40. N. C. Klapoetke et al., Independent optical excitation of distinct neural populations. Nature Methods 11, 338-346 (2014).

41. R. M. Martin et al., Highly Efficient and Marker-free Genome Editing of Human Pluripotent Stem Cells by CRISPR-Cas9 RNP and AAV6 Donor-Mediated Homologous Recombination. Cell stem cell 24, 821-828 e825 (2019).

42. D. A. McCormick, D. Contreras, On the cellular and network bases of epileptic seizures. Annual review of physiology 63, 815-846 (2001).

43. M. Giraldo-Chica, N. D. Woodward, Review of thalamocortical resting-state fMRI studies in schizophrenia. Schizophrenia research 180, 58-63 (2017).

44. B. Baran et al., Increased thalamocortical connectivity in schizophrenia correlates with sleep spindle deficits: evidence for a common pathophysiology. Biological Psychiatry: Cognitive Neuroscience and Neuroimaging 4, 706-714 (2019).

45. P. Chen, E. Ye, X. Jin, Y. Zhu, L. Wang, Association between thalamocortical functional connectivity abnormalities and cognitive deficits in schizophrenia. Scientific reports 9, 2952 (2019).

46. B. Yao, S. F. Neggers, R. S. Kahn, K. N. Thakkar, Altered thalamocortical structural connectivity in persons with schizophrenia and healthy siblings. NeuroImage: Clinical 28, 102370 (2020).

47. V. Moreno-Juan et al., Prenatal thalamic waves regulate cortical area size prior to sensory processing. Nature communications 8, 14172 (2017).

48. N. Antón-Bolaños et al., Prenatal activity from thalamic neurons governs the emergence of functional cortical maps in mice. Science 364, 987-990 (2019).

49. O. Revah et al., Maturation and circuit integration of transplanted human cortical organoids. Nature 610, 319-326 (2022).

50. S. P. Paşca et al., Using iPSC-derived neurons to uncover cellular phenotypes associated with Timothy syndrome. Nature medicine 17, 1657-1662 (2011).

51. E. A. Susaki et al., Versatile whole-organ/body staining and imaging based on electrolyte-gel properties of biological tissues. Nature communications 11, 1982 (2020).

52. S. A. Sloan, J. Andersen, A. M. Pasca, F. Birey, S. P. Pasca, Generation and assembly of human brain region-specific three-dimensional cultures. Nature protocols 13, 2062-2085 (2018).

53. A. E. Trevino et al., Chromatin and gene-regulatory dynamics of the developing human cerebral cortex at single-cell resolution. Cell 184, 5053-5069. e5023 (2021).

54. H. Wu et al., Mapping the single-cell transcriptomic response of murine diabetic kidney disease to therapies. Cell metabolism 34, 1064-1078. e1066 (2022).

55. T. Stuart et al., Comprehensive integration of single-cell data. Cell 177, 1888-1902. e1821 (2019).

56. N. C. Klapoetke et al., Independent optical excitation of distinct neural populations. Nature methods 11, 338-346 (2014).

57. A. P. Buccino et al., SpikeInterface, a unified framework for spike sorting. Elife 9, e61834 (2020).

58. F. Birey et al., Dissecting the molecular basis of human interneuron migration in forebrain assembloids from Timothy syndrome. Cell stem cell 29, 248-264. e247 (2022).

Acknowledgements

We thank members of the Paşca laboratory at Stanford University for scientific inputs and the Stanford Wu Tsai Neurosciences Virus Core for production of AAVs. This work was supported by the Welcome Leap 1kD Program (to S.P.P.), Stanford Brain Organogenesis Big Idea Grant from the Wu Tsai Neurosciences Institute (to S.P.P.), the NYSCF Robertson Stem Cell Investigator Award (to S.P.P.), the Kwan Research Fund (to S.P.P.), the Senkut Research Funds (to S.P.P.), the Ludwig Foundation (to S.P.P.), the Chan Zuckerberg Initiative Ben Barres Investigator Award (to S.P.P.), the CZ BioHub Investigator Program (to S.P.P.), a fellowship from the National Research Foundation of Korea (to J.K), Stanford Medicine Dean's Fellowships (to Y.M., F.B. and X.M.), and Stanford Maternal & Child Health Research Institute (MCHRI) Postdoctoral Fellowships (to Y.M., O.R., F.B.). This paper was typeset with the bioRxiv word template by @Chrelli: www.github.com/chrelli/bioRxiv-word-template

Author contributions

J.K., Y.M. and S.P.P. conceived the project and designed experiments. J.K. carried out the differentiation experiments, characterization of hDiO, extracellular recording and disease modeling. Y.M. performed the differentiation experiments, single cell transcriptomics analyses and calcium imaging with optogenetics. M.-Y.L. performed whole cell recordings and analyses. O.R. and J.R.H. contributed to the development of the extracellular recording platform and the analysis pipeline. S.S. and M.H.P. generated and validated CACNA1G variant hiPS cell lines. F.B. and X.M. contributed to optimizing the protocol of regionalized organoids. M.V.T. performed the differentiation experiments and characterization of regionalized organoids. S.D.P analyzed optically-evoked calcium imaging results. J.A. and A.M.P. contributed to primary tissue experiments. J.K., Y.M. and S.P.P. wrote the manuscript with input from all authors.

Competing interest statement

Stanford University has filed a provisional patent application covering the generation of multi-region assembloids. M.H.P. is on the Board of Directors and holds equity in Graphite Bio. M.H.P. serves on the SAB of Allogene Tx and is an advisor to Versant Ventures.

Materials and Methods

Characterization and maintenance of hiPS cells

Human induced pluripotent stem (hiPS) cell lines used in this study were validated using methods described in previous studies (36, 50). Genome-wide SNP genotyping was performed using the Illumina genome-wide GSAMD-24v2-0 SNP microarray at the Children's Hospital of Philadelphia. Cultures were tested for and maintained Mycoplasma free. A total of 5 control hiPS cell lines derived from fibroblasts collected from 5 healthy subjects were used for experiments. One of the control hiPS cell lines was used to generate CACNA1G mutant hiPS cell lines (**Supplementary Table 1**). Approval for these experiments was obtained from the Stanford IRB panel and informed consent was obtained from all subjects.

Human primary tissue

Human brain tissue samples were obtained under a protocol approved by the Research Compliance Office at Stanford University.

Generation of hDiO and hCO from hiPS cells

For neural differentiation, hiPS cells were cultured on vitronectin-coated plates (5 µg/mL, Thermo Fisher Scientific, A14700) in Essential 8 medium (Thermo Fisher Scientific, A1517001). Cells were passaged every 4–5 days with UltraPure™ 0.5 mM EDTA, pH 8.0 (Thermo Fisher Scientific, 15575). For the generation of regionalized neural organoids, hiPS cells were incubated with Accutase® (Innovative Cell Technologies, AT104) at 37°C for 7 min and dissociated into single cells. Optionally, 1–2 days before organoid formation, hiPS cells were exposed to 1% dimethylsulfoxide (DMSO) (Sigma-Aldrich, 472301) in Essential 8 medium. For aggregation into organoids, approximately 3 x 10^6 single hiPS cells were added per AggreWell-800 well in Essential 8 medium supplemented with the ROCK inhibitor Y27632 (10 µM, Selleckchem, S1049), centrifuged at 100 *g* for 3 min, and then incubated at 37°C in 5% CO₂. After 24 hours, organoids consisting of approximately 10,000 cells were collected from each microwell by pipetting medium in the well up and down with a cut P1000 pipet tip and transferred into ultra-low attachment plastic dishes (Corning, 3262) in Essential 6 medium (Thermo Fisher Scientific, A1516401) supplemented with the SMAD pathway inhibitors dorsomorphin (2.5 µM, Sigma-Aldrich, P5499) and SB-431542 (10 µM, R&D Systems, 1614). For the first 5 days, Essential 6 medium was changed every day and supplemented with dorsomorphin and SB-431542. To generate hDiO, medium was supplemented with 1 µM CHIR (Selleckchem, S1263) on day 5. On day 6, neural organoids were transferred to neural media containing Neurobasal™-A Medium (Thermo Fisher Scientific, 10888022), B-27™ Supplement, minus vitamin A (Thermo Fisher Scientific, 12587010), GlutaMAX™ Supplement (1:100, Thermo Fisher Scientific, 35050079), Penicillin-Streptomycin (1:100, Thermo Fisher Scientific, 15070063), and supplemented with 1 µM CHIR (Selleckchem, S1263). On day 9 of differentiation, medium was supplemented with 100 nM SAG (day 9–15, Millipore Sigma, 566660-1MG), in addition to the compounds described above. Furthermore, on day 12 of differentiation, organoids were supplemented with 30 ng/mL BMP7 (PeproTech, 120-03P), in addition to the compounds described above. From day 19, to promote differentiation of the neural progenitors into neurons of hDiO, the neural medium was supplemented with brain-derived neurotrophic factor (BDNF; 20ng mL⁻¹, PeproTech, 450-02), NT3 (20 ng mL⁻¹, PeproTech, 450-03), L-Ascorbic Acid 2-phosphate Trisodium Salt (AA; 200 µM, FUJIFILM Wako Chemical Corporation, 323-44822), N6, 2'-O-Dibutyryladenosine 3', 5' -cyclic monophosphate sodium salt (cAMP; 50 µM, Millipore Sigma, D0627), cis-4, 7, 10, 13, 16, 19-Docosahexaenoic acid (DHA; 10 µM, Millipore Sigma, D2534), and 2.5 µM DAPT (only from day 19-25, STEMCELL Technologies, 72082). From day 43, only neural medium containing B-27™ Plus Supplement (Thermo Fisher Scientific, A3582801) was used and changed every 4 days. hCO were generated as previously described (*36*). From day 22, the neural media was supplemented with BDNF NT3, AA, cAMP, and DHA. From day 46, only neural media containing B-27™ Plus Supplement was used for media changes (every 4 days).

3D clearing and staining of a human primary brain sample

To optically clear and stain human primary tissue, the hydrophilic chemical cocktail-based CUBIC protocol was used (51). A dissected primary brain sample was fixed with a 4% PFA/PBS solution at 4°C overnight. The next day, the tissue was washed twice with PBS and incubated in Tissue-Clearing Reagent CUBIC-L (TCI, T3740) at 37°C for 2 days. The sample was washed three times with PBS, then nuclei were stained with RedDot2 (Biotium, #40061, 1:200 dilution) in PBS containing 500 mM NaCl at 37°C overnight. The tissue was washed twice with PBS, and once with a solution containing 10 mM HEPES, 10% TritonX-100, 200 mM NaCl and 0.5% BSA (HEPES-TSB) at 37°C for 2 hours, and then stained with anti-TCF7L2 (rabbit, Cell Signaling Technology, 2569S, 1:500 dilution) antibody in HEPES-TSB solution at 37°C for 2 days. Stained tissue was washed twice with 10% Triton X-100 in PBS and once with HEPES- TSB solution for 2 hours, and then incubated with a donkey anti-rabbit IgG (H&L) highly cross-adsorbed secondary antibody, Alexa Fluor 568 (Thermo Fisher Scientific, A10042, 1:300 dilution) in HEPES-TSB solution at 37°C for 2 days. The sample was washed twice with 10% Triton X-100 in PBS for 30 min and once with PBS for 1 hour. After washing with PBS, the sample was incubated with Tissue-Clearing Reagent CUBIC-R⁺ (TCI, T3741) at room temperature for 2 days for refractive index matching. The CUBIC-cleared tissue was imaged using a 10x objective on a Leica TCS SP8 confocal microscope.

Cryosection and immunocytochemistry

hCO, hDiO and assembloids were fixed in 4% paraformaldehyde (PFA)/phosphate buffered saline (PBS) overnight at 4°C. They were then washed in PBS and transferred to 30% sucrose/PBS for 2–3 days until the organoids/assembloids sank into the solution. Subsequently, they were rinsed in optimal cutting temperature (OCT) compound (Tissue-Tek OCT Compound 4583, Sakura Finetek) and 30% sucrose/PBS (1:1), embedded, and snap-frozen using dry ice. For immunofluorescence staining, 30 µm-thick sections were cut using a Leica Cryostat (Leica, CM1860). Cryosections were washed with PBS to remove excess OCT from the sections and blocked in 10% Normal Donkey Serum (NDS, Millipore Sigma, S30-100ML), 0.3% Triton X- 100 (Millipore Sigma, T9284-100ML), and 0.1% BSA diluted in PBS for 1 hour at room temperature. The sections were then incubated overnight at 4°C with primary antibodies diluted in PBS containing 2% NDS and 0.1% Triton X-100. PBS was used to wash the primary antibodies and the cryosections were incubated with secondary antibodies in PBS with the PBS containing 2% NDS and 0.1% Triton X-100 for 1 hour. The following primary antibodies were used for staining: anti-TCF7L2 (rabbit, Cell Signaling Technology, 2569S, 1:500 dilution), anti-GFP (chicken, GeneTex, GTX13970, 1:1000 dilution), anti-VGLUT2 (mouse, MilliporeSigma, MAB5540, 1:200 dilution), and anti-mCherry (rabbit, GeneTex, GTX128508, 1:1000 dilution). Alexa Fluor dyes (Life Technologies) were used at 1:1,000 dilution and nuclei were visualized with the Hoechst 33258 dye (Life Technologies, H3549, 1:10,000 dilution). Cryosections were mounted for microscopy on glass slides using Aqua-Poly/Mount (Polysciences, 18606), and imaged on a Leica TCS SP8 confocal microscope. Images were processed and analyzed using Fiji (NIH) and IMARIS (Oxford Instruments). hiPS cell cultures on vitronectin-coated glass coverslips were fixed in 4% PFA in PBS for 20 min and then rinsed twice for 5 min with PBS. Coverslips were blocked in 10% NDS, 0.3% Triton X-100, and 0.1% BSA diluted in PBS for 1 hour at room temperature and then incubated overnight at 4°C with a primary antibody, anti-OCT4 (C30A3) (rabbit, Cell Signaling Technology, 2840, 1:200 dilution) diluted in PBS containing 2% NDS and 0.1% Triton X-100. The next day, the solution (including the primary antibody) was washed with PBS three times, and the coverslips were incubated with secondary antibodies including Rhodamin Phalloidin 555 (Thermo Fisher Scientific, #R415, 1:400 dilution) in PBS with 2% NDS and 0.1% Triton X-100 for 1 hour. After PBS washing three times, nuclei were visualized with Hoechst 33258 (Life Technologies, H3549, 1:10,000 dilution), and the coverslips were mounted on glass slides using Aqua-Poly/Mount (Polysciences, 18606), and imaged on a Leica Stellaris 5 confocal microscope.

Real-time qPCR

mRNA from hCO and hDiO were isolated using the RNeasy Mini kit (Qiagen, 74106) with DNase I, Amplification Grade (Thermo Fisher Scientific, 18068-015). Template cDNA was prepared by reverse transcription using the SuperScript™ III First-Strand Synthesis SuperMix for qRT-PCR (Thermo Fisher Scientific, 11752250). qPCR was performed using the SYBR™ Green PCR Master Mix (Thermo Fisher Scientific, 4312704) on a QuantStudio™ 6 Flex Real- Time PCR System (Thermo Fisher Scientific, 4485689). Primers used in this study are listed in **Supplementary Table 2**.

Single cell RNA-seq library preparation and data analysis

Dissociation of hDiO and human tissue was performed as described previously (*52*, *53*). For the organoid dissociation, 4–5 randomly selected organoids were pooled to obtain single cell suspension from hDiO and then incubated in 30 U/mL papain enzyme solution (Worthington Biochemical, LS003126) and 0.4% DNase (12,500 U/mL, Worthington Biochemical, LS2007) at 37°C for 45 minutes. After enzymatic dissociation, organoids were washed with a solution including protease inhibitor and gently triturated to achieve a single cell suspension. Cells were resuspended in 0.04% BSA/PBS (Millipore-Sigma, B6917-25MG) and filtered through a 70 µm Flowmi Cell Strainer (Bel-Art, H13680-0070) and the number of cells were counted. To target 7,000 cells (for hDiO) and 10,000 cells (for primary sample) after recovery, approximately 11,600 and 16,000 cells were loaded, respectively, per lane on a Chromium Single Cell 3′ chip (Chromium Next GEM Chip G Single Cell Kit, 10x Genomics, PN-1000127) and cDNA libraries were generated with the Chromium Next GEM Single Cell 3′ GEM, Library & Gel Bead Kit v3.1 (10x Genomics, PN-1000128), according to the manufacturer's instructions. Each library was sequenced using the Illumina NovaSeq S4 2 × 150 bp by Admera Health. Quality control, UMI counting of Ensembl genes and aggregation of samples were performed by the 'count'

and 'aggr' functions in Cell Ranger software (version, Cellranger-6.0.1). Further downstream analyses were performed using the R package Seurat. Genes on the X or Y chromosome were removed from the count matrix to avoid biases in clustering due to the sex of the hiPS cell lines. Cells with more than 7,500 or less than 200 detected genes or with mitochondrial content higher than 15% were excluded. Genes that were not expressed in at least three cells were not included in the analysis. Gene expression was normalized using a global-scaling normalization method (normalization method, 'LogNormalize'; scale factor, 10,000), and the 2,000 most variable genes were selected (selection method, 'vst') and scaled (mean = 0 and variance = 1 for each gene) before principal component analysis. The top 10 principal components were used for clustering (resolution of 1.0), using the 'FindNeighbors' and 'FindClusters' functions, and for visualization with UMAP. Clusters were grouped based on the expression of known marker genes. Unbiased spatial mapping of the whole combined cluster was performed using VoxHunt (version, VoxHunt1.0.0). Briefly, the 50 most variable features from the ISH Allen Brain Atlas data of the E15.5 mouse brain were selected, and similarity maps were calculated. These maps were then plotted in the sagittal views. Comparison to BrainSpan transcriptomic data of micro-dissected human brain tissue at postconceptional weeks (PCW) 15–24 was also performed using VoxHunt with default settings. A circular plot was generated using the R package plot1cell (54). The R package scCustomize (v1.1.1) (https://samuel-marsh.github.io/scCustomize) was used to generate the UMAP for integrated samples ('FeaturePlot_scCustom' function), and to calculate the percentage of clusters per sample ('Percent_Expressing' function). Integration of hDiO and primary tissue samples scRNA-seq datasets was performed using a standard Seurat V3 integration workflow (55).

Viral labeling and live cell imaging

Viral infection of 3D neural organoids was performed as previously described (36). Briefly, neural organoids were transferred into a 1.5 mL Eppendorf tube containing 200 μL of neural media and incubated with the virus overnight at 37°C, 5% CO$_2$. Next day, 800 μL of fresh culture media was added. The following day, neural organoids were transferred into fresh culture media in ultra-low attachment plates (Corning, 3471, 3261). For live cell imaging, labeled hCO or hDiO or assembloids were transferred into a well of a Corning^{TM} 96-Well microplate (Corning, 4580) in 150 μL of neural media or on a 20 mm glass coverslip in a 35 mm glass bottom well (Cellvis, D35-20-0-N) and incubated in an environmentally controlled chamber for 15–30 min before imaging on a Leica TCS SP8 or Leica Stellaris 5 confocal microscope. The viruses used in this study were: AAV-DJ-hSYN1::EYFP (Stanford University Neuroscience Gene Vector and Virus Core, GVVC-AAV-16), AAV-DJ-hSYN1::mCherry (Stanford University Neuroscience Gene Vector and Virus Core, GVVC-AAV-17), AAV-DJ-EF1-DIO-tdTomato (Stanford University Neuroscience Gene Vector and Virus Core, GVVC-AAV-169), AAV1-hSYN1::Cre (Addgene, #105553), AAV1-hSYN1::ChrimsonR-tdTomato (56) (Addgene, #59171-AAV1) and AAV-DJ-hSYN1::GCaMP6f (Stanford University Neuroscience Gene Vector and Virus Core, GVVC-AAV-96).

Generation of thalamo-cortical assembloids

To generate thalamo-cortical assembloids, hDiO and hCO were separately generated from hiPS cells and assembled 2–3 days later by placing them in proximity in 1.5 mL Eppendorf tubes (for 3 days) in an incubator. Medium was gently replaced on day 2 and on the third day, assembloids were transferred using a cut P1000 pipette tip in 24-well ultra-low attachment plates in the neural media described above. Media was changed every 4 days.

Axon projection imaging in thalamo-cortical assembloids

AAV-DJ-hSYN1::EYFP$^+$ cells projecting into hCOs from hDiOs or AAV-DJ-hSYN1::mCherry$^+$ cells projecting into hDiOs from hCOs were imaged under environmentally controlled conditions in intact thalamo-cortical assembloids using a Leica TCS SP8 or Leica Stellaris 5 confocal microscope with a motorized stage. Assembloids were transferred to a well in a 24-well glass bottom plate (Cellvis, P24-0-N) in culture medium, and incubated in an environmentally controlled chamber for 15–30 min before imaging. The percentages of EYFP coverage in hCOs and that of mCherry coverage in hDiOs were quantified with Fiji (ImageJ, version 2.1.0, NIH) from projected z-stack images.

Viral transsynaptic tracing in thalamo-cortical assembloids

For anterograde tracing in thalamo-cortical assembloids, hCO were infected with AAV-DJ-EF1α::DIO-tdTomato and AAV-DJ-hSYN1::EYFP, while hDiO were infected with AAV1-hSYN1::Cre. For retrograde tracing in thalamo-cortical assembloids, hCO were infected with rabies-ΔG-Cre-EGFP and AAV-DJ-EF1α::CVS-G-WPRE-pGHpA, while hDiO were infected with AAV-DJ-EF1α::DIO-tdTomato. After viral infection, hCO and hDiO were assembled and maintained in culture with medium changes every 4 days. After 15–38 days after assembly, z-stack images from assembloids were captured using a Leica TCS SP8 or Leica Stellaris 5 confocal microscope and analyzed in Fiji (ImageJ, version 2.1.0, NIH).

Optogenetic stimulation and calcium imaging

Thalamo-cortical assembloids expressing AAV1-hSYN1::ChrimsonR-tdTomato in hDiO and AAV-DJ-hSYN1::GCaMP6f in hCO were placed on a 20 mm coverslip glass (in a 35 mm glass-bottom plate) in neural medium, and imaged using a 10x objective on a Leica TCS SP8 confocal microscope. For optogenetic stimulation, ChrimsonR-tdTomato$^+$ cells in hDiO were activated with 590 nm light using an optical fiber-coupled LED (Thorlabs, M590F3). GCaMP6f was imaged at a frame rate of 14.7 frames/sec. Imaging experiments were 1500 frame-long and one 590 nm of LED (71 ms) was applied every 150 frames. The pulse was generated by a Cyclops LED connected to the Leica TCS SP8. Results were analyzed with Fiji (NIH) and MATLAB (version R2022a, MathWorks). After ROI registration, raw time series were transformed to relative changes in fluorescence: $\Delta F/F_{(t)} = (F_{(t)}-F_0)/F_0$. To verify whether the optically-evoked responses were time-locked to the LED stimulation, $\Delta F/F$ responses were compared to $\Delta F/F$ values at randomly selected time points. For time-locked $\Delta F/F$, the amplitudes of $\Delta F/F$ from each cell were calculated as the maximum $\Delta F/F$ value within a 20–30 frames window (1,360 to 2,040 ms) following LED stimulation (minus the mean of the baseline 1 sec before the stimulation). The median was used to exclude shape-dependent artifacts.

Extracellular recording and analyses

Neural organoids or assembloids were embedded into 3% low-melting gel agarose (IBI Scientific, IB70056) and transferred into artificial cerebrospinal fluid (aCSF) containing 124 mM NaCl, 3 mM KCl, 1.25 mM NaH$_2$PO$_4$, 1.2 mM MgSO$_4$, 1.5 mM CaCl$_2$, 26 mM NaHCO$_3$ and 10 mM D-(+)-glucose with addition of GlutaMAX (Gibco, 35050061). Embedded neural organoids or assembloids were placed on a Brain Slice Chamber-2 (Scientific Systems Design Inc, S- BSC2) and perfused with aCSF (bubbled with 95% O$_2$ and 5% CO$_2$) at 37°C. Temperature was controlled and retained at 37°C by connecting to a Proportional Temperature Controller PTC03 (AutoMate Scientific, S-PTC03). Acute 32 channel P-1 probe with 2 shanks (Cambridge Neuro-Tech, ASSY-37 P-1) were connected to an Acute probe adaptor; 32 channels Samtec to Omnetics (Cambridge NeuroTech, ADPT A32-Om32). Signals were acquired using the Intan 1024ch recording controller (Intan Technologies) at 30000 Hz. For optogenetic stimulation, the pulses were generated by the pulse stimulator (Grass Instrument, S48) connected to a Cyclops LED driver and a fiber cable (Thorlabs, M126L01). A 20 sec recording protocol (a 19 sec-long light-off phase followed by a 1 sec-long constant 590 nm illumination) was repeated in 9 trials. Raw recording data were processed using Intan Technologies code and analyzed by MATLAB (version R2022a, MathWorks). To count hypersynchronous events, the continuous wavelet transform analysis (CWT) function in MATLAB was applied to median signal values obtained from 32 channels; this produced time-frequency decomposition and averaged absolute values of frequency band for spike activity (500 ~ 2000 Hz), referred as spike power, were z-scored. The hypersynchronous events were defined as having a z-scored spike energy was greater than 10 and a median value of the next 250 ms greater than 2.2. The inter-activity interval was set to 5 sec. Local field potential (LFP) signals were obtained by bandpass filtering from 1 Hz to 100 Hz with 2nd-order Butterworth filter. To sort single units, high-pass (350 ~ 2000 Hz) filtering and common median referencing were applied to raw recording data. Single unit sorting processes were performed with Tridesclous (version 1.6.4, https://tridesclous.readthedocs.io/en/latest) using Spikeinterface package (57) (version 0.7.6, https://github.com/SpikeInterface) and sorted units were manually curated using Phy2 (https://github.com/cortex-lab/phy). To compare the evoked activity in a-hCO by optogenetic stimulation, spike power of a-hCO was extracted as described above, and the first 250 ms of each light stimulation were averaged from 9 trials (per assembloids). Units were considered

responding if the mean firing rates during the light-on phase was significantly (by paired *t*-test) increased over the light-off phase over 9 trials.

Patch-clamp recordings

Patch-clamp recordings were performed as described previously (*58*). Briefly, hDiOs were infected with AAV-DJ-hSYN1::EYFP and placed on top of cell culture inserts (0.4 mm pore size; Corning, 353090) that were positioned in 6-well plates to form flattened organoids. Recordings were typically made on day 128–135. EYFP[+] neurons were identified with an upright slice scope microscope (Scientifica) equipped with an Infinity2 CCD camera and Infinity Capture software (Teledyne Lumenera). Recordings were performed with borosilicate glass electrodes with a resistance of 7–10 MΩ. BrainPhys neuronal media (STEMCELL Technologies, 05790) was used as an external solution. The internal solution contained 127 mM K-gluconate, 8 mM NaCl, 4 mM MgATP, 0.3 mM Na_2GTP, 10 mM HEPES and 0.6 mM EGTA, pH adjusted to 7.2 with KOH (290 mOsm). For hyperpolarization–induced rebound spikes, cells were held at –50 mV and hyperpolarizing current steps (1 sec) with an increment of 10 pA from –100 pA to 0 pA were tested. For barium current recordings, cells were recorded in the presence of TTX (0.5 μM) to block sodium currents, and were held at –70 mV in voltage-clamp and voltage steps (1 sec, from –90 mV to +40 mV) were given with an increment of 10 mV. The external solution contained 100 mM NaCl, 3 mM KCl, 2 mM $MgCl_2$, 4 mM $BaCl_2$, 25 mM TEA-Cl, 4 mM 4-AP, 10 mM HEPES, 20 mM glucose, pH 7.4 with NaOH, 300 mOsm. The internal solution contained 110 mM $CsMethylSO_3$, 30 mM TEA-Cl, 10 mM EGTA, 4 mM MgATP, 0.3 mM Na_2GTP, 10 mM HEPES, 5 mM QX314-Cl, pH 7.2 with CsOH, 290 mOsm. Leak subtraction was implemented in MultiClamp 700B software. *I-V* curves were fitted in Origin (OriginPro 2021b, OriginLab) with a Boltzmann exponential function: $I = G_{max} * (V - E_{Ba}) / \{1 + \exp[(V_{0.5} - V) / K]\}$, where G_{max} is the maximal conductance of the calcium channels, E_{Ba} is the reversal potential of barium estimated by the curve-fitting program, $V_{0.5}$ is the potential for half-maximal steady-state activation of the barium current, and K is a voltage-dependent slope factor. Data were acquired with a MultiClamp 700B Amplifier (Molecular Devices) and a Digidata 1550B Digitizer (Molecular Devices), low-pass filtered at 2 kHz, digitized at 20 kHz and analyzed with pCLAMP (version 10.6, Molecular Devices). Cells were given a –10 mV hyperpolarization (100 ms) every 10 sec to monitor input resistance and access resistance. Cells were excluded for analysis if either value changed by > 30%. The liquid junction potential was not corrected.

Genome editing of hiPS cells to generate *CACNA1G* variant hiPS cell lines

For generating *CACNA1G* knockout cells, hiPS cells were genome edited with dual sgRNA:Cas9 RNP complexes to create a frameshift deletion in exon 1 of the *CACNA1G* gene. For this, 24 hours prior to gene editing, hiPS cells were treated with 10 μM ROCK inhibitor (Y-27632). hiPS cells (at 70–80% confluence) were detached from the culture plate and dissociated into single cells by following incubation with Accutase (Innovative Cell Technologies) for 5 min at 37°C. ROCK inhibitor-supplemented mTeSR1 media was added to neutralize Accutase. Concurrent with the cell dissociation process, the Cas9 RNP complex was prepared. The RNP complex for each of the two sgRNA was formed separately by combining 5 μg of HiFi Cas9 (Aldevron) and 1.75 μg of sgRNA (Synthego) (molar ratio of 1:2.5 (Cas9: sgRNA)) for 10 min at room temperature. The two RNP complexes were then combined and diluted in 20 μL of P3 Primary Cell Nucleofector Solution (Lonza). To electroporate the Cas9 RNP complexes into hiPS cells, 500,000 cells were mixed with the nucleofection solution containing the RNPs and were electroporated in one well of a 16-well Nucleocuvette Strip using the program CA137 in 4D Nucleofector system (Lonza). Following electroporation, cells were plated into two wells of a Matrigel-coated 24-well plate containing 500 μL of mTeSR1 media supplemented with 10 μM Y-27632. The next day, the cells were switched to fresh 10 μM Y-27632

supplemented mTeSR1 medium. After 48 hours of electroporation, cells were cultured in mTeSR1 medium without Y-27632. To isolate single cell clones, edited hiPS cells were detached and dissociated into single cells using Accutase. After dissociation, Accutase was neutralized with mTeSR1 medium and cells were counted. Single cells were then plated at a density of 250 cells per well of a 6-well plate in mTeSR1 medium supplemented with 1X CloneR solution (Stem Cell 28 Technologies) and incubated for 48 hours. At day 2, cells were switched to fresh mTeSR1 with 1X CloneR and incubated for 2 more days. From day 4, cells were cultured in mTeSR1 medium without CloneR. At day 7–10, single cell colonies were picked by scraping and were then plated in one well of a 48-well plate in mTeSR1 medium with 10 μM Y-27632. From the next day, single cell clones were maintained in mTeSR1 medium. The genotype of edited clones was determined by PCR with the primer set FW, 5'-GAAGCGAAGAAGCCGGAACAA-3'; RV, 5'-CCCGGGTTACCGACGAGA-3', and Sanger sequencing (Mclab or Genewiz) was performed with the following primers: FW, 5'- TAGAGCCCAC-CAGATGTGCC-3', RV, 5'- GGGCTCACTTCGACTCACC-3'. ICE analysis (https://ice.synthego.com/) was performed on sequencing chromatograms to determine the genotype. Synthetic sgRNA (with 2'-O-methyl 3'- phosphorothioate modified nucleotides at the three terminal positions at both the 5' and 3' ends) were purchased from Synthego. The genomic target sequences of the sgRNAs were G2: 5'-GATATGGGTTACAGACCGTG-3' and G3: 5'-TTGAGCCGCATGAAGCTCCG-3'. sgRNAs with high specificity scores were selected based on the COSMID tool (https://crispr.bme.gatech.edu). For generating the *CACNA1G* M1531V isogenic hiPS cells, AAV6/Cas9 RNP genome editing was performed, as described previously (*41*). Cas9:sgRNA RNP complex was nucleofected into hiPS cells as above. For this gene editing, an sgRNA with a target site close to the genomic region encoding for M1531 was used. After nucleofection, the cells were plated in 10 μM Y-27632-supplemented mTeSR1 medium with a AAV6 homology directed repair donor template at an MOI of 5K, and incubated at 37°C for 24 hours. Cells were then switched to fresh mTeSR1 medium with 10 μM Y-27632. Next day forward, cells were maintained in mTeSR1 medium. Genome edited hiPS cells were subjected to single cell cloning as described above. The single cell clones were genotyped using PCR, Sanger sequencing and ICE analysis. The primer set used for PCR was FW: 5'-CATGGGGTCTCTAGCCGTC-3' and RV: 5'- GAAAGACTAC-CGACCAGCCA-3' and the Sanger sequencing primer used was, 5'-CTGTGGGCAGCTCCTAAGACA-3'. ICE analysis was done on the sequencing chromatograms to determine the frequency of control, INDEL and HDR in the single cell clones. The genomic target sequence of the sgRNA used was: 5'-CATCTCGTTCCTGCTCATTG-3'. AAV6 donor template was designed and constructed with 1.2 kb homology arms, incorporating the intended M1531V variant and silent variants on the sgRNA target site to avoid recutting post-HDR. The donor template sequence was cloned into a pAAV-MCS backbone using NEB hifi DNA assembly kit. AAV6 was packaged in 293T cells, purified using the AAVpro purification kit (Takara), and the titer was determined using ddPCR, as described previously (*41*).

Statistics

Data are presented as mean ± s.e.m. unless described otherwise. Raw data were tested for normality of distribution and statistical analyses were performed by paired and unpaired *t*-test (two-tailed), Mann-Whitney test, Wilcoxon test, one-way *ANOVA*, or Kruskal-Wallis test with multiple comparison tests depending on the dataset. Sample sizes were estimated empirically. GraphPad Prism Version 9.3.1 or MATLAB (version R2022a, MathWorks) were used for statistical analyses.

Code availability

The codes used for calcium imaging and electrophysiology analyses in this study are available on request from the corresponding author.

Supplementary Figure 1. Generation of human diencephalic organoids (hDiO). (**A**) Protocols to generate hCO and hDiO from human pluripotent stem cells. (**B**) Normalized expression of *TCF7L2*, *OTX2* and *FOXG1* from hCO and hDiO at day 18 of differentiation as measured by qPCR ($n = 24$ organoids from 8 differentiation of 4 hiPS cell lines; Mann-Whitney test, ***$P = 0.0006$, ****$P < 0.0001$). (**C**) UMAP of hDiO cells by hiPS cell line and distribution of cell clusters. (**D**) UMAP showing cell type markers. (**E**) VoxHunt analysis per cluster. (**F**) UMAP indicating regional thalamic markers. (**G**) *In vitro* and *in vivo* counterpart comparison of scRNAseq data. UMAP including hDiO and PCW16–20 primary thalamus. Abbreviations: VFC; ventrolateral frontal cortex, V1C; primary visual cortex, STR; striatum, STC; Posterior superior temporal cortex, S1C; primary somatosensory cortex, OFC; orbital frontal cortex, MFC; medial frontal cortex, MD; mediodorsal nucleus of thalamus, M1C; primary motor cortex, ITC; inferior temporal cortex, IPC; posterior inferior parietal cortex, HIP; hippocampus, DFC; dorsolateral prefrontal cortex, CBC; cerebellar cortex, AMY; amygdala, A1C; primary auditory temporal cortex.

Supplementary Figure 2. Electrophysiological characterization of hDiO neurons. (**A**) Representative traces showing spontaneous firing of EYFP-labeled hDiO neurons in cell-attached (left, voltage clamp) and whole-cell recordings (right, current clamp). (**B**) Representative examples and *I–V* curves (barium currents) of hDiO neurons (held at −70 mV, voltage steps from −90 mV to +40 mV with an increment of 10 mV). (**C**) Representative barium current traces (left) and zoomed-in traces (right). Traces for individual trials and the average are shown in grey and pink respectively. (**D**) Representative barium current traces for depolarization at −30 mV at baseline (left), after TTA-P2 treatment (right) and quantitation of rebound spikes (*n* = 5 cells for baseline and TTA-P2; paired *t*-test, *P = 0.041).

Supplementary Figure 3. Optogenetics and extracellular recording of thalamo-cortical assembloids. (**A**) Image of a custom extracellular recording platform for organoids and assembloids (left). Organoids are embedded into 3% agarose (top, left and right) and then placed on the Brain Slice Chamber (bottom, right). (**B**) Design of high-density probe that includes 32 channels in 2 shanks. (**C**) Process for single unit sorting. (**D**) Schematic illustration extracellular recordings combined with optogenetic stimulation of hDiO expressing ChrimsonR-tdTomato. (**E**) Representative trace showing optogenetic stimulation in artificial CSF (baseline; top) and TTX treatment (bottom), and (**F**) quantification of firing rates in hCO, before and after TTX treatment (n = 7 units; paired t-test, *P = 0.0213). (**G**) Schematic showing a control experiment with extracellular recording of a-hCO following light application of to a-hDiO not expressing an opsin in thalamo-cortical assembloids. (**H**) Representative raster plot aligned with light stimulation (top) and randomly selected time point (bottom). (**I**) Quantification of firing rates in a-hCO during light on and off phases (n = 13 units from 1 differentiation experiment of 2 hiPS cell lines; paired t-test, n.s., not significant).

Supplementary Figure 4. CRISPR/Cas9 based generation of hiPS cell lines carrying *CACNA1G* variants. (**A**) UMAP showing expression of *CACNA1G* (top) and *NTNG1* (bottom) in hDiO at day 100 of differentiation. (**B**) UMAP showing expression of *CACNA1G* (top) and *SNAP25* (bottom) in combined hDiO and human primary thalamus. (**C**) Alignment of the *CACNA1G* control and donor template (AAV6) sequences around the M1531V variant site. AAV6 donor sequence was designed to introduce the desired M1531V variant (in red) and silent variants (in green) to prevent recutting after gene editing. G11 denotes the target site of the sgRNA used for gene editing. (**D**) Allelic distribution of control, INDEL and HDR (M1531V) frequencies based on ICE analysis in the gene edited single cell hiPS cell clones. Clone#7 was used in this study. (**E**) Sanger sequencing results of parental hiPS cells (top) and genome edited *CACNA1G* M1531V hiPS cells (bottom). (**F**) PCR on gene edited hiPS cell clones amplifying exon 1 of *CACNA1G*. Clones#1, #2 and #3 demonstrated a deletion in exon 1. (**G**) Allelic frequency of INDEL and KO based on ICE analysis on the Sanger sequences of PCR amplicons from **F**. This analysis confirmed homozygous KO due to a frameshift deletion of 178/179-bp in clones #1 and #2, which were used in this study. (**H**) Uncropped gel image of **F**. (**I**) Sanger sequencing results of parental hiPS cells (top) and *CACNA1G* KO hiPS cell lines, clone#1 and #2 (middle and bottom). (**J**) Immunostaining of pluripotency marker OCT4 (yellow), F-actin (magenta) and Hoechst (cyan). Scale bar: 100 μm. (**K**) Regionalized organoids generated from the edited hiPS cells. Scale bar: 1 mm.

Supplementary Figure 5. Gain- and loss-of-function *CACNA1G* variants in thalamo-cortical assembloids. (**A**) Representative and average barium current traces induced by depolarization (from –70 mV to –30 mV) from control and *CACNA1G* KO EYFP+ neurons in flattened hDiOs (control, $n = 4$ cells; *CACNA1G* KO, $n = 6$ cells; Mann-Whitney test, **$P = 0.0095$). (**B**) Percentage of hDiOs showing hypersynchronous activity in control or M1531V hDiO (left), and TTA-P2 treated hDiO (right). (**C**) Percentage of hCO-hDiOs showing hypersynchronous activity in control or M1531V assembloids (left: a-hCO and right: a-hDiO). (**D**) Representative traces showing co-occurrence of hypersynchronous activity in M1531V assembloids. Arrows indicate the hypersynchronous activity. (**E**) Quantification of spike power aligned to onset of hypersynchronous activity and aligned to random time point (top: a-hCO, bottom: a-hDiO). (**F**) Schematic illustration of the experimental design for hDiO optogenetic and extracellular recording (left). Quantification of z-scored power of spike band frequency in a-hDiO during light stimulation (control, $n = 9$ assembloids; *CACNA1G* KO, $n = 10$ organoids; Mann-Whitney test, n.s., not significant).